



Document downloaded from the institutional repository of the University of Alcalá: <https://ebuah.uah.es/dspace/>

This is a postprint version of the following published document:

Barriopedro, P. et al., 2020. Successive Protonation and Methylation of Bridging Imido and Nitrido Ligands at Titanium Complexes. *Inorganic chemistry*, 59(11), pp.7631–7643.

Available at <https://doi.org/10.1021/acs.inorgchem.0c00550>

© 2020 American Chemical Society

(Article begins on next page)



This work is licensed under a

Creative Commons Attribution-NonCommercial-NoDerivatives
4.0 International License.

Successive Protonation and Methylation of Bridging Imido and Nitrido Ligands at Titanium Complexes


Paula Barriopedro,[†] Jorge Caballo,[†] Miguel Mena,^{†,‡} Adrián Pérez-Redondo,^{†,‡} and Carlos Yélamos^{*,†,‡}

[†] Departamento de Química Orgánica y Química Inorgánica, Universidad de Alcalá. 28805 Alcalá de Henares-Madrid (Spain). E-mail: carlos.yelamos@uah.es

[‡] Instituto de Investigación Química “Andrés M. del Río” (IQAR), Universidad de Alcalá. 28805 Alcalá de Henares-Madrid (Spain).

Synopsis:

Successive electrophilic attacks of ROTf (R = H, Me; OTf = OSO₂CF₃) reagents at imido and nitrido moieties of polynuclear titanium(IV) complexes enable isolation and structural characterization of a series of intermediates with nitrogen-based ligands (μ_n -N, μ_n -NH, μ -NH₂, NH₃) en route to ammonium salts [NR₄]OTf or [NR₄][Ti(η^5 -C₅Me₅)(OTf)₄] formation.



Abstract:

The reactions of nitrido complexes $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)(\mu\text{-NH})\}_3(\mu_3\text{-N})]$ (**1**) and $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-N})_4]$ (**2**) with electrophilic reagents ROTf (R = H, Me; OTf = OSO₂CF₃) in different molar ratios have allowed the structural characterization of a series of titanium intermediates en route to ammonium salts [NR₄]OTf or [NR₄][Ti(η⁵-C₅Me₅)(OTf)₄] formation. The treatment of the trinuclear imido-nitrido complex **1** with 5.5 equiv of triflic acid in toluene at room temperature led to the dinuclear complex $[\text{Ti}_2(\eta^5\text{-C}_5\text{Me}_5)_2(\mu\text{-N})(\text{NH}_3)(\mu\text{-O}_2\text{SO}_2\text{CF}_3)_2(\text{OTf})]$ (**3**) and [NH₄]OTf. Compound **3**, along with the ammonium salts [NMe₄]OTf and [NMe₄][Ti(η⁵-C₅Me₅)(OTf)₄] (**5**), was also obtained in the reaction of **1** with 8 equiv of methyl triflate in toluene at 100 °C. A trinuclear complex $[\text{Ti}_3(\eta^5\text{-C}_5\text{Me}_5)_3(\mu\text{-N})(\mu\text{-NH})_2(\mu\text{-O}_2\text{SO}_2\text{CF}_3)(\text{OTf})]$ (**4**), intermediate in the formation of **3**, was isolated in the treatment of **1** with 4 equiv of MeOTf, although compound **4** was prepared in better yield by treatment of **1** with Me₃SiOTf (2 equiv). Addition of a large excess of MeOTf or HOTf reagents to solutions of **3** resulted in the clean formation of ammonium salts [NR₄][Ti(η⁵-C₅Me₅)(OTf)₄] (R = Me (**5**), H (**6**)). Treatment of the tetranuclear nitrido complex $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-N})_4]$ (**2**) with 1 equiv of ROTf in toluene afforded the precipitation of the ionic compounds $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-N})_3(\mu_3\text{-NR})][\text{OTf}]$ (R = H (**8**), Me (**9**)), while a large excess of HOTf led to the formation of $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-N})_3(\mu_3\text{-NH})][\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)(\text{OTf})_4(\text{NH}_3)]$ (**10**) by rupture of a fraction of tetranuclear molecules. Complex **2** reacted with 1 equiv of $[\text{M}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\text{H}]$ (M = Mo, Cr) via hydrogenation of one nitrido ligand to give the molecular derivative $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-N})_3(\mu_3\text{-NH})]$ (**11**) and $[\{\text{M}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\}_2]$, while a second 1 equiv of $[\text{M}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\text{H}]$ produced

ionic compounds $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-N})_2(\mu_3\text{-NH})_2][\text{M}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3]$ (M = Mo (**12**), Cr (**13**)) by protonation of other nitrido group. The X-ray crystal structures of **3**, **4**, **5**, **9**, **10**, and **13** were determined.

Introduction

Transition metal nitrido and imido species are of interest because of their implications as key intermediates in biological and industrial dinitrogen fixation.^{1,2} Since the seminal contributions of Cummins and co-workers with molybdenum(III) trisamido complexes,³ the reductive cleavage of dinitrogen to form nitrido ligands by low-valent transition metal species is well-documented in the literature.^{1,4} Most of the mid-transition metal (Groups 6–8) complexes show the nitrido ligands derived from N₂ as a terminal functionality M≡N,⁵ although singular polynuclear iron species with bridging nitrido/imido ligands have been reported by Holland⁶ and Murray⁷ groups. In contrast, early transition metal (Groups 4 and 5) systems typically contain bridging nitrido (μ_n -N) moieties,^{8,9,10} although several examples with the M≡N functionality (M = Ti,¹¹ Zr,¹² V,¹³ and Nb¹⁴) have been described. In early transition metal nitrido complexes, the high stability of the resulting metal–nitrogen bonds makes difficult the subsequent functionalization and eventual release of nitrogen-containing products from the coordination sphere of the azaphilic transition element.¹⁵ In general, those nitrido derivatives do not react with H₂, and strong acids or silylating agents are required for release of the nitrogen material. For instance, nitrido complexes are typically reacted with excess ethereal HCl to give ammonium chloride NH₄Cl as diagnostic evidence of nitrogen incorporation from N₂. Alternative methods based on hydrogenolysis of metal–nitrogen bonds by proton-coupled electron transfer (PCET) have been recently explored by Chirik.¹⁶


Noteworthy, Hou and co-workers have described a series of low-valent polyhydride titanium complexes capable of performing the cleavage of the N₂ molecule under mild conditions and the subsequent partial hydrogenation of the resulting μ_n -nitrido ligands to

give μ_n -NH moieties.¹⁷ The reactivity of bridging nitrido μ_n -N and imido μ_n -NH functionalities on multimetallic complexes is important to understand in the context of the mechanism of Haber-Bosch ammonia synthesis from N_2 and H_2 .^{6,17} In this heterogeneous catalyzed process, the rate-determining step is the $N\equiv N$ bond cleavage in the surface of the catalyst to form two nitrides bridging several metal centers.¹⁸ Subsequent hydrogenation of the nitrides produces imide NH, amide NH_2 , and ammine NH_3 groups adsorbed on the metallic surfaces before ammonia elimination occurs. Well-defined species that resemble the proposed intermediates at the molecular level are relevant to the understanding of these elementary steps.

Polynuclear nitrido complexes of titanium,¹⁹ zirconium,²⁰ and tantalum²¹ with bridging nitrido μ_n -N, imido μ_n -NH, or amido μ - NH_2 ligands are usually obtained by the reaction of organometallic derivatives with excess ammonia. Representative examples for titanium(IV) derivatives are the trinuclear imido-nitrido [$\{Ti(\eta^5-C_5Me_5)(\mu-NH)\}_3(\mu_3-N)$]²² (**1**) and the tetranuclear [$\{Ti(\eta^5-C_5Me_5)\}_4(\mu_3-N)_4$] (**2**) species,²³ which are easily prepared by ammonolysis of $[Ti(\eta^5-C_5Me_5)X_3]$ ($X = Me, NMe_2$) mononuclear complexes. Compounds **1** and **2** are structurally related to the systems isolated by Hou and co-workers from dinitrogen splitting,¹⁷ as well as to those reported by us in the exposure of $[Ti(\eta^5-C_5Me_5)Me_3]$ to forming gas (H_2/N_2 mixture) under ambient conditions.²⁴


As part of a project devoted to the development of new reactivity patterns of the bridging ligands of the polynuclear complexes **1** and **2**,^{25,26} we became interested in studying their reaction with electrophiles. In particular, we reported on the reaction of **1** with one equivalent of electrophilic reagents ROTf ($R = H, Me, Me_3Si$) to generate polynuclear complexes by selective functionalization of the imido groups (Scheme 1).^{27,28}

As a logical continuation of this study, herein we describe the reactivity of **1** with higher molar ratios of ROTf reagents and the characterization of titanium nitrido intermediates en route to the ammonium salts $[\text{NR}_4]\text{OTf}$ or $[\text{NR}_4][\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)(\text{OTf})_4]$ formation.



Scheme 1. Reaction of **1** with 1 equiv of ROTf

This study with electrophiles ROTf has been also extended to the tetranuclear nitrido complex **2**, following preliminary results on the protonation or hydrogenation of a $\mu_3\text{-N}$ nitrido ligand of this compound with one equivalent of HOTf or two equivalents of ammonia borane (Scheme 2).²⁹ In addition, here we report on the use of group 6 hydride derivatives $[\text{M}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\text{H}]$ in the hydrogenation and protonation of the nitrido ligands in compound **2**.



Scheme 2. Protonation and hydrogenation of **2**

Experimental Section

General Considerations. All manipulations were carried out under argon atmosphere using Schlenk line or glovebox techniques. Toluene and hexane were distilled from Na/K alloy just before use. Dichloromethane was dried with P₂O₅ and distilled prior to use. Fluorobenzene was dried with CaH₂ and distilled prior to use. NMR solvents were dried with Na/K alloy (C₆D₆) or calcium hydride (CDCl₃, (CD₃)₂SO, C₅D₅N) and distilled before use. Oven-dried glassware was repeatedly evacuated with a pumping system (ca. 1 × 10⁻³ Torr) and subsequently filled with inert gas. ROSO₂CF₃ (R = H, Me, Me₃Si) were purchased from Aldrich and used as received. [{Ti(η⁵-C₅Me₅)(μ-NH)}₃(μ₃-N)] (**1**),²² [{Ti(η⁵-C₅Me₅)₄(μ₃-N)₄] (**2**),²³ [{Ti(η⁵-C₅Me₅)₄(μ₃-N)₃(μ₃-NH)][OTf] (**8**),²⁹ [Ti(η⁵-C₅Me₅)Me₃],³⁰ and [M(η⁵-C₅H₅)(CO)₃H] (M = Mo, Cr)³¹ were prepared according to published procedures.

Samples for infrared spectroscopy were prepared as KBr pellets, and the spectra were obtained using an FT-IR Perkin-Elmer SPECTRUM 2000 or FT-IR Perkin-Elmer FRONTIER spectrophotometers. ¹H, ¹³C{¹H} and ¹⁹F NMR spectra were recorded on a Varian Unity-300, Mercury-300 and/or Unity-500 spectrometers. Chemical shifts (δ, ppm) in the ¹H and ¹³C{¹H} NMR spectra are given relative to residual protons or to carbon of the solvent, C₆D₆ (¹H: δ = 7.15; ¹³C: δ = 128.0), CDCl₃ (¹H: δ = 7.24; ¹³C: δ = 77.0), (CD₃)₂SO (¹H: δ = 2.49; ¹³C: δ = 39.0) or C₅D₅N (¹H: δ = 8.71; ¹³C: δ = 149.9). Chemical shifts (δ, ppm) in the ¹⁹F NMR spectra are given relative to CFCl₃ as external reference. The effective magnetic moments were determined by the Evans NMR method at 293 K (using a 300 MHz instrument with a field strength of 7.05 Tesla).³² Microanalyses (C, H,

N, S) were performed in a Leco CHNSO-932 or Perkin Elmer CHNS/O 2400 microanalyzers.

Synthesis of $[\text{Ti}_2(\eta^5\text{-C}_5\text{Me}_5)_2(\mu\text{-N})(\text{NH}_3)(\mu\text{-O}_2\text{SOCF}_3)_2(\text{OTf})]$ (3**).** *Method A:* A 100 mL Schlenk tube was charged with **1** (0.20 g, 0.33 mmol), HOTf (0.25 g, 1.67 mmol), and toluene (25 mL). The reaction mixture was stirred at room temperature for 2 days to give a dark orange solution and a fine white solid. The solid (0.043 g) was isolated by filtration onto a glass frit and characterized as $[\text{NH}_4]\text{OTf}$ by ^1H NMR spectroscopy in DMSO-d_6 , $\delta = 7.06$ (t, $^1J(\text{H},^{14}\text{N}) = 51.2$ Hz, 4H; NH_4). The volatile components were removed from the filtrate under reduced pressure to afford **3** as an orange solid (0.34 g, 81%). *Method B:* A 100 mL ampule (Teflon stopcock) was charged with **1** (0.20 g, 0.33 mmol), MeOTf (0.43 g, 2.62 mmol), and toluene (20 mL). The reaction mixture was stirred at 100 °C for 2 days to give a dark orange solution and a light violet solid. The solid (0.31 g) was isolated by filtration onto a glass frit and subsequently characterized as a 1:1 mixture of $[\text{NMe}_4]\text{OTf}$ and $[\text{NMe}_4][\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)(\text{OTf})_4]$ (**5**), according to analysis by ^1H NMR spectroscopy of a solution of this solid in DMSO-d_6 . The volatile components were removed from the filtrate under reduced pressure to afford **3** as an orange solid (0.26 g, 93%). IR (KBr, cm^{-1}): $\tilde{\nu}$ 3344 (w), 3267 (w), 3183 (w), 2962 (w), 2917 (m), 1607 (w), 1489 (w), 1431 (m), 1382 (m), 1331 (vs), 1312 (vs), 1230 (vs), 1208 (vs), 1187 (vs), 1145 (s), 1029 (vs), 1008 (vs), 879 (vs), 857 (vs), 791 (m), 731 (w), 632 (vs), 513 (m), 444 (m). ^1H NMR (300 MHz, C_6D_6 , 20 °C): δ 2.07 (s, 15H; C_5Me_5), 1.76 (s, 15H; C_5Me_5), 1.68 (s br., 3H; NH_3). $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, C_6D_6 , 20 °C): δ 131.9 (C_5Me_5), 128.9 (C_5Me_5), 12.4 (C_5Me_5), 12.0 (C_5Me_5), the CF_3 carbon atom resonances were not detected. ^{19}F NMR (282 MHz, C_6D_6 , 20 °C): δ -76.8

(s), -77.6 (s). Anal. Calcd for $C_{23}H_{33}F_9N_2O_9S_3Ti_2$ ($M_w = 844.43$): C 32.71, H 3.94, N 3.32, S 11.39. Found: C 33.03, H 3.79, N 3.52, S 11.24.

Synthesis of $[Ti_3(\eta^5-C_5Me_5)_3(\mu-N)(\mu-NH)_2(\mu-O_2SO_2CF_3)(OTf)]$ (4**).** A 100 mL ampule (Teflon stopcock) was charged with **1** (0.30 g, 0.49 mmol), Me_3SiOTf (0.22 g, 0.99 mmol), and toluene (30 mL). After stirring at room temperature for 24 h, the reaction mixture was heated at $85\text{ }^\circ\text{C}$ for 2 d to give a dark orange solution. After filtration, the volatile components of the solution were removed under reduced pressure to afford **4** as an orange solid (0.31 g, 70%). IR (KBr, cm^{-1}): $\tilde{\nu}$ 3358 (m), 3261 (w), 3184 (w), 2915 (s), 2863 (m), 2731 (w), 1614 (w), 1495 (w), 1434 (m), 1382 (m), 1337 (vs), 1312 (vs), 1236 (vs), 1196 (vs), 1139 (s), 1068 (w), 1011 (vs), 927 (vs), 906 (vs), 788 (s), 720 (m), 633 (vs), 571 (w), 518 (m), 435 (m). 1H NMR (300 MHz, C_6D_6 , $20\text{ }^\circ\text{C}$): δ 14.19 (s br., 2H; NH), 2.04 (s, 30H; C_5Me_5), 1.87 (s, 15H; C_5Me_5). $^{13}C\{^1H\}$ NMR (125 MHz, C_6D_6 , $20\text{ }^\circ\text{C}$): δ 124.2 (C_5Me_5), 122.9 (C_5Me_5), 11.7 (C_5Me_5), 11.6 (C_5Me_5), the CF_3 carbon atom resonances were not detected. ^{19}F NMR (282 MHz, C_6D_6 , $20\text{ }^\circ\text{C}$): δ -76.4 (s). Anal. Calcd for $C_{32}H_{47}F_6N_3O_6S_2Ti_3$ ($M_w = 891.47$): C 43.11, H 5.31, N 4.71, S 7.19. Found: C 43.07, H 5.63, N 4.34, S 6.96.

Reaction of **1 with 4 equiv of $MeOTf$.** A 100 mL ampule (Teflon stopcock) was charged with **1** (0.20 g, 0.33 mmol), $MeOTf$ (0.22 g, 1.34 mmol), and toluene (20 mL). The reaction mixture was stirred at $60\text{ }^\circ\text{C}$ for 16 h to give a dark orange solution and a fine white solid. The solid (0.063 g) was isolated by filtration onto a glass frit and characterized as $[NMe_4]OTf$. The volatile components were removed from the filtrate under reduced pressure to afford an orange solid (0.28 g). Analysis by NMR spectroscopy of a solution of this solid in benzene- d_6 showed it to be a 1:1 mixture of compounds **3** and **4**.

Spectroscopic and analytical data for [NMe₄]OTf:³³ IR (KBr, cm⁻¹): $\tilde{\nu}$ 3047 (w), 2973 (m), 1493 (s), 1420 (w), 1252 (vs), 1229 (vs), 1154 (vs), 1033 (vs), 953 (s), 793 (w), 757 (w), 639 (vs), 573 (m), 519 (m), 459 (w). ¹H NMR (300 MHz, DMSO-d₆, 20 °C): δ 3.08 (s; NMe₄). ¹³C{¹H} NMR (75 MHz, DMSO-d₆, 20 °C): δ 120.2 (q, ¹J(C,F) = 320.3 Hz; CF₃), 53.9 (t, ¹J(C,¹⁴N) = 4.0 Hz; NMe₄). ¹⁹F NMR (282 MHz, DMSO-d₆, 20 °C): δ -81.9 (s). Anal. Calcd for C₅H₁₂F₃NO₃S (*M_w* = 223.21): C 26.91, H 5.42, N 6.28, S 14.37. Found: C 26.36, H 5.02, N 6.31, S 14.60.

Synthesis of [NMe₄][Ti(η^5 -C₅Me₅)(OTf)₄] (5). A 100 mL Schlenk tube was charged with **7** (0.10 g, 0.16 mmol), [NMe₄]OTf (0.035 g, 0.16 mmol), and toluene (25 mL). The reaction mixture was stirred at room temperature for 24 h to give a dark violet solid and a violet solution. The solid was isolated by filtration onto a glass frit and vacuum-dried to afford **5** as a dark violet powder (0.080 g, 57%). IR (KBr, cm⁻¹): $\tilde{\nu}$ 2923 (w), 1491 (m), 1364 (m), 1243 (s), 1201 (vs), 1197 (vs), 1193 (vs), 1148 (m), 1033 (m), 1009 (m), 974 (s), 766 (w), 631 (s), 596 (w), 508 (w), 451 (w), 412 (w). ¹H NMR (300 MHz, DMSO-d₆, 20 °C): δ 3.08 (s, 12H; NMe₄), 2.08 (s, 15H; C₅Me₅). ¹³C{¹H} NMR (75 MHz, DMSO-d₆, 20 °C): δ 140.0 (C₅Me₅), 120.2 (q, ¹J(C,F) = 320.3 Hz; CF₃), 53.9 (t, ¹J(C,¹⁴N) = 3.8 Hz; NMe₄), 13.3 (C₅Me₅). ¹⁹F NMR (282 MHz, DMSO-d₆, 20 °C): δ -77.7 (s). Anal. Calcd for C₁₈H₂₇F₁₂NO₁₂S₄Ti (*M_w* = 853.51): C 25.33, H 3.19, N 1.64, S 15.03. Found: C 24.89, H 3.01, N 2.05, S 15.46.

Synthesis of [NH₄][Ti(η^5 -C₅Me₅)(OTf)₄] (6). A 100 mL Schlenk tube was charged with **3** (0.20 g, 0.24 mmol), HOTf (0.19 g, 1.27 mmol), and toluene (30 mL). The reaction mixture was stirred at room temperature for 24 h to give a dark violet solid and a violet solution. The solid was isolated by filtration onto a glass frit and vacuum-dried to afford **6** as a dark

violet powder (0.34 g, 89%). IR (KBr, cm^{-1}): $\tilde{\nu}$ 3259 (m, broad), 1489 (w), 1435 (m), 1351 (s), 1248 (vs), 1211 (vs), 1192 (vs), 1157 (s), 1033 (s), 988 (vs), 769 (w), 634 (s), 623 (s), 598 (w), 506 (w), 449 (w). ^1H NMR (300 MHz, DMSO-d_6 , 20 °C): δ 7.06 (t, $^1J(\text{H},^{14}\text{N}) = 51.2$ Hz, 4H; NH_4), 2.08 (s, 15H; C_5Me_5). $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, DMSO-d_6 , 20 °C): δ 140.0 (C_5Me_5), 120.2 (q, $^1J(\text{C},\text{F}) = 320.0$ Hz; CF_3), 13.3 (C_5Me_5). ^{19}F NMR (282 MHz, DMSO-d_6 , 20 °C): δ -77.8 (s). Anal. Calcd for $\text{C}_{14}\text{H}_{19}\text{F}_{12}\text{NO}_{12}\text{S}_4\text{Ti}$ ($M_w = 797.40$): C 21.09, H 2.40, N 1.76, S 16.08. Found: C 20.84, H 2.50, N 1.83, S 16.15.

Synthesis of $[\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)(\text{OTf})_3]$ (7). A 100 mL Schlenk tube was charged with $[\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\text{Me}_3]$ (0.20 g, 0.88 mmol), HOTf (0.40 g, 2.67 mmol), and toluene (40 mL). The reaction mixture was stirred at room temperature for 24 h to give a violet solid and a violet solution. The solid was isolated by filtration onto a glass frit and vacuum-dried to afford **7** as a violet powder (0.37 g, 67%). IR (KBr, cm^{-1}): $\tilde{\nu}$ 2923 (w), 1489 (w), 1432 (w), 1374 (s), 1358 (s), 1340 (s), 1238 (vs), 1198 (vs), 1141 (vs), 1013 (s), 981 (vs), 947 (s), 769 (w), 636 (s), 622 (s), 598 (m), 510 (w), 453 (w), 419 (w). ^1H NMR (300 MHz, CDCl_3 , 20 °C): δ 2.63 (s, C_5Me_5). ^1H NMR (300 MHz, DMSO-d_6 , 20 °C): δ 2.08 (s; C_5Me_5). $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3 , 20 °C): δ 149.9 (C_5Me_5), 118.3 (q, $^1J(\text{C},\text{F}) = 320.1$ Hz; CF_3), 14.7 (C_5Me_5). $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, DMSO-d_6 , 20 °C): δ 140.0 (C_5Me_5), 120.2 (q, $^1J(\text{C},\text{F}) = 320.1$ Hz; CF_3), 13.3 (C_5Me_5). ^{19}F NMR (282 MHz, CDCl_3 , 20 °C): δ -75.9 (s). ^{19}F NMR (282 MHz, DMSO-d_6 , 20 °C): δ -77.7 (s). Anal. Calcd for $\text{C}_{13}\text{H}_{15}\text{F}_9\text{O}_9\text{S}_3\text{Ti}$ ($M_w = 630.30$): C 24.77, H 2.40, S 15.26. Found: C 24.72, H 2.50, S 15.16.

Synthesis of $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-N})_3(\mu_3\text{-NMe})][\text{OTf}]$ (9). A 100 mL ampule (Teflon stopcock) was charged with **2** (0.20 g, 0.25 mmol), MeOTf (0.042 g, 0.26 mmol), and toluene (20 mL). The system was allowed to react at 100 °C without any stirring for 3 days.

After decantation, the resultant dark red needles were vacuum-dried to afford **9** (0.18 g, 72%). IR (KBr, cm^{-1}): $\tilde{\nu}$ 2909 (s), 1602 (w), 1489 (m), 1437 (s), 1378 (s), 1275 (vs), 1221 (s), 1144 (vs), 1062 (w), 1032 (vs), 953 (w), 869 (w), 802 (m), 750 (w), 732 (w), 648 (vs), 619 (vs), 571 (s), 548 (m), 517 (m), 463 (vs), 434 (m). ^1H NMR (300 MHz, CDCl_3 , 20 $^\circ\text{C}$): δ 4.39 (s, 3H; NMe), 2.08 (s, 45H; C_5Me_5), 2.07 (s, 15H; C_5Me_5). $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3 , 20 $^\circ\text{C}$): δ 126.6 (C_5Me_5), 125.3 (C_5Me_5), 55.3 (NMe), 12.7 (C_5Me_5), 12.5 (C_5Me_5), the CF_3 carbon atom resonance was not detected. ^{19}F NMR (282 MHz, CDCl_3 , 20 $^\circ\text{C}$): δ -78.0 (s). Anal. Calcd for $\text{C}_{42}\text{H}_{63}\text{F}_3\text{N}_4\text{O}_3\text{STi}_4$ ($M_w = 952.51$): C 52.96, H 6.67, N 5.88, S 3.37. Found: C 53.04, H 6.38, N 6.06, S 3.19.

Synthesis of $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-N})_3(\mu_3\text{-NH})][\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)(\text{OTf})_4(\text{NH}_3)]$ (10**).** *Method*

A: A 100 mL Schlenk tube was charged with **2** (0.20 g, 0.25 mmol), HOTf (0.16 g, 1.07 mmol), and toluene (20 mL). The reaction mixture was stirred at room temperature for 24 h to give a dark red solid and a red solution. The solid was isolated by filtration onto a glass frit and vacuum-dried to afford **10** as a dark red powder (0.29 g, 73%). *Method B:* A 100 mL Schlenk tube was charged with **2** (0.10 g, 0.13 mmol), **6** (0.10 g, 0.13 mmol), and toluene (25 mL). The reaction mixture was stirred at room temperature for 24 h to give a dark red solid and a red solution. The solid was isolated by filtration onto a glass frit and vacuum-dried to afford **10** as a dark red powder (0.16 g, 80%). IR (KBr, cm^{-1}): $\tilde{\nu}$ 3360 (w), 3215 (w, broad), 2916 (m), 1653 (w), 1620 (w), 1494 (w), 1442 (m), 1381 (m), 1348 (s), 1236 (s), 1200 (vs), 1032 (s), 1001 (vs), 790 (m), 736 (m), 636 (vs), 514 (w), 468 (w), 446 (w). ^1H NMR (300 MHz, CDCl_3 , 20 $^\circ\text{C}$): δ 12.42 (s br., 1H; NH), 2.40 (s br., 15H; C_5Me_5), 2.06 (s, 15H; C_5Me_5), 2.05 (s, 45H; C_5Me_5), the resonance signal for the NH_3 ligand was not detected. $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3 , 20 $^\circ\text{C}$): δ 147.2 (C_5Me_5), 126.2 (C_5Me_5),

125.1 (C_5Me_5), 13.7 (C_5Me_5), 12.7 (C_5Me_5), 12.2 (C_5Me_5), the CF_3 carbon atom resonance was not detected. ^{19}F NMR (282 MHz, $CDCl_3$, 20 °C): δ -76.1 (s). Anal. Calcd for $C_{54}H_{79}F_{12}N_5O_{12}S_4Ti_5$ ($M_w = 1585.81$): C 40.90, H 5.02, N 4.42, S 8.09. Found: C 40.22, H 4.72, N 4.51, S 7.90.

Synthesis of $[Ti(\eta^5-C_5Me_5)_4(\mu_3-N)_3(\mu_3-NH)]$ (11**).**²⁹ A 100 mL Schlenk tube was charged with **2** (0.30 g, 0.38 mmol), $[Mo(\eta^5-C_5H_5)(CO)_3H]$ (0.094 g, 0.38 mmol), and toluene (25 mL). The reaction mixture was stirred at room temperature for 24 h to give a dark blue solution and a fine dark blue solid. The solid was isolated by filtration onto a glass frit and vacuum-dried to afford **11** as a dark blue powder (0.26 g, 88%). IR (KBr, cm^{-1}): $\tilde{\nu}$ 3333 (w) (ν_{NH}). 1H NMR (300 MHz, $CDCl_3$, 20 °C): δ 10.0 (s br., $\Delta\nu_{1/2} = 45$ Hz, 15H; C_5Me_5), 8.7 (s br., $\Delta\nu_{1/2} = 33$ Hz, 45H; C_5Me_5). The effective magnetic moment of **11** was determined to be 1.70 μ_B (based on a unit formula of $C_{40}H_{61}N_4Ti_4$) on a $CDCl_3$ solution.

Synthesis of $[Ti(\eta^5-C_5Me_5)_4(\mu_3-N)_2(\mu_3-NH)_2][Mo(\eta^5-C_5H_5)(CO)_3]$ (12**).** A 100 mL Schlenk tube was charged with **2** (0.25 g, 0.32 mmol), $[Mo(\eta^5-C_5H_5)(CO)_3H]$ (0.078 g, 0.32 mmol), and toluene (20 mL). The reaction mixture was stirred at room temperature for 2 h and a second 1 equiv of $[Mo(\eta^5-C_5H_5)(CO)_3H]$ (0.078 g, 0.32 mmol) was added to the resultant dark suspension. After stirring at ambient temperature for 16 h, a dark green precipitate was isolated by filtration onto a glass frit and vacuum-dried to afford **12** (0.27 g, 82%). IR (KBr, cm^{-1}): $\tilde{\nu}$ 3277 (m), 3264 (m), 2910 (s), 2858 (m), 1895 (vs), 1781 (vs), 1747 (vs), 1494 (m), 1427 (s), 1377 (s), 1023 (w), 788 (vs), 782 (m), 697 (w), 642 (w), 617 (w), 515 (m), 496 (m), 445 (m), 421 (m). 1H NMR (300 MHz, C_5D_5N , 20 °C): δ 9.3 (s br., $\Delta\nu_{1/2} = 61$ Hz, 30H; C_5Me_5), 8.0 (s br., $\Delta\nu_{1/2} = 55$ Hz, 30H; C_5Me_5), 5.51 (s, 5H; C_5H_5). Anal. Calcd for $C_{48}H_{67}MoN_4O_3Ti_4$ ($M_w = 1035.48$): C 55.68, H 6.52, N 5.41. Found: C 55.45, H

6.12, N 5.37. The effective magnetic moment of **12** was determined to be $1.64 \mu_B$ (based on a unit formula of $C_{48}H_{67}MoN_4O_3Ti_4$) on a C_5D_5N solution.

Synthesis of $[Ti(\eta^5-C_5Me_5)]_4(\mu_3-N)_2(\mu_3-NH)_2[Cr(\eta^5-C_5H_5)(CO)_3]$ (13**).** In a fashion similar to the preparation of **12**, the treatment of **2** (0.20 g, 0.25 mmol) with $[Cr(\eta^5-C_5H_5)(CO)_3H]$ (0.10 g, 0.50 mmol) in toluene (15 mL) for 2 days afforded **13** as a dark green powder (0.19 g, 38%). IR (KBr, cm^{-1}): $\tilde{\nu}$ 3289 (m), 3266 (m), 3101 (w), 2911 (s), 2858 (m), 1936 (m), 1890 (vs), 1778 (vs), 1746 (vs), 1487 (m), 1427 (s), 1377 (s), 1261 (w), 1112 (w), 1066 (w), 1023 (m), 786 (s), 747 (s), 695 (m), 652 (s), 509 (s), 445 (s). 1H NMR (300 MHz, C_5D_5N , 20 °C): δ 9.3 (s br., $\Delta\nu_{1/2} = 70$ Hz, 30H; C_5Me_5), 8.0 (s br., $\Delta\nu_{1/2} = 57$ Hz, 30H; C_5Me_5), 5.02 (s, 5H, C_5H_5). Anal. Calcd for $C_{48}H_{67}CrN_4O_3Ti_4$ ($M_w = 991.53$): C 58.14, H 6.81, N 5.65. Found: C 57.46, H 6.61, N 5.92. The effective magnetic moment of **13** was determined to be $1.62 \mu_B$ (based on a unit formula of $C_{48}H_{67}CrN_4O_3Ti_4$) on a C_5D_5N solution.

X-ray structure determination of **3, **4**, **5**, **9**, **10**, and **13**.** Orange crystals of **3** were grown at room temperature by slow diffusion of hexane into a fluorobenzene solution of this compound. Orange crystals of $4 \cdot C_7H_8$ and dark red crystals of $9 \cdot C_7H_8$ were obtained from toluene solutions at -30 °C. Dark violet crystals of **5** were grown by slow cooling at room temperature of a toluene solution heated at 110 °C. Dark red crystals of **10** were obtained at room temperature by diffusion of hexane into a dichloromethane solution of this compound. Dark green crystals of $13 \cdot C_6D_6$ were grown by slow cooling at room temperature of a benzene- d_6 solution heated at 80 °C in a NMR tube. The crystals were removed from the Schlenk or NMR tubes and covered with a layer of a viscous perfluoropolyether (Fomblin®Y). A suitable crystal was selected with the aid of a

microscope, mounted on a cryoloop, and immediately placed in the low temperature nitrogen stream of the diffractometer. The intensity data sets were collected at 200 or 150 K on a Bruker-Nonius KappaCCD diffractometer equipped with an Oxford Cryostream 700 unit. Crystallographic data for all the complexes are presented in Table S1 of the Supporting Information.

The structures were solved, using the WINGX package,³⁴ by direct methods (**13**) (SHELXS-2013)³⁵ or intrinsic phasing methods (the rest) (SHELXT)³⁶, and refined by least-squares against F^2 (SHELXL-2014/7).³⁷ It was not possible to obtain crystals of better quality for compounds **3–5** and **9**, and therefore some disorder problems could not be solved in those structures.

The asymmetric unit of **3** was formed by two independent molecules associated through hydrogen bonds, but there were no significant differences between these molecules. Crystals of **3** showed disorder for one bridging triflate ligand (S(2), O(21), O(22), O(23), C(2), F(21), F(22) and F(23)). This disorder was treated by using the PART tool of the SHELXL program and allowing free refinement of the occupancy factor with the FVAR command. The final values of occupancy were 89.9 and 10.1%. All non-hydrogen atoms were anisotropically refined, except the atoms corresponding to the minor position of the disordered triflate group (S(2)', O(21)', O(22)', O(23)', C(2)', F(21)', F(22)' and F(23)'), which were refined isotropically. All the hydrogen atoms were placed geometrically and refined by using a riding model, except those of the ammine ligands (H(21), H(22), H(23), H(41), H(42) and H(43)), which were found in the difference Fourier map and refined isotropically.

In the difference Fourier map for compound **4**, a distorted molecule of toluene was found, so it was constrained to be constituted by a regular hexagon. Crystals of **4** presented

disorder for the non-bridging triflate ligand (S(3), O(31), O(32), O(33), C(3), F(31), F(32) and F(33)). This disorder was also treated by using the PART tool with final values of 75.6 and 24.4%. Additionally, SADI restraints were employed for the CF₃ moiety of the minor position of this disordered triflate group. All non-hydrogen atoms were anisotropically refined, whereas all the hydrogen atoms were positioned geometrically and refined using a riding model. Moreover, the atoms of the pentamethylcyclopentadienyl ring linked to Ti(2) were restrained with DELU and SIMU instructions.

In the crystallographic study of **5**, all non-hydrogen atoms were anisotropically refined, whereas all the hydrogen atoms were positioned geometrically and refined by using a riding model.

Complex **9** crystallized with a toluene molecule. Crystals of **9** presented disorder for the atoms of the triflate group (S(1), O(1), O(2), O(3), C(2), F(1), F(2) and F(3)) and the carbon atoms of the pentamethylcyclopentadienyl ligand linked to Ti(3) (C(31)-C(40)). These disorders were also treated conventionally affording occupancy factors of 60.2 and 39.8% for the triflate anion, and 79 and 21% for the C₅Me₅ moiety. All non-hydrogen atoms were anisotropically refined, while all the hydrogen atoms were positioned geometrically and refined using a riding model. The minor position for the disordered C₅Me₅ ring was constrained to retain a regular geometry, and DELU and SIMU restraints were used with the carbon atoms of this fragment. Atoms N(1) and C(1) were also restrained with DELU and SIMU commands, and SADI restraints were employed with the triflate group. Additionally, FREE instruction was applied to the atoms C(1) and O(1)' to avoid problems on calculated hydrogen atoms.


In the study of compound **10**, all non-hydrogen atoms were anisotropically refined. All hydrogen atoms were placed geometrically and refined using a riding model, except

those linked to nitrogen (H(11), H(51), H(52) and H(53)), which were found in the difference Fourier map and refined isotropically. Moreover, distance restraints were applied to constrain the distance N(1)–H(11).

Compound **13** crystallized with a benzene molecule, which was located in the difference Fourier map, although it was constrained to be a regular hexagon. All non-hydrogen atoms were anisotropically refined. All hydrogen atoms were positioned geometrically and refined by using a riding model, except hydrogen H(1) and H(3) which were also found in the Fourier map, and refined isotropically.

Results and Discussion

Reaction of complex 1 with excess ROTf electrophilic reagents. The treatment of $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)(\mu\text{-NH})\}_3(\mu_3\text{-N})]$ (**1**) with 5.5 equiv of triflic acid in toluene at room temperature led to the precipitation of ammonium triflate and a dark orange solution from which the dinuclear complex $[\text{Ti}_2(\eta^5\text{-C}_5\text{Me}_5)_2(\mu\text{-N})(\text{NH}_3)(\mu\text{-O}_2\text{SOCF}_3)_2(\text{OTf})]$ (**3**) was isolated in 81% yield (Scheme 3). Reactions of **1** with lower ratios (1:2, 1:3 or 1:4) of HOTf gave complicated mixtures of products, where only compounds $[\text{Ti}_3(\eta^5\text{-C}_5\text{Me}_5)_3(\mu_3\text{-N})(\mu\text{-NH})_2(\mu\text{-NH}_2)(\text{OTf})]$ (Scheme 1)²⁸ and **3** were identified by NMR spectroscopy.



Scheme 3. Synthesis of the dinuclear complex **3**.

Complex **3** was characterized by analytical and spectroscopic methods, as well as by an X-ray crystal structure determination. The IR spectrum (KBr) of **3** shows three bands for the ν_{NH} vibrations, between 3344 and 3183 cm^{-1} , and one absorption at 1607 cm^{-1} assignable to the NH_3 bending mode.³⁸ In addition, the spectrum reveals several strong absorptions in the range 1331-1008 cm^{-1} for the coordinated triflate ligands.³⁹ The ^1H NMR spectrum of **3** in benzene- d_6 at room temperature shows two singlets in a 1:1 ratio for the $\eta^5\text{-C}_5\text{Me}_5$ groups, and a broad resonance for the ammine ligand at $\delta = 1.68$. The ^{19}F NMR spectrum of **3** in benzene- d_6 reveals two resonance signals, assigned to bridging and

terminal triflate ligands, in accord with its solid-state structure. The solid-state structural study of a single crystal of **3** shows two molecules associated through hydrogen bonding interactions between the terminal triflate group of one molecule and the ammine ligand of the other molecule in the asymmetric unit (*vide infra*). There are no substantial differences between the two molecules (see Figure 1 and Table 1 for one molecule, and Figure S1 and Table S2 in the Supporting Information for the second one). Compound **3** shows a dinuclear structure with two asymmetric moieties “Ti(η^5 -C₅Me₅)L” held together by one bridging nitrido ligand and two μ -O₂SOCF₃ groups. The titanium atoms have classical four-legged piano-stool arrangements, where the legs are occupied by one nitrido, one terminal triflate, and two bridging triflate ligands for Ti(1) and Ti(3); and one nitrido, one ammine, and two bridging triflate ligands for Ti(2) and Ti(4). The bridging nitrido ligand of **3** is characterized by titanium–nitrogen bond lengths of averaged 1.797(2) Å, which is consistent with a Ti=N=Ti unit, based in the few examples found in the literature containing that fragment.^{19a,40} The narrow Ti–N(nitrido)–Ti angle (av. 133.4(1)°) is likely due to the existence of two additional triflate ligands bridging the metal centers. The titanium–oxygen distances of 2.100(3)–2.185(3) Å are in the normal range for triflate ligands in complexes such as [Ti(η^5 -C₅Me₅)(μ -OH)(μ -O₂SOCF₃)(OTf)]₂,⁴¹ and the titanium–nitrogen(ammine) bond length (av. 2.194(9) Å) compares well with those found in the few examples of titanium organometallic complexes with NH₃ ligands.^{16a,19 a,42}




Figure 1. Perspective view of one of the two associated molecules in the asymmetric unit of complex **3** (thermal ellipsoids at the 50% probability level). The methyl groups of the pentamethylcyclopentadienyl ligands are omitted for clarity.

Table 1. Selected Lengths (Å) and Angles (deg) for One of the Two Associated Molecules of **3**

Ti(1)–N(12)	1.796(3)	Ti(1)–O(11)	2.100(3)
Ti(1)–O(21)	2.173(5)	Ti(1)–O(31)	2.149(3)
Ti(2)–N(12)	1.798(3)	Ti(2)–N(2)	2.188(4)
Ti(2)–O(22)	2.159(6)	Ti(2)–O(32)	2.127(3)
S(2)–O(21)	1.439(6)	S(2)–O(22)	1.458(6)
S(3)–O(31)	1.465(3)	S(3)–O(32)	1.460(3)
Ti(1)···Ti(2)	3.301(1)		
N(12)–Ti(1)–O(11)	134.8(1)	N(12)–Ti(1)–O(21)	85.6(2)
N(12)–Ti(1)–O(31)	86.0(1)	O(11)–Ti(1)–O(21)	77.9(2)
O(11)–Ti(1)–O(31)	78.2(1)	O(21)–Ti(1)–O(31)	136.3(2)
N(12)–Ti(2)–N(2)	125.2(2)	N(12)–Ti(2)–O(22)	86.1(2)
N(12)–Ti(2)–O(32)	86.3(1)	N(2)–Ti(2)–O(22)	77.0(2)
N(2)–Ti(2)–O(32)	75.8(2)	O(22)–Ti(2)–O(32)	140.1(2)
Ti(1)–N(12)–Ti(2)	133.4(2)	Ti(1)–O(21)–S(2)	125.2(4)
Ti(1)–O(31)–S(3)	128.4(2)	Ti(2)–O(22)–S(2)	125.0(3)
Ti(2)–O(32)–S(3)	129.0(2)	O(21)–S(2)–O(22)	113.6(3)
O(31)–S(3)–O(32)	113.0(2)		

As mentioned above, analysis of the crystal structure of **3** reveals the existence of several N–H···O and N–H···F hydrogen bonds (Figure 2, Table 2). Within each molecule, there is an intramolecular hydrogen bonding interaction between an oxygen atom of the terminal triflate and one hydrogen atom of the ammine ligand, which can be classified as strong according to the criteria on the donor–acceptor distances.⁴³ In addition, the two molecules of the asymmetric unit are associated by weaker hydrogen bonding interactions between oxygen or fluorine atoms of the terminal triflate ligand of one molecule and the hydrogen atoms on the ammine group of the other molecule.





Figure 2. Simplified view of the most relevant intra- and intermolecular hydrogen bonding interactions present in complex **3** (N–H···O hydrogen bonds are drawn in orange, whereas N–H···F interactions are in green).

Table 2. Relevant Hydrogen Bonds^a for Compound **3**

D–H···A	D···A/Å	H···A/Å	D–H···A/deg
N(2)–H(21)···O(42)	3.275(6)	2.77(7)	124(6)
N(2)–H(22)···F(41)	3.548(8)	2.75(6)	154(6)
N(2)–H(23)···O(12)	2.961(6)	2.09(6)	165(6)
N(4)–H(41)···O(43)	2.972(7)	2.18(6)	164(5)
N(4)–H(42)···O(13)	3.164(6)	2.60(7)	126(6)
N(4)–H(43)···F(21)	3.259(7)	2.34(5)	159(5)

^aA = acceptor; D = donor.

Compound **3** was also isolated in 93% yield from the orange solution resulting of the treatment of **1** with 8 equiv of methyl triflate in toluene at 100 °C for 2 days (Scheme 4). Along with this solution, a light violet precipitate was obtained in the reaction. This solid was only soluble in dimethylsulfoxide-d₆, and its composition was analyzed by NMR spectroscopy. The spectra revealed a 1:1 mixture of the tetramethylammonium salts [NMe₄]OTf and [NMe₄][Ti(η⁵-C₅Me₅)(OTf)₄] (**5**), which were independently synthesized by other procedures. Thus, the reaction of **1** with 4 equiv of MeOTf in toluene at 60 °C for 16 h afforded an orange solution and a white precipitate. Spectroscopic and analytical data of this solid were consistent with the ionic compound [NMe₄]OTf,³³ whereas the orange solution obtained in the reaction was shown to contain a 1:1 mixture of **3** and the trinuclear derivative [Ti₃(η⁵-C₅Me₅)₃(μ-N)(μ-NH)₂(μ-O₂SO₂CF₃)(OTf)] (**4**) according to ¹H NMR spectroscopy. Compounds **3** and **4** exhibit a similar solubility in toluene, but crystallization at –30 °C gave a few orange crystals of **4**·C₇H₈ suitable for a single crystal X-ray diffraction determination (vide infra). Interestingly, addition of 4 equiv of MeOTf to the toluene solution containing derivatives **3** and **4**, and subsequent heating at 110 °C for 16 h, cleanly afforded an orange solution of complex **3** along with violet crystals characterized as [NMe₄][Ti(η⁵-C₅Me₅)(OTf)₄] (**5**) by X-ray crystallography.



Scheme 4. Reactions of **1** with excess ROTf (R = Me, SiMe₃).

The synthesis of **4** in a pure form and good yield (70%) was achieved by treatment of **1** with 2 equiv of trimethylsilyl triflate in toluene at 85 °C (Scheme 4). The first step in the formation of **4** involves the reaction of one equivalent of Me₃SiOTf with **1** at room temperature to give the previously reported complex [Ti₃(η⁵-C₅Me₅)₃(μ₃-N)(μ-NH)₂(μ-NHSiMe₃)] [OTf] (Scheme 1).²⁸ Subsequent silylation of the NHSiMe₃ amido ligand of this ionic compound with an additional equivalent of Me₃SiOTf at higher temperatures would result in formation of the amine NH(SiMe₃)₂, which is eventually released with concomitant coordination of the triflate groups to give compound **4**.

The ¹H NMR spectrum of **4** in benzene-d₆ at room temperature shows two sharp singlets in a 2:1 ratio for the η⁵-C₅Me₅ ligands and a broad resonance at δ = 14.19 for the imido groups. While the ¹H and ¹³C{¹H} NMR spectra are consistent with a C_s symmetry in solution, the ¹⁹F NMR spectrum showed only one resonance for the triflate groups, even though the solid-state structure reveals inequivalent terminal and bridging triflate ligands. This could be due to coincidence of the resonance signals or the existence of a rapid

dynamic exchange involving terminal and bridging sites of the triflate groups in solution. The X-ray crystal structure consists of a [Ti₃N₃] six-membered ring in boat conformation (Figure 3). The titanium atoms have classical three-legged piano-stool arrangements, where the legs are occupied by two imido and a terminal triflate ligands in the case of Ti(3), and one of the oxygen atoms of the bridging triflate group, one imido and one nitrido ligands for Ti(1) and Ti(2). The trifluoromethanesulfonato ligand that links these two titanium atoms closes a second six-membered [Ti₂NO₂S] ring in chair conformation. The Ti–N(imido) distances (Table 3) in **4** range from 1.880(5) to 1.894(5) Å, and resemble the values found in complexes **1** (av. 1.930(7) Å)²² and [Ti₃(η⁵-C₅Me₅)₃(μ₃-N)(μ-NH)(μ-NH₂)(μ-NR)(OTf)] (1.831(3)–1.858(3) Å)²⁸. However, the Ti–N(nitrido) distances of 1.791(5) and 1.797(4) Å are clearly shorter and agree with the higher titanium-nitrogen bond order expected for a Ti=N=Ti unit.⁴⁰ Those Ti–N lengths compare well with the averaged value of 1.797(2) Å found in the dinuclear compound **3**. In a fashion similar to this derivative, the narrow Ti–N(nitrido)–Ti angle (125.1(3)°) in complex **4** can be explained by the existence of two additional bridging systems, a triflate ligand and a [(μ-NH)₂Ti(η⁵-C₅Me₅)(OTf)] fragment, between the Ti(1) and Ti(2) atoms. Lastly, the crystal structure of **4** shows two different types of bonded triflate ligands. The terminal triflate ligand is coordinated to Ti(3) with a titanium–oxygen bond distance of 2.038(10) Å, which compares well with those found in other compounds such as [Ti(η⁵-C₅Me₅)(OMe)(μ-O₂SOCF₃)(OTf)]₂ (2.039(4) Å) and [Ti(η⁵-C₅Me₅)(μ-OH)(μ-O₂SOCF₃)(OTf)]₂ (2.071(3) Å).^{41a} The Ti–O distances associated with the bridging triflate ligand (averaged 2.088(9) Å) are logically longer than that of the terminal ligand, but they are shorter than those (2.125(2)–2.164(3) Å) found in the bridging ligands of the aforementioned examples.^{41a}





Figure 3. Perspective view of complex $4 \cdot C_7H_8$ (thermal ellipsoids at the 50% probability level). The methyl groups of the $\eta^5-C_5Me_5$ ligands and the toluene solvent molecule are omitted for clarity.

Table 3. Selected Lengths (Å) and Angles (deg) for Complex **4**

Ti(1)–N(12)	1.797(4)	Ti(1)–N(13)	1.880(5)
Ti(1)–O(11)	2.082(3)	Ti(2)–N(12)	1.791(5)
Ti(2)–N(23)	1.888(4)	Ti(2)–O(12)	2.095(3)
Ti(3)–N(13)	1.894(5)	Ti(3)–N(23)	1.892(4)
Ti(3)–O(31)	2.038(10)	Ti(1)···Ti(2)	3.184(1)
Ti(1)···Ti(3)	3.276(1)	Ti(2)···Ti(3)	3.198(1)
N(12)–Ti(1)–N(13)	101.6(3)	N(12)–Ti(1)–O(11)	100.4(2)
N(13)–Ti(1)–O(11)	101.8(2)	N(12)–Ti(2)–N(23)	104.7(3)
N(12)–Ti(2)–O(12)	99.6(2)	N(23)–Ti(2)–O(12)	103.5(2)
N(13)–Ti(3)–N(23)	116.5(2)	N(13)–Ti(3)–O(31)	100.3(3)
N(23)–Ti(3)–O(31)	107.9(2)	Ti(1)–N(12)–Ti(2)	125.1(3)
Ti(1)–N(13)–Ti(3)	120.5(3)	Ti(2)–N(23)–Ti(3)	115.6(2)

compound **5** in high purity. While ionic compounds **5** and **6** are only soluble in dimethylsulfoxide- d_6 , the molecular complex **7** is also soluble in chloroform- d_1 . Complexes **5-7** were characterized by spectroscopic and analytical methods, as well as by an X-ray crystal structure determination for **5**. IR spectra (KBr) show several strong absorptions between 1364 and 1009 cm^{-1} for the coordinated trifluoromethanesulfonato groups.³⁹ In addition, the IR spectrum of **6** displays a very broad band at 3259 cm^{-1} for the ν_{NH} vibration of the $[\text{NH}_4]^+$ cation. The ^1H NMR spectrum of a violet solution of **7** in chloroform- d_1 exhibits a singlet at $\delta = 2.63$ for the $\eta^5\text{-C}_5\text{Me}_5$ group. The resonance for the *ipso* carbons of the C_5Me_5 group ($\delta = 149.9$) in the $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum of **7** in chloroform- d_1 is shifted downfield with respect those found in analogous 12-electron complexes $[\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\text{X}_3]$ ($\text{X} = \text{Cl}$, $\delta = 137.2$; Br , $\delta = 138.2$; OEt , $\delta = 120.5$; OPh , $\delta = 126.5$).⁴⁴ This deshielding effect is consistent with the strong electron withdrawing character of the triflate ligand. Noteworthy, the violet compound **7** readily dissolves in dimethylsulfoxide- d_6 to give a brown solution, and the $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum shows a resonance signal at $\delta = 140.0$ for the *ipso* carbons of the C_5Me_5 group. The color change and significant high-field shift of this resonance signal when compared with those in chloroform- d_1 suggest the coordination of dimethylsulfoxide molecules to the titanium atom of compound **7** (Scheme 6). Furthermore, the ^1H and $^{13}\text{C}\{^1\text{H}\}$ NMR spectra of brown solutions of compounds **5** and **6** in dimethylsulfoxide- d_6 show resonance signals for the $\eta^5\text{-C}_5\text{Me}_5$ ligands ($\delta_{\text{H}} = 2.08$, and $\delta_{\text{C}} = 140.0$ and 13.3) which are identical to those of complex **7** in that solvent. In addition, the NMR spectra of **5** and **6** shows the same resonance signals assignable to the $[\text{NR}_4]^+$ groups that those found for the $[\text{NR}_4]\text{OTf}$ salts. These data suggest that compounds **5-7** reacts with dimethylsulfoxide- d_6 to give the same species in solution via displacement of

the corresponding salts $[\text{NR}_4]\text{OTf}$ in compounds **5** and **6**. This species could be tentatively formulated as the 16-electron complex $[\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)(\text{OTf})_3\{\text{OS}(\text{CD}_3)_2\}_2]$ (Scheme 6), but ionic derivatives with triflate anions (e.g., $[\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)(\text{OTf})_2\{\text{OS}(\text{CD}_3)_2\}_3][\text{OTf}]$) cannot be ruled out.



Scheme 6. Synthesis of **5** and plausible behavior of **5-7** in DMSO-*d*₆.

The solid-state structure of **5** shows well-separated tetrahedral $[\text{NMe}_4]^+$ cations and mononuclear titanium anions $[\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)(\text{OTf})_4]^-$. The titanium atom exhibits a four-legged piano-stool arrangement, with the legs occupied by an oxygen atom of the terminal trifluoromethanesulfonato groups (Figure 4 and Table S3). The Ti–O bond lengths of averaged 2.014(7) Å are similar to those found in complexes **3**, **4** and other examples with terminal triflate ligands.⁴¹ Furthermore, the distance from the titanium atom to the centroid

of the pentamethylcyclopentadienyl group of 2.045 Å is comparable to those reported for the analogous derivatives $[\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\text{Cl}_3]^{45}$ and $[\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)(\text{OC}_6\text{F}_5)_3]^{46}$.





Figure 4. Perspective view for the anion of compound **5** (thermal ellipsoids at the 50% probability level). Hydrogen atoms of the methyl groups are omitted for clarity. Selected averaged bond lengths (Å) and angles (deg): Ti(1)–O 2.014(7), Ti(1)–Cm 2.045, O–Ti(1)–Cm 109(1), O–Ti(1)–O_{cis} 83.7(7), O–Ti(1)–O_{trans} 141(2). Cm = Centroid of the pentamethylcyclopentadienyl ring.

Reactions of complex **2** with ROTf and $[\text{M}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\text{H}]$ reagents.

We have also studied the reactivity of the tetranuclear nitrido complex $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-N})_4]$ (**2**) with electrophilic ROTf (R = H, Me, Me₃Si) reagents (Scheme 7). Treatment of **2** with 1 equiv of triflic acid or methyl triflate in toluene afforded the precipitation of the ionic complexes $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-N})_3(\mu_3\text{-NR})][\text{OTf}]$ (R = H (**8**),²⁹ Me (**9**)). Interestingly, no reaction was observed in the treatment of **2** with the larger

trimethylsilyl fragment of Me₃SiOTf even at high temperatures. While compound **9** did not react with higher ratios of MeOTf even upon heating the reaction mixture at 100 °C, the treatment of **2** with excess HOTf in toluene at room temperature led to the formation of $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-N})_3(\mu_3\text{-NH})][\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)(\text{OTf})_4(\text{NH}_3)]$ (**10**). We speculate that triflic acid could produce the complete rupture of a fraction of a tetranuclear aggregate to form 4 equiv of $[\text{NH}_4][\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)(\text{OTf})_4]$ (**6**), and this salt subsequently reacts with the remaining tetranuclear molecules to give **10**. Indeed, compound **10** was isolated in 80% yield by treatment of a toluene solution of **2** with 1 equiv of the ionic derivative **6** at room temperature (Scheme 7).



Scheme 7. Reactions of **2** with ROTf (R = H, Me).

Compounds **8-10** were obtained in good yields (72-80%) as red solids, which are not soluble in hydrocarbon solvents but exhibit a good solubility and stability in halogenated solvents. The compounds were characterized by spectroscopic and analytical methods, as

well as by X-ray crystal structure determinations. The IR spectra (KBr) of derivatives **8** and **10** show one and two bands, respectively, for the ν_{NH} vibrations of the NH and NH_3 ligands. In addition, the IR spectrum of **10** reveals one band at 1620 cm^{-1} assignable to the NH_3 bending mode. However, the most important difference among the IR spectra of the three compounds is found in the trifluoromethanesulfonato group $\nu_{\text{as}}(\text{SO}_3)$ vibrations. In complex **9** there is only one band at 1275 cm^{-1} assignable to the CF_3SO_3^- ion,^{39b} whereas in **8** this vibration splits into two bands, at 1285 and 1255 cm^{-1} , due to the hydrogen bonding interaction between the triflate ion and the NH imido group of the cation $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)_4(\mu_3\text{-N})_3(\mu_3\text{-NH})\}]^+$.^{39c} The ^1H NMR spectra in chloroform- d_1 at room temperature of compounds **8-10** reveal two singlets in a 3:1 ratio for the C_5Me_5 ligands of the cation, which are consistent with C_{3v} -symmetric structures in solution. In addition to the resonances due to the C_5Me_5 ligands of the cation, the $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum of **10** displays one resonance signal at $\delta = 147.2$ for the *ipso* carbons of the $\eta^5\text{-C}_5\text{Me}_5$ group of the anionic fragment. This value is slightly shifted to high-field compared to that of the neutral complex $[\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)(\text{OTf})_3]$ (**7**) in chloroform- d_1 ($\delta = 149.9$), but clearly downfield with respect to that of complexes **5-7** in dimethylsulfoxide- d_6 ($\delta = 140.0$). These data further support the incorporation of dimethylsulfoxide molecules into the coordination sphere of the titanium center in complexes **5-7** in that solvent (see above), because the 16-electron complex $[\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)(\text{OTf})_4(\text{NH}_3)]^-$ should give a resonance at higher field than the 14-electron anion $[\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)(\text{OTf})_4]^-$.

The X-ray diffraction studies on complexes **8-10** confirm their ionic nature. There are no substantial differences in the cations of these compounds, except the substitution of the $\mu_3\text{-NH}$ group in **8** and **10** for a $\mu_3\text{-NMe}$ ligand in **9**. The solid-state structure of **8** shows a

weak⁴³ N–H···O hydrogen bonding interaction between the triflate anion and the NH imido group of the cationic fragment.²⁹ The structure for the cation of **10** is shown in Figure 5 and selected averaged distances and angles for the cations of the three complexes are compared in Table 4. The crystal structure of these ions consist of almost perfect [Ti₄N₄] cube cores, with the Ti–N–Ti and the N–Ti–N angles very close to 90°, in an analogue disposition of the parent complex **2**.²³ Each titanium center is bound to one pentamethylcyclopentadienyl ligand and three bridging nitrogen atoms in a three-legged piano-stool arrangement. The protonation or methylation of one nitrido ligand results in a lengthening of ca. 0.1 Å of the Ti–N distances of the {Ti₃(μ₃-NR)} fragments compared with the Ti–N separations of the {Ti₃(μ₃-N)} units. An analogous lengthening in those distances has been found in the {Ti₃(μ₄-N)MX} fragments of the Lewis acid-base adducts [{Ti(η⁵-C₅Me₅)}₄(μ₃-N)_{4-n}{(μ₄-N)MX}_n] previously prepared in our group.²⁶




Figure 5. Perspective view for the cation of **10** (thermal ellipsoids at the 50% probability level). Hydrogen atoms of the η⁵-C₅Me₅ ligands are omitted for clarity.

Table 4. Selected Averaged Lengths (Å) and Angles (deg) for the Cations of **8–10**

	8 ·3C ₆ H ₅ F ²⁹	9 ·C ₇ H ₈	10
Ti–N _{imido}	2.044(6)	2.040(2)	2.010(8)
Ti–N _{nitrido}	1.937(12)	1.932(13)	1.944(11)
Ti···Ti	2.782(1)–2.876(1)	2.782(1)–2.867(1)	2.797(1)–2.862(1)
N(1)–C(1)	–	1.394(10)	–
Ti–N _{imido} –Ti	89.4(2)	89.2(2)	90.7(1)
Ti–N _{nitrido} –Ti	91.3(1)–96.7(1)	91.6(2)–96.3(2)	91.8(1)–95.2(1)
N–Ti–N	87.6(7)	87.6(6)	87.5(6)
C(1)–N(1)–Ti	–	123.7(5)–127.9(5)	–

The anionic fragment of complex **10** (Figure 6) is similar to the titanate anion of [NMe₄][Ti(η⁵-C₅Me₅)(OTf)₄] (**5**) with the incorporation of an ammine ligand. If the centroid of the pentamethylcyclopentadienyl ring is considered, the titanium atom exhibits a distorted octahedral geometry, with the η⁵-C₅Me₅ and ammine groups located in *trans* positions and four oxygen atoms of the triflate ligands occupying the rest of the sites in the octahedron. The Ti–C_m distance of 2.128 Å in **10** is clearly longer than that found in the anion of **5** (2.045 Å) most likely due to the electron donation of the NH₃ ligand and the steric repulsions about the more crowded titanium center. The Ti(5)–N(5) bond length of 2.231(4) Å is similar to that found in the Ti–NH₃ fragment of the dinuclear complex **3** (av. 2.194(9) Å), while the Ti(5)–O bond lengths have an averaged value of 2.073(4) Å and are slightly longer than in **5** (av. Ti–O 2.014(7) Å).





Figure 6. Perspective view for the anion of **10** (thermal ellipsoids at the 50% probability level). Hydrogen atoms of the η^5 -C₅Me₅ ligands are omitted for clarity. Selected averaged bond lengths (Å) and angles (deg) for the anion: Ti(5)–N(5) 2.231(4), Ti(5)–O 2.073(4), Ti(5)–C_m 2.128, N(5)–Ti(5)–C_m 177.7, O–Ti(5)–C_m 101.7(5), O–Ti(5)–O_{cis} 88(2), O–Ti(5)–O_{trans} 156.6(9), N(5)–Ti(5)–O 78(2). C_m = Centroid of the pentamethylcyclopentadienyl ring.

We have also explored the reaction of complex [$\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-N})_4$] (**2**) with weakly acidic group 6 hydride derivatives $[\text{M}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\text{H}]$ (M = Mo, pK_a = 13.9; M = Cr, pK_a = 13.3)⁴⁷ (Scheme 8). The chromium hydride compound is a weak acid and reducing agent that has been applied to N–H bond formation in titanium compounds by PCET pathways.^{16b} Indeed, complex **2** reacted with $[\text{M}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\text{H}]$ (1 equiv) in toluene at room temperature to give the paramagnetic molecular derivative [$\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-N})_3(\mu_3\text{-NH})$] (**11**) via hydrogenation of one nitrido ligand. Compound **11** has been previously prepared by treatment of **2** with 2 equiv of ammonia borane (65% yield) or

by stepwise proton and electron transfer with HOTf and $[\text{K}(\text{C}_5\text{Me}_5)]$ (Scheme 2).²⁹ However, the synthesis with $[\text{Mo}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\text{H}]$ gave **11** in a higher isolated yield (88%) since the $[\{\text{Mo}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\}]_2$ by-product of this reaction is easily separated from **11** due to its enhanced solubility in hydrocarbon solvents.



Scheme 8. Reactions of **2** with $[\text{M}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\text{H}]$ ($\text{M} = \text{Mo}, \text{Cr}$).

In contrast to this hydrogenation reaction of a nitrido group in **2**, the treatment of **11** with an additional equivalent of $[\text{M}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\text{H}]$ afforded the ionic compounds $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-N})_2(\mu_3\text{-NH})_2][\text{M}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3]$ ($\text{M} = \text{Mo}$ (**12**), Cr (**13**)) by protonation of other nitrido group. Compounds **12** and **13** were isolated in 82 and 38% yields as dark green solids, which are not soluble in hydrocarbon solvents but exhibit a good solubility in pyridine- d_5 . The compounds are also soluble in chloroform- d_1 but readily react with this solvent to give $[\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\text{Cl}_3]$ and other unidentified species. Complexes **12** and **13** were characterized by spectroscopic and analytical methods, as well as by an X-ray crystal structure determination for **13**. The IR spectra (KBr) show two N–H stretching modes, between 3289 and 3264 cm^{-1} , as the absorptions expected for C_{2v} symmetry (A_1 and

B₂ vibrations) of the cationic fragment. These two bands are similar to those found in the neutral complex $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-CH})_2(\mu_3\text{-NH})_2]$,²⁴ while one absorption is observed in the spectra of **11** (3333 cm⁻¹) and **8** (3237 cm⁻¹).²⁹ In addition, the IR spectra of compounds **12** and **13** show strong bands, between 1936 and 1746 cm⁻¹, for the $\nu(\text{CO})$ vibrations of the $[\text{M}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3]^-$ anions. The ¹H NMR spectra of **12** and **13** in pyridine-d₅ reveal two far-downfield ($\delta = 9.3$ and 8.0) and broad resonances for the $\eta^5\text{-C}_5\text{Me}_5$ ligands in a 1:1 ratio in accord with a C_{2v} symmetry in solution. These resonance signals are comparable to those found in the spectrum of complex **11** in chloroform-d₁ ($\delta = 10.0$ and 8.7 in a 1:3 ratio) which is paramagnetic with a $\mu_{\text{eff}} = 1.70 \mu_{\text{B}}$ according to the Evans method³² determination of the magnetic susceptibility in this solvent. Similarly, compounds **12** and **13** in pyridine-d₅ solutions gave μ_{eff} values of 1.64 and 1.62 μ_{B} , which are consistent with the presence of one unpaired electron in the complexes.

The crystal structure of compound **13** contains $[\text{Cr}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3]^-$ and $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-N})_2(\mu_3\text{-NH})_2]^+$ ions. The cationic fragment shows an almost perfect [Ti₄N₄] cube core (Figure 7), with the Ti–N–Ti and N–Ti–N angles very close to 90°, similar to those found in the neutral complexes $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-N})_4]$ ²³ (**2**) and $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-N})_3(\mu_3\text{-NH})]$ (**11**).²⁹ The titanium–nitrogen(imido) lengths (av. 2.02(2) Å) are longer than titanium–nitrogen(nitrido) separations (av. 1.964(8) Å). The averaged Ti⋯Ti (2.85(3) Å) and Ti–N (1.99(3) Å) separations in **13** are slightly longer than those found in complex **2** (2.783(2) and 1.938(7) Å).




Figure 7. Perspective view of the cation of **13** (thermal ellipsoids at the 50% probability level). Hydrogen atoms of the η^5 -C₅Me₅ ligands are omitted for clarity. Selected averaged lengths (Å) and angles (deg): Ti–N(1) 2.032(8), Ti–N(2) 1.962(5), Ti–N(3) 1.998(3), Ti–N(4) 1.965(12), Ti⋯Ti 2.824(1)–2.904(1), N–Ti–N 87.8(1)–89.2(1), Ti–N–Ti 88.8(1)–93.6(1).

Conclusion

We have shown the isolation and structural characterization of several polynuclear nitrido titanium(IV) derivatives through the reaction of complexes $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)(\mu\text{-NH})\}_3(\mu_3\text{-N})]$ and $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-N})_4]$ with different molar ratios of electrophilic reagents ROTf (R = H, Me, Me₃Si; OTf = OSO₂CF₃). The electrophilic attacks with low ratios of electrophiles occur selectively at the imido ligands of $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)(\mu\text{-NH})\}_3(\mu_3\text{-N})]$ to give trinuclear or dinuclear species while the nitrido ligand remains unaltered. The functionalization of the bridging nitrido unit was only observed with a large

excess of electrophiles to afford the ammonium salts $[\text{NR}_4][\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)(\text{OTf})_4]$ ($\text{R} = \text{H}, \text{Me}$). Similarly, one of the nitrido groups of $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-N})_4]$ reacts with triflic acid or methyl triflate to form a $\mu_3\text{-NR}$ imido ligand, but a higher ratio of triflic acid produces the rupture of a fraction of the tetranuclear complex with formation of the ionic species $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-N})_3(\mu_3\text{-NH})][\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)(\text{OTf})_4(\text{NH}_3)]$. Initial stoichiometric studies of the tetranitrido $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)\}_4(\mu_3\text{-N})_4]$ with group 6 hydride complexes $[\text{M}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\text{H}]$ ($\text{M} = \text{Mo}, \text{Cr}$) results in the hydrogenation of one nitrido ligand but a second equivalent of $[\text{M}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\text{H}]$ leads to protonation of other nitrido group and not electron transfer occurs. We are currently evaluating the redox potentials of our titanium polynuclear species as well as the use of other hydride compounds in order to establish mild pathways for N–H bond formation.

Associated Content

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.inorg-chem.xxxxxx.

Experimental crystallographic data of complexes **3**, **4**, **5**, **9**, **10**, and **13**; perspective view of the crystal structures of complexes **3** and **9**; tables for selected lengths and angles of the crystal structures of **3**, **5**, **9**, **10** and **13**; selected NMR spectra (PDF).

Accession Codes

CCDC 1977628-1977633 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

Author Information:

Corresponding Author

*E-mail: carlos.yelamos@uah.es.

ORCID

Miguel Mena: 0000-0001-7456-1212

Adrián Pérez-Redondo: 0000-0002-0086-8825

Carlos Yélamos: 0000-0003-0425-4799.

Notes

The authors declare no competing financial interest.

Acknowledgments.

We are grateful to the Spanish MCIU (PGC2018-094007-B-I00) and the Universidad de Alcalá (CCG2018/EXP-008) for financial support of this research. P.B. and J.C. thank the UAH and MEC for fellowships.

References

- (1) Nishibayashi, Y. *Transition Metal-Dinitrogen Complexes: Preparation and Reactivity*; Wiley-VCH: Weinheim, 2019.
- (2) (a) Nishibayashi, Y., Ed. *Nitrogen Fixation in Top. Organomet. Chem.*, Vol. 60; Springer Verlag: Berlin/Heidelberg, 2017. (b) Schrock, R. R. Catalytic Reduction of Dinitrogen to Ammonia at a Single Molybdenum Center. *Acc. Chem. Res.* **2005**, *38*, 955–962. (c) Hoffman, B. M.; Lukoyanov, D.; Yang, Z.; Dean, D. R.; Seefeldt, L. C. Mechanism of Nitrogen Fixation by Nitrogenase: The Next Stage. *Chem. Rev.* **2014**, *114*, 4041–4062. (d) Jia, H.-P.; Quadrelli, E. A. Mechanistic Aspects of Dinitrogen Cleavage and Hydrogenation to Produce Ammonia in Catalysis and Organometallic Chemistry: Relevance of Metal Hydride Bonds and Dihydrogen. *Chem. Soc. Rev.* **2014**, *43*, 547–564.
- (3) (a) Laplaza, C. E.; Cummins, C. C. Dinitrogen Cleavage by a Three-Coordinate Molybdenum(III) Complex. *Science* **1995**, *268*, 861–863. (b) Laplaza, C. E.; Johnson, M. J. A.; Peters, J. C.; Odom, A. L.; Kim, E.; Cummins, C. C.; George, G. N.; Pickering, I. J. Dinitrogen Cleavage by Three-Coordinate Molybdenum(III) Complexes: Mechanistic and Structural Data. *J. Am. Chem. Soc.* **1996**, *118*, 8623–8638.
- (4) For a recent review on N₂ splitting, see: Klopsch, I.; Yuzik-Klimova, E. Y.; Schneider, S. Functionalization of N₂ by Mid to Late Transition Metals via N–N Bond Cleavage. *Top. Organomet. Chem.* **2017**, *60*, 71–112.
- (5) For general references on complexes with terminal nitrido ligands, see: (a) Nugent, W. A.; Mayer, J. M. *Metal-Ligand Multiple Bonds*; John Wiley & Sons: New York, 1988. (b) Eikey, R. A.; Abu-Omar, M. M. Nitrido and Imido Transition Metal Complexes of Groups 6–8. *Coord. Chem. Rev.* **2003**, *243*, 83–124. (c) Berry, J. F. Terminal Nitrido and Imido

-
- Complexes of the Late Transition Metals. *Comments Inorg. Chem.* **2009**, *30*, 28–66. (d) Smith, J. M. Reactive Transition Metal Complexes. *Prog. Inorg. Chem.* **2014**, *58*, 417–470.
- (6) (a) Rodríguez, M. M.; Bill, E.; Brennessel, W. W.; Holland, P. L. N₂ Reduction and Hydrogenation to Ammonia by a Molecular Iron-Potassium Complex. *Science* **2011**, *334*, 780–783. (b) Grubel, K.; Brennessel, W. W.; Mercado, B. Q.; Holland, P. L. Alkali Metal Control over N–N Cleavage in Iron Complexes. *J. Am. Chem. Soc.* **2014**, *136*, 16807–16816. (c) MacLeod, K. C.; McWilliams, S. F.; Mercado, B. Q.; Holland, P. L. Stepwise N–H Bond Formation from N₂-Derived Iron Nitride, Imide and Amide Intermediates to Ammonia. *Chem. Sci.* **2016**, *7*, 5736–5746. (d) Bhutto, S. M.; Holland, P. L. Dinitrogen Activation and Functionalization Using β -Diketiminato Iron Complexes. *Eur. J. Inorg. Chem.* **2019**, 1861–1869.
- (7) Lee, Y.; Sloane, F. T.; Blondin, G.; Abboud, K. A.; García-Serres, R.; Murray, L. J. Dinitrogen Activation Upon Reduction of a Triiron(II) Complex. *Angew. Chem., Int. Ed.* **2015**, *54*, 1499–1503.
- (8) For reviews on polynuclear nitrido complexes, see: (a) Dehnicke, K.; Strähle, J. The Transition Metal-Nitrogen Multiple Bond. *Angew. Chem., Int. Ed. Engl.* **1981**, *20*, 413–426. (b) Dehnicke, K.; Strähle, J. Nitrido Complexes of Transition Metals. *Angew. Chem., Int. Ed. Engl.* **1992**, *31*, 955–978. (c) Mena, M.; Pérez-Redondo, A.; Yélamos C. Heterometallic Cube-Type Molecular Nitrides. *Eur. J. Inorg. Chem.* **2016**, 1762–1778.
- (9) For Group 4 metal complexes with bridging nitrido ligands generated by N₂ cleavage, see: (a) Pool, J. A.; Lobkovsky, E.; Chirik, P. J. Hydrogenation and Cleavage of Dinitrogen to Ammonia with a Zirconium Complex. *Nature* **2004**, *427*, 527–530. (b) Nikiforov, G. B.; Vidyaratne, I.; Gambarotta, S.; Korobkov, I. Titanium-Promoted Dinitrogen Cleavage,

-
- Partial Hydrogenation, and Silylation. *Angew. Chem., Int. Ed.* **2009**, *48*, 7415–7419. (c) Semproni, S. P.; Milsman, C.; Chirik, P. J. Structure and Reactivity of a Hafnocene μ -Nitrido Prepared From Dinitrogen Cleavage. *Angew. Chem., Int. Ed.* **2012**, *51*, 5213–5216.
- (d) Semproni, S. P.; Chirik, P. J. Synthesis of a Base-Free Hafnium Nitride from N_2 Cleavage: A Versatile Platform for Dinitrogen Functionalization. *J. Am. Chem. Soc.* **2013**, *135*, 11373–11383.
- (10) For a recent review on Group 5 metal complexes with bridging nitrido ligands generated by N_2 cleavage, see: Ishida, Y.; Kawaguchi, H. Reactivity of Group 5 Element Dinitrogen Complexes and N_2 -Derived Nitrides. *Top. Organomet. Chem.* **2017**, *60*, 45–70.
- (11) (a) Fuller, A.-M.; Clegg, W.; Harrington, R. W.; Hughes, D. L.; Lancaster, S. J. Mononuclear $Ti\equiv N$ Complexes Formed by the Facile Multiple Deprotonation of $H_3N\cdot B(C_6F_5)_3$: The Importance of Chloride Ligands. *Chem. Commun.* **2008**, 5776–5778.
- (b) Fuller, A.-M.; Hughes, D. L.; Jones, G. A.; Lancaster, S. J. The Structure and Chemistry of Tris(pentafluorophenyl)borane Protected Mononuclear Nitridotitanium Complexes. *Dalton Trans.* **2012**, *41*, 5599–5609. (c) Thompson, R.; Chen, C.-H.; Pink, M.; Wu, G.; Mindiola, D. J. A Nitrido Salt Reagent of Titanium. *J. Am. Chem. Soc.* **2014**, *136*, 8197–8200. (d) Carroll, M. E.; Pinter, B.; Carroll, P. J.; Mindiola, D. J. Mononuclear and Terminally Bound Titanium Nitrides. *J. Am. Chem. Soc.* **2015**, *137*, 8884–8887. (e) Grant, L. N.; Pinter, B.; Kurogi, T.; Carroll, M. E.; Wu, G.; Manor, B. C.; Carroll, P. J.; Mindiola, D. J. Molecular Titanium Nitrides: Nucleophiles Unleashed. *Chem. Sci.* **2017**, *8*, 1209–1224.
- (12) Grant, L. N.; Pinter, B.; Gu, J.; Mindiola, D. J. Molecular Zirconium Nitride Super Base from a Mononuclear Parent Imide. *J. Am. Chem. Soc.* **2018**, *140*, 17399–17403.

-
- (13) (a) Willing, W.; Christophersen, R.; Müller, U.; Dehnicke, K. $\text{VNCl}_2(\text{Pyridin})_2$ Synthese, IR-Spektrum und Kristallstruktur. *Z. Anorg. Allg. Chem.* **1987**, *555*, 16–22. (b) Critchlow, S. C.; Lerchen, M. E.; Smith, R. C.; Doherty, N. M. Vanadium Nitride Linear Chain Polymers and Monomers. Synthesis and Structures of $[\text{V}(\mu\text{-N})\text{Cl}_2(\text{py})_2]_\infty$ and $\text{V}(\text{N})\text{Cl}_2(\text{quin})_2$. *J. Am. Chem. Soc.* **1988**, *110*, 8071–8075. (c) Tran, B. L.; Pink, M.; Gao, X.; Park, H.; Mindiola, D. J. Low-Coordinate and Neutral Nitrido Complexes of Vanadium. *J. Am. Chem. Soc.* **2010**, *132*, 1458–1459. (d) Johnson, C. E.; Kysor, E. A.; Findlater, M.; Jasinski, J. P.; Metell, A. S.; Queen, J. W.; Abernethy, C. D. The Synthesis and Characterization of $[\text{IMesH}]^+[(\eta^3\text{-C}_5\text{H}_5)\text{V}(\text{N})\text{Cl}_2]^-$: An Anionic Vanadium(V) Complex with a Terminal Nitrido Ligand. *Dalton Trans.* **2010**, *39*, 3482–3488. (e) Groysman, S.; Villagrán, D.; Freedman, D. E.; Nocera, D. G. Dinitrogen Binding at Vanadium in a Tris(alkoxide) Ligand Environment. *Chem. Commun.* **2011**, *47*, 10242–10244. (f) Tran, B. L.; Pinter, B.; Nichols, A. J.; Konopka, F. T.; Thompson, R.; Chen, C.-H.; Krzystek, J.; Ozarowski, A.; Telser, J.; Baik, M.-H.; Meyer, K.; Mindiola, D. J. A Planar Three-Coordinate Vanadium(II) Complex and the Study of Terminal Vanadium Nitrides from N_2 : A Kinetic or Thermodynamic Impediment to N–N Bond Cleavage? *J. Am. Chem. Soc.* **2012**, *134*, 13035–13045.
- (14) (a) Mindiola, D. J.; Meyer, K.; Cherry, J.-P. F.; Baker, T. A.; Cummins, C. C. Dinitrogen Cleavage Stemming from a Heterodinuclear Niobium/Molybdenum N_2 Complex: New Nitridoniobium Systems Including a Niobazene Cyclic Trimer. *Organometallics* **2000**, *19*, 1622–1624. (b) Figueroa, J. S.; Piro, N. A.; Clough, C. R.; Cummins, C. C. A Nitridoniobium(V) Reagent That Effects Acid Chloride to Organic Nitrile Conversion: Synthesis via Heterodinuclear (Nb/Mo) Dinitrogen Cleavage, Mechanistic Insights, and

-
- Recycling. *J. Am. Chem. Soc.* **2006**, *128*, 940–950. (c) Camp, C.; Grant, L. N.; Bergman, R. G.; Arnold, J. Photo-Activation of d^0 Niobium Imido Azides: *en Route* to Nitrido Complexes. *Chem. Commun.* **2016**, *52*, 5538–5541.
- (15) (a) Shaver, M. P.; Fryzuk, M. D. Activation of Molecular Nitrogen: Coordination, Cleavage and Functionalization of N_2 Mediated By Metal Complexes. *Adv. Synth. Catal.* **2003**, *345*, 1061–1076. (b) Fryzuk, M. D. Side-on End-on Bound Dinitrogen: An Activated Bonding Mode That Facilitates Functionalizing Molecular Nitrogen. *Acc. Chem. Rev.* **2009**, *42*, 127–133. (c) Andino, J. G.; Mazumder, S.; Pal, K.; Caulton, K. G. New Approaches to Functionalizing Metal-Coordinated N_2 . *Angew. Chem., Int. Ed.* **2013**, *52*, 4726–4732.
- (16) (a) Pappas, I.; Chirik, P. J. Ammonia Synthesis by Hydrogenolysis of Titanium–Nitrogen Bonds Using Proton Coupled Electron Transfer. *J. Am. Chem. Soc.* **2015**, *137*, 3498–3501. (b) Pappas, I.; Chirik, P. J. Catalytic Proton Coupled Electron Transfer from Metal Hydrides to Titanocene Amides, Hydrazides and Imides: Determination of Thermodynamic Parameters Relevant to Nitrogen Fixation. *J. Am. Chem. Soc.* **2016**, *138*, 13379–13389. (c) Bezdek, M. J.; Pappas, I.; Chirik, P. J. Determining and Understanding N–H Bond Strengths in Synthetic Nitrogen Fixation Cycles. *Top. Organomet. Chem.* **2017**, *60*, 1–22.
- (17) (a) Shima, T.; Hu, S.; Luo, G.; Kang, X.; Luo, Y.; Hou, Z. Dinitrogen Cleavage and Hydrogenation by a Trinuclear Titanium Polyhydride Complex. *Science* **2013**, *340*, 1549–1552. (b) Guru, M. M.; Shima, T.; Hou, Z. Conversion of Dinitrogen to Nitriles at a Multinuclear Titanium Framework. *Angew. Chem., Int. Ed.* **2016**, *55*, 12316–12320. (c) Wang, B.; Luo, G.; Nishiura, M.; Hu, A.; Shima, T.; Luo, Y.; Hou, Z. Dinitrogen Activation by Dihydrogen and a PNP-Ligated Titanium Complex. *J. Am. Chem. Soc.* **2017**, *139*, 1818–1821. (d) Shima, T.; Hou, Z. Dinitrogen Fixation by Transition Metal Hydride

-
- Complexes. *Top. Organomet. Chem.* **2017**, *60*, 23–43. (e) Shima, T.; Luo, G.; Hu, S.; Luo, Y.; Hou, Z. Experimental and Computational Studies of Dinitrogen Activation and Hydrogenation at a Tetranuclear Titanium Imide/Hydride Framework. *J. Am. Chem. Soc.* **2019**, *141*, 2713–2720.
- (18) (a) Schlögl, R. in *Handbook of Heterogeneous Catalysis*, 2nd ed. (Eds. G. Ertl, G. Knözinger, F. Schüth, J. Weitkamp), Wiley-VCH: Weinheim, 2008, vol. 5, pp. 2501–2575. (b) Honkala, K.; Hellman, A.; Remediakis, I. N.; Logadottir, A.; Carlsson, A.; Dahl, S.; Christensen, C. H.; Nørskov, J. K. Ammonia Synthesis from First-Principles Calculations. *Science* **2005**, *307*, 555–558. (c) Ertl, G. Reactions at Surfaces: From Atoms to Complexity (Nobel Lecture). *Angew. Chem., Int. Ed.* **2008**, *47*, 3524–3535.
- (19) (a) Abarca, A.; Gómez-Sal, P.; Martín, A.; Mena, M.; Poblet, J.-M.; Yélamos, C. Ammonolysis of Monopentamethylcyclopentadienyl Titanium(IV) Derivatives. *Inorg. Chem.* **2000**, *39*, 642–651. (b) Bai, G.; Roesky, H. W.; Noltemeyer, M.; Hao, H.; Schmidt, H.-G. Synthesis of the First Compound with a Rhombohedral $\text{Ti}_6(\mu_3\text{-NH})_6(\mu_3\text{-N})_2$ Core Structure by Ammonolysis of a Titanium Chelate in a Two-Phase System. *Organometallics* **2000**, *19*, 2823–2825. (c) Aguado-Ullate, S.; Carbó, J. J.; González-del Moral, O.; Martín, A.; Mena, M.; Poblet, J.-M.; Santamaría, C. Ammonia Activation by μ_3 -Alkyldiynide Fragments Supported on a Titanium Molecular Oxide Model. *Inorg. Chem.* **2011**, *50*, 6269–6279.
- (20) (a) Banaszak Holl, M. M.; Wolczanski, P. T. Polyamidoimidonitride Clusters of Zirconium. *J. Am. Chem. Soc.* **1992**, *114*, 3854–3858. (b) Bai, G.; Müller, P.; Roesky, H. W.; Usón, I. Intramolecular Coupling of Two Cyclopentadienyl Ring Systems of Zirconium—Unprecedented Formation of a Dihydride and Preparation of the $\{[(\text{MeC}_5\text{H}_4\text{Me})\text{Zr}]\}_5(\mu_5\text{-$

-
- N)(μ_3 -NH) $_4$ (μ -NH $_2$) $_4$] Cluster in a Two-Phase System. *Organometallics* **2000**, *19*, 4675–4677. (c) Bai, G.; Roesky, H. W.; Müller, P. Ammonolysis of M–Cl Bonds of Organozirconium(IV) and Titanium(III) Chlorides in a Liquid Ammonia/Toluene Two Phase System. *Bull. Pol. Acad. Sci. Chem.* **2002**, *50*, 1–10. (d) Bai, G.; Vidovic, D.; Roesky, H. W.; Magull, H. A Novel Potassium-Centered Highly Symmetrically Polynuclear Zirconium Complex: $\text{K}[\{(\text{Cp}^*\text{Zr})_3(\mu_3\text{-N})(\mu_3\text{-NH})(\mu\text{-NH}_2)_3\}_4(\text{NH}_2)_5(\text{NH}_3)_7]$. *Polyhedron* **2004**, *23*, 1125–1129.
- (21) (a) Banaszak Holl, M. M.; Wolczanski, P. T.; Van Duyne, G. D. Ladder Structure of $[(^t\text{BuCH}_2)_2\text{TaN}]_5 \cdot \text{NH}_3 \cdot 2\text{C}_7\text{H}_8$ and Its Relationship to Cubic TaN. *J. Am. Chem. Soc.* **1990**, *112*, 7989–7994. (b) Banaszak Holl, M. M.; Kersting, M.; Pendley, B. D.; Wolczanski, P. T. Ammonolysis of Tantalum Alkyls: Formation of Cubic TaN and a Trimeric Nitride, $[\text{Cp}^*\text{MeTaN}]_3$. *Inorg. Chem.* **1990**, *29*, 1518–1526.
- (22) Roesky, H. W.; Bai, Y.; Noltemeyer, M. Synthesis and Structure of $[\{(\eta^5\text{-C}_5\text{Me}_5)\text{Ti}(\text{NH})\}_3\text{N}]$, a Titanium Imide Nitride. *Angew. Chem., Int. Ed. Engl.* **1989**, *28*, 754–755.
- (23) Gómez-Sal, P.; Martín, A.; Mena, M.; Yélamos, C. Synthesis and Molecular Structure of the First Organometallic Nitride Cubane: $[\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)]_4(\mu_3\text{-N})_4$. *J. Chem. Soc. Chem. Commun.* **1995**, *39*, 2185–2186.
- (24) González-Moreiras, M.; Mena, M.; Pérez-Redondo, A.; Yélamos, C. Cleavage of Dinitrogen from Forming Gas by a Titanium Molecular System under Ambient Conditions. *Chem. Eur. J.* **2017**, *23*, 3558–3561.

-
- (25) Carbó, J. J.; Martín, A.; Mena, M.; Pérez-Redondo, A.; Poblet, J.-M.; Yélamos, C. Addition of Terminal Alkynes to a Molecular Titanium-Zinc Nitride. *Angew. Chem., Int. Ed.* **2007**, *46*, 3095–3098.
- (26) Carbó, J. J.; Martínez-Espada, N.; Mena, M.; Mosquera, M. E. G.; Poblet, J.-M.; Yélamos, C. Lewis Base Behavior of Bridging Nitrido Ligands of Titanium Polynuclear Complexes. *Chem. Eur. J.* **2009**, *15*, 11619–11631.
- (27) Caballo, J.; González-Moreiras, M.; Mena, M.; Pérez-Redondo, A.; Yélamos, C. Electrophilic Attack on Trinuclear Titanium Imido-Nitrido Systems. *Dalton Trans.* **2012**, *41*, 6069–6071.
- (28) Caballo, J.; González-Moreiras, M.; Mena, M.; Pérez-Redondo, A.; Yélamos, C. Reactivity with Electrophiles of Imido Groups Supported on Trinuclear Titanium Systems. *Inorg. Chem.* **2013**, *52*, 11519–11529.
- (29) Caballo, J.; González-Moreiras, M.; Greño, M.; Mena, M.; Pérez-Redondo, A.; Yélamos, C. Partial Hydrogenation of a Tetranuclear Titanium Nitrido Complex with Ammonia Borane. *Inorg. Chem.* **2014**, *53*, 8851–8853.
- (30) Mena, M.; Royo, P.; Serrano, R.; Pellinghelli, M. A.; Tiripicchio, A. Electron-Deficient (Pentamethylcyclopentadienyl)titanium Trialkyls: Evidence of Ti···H–C and Ti···C–C Interactions. Crystal and Molecular Structure of μ -[*o*-(CH₂)₂C₆H₄]{(η^5 -C₅Me₅)Ti[*o*-(CH₂)₂C₆H₄]}₂. *Organometallics* **1989**, *8*, 476–482.
- (31) (a) Fischer, E. O. Cyclopentadienyl Tricarbonyl Hydrides of Chromium, Molybdenum, and Tungsten. *Inorg. Synth.* Vol. VII, 1963, 136–139. (b) Burchell, R. P. L.; Sirsch, P.; Decken, A.; McGrady, G. S. A Structural Study of [CpM(CO)₃H] (M = Cr, Mo and W) by Single-

Crystal X-ray Diffraction and DFT Calculations: Sterically Crowded yet Surprisingly Flexible Molecules. *Dalton Trans.* **2009**, 5851–5857.

- (32) (a) Evans, D. F. The Determination of the Paramagnetic Susceptibility of Substances in Solution by Nuclear Magnetic Resonance. *J. Chem. Soc.* **1959**, 2003–2005. (b) Sur, S. K. Measurement of Magnetic Susceptibility and Magnetic Moment of Paramagnetic Molecules in Solution by High-Field Fourier Transform NMR Spectroscopy. *J. Magn. Reson.* **1989**, 169–173. (c) Bain, G. A.; Berry, J. F. Diamagnetic Corrections and Pascal's Constants. *J. Chem. Educ.* **2008**, *85*, 532–536.
- (33) (a) Chin, C. S.; Chong, D.; Lee, B.; Jeong, H.; Won, G.; Do, Y.; Park, Y. J. Activation of Acetonitrile in $[\text{Cp}^*\text{Ir}(\eta^3\text{-CH}_2\text{CHCHPh})(\text{NCMe})]^+$: Crystal Structures of Iridium–Amidine, Imino–Ether, Amido, and Amide Complexes. *Organometallics* **2000**, *19*, 638–648. (b) Bartoli, S.; Roelens, S. Binding of Acetylcholine and Tetramethylammonium to a Cyclophane Receptor: Anion's Contribution to the Cation– π Interaction. *J. Am. Chem. Soc.* **2002**, *124*, 8307–8315.
- (34) Farrugia, L. J. *WinGX* and *ORTEP for Windows*: An Update. *J. Appl. Crystallogr.* **2012**, *45*, 849–854.
- (35) Sheldrick, G. M. A Short History of *SHELX*. *Acta Crystallogr., Sect. A: Found. Crystallogr.* **2008**, *64*, 112–122.
- (36) Sheldrick, G. M. *SHELXT* – Integrated Space–Group and Crystal-Structure Determination. *Acta Crystallogr., Sect. A: Found. Adv.* **2015**, *71*, 3–8.
- (37) Sheldrick, G. M. Crystal Structure Refinement with *SHELXL*. *Acta Crystallogr., Sect. C: Struct. Chem.* **2015**, *71*, 3–8.

-
- (38) Nakamoto, K. *Infrared and Raman Spectra of Inorganic and Coordination Compounds*, 5th ed.; Wiley: New York, 1997.
- (39) (a) Lawrance, G. A. Coordinated Trifluoromethanesulfonate and Fluorosulfate. *Chem. Rev.* **1986**, *86*, 17–33. (b) Johnston, D. H.; Shriver, D. F. Vibrational Study of the Trifluoromethanesulfonate Anion: Unambiguous Assignment of the Asymmetric Stretching Modes. *Inorg. Chem.* **1993**, *32*, 1045–1047. (c) Huang, W.; Frech, R.; Wheeler, R. A. Molecular Structures and Normal Vibrations of CF_3SO_3^- and Its Lithium Ion Pairs and Aggregates. *J. Phys. Chem.* **1994**, *98*, 100–110.
- (40) (a) Duan, Z.; Verkade, J. G. A Novel Bis(azatitanatrane) with a T-Shaped Nitrido Bridging Ligand: Synthesis and Structure of $\{[\text{N}(\text{CH}_2\text{CH}_2\text{N-}i\text{-Pr})_3\text{Ti}]_2(\mu_3\text{-N})\text{Na}(\text{THF})\}$. *Inorg. Chem.* **1996**, *35*, 5325–5327. (b) Carmalt, C. J.; Milehan, J. D.; White, A. J. P.; Williams, D. J. Synthesis and Characterisation of a Bridging Nitrido Complex of Titanium. *New J. Chem.* **2000**, *24*, 929–930.
- (41) (a) Ngo, S. C.; Toscano, P. J.; Welch, J. T. Preparation and Characterization of Half-Sandwich (Pentamethylcyclopentadienyl)triflatotitanium(IV) Complexes: Solid-State Structural Motifs and Catalysis Studies. *Helv. Chim. Acta* **2002**, *85*, 3366–3382. (b) Kessler, M.; Hansen, S.; Hollmann, D.; Klahn, M.; Beweries, T.; Spannenberg, A.; Brückner, A.; Rosenthal, U. Synthesis of $\text{Cp}^*\text{Ti}(\text{OTf})$ and Its Reaction with Water. *Eur. J. Inorg. Chem.* **2011**, 627–631.
- (42) (a) García-Castro, M.; Martín, A.; Mena, M.; Poblet, J.-M.; Yélamos, C. Iodine Attack on the Metalloligand $[\{\text{Ti}(\eta^5\text{-C}_5\text{Me}_5)(\mu\text{-NH})\}_3(\mu_3\text{-N})]$: Surprising Formation of the $[\text{Ti}_3(\eta^5\text{-C}_5\text{Me}_5)_3\text{I}_2(\mu\text{-NH})_3(\text{NH}_3)]^+$ Cation. *Eur. J. Inorg. Chem.* **2006**, 1155–1160. (b) Benda, C. B.; Waibel, M.; Fässler, T. F. On the Formation of Intermetalloid Clusters:

Titanocene(III)diammin as a Versatile Reactant Toward Nonastannide Zintl Clusters. *Angew. Chem., Int. Ed.* **2015**, *54*, 522–526.

- (43) (a) Jeffrey, G. A. *An Introduction to Hydrogen Bonding*; Oxford University Press: New York, 1997. (b) Steiner, T. The Hydrogen Bond in the Solid State. *Angew. Chem., Int. Ed.* **2002**, *41*, 48–76.
- (44) Nesmeyanov, A. N.; Fedin, E. I.; Petrovskii, P. V.; Dubovitskii, V. A.; Nogina, O. V.; Kochetkova, N. S. ¹³C NMR Spectra and Electronic Effects in Cyclopentadienyl Derivatives of Metals II. Reference Point for Chemical Shifts of the Key Carbon Atom in the Series of C Substituents. *Zh. Strukt. Khim.* **1975**, *16*, 759–764.
- (45) Pevec, A. Crystal Structures of (η⁵-C₅Me₅)TiCl₃ and (η⁵-C₅Me₄H)TiCl₃. *Acta Chim. Slov.* **2003**, *50*, 199–206.
- (46) Lee, J.; Do, Y.; Kim, Y. Facile Synthesis and X-Ray Structures of (η⁵-C₅Me₅)Ti(OAr^F)₃ (OAr^F = OC₆F₅, OCH₂C₆F₅, and OCH₂C₆F₂H₃). *J. Organomet. Chem.* **2007**, *692*, 3593–3598.
- (47) Jordan, R. F.; Norton, J. R. Kinetic and Thermodynamic Acidity of Hydrido Transition-Metal Complexes. 1. Periodic Trends in Group 6 Complexes and Substituent Effects in Osmium Complexes. *J. Am. Chem. Soc.* **1982**, *104*, 1255–1263.