Northeastern Illinois University

NEIU Digital Commons

NEIU Student Research and Creative Activities Symposium

2021 NEIU Student Research and Creative Activities Symposium

Apr 23rd, 10:45 AM

Characterizing the Molecular Function of the Mutagen Sensitivity Gene, mus109

Vada Becker Northeastern Illinois University

Chandini Mitchell Winthrop University

Jordan DeLoach Winthrop University

Kathryn Kohl Winthrop University

Elyse Bolterstein Northeastern Illinois University

Follow this and additional works at: https://neiudc.neiu.edu/srcas

Becker, Vada; Mitchell, Chandini; DeLoach, Jordan; Kohl, Kathryn; and Bolterstein, Elyse, "Characterizing the Molecular Function of the Mutagen Sensitivity Gene, mus109" (2021). *NEIU Student Research and Creative Activities Symposium*. 3.

https://neiudc.neiu.edu/srcas/2021/s07/3

This Event is brought to you for free and open access by the Conferences and Symposia at NEIU Digital Commons. It has been accepted for inclusion in NEIU Student Research and Creative Activities Symposium by an authorized administrator of NEIU Digital Commons. For more information, please contact h-owen3@neiu.edu,l-wallis@neiu.edu.

CHARACTERIZING THE MOLECULAR FUNCTION OF THE MUTAGEN SENSITIVITY GENE, *MUS109*

Vada Becker¹, Chandani Mitchell², Jordan DeLoach², Kathryn Kohl Ph.D.², Elyse Bolterstein Ph.D.¹ ¹Department of Biology, Northeastern Illinois University, Chicago, Illinois 60625 ²Department of Biology, Winthrop University, Rock Hill, SC 29733

DNA repair pathways are essential in repairing damage that otherwise could cause genomic instability and cancer. Because mutations in DNA repair genes are linked to numerous human diseases, elucidating the molecular functions of DNA repair pathways will improve our understanding of disease mechanisms. Drosophila melanogaster has orthologs to most human DNA repair genes, so we can study mutants with reduced DNA repair function to investigate mechanism. Our work investigates the function of mus109, which is thought to be involved in DNA repair because allelic mutants are sensitive to various DNA damaging reagents. There are three available mus109 mutant alleles: the lethal mutant loss of function allele mus109^{ls}, and the hypomorphic alleles *mus109^{D1}* and *mus109^{D2}*. Our collaborators at Winthrop University used complementation crosses of these mutant alleles to map the location of *mus109* on the X chromosome and confirmed mutant sensitivity to the DNA alkylating reagent methyl methansulfonate (MMS). We performed a protein sequence alignment comparing *mus109* mutant alleles and found that mutations in each allele result in a loss of nuclease and/or helicase domains. To elucidate mus109 involvement in various DNA repair pathways, we tested larval sensitivity of different allele combinations to DNA damaging reagents. We have preliminary data suggesting that different mutant alleles have different responses to bleomycin, a chemotherapeutic that causes double strand breaks. Hypomorph mus109^{D1}/mus109^{D2} larvae are not sensitive to bleomycin whereas mus109¹⁵/mus109^{D2} larvae are mildly sensitive. We also have preliminary data suggesting mus109^{D1}/mus109^{D2} larvae are not sensitive to the single-strand break-inducer, camptothecin. These data indicate that *mus109* is involved in DNA repair pathways that resolve double-strand breaks and that *mus109* alleles have variable sensitivities to different mutagens, which could be explained by the variation in protein output for each allele. We plan to continue to assess *mus109* mutant larval sensitivity to reagents such as the replication fork staller, hydroxyurea, to further characterize the function of the gene in DNA repair.