ANNUAL REVIEWS

Annual Review of Public Health

Climate Change, Food Supply, and Dietary Guidelines

Colin W. Binns,¹ Mi Kyung Lee,² Bruce Maycock,^{3,4} Liv Elin Torheim,⁵ Keiko Nanishi,⁶ and Doan Thi Thuy Duong⁷

- ¹ School of Public Health, Curtin University, Perth, Western Australia 6845, Australia; email: c.binns@curtin.edu.au
- ²College of Science, Health, Engineering and Education, Murdoch University, Murdoch, Western Australia 6150, Australia; email: m.k.lee@murdoch.edu.au
- ³College of Medicine and Health, University of Exeter, Exeter EX1 2LU, United Kingdom
- ⁴Asia-Pacific Academic Consortium of Public Health (APACPH), APACPH KL Secretariat Office, Department of Social and Preventive Medicine, Faculty of Medicine, University of Malaya, 50603 Kuala Lumpur, Malaysia; email: bmaycock@iinet.net.au
- ⁵Faculty of Health Sciences, Department of Nursing and Health Promotion, Oslo Metropolitan University, NO-0130 Oslo, Norway, email: livtor@oslomet.no
- ⁶Office of International Academic Affairs, Graduate School of Medicine, The University of Tokyo, Bunkyo-Ku, Tokyo 113-0033, Japan; email: keikonanishi@yahoo.co.jp
- ⁷Faculty of Social Sciences, Behavior and Health Education, Hanoi University of Public Health, Bac Tu Liem District, Hanoi 100000, Vietnam; email: dttd@huph.edu.vn



www.annualreviews.org

- Download figures
- Navigate cited references
- · Keyword search
- · Explore related articles
- Share via email or social media

Annu. Rev. Public Health 2021. 42:233-55

First published as a Review in Advance on January 26, 2021

The Annual Review of Public Health is online at publhealth.annualreviews.org

https://doi.org/10.1146/annurev-publhealth-012420-105044

Copyright © 2021 by Annual Reviews. This work is licensed under a Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See credit lines of images or other third-party material in this article for license information.



Keywords

climate change, food supply, human nutrition, dietary guidelines, breastfeeding, Sustainable Development Goals

Abstract

Food production is affected by climate change, and, in turn, food production is responsible for 20–30% of greenhouse gases. The food system must increase output as the population increases and must meet nutrition and health needs while simultaneously assisting in achieving the Sustainable Development Goals. Good nutrition is important for combatting infection, reducing child mortality, and controlling obesity and chronic disease throughout the life course. Dietary guidelines provide advice for a healthy diet, and the main principles are now well established and compatible with sustainable development. Climate change will have a significant effect on food supply; however, with political commitment and substantial investment, projected improvements will be sufficient to provide food for the healthy

diets needed to achieve the Sustainable Development Goals. Some changes will need to be made to food production, nutrient content will need monitoring, and more equitable distribution is required to meet the dietary guidelines. Increased breastfeeding rates will improve infant and adult health while helping to reduce greenhouse gases.

1. INTRODUCTION

This article is the third review on nutrition and climate change in the Annual Review of Public Health (ARPH), which reflects its importance to global health and extends the discussion of food for population health and changes brought on by climate change (63, 102). A decade ago, the Lancet Commission concluded that anthropogenic climate change threatened to undermine the healthy lives of billions and to undo the past 50 years of gains in public health. An effective response to climate change could be the greatest global health opportunity of the 21st century (36). Food production is affected by climate change, which in turn is responsible for 20-30% of greenhouse gases (GHGs). The food system must increase output as the population increases and must meet nutrition and health needs while simultaneously assisting in the achievement of the Sustainable Development Goals (SDGs) (15, 99, 136). The climate crisis comes from a rapid increase in human use of natural resources to satisfy rising standards of living, increased energy use, population growth, and dietary shifts toward higher consumption of animal products (45). The effects of climate change on nutrition and health have been discussed for at least six decades (49, 87). Many of the causes and outcomes identified decades ago have since worsened, with only limited international cooperation emerging to change global trajectory. Substantial increases in the production of CO₂ and other GHGs have resulted in overall global warming during the Anthropocene (1). While a proportion of the CO₂ generated is utilized by plants, excess accumulates in the atmosphere or oceans, the latter being the great CO₂ sink. The result is increasing ocean acidification, altering its ecology (35, 116). Water vapor is also a GHG, levels of which increase by rising temperatures and in turn increase rainfall in some areas, contributing to adverse weather events. If climate change continues unabated, in a generation 1-3 billion people will be attempting to live in temperatures experienced now by only small populations in the Sahara (154).

Since the publication of the previous ARPH articles on this subject, world events have illustrated the fragility of global initiatives against climate change; efforts to reduce the rate of change stall as major countries withdraw or do not meet commitments outlined in the Paris Accord (85). The rate of climate change is increasing: 2015–2019 was the hottest five-year period on record, resulting from ever-increasing CO₂ emissions (152). There has been increased crop damage from storms, droughts, floods, salination, landslides, and wildfires on all continents. The loss of plants and animals from these events, together with the concentration of cropping on fewer hectares, reduces biodiversity (62). These changes will increase climate-related strain on food supplies and nutrition and health, with the coronavirus disease 2019 (COVID-19) pandemic imposing additional health and economic burdens.

The aim of this review is to consider the ways in which climate change is altering food supply and how these changes will relate to dietary guidelines in the future.

1.1. Progress and Challenges in Global Health, Food, and Nutrition

Substantial improvements have been made to health and nutrition in recent decades, even though a number of the Millennium Development Goals targets were not fully met. The global under-5 child mortality rate decreased from 118/1,000 live births in 1980 to 39/1,000 in 2018. In West and Central Africa, the corresponding values were 217/1,000 down to 97/1,000. Changes in China

are even more dramatic; infant mortality declined from an estimated 300/1,000 (in 1941) to 7 per 1,000 (in 2018), demonstrating concerted efforts to improve nutrition and health (139). The global prevalence of underweight due to insufficient energy intake has been reduced by nearly 50% since 1990, driven largely by increased yields of the major crops, including wheat, rice, and maize (53). The Intergovernmental Panel on Climate Change (IPCC) has estimated that, since 1961, per capita availability of food has increased by one-third (96). This increase has required greater use of nitrogen fertilizers (up by 800%) and water resources for irrigation (by at least double). Despite the improvements in the proportion of the population suffering from nutrition problems, in absolute numbers an estimated 821 million people are currently undernourished, 150 million children under five are stunted, and 613 million women aged 15–49 suffer from iron deficiency (96). The world is still faced with the tragedy of nearly 5 million child deaths annually. Micronutrient deficiencies affect an estimated 2 billion people worldwide, which could increase in the coming decades as food diversity decreases (133). Improvements in food supply resulted in a shift to the right of the distribution curve of food consumption, decreasing undernutrition. An unwanted side effect was an increase in the global prevalence of overweight and obesity.

Climate change and malnutrition in all its forms, including obesity and undernutrition, constitute two of the greatest threats to planetary and human health (44). Risks to human health include increases in infectious diseases, heat-related chronic disease, allergies, and injuries and stress from adverse weather events (30, 63, 79, 144). Pollution occurring concurrently with climate change may affect health (98). The exact impacts will vary with location and national wealth, but most impact will still fall on children and poorer populations. Children should be the focus of climate change discussion and implementation of the SDGs (34, 157). This approach is consistent with the legal obligations to which most countries have committed by their ratification of the Convention of the Rights of the Child, which includes obligations to protect children's right to a healthy food environment (140).

The Global Burden of Disease (GBD) study provides a comprehensive description of the impact of suboptimal diets on chronic disease morbidity and mortality, around 60% of the risk of disability-adjusted life years (DALYs), and demonstrates the need for improving nutrition in all regions and at all levels of development to control disease burden (2, 61, 127). Despite the decrease in world poverty levels and increase in world food production, both in absolute terms and in per capita availability, dietary inequalities persist, a condition all countries have committed to ending by 2030 in accord with the SDGs. Overall food energy supply by region, with inequities between rich and poor populations, is illustrated in **Figure 1** (54).

While food production has increased, a large amount of food is wasted, estimated at 35% (11). Food waste is often underestimated and has a double or even triple effect, as energy is required for production, by humans in agriculture labor, and in food preparation, all contributing to GHGs (143). Wastage occurs at many stages of food production, including mechanical damage and spillage during harvest, sorting, handling, storage, transportation, processing, retailing, and home use (14, 58). Wastage at final consumption varies, depending on economic level; for fruit and vegetables, it is estimated to be 2–5% in Africa, Asia, and Latin America compared with 12–17% in Europe, North America, Oceania, and industrialized parts of Asia (51, 55).

1.2. Food Security and the United Nations Sustainable Development Goals (SDGs)

The Food and Agricultural Organization (FAO) lists four dimensions of food security (50):

- Sufficient food is available.
- Everyone has access to it.




Figure 1

Average dietary energy supply by region. Figure adapted from Reference 35.

- The food supply is well utilized.
- All three of these dimensions are stable over time.

Sustainable diets are defined as being "protective and respectful of biodiversity and ecosystems, culturally acceptable, accessible, economically fair and affordable; nutritionally adequate, safe and healthy; while optimizing natural and human resources" (82, p. 641; 113). To the economist, that real prices of agricultural products have been trending downward for the past 50 years suggests that food production is sufficient, but the prevalence of undernutrition, overnutrition (obesity), and micronutrient deficiency shows that distribution is inequitable. The global food system faces an ambitious challenge in meeting nutritional needs for all, while reducing GHGs. Goal 2 of the SDGs emphasizes nutrition, "End hunger, achieve food security and improved nutrition and promote sustainable agriculture," but the majority of the SDGs affect food production and nutrition in some way (19). This is illustrated in **Figure 2**, using as its basis the UNICEF framework on nutrition, its antecedents, and outcomes.

1.3. Population Trends Affecting Food Requirements and Production

Additional changes in human ecology are occurring in parallel with climate change. The world population growth rate has slowed to 1.1%, but the population is still increasing by 83 million annually. The globe is becoming more urbanized, and vast numbers of migrants and refugees are changing population structures and water, food, employment, and housing requirements. The United National Development Program (UNDP) has summarized population trends that will affect the production of, and requirements for, food (142). The world's population has grown from 2.6 billion in 1950 to 7.7 billion in 2019 and will reach 8.5 billion in 2030 and 9.7 billion in 2050. By 2050, the world will have 9 billion people, and providing them with enough food, despite climate change and environmental pressures, will be the greatest challenge of the twenty-first century (40). The population will continue to age (>60 years) from 901 million in 2015 to 1.4 billion in 2030 and will continue to urbanize, with cities (>1 million) increasing from 23% of the world's population to 28% in 2030. The rural population will decline from 45% to 40%, decreasing the workforce for food production (137, 141). While the population of Sub-Saharan Africa will double by 2050, many higher-income countries (HICs) will continue to have declining




Figure 2

Relationship between UNICEF conceptual framework of undernutrition and the Sustainable Development Goals (SDGs). Numbers correspond to the relevant SDGs. Figure adapted from Reference 44.

populations. Infants, young children, pregnant women, and the elderly will be more impacted by climate change (28, 34, 131).

2. IMPACT OF CHANGES IN CLIMATE ON FOOD PRODUCTION AND SUPPLY

2.1. Changes to Global Food Supply

In a series of reports over the past 15 years, the IPCC has provided scientific evidence of the effect of climate change on global food supplies (2, 50–55, 57, 59–71).

- Hotter climates will shift production toward the poles and will also cause faster plant growth
 and ripening and decrease nutrient density.
- Areas of dry land will increase, while some regions will have increased rainfall (80). The amount of arable land in use is almost at a maximum. Increasing environmental degradation, desertification, soil depletion, overgrazing, rising sea levels, urban development, roads,

- and industrial use may reduce land further. Saltwater encroachment will affect some particularly low-lying, but highly productive rice-growing areas of Asia. As a result, agricultural productivity will have to increase.
- Adverse weather events including storms, hurricanes, droughts, flooding, landslides, and erosion will increase in frequency and severity, which will damage crops and disrupt harvesting, transportation, and storage.
- Transport will be disrupted owing to adverse weather events, increased fuel costs, conflict, and political issues.
- Spoilage and bacterial damage, including *Listeria*, *Salmonella*, and mycotoxin contamination, increase with rising temperatures and increased numbers of extreme heat days (100). Mitigating this damage will require improved agricultural practices and better processing, packaging, and storage.
- Agricultural yields depend on crop protection measures. The main purpose of pesticide use is to increase food security by controlling pests and weeds, and these will increase with climate change. The increased number of insects, including locust plagues, will cause crop damage, decrease crop yield, and result in greater costs with increased chemical residues (43, 153). Pesticide use will increase, and higher temperatures and adverse weather may cause faster dissipation (41).
- Contaminants may enter food at several stages of agriculture, processing, packaging, transport, or storage. New toxic residues (emerging contaminants) in food are increasing as a consequence of changes in industrial processes, intensifying agricultural practices, environmental pollution, and climate change (57). Chemical contaminants have become a food safety concern, owing to pesticide residues and environmental contaminants (74).
- Climate change will have a serious negative impact on crop productivity as the level of warming progresses (80). Productivity of both commercial crops (maize, rice, and wheat) and crops such as millet and sorghum will be affected. Impacts will vary depending on CO₂ concentrations, fertility levels, and region (80). A meta-analysis shows that adoption of improved farming practices and technologies such as improved varieties, planting at optimal times, and improved water and fertilizer management has the potential to reduce the negative impact on crop yield (3). With this scenario, climate change may not add significantly to the challenge of food production for the majority of countries except for some potential hot spots around the world. However, massive investment, policy, and institutional support will be required to facilitate adoption and scaling-out of such practices and to address climatic variability (3).
- Food variety will decrease, as measured by the "food diversity index" (47). Diverse food systems are more resilient in enhancing food security in the face of climate change. They are important for nutrient cycling, carbon sequestration, control of soil erosion, reduction of GHG emissions, and control of hydrological processes (97). For human nutrition, food diversity is important, especially for children, because it increases the likelihood of meeting nutritional needs, including intakes of phytochemicals, and decreases the impact of contaminants and toxicants (83, 123).
- Potable water supplies will decline owing to decreased rainfall, salinization, pollution, increasing population, and industrial use. This decline will affect water-intensive production systems, and dairying will be the most affected (67, 73). Decreases in milk production will reduce the availability of an important source of calcium and high-quality protein. At the same time, osteoporosis will become more prevalent in aging populations, which may require additional interventions. Decreases in the dairy herd may have a public health benefit

- by reducing the production and use of infant formula. More irrigation is a strategy to increase food production, but the lack of water will limit its expansion (38).
- The global food system contributes 25% of GHG emissions, with the largest source being livestock production. The food and agriculture industries project that an increase in demand for meat is likely to undermine efforts to keep global average warming below a 2°C target (114). How GHGs are accounted for may differ in the country of production compared with the country of consumption. For example, the GHG production of Hong Kong is underestimated as imports of meat are ignored. GHG emissions hidden in meat and dairy products, which are all imported into Hong Kong, demonstrate that consumption is about 59% higher than the city's total GHG emissions using conventional production-based calculations (155).
- Nuts have important benefits for nutrition, but yields will decrease with climate change (4).

2.2. Oceans, Fish, and Seafood

Fish and seafood, containing protein, a high level of omega-3 fatty acids in fatty fish, and micronutrients, are important components of a healthy diet (148). Fish provide at least 15% of the daily average intake of animal protein for 4.5 billion people, and in 2010 it provided more protein than cattle and poultry combined (13). For some populations in West Africa, island states, and coastal Asia, fish consumption is over 50% of the total animal protein consumed; small fish, eaten whole, are also an important source of many micronutrients (76). As ocean health changes, sustainable production will decrease, requiring more effort to reduce pollution (42). The world's oceans are changing because of increasing acidification (dissolved CO₂), increased fishing, minerals and petroleum extraction, and increasing use for aquaculture (35, 106). Ocean acidification has the greatest effect on calcifying organisms, including mollusks, corals, and plankton (25), which affects species higher up the food chain. Acidification can affect the growth and life span of some fish, such as sea bream, and can reduce plankton size, which decreases the growth of species such as anchovy and sardines (6, 31).

Catching and consuming fish are generally localized activities. Nine of the top 15 countries for marine capture are in the Asia-Pacific region, accounting for 87% of the global catch (52, 78). The EAT-Lancet Commission on healthy diets recommended 28 g per day of fatty fish for the protective effect of omega-3, but only the East Asia-Pacific region currently achieves this goal. The 2010 dietary guidelines for Americans encourage individuals to double their intake of fish (94), which would require a doubling of the current production to meet these recommendations (148). The oceans cannot supply these recommended levels of fish; the maximum sustainable catch has already been exceeded, with 30% of ocean wild fish stocks overfished and 60% fully fished (52). Wild fish stocks face compounding pressures from pollution, loss of habitat from coastal developments, and increased eutrophication. Increasing sea temperatures will result in the migration of stocks toward the poles (59). Aquaculture is the only method available to increase fish production, and currently two-thirds of production comes from Asia (excluding China). Changes to fish feeding methods and growth patterns may alter nutritional content, with reduced lipids, vitamin D, omega-3 fatty acids, and proteins (59, 81, 103, 134, 148). These changes will need monitoring and research to reduce the environmental impacts of mariculture (52).

2.3. Interactions Between Climate and Food Production

The global food production system is influenced by climate change, while the system itself in turn influences GHGs (see **Figure 3**). As production increases, land use will intensify and biodiversity will be lost, as fewer cultivars are used to increase yields. Land and sea pollution will increase, particularly in lower-income countries (LICs), where the majority of farmers have small




Figure 3

Food production, dietary guidelines, and healthy diet. Wastage is depicted as a negative impact (red arrows), but if wastage can be reduced, it will have a positive impact on food supply (green arrows). Abbreviations: NRV, nutrient reference value; RDA, recommended dietary allowance; RDI, recommended dietary intake; SSB, sugar-sweetened beverages.

holdings and will find it difficult to invest in technological solutions (68, 126). Some proposed climate mitigation strategies could have adverse side effects on food production; for example, using land for afforestation may reduce the land available for food production (65). The opposite effect may occur in some countries, with forests being destroyed to increase agricultural land. Another important issue for nutrition is a decrease in micronutrient density, as yield increases require improved monitoring and efforts to increase the nutrient content in food crops (90). Dietary choices affect food production, which in turn has an effect on the environment. Dietary guidelines recommend eating fewer animal products. If this guidance is followed, the decrease in production will lessen GHG emissions (115). Some of the interactions between climate change, dietary guidelines, and food production are shown in **Figure 3**.

3. BRINGING TOGETHER CLIMATE CHANGE, FOOD, AND HEALTH

3.1. A Diet That Is Good for the People and the Planet

Recommendations for healthy eating go back millennia to Hippocrates and earlier. Scientific nutrition recommendations began with the era of analytical chemistry in the nineteenth century, which led to the measurement of protein and subsequently energy, vitamins, and micronutrients. Recommendations were initially made for nutrient deficiencies in vitamin C and thiamin, followed by additional micronutrients, including iron and iodine (101, 121). Beginning in the 1950s, the infectious disease burden of undernutrition was documented, and, more recently, the burden of chronic disease, including obesity, diabetes, heart disease, and cancer, has come to the fore. Other important areas include the effect of early-life nutrition (the first 1,000 days and developmental origins of health and disease hypothesis) and the maintenance of a healthy human microbiome for good health (10, 122, 138). The promotion of sustainable, resilient food systems for healthy diets is the first principle for action during the UN Decade of Action on Nutrition (86).

3.2. Food-Based Dietary Guidelines

Most food-based dietary guidelines (FBDG) are based on the following principles: address significant public health issues, both acute and chronic; be food-based to allow for easier translation into

practical dietary advice; achieve required nutrients; maintain healthy growth and body weight; and optimize whole of life health. In addition, guidelines should meet the SDGs and reduce GHGs (22). A review of food-based dietary guidelines from 34 European countries and their sustainability credentials concluded that shifting from current Western diets to more sustainable dietary patterns produces benefits. Consuming more plant-based foods (e.g., vegetables, fruit, legumes, seeds, nuts, whole grains) and fewer animal-based foods (especially red meat), snack foods, and beverages would improve health and have environmental benefits (12). Recent regional dietary guidelines recommend similar changes and usually include physical activity, healthy body weight, breastfeeding, clean water, safe food, less sugar and salt, decreases in saturated fats, and elimination of *trans* fats (22). Another example of regional recommendations is the Nordic Nutrition Guidelines, which would produce health benefits such as a reduction in cardiovascular disease and cancer and would lessen environmental stress (117). In **Table 1**, the general recommendations in dietary guidelines are shown for the major food groups, current production levels, and suggested changes to implement by 2030.

3.3. Dietary Guidelines and Changes Needed to Ensure Sustainable Food Production

In a summary of 12 major reviews of DGs and climate change, all the diets optimized for sustainability and nutrition were more plant based, with reductions in meat, particularly ruminant meats (beef and lamb). Six studies recommended increased fish intake, while the majority suggested fewer dairy products. Other foods to be reduced included sweet foods (biscuits, cakes, desserts), savory snacks, white bread, and both alcoholic beverages and sugar-sweetened beverages (SSBs). These findings are similar to those in 7 out of 8 recent review articles on the sustainability of diets (150). Examining the 14 discrete environmental areas of concern identified in the SDGs, most studies are on GHG emissions and, to a lesser extent, land and water use. In the case of GHG emissions, changes in land use and soil carbon stocks were seldom considered, which represents a disconnect between science informing strategic climate action in the agricultural sector and science informing public health nutrition. In the case of land and water use, few studies used metrics that are appropriate in a life-cycle context. The evidence available shows that recommended diets have lower environmental impacts than do typical diets (112). A consistent scientific approach to estimating the impact of climate on diet and vice versa is required (68, 71, 72).

Many national DGs already include recommendations related to changes that would improve health and would be beneficial to the environment and sustainability. For example, Argentina Australia, China, New Zealand, Sweden, Switzerland, the United Kingdom, and the United States suggest increasing consumption of whole plant foods, vegetables, fruits, legumes, and fewer animal foods (7, 29, 32, 48, 66, 95). Canada is representative of a number of countries who also recommend reducing SSBs and eliminating *trans* fats (8).

3.4. Special Population Groups

The special needs of some population groups, including infants, children, pregnant women, and the elderly, should be considered in planning responses to climate changes and healthy eating. In many countries, complementary diets given to infants after the age of six months are high volume and often low nutrient density, limiting nutrient intake (e.g., iron) (18). While nutrient density is important in infants and young children, a diet high in fiber becomes more important in adults for the prevention of chronic disease.

Children are the population segment most susceptible to the effects of climate change, bearing an estimated 88% of the increased burden of disease (108, 110, 128). Food security for infants

Table 1 Changes to food production to meet sustainable dietary guidelines

Food group	Projected changes in food production (IPCC)	DGs recommen- dations ^a	WHO nutrition recommen- dations	Nordic recommen- dations	HICs, current availability	LICs, current availability	Conclusion: changes needed by 2030
Nuts	Decrease	Increase	Increase	Increase	Insufficient	Insufficient	Increase needed
Meat, animal products	Decrease	Decrease red and processed	Decrease	Decrease	High (reduce)	Sufficient	Decrease red and processed
Fish	Insufficient	Increase	Increase	Increase	Insufficient	Coastal areas vary; often sufficient	Increase (aquaculture)
Grains	Increase	Increase whole grain	Increase	Increase	Insufficient	Usually sufficient	Increase
Vegetables, pulses	Increase	Increase	Increase	Increase	Insufficient	Sufficient	Increase
Fruits, berries	Decrease	Increase	Increase	Increase not enough	Insufficient	Insufficient	Increase
Dairy		Decrease; use low fat		Decrease; use low fat	Too high	Insufficient	Decrease
Fats and oils		Decrease overall, especially saturated and trans	Avoid animal fats; limit saturated; No <i>trans</i>		Decrease	Decrease fats	Decrease fats
		Increase un- saturated	Prefer unsat- urated			Increase unsaturated from fish, plants	Increase unsaturated from fish, plants
Sugar, SSB	Decrease	Decrease	Decrease	Decrease	Too high	High	Decrease
Salt		Decrease	Decrease; use iodized		Too high	Too high	Decrease
Alcohol		Limit	Limit	Limit	Limit	Limit	Decrease
Breastfeeding		Increase	Increase	Increase	Insufficient EBF	Insufficient EBF	Increase EBF
Food processing ^b		Decrease ultra processing					Implement DGs in processing
Wastage	Decrease	Decrease			Too high 35%	Less than 20%	Reduce wastage

Abbreviations: DG, dietary guidelines; EBF, exclusive breastfeeding; HICs, high-income countries; IPCC, Intergovernmental Panel on Climate Change; LICs, low-income countries; SSB, sugar-sweetened beverages; WHO, World Health Organization.

Colors: green, increase; red, decrease or limit; yellow, insufficient or change needed.

and young children is not possible without promoting and achieving high rates of breastfeeding. Despite endorsements by all international professional organizations, the World Health Organization (WHO), and governments, breastfeeding targets have never been achieved on a global scale (118). The health benefits of breastfeeding, both during infancy and throughout life, including

^aRecommendations common to most dietary guidelines.

^bImportant for reducing wastage, improving bioavailability; aim to achieve DGs.

the preservation of a healthy microbiome, have been extensively studied and will provide some protection against the effects of climate change (16, 17, 149). Even in disaster situations, including famine, breastfeeding remains the best option for meeting infants' nutritional needs (23). The alternative to breastfeeding is infant formula, which is usually based on cow's milk. While milk production in the twenty-first century is more efficient than in the past, the total impact is greater owing to increased production (27). When compared with breastmilk, milk production uses large amounts of water, which will be in scarce supply with climate change, and has a high carbon footprint (124, 130).

Vietnam is an example of a low- to middle-income country (LMIC) that will experience water stress owing to climate change affecting the Mekong system; at the same time, Vietnam's milk production has increased by 12% per annum over the past decade (146). In LMICs, infant formula is increasingly advertised on the Internet, which is accessible almost everywhere by smartphone and is driving increased consumption and decreased breastfeeding. In recent years (2006–2011), imports of infant formula to Vietnam have increased by 150%, and the response of the local industry, with government support, is to boost the formula and dairy industries (125). Breastmilk is the ideal food for infants, with benefits that last into adulthood, which is important for achieving the SDGs; however, the global infant formula and dairy industries are formidable opponents (19, 21, 88, 124). Extensive promotion of breastfeeding would provide benefits for infant health, reduce health costs, and benefit the environment (46). The increase in adverse weather events along with rising population densities and uncontrolled distribution of infant formula heightens the risk to breastfeeding in disaster situations (i.e., the possibility of breastfeeding being discontinued) (23).

The elderly are vulnerable to increasing temperatures and require extra fluids and access to nutritious diets. They often move to urban environments away from traditional foods and extended family support. For them, food traditions are deeply engrained in culture and change is not easy. Traditional foods may no longer be available owing to agriculture shifts because of climate change, and meeting their nutrient needs is often compounded by dentition and mobility problems. Specific nutrients at risk include vitamin D, B₁₂, iron, and protein (105). In a Swedish study, the diets of the elderly were found to be more GHG intensive (129).

4. IMPLEMENTING CHANGE: WHAT WILL BE NEEDED?

Future objectives for the food and nutrition sector will be to improve the quality of the food supply for everyone (food and health for all) while navigating the changes expected in food production over the next several decades. Nutrient deficiencies may increase in LICs and recur in HICs, as food diversity decreases and changes in nutrient content occur. The changes in crops and farmed seafood (aquaculture) will require monitoring of nutrient composition and bioavailability, phytochemicals, and dietary content. The predominant guiding philosophy of public health nutrition has been to meet all nutrient needs through eating a variety of healthy foods. This approach may no longer be possible and may have to be met by fortification and supplements.

Micronutrient deficiencies, particularly iron, zinc, and vitamins, are of great importance (39). Issues around interventions are complex because it is difficult to ensure adequate amounts without a risk of toxicity (84). Biofortification with iron and vitamin A has proved challenging in delivering consistently safe doses while minimizing side effects, including infections (24). The preferred option is always to eat a varied diet of nutrient-dense foods.

With improved food production in recent decades, the distribution of food consumption has shifted to the right. Too little food, the left of the curve, results in undernutrition with a substantial burden of morbidity and mortality, whereas at the opposite end of the spectrum obesity and its sequelae cause major health problems for the two billion adults who are now overweight or obese.

India is an example of an LMIC that succeeded in shifting the food distribution curve to the right, with less undernutrition but more obesity (5, 156). Reduction of both undernutrition and overconsumption (obesity) simultaneously implies a compression of the food consumption curve by balancing increased physical activity with energy intake. This approach requires improvements in food and nutrition equity within and between countries (18, 109, 145). Swinburn et al. (132) describe an international pandemic of obesity, undernutrition, and climate change causing major health problems for humanity as a "syndemic." The obesity epidemic is also affecting infants and young children, owing at least in part to declines in breastfeeding, increased use of infant formula and SSBs, and disruption of the microbiome (60, 120) (see Figure 3).

Consumption of food groups such as fruits, berries, nuts, and unsaturated oils should be increased. However, these groups are projected to decrease, and research is required to increase yields. Ensuring potable water for all (for drinking, food preparation, cleaning, washing) rather than using it for increased irrigation and other projects will be difficult. The inevitable decrease in meat supplies (and other animal foods) will cause tension as decisions are made either to distribute meat more evenly around the world to meet iron and protein deficits or to supply steaks and hamburgers to wealthy countries. Additional promotion is needed for breastfeeding and for the reduction of wastage, food contamination (chemical and microbiological), and spoilage. Better packaging and storage are needed to achieve these aims.

Meeting future food needs (up to 50% more by 2050) is possible and can result in healthier diets (91). However, improved diets will not just happen from changes made to food production to moderate climate change. Health promotion will be needed to actively drive changes in eating practices (75). The strengthening of health promotion and public health programs, aimed at preventing overweight and obesity and treating chronic disease with nutrition interventions, will be an effective climate change adaptation strategy (127).

Some nutritional needs will be met with fortification or supplementation (e.g., sprinkles) rather than through an increased variety of foods (37, 111). Aquaculture and mariculture of fish with monitoring for nutrient content and improved food stocks will be needed. Nutrient content in and technological modification of cultivars will need ongoing monitoring to maintain and improve nutritional quality. Toxicants may be produced by plants in response to climate change and must be monitored. Research is needed for improving the production of fruits, nuts, berries, etc. With the loss of biodiversity, nutritional deficiencies may emerge, and it may be many decades before all nutrient functions are discovered. Potential interactions with the use of supplements and fortification should be monitored. Continuing research into the promotion of dietary guidelines and better food storage, packaging, and preservation is required. There may be a loss of cultural and culinary traditions. While change has always been a part of human society, the impact of the relatively rapidly changing climate is unknown.

Generally, the dietary guidelines that already exist are more environmentally friendly than are existing food consumption patterns, and, if applied uniformly, they will improve health and enable the SDGs to be met. A carefully selected diet that meets environmental needs can meet all nutrient requirements (9). Overcoming the barriers will depend on health promotion and motivation to change, equity in food distribution, and international collaboration to meet the threat to human life as we know it. Actions needed to improve and apply dietary guidelines in the climate change era are as follows:

- Global commitment to international collaboration on environmental action and to carbonneutral economies;
- 2. Education on nutrition and health;
- 3. Improved monitoring of food production and safety and nutrient content;

- Control on the reckless promotion of unhealthy foods by industry (and by governments, e.g., the US government pressuring countries to accept infant formula);
- Elimination of government subsidies for the production of unhealthy foods and foods that are detrimental to the environment;
- 6. Increases in the production of fruit, nuts, and vegetables (how do we do this when forests and orchards are disappearing?);
- 7. Reductions in the use of discretionary foods that are generally high in sugar, fat, or salt;
- 8. Taxes on sugar, salt, and saturated and trans fats, with incentives for nuts, etc. (107);
- 9. Breastfeeding promotion, control on infant formula sales, and implementation of WHO code (107, 147);
- 10. Improved distribution of food across countries at all levels of development along with a reduction in food wastage (104); and
- 11. More research on implementing change, including on determining how to get wealthy nations to reduce excessive intakes of environmentally costly foods.

We recommend that research focuses on the following three considerations. First, nudge theory may produce gradual change over time (20). Because healthy diets alone may not produce substantial reductions in GHG emissions, DGs need to emphasize recommendations for environmental sustainability. Minimizing the shift from current foods is likely to make changes more achievable (77). Second, changing populations' ambitions is a concern; when a country becomes richer, its population adopts Western diets. Why does this happen when Western diets are less healthy and have environmental and social disadvantages? Third, why are fat chubby babies perceived as healthy and desirable in Asia (33, 93)?

4.1. Threats to Progress

A range of responses to the simultaneous climate and COVID-19 crises are possible. Governments may seek rapid recovery from the COVID-induced economic crisis and use cheap fossil fuels to pursue economic growth regardless of the environmental costs. An optimistic scenario would see governments working together with low-carbon technologies to reduce GHGs while reducing poverty and inequality (135).

4.2. Food Equity

In nutrition, equity implies enough food to grow and develop optimally and to avoid deficiencies, stunting, obesity, infectious diseases, and excess chronic disease. It will never mean that everyone will eat exactly the same food prepared in the same way—a monotonous uniformity that ignores millennia of culinary traditions. It also means adequate amounts of potable water for drinking, food preparation, and washing.

4.3. Pandemics

Public health workers have been dreading another influenza pandemic. But the risk of other epidemic infectious diseases (EIDs), including newly emerging diseases, is ever present. Few recent epidemics have the global significance of COVID-19. Around 70% of EIDs, and almost all recent pandemics, have originated in animals (the majority in wildlife), and their emergence stems from complex interactions among wild and/or domestic animals and humans (45). The COVID-19 epidemic is imposing strains on the world food supply chain, including labor shortages, transport constraints, and the closure of food processing plants (151). People who live in areas of food crises have higher rates of malnutrition (acute, chronic, and micronutrient

deficiencies), weakened immune systems, and increased rates of chronic disease. If they contract COVID-19, the consequences will be more severe. Increased illness and death will create further food shortages owing to labor shortages, which will exacerbate the effects of climate change on food production. Individuals with obesity and noncommunicable diseases have an increased risk of a more severe course of COVID-19 infection, which should give more impetus to governments to tackle and prevent obesity and related noncommunicable diseases (119).

4.4. Conflict

Shortages of water and/or food have historically been major sources of conflict, and these conflicts continue to this day (92). Conflict or other security issues were the main cause of food crises in 2019, but weather extremes and economic shocks became increasingly significant compared with 2018. Most of the countries with food insecurity have limited resources to deal with epidemics or severe adverse weather events. The Global Report on Food Crises documents that during 2019 135 million people were classified as acutely food-insecure (up by 12% from the previous year) (70). Food insecurity along with water shortages, poor sanitation facilities, and lack of access to quality health care have contributed to declines in child nutrition. Most instances of severe food and water shortages in Africa are often related to conflict (64, 70). A long-term association has been demonstrated between food crises and conflict (26). Rapid increases in food prices in urban areas can result in social and political unrest and, at a regional level, can be a trigger for conflict. Changes in climatic conditions, including long-term climate changes, increase the risk of violence and conflict (26).

4.5. Political Inertia and Short-Term Planning

Both climate change and the emergence of overweight and obesity are slowly evolving issues, and humans have difficulty in appreciating the need for urgent action in these situations (69). Too much time has already been lost, and this effort needs to begin now and to be integrated across all disciplines (68, 72, 132). As consumers become richer, they tend to purchase more varied, higher priced, and highly processed foods, or they eat out (72). The rich pay higher prices per nutrient and outbid poorer countries for foods that provide essential nutrients, such as iron from meat. Food prices, especially of nutritious foods, may increase as a consequence of reduced productivity, putting further strains on low-income households (56). Policies aimed at reducing poverty and income inequality, while enhancing employment and income-generating activities, are key to raising people's incomes and ensuring the affordability of healthy diets.

4.6. Multinational Businesses and Climate Change

The agriculture, transport, food processing, retail, and marketing sectors need to orient their goals toward improving health and the environment by producing and advertising foods that meet the SDGs while consuming less energy. There is an increasing need for responsible corporate policy and investment to contribute toward improving the environment (89). Can the corporate world change its ways? Can they be encouraged to produce foods that minimize climate impact, are environmentally friendly, and are health oriented?

5. CONCLUSION

Substantial progress has been made in improving global nutrition. Obesity and chronic disease now represent new global challenges, even while undernutrition persists in some populations.

Reducing and even reversing anthropogenic climate change have so far proved extremely difficult or impossible. GHGs, which are environmentally detrimental, are still increasing, driven by higher standards of living in modern consumer societies. Getting people to change individual habits and cultures and to share with other nations who are less well-off has not proved possible to date.

Many studies have shown how following dietary guidelines could improve overall health and reduce obesity and chronic disease if food production, processing, distribution, and consumption follow this advice. Research has offered consistent predictions on how food production will change in the future and how these changes could be compatible with healthy diets. As the climate changes, improved technology will be needed to maintain food supplies and ensure that all nutrition requirements are met. If food variety declines, additional research will be needed to improve micronutrient content to meet nutritional needs.

Some foods that should be made more accessible and are associated with improved health, such as whole grain cereals, fruits, vegetables, legumes, nuts, berries, and olive oil, have low environmental impacts. Red meat and dairy products have the largest negative environmental impacts and are generally associated with a higher disease risk. Increases in supplies of fish are needed, while ensuring that their nutritional quality is maintained. Increasing breastfeeding rates is important and achievable for improving health and reducing the environmental impact of alternatives. Dietary transitions toward greater consumption of healthier foods would improve environmental sustainability and health. However, considerable change in the food industry and consumer behavior will be required, as, so far, no country has successfully reversed the obesity epidemic because the systemic and institutional drivers of obesity remain largely unabated.

The evidence, to date, demonstrates that major advances in sustainable food production and availability can be achieved if all the political efforts and science can be mobilized to apply improved technologies.

SUMMARY POINTS

- 1. Nutrition improved in the last few decades, but undernutrition is still prevalent in some regions while obesity is increasing in others.
- Climate change will affect health and will substantially change food production. For the medium term, it will be possible to produce enough food to maintain adequate intakes, using improved farming practices and technology and more equity in distribution.
- Following dietary guidelines would improve health, help reduce GHGs, and meet the SDGs.
- 4. Monitoring of nutrients in food will be essential.
- 5. Increasing breastfeeding has benefits for health and the environment.
- The combination of climate change and improvements in the quality of nutrition is the major public health challenge of this decade and, indeed, this century.

FUTURE ISSUES

- Ongoing monitoring of and research on food supply (quantity and quality) and nutrient content are required.
- 2. Monitor and promote breastfeeding and the marketing of artificial substitutes.

- 3. Research into the environmental footprint of foods is needed.
- 4. Continuing efforts should focus on understanding ways of encouraging populations to follow DGs. Monitor the DGs in the light of climate and health changes.

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

AUTHOR CONTRIBUTIONS

All authors have read the final manuscript and approved its contents. C.W.B. and M.K.L. undertook the initial literature search. All authors wrote sections of the manuscript. C.W.B. and M.K.L. undertook final editing and formatting before final approval by all authors.

ACKNOWLEDGMENTS

This research is funded in part by a grant from the Vietnam National Foundation for Science and Technology Development (NAFOSTED) and the National Health and Medical Research Council of Australia under grant number NHMRC.108.03-2018.09.

We are grateful to the following colleagues who reviewed the paper: Alice Lakati, AMREF University, Nairobi, Kenya; Liqian Qiu, Zhejiang University PR China; Li Tang, Chengdu Women's and Children's Central Hospital, School of Medicine, University of Electronic Science and Technology of China, Chengdu, China; Jessica Lin, Taipei Medical University; Minh Pham, Thai Nguyen University, Vietnam; Tomiko Hokama, University of the Ryukyus, Japan; Mahnaz Zarshenas, Shiraz University of Medical Science, Iran; Masaharu Kagawa, Kagawa Nutrition University, Japan; Wah Yun Low, University of Malaya, Malaysia; Raheema Abdul Raheem, The Research Centre, Maldives National University, Male', Maldives; Philip Baker, Queensland University of Technology, Australia; Hazreen Abdul Majid, University of Malaya; Noran Naqiah Hairi, University of Malaya; Awang Bulgiba Awang Mahmud, Academy of Sciences, Malaysia; and Indika Karunathilake, Faculty of Medicine, University of Colombo, Sri Lanka.

LITERATURE CITED

- Abram N, Gattuso J-P, Prakash A, Cheng L, Chidichimo MP, et al. 2019. Framing and context of the report. In IPCC Special Report on the Ocean and Cryosphere in a Changing Climate, ed. H-O Pörtner, DC Roberts, V Masson-Delmotte, P Zhai, M Tignor, et al. New York: Intergov. Panel Clim. Change (IPCC)
- Afshin A, Sur PJ, Fay KA, Cornaby L, Ferrara G, et al. 2019. Health effects of dietary risks in 195 countries, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. Lancet 393:1958–72
- Aggarwal P, Vyas S, Thornton P, Campbell BM. 2019. How much does climate change add to the challenge of feeding the planet this century? *Environ. Res. Lett.* 14:043001
- Alae-Carew C, Nicoleau S, Bird FA, Hawkins P, Tuomisto HL, et al. 2020. The impact of environmental changes on the yield and nutritional quality of fruits, nuts and seeds: a systematic review. *Environ. Res.* Lett. 15:023002
- Aleksandrowicz L, Green R, Joy EJM, Harris F, Hillier J, et al. 2019. Environmental impacts of dietary shifts in India: a modelling study using nationally-representative data. Environ. Int. 126:207–15

- Araújo JE, Madeira D, Vitorino R, Repolho T, Rosa R, Diniz M. 2018. Negative synergistic impacts
 of ocean warming and acidification on the survival and proteome of the commercial sea bream, Sparus
 aurata. 7. Sea Res. 139:50–61
- Arrieta EM, González AD. 2018. Impact of current, National Dietary Guidelines and alternative diets on greenhouse gas emissions in Argentina. Food Policy 79:58–66
- Bacon SL, Campbell NRC, Raine KD, Tsuyuki RT, Khan NA, et al. 2019. Canada's new Healthy Eating Strategy: implications for health care professionals and a call to action. Can. Fam. Phys. 65:393–98
- Bälter K, Sjörs C, Sjölander A, Gardner C, Hedenus F, Tillander A. 2017. Is a diet low in greenhouse gas emissions a nutritious diet? Analyses of self-selected diets in the LifeGene study. Arch. Public Health 75:17
- 10. Barker DJP. 2004. The developmental origins of chronic adult disease. Acta Paediatr. 93:26-33
- Barrington S, Adhikari B. 2018. Production vs. consumption management for sustainable agricultural resources. Int. 7. Innov. Sustain. Dev. 12:183–200
- Bechthold A, Boeing H, Tetens I, Schwingshackl L, Nöthlings U. 2018. Perspective: Food-based dietary guidelines in Europe—scientific concepts, current status, and perspectives. Adv. Nutr. 9:544–60
- Béné C, Barange M, Subasinghe R, Pinstrup-Andersen P, Merino G, et al. 2015. Feeding 9 billion by 2050—putting fish back on the menu. Food Secur. 7:261–74
- Bernstad AK, Canovas A, Valle R. 2017. Consideration of food wastage along the supply chain in lifecycle assessments: a mini-review based on the case of tomatoes. Waste Manag. Res. 35:29–39
- Biesbroek S, Verschuren WM, Boer JM, van der Schouw YT, Sluijs I, Temme EH. 2019. Are our diets getting healthier and more sustainable? Insights from the European Prospective Investigation into Cancer and Nutrition—Netherlands (EPIC-NL) cohort. Public Health Nutr. 22:2931–40
- Binns C, Lee M, Low WY. 2016. The long-term public health benefits of breastfeeding. Asia Pac. J. Public Health 28:7–14
- Binns C, Lee M, Scott JA. 2020. Transitioning to good health and well-being: the essential role of breast-feeding. In Sustainable Development Goals Commentaries. Vol. 3: Transitioning to Good Health and Well-Being, ed. PDA Flahault. Geneva: MDPI Books
- Binns C, Lee MK, Low WY, Baker P, Bulgiba A, et al. 2020. Guidelines for complementary feeding of infants in the Asia Pacific Region: APACPH Public Health Nutrition Group. Asia Pac. J. Public Health. 32:179–87
- Binns C, Lee MK, Low WY, Zerfas A. 2017. The role of public health nutrition in achieving the Sustainable Development Goals in the Asia Pacific Region. Asia Pac. J. Public Health 29:617–24
- Binns C, Low WY. 2017. Nobel Prizes, nudge theory, and public health. Asia Pac. J. Public Health 29:632– 34
- Binns CW, Lee MK. 2014. Exclusive breastfeeding for six months: the WHO six months recommendation in the Asia Pacific Region. Asia Pac. J. Clin. Nutr. 23:344–50
- Binns CW, Lee MK, Kagawa M, Low WY, Liqian Q, et al. 2017. Dietary Guidelines for the Asia Pacific Region. Asia Pac. J. Public Health 29:98–101
- Binns CW, Lee MK, Tang L, Yu C, Hokama T, Lee A. 2012. Ethical issues in infant feeding after disasters. Asia Pac. J. Public Health 24:672–80
- Blanco-Rojo R, Vaquero MP. 2019. Iron bioavailability from food fortification to precision nutrition. A review. *Innov. Food Sci. Emerg. Technol.* 51:126–38
- Browman HI. 2016. Applying organized scepticism to ocean acidification research. ICES J. Mar. Sci. 73:529–36
- Bruck T, d'Errico M. 2019. Food security and violent conflict: introduction to the special issue. World Dev. 117:167–71
- Capper JL, Cady RA, Bauman DE. 2009. The environmental impact of dairy production: 1944 compared with 2007. J. Anim. Sci. 87:2160–67
- Carter TR, Fronzek S, Inkinen A, Lahtinen I, Lahtinen M, et al. 2016. Characterising vulnerability of the elderly to climate change in the Nordic region. Reg. Environ. Change 16:43–58
- Castiglione C, Mazzocchi M. 2019. Ten years of five-a-day policy in the UK: Nutritional outcomes and environmental effects. Ecol. Econ. 157:185–94

- Chan AWM, Hon KL, Leung TF, Ho MHK, Duque JSR, Lee TH. 2018. The effects of global warming on allergic diseases. Hong Kong Med. 7. 24:277–84
- Checkley DM Jr., Asch RG, Rykaczewski RR. 2017. Climate, anchovy, and sardine. Annu. Rev. Mar. Sci. 9:469–93
- Chen CX, Chaudhary A, Mathys A. 2019. Dietary change scenarios and implications for environmental, nutrition, human health and economic dimensions of food sustainability. Nutrients 11:856
- Chen S, Binns CW, Maycock B, Zhao Y, Liu Y. 2014. Chinese mothers' perceptions of their child's weight and obesity status. Asia Pac. 7. Clin. Nutr. 23:452–58
- Clark H, Coll-Seck AM, Banerjee A, Peterson S, Dalglish SL, et al. 2020. A future for the world's children? A WHO-UNICEF-Lancet Commission. Lancet 395:605–58
- Cleugh H, Stafford Smith M, Battaglia M, Graham P. 2011. Climate Change: Science and Solutions for Australia. Melbourne, Aust.: CSIRO
- Costello A, Abbas M, Allen A, Ball S, Bell S, et al. 2009. Managing the health effects of climate change. Lancet 373:1693–733
- D'Agostino A, Ssebiryo F, Murphy H, Cristello A, Nakiwala R, et al. 2019. Facility- and communitybased delivery of micronutrient powders in Uganda: opening the black box of implementation using mixed methods. *Matern. Child Nutr.* 15(S5):e12798
- Darko RO, Yuan SQ, Hong L, Liu JP, Yan HF. 2016. Irrigation, a productive tool for food security—a review. Acta Agric. Scand. Sect. B-Soil Plant Sci. 66:191–206
- Darnton-Hill I. 2019. Public health aspects in the prevention and control of vitamin deficiencies. Curr. Dev. Nutr. 3:nzz075
- De Laurentiis V, Hunt DVL, Rogers CDF. 2016. Overcoming food security challenges within an energy/ water/food nexus (EWFN) approach. Sustainability 8:95
- Delcour I, Spanoghe P, Uyttendaele M. 2015. Literature review: impact of climate change on pesticide use. Food Res. Int. 68:7–15
- 42. Depledge MH, White MP, Maycock B, Fleming LE. 2019. Time and tide. BMJ 366:14671
- Devi D, Verma S, Sharma P, Sharma H, Gupta N, Thakur P. 2019. Effect of climate change on insect pests of fruit crops and adaptation and mitigation strategies: a review. *J. Entomol. Zool. Stud.* 7:507–12
- 44. Dietz WH. 2020. Climate change and malnutrition: We need to act now. J. Clin. Invest. 130:556-58
- Di Marco M, Baker ML, Daszak P, De Barro P, Eskew EA, et al. 2020. Sustainable development must account for pandemic risk. PNAS 117:3888–92
- Doan TTD, Binns C, Pham NM, Zhao Y, Dinh TPH, et al. 2020. Improved breastfeeding by empowering mothers in Vietnam: a randomised controlled trial of a mobile app. *Int. J. Environ. Res. Public Health* 17(15):5552
- Drescher LS, Thiele S, Mensink GB. 2007. A new index to measure healthy food diversity better reflects a healthy diet than traditional measures. 7. Nutr. 137:647–51
- Drew J, Cleghorn C, Macmillan A, Mizdrak A. 2020. Healthy and climate-friendly eating patterns in the New Zealand context. *Environ. Health Perspect.* 128:017007
- Dunk JH, Jones DS. 2020. Sounding the alarm on climate change, 1989 and 2019. N. Engl. J. Med. 382:205–7
- 50. FAO (Food Agric, Organ.). 1996. Documents. World Food Summit. http://www.fao.org/WFS/
- FAO (Food Agric. Organ.). 2011. Global food losses and food waste: extent, causes and prevention. Rep., FAO, Rome. http://www.fao.org/3/a-i2697e.pdf
- FAO (Food Agric. Organ.). 2018. The state of world fisheries and aquaculture 2018: meeting the Sustainable Development Goals. Rep., FAO, Rome. http://www.fao.org/3/i9540en/i9540en.pdf
- 53. FAO (Food Agric. Organ.). 2019. The state of food security and nutrition in the world: safeguarding against economic slowdowns and downturns. Rep., FAO, Rome. http://www.fao.org/3/ca5162en/ca5162en.pdf
- FAO (Food Agric. Organ.). 2019. World food and agriculture: statistical pocketbook. Rep., FAO, Rome. http://www.fao.org/3/ca6463en/ca6463en.pdf
- FAO (Food Agric, Organ.). 2020. Global food loss and waste. Sustainable Development Goals. http://www.fao.org/sustainable-development-goals/indicators/12.3.1/en/

- FAO (Food Agric. Organ.), IFAD (Int. Fund Agric. Dev.), UNICEF, WFP (World Food Progr.), WHO
 (World Health Organ.). 2020. The State of Food Security and Nutrition in the World 2020. Transforming food
 systems for affordable healthy diets. Rome: FAO. https://doi.org/10.4060/ca9692en
- Farré M, Barceló D. 2013. Analysis of emerging contaminants in food. TrAC Trends Anal. Chem. 43:240–
- Farr-Wharton G, Foth M, Choi JHJ. 2014. Identifying factors that promote consumer behaviours causing expired domestic food waste. 7. Consum. Behav. 13:393–402
- Fleming L, Maycock B, White MP, Depledge MH. 2019. Fostering human health through ocean sustainability in the 21st century. People Nat. 1:276–83
- 60. Forbes JD, Azad MB, Vehling L, Tun HM, Konya TB, et al. 2018. Association of exposure to formula in the hospital and subsequent infant feeding practices with gut microbiota and risk of overweight in the first year of life. JAMA Pediatr. 172:e181161
- 61. Forouzanfar MH, Afshin A, Alexander LT, Anderson HR, Bhutta ZA, et al., GBD Risk Factors Collab. 2016. Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 388:1659–724
- Friel S. 2020. Climate change and the people's health: the need to exit the consumptagenic system. Lancet 395:666–68
- Frumkin H, Haines A. 2019. Global environmental change and noncommunicable disease risks. Annu. Rev. Public Health 40:261–82
- FSIN (Food Secur. Inf. Netw.). 2019. Global Report on Food Crises: Joint Analysis for Better Decisions. Rep.,
 FSIN UN, New York
- Fujimori S, Hasegawa T, Krey V, Riahi K, Bertram C, et al. 2019. A multi-model assessment of food security implications of climate change mitigation. *Nat. Sustain.* 2:386–96
- 66. Gardner CD, Hartle JC, Garrett RD, Offringa LC, Wasserman AS. 2019. Maximizing the intersection of human health and the health of the environment with regard to the amount and type of protein produced and consumed in the United States. Nutr. Rev. 77:197–215
- Gauly M, Ammer S. 2020. Review: Challenges for dairy cow production systems arising from climate changes. *Animal* 14:S196–203
- Ghosh A, Misra S, Bhattacharyya R, Sarkar A, Singh AK, et al. 2020. Agriculture, dairy and fishery farming practices and greenhouse gas emission footprint: a strategic appraisal for mitigation. *Environ. Sci. Pollut. Res.* 27:10160–84
- Gills B, Morgan J. 2020. Global Climate Emergency: after COP24, climate science, urgency, and the threat to humanity. Globalizations 17:885–902
- Glob. Netw. Against Food Crises. 2020. Global report on food crises. Rep., Food Secur. Inf. Netw., Rome. https://docs.wfp.org/api/documents/WFP-0000114546/download/?_ga=2.149815478.146837182.1587518448%E2%80%931146868816.1587518448
- 71. Godfray HCJ. 2015. The debate over sustainable intensification. Food Secur. 7:199-208
- Godfray HCJ, Crute IR, Haddad L, Lawrence D, Muir JF, et al. 2010. The future of the global food system. Philos. Trans. R. Soc. B 365:2769–77
- Grout L, Baker MG, French N, Hales S. 2020. A review of potential public health impacts associated with the global dairy sector. Geobealth 4:e2019GH000213
- Guo WJ, Pan BH, Sakkiah S, Yavas G, Ge WG, et al. 2019. Persistent organic pollutants in food: contamination sources, health effects and detection methods. Int. J. Environ. Res. Public Health 16:4361
- He P, Baiocchi G, Feng KS, Hubacek K, Yu Y. 2019. Environmental impacts of dietary quality improvement in China. 7. Environ. Manag. 240:518–26
- HLPE (High Level Panel Experts Food Secur. Nutr.). 2014. Sustainable fisheries and aquaculture for food security and nutrition. Rep., HLPE, Comm. World Food Secur., FAO, Rome. http://www.fao.org/3/ai3844e.pdf
- Horgan GW, Perrin A, Whybrow S, Macdiarmid JI. 2016. Achieving dietary recommendations and reducing greenhouse gas emissions: modelling diets to minimise the change from current intakes. *Int. J. Behav. Nutr. Phys. Act.* 13:46

- Huelsenbeck M. 2012. Ocean-based food security threatened in a high CO₂ world. Rep., Oceana, Washington,
 DC. https://oceana.org/sites/default/files/reports/Ocean-Based_Food_Security_Threatened_in_a_High_CO2_World.pdf
- IPCC (Intergov. Panel Clim. Change). 2014. Climate Change 2014 Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, eds. RK Pachauri, LA Meyer. Geneva: IPCC. https://www.ipcc.ch/site/assets/uploads/2018/05/SYR_AR5_FINAL_full_wcover.pdf
- IPCC (Intergov. Panel Clim. Change). 2019. Climate Change and Land: An IPCC Special Report on Climate Change, Desertification, Land Degradation, Sustainable Land Management, Food Security, and Greenbouse Gas Fluxes in Terrestrial Ecosystems, ed. PR Shukla, J Skea, EC Buendía, V Masson-Delmotte, H-O Pörtner, et al. New York: IPCC. https://www.ipcc.ch/site/assets/uploads/sites/4/2020/08/200730-IPCCJ7230-SRCCL-Complete-BOOK-HRES.pdf
- 81. Jakobsen J, Smith C, Bysted A, Cashman KD. 2019. Vitamin D in wild and farmed Atlantic salmon (Salmo Salar)—What do we know? Nutrients 11:982
- 82. Jones AD, Hoey L, Blesh J, Miller L, Green A, Shapiro LF. 2016. A systematic review of the measurement of sustainable diets. *Adv. Nutr.* 7:641–64
- 83. Khamis AG, Mwanri AW, Ntwenya JE, Kreppel K. 2019. The influence of dietary diversity on the nutritional status of children between 6 and 23 months of age in Tanzania. *BMC Pediatr*. 19:518
- 84. Kihara J, Bolo P, Kinyua M, Rurinda J, Piikki K. 2020. Micronutrient deficiencies in African soils and the human nutritional nexus: opportunities with staple crops. *Environ. Geochem. Health* 42:3015–33
- Lang T, Mason P. 2018. Sustainable diet policy development: implications of multi-criteria and other approaches, 2008–2017. Proc. Nutr. Soc. 77:331–46
- Lawrence MA, Baker PI, Pulker CE, Pollard CM. 2019. Sustainable, resilient food systems for healthy diets: the transformation agenda. *Public Health Nutr.* 22:2916–20
- Leaf A. 1989. Potential health effects of global climatic and environmental changes. N. Engl. J. Med. 321:1577–83
- Lee MK, Binns C. 2019. Breastfeeding and the risk of infant illness in Asia: a review. Int. J. Environ. Res. Public Health 17:186
- 89. Leins S. 2020. 'Responsible investment': ESG and the post-crisis ethical order. Econ. Soc. 49:71-91
- Leisner CP. 2020. Review: Climate change impacts on food security—focus on perennial cropping systems and nutritional value. *Plant Sci.* 293:110412
- Le Mouël C, Forslund A. 2017. How can we feed the world in 2050? A review of the responses from global scenario studies. Eur. Rev. Agric. Econ. 44:541–91
- Levy BS. 2019. Increasing risks for armed conflict: climate change, food and water insecurity, and forced displacement. Int. J. Health Serv. 49:682–91
- Li B, Adab P, Cheng KK. 2015. The role of grandparents in childhood obesity in China—evidence from a mixed methods study. Int. J. Behav. Nutr. Phys. Act. 12:91
- Love DC, Fry JP, Milli MC, Neff RA. 2015. Wasted seafood in the United States: quantifying loss from production to consumption and moving toward solutions. Glob. Environ. Change 35:116–24
- Martin M, Brandão M. 2017. Evaluating the environmental consequences of Swedish food consumption and dietary choices. Sustainability 9:2227
- Mbow C, Rosenzweig C, Barioni G, Benton T, Herrero M, et al. 2019. Food security and insecurity, the food system and climate change. See Ref. 80, pp. 442–44. https://www.ipcc.ch/site/assets/uploads/ 2019/11/08_Chapter-5.pdf
- Mbow C, Rosenzweig C, Barioni G, Benton T, Herrero M, et al. 2019. Role of agroecology and diversification. See Ref. 80, pp. 468–70. https://www.ipcc.ch/site/assets/uploads/2019/11/08_Chapter-5.pdf
- McKinney A. 2019. A planetary health approach to study links between pollution and human health. Curr. Pollut. Rep. 5:394–406
- Moon BK. 2017. The Sustainable Development Goals from the perspective of the former Secretary General of the UN. Keynote address presented at the 49th Asia-Pacific Academic Consortium for Public Health Conference, Incheon, South Korea, Aug. 17

- Moretti A, Pascale M, Logrieco AF. 2019. Mycotoxin risks under a climate change scenario in Europe. Trends Food Sci. Technol. 84:38–40
- Mozaffarian D, Rosenberg I, Uauy R. 2018. History of modern nutrition science—implications for current research, dietary guidelines, and food policy. BMJ 361:k2392
- 102. Myers SS, Smith MR, Guth S, Golden CD, Vaitla B, et al. 2017. Climate change and global food systems: potential impacts on food security and undernutrition. Annu. Rev. Public Health 38:259–77
- 103. Nichols PD, Glencross B, Petrie JR, Singh SP. 2014. Readily available sources of long-chain omega-3 oils: Is farmed Australian seafood a better source of the good oil than wild-caught seafood? *Nutrients* 6:1063–79
- 104. Nooghabi SN, Burkart S, Mahmoudi H, Taheri F, Damghani AM, et al. 2018. More food or better distribution? Reviewing food policy options in developing countries. Food Rev. Int. 34:566–80
- Obeid R, Heil SG, Verhoeven MMA, van den Heuvel E, de Groot L, Eussen S. 2019. Vitamin B12 intake from animal foods, biomarkers, and health aspects. Front. Nutr. 6:93
- 106. Olsen E, Kaplan IC, Ainsworth C, Fay G, Gaichas S, et al. 2018. Ocean futures under ocean acidification, marine protection, and changing fishing pressures explored using a worldwide suite of ecosystem models. Front. Mar. Sci. 5:64
- Perez-Escamilla R, Bermudez O, Buccini GS, Kumanyika S, Lutter CK, et al. 2018. Nutrition disparities and the global burden of malnutrition. BM7 361:k2252
- 108. Philipsborn RP, Chan K. 2018. Climate change and global child health. Pediatrics 141:e20173774
- Popkin BM, Corvalan C, Grummer-Strawn LM. 2020. Dynamics of the double burden of malnutrition and the changing nutrition reality. *Lancet* 395:65–74
- Ragavan MI, Marcil LE, Garg A. 2020. Climate change as a social determinant of health. *Pediatrics* 145:e20193169
- 111. Reerink I, Namaste SML, Poonawala A, Dhillon CN, Aburto N, et al. 2017. Experiences and lessons learned for delivery of micronutrient powders interventions. *Matern. Child Nutr.* 13:e12495
- Ridoutt BG, Hendrie GA, Noakes M. 2017. Dietary strategies to reduce environmental impact: a critical review of the evidence base. Adv. Nutr. 8:933–46
- Ritchie H, Reay DS, Higgins P. 2018. The impact of global dietary guidelines on climate change. Glob. Environ. Change 49:46–55
- Ritchie H, Reay DS, Higgins P. 2018. Potential of meat substitutes for climate change mitigation and improved human health in high-income markets. Front. Sustain. Food Syst. 2. https://doi.org/10.3389/ fsufs.2018.00016
- Rose D, Heller MC, Roberto CA. 2019. Position of the Society for Nutrition Education and Behavior: the importance of including environmental sustainability in dietary guidance. J. Nutr. Educ. Behav. 51:3– 15.e1
- 116. Sage RF. 2020. Global change biology: a primer. Glob. Change Biol. 26:3–30
- 117. Saha S, Nordström J, Mattisson I, Nilsson PM, Gerdtham UG. 2019. Modelling the effect of compliance with Nordic nutrition recommendations on cardiovascular disease and cancer mortality in the Nordic countries. Nutrients 11:1434
- Salmon L. 2015. Food security for infants and young children: an opportunity for breastfeeding policy?
 Int. Breastfeed 7. 10:7
- Sattar N, McInnes IB, McMurray JJV. 2020. Obesity is a risk factor for severe COVID-19 infection: multiple potential mechanisms. Circulation 142:4–6
- Scharf RJ, DeBoer MD. 2016. Sugar-sweetened beverages and children's health. Annu. Rev. Public Health 37:273–93
- 121. Schreier P, Binns CW, Högger P, Wu D. 2013. It began with citrus. Nutr. Med. 1:5-7
- Sdona E, Briana DD, Malamitsi-Puchner A. 2020. Impact of economic crises on offspring health and the developmental origins of health and disease concept. Acta Paediatr. 109:453–59
- Sebayang SK, Dibley MJ, Astutik E, Efendi F, Kelly PJ, Li M. 2020. Determinants of age-appropriate breastfeeding, dietary diversity, and consumption of animal source foods among Indonesian children. *Matern. Child Nutr.* 16:e12889
- Smith JP. 2019. A commentary on the carbon footprint of milk formula: harms to planetary health and policy implications. *Int. Breastfeed. 7.* 14:49

- 125. Son M. 2017. Vietnam's appetite for foreign baby formula is making US giants rich. VN Express, May 26. https://e.vnexpress.net/news/business/vietnam-s-appetite-for-foreign-baby-formula-is-making-us-giant-abbott-rich-3590559.html
- 126. Springmann M, Clark M, Mason-D'Croz D, Wiebe K, Bodirsky BL, et al. 2018. Options for keeping the food system within environmental limits. *Nature* 562:519–25
- Springmann M, Mason-D'Croz D, Robinson S, Garnett T, Godfray HCJ, et al. 2016. Global and regional health effects of future food production under climate change: a modelling study. *Lancet* 387:1937–46
- Stanberry LR, Thomson MC, James W. 2018. Prioritizing the needs of children in a changing climate. PLOS Med. 15:e1002627
- Strid A, Hallström E, Hjorth T, Johansson I, Lindahl B, et al. 2019. Climate impact from diet in relation to background and sociodemographic characteristics in the Vasterbotten Intervention Programme. Public Health Nutr. 22:3288–97
- Sultana MN, Uddin MM, Ridoutt BG, Peters KJ. 2014. Comparison of water use in global milk production for different typical farms. Agric. Syst. 129:9–21
- 131. Suzuki K. 2018. The developing world of DOHaD. J. Dev. Orig. Health Dis. 9:266-69
- Swinburn BA, Kraak VI, Allender S, Atkins VJ, Baker PI, et al. 2019. The global syndemic of obesity, undernutrition, and climate change: the Lancet Commission report. *Lancet* 393:791–846
- Titcomb TJ, Tanumihardjo SA. 2019. Global concerns with B vitamin statuses: biofortification, fortification, hidden hunger, interactions, and toxicity. Compr. Rev. Food Sci. Food Sci. 18:1968–84
- Tlusty MF, Tyedmers P, Bailey M, Ziegler F, Henriksson PJG, et al. 2019. Reframing the sustainable seafood narrative. Glob. Environ. Change 59:101991
- 135. Tollefson J. 2020. How hot will Earth get by 2100? Nature 580:443-45
- UN. 2019. The Sustainable Development Goals report 2019. Rep., UN, New York. https://unstats.un.org/sdgs/report/2019/The-Sustainable-Development-Goals-Report-2019.pdf
- UN Dep. Econ. Soc. Aff. 2018. World population ageing. Rep., UN, New York. https://www.un.org/en/development/desa/population/publications/pdf/ageing/WPA2017_Highlights.pdf
- 138. UNICEF (UN Child. Fund). 2017. The first 1000 days. UNICEF. http://1000days.unicef.ph/
- UNICEF (UN Child. Fund). 2019. The State of the World's Children 2019. Children, Food and Nutrition: Growing well in a changing world. Flagship Rep., UNICEF, New York. https://www.unicef.org/reports/state-of-worlds-children-2019
- 140. UNICEF (UN Child. Fund), UN Spec. Rapp. Right Food. 2019. Protecting children's right to a healthy food environment. Rep., UNICEF, UN Hum. Rights Counc., Geneva. https://www.unicef.nl/files/Advocacy-brief-healthy-food-enviro-final.pdf
- 141. UN Popul. Div. 2019. The world's cities in 2018. Data Bookl., UN, New York. https://www.un.org/en/events/citiesday/assets/pdf/the_worlds_cities_in_2018_data_booklet.pdf
- 142. UN Popul. Div. 2020. Population trends. United Nations. https://population.un.org/wpp/
- 143. Vittuari M, De Menna F, Pagani M. 2016. The hidden burden of food waste: the double energy waste in Italy. Energies 9:660
- 144. Watts N, Adger WN, Ayeb-Karlsson S, Bai Y, Byass P, et al. 2017. The Lancet Countdown: tracking progress on health and climate change. Lancet 389:1151–64
- Wells JC, Sawaya AL, Wibaek R, Mwangome M, Poullas MS, et al. 2020. The double burden of malnutrition: aetiological pathways and consequences for health. *Lancet* 395:75–88
- 146. Whitehead R. 2020. Dairy majors investing heavily as Vietnam ramps up milk production. DAIRY Reporter, Jan. 8. https://www.dairyreporter.com/Article/2020/01/08/Dairy-majors-investing-as-Vietnam-ramps-up-milk-production
- WHO (World Health Organ.), UNICEF, IBFAN. 2020. Marketing of breast-milk substitutes: national implementation of the international code. Status report 2020. Rep., WHO, Geneva. https://www.who.int/ publications/i/item/9789240006010
- 148. Willett W, Rockström J, Loken B, Springmann M, Lang T, et al. 2019. Food in the Anthropocene: the EAT-Lancet Commission on healthy diets from sustainable food systems. Lancet 393:447–92
- 149. Wilson AS, Koller KR, Ramaboli MC, Nesengani LT, Ocvirk S, et al. 2020. Diet and the human gut microbiome: an international review. Dig. Dis. Sci. 65:723–40

- Wilson N, Cleghorn CL, Cobiac LJ, Mizdrak A, Nghiem N. 2019. Achieving healthy and sustainable diets: a review of the results of recent mathematical optimization studies. Adv. Nutr. 10:S389

 –403
- 151. Wise J. 2020. Covid-19: pandemic exposes inequalities in global food systems. BM7 369:m1915
- WMO (World Meteorol. Organ.). 2019. The global climate in 2015–2019. Press Release, Sept. 22, World Meteorol. Organ., Geneva
- 153. WMO (World Meteorol. Organ.), FAO (Food Agric. Organ.). 2016. Weather and desert locusts. WMO 1175, WMO, Geneva. https://library.wmo.int/doc_num.php?explnum_id=3213
- 154. Xu C 徐, Kohler TA, Lenton TM, Svenning J-C, Scheffer M. 2020. Future of the human climate niche. *PNAS* 117:11350–55
- 155. Yau YY, Thibodeau B, Not C. 2018. Impact of cutting meat intake on hidden greenhouse gas emissions in an import-reliant city. Environ. Res. Lett. 13:064005
- 156. Young MF, Nguyen P, Tran LM, Avula R, Menon P. 2020. A double edged sword? Improvements in economic conditions over a decade in India led to declines in undernutrition as well as increases in overweight among adolescents and women. J. Nutr. 150:364–72
- 157. Zivin JG, Shrader J. 2016. Temperature extremes, health, and human capital. Future Child. 26:31-50

Contents

Commencary	
Climate Change Disinformation and How to Combat It Stephan Lewandowsky	1
Epidemiology and Biostatistics	
Considerations for Developing an Agenda for Gun Violence Prevention Research Mark Rosenberg	23
Influenza Virus: Tracking, Predicting, and Forecasting Sheikh Taslim Ali and Benjamin J. Cowling	43
Prediabetes and What It Means: The Epidemiological Evidence *Justin B. Echouffo-Tcheugui and Elizabeth Selvin	59
Reproducible Research: A Retrospective Roger D. Peng and Stephanie C. Hicks	79
Vaccine Hesitancy, Acceptance, and Anti-Vaccination: Trends and Future Prospects for Public Health Ève Dubé, Jeremy K. Ward, Pierre Verger, and Noni E. MacDonald	175
Declining Life Expectancy in the United States: Missing the Trees for the Forest Sam Harper, Corinne A. Riddell, and Nicholas B. King	381
Trends in Abortion Policies in Low- and Middle-Income Countries Pascale Allotey, T.K. Sundari Ravindran, and Vithiya Sathivelu	
Social Environment and Behavior	
Prediabetes and What It Means: The Epidemiological Evidence	50

Overdose Epidemic Magdalena Cerdá, Noa Krawczyk, Leah Hamilton, Kara E. Rudolph, Samuel R. Friedman, and Katherine M. Keyes	95
Early Childhood Adversity, Toxic Stress, and the Impacts of Racism on the Foundations of Health Jack P. Shonkoff, Natalie Slopen, and David R. Williams	15
Expanding Implementation Research to Prevent Chronic Diseases in Community Settings Stephanie Mazzucca, Elva M. Arredondo, Deanna M. Hoelscher, Debra Haire-Joshu, Rachel G. Tabak, Shiriki K. Kumanyika, and Ross C. Brownson 13	35
Understanding and Responding to Health Literacy as a Social Determinant of Health Don Nutbeam and Jane E. Lloyd	59
Vaccine Hesitancy, Acceptance, and Anti-Vaccination: Trends and Future Prospects for Public Health Ève Dubé, Jeremy K. Ward, Pierre Verger, and Noni E. MacDonald	75
Environmental and Occupational Health	
Air Quality in Africa: Public Health Implications Asmamaw Abera, Johan Friberg, Christina Isaxon, Michael Jerrett,	93
Ebba Malmqvist, Cheryl Sjöström, Tahir Taj, and Ana Maria Vargas	, ,
At the Water's Edge: Coastal Settlement, Transformative Adaptation, and Well-Being in an Era of Dynamic Climate Risk William Solecki and Erin Friedman	
At the Water's Edge: Coastal Settlement, Transformative Adaptation, and Well-Being in an Era of Dynamic Climate Risk	11
At the Water's Edge: Coastal Settlement, Transformative Adaptation, and Well-Being in an Era of Dynamic Climate Risk William Solecki and Erin Friedman 21 Climate Change, Food Supply, and Dietary Guidelines Colin W. Binns, Mi Kyung Lee, Bruce Maycock, Liv Elin Torheim,	11

Extreme Weather and Climate Change: Population Health and Health System Implications	
Kristie L. Ebi, Jennifer Vanos, Jane W. Baldwin, Jesse E. Bell,	
David M. Hondula, Nicole A. Errett, Katie Hayes, Colleen E. Reid,	
Shubhayu Saha, June Spector, and Peter Berry2	293
Green Infrastructure and Health Mark J. Nieuwenhuijsen	317
Public Health Practice and Policy	
Climate Change Disinformation and How to Combat It Stephan Lewandowsky	1
Expanding Implementation Research to Prevent Chronic Diseases in Community Settings	
Stephanie Mazzucca, Elva M. Arredondo, Deanna M. Hoelscher, Debra Haire-Joshu, Rachel G. Tabak, Shiriki K. Kumanyika, and Ross C. Brownson	135
Addressing Social Needs in Health Care Settings: Evidence, Challenges, and Opportunities for Public Health Matthew W. Kreuter, Tess Thompson, Amy McQueen, and Rachel Garg	
Benchmarking as a Public Health Strategy for Creating Healthy Food Environments: An Evaluation of the INFORMAS Initiative (2012–2020)	
Gary Sacks, Janelle Kwon, Stefanie Vandevijvere, and Boyd Swinburn	345
Cash Transfers and Health	272
Sicong Sun, Jin Huang, Darrell L. Hudson, and Michael Sherraden	505
Declining Life Expectancy in the United States: Missing the Trees for the Forest	
Sam Harper, Corinne A. Riddell, and Nicholas B. King	381
Enhancing Community Engagement by Schools and Programs of Public Health in the United States Mindi B. Levin, Janice V. Bowie, Steven K. Ragsdale, Amy L. Gawad,	
Lisa A. Cooper, and Joshua M. Sharfstein	405
Progress in National Policies Supporting the Sustainable Development Goals: Policies that Matter to Income and Its Impact on Health Amy Raub and Jody Heymann	423
Sugar-Sweetened Beverage Reduction Policies: Progress and Promise	
James Krieger, Sara N. Bleich, Stephanie Scarmo, and Shu Wen Ng	439

Health Services

Extreme Weather and Climate Change: Population Health and Health	
System Implications	
Kristie L. Ebi, Jennifer Vanos, Jane W. Baldwin, Jesse E. Bell,	
David M. Hondula, Nicole A. Errett, Katie Hayes, Colleen E. Reid,	
Shubhayu Saha, June Spector, and Peter Berry	293
Addressing Social Needs in Health Care Settings: Evidence,	
Challenges, and Opportunities for Public Health	
Matthew W. Kreuter, Tess Thompson, Amy McQueen, and Rachel Garg	329
Improving Access to Care: Telemedicine Across Medical Domains	
William Barbosa, Kina Zhou, Emma Waddell, Taylor Myers, and E. Ray Dorsey	463
The Effects of Home Care Provider Mix on the Care Recipient: An	
International, Systematic Review of Articles from 2000 to 2020	
Norma B. Coe, R. Tamara Konetzka, Melissa Berkowitz, Emily Blecker,	
and Courtney H. Van Houtven	483
Trends in Abortion Policies in Low- and Middle-Income Countries	
Pascale Allotey, T.K. Sundari Ravindran, and Vithiya Sathivelu	505
Indexes	
Cumulative Index of Contributing Authors, Volumes 33–42	519
Cumulative Index of Article Titles, Volumes 33-42	526

Errata

An online log of corrections to *Annual Review of Public Health* articles may be found at http://www.annualreviews.org/errata/publhealth