

ENVIRONMENTAL RESEARCH LETTERS

LETTER • **OPEN ACCESS**

Combined heat and drought suppress rainfed maize and soybean yields and modify irrigation benefits in the USA

To cite this article: Xiangyu Luan *et al* 2021 *Environ. Res. Lett.* **16** 064023

View the [article online](#) for updates and enhancements.

ENVIRONMENTAL RESEARCH
LETTERS

LETTER

Combined heat and drought suppress rainfed maize and soybean yields and modify irrigation benefits in the USA

OPEN ACCESS

RECEIVED

22 October 2020

REVISED

6 April 2021

ACCEPTED FOR PUBLICATION

28 April 2021

PUBLISHED

21 May 2021

Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Xiangyu Luan¹ , Riccardo Bommarco² , Anna Scaini^{3,4} and Giulia Vico¹ ¹ Swedish University of Agricultural Sciences, Department of Crop Production Ecology, SE 750 07 Uppsala, Sweden² Swedish University of Agricultural Sciences, Department of Ecology, SE 750 07 Uppsala, Sweden³ Stockholm University, Department of Physical Geography, SE 106 91 Stockholm, Sweden⁴ Stockholm University, Bolin Centre for Climate Research, SE 106 91 Stockholm, SwedenE-mail: giulia.vico@slu.se**Keywords:** agriculture, crop yield, climate change, drought, rain, seasonal extremesSupplementary material for this article is available [online](#)**Abstract**

Heat and water stress can drastically reduce crop yields, particularly when they co-occur, but their combined effects and the mitigating potential of irrigation have not been simultaneously assessed at the regional scale. We quantified the combined effects of temperature and precipitation on county-level maize and soybean yields from irrigated and rainfed cropping in the USA in 1970–2010, and estimated the yield changes due to expected future changes in temperature and precipitation. We hypothesized that yield reductions would be induced jointly by water and heat stress during the growing season, caused by low total precipitation (P_{GS}) and high mean temperatures (T_{GS}) over the whole growing season, or by many consecutive dry days (CDD_{GS}) and high mean temperature during such dry spells (T_{CDD}) within the season. Whole growing season (T_{GS} , P_{GS}) and intra-seasonal climatic indices (T_{CDD} , CDD_{GS}) had comparable explanatory power. Rainfed maize and soybean yielded least under warm and dry conditions over the season, and with longer dry spells and higher dry spell temperature. Yields were lost faster by warming under dry conditions, and by lengthening dry spells under warm conditions. For whole season climatic indices, maize yield loss per degree increase in temperature was larger in wet compared with dry conditions, and the benefit of increased precipitation greater under cooler conditions. The reverse was true for soybean. An increase of 2 °C in T_{GS} and no change in precipitation gave a predicted mean yield reduction across counties of 15.2% for maize and 27.6% for soybean. Irrigation alleviated both water and heat stresses, in maize even reverting the response to changes in temperature, but dependencies on temperature and precipitation remained. We provide carefully parameterized statistical models including interaction terms between temperature and precipitation to improve predictions of climate change effects on crop yield and context-dependent benefits of irrigation.

1. Introduction

The harvest we reap from our crop fields depend to large extent on the climatic conditions and their fluctuation (Porter and Semenov 2005). Temperature and precipitation explain one-third of global yield variation (Ray *et al* 2015). Climate change is expected to put pressure on crop production (IPCC 2019) and has already caused yield losses (Lobell *et al* 2011). Increased frequency of co-occurring high temperatures and low precipitation (Mazdiyasi and AghaKouchak 2015, Alizadeh *et al* 2020) suppresses

crop yields by causing heat and water stress in the crop plants (Lesk *et al* 2016, Zscheischler *et al* 2017). Carefully parameterized yield models that include interaction terms between temperature and precipitation hold potential to improve predictions of climate change impacts on crop yield (Carter *et al* 2018).

Plant physiology and small-scale field and controlled-environment experiments tell us that stress combinations can have synergistic effects on yield formation (Suzuki *et al* 2014). Reduced water availability limits the plant's ability to regulate its temperature via evaporative cooling, thereby increasing

its vulnerability to high temperatures (Siebert *et al* 2014, Neukam *et al* 2016). Experiments demonstrate how yields are more suppressed when heat and water stresses are combined than their summed effects when occurring in isolation (Prasad *et al* 2011, Mahrookashani *et al* 2017, Cohen *et al* 2021). Analyses of yields from arable fields with detailed climatic data show that interactive effects of precipitation and temperature or precipitation and vapor pressure deficit are needed to explain yield variation (Urban *et al* 2015, Carter *et al* 2018). Yet, the combined effects of heat and water stress on yields have often been overlooked in field-scale experiments and modeling (Rötter *et al* 2018).

The combined effects of precipitation and temperature have been examined for national crop yields. Yields across Europe were better explained with the bivariate return period of warm or cold temperatures and high or low precipitation total, compared with models relying only on temperature and precipitation (Zscheischler *et al* 2017). National maize and soybean yield losses were exacerbated in hot and dry seasons globally, in the USA and India (Matiu *et al* 2017), the American Midwest (Carter *et al* 2018, Kukul and Irmak 2018), and France (Hawkins *et al* 2013). But aggregating yields over vast geographical areas, such as with national data, can mask and average out adverse climatic conditions that often have more limited geographic range (Matiu *et al* 2017).

Crops respond to both seasonal average conditions and unfavorable conditions of shorter duration. Climatic conditions integrated over the growing season, such as precipitation totals and temperature means, are well correlated with crop yield across nations and regions (Lobell *et al* 2011, Osborne and Wheeler 2013, Challinor *et al* 2014, Zhao *et al* 2017). But averaging over the growing season can mask short-term but potentially damaging conditions, which can cause substantial yield losses (Lobell *et al* 2012, 2013, Troy *et al* 2015, Lesk *et al* 2016, Vogel *et al* 2019) depending on timing, duration, and intensity of the unfavorable conditions (Tack *et al* 2017). Analyses with a finer spatial and temporal resolution across large geographic areas and long time series, with climatic indices based on knowledge of plant physiological response to environmental stressors, could improve our understanding and predictions of climate impacts on yield.

Although crops are exposed to the same climatic conditions, yield can be differently affected under irrigated and rainfed cropping (Siebert *et al* 2017). Irrigation can, at least partially, mitigate negative effects from adverse climatic conditions on crop yields (Troy *et al* 2015, Leng 2017, Li and Troy 2018, Zhu *et al* 2019), directly by alleviating crop water stress (Zipper *et al* 2016) and indirectly by reducing heat stress through evaporative cooling (Siebert *et al* 2017, Tack *et al* 2017, Luan and Vico 2021). However,

yield data collected across nations and regions are usually not separated into irrigated and rainfed yields. And, even where these data are available (as in parts of the USA), there is a lack of large-scale analysis of the interactive effects of temperature and precipitation on yield, explicitly considering the role of irrigated or rainfed cropping. Analyzing irrigated and rainfed crop yields from the same geographic location will elucidate the mitigating effects of irrigation on yields and how these effects might be linked to temperature and precipitation.

We explored the interactive effects of temperature and precipitation and the role of irrigation on fine-resolution (i.e. county-level) crop yields from the USA 1970–2010. We combined, for the first time, all of these factors in the same analysis, using county-level data, distinguishing irrigated and rainfed yields, and considering the interactive effects of temperature- and precipitation-related climatic indices. We considered two sets of climatic indices to capture different physiological mechanisms: (a) mean climatic conditions during the whole growing season (subscript GS), i.e. growing season precipitation total, P_{GS} , and mean air temperature, T_{GS} , and (b) shorter-term intra-seasonal conditions, as represented by maximum number of consecutive days with precipitation <2 mm during the growing season, CDD_{GS} , and mean daily air temperature during this dry period, T_{CDD} . We hypothesized that (a) there are compounded damaging impacts of combined high temperature and reduced water availability, which correspond to potential negative effects of heat and water stress, (b) irrigation reduces negative impacts of the two climatic stresses, both when occurring separately and when combined, and (c) measures of within-season conditions explain crop responses to heat and water stress at least as well as the whole season indices.

2. Materials and methods

2.1. Crop yields

We selected two staple crops, soybean and maize, grown both rainfed and irrigated, with a wide geographical distribution in the USA (supplementary information, SI, figure S1 (available online at stacks.iop.org/ERL/16/064023/mmedia)). County-level grain yield data for the period 1970–2010, separated for rainfed and irrigated cropping, were obtained from the US Department of Agriculture National Agricultural Statistical Service (Quick Stats; for details see SI S1.1).

2.2. Climatic indices and their calculations

We selected four climatic indices as candidate explanatory variables for crop yields to represent variation in air temperature and plant soil water availability during the growing season. The growing season was

defined as the period between the local mean planting and harvesting dates (SI S1.3).

Two climatic indices were chosen to reflect air temperature and soil water availability during the whole growing season (subscript GS): mean daily air temperature averaged over the growing season (T_{GS}), and total precipitation over the growing season (P_{GS}), reflecting the input to the soil water balance.

Two climatic indices were chosen to represent conditions of shorter duration but linked to potential water and heat stresses within the growing season. The maximum number of consecutive dry days with daily precipitation less than 2 mm within the growing season (CDD_{GS}) reflects the length of each season's longest dry spell, during which soil water availability is gradually reduced: the longer the period, the more likely the occurrence of water stress. The second intra-seasonal index was the mean daily air temperature during CDD_{GS} (T_{CDD}). While CDD_{GS} can also fall in periods other than the warmest part of the growing season, CDD_{GS} and T_{CDD} describe conditions occurring at the same time, including the combination of potentially damaging heat and water stress. We repeated analyses described below also setting as threshold for a dry day daily precipitation of 0 and 1 mm, reaching similar conclusions.

To calculate the climatic indices, we used daily gridded data of precipitation and air temperature at $1/8^\circ$ spatial resolution (Maurer *et al* 2002) and information on the timing and duration of the growing season at spatial resolution of $1/2^\circ$ (Sacks *et al* 2010). These data were spatially aggregated at the county scale, before calculating the seasonal and intra-seasonal climatic indices (SI S1.2).

2.3. Statistical analyses and model predictions

We explored how crop yields varies with selected climatic indices and management (i.e. irrigated or rainfed cropping), via mixed effects statistical models explicitly including interactions of these drivers (e.g. Gałecky and Burzykowski 2013).

For each crop, the seasonal and intra-seasonal indices were analyzed in two separate statistical models, with yield (either maize or soybean yield) as dependent variable. Yields and climatic indices were not de-trended prior to analysis. Instead, the effect of likely trends from climate change and technological advances were considered by including time as the continuous variable t , as years elapsed from 1969. The fixed factors also included the precipitation-related index P_{GS} for the seasonal and CDD_{GS} for the intra-seasonal analysis, and the temperature-related index T_{GS} for the seasonal and T_{CDD} for the intra-seasonal analysis. Management M was included as categorical variable, for either irrigation or rainfed cropping. The two- and three-way interactions among the factors temperature, precipitation and management

were added as fixed parts to the model. The resulting fixed part of the model reads:

$$\begin{aligned} \text{Yield} = & \beta_0 + \beta_P x_P + \beta_T x_T + \beta_t t \\ & + \beta_{PT} x_P x_T + \Delta_0 M + \Delta_P M x_P \\ & + \Delta_T M x_T + \Delta_{PT} M x_P x_T \end{aligned} \quad (1)$$

where t is years elapsed from 1969, M is the management ($M = 0$ for rainfed and $M = 1$ for irrigated), and $x_P = P_{GS}$ and $x_T = T_{GS}$ for the seasonal statistical model and $x_P = CDD_{GS}$ and $x_T = T_{CDD}$ for the intra-seasonal one, i.e. two models were fitted for each crop. Regarding the coefficients, β_0 is the model intercept and Δ_0 the change in intercept from a shift from rainfed to irrigated cropping. Further, β_i and Δ_i are the slopes describing the changes in yield explained by precipitation- and temperature-related indices ($i = P$ or T respectively) and their interactions ($i = PT$), for rainfed cropping (β_i) and how these slopes are changed by a shift from rainfed to irrigated cropping (Δ_i). To these models, we added as random factors the interaction between county and management M , to allow for different responses to irrigation and rainfed cropping in each county. We also added year as categorical variable to the random part of the model, to account for the covariance across large geographic areas in climatic conditions within a year. The model was fitted to the data across all counties, as detailed in SI, S2.1. No model simplification was done, because treatment interactions were an inherent part of this study.

The model estimates in table 1 enable calculations of yield outcomes under any temperature and precipitation within the explored range of climatic indices (SI S2.2). Below, we first explored the sensitivity of yield to changes in temperature- and precipitation-related indices. Second, we showed how the model can be used to predict impacts on yield from changes of climatic conditions in line with climate change projections. We show results for a 2°C increase in both temperature indices, accompanied by reduced, unchanged or increased precipitation indices. We used currently observed climatic conditions each year and county as baseline. The percentage change was averaged over the years within each county.

3. Results

As expected, increasing temperature reduced rainfed yields (negative coefficient for T_{GS} in table 1(a)). Precipitation increased yields in both crops, but more so in maize than soybean (table 1, figure 1). Irrigated crops yielded consistently more than rainfed crops (figure 2). There were also several significant two- and three-way interactions among management, precipitation and temperature, for both whole season and intra-seasonal climatic indices (table 1). Hence,

Table 1. Model estimates, their standard errors (SE) and significance (p) from the linear mixed effects model for maize and soybean, for (a) whole growing season and (b) intra-seasonal indices, where M is water management and t is years elapsed from 1969. The model estimates (equation (1)) are presented with rainfed cropping as the baseline level for the categorical variable of management M . The marginal and conditional R^2 give information on the goodness of fit of the model to data, i.e. the variation explained by the fixed factors alone and by the entire model including fixed and random factors, respectively.

Predictor	Maize			Soybean		
	Symbol in equation (1) (units)	Estimate	SE	Estimate	SE	p
(a) Whole growing season results where $x_P = P_{GS}$ and $x_T = T_{GS}$						
Intercept		4.184	5.912×10^{-1}	6.214	3.134×10^{-1}	<0.001
x_P	β_0 (ton ha ⁻¹)	7.772×10^{-2}	1.055×10^{-2}	-3.908×10^{-3}	6.188×10^{-3}	0.528
x_T	β_P (ton ha ⁻¹ cm ⁻¹)	-1.921×10^{-1}	2.720×10^{-2}	-2.463×10^{-1}	1.348×10^{-2}	<0.001
t	β_T (ton ha ⁻¹ yr ⁻¹)	1.057×10^{-1}	8.030×10^{-3}	2.788×10^{-2}	2.486×10^{-3}	<0.001
M	β_t (ton ha ⁻¹ yr ⁻¹)	2.612	7.201×10^{-1}	-9.421×10^{-1}	3.696×10^{-1}	0.0108
$x_P \times M$	Δ_0 (ton ha ⁻¹)	-1.082×10^{-1}	1.431×10^{-2}	-1.355×10^{-2}	8.076×10^{-3}	0.0935
$x_T \times M$	Δ_P (ton ha ⁻¹ cm ⁻¹)	1.551×10^{-1}	3.473×10^{-2}	1.108×10^{-1}	1.618×10^{-2}	<0.001
$x_P \times x_T$	Δ_T (ton ha ⁻¹ °C ⁻¹)	-1.925×10^{-3}	5.051×10^{-4}	7.125×10^{-4}	2.724×10^{-4}	0.00892
$x_P \times x_T \times M$	β_{PT} (ton ha ⁻¹ cm ⁻¹ °C ⁻¹)	3.575×10^{-3}	6.884×10^{-4}	4.751×10^{-6}	3.580×10^{-4}	0.989
Number of observations	Δ_{PT} (ton ha ⁻¹ cm ⁻¹ °C ⁻¹)	15 580		10 148		
Marginal R^2		0.723		0.657		
Conditional R^2		0.883		0.829		
(b) Intra-seasonal results where $x_P = CDD_{GS}$ and $x_T = T_{CDD}$						
Intercept		2.575	2.526×10^{-1}	1.662	1.052×10^{-1}	<0.001
x_P	β_0 (ton ha ⁻¹)	-9.323×10^{-3}	4.923×10^{-3}	-7.997×10^{-3}	3.871×10^{-3}	0.0389
x_T	β_P (ton ha ⁻¹ d ⁻¹)	-1.660×10^{-2}	6.535×10^{-3}	-8.702×10^{-3}	3.660×10^{-3}	0.0174
t	β_T (ton ha ⁻¹ °C ⁻¹)	1.087×10^{-1}	8.606×10^{-3}	2.836×10^{-2}	2.410×10^{-3}	<0.001
M	β_t (ton ha ⁻¹ yr ⁻¹)	4.296	1.992×10^{-1}	7.095×10^{-1}	1.184×10^{-1}	<0.001
$x_P \times M$	Δ_0 (ton ha ⁻¹)	-2.323×10^{-2}	6.731×10^{-3}	2.198×10^{-3}	5.179×10^{-3}	0.671
$x_T \times M$	Δ_P (ton ha ⁻¹ d ⁻¹)	-1.754×10^{-2}	9.048×10^{-3}	6.161×10^{-4}	4.978×10^{-3}	0.902
$x_P \times x_T$	Δ_T (ton ha ⁻¹ °C ⁻¹)	-3.342×10^{-4}	2.559×10^{-4}	-3.523×10^{-4}	1.672×10^{-4}	0.0352
$x_P \times x_T \times M$	β_{PT} (ton ha ⁻¹ d ⁻¹ °C ⁻¹)	1.807×10^{-3}	3.519×10^{-4}	5.607×10^{-4}	2.252×10^{-4}	0.0128
Number of observations	Δ_{PT} (ton ha ⁻¹ d ⁻¹ °C ⁻¹)	15 580		10 148		
Marginal R^2		0.695		0.577		
Conditional R^2		0.874		0.794		





Figure 1. Crop yields as a function of temperature- and precipitation-related climatic indices for (a) whole growing season total precipitation (P_{GS}) and mean daily temperature (T_{GS}), and (b) for intra-seasonal largest number of consecutive dry days (CDD_{GS}) and mean daily temperature during this period (T_{CDD}). The contour plots are based on the fixed part of the statistical model (equation (1)), with coefficient estimates for maize and soybean (table 1), and are relative to rainfed (left) and irrigated (right) cropping. Yields refer to year 1991, i.e. the middle-point of the period considered. The ranges of the climatic indices correspond to those of the observations (SI, figure S3).




yields depended on the combination of temperature and precipitation, which also altered the benefit from irrigation (figures 3 and S7 in the SI). Yields increased over the years (positive coefficient for t in table 1). The explanatory powers of the whole season and intra-seasonal models were similar, although models based on whole season climatic indices explained somewhat more of the yield variation (marginal R^2 in table 1).

3.1. Rainfed yield responses to whole growing season climatic conditions

The mean seasonal daily temperature (T_{GS}) and seasonal accumulated precipitation (P_{GS}) affected rainfed yields in both maize and soybean (table 1(a)), where high temperature and low precipitation gave the lowest yields (table 1(a), figure 1(a)). The interaction $x_P \times x_T$ was significant, i.e. the effects on yield of temperature and precipitation were not equal across their respective ranges (table 1(a), figure 3). For instance, at a seasonal precipitation of 280 mm there was a loss of 0.25 ton ha⁻¹ of maize per °C rise in

temperature, while at 490 mm precipitation the loss increased to 0.29 tons ha⁻¹ °C⁻¹ (solid green arrows in figure 3(a)). Hence, a 1 °C rise in temperature had a larger negative impact on maize yields at higher precipitation totals. The reverse was true for soybean. At 280 mm precipitation 0.23 ton ha⁻¹ soybean were lost for °C increase in temperature, but at 490 mm precipitation the loss was 0.21 ton ha⁻¹ °C⁻¹ (dotted green arrows in figure 3(a)).

Precipitation increased maize yields more at low than at high temperatures: a 10 mm increase in precipitation at 19 °C increased maize yield by 0.041 ton ha⁻¹, while the same precipitation increase at 23 °C increased the yield by 0.033 ton ha⁻¹ (solid green arrows in figure 3(b)). Again, the reverse was true for soybean, where 10 mm more precipitation increased yield only by 0.0096 ton ha⁻¹ at a low temperature, but 0.012 ton ha⁻¹ at a high temperature. The difference between maize and soybean can also be seen in the curvature in the surface plots, convex in maize and concave in soybean (figure 1(a)).



3.2. Rainfed yield responses to intra-seasonal climatic conditions

Long dry spells (high CDD_{GS}) and high mean temperatures during the dry spell (high T_{CDD}) reduced yields in both soybean and maize (table 1(b), figure 1(b)). For each $^{\circ}\text{C}$ rise during dry spell, yields were reduced more during longer dry spells for soybean, but not significantly so for maize (table 1(b), green arrows in figure 3(c)).

A lengthening of the dry spell by 1 d reduced yields at both low and high temperatures (green solid arrow figure 3(d)), but yields were slightly more reduced per day of dry spell lengthening at high compared with low temperatures during the dry spell for maize, but not significantly so for soybean ($x_P \times x_T$ in table 1(b), green dotted arrow figure 3(d)).

3.3. Irrigated yield responses to whole growing season climatic conditions

Irrigation increased both maize and soybean yields (figures 1 and 2, table 1(a)), except under the lowest and wettest conditions (dark red area figure 2). Irrigation dampened and sometimes reversed the effects

of temperature and precipitation changes on yield, but the interactive effects of temperature and precipitation remained for maize, but not for soybeans ($x_P \times x_T \times M$ in table 1(a); figures 2 and 3). For instance, at a seasonal precipitation of 280 mm there was an increase of $0.0092 \text{ ton ha}^{-1}$ of irrigated maize per $^{\circ}\text{C}$ rise in temperature, but a $0.044 \text{ ton ha}^{-1} \text{ } ^{\circ}\text{C}^{-1}$ increase at 490 mm precipitation (solid blue arrows in figure 3(a)). Hence, irrigation reverted negative impacts of high temperature in rainfed maize. Irrigated soybean continued to lose yield with increasing temperature, but less so compared with rainfed cropping. Irrigation benefits were larger at high precipitation (figure 2(b), dotted blue arrows in figure 3(a)).

Irrigation reduced the effects of precipitation changes. A 10 mm increase in precipitation resulted in a maize yield increase smaller than in rainfed cropping, and the increase was higher at warmer temperatures (solid blue arrows in figure 3(b)). Irrigated soybean yields instead declined by $0.0038 \text{ ton ha}^{-1}$ at low and $0.00096 \text{ ton ha}^{-1}$ at high temperatures per 10 mm added precipitation (dotted blue arrows in figure 3(b)).

3.4. Irrigated yield responses to intra-seasonal climatic conditions

Irrigation enhanced yields in both maize and soybean across dry spell lengths and temperatures (table 1(b), figure 2), and dampened and sometimes reversed effects of changes in dry spell length and temperature on yield, with significant three-way ($x_P \times x_T \times M$) interactions in both crops (table 1(b), figure 3). In maize, irrigation increased yields with a rise in temperature during longer, but not shorter, dry spells (blue solid arrows in figure 3(c)). Irrigation reduced soybean yield loss per °C increase during the dry spell slightly more at long compared with short dry spells (blue dotted arrows in figure 3(c)). The patterns were the same for a lengthening of the dry spell, at low or high dry spell temperatures (figure 3(d)).

4. Discussion

Based on detailed crop yield information from counties across the USA, in which rainfed and irrigated cropping co-exist, we provide climatic models that explain up to 72% and 66% of maize and soybean yield variability. This is an improvement compared with other analyses (e.g. Ray *et al* 2015, Zampieri *et al* 2017, Zhu *et al* 2019), we believe at least partially thanks to the use of county-level data, separately observed rainfed and irrigated yields, and explicitly including the interactions between precipitation- and air temperature-related climatic indices. As expected, high temperatures reduced rainfed maize and soybean yields (Schlenker and Roberts 2009, Schauburger *et al* 2017). These declines were accelerated by low precipitation (Lobell and Field 2007, Zscheischler *et al* 2017, Feng and Hao 2020). Maize and soybean yields decreased particularly under warm and dry weather. In addition, interaction terms between whole season temperature and precipitation indices were significant for rainfed yields, except for soybean yields explained by intra-seasonal climatic indices (table 1). This was in agreement with previous results in four states in the USA (Carter *et al* 2018), but not with an analysis based on national yield data, where interactions were found for some crops and nations, but not for soybean and maize in the USA (Matiu *et al* 2017). The interaction terms signify that the yield change due to rising temperature or precipitation differs depending on how warm and dry the conditions are (figure 3). For instance, rainfed soybean yields declined faster with rising temperature in dry conditions and benefitted more by precipitation increase under warmer conditions.

Compared with soybean, maize has higher total evapotranspiration (Suyker and Verma 2009) and optimal precipitation requirement (Dietzel *et al* 2016), but lower optimal temperature for grain filling and critical temperature for yield reduction (Schlenker and Roberts 2009, Hoffman *et al* 2020). Soybean yield was indeed less responsive than maize

to an increase in growing season temperature and precipitation (figures 3(a) and (b)), but responses were similar for changes in intra-seasonal indices (figures 3(c) and (d)). In relative terms, the mean yield reduction for the USA per degree warming has been estimated to 10.3% for maize and 6.8% for soybean (Zhao *et al* 2017). Other analyses suggest losses up to 30% (Rose *et al* 2017) and extreme variability across states (Mourtzinis *et al* 2015). For maize, the first set of estimates fits well our predicted mean yield reduction across counties of 15.2% with a 2 °C temperature increase given no change in precipitation (figure 4 gray boxes). However, the corresponding relative loss in rainfed soybean amounted to 27.6%. This loss is larger than the estimates mentioned above, but our estimation does not account for the potential mitigation from enhanced air CO₂ concentration, which will be greater in C3 species, such as soybean, than in C4 species (Makowski *et al* 2020). For both crops, the losses due to an increase in temperature were exacerbated by a concurrent reduction in precipitation or lengthening of dry spells (figure 4 orange boxes). The exact mechanism for the different sensitivities to combined changes in temperature and precipitation are complex and mediated by the timing of heat and water stress, with some particularly sensitive crop phenological stages (Hoffman *et al* 2020).

The two analyses, based on whole growing season or intra-seasonal climatic indices, provide complementary information. The intra-seasonal model had high explanatory power, just slightly lower than the whole season model (70% vs 72% variation explained for maize and 58% vs 66% for soybean, table 1), showing that unfavorable conditions with short duration are important determinants of yield. The slightly lower performance can be partially explained by the larger role of soil moisture at the beginning of the longest dry spell compared with that at the beginning of the growing season. Further, the timing of the longest dry spell was spread during the growing season, i.e. the longest dry spell did not necessarily co-occur with the warmest seasonal temperatures (SI, figure S4). Considering the timing and shift in timing of dry spells (Breinl *et al* 2020) might improve predictions and support management decisions (Zipper *et al* 2016).

The choice of whether to irrigate or not has large impacts on crop yields and their dependence on climatic conditions, underlining the importance of explicitly evaluating the outcomes of rainfed and irrigated agriculture when assessing the role of climatic conditions. In general, irrigation increases yields by reducing the negative effects of dry and hot weather conditions or both (Zhang *et al* 2015, Leng 2017, Tack *et al* 2017, Li and Troy 2018). Here, irrigation was most effective in enhancing yields under conditions of low precipitation or extended dry spells and high temperatures (figure 2), when water shortage is most likely and extreme and rainfed yields




Figure 4. Distribution of county-average percent changes in yield, as calculated based on the fixed part of the statistical model (equation (1), estimates in table 1) for three expected effects of climate change on temperature and precipitation for maize (top) and soybean (bottom), based on climatic indices describing growing season (left, P_{GS} and T_{GS}) and intra-seasonal (right, CDD_{GS} and T_{CDD}) conditions. We present yield changes predicted by the statistical model in equation (1) from a 2°C increase in temperature-related indices (T_{GS} in the left column and T_{CDD} in the right column). To this warming, we superimpose three shifts in precipitation: (a) a 10% reduction in P_{GS} and increase in CDD_{GS} (orange box, left), (b) unaltered P_{GS} and CDD_{GS} (gray box, center), and (c) a 10% increase in P_{GS} and reduction in CDD_{GS} (blue box, right). In all cases, the relative change in yield is calculated using as baseline the yield predicted by the model under currently observed climatic conditions in the same year and county. A single data point per county is included in the box plots, representing the percentage change averaged over all the years for which data are available. Horizontal bars are means, boxes extend from the 1st to the 3rd quartile, whiskers from the 5th to the 95th percentiles, and symbols are points outside this range.

were low (figure 1). In the central USA, irrigation was able to reduce the combined effects of precipitation and accumulated extreme temperatures in maize and soybean (Zhang *et al* 2015), as well as the effects of dry spells, precipitation totals and several types of temperature extremes in isolation (Troy *et al* 2015). In another analysis, the marginal gain of irrigation was reduced as seasonal precipitation increased in both maize and soybean, but temperature had a less definite effect (Li and Troy 2018).

Irrigation also reduced the dependence of crop yields on climatic conditions (shorter blue than green arrows in figure 3) and in some cases even reverted the direction of response. Irrigation mitigated and in the case of maize reverted the effects of increased temperature, pointing to the importance of water availability to alleviate also heat stress by supporting evaporative cooling. In the case of soybean, irrigation mitigated, but did not cancel the negative effects of increased temperature (figure 3(a)), in agreement with local and global analyses (Tack *et al* 2017, Agnolucci *et al* 2020). This underlines the importance of reducing

water stress to increase the optimal temperature for crop yield and stave off the negative effects of high temperatures (Siebert *et al* 2017, Agnolucci *et al* 2020). Irrigation also reduced the positive effects of an increase in precipitation in maize, while a shift from positive to negative dependence of yields on precipitation when shifting from rainfed to irrigation cropping emerged in soybean (figure 3(b)). We speculate that maize, with its higher water demands and lower optimal temperature for grain filling, always benefited from enhanced water availability. Conversely, irrigation in soybean fulfilled its lower water demands and allowed it to exploit the additional sunshine provided by reduced precipitation (Zhang *et al* 2015).

Although being an effective adaptation strategy to increasingly dry and hot climates (figures 2–4), irrigation is in many regions unsustainable to expand or impossible to implement due to water scarcity (Wada *et al* 2012, Rosa *et al* 2020). In the American Midwest, for instance, irrigation relies on groundwater, which is already overexploited (Scanlon *et al* 2012). Irrigation

can also lead to soil salinization and exacerbate pollution of surface- and ground-water via salt mobilization and nutrient leaching (Scanlon *et al* 2007). To avoid these issues, and with expected higher temperatures, more intermittent precipitation and reductions in summer precipitation totals, soil and crop management other than irrigation will be required to minimize the need for migration from exposed crop cultivation areas (Sloat *et al* 2020). Adaptation practices include growing drought- and heat-tolerant varieties and crops (Tack *et al* 2016), altering sowing dates (Lobell *et al* 2014, Mourtzinis *et al* 2019), enhancing soil water retention capacity through, for instance, conservation tillage, cover crops and organic soil amendments (e.g. Lal 2004, Pittelkow *et al* 2015, Kaye and Quemada 2017) and diversifying crop rotations (Bowles *et al* 2020, Marini *et al* 2020).

5. Conclusions

We confirm drastic yield reductions of maize and soybean under combined hot and dry conditions, which are likely to induce heat and water stress in the crops. Both maize and soybean yields declined with increased temperature and decreased precipitation. The interaction terms between temperature and precipitation in the statistical models showed that yield changes from increasing temperature or decreasing precipitation differed depending on climatic conditions, where rainfed yields were reduced more rapidly with warmer temperatures in dry conditions for soybean and wet conditions for maize. Irrigation increased and stabilized yields and alleviated both water deficit and high temperatures.

Data availability statement

No new data were created or analyzed in this study.

Acknowledgments

This research was funded by the Swedish Research Council Vetenskapsrådet Grant 2016-04910 to G V and X L, and Grant 2016-06313 to G V and A S; by the Swedish Research Council for Sustainable Development FORMAS Grant 2018-02872 to G V and R B; and by the Bolin Centre for Climate Research (RA7) to A S.

ORCID iDs

Xiangyu Luan  <https://orcid.org/0000-0002-0244-0670>

Riccardo Bommarco  <https://orcid.org/0000-0001-8888-0476>

Anna Scaini  <https://orcid.org/0000-0002-3527-0241>

Giulia Vico  <https://orcid.org/0000-0002-7849-2653>

References

- Agnolucci P, Rapti C, Alexander P, De Lipsis V, Holland R A, Eigenbrod F and Ekins P 2020 Impacts of rising temperatures and farm management practices on global yields of 18 crops *Nat. Food* **1** 562–71
- Alizadeh M R, Adamowski J, Nikoo M R, AghaKouchak A, Dennison P and Sadegh M 2020 A century of observations reveals increasing likelihood of continental-scale compound dry–hot extremes *Sci. Adv.* **6** aaz4571
- Bowles T M *et al* 2020 Long-term evidence shows that crop-rotation diversification Feng on increases agricultural resilience to adverse growing conditions in North America *One Earth* **2** 284–93
- Breiln K, Mazzoleni M, Di Baldassarre G, Lun D and Vico G 2020 Extreme dry and wet spells face changes in their duration and timing *Environ. Res. Lett.* **15** 074040
- Carter E K, Melkonian J, Steinschneider S and Riha S J 2018 Rainfed maize yield response to management and climate covariability at large spatial scales *Agric. For. Meteorol.* **256–257** 242–52
- Challinor A, Watson J, Lobell D, Howden S M, Smith D R and Chhetri N 2014 A meta-analysis of crop yield under climate change and adaptation *Nat. Clim. Change* **4** 287–91
- Cohen I, Zandalinas S I, Huck C, Fritschi F B and Mittler R 2021 Meta-analysis of drought and heat stress combination impact on crop yield and yield components *Physiol. Plant.* **171** 66–76
- Dietzel R, Liebman M, Ewing R, Helmers M, Horton R, Jarchow M and Archontoulis S 2016 How efficiently do corn- and soybean-based cropping systems use water? A systems modeling analysis *Glob. Change Biol.* **22** 666–81
- Feng S and Hao Z 2020 Quantifying likelihoods of extreme occurrences causing maize yield reduction at the global scale *Sci. Total Environ.* **704** 135250
- Galecki A and Burzykowski T 2013 *Linear Mixed-Effects Models Using R* (Berlin: Springer) pp xxxii + 542
- Hawkins E, Fricker T E, Challinor A J, Ferro C A, Ho C K and Osborne T M 2013 Increasing influence of heat stress on French maize yields from the 1960s to the 2030s *Glob. Change Biol.* **19** 937–47
- Hoffman A L, Kemanian A R and Forest C E 2020 The response of maize, sorghum, and soybean yield to growing-phase climate revealed with machine learning *Environ. Res. Lett.* **15** 094013
- IPCC 2019 Climate change and land: an IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems (available at: www.ipcc.ch/srccl/)
- Kaye J P and Quemada M 2017 Using cover crops to mitigate and adapt to climate change. A review *Agron. Sustain. Dev.* **37** 4
- Kukul M S and Irmak S 2018 Climate-driven crop yield and yield variability and climate change impacts on the US Great Plains agricultural production *Sci. Rep.* **8** 3450
- Lal R 2004 Soil carbon sequestration impacts on global climate change and food security *Science* **11** 1623–7
- Leng G 2017 Recent changes in county-level corn yield variability in the United States from observations and crop models *Sci. Total Environ.* **607** 683–90
- Lesk C, Rowhani P and Ramankutty N 2016 Influence of extreme weather disasters on global crop production *Nature* **529** 84–7
- Li X and Troy T J 2018 Changes in rainfed and irrigated crop yield response to climate in the western US *Environ. Res. Lett.* **13** 064031
- Lobell D B and Field C B 2007 Global scale climate–crop yield relationships and the impacts of recent warming *Environ. Res. Lett.* **2** 014002
- Lobell D B, Hammer G L, McLean G, Messina C, Roberts M J and Schlenker W 2013 The critical role of extreme heat for maize

- production in the United States *Nat. Clim. Change* **3** 497–501
- Lobell D B, Roberts M J, Schlenker W, Braun N, Little B B, Rejesus R M and Hammer G L 2014 Greater sensitivity to drought accompanies maize yield increase in the US Midwest *Science* **344** 516–9
- Lobell D B, Schlenker W and Costa-Roberts J 2011 Climate trends and global crop production since 1980 *Science* **333** 616–20
- Lobell D B, Sibley A and Ortiz-Monasterio J I 2012 Extreme heat effects on wheat senescence in India *Nat. Clim. Change* **2** 186–9
- Luan X and Vico G 2021 Canopy temperature and heat stress are increased by compound high air temperature and water stress and reduced by irrigation—a modeling analysis *Hydrol. Earth Syst. Sci.* **25** 1411–23
- Mahrookashani A, Siebert S, Hugging H and Ewert F 2017 Independent and combined effects of high temperature and drought stress around anthesis on wheat *J. Agron. Crop Sci.* **203** 453–63
- Makowski D, Marajo-Petizon E, Durand J-L and Ben-Ari T 2020 Quantitative synthesis of temperature, CO₂, rainfall, and adaptation effects on global crop yields *Eur. J. Agron.* **115** 126041
- Marini L, St Martin A, Vico A, Baldoni G, Berti A, Blecharczyk A, Malecka-Jankowiak I, Morari F, Sawinska Z and Bommarco R 2020 Crop rotations sustain cereal yields under a changing climate *Environ. Res. Lett.* **15** 124011
- Matiu M, Ankerst D P and Menzel A 2017 Interactions between temperature and drought in global and regional crop yield variability during 1961–2014 *PLoS One* **12** e0178339
- Maurer E P, Wood A, Adam J, Lettenmaier D P and Nijssen B 2002 A long-term hydrologically based dataset of land surface fluxes and states for the conterminous United States *J. Clim.* **15** 3237–51
- Mazdiyasi O and AghaKouchak A 2015 Substantial increase in concurrent droughts and heatwaves in the United States *Proc. Natl Acad. Sci. USA* **112** 11484–9
- Mourtzinis S et al 2015 Climate-induced reduction in US-wide soybean yields underpinned by region- and in-season-specific responses *Nat. Plant* **1** 14026
- Mourtzinis S, Specht J E and Conley S P 2019 Defining optimal soybean sowing dates across the US *Sci. Rep.* **9** 2800
- Neukam D, Ahrends H, Luig A, Manderscheid R and Kage H 2016 Integrating wheat canopy temperatures in crop system models *Agronomy* **6** 1
- Osborne T M and Wheeler T R 2013 Evidence for a climate signal in trends of global crop yield variability over the past 50 years *Environ. Res. Lett.* **8** 024001
- Pittelkow C M, Linquist B A, Lundy M E, Liang X, van Groenigen K J, Lee J, van Gestel N, Six J, Venterea R T and van Kessel C 2015 When does no-till yield more? A global meta-analysis *Field Crops Res.* **183** 156–68
- Porter J R and Semenov M A 2005 Crop responses to climatic variation *Phil. Trans. R. Soc. B* **360** 2021–35
- Prasad P V V, Pisipati S R, Momcilovic I and Ristic Z 2011 Independent and combined effects of high temperature and drought stress during grain filling on plant yield and chloroplast EF-Tu expression in spring wheat *J. Agron. Crop Sci.* **197** 430–41
- Ray D K, Gerber J S, MacDonald G K and West P C 2015 Climate variation explains a third of global crop yield variability *Nat. Commun.* **6** 5989
- Rosa L, Chiarelli D D, Rulli M C, Dell'Angelo J and D'Odorico P 2020 Global agricultural economic water scarcity *Sci. Adv.* **6** eaaz6031
- Rose G, Osborne T, Greatrex H and Wheeler T 2017 Impact of progressive global warming on the global-scale yield of maize and soybean *Clim. Change* **134** 417–28
- Rötter R P, Appiah M, Fichtler E, Kersebaum K C, Trnka M and Hoffmann M P 2018 Linking modelling and experimentation to better capture crop impacts of agroclimatic extremes—a review *Field Crops Res.* **221** 142–56
- Sacks W J, Deryng D, Foley J A and Ramankutty N 2010 Crop planting dates: an analysis of global patterns *Glob. Ecol. Biogeogr.* **19** 607–20
- Scanlon B R, Faunt C C, Longueueve L, Reedy R C, Alley W M, McGuire V L and McMahon P B 2012 Groundwater depletion and sustainability of irrigation in the US High Plains and Central Valley *Proc. Natl Acad. Sci. USA* **109** 9320–5
- Scanlon B R, Jolly I, Sophocleous M and Zhang L 2007 Global impacts of conversions from natural to agricultural ecosystems on water resources: quantity versus quality *Water Resour. Res.* **43** W03437
- Schauberger B et al 2017 Consistent negative response of US crops to high temperatures in observations and crop models *Nat. Commun.* **8** 13931
- Schlenker W and Roberts M J 2009 Nonlinear temperature effects indicate severe damages to US crop yields under climate change *Proc. Natl Acad. Sci. USA* **106** 15594–8
- Siebert S, Ewert F, Rezaei E E, Kage H and Grass R 2014 Impact of heat stress on crop yield—on the importance of considering canopy temperature *Environ. Res. Lett.* **9** 044012
- Siebert S, Webber H, Zhao G and Ewert F 2017 Heat stress is overestimated in climate impact studies for irrigated agriculture *Environ. Res. Lett.* **12** 054023
- Sloat L L, Davis S J, Gerber J S, Moore F C, Ray D K, West P C and Mueller N D 2020 Climate adaptation by crop migration *Nat. Commun.* **11** 1243
- Suyker A E and Verma S B 2009 Evapotranspiration of irrigated and rainfed maize–soybean cropping systems *Agric. For. Meteorol.* **149** 443–52
- Suzuki N, Rivero R M, Shulaev V, Blumwal E and Mittler R 2014 Abiotic and biotic stress combinations *New Phytol.* **203** 32–43
- Tack J, Barkley A and Hendricks N 2017 Irrigation offsets wheat yield reductions from warming temperatures *Environ. Res. Lett.* **12** 114027
- Tack J, Barkley A, Rife T W, Poland J A and Nalley L L 2016 Quantifying variety-specific heat resistance and the potential for adaptation to climate change *Glob. Change Biol.* **22** 2904–12
- Troy T J, Kipgen C and Pal I 2015 The impact of climate extremes and irrigation on US crop yields *Environ. Res. Lett.* **10** 054013
- Urban D W, Sheffield J and Lobell D B 2015 The impacts of future climate and carbon dioxide changes on the average and variability of US maize yields under two emission scenarios *Environ. Res. Lett.* **10** 045003
- Vogel E, Donat M G, Alexander L V, Meinshausen M, Ray D K, Karoly D, Meinshausen N and Frieler K 2019 The effects of climate extremes on global agricultural yields *Environ. Res. Lett.* **14** 054010
- Wada Y, van Beek L P H and Bierkens M F P 2012 Nonsustainable groundwater sustaining irrigation: a global assessment *Water Resour. Res.* **48** W00L06
- Zampieri M, Ceglar A, Dentener F and Toreti A 2017 Wheat yield loss attributable to heat waves, drought and water excess at the global, national and subnational scales *Environ. Res. Lett.* **12** 064008
- Zhang T, Lin X and Sassenrath G F 2015 Current irrigation practices in the central United States reduce drought and extreme heat impacts for maize and soybean, but not for wheat *Sci. Total Environ.* **508** 331–42
- Zhao C et al 2017 Temperature increase reduces global yields of major crops in four independent estimates *Proc. Natl Acad. Sci. USA* **114** 9326–31
- Zhu X, Troy T J and Devineni N 2019 Stochastically modeling the projected impacts of climate change on rainfed and irrigated US crop yields *Environ. Res. Lett.* **14** 074021
- Zipper S C, Qiu J and Kucharik C J 2016 Drought effects on US maize and soybean production: spatiotemporal patterns and historical changes *Environ. Res. Lett.* **11** 094021
- Zscheischler J, Orth R and Seneviratne S I 2017 Bivariate return periods of temperature and precipitation explain a large fraction of European crop yields *Biogeosciences* **14** 3309–20