



Analytical study of the sth-order perturbative corrections to the solution to a one-dimensional harmonic oscillator perturbed by a spatially power-law potential $V_{per}(x) = X$






Author	Tran Duong Anh-Tai, Duc T. Hoang, Thu D. H. Truong, Chinh Dung Nguyen, Le Ngoc Uyen, Do Hung Dung, Nguyen Duy Vy, Vinh N. T. Pham
journal or publication title	AIP Advances
volume	11
number	085310
year	2021-08-06
Publisher	American Institute of Physics
Rights	(C) 2021 The Author(s)
Author's flag	publisher
URL	http://id.nii.ac.jp/1394/00002044/

doi: info:doi/10.1063/5.0059800

Analytical study of the sth-order perturbative corrections to the solution to a one-dimensional harmonic oscillator perturbed by a spatially power-law potential $V_{per}(x) = \lambda x^\alpha$

Cite as: AIP Advances 11, 085310 (2021); <https://doi.org/10.1063/5.0059800>

Submitted: 12 June 2021 . Accepted: 24 July 2021 . Published Online: 06 August 2021

 Tran Duong Anh-Tai, Duc T. Hoang,  Thu D. H. Truong, Chinh Dung Nguyen,  Le Ngoc Uyen, Do Hung Dung,  Nguyen Duy Vy,  Vinh N. T. Pham, et al.



View Online



Export Citation



CrossMark

ARTICLES YOU MAY BE INTERESTED IN

[A simpler graphical solution and an approximate formula for energy eigenvalues in finite square quantum wells](#)

American Journal of Physics **88**, 1019 (2020); <https://doi.org/10.1119/10.0001694>

[Multi-object aerodynamic design optimization using deep reinforcement learning](#)

AIP Advances **11**, 085311 (2021); <https://doi.org/10.1063/5.0058088>

[A short introduction to the Lindblad master equation](#)

AIP Advances **10**, 025106 (2020); <https://doi.org/10.1063/1.5115323>

Call For Papers!

AIP Advances

SPECIAL TOPIC: Advances in
Low Dimensional and 2D Materials

Analytical study of the s th-order perturbative corrections to the solution to a one-dimensional harmonic oscillator perturbed by a spatially power-law potential $V_{per}(x) = \lambda x^\alpha$

Cite as: AIP Advances 11, 085310 (2021); doi: 10.1063/5.0059800

Submitted: 12 June 2021 • Accepted: 24 July 2021 •

Published Online: 6 August 2021









View Online



Export Citation



CrossMark

Tran Duong Anh-Tai,^{1,a)}  Duc T. Hoang,²  Thu D. H. Truong,²  Chinh Dung Nguyen,^{3,4}  Le Ngoc Uyen,⁵ 
Do Hung Dung,⁶  Nguyen Duy Vy,^{7,8,b)}  and Vinh N. T. Pham^{2,c)} 

AFFILIATIONS

¹Quantum Systems Unit, Okinawa Institute of Science and Technology Graduate University, Onna, Okinawa 904-0495, Japan

²Department of Physics, Ho Chi Minh City University of Education, Ho Chi Minh City, Vietnam

³Institute of Fundamental and Applied Sciences, Duy Tan University, 6 Tran Nhat Duat St., District 1, Ho Chi Minh City 700000, Vietnam

⁴Faculty of Natural Sciences, Duy Tan University, 03 Quang Trung St., Hai Chau District, Danang 550000, Vietnam

⁵Department of Engineering Science, The University of Electro-Communications, Chofu, Tokyo 182-8585, Japan

⁶Department of Natural Science, Dong Nai University, Dong Nai, Vietnam

⁷Laboratory of Applied Physics, Advanced Institute of Materials Science, Ton Duc Thang University, Ho Chi Minh City, Vietnam

⁸Faculty of Applied Sciences, Ton Duc Thang University, Ho Chi Minh City, Vietnam

^{a)}Electronic mail: tai.tran@oist.jp

^{b)}Electronic mail: nguyenduyvy@tdtu.edu.vn

^{c)}Author to whom correspondence should be addressed: vinhpnt@hcmue.edu.vn

ABSTRACT

In this work, we present a rigorous mathematical scheme for the derivation of the s th-order perturbative corrections to the solution to a one-dimensional harmonic oscillator perturbed by the potential $V_{per}(x) = \lambda x^\alpha$, where α is a positive integer, using the non-degenerate time-independent perturbation theory. To do so, we derive a generalized formula for the integral $I = \int_{-\infty}^{+\infty} x^\alpha \exp(-x^2) H_n(x) H_m(x) dx$,

where $H_n(x)$ denotes the Hermite polynomial of degree n , using the generating function of orthogonal polynomials. Finally, the analytical results with $\alpha = 3$ and $\alpha = 4$ are discussed in detail and compared with the numerical calculations obtained by the Lagrange-mesh method.

© 2021 Author(s). All article content, except where otherwise noted, is licensed under a Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>). <https://doi.org/10.1063/5.0059800>

I. INTRODUCTION

Approximation methods play a crucial role in quantum mechanics since the number of problems that are exactly solvable is small in comparison to those that must be solved approximately. To our knowledge, the hydrogen atom, harmonic oscillators, and quantum particles in some specific potential wells have exact

solutions,¹⁻⁴ and two cold atoms interacting through a point-like force in a three-dimensional harmonic oscillator potential⁵ can also be solved analytically. Recently, Jafarov *et al.* reported an exact solution to the position-dependent effective mass harmonic oscillator model.⁶ Because of this, approximation methods have been developed early since the dawn of quantum mechanics. One of the essential approximation methods is the perturbation theory

(PT) established by Schrödinger in 1926.⁷ Later, it was immediately used to interpret the LoSurdo–Stark effect of the hydrogen atom by Epstein.⁸ Although the PT is not well convergent at higher-order corrections,^{1–3,9,10} many more efficient approximation methods have been developed to treat quantum-mechanically complex problems,¹¹ it is still a paramount and elementary approximation method in quantum physics. For instance, the PT contributes significantly to quantum optics and quantum field theory, as discussed in Ref. 12. For this reason, the PT is usually presented clearly and discussed in detail in quantum mechanics textbooks for undergraduate students.^{1–3,9}

The one-dimensional harmonic oscillator is not only a rich pedagogical example for approximation theories in quantum mechanics^{13–24} but also an excellent candidate for numerical^{25,26} and analytical or algebraic methods,^{6,27–30} owing to its simple calculation and exact solution. Moreover, the one-dimensional harmonic oscillator potential has also played a key role in studies of ultracold atomic quantum gases for the last two decades.^{5,31–51} Moshinsky and Smirnov⁵² provided a deep review of the role of quantum harmonic oscillators in modern physics. In standard quantum mechanics textbooks,^{1–3} a one-dimensional anharmonic oscillator is usually presented as an application of PT because the solutions can be constructed based on the exact solution to a one-dimensional harmonic oscillator. The two perturbative potentials that are usually considered are $V_{per}(x) = \lambda x$ and $V_{per}(x) = \lambda x^4$. It is common to present the calculation of the first- and second-order corrections of the energy; however, the corrections to the wave function are usually not provided in detail.^{1–3} Moreover, physicists have also studied the generalized case $V_{per}(x) = \lambda x^{2\beta}$, where β is a positive integer. The case $\beta = 2$ has been studied using the WKB method,⁵³ the intermediate Hamiltonian,⁵⁴ the Padé approximation,^{55–57} the Heisenberg matrix mechanics,⁵⁸ and the variational perturbation.⁵⁹ In addition, the cases $\beta = 3$ and $\beta = 4$ were studied in Refs. 60 and 61, and the considered potential of the harmonic oscillator is $V(x) = x^2$ instead of $V(x) = 0.5x^2$. The authors intended to establish efficient methods to solve the problem mathematically, regardless of their physical meaning. Recently, the problem has been extended to sextic (x^6) and decatic (x^{10}) potentials using polynomial solutions^{62,63} and a polynomial perturbative potential.⁶⁴ Interestingly, we realized that in the above-mentioned works, the authors only considered even values of the power of x , while the odd cases have not been studied. It is also interesting to note that the wave function was not considered in the above-mentioned articles. Essentially, the applications of the one-dimensional anisotropic oscillator can be found in chemistry, in which the perturbative potential is used to study the vibration in molecules.^{65–69} In addition, the perturbative potential λx^4 has recently been used to model the Brownian motion of particles in optical tweezers.⁷⁰ Consequently, it is necessary to compute the approximated wave function and the energy of a one-dimensional harmonic oscillator perturbed by the potential $V_{per}(x) = \lambda x^\alpha$ for arbitrary eigenstates with arbitrary values of α .

The goal of this study is to present a systematic and complete treatment of the s th-order perturbative corrections to the solution to a one-dimensional harmonic oscillator perturbed by the potential $V_{per}(x) = \lambda x^\alpha$. To achieve this goal, we derived a formula for $I = \int_{-\infty}^{+\infty} x^\alpha \exp(-x^2) H_n(x) H_m(x) dx$, with $H_n(x)$ being the Hermite polynomial of degree n . Our scheme is based on the so-called

generating functions of orthogonal polynomials.⁷¹ Because the potential depends solely on the spatial coordinate and the states are non-degenerate, the non-degenerate time-independent PT is used to derive the corrections to the wave function and energy. Note that our results can be used for arbitrary eigenstates of a one-dimensional anharmonic oscillator with an arbitrary power coefficient α . This is significantly different from previous works, as discussed above.

The remainder of this paper is organized as follows: Sec. II briefly outlines the time-independent PT for non-degenerate states. Section III presents the main results and discussion. Finally, conclusions are presented in Sec. IV. For simplicity, we use atomic units in which $\hbar = m = \omega = 1$ throughout this study. In addition, the notation $X_{n,s}^\alpha$ denotes the s th-order perturbative correction to the physical quantity X in the state with the quantum number n and the power coefficient α .

II. NON-DEGENERATE TIME-INDEPENDENT PERTURBATION THEORY AND THE $2s + 1$ RULE

This section presents the time-independent PT for non-degenerate states and a recurrence relation to obtain higher-order corrections to the wave function and energy. We followed the procedure of Fernandez.⁹ The Schrödinger equation describing a one-dimensional quantum system is as follows:

$$\hat{H}\psi_n = E_n\psi_n, \quad (1)$$

where \hat{H} is the Hamiltonian operator and E_n is the eigenvalue corresponding to the eigenfunction ψ_n . The Hamiltonian can be split into two parts,

$$\hat{H} = \hat{H}_0 + \lambda\hat{H}', \quad (2)$$

with \hat{H}_0 being the Hamiltonian operator, whose eigenvalues and eigenfunctions are analytically solvable and satisfy the equation

$$\hat{H}_0\psi_{n,0} = E_{n,0}\psi_{n,0}, \quad (3)$$

and \hat{H}' being sufficiently small and considered as a small perturbation with parameter λ . The Taylor formula is used to expand the energy and the eigenfunction of the Hamiltonian \hat{H} as a function of perturbation parameter λ ,

$$E_n = \sum_{s=0}^{\infty} E_{n,s}\lambda^s, \quad \psi_n = \sum_{s=0}^{\infty} \psi_{n,s}\lambda^s, \quad (4)$$

where s is the order of the perturbative correction. Substituting Eq. (4) into Eq. (1), we obtain a recurrence equation expressing the relation between the corrections to the eigenfunction $\psi_{n,s}$ and the energy $E_{n,s}$ as follows:

$$[\hat{H}_0 - E_{n,0}]\psi_{n,s} = \sum_{j=1}^s E_{n,j}\psi_{n,s-j} - \hat{H}'\psi_{n,s-1}. \quad (5)$$

The perturbation correction to the wave function, $\psi_{n,s}$, is then expanded as a linear combination of the eigenfunctions of the non-perturbative Hamiltonian, \hat{H}_0 , as follows:

$$\psi_{n,s} = \sum_m c_{m,s}\psi_{m,0}, \quad (6)$$

where $c_{mn,s} = \langle \psi_{m,0} | \psi_{n,s} \rangle$ is the expanding coefficient. Substituting Eq. (6) into Eq. (5) and then integrating over the whole space after multiplying both sides by $\psi_{m,0}^*$ yield

$$[E_{m,0} - E_{n,0}]c_{mn,s} = \sum_{j=1}^s E_{n,j}c_{mn,s-j} - \sum_k \hat{H}'_{mk}c_{kn,s-1}, \quad (7)$$

where $\hat{H}'_{mk} = \langle \psi_{m,0} | \hat{H}' | \psi_{k,0} \rangle$. For non-degenerate states, $E_{m,0} \neq E_{n,0}$ if and only if $m \neq n$, we can derive the general expression for the correction to the energy by letting $m = n$ in Eq. (7); hence, we obtain

$$E_{n,s} = \langle \psi_{n,0} | \hat{H}' | \psi_{n,s-1} \rangle - \sum_{j=1}^{s-1} E_{n,j}c_{nn,s-j}. \quad (8)$$

The normalization condition $\langle \psi_n | \psi_n \rangle = 1$ results in a constraint on the correction to the wave function,

$$\sum_{j=0}^s \langle \psi_{n,j} | \psi_{n,s-j} \rangle = \delta_{s0}. \quad (9)$$

Finally, in the case of $m \neq n$, we can derive the expanding coefficients for the correction to the wave function as follows:

$$c_{mn,s} = \frac{1}{E_{n,0} - E_{m,0}} \left(\sum_k \hat{H}'_{mk}c_{kn,s-1} - \sum_{j=1}^s E_{n,j}c_{mn,s-j} \right) \quad (10)$$

and

$$c_{nn,1} = 0, c_{nn,s} = -\frac{1}{2} \sum_{j=1}^{s-1} \sum_m c_{mn,j}c_{mn,s-j}, s > 1 \quad (11)$$

for the case of $m = n$. Substituting $s = 1$ into Eq. (8) derives the formula for the first-order correction to the energy, which is the average of the perturbation potential with respect to the eigenfunction $\psi_{n,0}$,

$$E_{n,1} = \langle \psi_{n,0} | \hat{H}' | \psi_{n,0} \rangle, \quad (12)$$

and the expanding coefficient for the first-order correction to the wave function is then derived as follows:

$$c_{mn,1} = \frac{\langle \psi_{m,0} | \hat{H}' | \psi_{n,0} \rangle}{E_{n,0} - E_{m,0}}. \quad (13)$$

Obtaining the second-order correction to the energy is also straightforward. It is obtained as follows:

$$E_{n,2} = \langle \psi_{n,0} | \hat{H}' | \psi_{n,1} \rangle = \sum_{m \neq n} \frac{|\langle \psi_{m,0} | \hat{H}' | \psi_{n,0} \rangle|^2}{E_{n,0} - E_{m,0}}. \quad (14)$$

These results can be found in standard quantum mechanics textbooks.¹⁻³ To derive higher-order corrections to the energy, it is obvious that one can use the recurrence given by Eq. (8). However, there is another way to quickly compute the corrections to the energy. It is called the $2s + 1$ rule, in which, once we know the s order of the correction to the wave function, we are allowed to compute the corrections to the energy up to the $2s + 1$ order. For the detailed derivation of the rule, one should refer to the textbook.⁹ Below, we list the formula for the third-, fourth-, and fifth-order corrections to the energy used for calculations in Sec. III,

$$E_{n,3} = \langle \psi_{n,1} | \hat{H}' - E_{n,1} | \psi_{n,1} \rangle, \quad (15)$$

$$E_{n,4} = \langle \psi_{n,2} | \hat{H}' - E_{n,1} | \psi_{n,1} \rangle - E_{n,2}(\langle \psi_{n,2} | \psi_{n,0} \rangle + \langle \psi_{n,1} | \psi_{n,1} \rangle), \quad (16)$$

$$E_{n,5} = \langle \psi_{n,2} | \hat{H}' - E_{n,1} | \psi_{n,2} \rangle - E_{n,2}(\langle \psi_{n,1} | \psi_{n,2} \rangle + \langle \psi_{n,2} | \psi_{n,1} \rangle). \quad (17)$$

III. RESULTS AND DISCUSSION

A. Derivation of the s th-order perturbative corrections to the solution to a one-dimensional anharmonic oscillator

The Schrödinger equation describing a one-dimensional harmonic oscillator induced by a perturbation potential λx^α is

$$\left(-\frac{1}{2} \frac{d^2}{dx^2} + \frac{1}{2}x^2 + \lambda x^\alpha \right) \psi_n^\alpha(x) = E_n^\alpha \psi_n^\alpha(x), \quad (18)$$

where λ is the strength of the external field, which gives rise to the perturbation, and α is a positive integer. In the absence of the perturbation, Eq. (18) is the well-known equation describing a one-dimensional harmonic oscillator with a wave function

$$\psi_{n,0}^0(x) = A_n \exp\left(-\frac{x^2}{2}\right) H_n(x), \quad (19)$$

where $A_n = \frac{1}{\sqrt{2^n n! \sqrt{\pi}}}$ is the normalization constant, n is the quantum number, and $H_n(x)$ is the Hermite polynomial of degree n , and the energy is given by

$$E_{n,0}^0 = n + \frac{1}{2}. \quad (20)$$

Since Eq. (18) cannot be analytically solvable, the PT is then chosen to approximate the solutions. As discussed above, the first-order correction of the wave function is given by the following equation:

$$\psi_{n,1}^\alpha(x) = \sum_{m \neq n} c_{mn,1} \psi_{m,0}^0(x), \quad (21)$$

where

$$c_{mn,1} = \frac{\langle \psi_{m,0}^0 | x^\alpha | \psi_{n,0}^0 \rangle}{E_{n,0} - E_{m,0}} \quad (22)$$

is the expanding coefficient of the first-order correction to the wave function. Making use of Eq. (A12) (see the Appendix), we obtain the following equation:

$$\begin{aligned} \langle \psi_{m,0}^0 | x^\alpha | \psi_{n,0}^0 \rangle &= \sum_{k=0}^{k \leq \alpha/2} \sum_{\ell=0}^{\alpha-2k} \binom{\alpha}{2k} \binom{\alpha-2k}{\ell} \frac{n!m!2^{n-\ell} A_n A_m}{(m-\alpha+2k+\ell)!} \\ &\times \Gamma(k + \frac{1}{2}) \delta_{m,n+\alpha-2(k+\ell)}, \end{aligned} \quad (23)$$

where $\delta_{m,n}$ denotes the Kronecker delta, satisfying

$$\delta_{m,n} = \begin{cases} 1, & m = n, \\ 0, & m \neq n. \end{cases} \quad (24)$$

By substituting Eq. (23) into Eq. (21), the first-order correction to the wave function for arbitrary states is obtained by

$$\psi_{n,1}^\alpha(x) = \sum_{m \neq n} \sum_{k=0}^{k \leq \alpha/2} \sum_{\ell=0}^{\alpha-2k} \binom{\alpha}{2k} \binom{\alpha-2k}{\ell} \frac{n!m!2^{n-\ell} A_n A_m}{(m-\alpha+2k+\ell)!} \times \frac{\Gamma(k+\frac{1}{2})}{(n-m)} \psi_{m,0}^0(x) \delta_{m,n+\alpha-2(k+\ell)}. \tag{25}$$

The first-order correction to the energy is then computed by the following equation:

$$E_{n,1}^\alpha = \langle \psi_{n,0}^0 | x^\alpha | \psi_{n,0}^0 \rangle. \tag{26}$$

Combining the known wave function of a one-dimensional harmonic oscillator and Eq. (A12), the first-order correction to the energy is obtained by

$$E_{n,1}^\alpha = \sum_{k=0}^{k \leq \alpha/2} \sum_{\ell=0}^{\alpha-2k} \binom{\alpha}{2k} \binom{\alpha-2k}{\ell} \frac{n!2^{-\ell}}{(n-\alpha+2k+\ell)!} \times \frac{\Gamma(k+\frac{1}{2})}{\sqrt{\pi}} \delta_{n,n+\alpha-2(k+\ell)}. \tag{27}$$

It is interesting to note that $E_{n,1}$ is non-zero only if

$$k + \ell = \frac{\alpha}{2}, \tag{28}$$

owing to the mathematical property of the Kronecker delta. Because k and ℓ are integers, the above equation has the solutions for even α solely. This means that in the case of odd α , the first-order correction to energy always equals to zero. Therefore, the anharmonic oscillator does not feel the presence of the external field in this case if

only the first-order approximation is considered. Therefore, it is necessary to compute higher-order corrections to the energy. Regarding the second-order correction of energy, it can be computed as follows:

$$E_{n,2}^\alpha = \langle \psi_{n,0}^0 | x^\alpha | \psi_{n,1}^\alpha \rangle. \tag{29}$$

By substituting Eqs. (19) and (25) into Eq. (29), we obtain the following equation:

$$E_{n,2}^\alpha = \sum_{m \neq n} \sum_{k=0}^{k \leq \alpha/2} \sum_{\ell=0}^{\alpha-2k} \binom{\alpha}{2k} \binom{\alpha-2k}{\ell} \times \frac{n!m!2^{n-\ell} A_n A_m}{(m-\alpha+2k+\ell)!} \frac{\Gamma(k+\frac{1}{2})}{(n-m)} \delta_{m,n+\alpha-2(k+\ell)} \times \int_{-\infty}^{+\infty} \psi_{m,0}^0(x) \psi_{n,0}^0(x) dx. \tag{30}$$

Once again, the integral can be treated by making use of Eq. (A12),

$$\int_{-\infty}^{+\infty} \psi_{m,0}^0(x) \psi_{n,0}^0(x) dx = \sum_{k'=0}^{k' \leq \alpha/2} \sum_{\ell'=0}^{\alpha-2k'} \binom{\alpha}{2k'} \binom{\alpha-2k'}{\ell'} \times \frac{n!m!2^{n-\ell'} A_n A_m}{(m-\alpha+2k'+\ell')!} \Gamma(k'+\frac{1}{2}) \times \delta_{m,n+\alpha-2(k'+\ell')}. \tag{31}$$

Substituting back into Eq. (30), the general second-order correction to the energy is obtained as follows:

$$E_{n,2}^\alpha = \sum_{m \neq n} \sum_{k=0}^{k \leq \alpha/2} \sum_{\ell=0}^{\alpha-2k} \binom{\alpha}{2k} \binom{\alpha-2k}{\ell} \frac{(n!m!)^2 2^{2(n-\ell)} A_n^2 A_m^2}{(m-\alpha+2k+\ell)!^2 (n-m)} \Gamma(k+\frac{1}{2})^2 \delta_{m,n+\alpha-2(k+\ell)}. \tag{32}$$

Using Eq. (15), the third-order correction to the energy for arbitrary states could be derived as follows:

$$E_{n,3}^\alpha = \langle \psi_{n,1}^\alpha | x^\alpha - E_{n,1}^\alpha | \psi_{n,1}^\alpha \rangle = \sum_{m_1 \neq n} \sum_{k_1=0}^{k_1 \leq \alpha/2} \sum_{\ell_1=0}^{\alpha-2k_1} \sum_{m_2 \neq n} \sum_{k_2=0}^{k_2 \leq \alpha/2} \sum_{\ell_2=0}^{\alpha-2k_2} \binom{\alpha}{2k_1} \binom{\alpha-2k_1}{\ell_1} \binom{\alpha}{2k_2} \binom{\alpha-2k_2}{\ell_2} \times \frac{n!m_1! \Gamma(k_1+\frac{1}{2}) \Gamma(k_2+\frac{1}{2})}{(m_1-\alpha+2k_1+\ell_1)! (n-m_1) \pi \sqrt{\pi}} \delta_{m_1, n+\alpha-2(k_1+\ell_1)} \left[\sum_{k'=0}^{k' \leq \alpha/2} \sum_{\ell'=0}^{\alpha-2k'} \binom{\alpha}{2k'} \binom{\alpha-2k'}{\ell'} \frac{m_2! 2^{n-m_2-\ell_1-\ell_2-\ell'} \Gamma(k'+\frac{1}{2})}{(m_2-\alpha+2k_2+\ell_2)! (m_2-\alpha+2k'+\ell')! (n-m_2)} \times \delta_{m_2, n+\alpha-2(k_2+\ell_2)} \delta_{m_2, m_1+\alpha-2(k'+\ell')} - \sum_{k=0}^{k \leq \alpha/2} \sum_{\ell=0}^{\alpha-2k} \binom{\alpha}{2k} \binom{\alpha-2k}{\ell} \frac{n! 2^{n-m_1-\ell_1-\ell_2-\ell} \Gamma(k+\frac{1}{2})}{(m_1-\alpha+2k_2+\ell_2)! (n-\alpha+2k+\ell)! (n-m_1)} \times \delta_{m_1, n+\alpha-2(k_2+\ell_2)} \delta_{n, n+\alpha-2(k+\ell)} \right]. \tag{33}$$

The wave function of a one-dimensional anharmonic oscillator with first-order correction is given by the following equation:

$$\psi_n^\alpha(x) = \psi_{n,0}^\alpha(x) + \lambda \psi_{n,1}^\alpha(x), \tag{34}$$

and the corresponding energy with third-order correction is as follows:

$$E_n^\alpha = E_{n,0}^\alpha + \lambda E_{n,1}^\alpha + \lambda^2 E_{n,2}^\alpha + \lambda^3 E_{n,3}^\alpha. \tag{35}$$

Obviously, the above results can be used to approximate the wave function and the energy for arbitrary states and arbitrary power α .

Since the calculation for the higher-order corrections is more tedious for the general case α , we constrained our calculation for the general power α at this point. In the following, we discuss two particular circumstances in which $\alpha = 4$ and $\alpha = 3$ in detail and then extend the calculation to the second-order correction to the wave function and the fourth- and fifth-order corrections to the energy for these cases.

B. Typical cases: $\alpha = 4$ and $\alpha = 3$

In this section, we discuss two particular circumstances where $\alpha = 4$ and $\alpha = 3$ as two typical examples of PT. In addition, to validate the significance of the higher-order corrections to the energy obtained in the textbook,⁹ we calculated the second-order correction to the wave function for these two cases and then computed the energy up to the fifth-order approximation. The numerical results obtained by the Lagrange-mesh method^{25,26} were then used as the benchmark to validate the applicable range of the analytically approximated results.

First, let us discuss the case where $\alpha = 4$ explicitly. According to Eq. (25), the first-order correction to the wave function is given by

the following equation:

$$\psi_{n,1}^4(x) = \frac{1}{4} \left[\frac{1}{4} \sqrt{(n-3)_4} \psi_{n-4,0}^0(x) - \frac{1}{4} \sqrt{(n+1)_4} \psi_{n+4,0}^0(x) + (2n-1) \sqrt{(n-1)_2} \psi_{n-2,0}^0(x) - (2n+3) \sqrt{(n+1)_2} \psi_{n+2,0}^0(x) \right], \tag{36}$$

where $(a)_n = a(a+1) \cdots (a+n-1)$ is the Pochhammer symbol. The first-, second-, and third-order corrections to the energy are obtained by using Eqs. (27), (32), and (33), respectively, with $\alpha = 4$,

$$E_{n,1}^4 = \frac{3}{4} (2n^2 + 2n + 1), \tag{37}$$

$$E_{n,2}^4 = -\frac{1}{8} (34n^3 + 51n^2 + 59n + 21), \tag{38}$$

$$E_{n,3}^4 = \left(\frac{375}{16} n^4 + \frac{375}{8} n^3 + \frac{177}{2} n^2 + \frac{1041}{16} n + \frac{333}{16} \right). \tag{39}$$

The second-order correction can be derived by the $2s+1$ rule, which was presented in Sec. II. The calculation shows that

$$\begin{aligned} \psi_{n,2}^4(x) = & \frac{1}{512} \sqrt{(n-7)_8} \psi_{n-8,0}^0(x) + \frac{1}{192} (6n-11) \sqrt{(n-5)_6} \psi_{n-6,0}^0(x) + \frac{1}{16} (2n-7)(n-1) \sqrt{(n-3)_4} \psi_{n-4,0}^0(x) \\ & + \frac{1}{2} \sqrt{(n-1)_2} \left(-\frac{1}{16} n^3 - \frac{129}{32} n^2 + \frac{107}{32} n - \frac{33}{16} \right) \psi_{n-2,0}^0(x) + \frac{1}{2} \sqrt{(n+1)_2} \left(-\frac{1}{16} n^3 + \frac{123}{32} n^2 + \frac{359}{32} n + \frac{75}{8} \right) \psi_{n+2,0}^0(x) \\ & + \frac{1}{16} (2n+9)(n+2) \sqrt{(n+1)_4} \psi_{n+4,0}^0(x) + \frac{1}{192} (6n+17) \sqrt{(n+1)_6} \psi_{n+6,0}^0(x) + \frac{1}{512} \sqrt{(n+1)_8} \psi_{n+8,0}^0(x) \\ & - \frac{1}{2} \left(\frac{65}{128} n^4 + \frac{65}{64} n^3 + \frac{487}{128} n^2 + \frac{211}{64} n + \frac{39}{32} \right) \psi_{n,0}^0(x). \end{aligned} \tag{40}$$

Consequently, the fourth- and fifth-order corrections to the energy are obtained, respectively, as follows:

$$E_{n,4}^4 = -\frac{10\,689}{64} n^5 - \frac{53\,445}{128} n^4 - \frac{71\,305}{64} n^3 - \frac{80\,235}{64} n^2 - \frac{111\,697}{128} n - \frac{30\,885}{128}, \tag{41}$$

$$E_{n,5}^4 = \frac{87\,549}{64} n^6 + \frac{262\,647}{64} n^5 + \frac{3\,662\,295}{256} n^4 + \frac{2\,786\,805}{128} n^3 + \frac{3\,090\,693}{128} n^2 + \frac{3\,569\,679}{256} n + \frac{916\,731}{256}. \tag{42}$$

Next, we compare the analytically approximated formulas with numerical calculations to validate the exactness and determine the applicable range of the approximated formulas. For this purpose, the relative deviation is given by the following equation:

$$\sigma = \left| \frac{E_{num} - E_{ana}}{E_{num}} \right|, \tag{43}$$

where E_{num} and E_{ana} are the numerical and analytical results, respectively. As shown in Fig. 1, in the small λ regime ($0 \leq \lambda \leq 0.05$), the approximated formulas match well to the numerical calculation. In this regime, the tenth-order corrected formula has the lowest relative deviation, approximately zero. However, the higher-order formulas diverge rapidly as the perturbative parameter increases. Nevertheless, in the large λ regime, the first-order corrected formula has, in general, a smaller relative deviation compared to others.

Finally, we explicitly present the corrections to the energy and wave function in the case of $\alpha = 3$. The first- and second-order corrections to the wave function of this case are given, respectively, as follows:

$$\begin{aligned} \psi_{n,1}^3(x) = & \frac{\sqrt{2}}{12} \sqrt{(n-2)_3} \psi_{n-3,0}^0(x) - \frac{\sqrt{2}}{12} \sqrt{(n+1)_3} \psi_{n+3,0}^0(x) \\ & + \frac{3\sqrt{2}}{4} n \sqrt{n} \psi_{n-1,0}^0(x) - \frac{3\sqrt{2}}{4} (n+1) \sqrt{n+1} \psi_{n+1,0}^0(x) \end{aligned} \tag{44}$$




FIG. 1. The first row shows the plots of the energy of the ground state and three excited states as a function of perturbative parameter λ with different orders of correction in the case of $\alpha = 4$. Meanwhile, the second row depicts the relative deviation between the analytical formulas and the numerical calculation obtained by the Lagrange-mesh method (black solid line). Note that the tenth-order corrected energy (pink star line) is taken from Ref. 9.

and

$$\begin{aligned} \psi_{n,2}^3(x) = & \frac{1}{144} \sqrt{(n+1)_6} \psi_{n+6,0}^0(x) + \frac{1}{32} \sqrt{(n+1)_4} (4n+7) \psi_{n+4,0}^0(x) \\ & + \frac{1}{16} \sqrt{(n+1)_2} (7n^2 + 33n + 27) \psi_{n+2,0}^0(x) \\ & + \frac{1}{144} \sqrt{(n-5)_6} \psi_{n-6,0}^0(x) \\ & + \frac{1}{32} \sqrt{(n-3)_4} (4n-3) \psi_{n-4,0}^0(x) \\ & + \frac{1}{16} \sqrt{(n-1)_2} (7n^2 - 19n + 1) \psi_{n-2,0}^0(x) \\ & - \frac{1}{2} \left(\frac{41}{18} n^3 + \frac{41}{12} n^2 + \frac{32}{9} n + \frac{29}{24} \right) \psi_{n,0}^0(x). \end{aligned} \quad (45)$$

Straightforwardly, we obtain

$$E_{n,1}^3 = 0, \quad (46)$$

$$E_{n,2}^3 = -\frac{15}{4} n^2 - \frac{15}{4} n - \frac{11}{8}, \quad (47)$$

$$E_{n,3}^3 = 0, \quad (48)$$

$$E_{n,4}^3 = -\frac{705}{16} n^3 - \frac{2115}{32} n^2 - \frac{1635}{32} n - \frac{465}{32}, \quad (49)$$

$$E_{n,5}^3 = 0. \quad (50)$$

Similarly, the analytically approximated formulas were compared to the numerical calculation. Because the odd-order corrections to




FIG. 2. Same as Fig. 1 but for $\alpha = 3$.

the energy are zero, the second- and fourth-order corrections are considered. The results are depicted in Fig. 2. The comparison shows that the second- and fourth-order corrected energies are relatively identical to the numerical results. The deviations of the ground and first-excited state are less than 1%, while those of higher-excited states are larger, ~5%, which is still acceptable.

IV. CONCLUSION

In summary, we have provided a mathematical procedure to derive the wave function and energy for any arbitrary states of a one-dimensional anharmonic oscillator with the general perturbation potential $V_{per}(x) = \lambda x^\alpha$ using the time-independent non-degenerate PT. Subsequently, the explicit results of two particular cases in which $\alpha = 3$ and $\alpha = 4$ are discussed in detail. In addition, the second-order correction to the wave function and up to fifth-order correction to the energy for these cases were computed. The analytical results were then compared with the numerical solutions obtained using the Lagrange-mesh method. The comparison indicates that in the regime in which the perturbation parameter is small, the analytical results agree well with those obtained numerically and then dramatically diverge as the perturbation parameter increases. The higher-order corrections to the energy diverge rapidly as the perturbative parameter increases. In addition, the fifth-order corrected energy in the case of $\alpha = 4$ is the same as that printed in Ref. 9; however, the procedure is more comfortable to approach. The results in the present article are anticipated to be a useful reference.

ACKNOWLEDGMENTS

This work was supported by the Ministry of Education and Training of Vietnam (Grant No. B2021-SPS-02-VL). Mr. Tran Duong Anh-Tai acknowledges support provided by the Okinawa Institute of Science and Technology Graduate University (OIST). The authors are thankful to Mr. Mathias Mikkelsen for introducing the Lagrange-mesh method to them.

APPENDIX: THE INTEGRAL

$$I = \int_{-\infty}^{+\infty} x^\alpha \exp(-x^2) H_n(x) H_m(x) dx$$

In this work, the Hermite polynomials of degrees m and n are expanded by generating function,⁷¹ respectively,

$$g(t, x) = \sum_{n=0}^{+\infty} H_n(x) \frac{t^n}{n!} = \exp(-t^2 + 2tx), \tag{A1}$$

$$f(z, x) = \sum_{m=0}^{+\infty} H_m(x) \frac{z^m}{m!} = \exp(-z^2 + 2zx). \tag{A2}$$

Then, multiplying both sides of the above two equations by $x^\alpha \exp(-x^2)$ and taking the integral from $-\infty$ to $+\infty$, we obtain

$$\begin{aligned} & \sum_{n=0}^{+\infty} \sum_{m=0}^{+\infty} \left[\int_{-\infty}^{+\infty} x^\alpha \exp(-x^2) H_m(x) H_n(x) dx \right] \frac{t^n z^m}{n! m!} \\ & = \exp(2tz) \int_{-\infty}^{+\infty} x^\alpha \exp[-(x - (t + z))^2] dx. \end{aligned} \tag{A3}$$

To simplify the calculation, let us introduce a new variable $y = x - (t + z)$, and then the right-hand side is rewritten as

$$A = \exp(2tz) \int_{-\infty}^{+\infty} [y + (t + z)]^\alpha \exp(-y^2) dy. \tag{A4}$$

The binomial in (A4) is expanded by the binomial formula

$$[y + (t + z)]^\alpha = \sum_{i=0}^{\alpha} \binom{\alpha}{i} (t + z)^{\alpha-i} y^i, \tag{A5}$$

and hence,

$$A = \exp(2tz) \sum_{i=0}^{\alpha} \binom{\alpha}{i} (t + z)^{\alpha-i} \int_{-\infty}^{+\infty} y^i \exp(-y^2) dy. \tag{A6}$$

The integral in Eq. (A6) can be straightforwardly deduced as

$$\int_{-\infty}^{+\infty} x^k \exp(-x^2) dx = \begin{cases} 0 & \text{if } k \text{ is odd,} \\ \Gamma\left(\frac{1+k}{2}\right) & \text{if } k \text{ is even,} \end{cases} \tag{A7}$$

therefore solely the integrals with even coefficients $i = 2k$ are considered

$$A = \exp(2tz) \sum_{k=0}^{k \leq \alpha/2} \binom{\alpha}{2k} \Gamma\left(k + \frac{1}{2}\right) (t + z)^{\alpha-2k}. \tag{A8}$$

It can be seen in (A3) that the value of the integral is the coefficient of $t^n z^m$; thus, we need to find that expanding coefficient. To do so, we expand

$$\exp(2tz) = \sum_{j=0}^{+\infty} \frac{(2tz)^j}{j!} \tag{A9}$$

by the Taylor formula and

$$(t + z)^{\alpha-2k} = \sum_{\ell=0}^{\alpha-2k} \binom{\alpha-2k}{\ell} t^\ell z^{\alpha-2k-\ell} \tag{A10}$$

by the binomial formula. Substituting into (A4), we obtain

$$A = \sum_{j=0}^{+\infty} \left[\sum_{k=0}^{k \leq \alpha/2} \sum_{\ell=0}^{\alpha-2k} \binom{\alpha}{2k} \binom{\alpha-2k}{\ell} \Gamma\left(k + \frac{1}{2}\right) \frac{2^j}{j!} \right] t^{j+\ell} z^{\alpha-2k-\ell+j}. \tag{A11}$$

Combining (A11) and (A3), the desired result is obtained,

$$\begin{aligned} & \int_{-\infty}^{+\infty} x^\alpha \exp(-x^2) H_m(x) H_n(x) dx \\ & = \sum_{k=0}^{k \leq \alpha/2} \sum_{\ell=0}^{\alpha-2k} \binom{\alpha}{2k} \binom{\alpha-2k}{\ell} \frac{n! m! 2^{n-\ell}}{(m - \alpha + 2k + \ell)!} \\ & \quad \times \Gamma\left(k + \frac{1}{2}\right) \delta_{m+n+\alpha-2(k+\ell)}. \end{aligned} \tag{A12}$$

DATA AVAILABILITY

The data that support the findings of this study are available from the corresponding author upon reasonable request.

REFERENCES

- ¹E. M. Lifshitz and L. D. Landau, *Quantum Mechanics: Non-Relativistic Theory* (Pergamon Press, 1965).
- ²H. A. Bethe and E. E. Salpeter, *Quantum Mechanics of One- and Two-Electron Atoms* (Springer, 1957).
- ³D. J. Griffiths, *Introduction to Quantum Mechanics* (Pearson, 2005).
- ⁴S.-H. Dong, *Factorization Method in Quantum Mechanics* (Springer Science & Business Media, 2007), Vol. 150.
- ⁵T. Busch, B.-G. Englert, K. Rzazewski, and M. Wilkens, "Two cold atoms in a harmonic trap," *Fund. Phys.* **28**, 549–559 (1998).
- ⁶E. I. Jafarov, S. M. Nagiyev, R. Oste, and J. Van der Jeugt, "Exact solution of the position-dependent effective mass and angular frequency Schrödinger equation: Harmonic oscillator model with quantized confinement parameter," *J. Phys. A: Math. Theor.* **53**, 485301 (2020).
- ⁷E. Schrödinger, "Quantisierung als eigenwertproblem," *Ann. Phys.* **385**, 437–490 (1926).
- ⁸P. S. Epstein, "The Stark effect from the point of view of Schrödinger's quantum theory," *Phys. Rev.* **28**, 695 (1926).
- ⁹F. M. Fernández, *Introduction to Perturbation Theory in Quantum Mechanics* (CRC Press, 2000).
- ¹⁰G. V. Dunne and M. Ünsal, "Generating nonperturbative physics from perturbation theory," *Phys. Rev. D* **89**, 041701 (2014).
- ¹¹M. Razavy, *Quantum Theory of Tunneling* (World Scientific, 2013).
- ¹²P. Meystre and M. Sargent, *Elements of Quantum Optics* (Springer Science & Business Media, 2007).
- ¹³F. Marsiglio, "The harmonic oscillator in quantum mechanics: A third way," *Am. J. Phys.* **77**, 253–258 (2009).
- ¹⁴M. Moriconi, "Hybrid derivation of the 1D harmonic oscillator propagator," *Am. J. Phys.* **88**, 573–575 (2020).
- ¹⁵M. Rushka and J. K. Freericks, "A completely algebraic solution of the simple harmonic oscillator," *Am. J. Phys.* **88**, 976–985 (2020).
- ¹⁶L. O. Castañón and A. Zuñiga-Segundo, "The forced harmonic oscillator: Coherent states and the RWA," *Am. J. Phys.* **87**, 815–823 (2019).
- ¹⁷M. Bhattacharya, H. Shi, and S. Preble, "Coupled second-quantized oscillators," *Am. J. Phys.* **81**, 267–273 (2013).
- ¹⁸F. D. Mazzitelli, M. D. Mazzitelli, and P. I. Soubelet, "Solutions of the Schrödinger equation for piecewise harmonic potentials: Remarks on the asymptotic behavior of the wave functions," *Am. J. Phys.* **85**, 750–756 (2017).
- ¹⁹R. Hauko and R. Repnik, "Damped harmonic oscillation: Linear or quadratic drag force?," *Am. J. Phys.* **87**, 910–914 (2019).
- ²⁰L. P. Gilbert, M. Belloni, M. A. Doncheski, and R. W. Robinett, "Piecewise zero-curvature energy eigenfunctions in one dimension," *Eur. J. Phys.* **27**, 1331 (2006).
- ²¹S. H. Patil, "Quadrupolar, triple δ -function potential in one dimension," *Eur. J. Phys.* **30**, 629 (2009).
- ²²S. H. Patil, "Harmonic oscillator with a δ -function potential," *Eur. J. Phys.* **27**, 899 (2006).
- ²³J. Viana-Gomes and N. M. R. Peres, "Solution of the quantum harmonic oscillator plus a delta-function potential at the origin: The oddness of its even-parity solutions," *Eur. J. Phys.* **32**, 1377 (2011).
- ²⁴I. Ghose and P. Sen, "The variational method applied to the harmonic oscillator in the presence of a delta function potential," *Eur. J. Phys.* **42**, 045406 (2021).
- ²⁵D. Baye and P.-H. Heenen, "Generalised meshes for quantum mechanical problems," *J. Phys. A: Math. Gen.* **19**, 2041 (1986).
- ²⁶D. Baye, "Lagrange-mesh method for quantum-mechanical problems," *Phys. Status Solidi B* **243**, 1095–1109 (2006).
- ²⁷E. I. Jafarov, N. I. Stoilova, and J. Van der Jeugt, "Finite oscillator models: The Hahn oscillator," *J. Phys. A: Math. Theor.* **44**, 265203 (2011).
- ²⁸E. I. Jafarov, N. I. Stoilova, and J. Van der Jeugt, "The Hahn oscillator and a discrete Fourier–Hahn transform," *J. Phys. A: Math. Theor.* **44**, 355205 (2011).
- ²⁹E. I. Jafarov and J. Van der Jeugt, "The oscillator model for the Lie superalgebra $sh(2|2)$ and Charlier polynomials," *J. Math. Phys.* **54**, 103506 (2013).
- ³⁰E. I. Jafarov and J. Van der Jeugt, "Discrete series representations for $sl(2|1)$, Meixner polynomials and oscillator models," *J. Phys. A: Math. Theor.* **45**, 485201 (2012).
- ³¹M. D. Girardeau, E. M. Wright, and J. M. Triscari, "Ground-state properties of a one-dimensional system of hard-core bosons in a harmonic trap," *Phys. Rev. A* **63**, 033601 (2001).
- ³²J. Goold, M. Krych, Z. Idziaszek, T. Fogarty, and T. Busch, "An eccentrically perturbed Tonks–Girardeau gas," *New J. Phys.* **12**, 093041 (2010).
- ³³J. Goold and T. Busch, "Ground-state properties of a Tonks–Girardeau gas in a split trap," *Phys. Rev. A* **77**, 063601 (2008).
- ³⁴T. Kinoshita, T. Wenger, and D. S. Weiss, "A quantum Newton's cradle," *Nature* **440**, 900–903 (2006).
- ³⁵T. Kinoshita, T. Wenger, and D. S. Weiss, "Observation of a one-dimensional Tonks–Girardeau gas," *Science* **305**, 1125–1128 (2004).
- ³⁶Y. J. Hao and S. Chen, "Ground-state properties of interacting two-component Bose gases in a one-dimensional harmonic trap," *Eur. Phys. J. D* **51**, 261–266 (2009).
- ³⁷L. T. Dat and V. N. T. Pham, "On the derivation of the entropy of ideal quantum gases confined in a three-dimensional harmonic potential," *Commun. Theor. Phys.* **72**, 045701 (2020).
- ³⁸H. H. Huy, T. D. Anh-Tai, T. Yamakoshi, and V. N. T. Pham, "Derivation of thermodynamic quantities of ideal Fermi gas in harmonic trap," *Hue Univ. J. Sci.: Nat. Sci.* **126**, 109–118 (2017).
- ³⁹V. N. T. Pham, T. D. Anh-Tai, H. H. Huy, N. H. Tung, N. D. Vy, and T. Yamakoshi, "A procedure for high-accuracy numerical derivation of the thermodynamic properties of ideal Bose gases," *Eur. J. Phys.* **39**, 055103 (2018).
- ⁴⁰T. Sowiński and M. Á. García-March, "One-dimensional mixtures of several ultracold atoms: A review," *Rep. Prog. Phys.* **82**, 104401 (2019).
- ⁴¹T. Sowiński, M. Brewczyk, M. Gajda, and K. Rzazewski, "Dynamics and decoherence of two cold bosons in a one-dimensional harmonic trap," *Phys. Rev. A* **82**, 053631 (2010).
- ⁴²B. Fang, P. Vignolo, M. Gattobigio, C. Miniatura, and A. Minguzzi, "Exact solution for the degenerate ground-state manifold of a strongly interacting one-dimensional Bose-Fermi mixture," *Phys. Rev. A* **84**, 023626 (2011).
- ⁴³I. Brouzos and P. Schmelcher, "Construction of analytical many-body wave functions for correlated bosons in a harmonic trap," *Phys. Rev. Lett.* **108**, 045301 (2012).
- ⁴⁴T. Bergeman, M. G. Moore, and M. Olshanii, "Atom-atom scattering under cylindrical harmonic confinement: Numerical and analytic studies of the confinement induced resonance," *Phys. Rev. Lett.* **91**, 163201 (2003).
- ⁴⁵A. D. Kerin and A. M. Martin, "Two-body quench dynamics of harmonically trapped interacting particles," *Phys. Rev. A* **102**, 023311 (2020).
- ⁴⁶M. A. García-March, B. Julia-Díaz, G. E. Astrakharchik, T. Busch, J. Boronat, and A. Polls, "Sharp crossover from composite fermionization to phase separation in microscopic mixtures of ultracold bosons," *Phys. Rev. A* **88**, 063604 (2013).
- ⁴⁷M. A. García-March and T. Busch, "Quantum gas mixtures in different correlation regimes," *Phys. Rev. A* **87**, 063633 (2013).
- ⁴⁸M. A. García-March, B. Julia-Díaz, G. E. Astrakharchik, T. Busch, J. Boronat, and A. Polls, "Quantum correlations and spatial localization in one-dimensional ultracold bosonic mixtures," *New J. Phys.* **16**, 103004 (2014).
- ⁴⁹M. Olshanii, "Atomic scattering in the presence of an external confinement and a gas of impenetrable bosons," *Phys. Rev. Lett.* **81**, 938 (1998).
- ⁵⁰M. Á. García-March, T. Fogarty, S. Campbell, T. Busch, and M. Paternostro, "Non-equilibrium thermodynamics of harmonically trapped bosons," *New J. Phys.* **18**, 103035 (2016).
- ⁵¹E. J. Lindgren, J. Rotureau, C. Forssén, A. G. Volosniev, and N. T. Zinner, "Fermionization of two-component few-fermion systems in a one-dimensional harmonic trap," *New J. Phys.* **16**, 063003 (2014).

- ⁵²M. Moshinsky and Y. F. Smirnov, *The Harmonic Oscillator in Modern Physics*, Contemporary Concepts in Physics (CRC Press, 1996).
- ⁵³C. M. Bender and T. T. Wu, "Analytic structure of energy levels in a field-theory model," *Phys. Rev. Lett.* **21**, 406 (1968).
- ⁵⁴N. W. Bazley and D. W. Fox, "Lower bounds for eigenvalues of Schrödinger's equation," *Phys. Rev.* **124**, 483 (1961).
- ⁵⁵J. J. Loeffel, A. Martin, B. Simon, and A. S. Wightman, "Padé approximants and the anharmonic oscillator," *Phys. Lett. B* **30**, 656–658 (1969).
- ⁵⁶B. Simon and A. Dicke, "Coupling constant analyticity for the anharmonic oscillator," *Ann. Phys.* **58**, 76–136 (1970).
- ⁵⁷S. Graffi, V. Grecchi, and B. Simon, "Borel summability: Application to the anharmonic oscillator," *Phys. Lett. B* **32**, 631–634 (1970).
- ⁵⁸R. R. Chasman, "Eigenvalue problems in matrix mechanics," *J. Math. Phys.* **2**, 733–735 (1961).
- ⁵⁹D. G. Truhlar, "Oscillators with quartic anharmonicity: Approximate energy levels," *J. Mol. Spectrosc.* **38**, 415–421 (1971).
- ⁶⁰J. Killingbeck, "The harmonic oscillator with λx^M perturbation," *J. Phys. A: Math. Gen.* **13**, 49 (1980).
- ⁶¹S. N. Biswas, K. Datta, R. P. Saxena, P. K. Srivastava, and V. S. Varma, "Eigenvalues of λx^{2m} anharmonic oscillators," *J. Math. Phys.* **14**, 1190–1195 (1973).
- ⁶²F. Maiz, M. M. Alqahtani, N. Al Sdran, and I. Ghnaim, "Sextic and decatic anharmonic oscillator potentials: Polynomial solutions," *Physica B* **530**, 101–105 (2018).
- ⁶³K. Manimegalai, S. Paul, M. M. Panja, and T. Sil, "Study of the sextic and decatic anharmonic oscillators using an interpolating scale function," *Eur. Phys. J. Plus* **135**, 133 (2020).
- ⁶⁴M. Jafarpour and D. Afshar, "Calculation of energy eigenvalues for the quantum anharmonic oscillator with a polynomial potential," *J. Phys. A: Math. Gen.* **35**, 87 (2001).
- ⁶⁵E. M. Loebel, "Physical chemistry: A series of monographs," in *Experimental Methods in Catalytic Research* (Elsevier, 1968).
- ⁶⁶J. David, R. Lide, A. P. Martin, G. L. Bir, and G. E. Pikus, *Critical Evaluation of Chemical and Physical Structural Information* (The National Academies Press, 1974).
- ⁶⁷J. Laane, "Vibrational potential energy surfaces and conformations of molecules in ground and excited electronic states," *Annu. Rev. Phys. Chem.* **45**, 179–211 (1994).
- ⁶⁸J. Laane, "Spectroscopic determination of ground and excited state vibrational potential energy surfaces," *Int. Rev. Phys. Chem.* **18**, 301–341 (1999).
- ⁶⁹J. Laane, "Experimental determination of vibrational potential energy surfaces and molecular structures in electronic excited states," *J. Phys. Chem. A* **104**, 7715–7733 (2000).
- ⁷⁰B. Suassuna, B. Melo, and T. Guerreiro, "Path integrals and nonlinear optical tweezers," *Phys. Rev. A* **103**, 013110 (2021).
- ⁷¹H. J. Weber and G. B. Arfken, *Essential Mathematical Methods for Physicists* (Elsevier, 2003).