



Prevalence and Risk Factors of Mental Health Symptoms and Suicidal Behavior Among University Students in Wuhan, China During the COVID-19 Pandemic

OPEN ACCESS

Edited by:

Haibo Yang,
Tianjin Normal University, China

Reviewed by:

Zeinab Abbas,
Lebanese International
University, Lebanon
Yun Li,
Shantou University Mental Health
Center, China

*Correspondence:

Lin Lu
linlu@bjmu.edu.cn
Yanping Bao
baoy@bjmu.edu.cn
Guofu Huang
531529128@qq.com

†These authors have contributed
equally to this work and share first
authorship

Specialty section:

This article was submitted to
Public Mental Health,
a section of the journal
Frontiers in Psychiatry

Received: 14 April 2021

Accepted: 17 June 2021

Published: 13 July 2021

Citation:

Xu Y, Su S, Jiang Z, Guo S, Lu Q,
Liu L, Zhao Y, Wu P, Que J, Shi L,
Deng J, Meng S, Yan W, Sun Y,
Yuan K, Lin X, Sun S, Ravindran AV,
Chen S, Wing YK, Tang X, Ran M,
Lu Y, Shi J, Huang G, Bao Y and Lu L
(2021) Prevalence and Risk Factors of
Mental Health Symptoms and Suicidal
Behavior Among University Students
in Wuhan, China During the COVID-19
Pandemic.
Front. Psychiatry 12:695017.
doi: 10.3389/fpsy.2021.695017

Yingying Xu^{1,2†}, Sizhen Su^{3†}, Zhendong Jiang^{4†}, Suihuai Guo⁴, Qingdong Lu^{1,2}, Lin Liu^{1,2}, Yimiao Zhao^{1,2}, Ping Wu¹, Jianyu Que³, Le Shi³, Jiahui Deng³, Shiqiu Meng¹, Wei Yan³, Yankun Sun³, Kai Yuan³, Xiao Lin³, Siwei Sun³, Arun V. Ravindran⁵, Sijing Chen⁶, Yun Kwok Wing⁶, Xiangdong Tang⁷, Maosheng Ran⁸, Yu Lu⁴, Jie Shi¹, Guofu Huang^{4*}, Yanping Bao^{1,2*} and Lin Lu^{3,9*}

¹ National Institute on Drug Dependence and Beijing Key Laboratory of Drug Dependence, Peking University, Beijing, China, ² School of Public Health, Peking University, Beijing, China, ³ Peking University Sixth Hospital, Peking University Institute of Mental Health, NHC Key Laboratory of Mental Health (Peking University), National Clinical Research Center for Mental Disorders (Peking University Sixth Hospital), Beijing, China, ⁴ Wuhan Wuchang Hospital, Wuhan University of Science and Technology, Wuhan, China, ⁵ Department of Psychiatry, University of Toronto, Toronto, ON, Canada, ⁶ Department of Psychiatry, Faculty of Medicine, Chinese University of Hong Kong, Hong Kong SAR, China, ⁷ Sleep Medicine Center, Department of Respiratory and Critical Care Medicine, Mental Health Center and Translational Neuroscience Center, State Key Laboratory of Biotherapy, West China Hospital, Sichuan University, Chengdu, China, ⁸ Department of Social Work and Social Administration, University of Hong Kong, Hong Kong, China, ⁹ Peking-Tsinghua Center for Life Sciences and PKU-IDG/McGovern Institute for Brain Research, Peking University, Beijing, China

Background: University students who are exposed to coronavirus disease 2019 (COVID-19) could be mentally distressed. We aimed to evaluate the pattern and risk factors of mental health and suicidal behavior among students who experienced long-term school closure due to the COVID-19 pandemic.

Methods: This large-sample, cross-sectional, online survey was conducted from June 29, 2020, to July 18, 2020. Eleven thousand two hundred fifty four participants were recruited from 30 universities located in Wuhan, Hubei Province, China. The prevalence of symptoms of depression, anxiety, insomnia, and posttraumatic stress disorder (PTSD) and suicidal behavior was evaluated using the Patient Health Questionnaire-9, Generalized Anxiety Disorder-7, Insomnia Severity Index, Posttraumatic Stress Disorder Checklist for DSM-5, and questions about suicidal ideation and attempts, respectively. Logistic regression was used to explore risk factors for mental health problems and suicidal behavior.

Results: The prevalence of mental health problems was 41.5% for depressive symptoms, 32.6% for anxiety symptoms, 35.0% for insomnia symptoms, 8.5% for PTSD symptoms, and 2.0% for suicidal behavior. Participants with high stress during the pandemic were at higher risk of symptoms of depression [adjusted odds ratio (OR) = 1.67, 95% confidence interval (CI) = 1.43–1.95, $p < 0.01$], anxiety (adjusted OR = 1.90, 95% CI = 1.63–2.23, $p < 0.01$), insomnia (adjusted OR = 1.64, 95%

CI = 1.44–1.87, $p < 0.01$), PTSD (adjusted OR = 1.71, 95% CI = 1.38–2.11, $p < 0.01$) and suicidal behavior (adjusted OR = 3.51, 95% CI = 2.28–5.40, $p < 0.01$). Distant relationship with parents, changes in lifestyle and alcohol use during the pandemic were associated with higher risk of mental health symptoms and suicidal behavior, whereas regular physical exercise reduced the risk of mental health problems.

Conclusions: The psychological symptoms and suicidal behavior were relatively high among students who attended university in Wuhan, China after 6 months of the COVID-19 outbreak in China. Comprehensive mental health services and suicide prevention strategies are essential for university students during the COVID-19 pandemic.

Keywords: university students, COVID-19, prevalence, mental health, suicidal behavior

INTRODUCTION

In December 2019, the coronavirus disease 2019 (COVID-19) outbreak began and was officially announced as a pandemic by the World Health Organization (1). People's lifestyles were profoundly changed by the serious health outcomes of COVID-19, and extremely strict containment measures were taken, including lockdown, quarantine, school closures, and social distancing (2, 3). The unpredictability and uncertainty of the COVID-19 pandemic that are associated with containment strategies and financial loss are among the major stressors that contribute to widespread emotional distress and a higher risk of psychiatric problems in vulnerable populations worldwide (4, 5), including COVID-19-infected cases (6), healthcare workers (7, 8), the elderly (9), and children and adolescents (10).

Mental health status and suicidal behavior among university students that are exacerbated due to long-term mandatory school closures during the pandemic require specific attention. University students might struggle with loneliness and isolation and experience severe psychological distress during the pandemic because of disconnections from friends and partners (11). Students who attended University in an epidemic area during the COVID-19 outbreak might suffer from stigma and discrimination, which were associated with a higher risk of mental health problems (2, 12). Additionally, mental health problem was strongly associated with suicidal ideation and attempts (13, 14). Individuals who had received counseling services on campus could no longer access counseling services, which may exacerbate their mental well-being and increase their risk of substance abuse or even suicidal behavior (11). Suicide is the second leading cause of death among individuals aged 15–29 years globally (15). A meta-analysis showed that the pooled prevalence of lifetime suicidal ideation and attempts among college students was 22.3 and 3.2%, respectively (16).

Several studies reported a potential rise of mental health problems and suicidal behavior among university students during the pandemic (2, 17–19). Major studies focused on a specific mental health problem and had a small sample size (20, 21), evaluated only specific groups (e.g., medical students) (19), or evaluated only one University (18). An exception was the online survey which assessed the prevalence of suicidal ideation, stress, and other mental symptoms among 69,054 university

students in France (22). However, most of these investigations including this survey of France were conducted during early stages of the pandemic (18, 23). To date, many countries are facing substantial threats from the ongoing pandemic and long-term quarantine. University students leaved school and have been changing their normal study habits and lifestyle for a prolonged period of time. Understanding their mental health status and related risk factors is vital for improving mental health, the development of public response strategies, and reopening schools in the future.

To better evaluate the impact of the long-term COVID-19 pandemic on psychological status and suicidal behavior among university students who experienced isolation due to returning home from Wuhan in early days of the pandemic, especially after long-term quarantine and school closure. We conducted a cross-sectional online survey to investigate the prevalence of mental health symptoms and suicidal behavior and potential risk factors among university students 6 months after the COVID-19 pandemic began in China.

METHODS

Study Design

The present study followed the American Association for Public Opinion Research reporting guidelines. Approval from the ethics committee of Peking University Sixth Hospital (Institute of Mental Health) was received before the study began. Informed consent was received online before the respondents began the questionnaire.

This cross-sectional online survey was conducted from June 29, 2020, to July 18, 2020. Based on convenience sampling method, in order to make our sample more representative, the survey involved universities including key universities, ordinary universities and vocational and technical colleges, and the types of subjects of universities were also taken into account. Finally, university students were recruited from 30 universities located in Wuhan, Hubei Province, China (24).

Self-administered questionnaires were sent to students through an online platform and all University classes level were invited to fill the questionnaire. Before the survey began, the details of the survey were given by the class instructor to the class wechat group of students or psychological teacher,

the students take part in the survey voluntarily and all of the participants, who then provided informed consent electronically. The investigation was anonymous, and the confidentiality of all information was ensured.

Study Population

A total of 65,845 students clicked on the survey link, and 11,325 individuals provided the informed consent, for a participation rate of 17.20%. The respondents came from 31 province-level regions in China and attended 30 universities in Wuhan, Hubei Province, China. They returned home from Wuhan for winter vacation when the COVID-19 outbreak began and experienced quarantine in their own home during the pandemic. A total of 11,325 individuals provided informed consent and completed the questionnaires, among whom 71 individuals were excluded because of invalid questionnaire. Participants were excluded if their Body Mass Index (BMI) were out of the range of 13–50, younger than 15 years, or contradictory options about the same questions, for example, their marital status was married while they were younger than 20 years or the type of student they fill in wasn't consistent with that their school enrolled.

Measurements and Covariates

The primary outcome was the prevalence and associated factors of symptoms of depression, anxiety, insomnia and PTSD and suicidal behavior. The survey lasted approximately 10 min. The first part gathered demographic information, including gender, age, nationality, province, city/town, level of education, accommodation in the school, satisfaction with major, academic performance, graduates, peer and teacher relationships, relationship with immediate family, and so on. The second part included questions about the pandemic. The third part focused on the individuals' frequent behaviors before and during the pandemic. The three parts of the survey are included in the **Supplementary Table 1**.

The fourth part of the questionnaire focused on mental health, including a family history of mental illness, questions about suicidal behavior, and validated measurement tools (in Chinese). We used the 9-item Patient Health Questionnaire (PHQ-9) (25), Generalized Anxiety Disorder-7 (GAD-7) (26), Insomnia Severity Index (ISI) (27), and Posttraumatic Stress Disorder Checklist for DSM-5 (PCL-5) (28) to assess symptoms of depression, anxiety, insomnia, and PTSD, respectively. Total scores on these scales were interpreted as the following: PHQ-9 (normal, 0–4; mild depression, 5–9; moderate depression, 10–14; severe depression, 15–27), GAD-7 (normal, 0–4; mild anxiety, 5–9; moderate anxiety, 10–14; severe anxiety, 15–21), ISI (normal, 0–7; subthreshold insomnia, 8–14; moderate insomnia, 15–21; severe insomnia, 22–28), PCL-5 (normal, 0–32; positive for PTSD, 33–80). The cutoff scores for detecting risk factors of symptoms of depression, anxiety, insomnia, and PTSD were 5, 5, 8, and 33, respectively. Suicidal behavior during lifetime included active suicidal ideation and suicide attempts, which were initially screened using a modified inventory (29) in this study. Participants who answered in the affirmative were asked about the occurrence of suicidal ideation and suicide attempts specifically after the COVID-19 outbreak [e.g., “I

thought about killing myself after the COVID-19 outbreak” [Yes/no]; “I deliberately tried to kill myself after the COVID-19 outbreak” (Yes/no)].

Statistical Analysis

Descriptive statistics were used to present demographic data. χ^2 tests were used to compare the prevalence of mild-to-severe mental health symptoms and suicidal ideation and attempts and multiple comparison corrections were conducted for χ^2 tests with more than 2 groups or 2 categories. To explore the factors that were potentially associated with depression, anxiety, insomnia, PTSD symptoms, and suicidal behavior, multiple logistic regression analyses were performed, and odds ratios (ORs) and 95% confidence intervals (CIs) are presented. All variables that were statistically significant in the unadjusted regression analysis were entered into the multivariable model, and then the backward method was applied to determine the variables that were statistically significant in the multivariable analysis. The variance inflation factor (VIF) of these variables > 10 indicates high collinearity (30). All of the data analyses were performed using SPSS 22 software. The level of significance was set to $p < 0.05$, and all tests were two-tailed.

RESULTS

Demographic Characteristics

A total of 11,254 eligible participants from 31 province-level regions in China were included in the final analysis. Of the total sample, 4,054 individuals (36.02%) were male, 8,139 individuals (72.32%) were 15–20 years old, 408 individuals (3.63%) were graduates. More than half of the respondents [6,960 (61.84%)] lived in Hubei province during the pandemic, and 1,402 (12.46%) lived in Wuhan. Of the total number, 6,455 (57.36%) had a harmonious relationship with their classmates and teachers, and most of them had a close relationship with their parents [7,797 (69.28%)]. Furthermore, 76 individuals (0.68%) had confirmed or suspected COVID-19 or were in close contact with confirmed cases, 2,235 individuals (19.86%) were under high stress, and 8,279 individuals (73.56%) experienced changes in lifestyle during the pandemic. A total of 663 individuals (5.89%) reported alcohol use during the pandemic, and 424 individuals (3.77%) reported tobacco use during the pandemic. Additional demographic pandemic-related characteristics are presented in **Table 1**.

Prevalence of Symptoms of Depression, Anxiety, Insomnia, and PTSD and Suicidal Behavior

In this survey, among the 11,254 university students, 5,931 (52.70%) reported at least one symptom of depression, anxiety, insomnia, or PTSD or suicidal behavior. The prevalence of depressive symptoms among the total sample was 41.52% [4,673, including 2,970 (26.39%) with mild depression and 1,703 (15.13%) with moderate-to-severe depression]. The prevalence of anxiety was 32.58% [3,666, including 2,633 (23.40%) with mild anxiety and 1,033 (9.18%) with moderate-to-severe anxiety]. The prevalence of insomnia was 35.00% [3,939, including

TABLE 1 | Descriptive statistics of demographic characteristics and pandemic-related information for students.

Factor	Participants (no. [%])
Overall	11,254 (100.00)
Gender	
Female	7,200 (63.98)
Male	4,054 (36.02)
Age (years)	
15–20	8,139 (72.32)
>20	3,115 (27.68)
Nationality	
Han	10,421 (92.60)
National minority	833 (7.40)
Universities	
Key universities	2,964 (26.34)
Ordinary universities	6,090 (54.11)
Vocational and technical colleges	2,200 (19.55)
Level of education	
Less than college	3,209 (28.51)
College degree or higher	8,005 (71.13)
Postgraduate	40 (0.36)
Living area	
Wuhan	1,402 (12.46)
Hubei Province outside Wuhan	5,558 (49.39)
Other provinces	4,294 (38.16)
Are you graduates?	
Yes	408 (3.63)
No	10,846 (96.37)
Relationship with classmates and teachers	
Strained relationship	41 (0.36)
General relationship	4,758 (42.28)
Harmonious relationship	6,455 (57.36)
Intimacy with parents	
Distant	132 (1.17)
General	3,325 (29.55)
Close	7,797 (69.28)
Positive individual history of mental illness	
Yes	2,863 (25.44)
No	8,391 (74.56)
Positive family history of mental illness	
Yes	377 (3.35)
Unknown	1,608 (14.29)
No	9,269 (82.36)
Are you infected with COVID-19 (confirmed or suspected cases or close contacts)?	
Yes	76 (0.68)
No	11,178 (99.32)
Stress during COVID-19 pandemic	
High	2,235 (19.86)
Moderate	4,025 (35.77)
Low	4,994 (44.38)
Changes in lifestyle during COVID-19 pandemic	
Yes	8,279 (73.56)

(Continued)

TABLE 1 | Continued

Factor	Participants (no. [%])
No	2,975 (26.44)
Alcohol use during COVID-19 pandemic	
Yes	663 (5.89)
No	10,591 (94.11)
Tobacco use during COVID-19 pandemic	
Yes	424 (3.77)
No	10,830 (96.23)
Have you learned about some mental health knowledge during COVID-19 pandemic?	
Yes	6,746 (59.94)
No	4,508 (40.06)
Do you engage in regular physical exercise during COVID-19 pandemic?	
Yes	4,696 (41.73)
No	6,558 (58.27)

3,181 (28.27%) with subthreshold insomnia and 758 (6.74%) with moderate-to-severe insomnia]. The prevalence of PTSD symptoms was 8.46% (952). The prevalence of suicidal behavior was 2.03% [229, including 218 (1.94%) with suicidal ideation and 11 (0.10%) with suicide attempts].

Table 2 shows the severity of mental health symptoms and the presence of suicidal behavior, stratified by demographic characteristics and COVID-19-related factors. Female university students had a significantly lower prevalence of PTSD symptoms (7.50 vs. 10.16%, $p < 0.01$) but a higher prevalence of suicidal behavior compared with males (2.39 vs. 1.41%, $p < 0.01$). The prevalence of mental health symptoms and suicidal behavior was higher among graduates, and students with strained relationships with their classmates/teachers and parents. Individuals who had a prior history of mental illness and positive family history of psychosis had a higher prevalence of mental health symptoms and suicidal behavior. Participants with confirmed or suspected COVID-19 or were in close contact with confirmed cases had a higher prevalence of mental health symptoms and suicidal behavior. The prevalence of mental health problems and suicidal behavior were higher among university students with high stress (depression: 61.83 vs. 26.75%, $p < 0.01$; anxiety: 52.44 vs. 18.46%, $p < 0.01$; insomnia: 52.89 vs. 22.63%, $p < 0.01$; PTSD: 19.37 vs. 3.08%, $p < 0.01$; suicidal behavior: 5.46 vs. 0.58%, $p < 0.01$).

Participants who experienced changes in lifestyle, alcohol use, and tobacco use during the pandemic had a significantly higher prevalence of mental health symptoms and suicidal behavior. Participants who learned about mental health knowledge and exercised regularly during the pandemic had a significantly lower prevalence of mental health symptoms. The detailed information are presented in **Table 2**.

Factors Associated With Symptoms of Depression, Anxiety, Insomnia, and PTSD and Suicidal Behavior

The results of the univariate logistic regression analysis of demographic and COVID-19 pandemic-related variables are

TABLE 2 | Prevalence of symptoms of depression, anxiety, insomnia, and PTSD and suicidal behavior in the students.

Variable	Depression ^a				Anxiety ^b				Insomnia ^c				PTSD ^d			Suicidal behavior	
	Participants (no. [%])				Participants (no. [%])				Participants (no. [%])				Participants (no. [%])			Participants (no. [%])	
	Normal	Mild	Moderate to severe	P-value ^e	Normal	Mild	Moderate to severe	P-value ^e	Normal	Subthreshold	Moderate to severe	P-value ^e	Normal	Positive	p value	p value	
Overall	6,581 (58.48)	2,970 (26.39)	1,703 (15.13)		7,588 (67.42)	2,633 (23.40)	1,033 (9.18)		7,315 (65.00)	3,181 (28.27)	758 (6.74)		10,302 (91.54)	952 (8.46)		229 (2.03)	
Gender																	
Female	4,122 (57.25)	1,978 (27.47)	1,100 (15.28)	< 0.01	4,802 (66.69)	1,745 (24.24)	653 (9.07)	0.03	4,676 (64.94)	2,050 (28.47)	474 (6.58)	0.87	6,660 (92.50)	540 (7.50)	< 0.01	172 (2.39)	< 0.01
Male	2,459 (60.66)	992 (24.47)	603 (14.87)		2,786 (68.72)	888 (21.90)	380 (9.37)		2,639 (65.10)	1,131 (27.90)	284 (7.01)		3,642 (89.84)	412 (10.16)		57 (1.41)	
Living area[#]																	
Wuhan	839 (59.84)	340 (24.25)	223 (15.91)	0.28	956 (68.19)	303 (21.61)	143 (10.20)	0.23	911 (64.98)	385 (27.46)	106 (7.56)	0.42	1,258 (89.73)	144 (10.27)	0.04	41 (2.92)	0.02
Hubei Province outside Wuhan	3,243 (58.35)	1,520 (27.35)	795 (14.30)	0.88	3,779 (67.99)	1,305 (23.48)	474 (8.53)	0.10	3,665 (65.94)	1,552 (27.92)	341 (6.14)	0.03	5,113 (91.99)	445 (8.01)	0.42	106 (1.91)	0.99
Other provinces	2,499 (58.20)	1,110 (25.85)	685 (15.95)		2,853 (66.44)	1,025 (23.87)	416 (9.69)		2,739 (63.79)	1,244 (28.97)	311 (7.24)		3,931 (91.55)	363 (8.45)		82 (1.91)	
Are you graduates?																	
Yes	226 (55.39)	99 (24.26)	83 (20.34)	0.20	263 (64.46)	84 (20.59)	61 (14.95)	0.19	243 (59.56)	124 (30.39)	41 (10.05)	0.02	340 (83.33)	68 (16.67)	< 0.01	17 (4.17)	< 0.01
No	6,355 (58.59)	2,871 (26.47)	1,620 (14.94)		7,325 (67.54)	2,549 (23.50)	972 (8.96)		7,072 (65.20)	3,057 (28.19)	717 (6.61)		9,962 (91.85)	884 (8.15)		212 (1.95)	
Relationship with classmates and teachers[#]																	
Strained relationship	12 (29.27)	9 (21.95)	20 (48.78)	< 0.01	18 (43.90)	11 (26.83)	12 (29.27)	< 0.01	12 (29.27)	16 (39.02)	13 (31.71)	< 0.01	27 (65.85)	14 (34.15)	< 0.01	6 (14.63)	< 0.01
General relationship	2,304 (48.42)	1,482 (31.15)	972 (20.43)	< 0.01	2,804 (58.93)	1,383 (29.07)	571 (12.00)	< 0.01	2,733 (57.44)	1,613 (33.90)	412 (8.66)	< 0.01	4,230 (88.90)	528 (11.10)	< 0.01	126 (2.65)	< 0.01
Harmonious relationship	4,265 (66.07)	1,479 (22.91)	711 (11.01)		4,766 (73.83)	1,239 (19.19)	450 (6.97)		4,570 (70.80)	1,552 (24.04)	333 (5.16)		6,045 (93.65)	410 (6.35)		97 (1.50)	
Intimacy with parents[#]																	
Distant	37 (28.03)	36 (27.27)	59 (44.70)	< 0.01	48 (36.36)	41 (31.06)	43 (32.58)	< 0.01	46 (34.85)	52 (39.39)	34 (25.76)	< 0.01	91 (68.94)	41 (31.06)	< 0.01	17 (12.88)	< 0.01
General	1,461 (43.94)	1,041 (31.31)	823 (24.75)	< 0.01	1,813 (54.53)	1,025 (30.83)	487 (14.65)	< 0.01	1,780 (53.53)	1,208 (36.33)	337 (10.14)	< 0.01	2,857 (85.92)	468 (14.08)	< 0.01	128 (3.85)	< 0.01
Close	5,083 (65.19)	1,893 (24.28)	821 (10.53)		5,727 (73.45)	1,567 (20.10)	503 (6.45)		5,489 (70.40)	1,921 (24.64)	387 (4.96)		7,354 (94.32)	443 (5.68)		84 (1.08)	

(Continued)

TABLE 2 | Continued

Variable	Depression ^a				Anxiety ^b				Insomnia ^c				PTSD ^d			Suicidal behavior	
	Participants (no. [%])				Participants (no. [%])				Participants (no. [%])				Participants (no. [%])			Participants (no. [%])	
	Normal	Mild	Moderate to severe	P-value ^e	Normal	Mild	Moderate to severe	P-value ^e	Normal	Subthreshold	Moderate to severe	P-value ^e	Normal	Positive	p value	p value	
Positive individual history of mental illness																	
Yes	915 (31.96)	1,033 (36.08)	915 (31.96)	< 0.01	1,229 (42.93)	1,017 (35.52)	617 (21.55)	< 0.01	1,130 (39.47)	1,248 (43.59)	485 (16.94)	< 0.01	2,316 (80.89)	547 (19.11)	< 0.01	143 (4.99)	< 0.01
No	5,666 (67.52)	1,937 (23.08)	788 (9.39)		6,359 (75.78)	1,616 (19.26)	416 (4.96)		6,185 (73.71)	1,933 (23.04)	273 (3.25)		7,986 (95.17)	405 (4.83)		86 (1.02)	
Positive family history of mental illness[#]																	
Yes	155 (41.11)	117 (31.03)	105 (27.85)	< 0.01	180 (47.75)	124 (32.89)	73 (19.36)	< 0.01	196 (51.99)	130 (34.48)	51 (13.53)	< 0.01	296 (78.51)	81 (21.49)	< 0.01	17 (4.51)	< 0.01
Unknown	663 (41.23)	541 (33.64)	404 (25.12)	< 0.01	848 (52.74)	491 (30.53)	269 (16.73)	< 0.01	866 (53.86)	565 (35.14)	177 (11.01)	< 0.01	1,367 (85.01)	241 (14.99)	< 0.01	57 (3.54)	< 0.01
No	5,763 (62.17)	2,312 (24.94)	1,194 (12.88)		6,560 (70.77)	2,018 (21.77)	691 (7.45)		6,253 (67.46)	2,486 (26.82)	530 (5.72)		8,639 (93.20)	630 (6.80)		155 (1.67)	
Are you infected with COVID-19 (confirmed or suspected cases or close contacts)?																	
Yes	21 (27.63)	18 (23.68)	37 (48.68)	< 0.01	25 (32.89)	24 (31.58)	27 (35.53)	< 0.01	26 (34.21)	27 (35.53)	23 (30.26)	< 0.01	48 (63.16)	28 (36.84)	< 0.01	5 (6.58)	0.02
No	6,560 (58.69)	2,952 (26.41)	1,666 (14.90)		7,563 (67.66)	2,609 (23.34)	1,006 (9.00)		7,289 (65.21)	3,154 (28.22)	735 (6.58)		10,254 (91.73)	924 (8.27)		224 (2.00)	
Stress during COVID-19 pandemic[#]																	
High	853 (38.17)	702 (31.41)	680 (30.43)	< 0.01	1,063 (47.56)	691 (30.92)	481 (21.52)	< 0.01	1,053 (47.11)	848 (37.94)	334 (14.94)	< 0.01	1,802 (80.63)	433 (19.37)	< 0.01	122 (5.46)	< 0.01
Moderate	2,070 (51.43)	1,311 (32.57)	644 (16.00)	< 0.01	2,453 (60.94)	1,224 (30.41)	348 (8.65)	< 0.01	2,398 (59.58)	1,384 (34.39)	243 (6.04)	< 0.01	3,696 (91.83)	329 (8.17)	< 0.01	78 (1.94)	< 0.01
Low	3,658 (73.25)	957 (19.16)	379 (7.59)		4,072 (81.54)	718 (14.38)	204 (4.08)		3,864 (77.37)	949 (19.00)	181 (3.62)		4,804 (96.20)	190 (3.80)		29 (0.58)	
Changes in lifestyle during COVID-19 pandemic?																	
Yes	4,409 (53.26)	2,433 (29.39)	1,437 (17.36)	< 0.01	5,224 (63.10)	2,185 (26.39)	870 (10.51)	< 0.01	4,988 (60.25)	2,631 (31.78)	660 (7.97)	< 0.01	7,477 (90.31)	802 (9.69)	< 0.01	200 (2.42)	< 0.01
No	2,172 (73.01)	537 (18.05)	266 (8.94)		2,364 (79.46)	448 (15.06)	163 (5.48)		2,327 (78.22)	550 (18.49)	98 (3.29)		2,825 (94.96)	150 (5.04)		29 (0.97)	
Alcohol use during COVID-19 pandemic																	
Yes	306 (46.15)	190 (28.66)	167 (25.19)	< 0.01	380 (57.32)	161 (24.28)	122 (18.40)	< 0.01	355 (53.54)	216 (32.58)	92 (13.88)	< 0.01	553 (83.41)	110 (16.59)	< 0.01	35 (5.28)	< 0.01
No	6,275 (59.25)	2,780 (26.25)	1,536 (14.50)		7,208 (68.06)	2,472 (23.34)	911 (8.60)		6,960 (65.72)	2,965 (28.00)	666 (6.29)		9,749 (92.05)	842 (7.95)		194 (1.83)	

(Continued)

TABLE 2 | Continued

Variable	Depression ^a				Anxiety ^b				Insomnia ^c				PTSD ^d			Suicidal behavior	
	Participants (no. [%])				Participants (no. [%])				Participants (no. [%])				Participants (no. [%])			Participants (no. [%])	
	Normal	Mild	Moderate to severe	P-value ^e	Normal	Mild	Moderate to severe	P-value ^e	Normal	Subthreshold	Moderate to severe	P-value ^e	Normal	Positive	p value	p value	
Tobacco use during COVID-19 pandemic																	
Yes	206 (48.58)	111 (26.18)	107 (25.24)	< 0.01	244 (57.55)	99 (23.35)	81 (19.10)	< 0.01	241 (56.84)	124 (29.25)	59 (13.92)	< 0.01	349 (82.31)	75 (17.69)	< 0.01	16 (3.77)	0.01
No	6,375 (58.86)	2,859 (26.40)	1,596 (14.74)		7,344 (67.81)	2,534 (23.40)	952 (8.79)		7,074 (65.32)	3,057 (28.23)	699 (6.45)		9,953 (91.90)	877 (8.10)		213 (1.97)	
Have you learned about some mental health knowledge during COVID-19 pandemic?																	
Yes	4,006 (59.38)	1,802 (26.71)	938 (13.90)	0.02	4,598 (68.16)	1,602 (23.75)	546 (8.09)	0.04	4,439 (65.80)	1,883 (27.91)	424 (6.29)	0.03	6,229 (92.34)	517 (7.66)	< 0.01	99 (2.20)	0.32
No	2,575 (57.12)	1,168 (25.91)	765 (16.97)		2,990 (66.33)	1,031 (22.87)	487 (10.80)		2,876 (63.80)	1,298 (28.79)	334 (7.41)		4,073 (90.35)	435 (9.65)		130 (1.93)	
Do you engage in regular physical exercise during COVID-19 pandemic?																	
Yes	3,018 (64.27)	1,130 (24.06)	548 (11.67)	< 0.01	3,380 (71.98)	970 (20.66)	346 (7.37)	< 0.01	3,299 (70.25)	1,157 (24.64)	240 (5.11)	< 0.01	4,365 (92.95)	331 (7.05)	< 0.01	84 (1.79)	0.12
No	3,563 (54.33)	1,840 (28.06)	1,155 (17.61)		4,208 (64.17)	1,663 (25.36)	687 (10.48)		4,016 (61.24)	2,024 (30.86)	518 (7.90)		5,937 (90.53)	621 (9.47)		145 (2.21)	
Symptoms of depression																	
Normal	—	—	—		6,219 (94.50)	346 (5.26)	16 (0.24)	< 0.01	5,597 (85.05)	934 (14.19)	50 (0.76)	< 0.01	6,550 (99.53)	31 (0.47)	< 0.01	28 (0.43)	< 0.01
Positive	—	—	—		1,369 (29.30)	2,287 (48.94)	1,017 (21.76)		1,718 (36.76)	2,247 (48.08)	708 (15.15)		3,752 (80.29)	921 (19.71)		201 (4.30)	
Symptoms of anxiety																	
Normal	6,219 (81.96)	1,190 (15.68)	179 (2.36)	< 0.01	—	—	—		6,103 (80.43)	1,358 (17.90)	127 (1.67)	< 0.01	7,549 (99.49)	39 (0.51)	< 0.01	46 (0.61)	< 0.01
Positive	362 (9.87)	1,780 (48.55)	1,524 (41.57)		—	—	—		1,212 (33.06)	1,823 (49.73)	631 (17.21)		2,753 (75.10)	913 (24.90)		183 (4.99)	
Symptoms of insomnia																	
Normal	5,597 (76.51)	1,431 (19.56)	287 (3.92)	< 0.01	6,103 (83.43)	1,063 (14.53)	149 (2.04)	< 0.01	—	—	—		7,207 (98.52)	108 (1.48)	< 0.01	63 (0.86)	< 0.01
Positive	984 (24.98)	1,539 (39.07)	1,416 (35.95)		1,485 (37.70)	1,570 (39.86)	884 (22.44)		—	—	—		3,095 (78.57)	844 (21.43)		166 (4.21)	
Symptoms of PTSD																	
Normal	6,550 (63.58)	2,809 (27.27)	943 (9.15)	< 0.01	7,549 (73.28)	2,356 (22.87)	397 (3.85)	< 0.01	7,207 (69.96)	2,690 (26.11)	405 (3.93)	< 0.01	—	—		126 (1.22)	< 0.01
Positive	31 (3.26)	161 (16.91)	760 (79.83)		39 (4.10)	277 (29.10)	636 (66.81)		108 (11.34)	491 (51.58)	353 (37.08)		—	—		103 (10.82)	

(Continued)

TABLE 2 | Continued

Variable	Depression ^a			Anxiety ^b			Insomnia ^c			PTSD ^d		Suicidal behavior	
	Normal	Mild	Moderate to severe	Normal	Mild	Moderate to severe	Normal	Subthreshold	Moderate to severe	Normal	Positive	Participants (no. [%])	<i>p</i> value
Suicidal behavior													
Normal	6,553 (59.44)	2,899 (26.29)	1,573 (14.27)	7,542 (68.41)	2,532 (22.97)	951 (8.63)	7,252 (65.78)	3,085 (27.98)	688 (6.24)	10,176 (92.30)	849 (7.70)	Participants (no. [%])	< 0.01
Positive	28 (12.23)	71 (31.00)	130 (56.77)	46 (20.09)	101 (44.10)	82 (35.81)	63 (27.51)	96 (41.92)	70 (30.57)	126 (55.02)	103 (44.98)	Participants (no. [%])	< 0.01

^aScores of 5 to 9 on the PHQ-9 were defined as mild depression, and scores of 10 or higher were defined as moderate-to-severe depression.

^bScores of 5 to 9 on the GAD-7 were defined as mild anxiety, and scores of 10 or higher were defined as moderate-to-severe anxiety.

^cScores of 8 to 14 on the ISI were defined as subthreshold insomnia, and scores of 15 or higher were defined as moderate-to-severe insomnia.

^dScores of 33 or higher on the PCL-5 were defined as PTSD symptoms.

^eχ² tests were used to compare the prevalence of mild-to-severe mental health symptoms in different populations.

This means the first two groups are compared to the last group respectively.

presented in the **Supplementary Table 2**. In the multivariable analysis, females were associated with a lower risk of PTSD (adjusted OR = 0.61, 95% CI = 0.51–0.72, *p* < 0.01) but a higher risk of suicidal behavior (adjusted OR = 1.93, 95% CI = 1.39–2.68, *p* < 0.01) compared with males. Participants who were under high stress had a higher risk of mental health symptoms and suicidal behavior (depression: adjusted OR = 1.67, 95% CI = 1.43–1.95, *p* < 0.01; anxiety: adjusted OR = 1.90, 95% CI = 1.63–2.23, *p* < 0.01; insomnia: adjusted OR = 1.64, 95% CI = 1.44–1.87, *p* < 0.01; PTSD: adjusted OR = 1.71, 95% CI = 1.38–2.11, *p* < 0.01; suicidal behavior: adjusted OR = 3.51, 95% CI = 2.28–5.40, *p* < 0.01) during the pandemic than participants who were under low stress. Participants who had symptoms of depression were at higher risk of other mental problems (anxiety: adjusted OR = 19.98, 95% CI = 17.50–22.82, *p* < 0.01; insomnia: adjusted OR = 3.73, 95% CI = 3.31–4.19, *p* < 0.01; PTSD: adjusted OR = 3.91, 95% CI = 2.60–5.88, *p* < 0.01) and suicidal behavior (adjusted OR = 2.58, 95% CI = 1.54–4.33, *p* < 0.01). Students who had symptoms of anxiety were at higher risk of other mental problems (depression: adjusted OR = 19.89, 95% CI = 17.41–22.72, *p* < 0.01; insomnia: adjusted OR = 1.94, 95% CI = 1.72–2.19, *p* < 0.01; PTSD: adjusted OR = 13.26, 95% CI = 9.27–18.97, *p* < 0.01) and suicidal behavior (adjusted OR = 1.66, 95% CI = 1.07–2.59, *p* = 0.02). Students with symptoms of insomnia were at higher risk of other mental problems (depression: adjusted OR = 3.68, 95% CI = 3.26–4.14, *p* < 0.01; anxiety: adjusted OR = 1.98, 95% CI = 1.75–2.23, *p* < 0.01; PTSD: adjusted OR = 4.43, 95% CI = 3.54–5.54, *p* < 0.01). Participants who had symptoms of PTSD were at higher risk of other mental problems (depression: adjusted OR = 3.75, 95% CI = 2.47–5.69, *p* < 0.01; anxiety: adjusted OR = 13.45, 95% CI = 9.37–19.30, *p* < 0.01; insomnia: adjusted OR = 4.36, 95% CI = 3.50–5.45, *p* < 0.01) and suicidal behavior (adjusted OR = 2.71, 95% CI = 1.98–3.71, *p* < 0.01). Participants with suicidal behavior were at higher risk of mental problems (depression: adjusted OR = 2.00, 95% CI = 1.17–3.43, *p* = 0.01; anxiety: adjusted OR = 1.56, 95% CI = 1.01–2.41, *p* = 0.04; PTSD: adjusted OR = 2.84, 95% CI = 2.05–3.93, *p* < 0.01). Students with a prior history of mental illness had a higher risk than students without a prior history of mental illness (depression: adjusted OR = 1.81, 95% CI = 1.59–2.07, *p* < 0.01; anxiety: adjusted OR = 1.52, 95% CI = 1.33–1.73, *p* < 0.01; insomnia: adjusted OR = 2.20, 95% CI = 1.98–2.44, *p* < 0.01; PTSD: adjusted OR = 1.53, 95% CI = 1.30–1.81, *p* < 0.01; suicidal behavior: adjusted OR = 1.94, 95% CI = 1.45–2.61, *p* < 0.01). Participants with a positive family history of psychosis had a significantly higher risk of mental health problems than participants without a family history of psychosis (anxiety: adjusted OR = 1.42, 95% CI = 1.04–1.95, *p* = 0.03; PTSD: adjusted OR = 1.89, 95% CI = 1.36–2.64, *p* < 0.01). Participants who engaged in regular exercise during the pandemic reported a lower risk of symptoms of depression (adjusted OR = 0.86, 95% CI = 0.77–0.97, *p* = 0.01), and insomnia (adjusted OR = 0.82, 95% CI = 0.75–0.91, *p* < 0.01). Participants who experienced changes in lifestyle and alcohol use during the pandemic also had a significantly higher risk of mental health problems and suicidal behavior. Since symptoms of depression, anxiety, insomnia, PTSD and suicidal behavior are possibly

correlated, we examined the collinearity in five final models and the VIF indicated that no collinearity existed. The highest VIF was 1.51 (symptoms of anxiety) for depression symptoms, 1.52 (symptoms of depression) for anxiety symptoms, 2.07 (symptoms of anxiety) for insomnia symptoms, 2.10 (symptoms of depression) for PTSD symptoms, 2.07 (symptoms of anxiety) for suicidal behavior in each final model. The detailed results of the multivariable analysis are shown in **Table 3**.

DISCUSSION

The present study investigated the prevalence and factors associated with mental health symptoms and suicidal behavior among university students after long-term quarantine and school closures during the COVID-19 pandemic in China. All of the students in this study had a high risk of COVID-19 exposure because they went home for winter vacation from Wuhan, Hubei Province, where COVID-19 was first identified in China. Compared to other university students in China, they experienced quarantine after returning home and might suffer from stigma and discrimination. Overall, more than half of the participants (52.7%) reported at least one symptom of depression, anxiety, insomnia, or PTSD or suicidal behavior. We identified several vulnerable students and risk factors, including graduates, distant relationships with parents, a past history of mental illness, a positive family history of psychosis, moderate to high stress, changes in lifestyle, and alcohol use during the pandemic. We also found that regular exercise during the pandemic was protective against mental health problems. Altogether, our findings highlight concerns about mental well-being among university students and may contribute to the development of better health policies and population-based long-term mental health management and intervention strategies for future pandemics and other public health emergencies.

The prevalence of anxiety (32.6%) in the present study was higher than the results of a previous epidemiological study (19) among medical college students during the pandemic, showing that 24.9% of participants had symptoms of anxiety. This difference may be attributable to long-term quarantine experience in the present survey and the fact that medical students are a special group who may more easily acquire mental health knowledge. Additionally, with ongoing worries about current academic studies, future employment, personnel relationships, and life stress, university students may be vulnerable to the development of mental and sleep problems (2, 11). The prevalence of symptoms of depression among students in this study was higher than in another online study (31) that was performed 1 month after the COVID-19 outbreak began among undergraduate students, which may be attributable to the impact of long-term quarantine during the pandemic.

However, the prevalence of suicidal ideation (1.9%) in the present study was lower than in a study in which 18.0% of a sample of United States college students reported suicidal thoughts, measured by the PHQ-9 (18). Some previous studies

showed that responses to items on the PHQ-9 that included self-harm or passive thoughts of death may not accurately reflect suicidal thoughts compared with questions that are designed specifically to assess suicidality, thus resulting in a higher rate of false positives (32–34). Moreover, the prevalence of suicidal ideation in the present study was lower than in a previous study that was conducted at the peak of the COVID-19 pandemic, which showed that 11.4% of university students reported suicidal thoughts (22). Our survey focused on the late sustained pandemic period, which is another possible reason for the lower prevalence of suicidal ideation in the present study (35). A previous study showed that suicidality was associated with fear of the infection and the experience of social isolation, which were more serious at the peak of the pandemic than at the late pandemic period (2, 35). Further studies with a longer follow-up period after the COVID-19 pandemic and large geographic coverage are warranted.

The present study identified several risk groups who were more likely to develop mental health problems and suicidal behavior. Females had a significantly higher prevalence of suicidal behavior, which is consistent with the results of another study of senior high school students during the COVID-19 pandemic (36). The present results also showed that strained relationships with classmates/teachers and parents increased the risk of mental health problems. For most students, isolation from social networks was associated with more depressive symptoms (37). Conflicts and tension between family members, which may have increased because of staying at home during the pandemic (38), may descend into domestic violence and lead to a worsening of mental health problems and suicidal behavior (39–41). An interesting finding is that we did not find difference of psychological problems between students from Wuhan/Hubei and that from other provinces in final model. The reason may be that all students in our study attended universities in Wuhan and the finding may also indicated that the mental health well-being among university students are needed in whole country, not in special high risk area of COVID-19 pandemic.

Participants who had a prior history of mental health problems reported more severe mental health symptoms and suicidal behavior. A positive history of a psychiatric disorder is highly associated with new-onset mental disorders during a stressful period (42). Additionally, the present study also found that participants with a positive family history of psychosis were more likely to suffer from anxiety, and PTSD. Furthermore, this study showed the issue of coexistent mental health symptoms and suicidal behavior during the pandemic which is consistent with other epidemiologic studies (43–45). Another cross-sectional survey of college students found a strong positive interaction effect between anxiety disorder and depression (44). A large cohort study also showed that 40% individuals who had mental disorders were diagnosed with more than one disorder type (46). Our findings indicated the importance of early detection and intervention of mental health symptoms and suicidal behavior among high risk population. It is necessary to establish mental health promotion strategies, such as health education, early response to public traumatic events to further prevent mental

TABLE 3 | Multivariable regression analysis of risk factors associated with mental health symptoms and suicidal behavior.

Variable	Depression ^a		Anxiety ^b		Insomnia ^c		PTSD ^d		Suicidal behavior	
	AOR (95% CI)	P-value	AOR (95% CI)	P-value	AOR (95% CI)	P-value	AOR (95% CI)	P-value	AOR (95% CI)	P-value
Gender (ref: male)										
Female	1.13 (1.00–1.27)	0.05	—	—	—	—	0.61 (0.51–0.72)	< 0.01	1.93 (1.39–2.68)	< 0.01
Are you graduates? (ref: no)										
Yes	—	—	—	—	—	—	2.04 (1.42–2.93)	< 0.01	1.67 (0.96–2.91)	0.07
Relationship with classmates and teachers (ref: harmonious)										
Strained	1.73 (0.64–4.66)	0.28	0.68 (0.26–1.80)	0.44	2.74 (1.16–6.47)	0.02	—	—	—	—
General	1.38 (1.23–1.54)	< 0.01	1.20 (1.07–1.35)	< 0.01	1.17 (1.06–1.29)	< 0.01	—	—	—	—
Intimacy with parents (ref: close)										
Distant	1.47 (0.82–2.63)	0.20	1.38 (0.81–2.36)	0.24	1.63 (1.04–2.57)	0.03	2.21 (1.36–3.58)	< 0.01	4.71 (2.57–8.64)	< 0.01
General	1.31 (1.15–1.48)	< 0.01	1.16 (1.03–1.32)	0.02	1.17 (1.06–1.30)	< 0.01	1.49 (1.26–1.75)	< 0.01	2.00 (1.49–2.68)	< 0.01
Positive individual history of mental illness (ref: no)										
Yes	1.81 (1.59–2.07)	< 0.01	1.52 (1.33–1.73)	< 0.01	2.20 (1.98–2.44)	< 0.01	1.53 (1.30–1.81)	< 0.01	1.94 (1.45–2.61)	< 0.01
Positive family history of mental illness (ref: no)										
Yes	1.02 (0.74–1.41)	0.88	1.42 (1.04–1.95)	0.03	—	—	1.89 (1.36–2.64)	< 0.01	—	—
Unknown	1.61 (1.37–1.89)	< 0.01	1.20 (1.02–1.40)	0.02	—	—	1.38 (1.14–1.66)	< 0.01	—	—
Stress during COVID-19 epidemic (ref: low)										
High	1.67 (1.43–1.95)	< 0.01	1.90 (1.63–2.23)	< 0.01	1.64 (1.44–1.87)	< 0.01	1.71 (1.38–2.11)	< 0.01	3.51 (2.28–5.40)	< 0.01
Moderate	1.42 (1.26–1.61)	< 0.01	1.75 (1.53–2.00)	< 0.01	1.44 (1.30–1.61)	< 0.01	0.98 (0.79–1.21)	0.84	1.99 (1.28–3.10)	< 0.01
Changes in lifestyle during COVID-19 pandemic (ref: no)										
Yes	1.40 (1.23–1.60)	< 0.01	1.19 (1.03–1.38)	0.02	1.55 (1.37–1.74)	< 0.01	—	—	—	—
Alcohol use during COVID-19 pandemic (ref: no)										
Yes	1.46 (1.15–1.87)	< 0.01	—	—	1.26 (1.03–1.53)	0.02	—	—	2.31 (1.53–3.49)	< 0.01
Tobacco use during COVID-19 pandemic (ref: no)										
Yes	—	—	—	—	—	—	1.34 (0.95–1.88)	0.10	—	—
Have you learned about some mental health knowledge during COVID-19 pandemic? (ref: no)										
Yes	—	—	—	—	—	—	0.86 (0.73–1.01)	0.06	—	—
Do you engage in regular physical exercise during COVID-19 pandemic? (ref: no)										
Yes	0.86 (0.77–0.97)	0.01	—	—	0.82 (0.75–0.91)	< 0.01	—	—	—	—
Symptoms of depression (ref: no)										
Yes	—	—	19.98 (17.50–22.82) ^f	< 0.01	3.73 (3.31–4.19) ^g	< 0.01	3.91 (2.60–5.88) ^h	< 0.01	2.58 (1.54–4.33) ⁱ	< 0.01
Symptoms of anxiety (ref: no)										
Yes	19.89 (17.41–22.72) ^e	< 0.01	—	—	1.94 (1.72–2.19) ^g	< 0.01	13.26 (9.27–18.97) ^h	< 0.01	1.66 (1.07–2.59) ⁱ	0.02
Symptoms of insomnia (ref: no)										
Yes	3.68 (3.26–4.14) ^e	< 0.01	1.98 (1.75–2.23) ^f	< 0.01	—	—	4.43 (3.54–5.54) ^h	< 0.01	—	—
Symptoms of PTSD (ref: no)										
Yes	3.75 (2.47–5.69) ^e	< 0.01	13.45 (9.37–19.30) ^f	< 0.01	4.36 (3.50–5.45) ^g	< 0.01	—	—	2.71 (1.98–3.71) ⁱ	< 0.01
Suicidal behavior (ref: no)										
Yes	2.00 (1.17–3.43) ^e	0.01	1.56 (1.01–2.41) ^f	0.04	—	—	2.84 (2.05–3.93) ^h	< 0.01	—	—

^aDepression was defined as PHQ-9 score ≥ 5 .

^bAnxiety was defined as GAD-7 score ≥ 5 .

^cInsomnia was defined as ISI score ≥ 8 .

^dPTSD was defined as PCL-5 score ≥ 33 .

^eThe highest VIF in the model of "depression" is 1.51 (symptoms of anxiety), 1.41 (symptoms of insomnia), 1.29 (symptoms of PTSD), and 1.06 (suicidal behavior).

^fThe highest VIF in the model of "anxiety" is 1.52 (symptoms of depression), 1.46 (symptoms of insomnia), 1.23 (symptoms of PTSD), and 1.06 (suicidal behavior).

^gThe highest VIF in the model of "insomnia" is 2.00 (symptoms of depression), 2.07 (symptoms of anxiety), and 1.24 (symptoms of PTSD).

^hThe highest VIF in the model of "PTSD" is 2.10 (symptoms of depression), 2.01 (symptoms of anxiety), 1.44 (symptoms of insomnia), and 1.04 (suicidal behavior).

ⁱThe highest VIF in the model of "suicidal behavior" is 1.98 (symptoms of depression) and 2.07 (symptoms of anxiety), and 1.25 (symptoms of PTSD).

health problems and suicidal behavior during the COVID-19 pandemic (2, 47–49). Furthermore, this study showed that suicidal behavior tends to increase when mental health status

is exacerbated during the pandemic, which indicate that the importance of establishing mental health promotion strategies to further prevent suicidal behavior (2, 47, 48). COVID-19

pandemic-related stress, and changes in health-related behaviors (e.g., exercise and alcohol consumption) were associated with a higher risk of mental health problems and suicidal behavior. Participants who were under moderate to high stress had a higher risk of mental health symptoms and suicidal behavior. This is consistent with findings that greater stress was significantly correlated with more negative psychological effects in both patients with severe acute respiratory syndrome and healthy control subjects (50). Previous studies also indicated that people under higher levels of stress, which might have come from work, family, or traumatic life events, were more vulnerable to suicidal behavior compared with control groups (51, 52). Moreover, a higher probability of mental health problems and suicidal behavior was also identified among individuals who experienced changes in lifestyle and who reported alcohol use during the pandemic. As a consequence of school closures, students' daily routines have changed dramatically, such as a reduction of social contacts and an increase of problem internet use, which can result in a higher incidence of mental health problems and suicidal behavior (53, 54). Furthermore, previous studies suggested that alcohol abuse was a major risk factor for mental health problems and suicidal behavior because of increases in impulsive and aggressive behaviors (51, 55–57). Conversely, limiting alcohol consumption might be associated with lower risk of mental illness and suicidal behavior (58). Thus, to avoid mental health problems during the COVID-19 pandemic, young people should maintain a healthy diet and adaptive lifestyle (59). We found that students who engaged in regular physical activities during the pandemic had a lower risk of depression, and insomnia. These findings are consistent with another study (60) of Chinese college students during the pandemic. Similarly, a previous study (22) of university students in France confirmed that more frequent physical activity was associated with less severe self-reported mental health symptoms. Thus, regular exercise is an effective strategy to promote mental well-being during the pandemic.

STRENGTHS AND LIMITATIONS

The strengths of this study included university students with high COVID-19 exposure experience in Wuhan, extensive geographic coverage in China, the large sample size, the evaluation of multiple risk factors, and the assessment of risk factors after long-term quarantine. The prevention and control of the pandemic in China is stabilizing, but sporadic cases are still being reported. Thus, issues that are related to prevention and control strategies still need to be addressed. We conducted this study with students who experienced long-term mandatory school closure (i.e., ~one semester of the school year). Students who experience long-term school closure may be more vulnerable than students who experience shorter-term closure (18, 19, 31, 61). The COVID-19 pandemic has persisted for nearly 1 year at the time of this writing, which has negatively impacted the daily lives of people worldwide. Its impact on mental health problems and suicidal behavior need to be explored. Our findings provide useful information for health policies, the identification of

at-risk students, and the development of population-specific psychological crisis management.

Our study also has several limitations. First, there were recall bias and information bias due to this survey included some questions about past situations, and selection bias since this was an voluntary online survey and the participation rate is low. Thus, the representativeness of the sample might be insufficient. Second, mental health symptoms and suicidal behavior were based on self-rating scales and items rather than clinical diagnoses. We were also unable to distinguish between preexisting mental health symptoms and new symptoms. Third, this was a cross-sectional study that reflected mental health status and suicidal behavior during the pandemic. Therefore, we were unable to identify associations between mental health problems and risk factors because of unclear chronology. More long-term longitudinal follow-up studies are warranted in the future.

CONCLUSIONS

The prevalence of symptoms of depression, anxiety, insomnia, and PTSD and suicidal behavior was high among university students who attended universities in Wuhan, China after long-term quarantine and school closures, especially among students with high stress, who experienced changes in lifestyle, and who reported alcohol use during the pandemic. Furthermore, graduation, distant relations with parents, and a personal or family history of mental illness were also associated with a higher risk of adverse mental health problems and suicidal behavior. These findings suggest that the COVID-19 pandemic may have severe and negative effects on mental health and suicidal behavior among vulnerable university students. Specific interventions that promote mental well-being among university students who are exposed to the long-term impact of the COVID-19 pandemic should be implemented.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by The ethics committee of Peking University Sixth Hospital (Institute of Mental Health). Written informed consent from the participants' legal guardian/next of kin was not required to participate in this study in accordance with the national legislation and the institutional requirements.

AUTHOR CONTRIBUTIONS

LLu, YB, and GH designed this survey, commented on and revised the manuscript, and wrote the final version. YX and SSu were responsible for data analysis and writing the initial draft. ZJ, QL, SG, YL, GH, and YB contributed to recruiting participants

and data collection. QL, LLi, YZ, LS, JQ, JD, SM, WY, YS, PW, KY, XL, SSun, and MR designed the questionnaire and commented on and revised the manuscript. AR, SC, YW, XT, and JS revised the manuscript. All of the authors contributed to the final draft of the manuscript.

FUNDING

This study was supported by grants from the National Key Research and Development Program of China (no. 2019YFA0706200), National Natural Science Foundation of China (no. 81761128036, 81821092, and 31900805), Special Research Fund of PKUHSC for Prevention and Control of

COVID-19, and Fundamental Research Funds for Central Universities (no. BMU2020HKYZZX008).

ACKNOWLEDGMENTS

We thank all of the participants for their willingness to participate in the study and the time that they devoted to the study.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpsy.2021.695017/full#supplementary-material>

REFERENCES

- Mahase E. Covid-19: WHO declares pandemic because of "alarming levels" of spread, severity, and inaction. *BMJ*. (2020) 368:m1036. doi: 10.1136/bmj.m1036
- Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet*. (2020) 395:912–20. doi: 10.1016/S0140-6736(20)30460-8
- Galea S, Merchant RM, Lurie N. The mental health consequences of COVID-19 and physical distancing: the need for prevention and early intervention. *JAMA Intern Med*. (2020) 180:817–18. doi: 10.1001/jamainternmed.2020.1562
- Shi L, Lu ZA, Que JY, Huang XL, Liu L, Ran MS, et al. Prevalence of and risk factors associated with mental health symptoms among the general population in china during the coronavirus disease 2019 pandemic. *JAMA Netw Open*. (2020) 3:e2014053. doi: 10.1001/jamanetworkopen.2020.14053
- Xin M, Luo S, She R, Yu Y, Li L, Wang S, et al. Negative cognitive and psychological correlates of mandatory quarantine during the initial COVID-19 outbreak in China. *Am Psychol*. (2020) 75:607–17. doi: 10.1037/amp0000692
- Duan L, Zhu G. Psychological interventions for people affected by the COVID-19 epidemic. *Lancet Psychiatry*. (2020) 7:300–2. doi: 10.1016/S2215-0366(20)30073-0
- Lai J, Ma S, Wang Y, Cai Z, Hu J, Wei N, et al. Factors associated with mental health outcomes among health care workers exposed to coronavirus disease 2019. *JAMA Netw Open*. (2020) 3:e203976. doi: 10.1001/jamanetworkopen.2020.3976
- Que J, Shi L, Deng J, Liu J, Zhang L, Wu S, et al. Psychological impact of the COVID-19 pandemic on healthcare workers: a cross-sectional study in China. *Gen Psychiatr*. (2020) 33:e100259. doi: 10.1136/gpsych-2020-100259
- Yang Y, Li W, Zhang Q, Zhang L, Cheung T, Xiang YT. Mental health services for older adults in China during the COVID-19 outbreak. *Lancet Psychiatry*. (2020) 7:e19. doi: 10.1016/S2215-0366(20)30079-1
- Liu JJ, Bao Y, Huang X, Shi J, Lu L. Mental health considerations for children quarantined because of COVID-19. *Lancet Child Adolesc Health*. (2020) 4:347–49. doi: 10.1016/S2352-4642(20)30096-1
- Zhai Y, Du X. Addressing collegiate mental health amid COVID-19 pandemic. *Psychiatry Res*. (2020) 288:113003. doi: 10.1016/j.psychres.2020.113003
- Qi R, Chen W, Liu S, Thompson PM, Zhang LJ, Xia F, et al. Psychological morbidities and fatigue in patients with confirmed COVID-19 during disease outbreak: prevalence and associated biopsychosocial risk factors. *medRxiv*. (2020). doi: 10.1101/2020.05.08.20031666
- Killgore WDS, Cloonan SA, Taylor EC, Fernandez F, Grandner MA, Dailey NS. Suicidal ideation during the COVID-19 pandemic: the role of insomnia. *Psychiatry Res*. (2020) 290:113134. doi: 10.1016/j.psychres.2020.113134
- Jefsen OH, Rohde C, Nørremark B, Østergaard SD. COVID-19-related self-harm and suicidality among individuals with mental disorders. *Acta Psychiatr Scand*. (2020) 142:152–53. doi: 10.1111/acps.13214
- World Health Organization. Mental health: suicide data. Available online at: https://www.who.int/mental_health/prevention/suicide/suicideprevent/en/ (accessed September 22, 2020).
- Mortier P, Cuijpers P, Kiekens G, Auerbach RP, Demyttenaere K, Green JG, et al. The prevalence of suicidal thoughts and behaviours among college students: a meta-analysis. *Psychol Med*. (2018) 48:554–65. doi: 10.1017/S0033291717002215
- Rubin GJ, Wessely S. The psychological effects of quarantining a city. *BMJ*. (2020) 368:m313. doi: 10.1136/bmj.m313
- Wang X, Hegde S, Son C, Keller B, Smith A, Sasangohar F. Investigating mental health of US college students during the covid-19 pandemic: cross-sectional survey study. *J Med Internet Res*. (2020) 22:e22817. doi: 10.2196/22817
- Cao W, Fang Z, Hou G, Han M, Xu X, Dong J, et al. The psychological impact of the COVID-19 epidemic on college students in China. *Psychiatry Res*. (2020) 287:112934. doi: 10.1016/j.psychres.2020.112934
- Jiang CX, Li ZZ, Chen P, Chen LZ. Prevalence of depression among college-goers in mainland china: a methodical evaluation and meta-analysis. *Medicine*. (2015) 94:e2071. doi: 10.1097/MD.0000000000002071
- Beiter R, Nash R, McCrady M, Rhoades D, Linscomb M, Clarahan M, et al. The prevalence and correlates of depression, anxiety, and stress in a sample of college students. *J Affect Disord*. (2015) 173:90–6. doi: 10.1016/j.jad.2014.10.054
- Wathelet M, Duhem S, Vaiva G, Baubet T, Habran E, Veerapa E, et al. Factors associated with mental health disorders among university students in france confined during the covid-19 pandemic. *JAMA Netw Open*. (2020) 3:e2025591. doi: 10.1001/jamanetworkopen.2020.25591
- Debowska A, Horecny B, Boduszek D, Dolinski D. A repeated cross-sectional survey assessing university students' stress, depression, anxiety, and suicidality in the early stages of the COVID-19 pandemic in Poland. *Psychol Med*. (2020) 2:1–4. doi: 10.1017/S003329172000392X
- Zhao Y, Jiang Z, Guo S, Wu P, Lu Q, Xu Y, et al. Association of Symptoms of Attention Deficit and Hyperactivity with Problematic Internet Use among University Students in Wuhan, China During the COVID-19 Pandemic. *J Affect Disord*. (2021) 286:220–27. doi: 10.1016/j.jad.2021.02.078
- Sun XY, Li YX, Yu CQ, Li LM. Reliability and validity of depression scales of Chinese version: a systematic review. *Chin J Epidemiol*. (2017) 38:110–16. doi: 10.3760/cma.j.issn.0254-6450.2017.01.021
- He XY, Li CB, Qian J, Cui HS, Wu WY. Reliability and validity of a generalized anxiety disorder scale in general hospital outpatients. *Shanghai Arch Psychiatry*. (2010) 22:200–3. doi: 10.1002/cpp.1802
- Bai CJ, Ji DH, Chen LX, Li L, Wang CX. Reliability and validity of insomnia severity index in clinical insomnia patients. *Chin J Prac Nurs*. (2018) 34:2182–86. doi: 10.3760/cma.j.issn.1672-7088.2018.28.005
- Wang L, Zhang L, Armour C, Cao C, Qing Y, Zhang J, et al. Assessing the underlying dimensionality of DSM-5 PTSD symptoms in Chinese adolescents surviving the 2008 Wenchuan earthquake. *J Anxiety Disord*. (2015) 31:90–7. doi: 10.1016/j.janxdis.2015.02.006

29. Meehan PJ, Lamb JA, Saltzman LE, O'Carroll PW. Attempted suicide among young adults: progress toward a meaningful estimate of prevalence. *Am J Psychiatry*. (1992) 149:41–4. doi: 10.1176/ajp.149.1.41
30. Marcoulides KM, Raykov T. Evaluation of Variance Inflation Factors in Regression Models Using Latent Variable Modeling Methods. *Educ Psychol Meas*. (2019) 79:874–82. doi: 10.1177/0013164418817803
31. Tang W, Hu T, Hu B, Jin C, Wang G, Xie C, et al. Prevalence and correlates of PTSD and depressive symptoms one month after the outbreak of the COVID-19 epidemic in a sample of home-quarantined Chinese university students. *J Affect Disord*. (2020) 274:1–7. doi: 10.1016/j.jad.2020.05.009
32. Wu Y, Levis B, Riehm KE, Saadat N, Levis AW, Azar M, et al. Equivalency of the diagnostic accuracy of the PHQ-8 and PHQ-9: a systematic review and individual participant data meta-analysis. *Psychol Med*. (2020) 50:1368–80. doi: 10.1017/S0033291719001314
33. Suarez L, Beach SR, Moore SV, Mastromauro CA, Januzzi JL, Celano CM, et al. Use of the Patient Health Questionnaire-9 and a detailed suicide evaluation in determining imminent suicidality in distressed patients with cardiac disease. *Psychosomatics*. (2015) 56:181–9. doi: 10.1016/j.psych.2014.12.005
34. Walker J, Hansen CH, Butcher I, Sharma N, Wall L, Murray G, et al. Thoughts of death and suicide reported by cancer patients who endorsed the “suicidal thoughts” item of the PHQ-9 during routine screening for depression. *Psychosomatics*. (2011) 52:424–7. doi: 10.1016/j.psych.2011.02.003
35. Leaune E, Samuel M, Oh H, Poulet E, Brunelin J. Suicidal behaviors and ideation during emerging viral disease outbreaks before the COVID-19 pandemic: a systematic rapid review. *Prev Med*. (2020) 141:106264. doi: 10.1016/j.ypmed.2020.106264
36. Hou TY, Mao XF, Dong W, Cai WP, Deng GH. Prevalence of and factors associated with mental health problems and suicidality among senior high school students in rural China during the COVID-19 outbreak. *Asian J Psychiatr*. (2020) 54:102305. doi: 10.1016/j.ajp.2020.102305
37. Elmer T, Mephram K, Stadtfeld C. Students under lockdown: Comparisons of students' social networks and mental health before and during the COVID-19 crisis in Switzerland. *PLoS ONE*. (2020) 15:e0236337. doi: 10.1371/journal.pone.0236337
38. Usher K, Bhullar N, Durkin J, Gyamfi N, Jackson D. Family violence and COVID-19: Increased vulnerability and reduced options for support. *Int J Ment Health Nurs*. (2020) 29:549–52. doi: 10.1111/inm.12735
39. Devries KM, Mak JY, Bacchus LJ, Child JC, Falder G, Petzold M, et al. Intimate partner violence and incident depressive symptoms and suicide attempts: a systematic review of longitudinal studies. *PLoS Med*. (2013) 10:e1001439. doi: 10.1371/journal.pmed.1001439
40. Soole R, Kólves K, De Leo D. Suicide in Children: A Systematic Review. *Arch Suicide Res*. (2015) 19:285–304. doi: 10.1080/13811118.2014.996694
41. Zatti C, Rosa V, Barros A, Valdivia L, Calegari VC, Freitas LH, et al. Childhood trauma and suicide attempt: a meta-analysis of longitudinal studies from the last decade. *Psychiatry Res*. (2017) 256:353–58. doi: 10.1016/j.psychres.2017.06.082
42. Bauer AE, Maegbaek ML, Liu X, Wray NR, Sullivan PF, Miller WC, et al. Familiarity of psychiatric disorders and risk of postpartum psychiatric episodes: a population-based cohort study. *Am J Psychiatry*. (2018) 175:783–91. doi: 10.1176/appi.ajp.2018.17111184
43. Wang W, Song W, Xia Z, He Y, Tang L, Hou J, et al. Sleep disturbance and psychological profiles of medical staff and non-medical staff during the early outbreak of COVID-19 in Hubei Province, China. *Front Psychiatry*. (2020) 11:733. doi: 10.3389/fpsy.2020.00733
44. Xiao H, Shu W, Li M, Li Z, Tao F, Wu X, et al. Social distancing among medical students during the 2019 coronavirus disease pandemic in China: disease awareness, anxiety disorder, depression, and behavioral activities. *Int J Environ Res Public Health*. (2020) 17:5047. doi: 10.3390/ijerph17145047
45. Zhang C, Yang L, Liu S, Ma S, Wang Y, Cai Z, et al. Survey of insomnia and related social psychological factors among medical staff involved in the 2019 novel coronavirus disease outbreak. *Front Psychiatry*. (2020) 11:306. doi: 10.3389/fpsy.2020.00306
46. Plana-Ripoll O, Musliner KL, Dalsgaard S, Momen NC, Weyer N, Christensen MK, et al. Nature and prevalence of combinations of mental disorders and their association with excess mortality in a population-based cohort study. *World Psychiatry*. (2020) 19:339–49. doi: 10.1002/wps.20802
47. Newby JM, O'Moore K, Tang S, Christensen H, Faasse K. Acute mental health responses during the COVID-19 pandemic in Australia. *PLoS ONE*. (2020) 15:e0236562. doi: 10.1371/journal.pone.0236562
48. Que J, Yuan K, Gong Y, Meng S, Bao Y, Lu L. Raising awareness of suicide prevention during the COVID-19 pandemic. *Neuropsychopharmacol Rep*. (2020) 40:392–5. doi: 10.1002/npr.12141
49. Fusar-Poli P, Correll CU, Arango C, Berk M, Patel V, Ioannidis JPA. Preventive psychiatry: a blueprint for improving the mental health of young people. *World Psychiatry*. (2021) 20:200–21. doi: 10.1002/wps.20869
50. Lee AM, Wong JG, McAlonan GM, Cheung V, Cheung C, Sham PC, et al. Stress and psychological distress among SARS survivors 1 year after the outbreak. *Can J Psychiatry*. (2007) 52:233–40. doi: 10.1177/070674370705200405
51. Park CHK, Lee JW, Lee SY, Moon J, Jeon DW, Shim SH, et al. Suicide risk factors across suicidal ideators, single suicide attempters, and multiple suicide attempters. *J Psychiatr Res*. (2020) 131:1–8. doi: 10.1016/j.jpsychires.2020.08.018
52. Lin W, Wang H, Gong L, Lai G, Zhao X, Ding H, et al. Work stress, family stress, and suicide ideation: A cross-sectional survey among working women in Shenzhen, China. *J Affect Disord*. (2020) 277:747–54. doi: 10.1016/j.jad.2020.08.081
53. Van Lancker W, Parolin Z. COVID-19, school closures, and child poverty: a social crisis in the making. *Lancet Public Health*. (2020) 5:e243–e44. doi: 10.1016/S2468-2667(20)30084-0
54. Calati R, Ferrari C, Brittner M, Oasi O, Olié E, Carvalho AF, et al. Suicidal thoughts and behaviors and social isolation: a narrative review of the literature. *J Affect Disord*. (2019) 245:653–67. doi: 10.1016/j.jad.2018.11.022
55. Borges G, Loera CR. Alcohol and drug use in suicidal behaviour. *Curr Opin Psychiatry*. (2010) 23:195–204. doi: 10.1097/YCO.0b013e3283386322
56. Darvishi N, Farhadi M, Haghtalab T, Poorolajal J. Alcohol-related risk of suicidal ideation, suicide attempt, and completed suicide: a meta-analysis. *PLoS ONE*. (2015) 10:e0126870. doi: 10.1371/journal.pone.0126870
57. Norström T, Rossow I. Alcohol consumption as a risk factor for suicidal behavior: a systematic review of associations at the individual and at the population level. *Arch Suicide Res*. (2016) 20:489–506. doi: 10.1080/13811118.2016.1158678
58. Xuan Z, Naimi TS, Kaplan MS, Bagge CL, Few LR, Maisto S, et al. Alcohol policies and suicide: a review of the literature. *Alcohol Clin Exp Res*. (2016) 40:2043–55. doi: 10.1111/acer.13203
59. Shah K, Kamrai D, Mekala H, Mann B, Desai K, Patel RS. Focus on Mental Health During the Coronavirus (COVID-19) Pandemic: Applying Learnings from the Past Outbreaks. *Cureus*. (2020) 12:e7405. doi: 10.7759/cureus.7405
60. Zhang Y, Zhang H, Ma X, Di Q. Mental health problems during the COVID-19 pandemics and the mitigation effects of exercise: a longitudinal study of college students in China. *Int J Environ Res Public Health*. (2020) 17:3722. doi: 10.3390/ijerph17103722
61. Cheng S, Jia C, Wang Y. Only children were associated with anxiety and depressive symptoms among college students in China. *Int J Environ Res Public Health*. (2020) 17:4035. doi: 10.3390/ijerph1714035

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Xu, Su, Jiang, Guo, Lu, Liu, Zhao, Wu, Que, Shi, Deng, Meng, Yan, Sun, Yuan, Lin, Sun, Ravindran, Chen, Wing, Tang, Ran, Lu, Shi, Huang, Bao and Lu. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.