



Undergraduate Research Conference at Missouri S&T

Apr 28th, 2021 - 10:30 AM

Tempered FBM with Reflecting Walls

Zachary Miller

Follow this and additional works at: <https://scholarsmine.mst.edu/ugrc>



Part of the [Physics Commons](#)

Miller, Zachary, "Tempered FBM with Reflecting Walls" (2021). *Undergraduate Research Conference at Missouri S&T*. 25.

<https://scholarsmine.mst.edu/ugrc/2021/full-schedule/25>

This Poster is brought to you for free and open access by Scholars' Mine. It has been accepted for inclusion in Undergraduate Research Conference at Missouri S&T by an authorized administrator of Scholars' Mine. This work is protected by U. S. Copyright Law. Unauthorized use including reproduction for redistribution requires the permission of the copyright holder. For more information, please contact scholarsmine@mst.edu.

Background

Anomalous diffusion:

- Random motion that violates linear relation between mean-square displacement and time, $\langle x^2 \rangle \sim t$
- Can be caused by long-time correlations between steps
- Can be modeled mathematically by Fractional Brownian Motion (FBM)

Reflected FBM

- Recent results [1]: unusual accumulation and depletion of particles close to reflecting walls (Normal diffusion=Flat)
- Probability density features power-law singularity $P(x) \sim x^{-\kappa}$ at the wall

Tempered FBM:

- In many applications, correlations between steps have a large but finite range
- For tempered FBM, power-law correlations are cut off (tempered) beyond a certain “tempering” time scale [2]

Objective

- Study tempered FBM on a one-dimensional interval with reflecting walls
- Analyze how tempering affects accumulation and depletion of particles near the walls
- Determine functional form of probability density close to a reflecting wall


Fractional Brownian Motion

- Gaussian stochastic process with stationary, power-law correlated increments
- Discrete FBM: $x_{n+1} = x_n + \xi_n$ (X_i)
- Covariance: $\langle \xi_m \xi_{m+n} \rangle = C_n = \frac{1}{2} \sigma^2 [|n-1|^\alpha - 2|n|^\alpha + |n+1|^\alpha]$ (behaves as power law at large n)
- α : anomalous diffusion exponent ($0 < \alpha < 2$) (Normal Diffusion speed)
- Tempering $\langle \xi_m \xi_{m+n} \rangle = C_n \exp(-\frac{t}{t_c})$ (Kills Correlations)

Monte Carlo methods


- Particles move on interval $[-L, L]$ with reflecting walls at both ends
- Particles start at origin at time $t = 0$ (Define Wall)
- Correlated random numbers representing ξ created via Fourier filtering technique [3]

Results: Mean-square displacement




- Mean-square displacement vs. time for different α and tempering times t_i (Measures square of displacement)

Results: Probability density



- Stationary probability density for different α . (Normal Diffusive Case in center due to tempering)

Results: Behavior near wall



- Stationary probability density vs distance from the wall
Dashed lines are power-law fits to $P(x) \sim (x-w)^\kappa$ (Log-Log, power law flat, all power laws same exponent)

Conclusions

- Studied impact of tempering time on FBM on one-dimensional interval with reflecting walls
- Particle accumulation and depletion near reflecting walls as for normal FBM
- Width of the accumulation/depletion region controlled by tempering time
- Power law singularity in probability density close to wall, $P(x) \sim (x-w)^\kappa$ with $\kappa = (2/\alpha) - 2$ (as in the untempered case)

References + Funding

- [1] A.H.O. Wada and T. Vojta, Phys. Rev E 97, 020102 (2018)
- [2] T. Guggenberger, et al., New J. Phys. 21, 022002 (2019)
- [3] D. Molina-Garcia et al, New J. Phys. 20, 103027 (2018)
- [3] H. A. Makse et al., Phys. Rev. E 53, 5445 (1996)

Seed award from Research Corporation, NSF OAC-1919789