



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
Main Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2021

Increasing awareness of degenerative cervical myelopathy: a preventative cause of non-traumatic spinal cord injury

Zipser, Carl M ; Margetis, Konstantinos ; Pedro, Karlo M ; Curt, Armin ; Fehlings, Michael ; Sadler, Iwan ; Tetreault, Lindsay ; Davies, Benjamin M

Abstract: Degenerative cervical myelopathy (DCM) is a common non-traumatic spinal cord disorder and characterized by progressive neurological impairment. Generally, it is still underdiagnosed and referral to spine specialists is often late, when patients already present with incomplete cervical spinal cord injury (SCI). To improve early diagnosis and accelerate referral, diagnostic criteria for DCM are required. Recently, AO Spine RECODE- DCM (REsearch Objectives and Common Data Elements for Degenerative Cervical Myelopathy) (aospine.org/recode), an international, interdisciplinary and interprofessional initiative, including patients with DCM, was funded with the aim to accelerate knowledge discovery that can change outcomes. In this perspective we advocate for the participation of SCI specialists in this process, where the expertise and perspective on this disorder and requirements for the diagnostic and therapeutic work up is well developed.

DOI: <https://doi.org/10.1038/s41393-021-00711-8>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-207241>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Zipser, Carl M; Margetis, Konstantinos; Pedro, Karlo M; Curt, Armin; Fehlings, Michael; Sadler, Iwan; Tetreault, Lindsay; Davies, Benjamin M (2021). Increasing awareness of degenerative cervical myelopathy: a preventative cause of non-traumatic spinal cord injury. *Spinal Cord*, 59(11):1216-1218.

DOI: <https://doi.org/10.1038/s41393-021-00711-8>

PERSPECTIVE OPEN



Increasing awareness of degenerative cervical myelopathy: a preventative cause of non-traumatic spinal cord injury

Carl M. Zipser¹✉, Konstantinos Margetis², Karlo M. Pedro³, Armin Curt¹, Michael Fehlings⁴, Iwan Sadler⁵, Lindsay Tetreault⁶, Benjamin M. Davies⁷, On behalf of the AO Spine RECODE DCM Steering Committee* and Members of the Diagnostic Criteria Working Group*

© The Author(s) 2021

Degenerative cervical myelopathy (DCM) is a common non-traumatic spinal cord disorder and characterized by progressive neurological impairment. Generally, it is still underdiagnosed and referral to spine specialists is often late, when patients already present with incomplete cervical spinal cord injury (SCI). To improve early diagnosis and accelerate referral, diagnostic criteria for DCM are required. Recently, AO Spine RECODE-DCM (REsearch Objectives and Common Data Elements for Degenerative Cervical Myelopathy) (aospine.org/recode), an international, interdisciplinary and interprofessional initiative, including patients with DCM, was funded with the aim to accelerate knowledge discovery that can change outcomes. In this perspective we advocate for the participation of SCI specialists in this process, where the expertise and perspective on this disorder and requirements for the diagnostic and therapeutic work up is well developed.

Spinal Cord; <https://doi.org/10.1038/s41393-021-00711-8>

PERSPECTIVE

Degenerative cervical myelopathy (DCM) (historically termed “cervical spondylotic myelopathy” [CSM]) is the most common non-traumatic, progressive spinal cord disorder with an estimated 2% prevalence [1]. The disorder is indeed imprecisely and insufficiently characterized by neck and radicular pain, fine motor dysfunction, gait instability, and bladder dysfunction and for most lacks common diagnostic criteria [2]. If not recognized and treated timely, patients may eventually present as incomplete cervical spinal cord injury (SCI). This letter aims to raise awareness of these shortcomings in the neurological community and emphasize an ongoing initiative to improve clinical care and foster global research [3]. The neurological field should not be left out in this effort.

Due to a variety of symptoms, patients eventually get referred to different specialists, commonly orthopedics, neurosurgeons, neurologists, or physiotherapists. While surgical decompression of the encroached spinal cord is recommended in patients experiencing already moderate/severe, or progressive symptoms, the goal of enabling earlier and/or preventative treatment has now been defined as a priority research need. In its early stages, DCM is frequently underdiagnosed or misdiagnosed as carpal tunnel syndrome or peripheral neuropathy, until patients develop more severe impairments of upper and lower limb function urging the consideration of incomplete cervical SCI. Given that DCM is a progressive but preventable neurological condition, the delayed diagnosis and late referral for evaluation of surgical decompression, can lead to poorer neurological outcomes [4]. In addition, the

pre-operative neurological status significantly influences post-operative recovery [5]. Therefore, an early diagnosis is important to achieve good clinical results. The diagnosis of DCM is currently based on clinical signs and symptoms, eventually complemented by cervical spine MRI. However, there is still no consensus on diagnostic criteria for DCM, leading to ambiguous, descriptive clinical diagnoses, and heterogeneous definitions of DCM applied in clinical studies [6, 7]. Furthermore, as demonstrated in the UK, DCM is rarely covered in the medical curriculum [8]. AO Spine RECODE-DCM (REsearch Objectives and Common Data Elements for Degenerative Cervical Myelopathy) (aospine.org/recode) is an international, interdisciplinary, and interprofessional initiative, including patients with DCM, which aims to accelerate knowledge discovery that can change outcomes [9]. This has included the formation of research priorities such as the development of common diagnostic criteria. Alongside its importance for clinical care, a sensitive and specific set of diagnostic criteria is required to foster research, particularly for those studies aiming at investigating neuroprotective strategies. Diagnostic criteria for DCM can help to overcome several shortcomings in patient care and research (Fig. 1). The development of diagnostic criteria for DCM would benefit from the experiences of the neurological community; a UK cohort study identified 45% of cases are initially diagnosed by Neurologists, whilst Neurologists are familiar with their development and implementation [10]. Common criteria will help the dialog between neurologists, general practitioners (primary care providers), and spine specialists, and propagate the knowledge of red flags in DCM that require timely and specific

¹Spinal Cord Injury Centre, Balgrist University Hospital, Zurich, Switzerland. ²Department of Neurosurgery, Icahn School of Medicine at Mount Sinai, New York, NY, USA. ³Division of Neurosurgery, Department of Neurosciences, University of the Philippines-Philippine General Hospital, Manila, Philippines. ⁴Division of Neurosurgery, University of Toronto, Toronto, ON, Canada. ⁵Myelopathy Support, Myelopathy.org, Cambridge, UK. ⁶Department of Neurology, NYU Langone Health, Graduate Medical Education, New York, NY, USA. ⁷Department of Neurosurgery, University of Cambridge, Cambridge, UK. *Lists of authors and their affiliations appear at the end of the paper. ✉email: carl.moritz.zipser@balgrist.ch

Received: 1 July 2021 Revised: 11 September 2021 Accepted: 14 September 2021
Published online: 09 October 2021



Fig. 1 Infographic providing an overview why diagnostic criteria for DCM are needed. The panels summarize shortcomings in patient care and research which diagnostic criteria for DCM can help to overcome.

actions, before established SCI. The development of diagnostic criteria involves both comprehensive diagnostic criteria for research and spine specialists, as well as easily applicable algorithms that speed up referral to cervical spine MRI and spine specialists. We have established an initial working group to act upon this opportunity (AO Spine RECODE DCM Diagnostic Criteria Incubator). If you are interested in contributing to this process, please contact us.

REFERENCES

- Smith SS, Stewart ME, Davies BM, Kotter MRN. The prevalence of asymptomatic and symptomatic spinal cord compression on magnetic resonance imaging: a systematic review and meta-analysis. *Glob Spine J.* 2021;11:597–607.
- Badhiwala JH, Ahuja CS, Akbar MA, Witiw CD, Nassiri F, Furlan JC, et al. Degenerative cervical myelopathy - update and future directions. *Nat Rev Neurol.* 2020;16:108–24.
- Grodzinski B, Bestwick H, Bhatti F, Durham R, Khan M, Partha Sarathi CI, et al. Research activity amongst DCM research priorities. *Acta Neurochir.* 2021;163: 1561–8.
- Pope DH, Mowforth OD, Davies BM, Kotter MRN. Diagnostic delays lead to greater disability in degenerative cervical myelopathy and represent a health inequality. *Spine.* 2020;45:368–77.
- Tetreault L, Kopjar B, Côté P, Arnold P, Fehlings MG. A clinical prediction rule for functional outcomes in patients undergoing surgery for degenerative cervical myelopathy: analysis of an international prospective multicenter data set of 757 subjects. *JBJS.* 2015;97:2038–46.
- Davies BM, McHugh M, Elgheriani A, Kolias AG, Tetreault L, Hutchinson PJA, et al. The reporting of study and population characteristics in degenerative cervical myelopathy: a systematic review. *PLoS ONE.* 2017;12:e0172564.
- Khan DZ, Khan MS, Kotter MR, Davies BM. Tackling research inefficiency in degenerative cervical myelopathy: illustrative review. *JMIR Res Protoc.* 2020;9:e15922-e.
- Waqar M, Wilcock J, Garner J, Davies B, Kotter M. Quantitative analysis of medical students' and physicians' knowledge of degenerative cervical myelopathy. *BMJ Open.* 2020;10:e028455.
- Davies BM, Khan DZ, Mowforth OD, McNair AGK, Gronlund T, Kolias AG, et al. RECODE DCM (REsearch Objectives and Common Data Elements for Degenerative Cervical Myelopathy): a consensus process to improve research efficiency in DCM, through establishment of a standardized dataset for clinical research and the definition of the research priorities. *Glob Spine J.* 2019;9:655–76S.
- Hilton B, Tempest-Mitchell J, Davies B, Kotter M. Route to diagnosis of degenerative cervical myelopathy in a UK healthcare system: a retrospective cohort study. *BMJ Open.* 2019;9:e027000.

AUTHOR CONTRIBUTIONS

CZ and BD: conceptualisation; CZ, KM, KP, AC, MF, IS, LT, BD: investigation; AC, MF, BD: supervision; CZ, IS, BD: visualisation; CZ and BD: writing—original draft; CZ, KM, KP, AC, MF, IS, LT, BD: writing—review and editing.

FUNDING

Open Access funding provided by Universität Zürich.

COMPETING INTERESTS

The authors declare no competing interests.

ADDITIONAL INFORMATION

Correspondence and requests for materials should be addressed to Carl M. Zipser.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the

article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021

ON BEHALF OF THE AO SPINE RECODE DCM STEERING COMMITTEE

Mark Kotter^{7,8}, Benjamin Davies^{7,8}, Brian Kwon⁹, Michael Fehlings⁴, Carl Zipser¹, Shekar Kurpad¹⁰, Vafa Rahimi-Movaghar¹¹, Bizhan Aarabi¹², James Harrop¹³, Armin Curt¹, James Guest¹⁴, Sukhvinder Kalsi-Ryan¹⁵, Angus McNair¹⁶, Julio Furlan^{17,18}, Iwan Sadler⁸, Delphine Houghton⁸, Ellen Sarewitz⁸, Julia Carter⁸, Margot Miller⁸, Timothy Boerger¹⁰, Paige Howard⁸, Shirley Widdop⁸, Carla Salzman⁸, Jamie Milligan¹⁹, Geno J. Merli²⁰, Robert Chen²¹, Jefferson R. Wilson⁴ and Ricardo Rodrigues-Pinto²²

⁸Myelopathy.org, International Charity for Degenerative Cervical Myelopathy, Bristol, Great Britain. ⁹Vancouver Spine Surgery Institute, Department of Orthopedics, The University of British Columbia, Vancouver, BC, Canada. ¹⁰Department of Neurosurgery, Medical College of Wisconsin, Wauwatosa, WI, USA. ¹¹Department of Neurosurgery, Sina Trauma and Surgery Research Center, Tehran University of Medical Sciences, Tehran, Iran. ¹²Department of Neurosurgery, University of Maryland School of Medicine, Baltimore, MD, USA. ¹³Department of Neurological Surgery, Thomas Jefferson University, Philadelphia, PA, USA. ¹⁴Department of Neurosurgery and The Miami Project to Cure Paralysis, The Miller School of Medicine, University of Miami, Miami, FL, USA. ¹⁵KITE|Toronto Rehab|University Health Network. Clinic/Scientific Lead - Rocket Family Upper Extremity Clinic-Lyndhurst Site, Toronto, ON, Canada. ¹⁶Bristol Centre for Surgical Research, University of Bristol, Bristol, Great Britain. ¹⁷KITE Research Institute, University Health Network, Toronto, ON, Canada. ¹⁸Department of Medicine, Division of Physical Medicine and Rehabilitation, University of Toronto, Toronto, ON, Canada. ¹⁹Department of Family Medicine, McMaster University, Hamilton, ON, Canada. ²⁰Jefferson Vascular Center, Thomas Jefferson University, Philadelphia, PA, USA. ²¹Krembil Research Institute, Division of Neurology, Department of Medicine, University of Toronto, Toronto, ON, Canada. ²²Spinal Unit (UVM), Department of Orthopaedics, Centro Hospitalar Universitário do Porto - Hospital de Santo António, Porto, Portugal

MEMBERS OF THE DIAGNOSTIC CRITERIA WORKING GROUP

Michael Fehlings⁴, Lindsay Tetreault⁶, Ricardo Rodrigues-Pinto²², Jamie R. F. Wilson²³, Nitin Kimmatkar²⁴, Jamie Milligan¹⁹, Benjamin Davies^{7,8,25}, Ratko Yurac²⁵, Lucy Cameron²⁶, Carl Zipser¹, Mohamed Abdel-Wanis²⁷, Ellen Sarewitz⁸, Bruno Lourenço Costa²⁸, Shirley Widdop⁸, Michael Betz²⁹, Oke Obadaseraye³⁰, Karlo Pedro³¹, Lianne Wood³² and Konstantinos Margetis²

²³Department of Neurosurgery, University of Nebraska Medical Center, Omaha, NE, USA. ²⁴Government Medical College Nagpur, Nagpur, India. ²⁵Universidad del Desarrollo, Santiago, Chile. ²⁶University Hospitals NHS Foundation Trust, Addenbrooke's or Rosie Hospital, Cambridge, Great Britain. ²⁷Sohag Faculty of Medicine, Sohag Governorate, Egypt. ²⁸Hospital de São Teotónio (Centro Hospitalar Tondela - Viseu, Epe), Coimbra, Portugal. ²⁹University Spine Center, Balgrist University Hospital, Zurich, Switzerland. ³⁰Orthopaedics and Trauma, Lagos, Nigeria. ³¹Philippine General Hospital, University of the Philippines, Manila, Philippines. ³²Nottingham University Hospital, Nottingham, Great Britain.