

EVALUATION OF THE PHOTOPROTECTIVE ROLE OF QUERCETIN TO
SELECTED LIGHT-ADAPTED AND SHADE-TOLERANT PLANT SPECIES

AISHA IDRIS

A thesis submitted in fulfilment of the requirement for the award of the Doctor of
Philosophy in Science

Faculty of Applied Sciences and Technology
Universiti Tun Hussein Onn Malaysia

JUNE 2019

DEDICATION

The thesis is dedicated to my parents Alhaji Idris Yusuf and Hajiya Hauwa Muhammad Babayo for their prayers, advice and encouragement, which guided me to this achievement. The thesis is also dedicated to my husband Zakiyyu, my siblings, and my children Muhammad, Iman, Hauwa, and Muhammad. May Allah bless them.



ACKNOWLEDGEMENT

Praise be to Allah, the Cherisher and Sustainer of the worlds. My special thanks go to my supervisor, Associate Professor Dr Alona C. Linatoc for her patience, guidance, encouragement, commitment, and advice during the duration of my study. I also appreciate my Co-supervisor, Associate Professor Dr Mohd Fadzelly Abu Bakar for his guidance and advice during the study.

My appreciation also goes to the staffs of Johor National Parks Corporation (JNPC) for their assistance. I also like to thank Mr Ishak Ayub and Mr Akmal for their help during laboratory analysis. I am grateful to the Office for Research, Innovation, Commercialization and Consultancy Management, Universiti Tun Hussein Onn Malaysia (UTHM) for the financial support under UTHM TIER 1 Vot no. 906.

Lastly. My appreciation goes to my parents (Alhaji Idris Yusuf and Hajiya Hauwa Muhammad Babayo), brothers (Hassan, Hussain, Ahmad, and Yusuf) sisters (Zainab, Hafsat, Amina, and Fatima), husband (Zakiyyu Ibrahim Takai), children (Muhammad, Iman, Hauwa, and Muhammad), mother-in-law (Halima Isa) and my brothers-in-law (Hussain Dadan Garba and Fareed L. Bello) for their help and support. May Almighty Allah bless and reward everyone abundantly.

ABSTRACT

Plants respond to different light intensities depending on their genetic make-up, mutation and other environmental conditions. *Mikania micrantha*, *Clidemia hirta*, and *Tetracera sarmentosa* were selected so as to analyse how varying light intensity affects some gas exchange characteristics, pigment and production of total flavonoid content. Besides, the study was aimed at evaluating the photoprotective role of quercetin to the selected plants, in addition to correlating how the production of flavonoids affects the plants photosynthesis. The photosynthetic rates of the selected plants were determined using LI-6400. The chlorophyll, carotenoid, anthocyanin, flavonoid, antioxidant enzymes, malondialdehyde, soluble sugar and soluble protein contents were quantified using spectroscopic techniques. Quercetin was quantified using high-performance liquid chromatography (HPLC). Sun-exposed plants were having the maximum photosynthesis and quercetin content compared with semi-shaded or fully shaded plants. The highest quercetin content was recorded for sun-exposed *C. hirta* ($0.950 \pm 0.023 \mu\text{g/ml}$) while the lowest was recorded for shaded *T. sarmentosa* ($0.13 \pm 0.007 \mu\text{g/ml}$). The highest oxidative stress was recorded for sun-exposed *T. sarmentosa* ($6.19 \pm 0.19 \mu\text{g/ml}$) which was also having the lowest quantum efficiency of photosystem II (0.509 ± 0.003). Superoxide dismutase activity was lowest under sun-exposed *C. hirta* ($1.86 \pm 0.06 \text{ U/mg protein}$), while peroxidase and catalase were lowest under Sun-exposed *T. sarmentosa* (59.59 ± 2.67 and 3.75 ± 0.17) U/mg protein respectively. The result obtained makes it possible to accept the generated hypothesis of the study because the quercetin content was higher when the antioxidant enzymes of the plants were low. This leads to a conclusion that increase in the production of secondary metabolites at high light intensity is not due to high CO₂ assimilation rate, but rather due to the production of photoprotective metabolites to conquer the light stress.

ABSTRAK

Tindak balas terhadap keamatan cahaya yang berlainan bergantung kepada genetik, mutasi dan keadaan alam sekitar yang lain. *Mikania micrantha*, *Clidemia hirta*, dan *Tetracera sarmentosa* dipilih untuk menganalisa bagaimana variasi keamatan cahaya mempengaruhi beberapa ciri pertukaran gas, pigmen dan pengeluaran jumlah kandungan flavonoid. Selain itu, penyelidikan ini bertujuan untuk menilai peranan fotoprotektif kuercetin terhadap tanaman pilihan, di samping mengaitkan bagaimana pengeluaran flavonoid mempengaruhi fotosintesis tumbuhan. Fotosintesis tumbuhan pilihan ditentukan menggunakan LI-6400. Klorofil, karotenoid, antosianin, flavonoid, enzim antioksidan, malondialdehid, gula larut dan kandungan protein larut dikira secara teknik spektroskopi. Kuercetin dikira menggunakan kromatografi cecair prestasi tinggi (HPLC). Tumbuhan yang terdedah kepada matahari mempunyai kandungan fotosintesis dan kuercetin yang maksimum berbanding dengan tumbuhan teduh separuh atau teduh penuh. Kandungan kuercetin tertinggi direkodkan oleh *C. hirta* ($0.950 \pm 0.023 \mu\text{g/ml}$) manakala yang paling rendah direkodkan oleh *T. sarmentosa* ($0.13 \pm 0.007 \mu\text{g/ml}$). Tekanan oksidatif tertinggi direkodkan oleh *T. sarmentosa* ($6.19 \pm 0.19 \mu\text{g / ml}$) yang terdedah kepada matahari yang juga mempunyai kecekapan terendah dalam fotosistem II (0.509 ± 0.003). Aktiviti superoxide dismutase adalah terendah pada *C. hirta* ($1.86 \pm 0.06 \text{ U / mg protein}$) manakala peroksidase dan catalase adalah terendah pada *T. sarmentosa* (59.59 ± 2.67 dan 3.75 ± 0.17) U / mg protein masing-masing. Keputusan yang diperoleh memungkinkan untuk menerima hipotesis kajian kerana kandungan kuercetin lebih tinggi apabila enzim antioksidan tumbuhan adalah rendah. Secara kesimpulannya, peningkatan pengeluaran metabolit sekunder semasa keamatan cahaya yang tinggi tidak disebabkan oleh kadar asimilasi CO₂ yang tinggi, tetapi disebabkan oleh pengeluaran metabolit pelindung cahaya untuk menangani tekanan cahaya.

TABLE OF CONTENTS

TITLE	i
DECLARATION	ii
DEDICATION	iii
ACKNOWLEDGEMENT	iv
ABSTRACT	v
ABSTRAK	vi
TABLE OF CONTENTS	vii
LIST OF TABLES	xi
LIST OF FIGURES	xiii
LIST OF SYMBOLS AND ABBREVIATIONS	xv
LIST OF APPENDICES	xviii
CHAPTER 1 INTRODUCTION	1
1.1 Background and rationale of the study	1
1.2 Problem statement	4
1.3 Aim and objectives of the study	5
1.4 Scope of the study	6
1.5 Significance of the study	6
CHAPTER 2 LITERATURE REVIEW	8
2.1 Light	8
2.1.1 Plant photoreceptors: how plants see the invisible	10
2.1.2 Plants adaptation to various light intensity	11
2.2 Photosynthesis	12
2.2.1 Photosynthetic light response curve	14
2.2.2 Photosynthetic carbon dioxide response curve	17

2.2.3	Photoinhibition	19
2.2.4	Chlorophyll <i>a</i> fluorescence	20
2.2.5	Effect of light on photosynthesis	23
2.3	Photosynthetic pigments	25
2.4	Flavonoids	30
2.4.1	Flavonoids biosynthesis	31
2.4.2	Functions of flavonoids	35
2.4.3	Effect of light on flavonoids accumulation	36
2.5	Anthocyanins	39
2.5.1	Biosynthesis of anthocyanin	41
2.5.2	Function of anthocyanin	42
2.5.3	Effect of light on anthocyanin accumulation	42
2.6	Quercetin	44
2.6.1	Quercetin biosynthesis	45
2.6.2	Functions of quercetin	46
2.6.3	Methods employed for quercetin quantification	47
2.6.4	Effect of light on quercetin accumulation	48
2.7	Studied plants	50
2.7.1	<i>Mikania micrantha</i>	50
2.7.2	<i>Clidemia hirta</i>	51
2.7.3	<i>Tetracera sarmentosa</i>	52
CHAPTER 3 METHODOLOGY		54
3.1	Study area	54
3.2	Plant identification and voucher specimen preparation	55
3.3	Selection of plants	56
3.4	Selection of leaves	56
3.5	Gas exchange measurement	57
3.5.1	Photosynthetic light response (A/Q)	

	curve	59
3.5.2	Photosynthetic carbon dioxide response (A/C) curve	60
3.5.3	Determination of chlorophyll <i>a</i> fluorescence parameters	60
3.6	Determination of photosynthetic pigment contents	61
3.7	Determination of flavonoid content	62
3.7.1	Determination of total flavonoid content (TFC)	62
3.7.2	Determination of the anthocyanin contents	63
3.7.3	Determination of quercetin content	64
3.8	Quantitative determination of lipid peroxidation	65
3.9	Determination of antioxidant enzymes activity	66
3.8.1	Determination of superoxide dismutase (SOD)	66
3.8.2	Determination of peroxidase (POD)	67
3.8.3	Determination of catalase (CAT)	68
3.10	Determination of total soluble sugar (TSS)	69
3.11	Determination of soluble protein content	69
3.12	Statistical analysis	70
CHAPTER 4 RESULTS AND DISCUSSION		72
4.1	Gas exchange measurements	72
4.1.1	Photosynthetic light response (A/Q) curve	72
4.1.2	Photosynthetic carbon dioxide response (A/C) curve	79
4.1.3	Quantum efficiency of photosystem II	82
4.2	Photosynthetic pigments	83

4.3	Flavonoid content	88
4.3.1	Total flavonoid content	88
4.3.2	Anthocyanin content	91
4.3.3	Quercetin content	92
4.4	Lipid peroxidation	100
4.5	Antioxidant enzymes	102
4.6	Total soluble sugar	105
4.7	Soluble protein content	107
4.8	Photoprotective role of quercetin	109
CHAPTER 5 CONCLUSION AND RECOMMENDATION		116
5.1	Conclusion	116
5.2	Recommendation	118
REFERENCES		119
APPENDICES		158
VITA		185



LIST OF TABLES

2.1	Plant photoreceptors	10
2.2	Differences between sun-exposed and shaded leaves	12
2.3	Effect of light on gas exchange characteristics	17
2.4	Effect of light on the quantum efficiency of Photosystem II (Fv/Fm)	22
2.5	Effect of light on plant photosynthesis	25
2.6	Effect of light on chlorophyll <i>a</i> and <i>b</i>	30
2.7	Different types of flavonoid	31
2.8	Effect of light quality, intensity, PAR and photoperiod of the accumulation of flavonoids in the selected plants	39
2.9	Different types of anthocyanins	41
2.10	Effect of light on anthocyanin accumulation	43
2.11	Quercetin glycosides	45
2.12	Effect of light on quercetin accumulation	49
4.1	Light compensation points (LCP)	74
4.2	Light saturation points (LSP)	75
4.3	Maximum net assimilation rate (A_{\max})	77
4.4	Apparent quantum yield (Φ)	78
4.5	Maximum carboxylation rate of RuBisCO (V_{cmax})	81
4.6	Maximum electron transport rate (J_{\max})	81
4.7	The quantum efficiency of photosystem II	82
4.8	Chlorophyll <i>a</i> content	84
4.9	Chlorophyll <i>b</i> content	84
4.10	Total chlorophyll <i>a+b</i> content	85
4.11	Carotenoid content	86
4.12	Chlorophyll <i>a/b</i> ratio	87

4.13	Total chlorophyll <i>a+b</i> /Carotenoid ratio	88
4.14	Total flavonoid content (TFC)	89
4.15	Anthocyanin content	91
4.16	Total quercetin content	98
4.17	Correlation coefficients among quercetin, anthocyanin and TFC	100
4.18	Malondialdehyde (MDA) content	101
4.19	Superoxide dismutase (SOD) activity	103
4.20	Peroxidase (POD) activity	103
4.21	Catalase (CAT) activity	104
4.22	Total soluble sugar content (TSS)	106
4.23	Soluble protein content (SP)	108
4.24	Correlation coefficients among MDA, SOD, POD, CAT and quercetin content	110
4.25	Correlation coefficients among quercetin contents, A_{net} , TSS and SP	115



LIST OF FIGURES

2.1	Electromagnetic radiations	9
2.2	Plant photoreceptors absorption zones	11
2.3	Z-scheme of photosynthesis	13
2.4	Photosystems	14
2.5	The photosynthetic light response curve	15
2.6	The photosynthetic carbon dioxide response curve	18
2.7	Photoinhibition of photosystem II	20
2.8	A: Photochemical (qP) and non-photochemical quenching (NPQ); B: Chlorophyll <i>a</i> fluorescence parameters	22
2.9	Photosynthetic pigments absorption spectrum	26
2.10	Structure of chlorophyll <i>a</i>	27
2.11	Structure of chlorophyll <i>b</i>	28
2.12	Structure of carotenoid	29
2.13	Basic structure of flavonoid	31
2.14	The flavonoid biosynthetic pathway	32
2.15	Shikimic acid pathway	33
2.16	Dihydroxy-B-ring-substituted flavonoid glycosides in sun-exposed <i>Ligustrum vulgare</i> leaves	34
2.17	Basic structure of anthocyanin	40
2.18	Molecular structure of quercetin	44
2.19	Quercetin biosynthetic pathway	46
2.20	<i>Mikania micrantha</i>	51
2.21	<i>Clidemia hirta</i>	52
2.22	<i>Tetracera sarmentosa</i>	53
3.1	Gunung Ledang Malaysia	55
3.2	Flowchart for the study	57

3.3	Portable photosynthesis system, LI-6400	58
4.1	Photosynthetic light response (A/Q) curves	73
4.2	Photosynthetic carbon dioxide response (A/C) curve	80
4.3	Quercetin standard calibration curve	89
4.4	FTIR spectra of the selected plants showing the O - quinone peak of quercetin; <i>A. M. micrantha</i> , <i>B. C. hirta</i> , <i>C. T. sarmentosa</i>	94
4.5	HPLC chromatograms of the selected plants showing quercetin peak	98
4.6	MDA standard calibration curve	101
4.7	TSS calibration curve	106
4.8	Soluble protein calibration curve	108
4.9	Daily variation in antioxidant enzyme and quercetin Accumulation	113



LIST OF SYMBOLS AND ABBREVIATIONS

%	-	Percent
°C	-	Degree Celsius
θ	-	Curve curvature
Φ	-	Apparent quantum yield
ϕ PSII	-	Efficiency of PSII photochemistry
μ g/ml	-	Microgram per milliliter
μ l	-	Microlitre
μ m	-	Micrometre
A/C	-	Photosynthetic CO ₂ response curve
A/Q	-	Photosynthetic light response curve
AlCl ₃	-	Aluminium chloride
A _{max}	-	Maximum net assimilation rate
A _{net}	-	Net assimilation rate
ANOVA	-	Analysis of variance
ATP	-	Adenosine triphosphate
C _{ar}	-	Carotenoid
CAT	-	Catalase
C _i	-	Intercellular CO ₂
CO ₂	-	Carbon dioxide
C _p	-	Crude protein
D _f	-	Dilution factor
DNA	-	Deoxyribonucleic acid
D _w	-	Dry weight
EDTA-Na ₂	-	Disodium ethylenediaminetetraacetate dehydrate
ETR	-	Electron transport rate
F _m	-	Maximal fluorescence
F _m ^l	-	Light adapted maximal fluorescence

F ₀	-	Minimal fluorescence
F ₀ ^l	-	Light adapted minimal fluorescence
F _s	-	Steady state fluorescence
F _t	-	Feet
F _v /F _m	-	Efficiency of photosynthesis
F _w	-	Fresh weight
g	-	Gram
H ₂ O	-	Water
H ₂ O ₂	-	Hydrogen peroxide
H ₂ SO ₄	-	Sulfuric acid
H ₃ PO ₄	-	Phosphoric acid
HPLC	-	High-performance liquid chromatography
IRGA	-	Infrared gas analyser
J _{max}	-	Maximum electron transport rate
LCP	-	Light compensation point
LED	-	Light-emitting diode
LSP	-	Light saturation point
m	-	Meter
M	-	Molar
MDA	-	Malondialdehyde
ml	-	Milliliter
mM	-	Millimolar
MW	-	Molecular weight
NBT	-	Nitro blue tetrazolium
nm	-	Nanometer
NPQ	-	Non-photochemical quenching
O ₂	-	Oxygen
PAL	-	Phenylalanine ammonia lyase
PAR	-	Photosynthetic active radiation
pH	-	- log [H ⁺]
PHOT	-	Phototropin photoreceptor
PPFD	-	Photosynthetic photon flux density
PSI	-	Photosystem I
PSII	-	Photosystem II

PTNJ	-	Perbadanan Taman Negara Johor
qp	-	Photochemical quenching
Rd	-	Dark respiration
ROS	-	Reactive oxygen species
RuBisCO	-	Ribulose-1, 5-bisphosphate carboxylase
SD	-	Standard deviation
SLA	-	Specific leaf area
SOD	-	Superoxide dismutase
SPSS	-	Statistical Package for the Social Science
TBA	-	Thiobarbituric acid
TCA	-	Trichloroacetic acid
TFC	-	Total flavonoid content
TSS	-	Total soluble sugar
UTHM	-	Universiti Tun Hussein Onn Malaysia
UV	-	Ultraviolet
UV-R8	-	UV-Resistance Locus 8
V_{cmax}	-	Maximum carboxylation rate of RuBisCO
W / V	-	Weight / Volume
YPF	-	Yield photon flux



LIST OF APPENDICES

APPENDIX	TITLE	PAGE
A	Research permit	158
B	Licor preparation checklist	159
C	Gas exchange measurements	160
D	Statistical analysis (D1 – D20)	161
E	Photoprotective role of quercetin	181
F	List of publications	182



PTTA UTHM
PERPUSTAKAAN TUNKU TUN AMINAH

CHAPTER 1

INTRODUCTION

1.1 Background and rationale of the study

The environment does not have a constant stable condition, it always changes, and these changes can lead to various effects in the structural, biochemical and gas exchange characteristics of a leaf including its shape, curling degree, and its surface characteristics. One of the most important environmental factor affecting plants is light. Light affects the growth and development of plants by influencing the morphology of a single leaf as well as the morphology of the whole plant. Plants can adapt to different light intensity depending on their environment. This adaptation would be possible if plant changes the distribution of its biomass and its morphology, in order to be able to utilize the amount of light they receive, so as to survive (Xue *et al.*, 2011).

Changes in the amount of light intensity received by a plant can affect the gas exchange characteristics. There is no fixed value or range of values that indicate the difference in photosynthesis of sun-exposed and shaded plants. This is because each plant responds to light differently. Also, some plants can have higher photosynthesis under higher light intensity (greater than 1000 $\mu\text{mol m}^{-2}\text{s}^{-1}$) while others have higher photosynthesis under lower light intensity (less than 1000 $\mu\text{mol m}^{-2}\text{s}^{-1}$). The compensation and saturation points of sun-exposed leaves exceeds that of shaded leaves (DeWalt, Denslow, & Hamrick, 2004; Lambers, Chapin, & Pons, 2008; Li *et al.*, 2016; Liu *et al.*, 2013; Qin *et al.*, 2012; Xue *et al.*, 2011), but in some plants like *Aeschynanthus longicaulus* (Li *et al.*, 2014), *Tetrastigma hemsleyanum* (Dai *et al.*, 2009), and *Lindera melissifolia* (Aleric & Kirkman, 2005), higher compensation and

saturation points were recorded at lower light intensity. Moreover, photosynthetic carboxylation capacity, electron transport capacity, quantum yield and curvature of light response curve varies depending on the level of light intensity received by a plant (Li *et al.*, 2017; Slattery *et al.*, 2017; Violet-Chabrand *et al.*, 2017). For instance, the maximum photosynthesis (A_{\max}) of *Chromera velia* was higher when the plant was grown at $406 \pm 30 \mu\text{mol O}_2 \text{ mg Chl}^{-1} \text{ h}^{-1}$ compared to when grown at $239 \pm 60 \mu\text{mol O}_2 \text{ mg Chl}^{-1} \text{ h}^{-1}$ (Belgio *et al.*, 2017) while *Changium smyrnioides* (Wang *et al.*, 2017) requires a semi-shaded environment for its maximum photosynthesis. Recent studies on *Eucheuma denticulatum* (Borlongan *et al.*, 2017), *Batrachospermum turfosum* (Aigner *et al.*, 2017), *Kappaphycus* sp (Borlongan *et al.*, 2017), *Typha latifolia* (Jespersen, Brix, & Sorrell, 2017), *Laminaria digitate* (Migné *et al.*, 2018), *Triticum aestivum* (Chen *et al.*, 2016), *Lactuca sativa* (Fu *et al.*, 2017), and *Vitis vinifera* (Li *et al.*, 2017) indicated that the level of light intensity received by a plant can affect the plant photosynthesis. The photosynthesis of *Oryza sativa* was higher under high light intensity (Gu *et al.*, 2017). Other plants with higher photosynthesis under high light intensity include *Brassica campestris* (Zhu *et al.*, 2017), *Betula pendula* (Niglas *et al.*, 2017), and *Phaseolus vulgaris* (Gauthier *et al.*, 2018).

Besides, the quality of light has an impact on photosynthesis (Dou *et al.*, 2017). For example, the photosynthesis rates, stomatal conductance and transpiration rates of *Dianthus caryophyllus* grown under blue light emitting diode (LED) was higher ($7.2 \mu\text{mol CO}_2 \text{ m}^{-2} \cdot \text{s}^{-1}$) than that grown under red LED ($4.1 \mu\text{mol CO}_2 \text{ m}^{-2} \cdot \text{s}^{-1}$) (Manivannan *et al.*, 2017). Likewise, the photosynthesis of *Solanum lycopersicum* increase significantly when the plant was grown under a combination of red and blue LED light (Li *et al.*, 2017). Other recent studies on effect of light quality on photosynthesis include that of *Populus trichocarpa* (Momayyezi & Guy, 2017), *Synechocystis* sp. (Oshita, Suzuki, & Kawono, 2018), *Cucumis sativus* (Särkkä *et al.*, 2017), *Cyclocarya paliurus* (Yang *et al.*, 2017), *Camptotheca acuminata* (Yu *et al.*, 2016) and *Lactuca sativa* (Zhen & van Iersel, 2017). Species cultivar (Slattery *et al.*, 2017), nutrient availability (Özer, 2017) and temperature (Lu *et al.*, 2017; Wu *et al.*, 2018) can influence the effect of light on plant photosynthesis.

Apart from variation in photosynthesis, shaded leaves differ from sun-exposed leaves in their structure and biochemical characteristics. The stomatal density (Jespersen *et al.*, 2017; Manivannan *et al.*, 2017; Sakhonwasee, Tummachai,

& Nimnoy, 2017; Sevik *et al.*, 2017; Zheng & Van Labeke, 2017), fresh and dry weight (Pedroso *et al.*, 2017, Rehana *et al.*, 2018), leaf area (Silva *et al.*, 2017), and leaf thickness (Violet-Chabrand *et al.*, 2017) differs among shaded and sun-exposed species.

In a shaded environment, the light received is weak and therefore the plant will produce more photosynthetic pigments (Belgio *et al.*, 2017; Chuyong & Acidri, 2017; Fu *et al.*, 2017; Jespersen *et al.*, 2017; Silva *et al.*, 2017; Wang *et al.*, 2017, Zhu *et al.*, 2017). Light quality can also influence the photosynthesis pigments of a plant (Rabara *et al.*, 2017; Singh *et al.*, 2017; Yu *et al.*, 2016; Zheng & Van Labeke, 2017) thereby affecting the chlorophyll *a/B* ratio.

Light intensity can affect the production of secondary metabolites like quercetin. Quercetin is a yellow flavonol which plays an important role in protecting plants against stress. It acts as an antioxidant molecule and UV-filter (Agati *et al.*, 2013; Brunetti *et al.*, 2013). It can protect plants against microorganisms; for example, *Vitis vinifera* accumulates quercetin for protection against *Plasmopara viticola* (Ali *et al.*, 2012). Moreover, quercetin plays a vital role in the interaction of a plant with its environment (Mierziak, Kostyn, & Kulma, 2014).

The total flavonoid content of a plant can be affected depending on the light intensity. For instance, *Glycine max* seedling (Yuan *et al.*, 2015) produces a high amount of flavonoids under high light intensity. *Lithocarpus litseifolius* (Li *et al.*, 2016) and *Zingiber officinale* (Ghasemzadeh *et al.*, 2010) accumulate high amount of flavonoid under lower light intensity (60 % of sunlight for the former and 310 $\mu\text{mol m}^{-2}\cdot\text{s}^{-1}$ for the latter). *Labisia pumila* accumulates more flavonoid when grown at 630 $\mu\text{mol m}^{-2}\cdot\text{s}^{-1}$ (Karimi *et al.*, 2013). *Anoectochilus roxburghii* produced a high amount of flavonoid under blue light (Ye *et al.*, 2017).

Other studies on the effect of light intensity and quality on accumulation of flavonoids include that of *Hyptis marrubioides* (Pedroso *et al.*, 2017), *Anacardium othonianum* (Gazolla *et al.*, 2017), *Chlorella miniata* (Sozmen *et al.*, 2018), *Berberis microphylla* (Arena, Postemsky, & Curvetto, 2017), *Lactuca sativa* (Bian *et al.*, 2018; Liu *et al.*, 2018; Pérez-López *et al.*, 2018), *Cannabis sativa* (Gianmaria, Gianpaolo, & Stiina, 2018), *Anoectochilus roxburghii* (Chen *et al.*, 2017), *Cyclocarya paliorus* (Liu *et al.*, 2018), *Abelmoschus esculentus* (Irshad *et al.*, 2018), *Elephantopus scaber* (Dawiyah *et al.*, 2018) and *Fagopyrum esculentum* (Nam *et al.*, 2018).

The variation in flavonoids accumulation among plant species can be due to the complex metabolism of flavonoids. In heliophytes, the activity of antioxidant enzymes decreases under lower light intensity (Li *et al.*, 2016), which increase reactive oxygen species (ROS). Due to an increase in ROS, more flavonoids would be synthesized in order to scavenge the ROS and protect the plant (Li *et al.*, 2016). The activity of antioxidant enzymes of a plant also varies depending on a plant species, and the amount of light received (Chen *et al.*, 2016). In previous studies, the activity of superoxide dismutase SOD and catalase (CAT) enzymes were higher in sun-exposed *Changium smyrnioides* (Wang *et al.*, 2017) and sun-exposed *Oryza sativa* (Gu *et al.*, 2017). In *Dianthus caryophyllus* (Manivannan *et al.*, 2017), and *Camptotheca acuminata* (Yu *et al.*, 2016), the antioxidant enzymes activity were higher under blue LED light. Variation in the antioxidant enzymes activities may be due to the level of stress encountered (acute or slow). Due to these differences, it is hypothesised in this study that “if the activity of antioxidant enzymes of a plant is low, then quercetin biosynthesis may increase”.

1.2 Problem statement

In plants, genetic make-up, mutation and environmental factors are the major characteristics that contribute to variation in response to light. The above characteristics make it difficult to come up with a general conclusion on how plants respond to light. *Labisia pumila* and *Ginkgo biloba* can accumulate a higher amount of quercetin under high light intensity while *Berberis microphylla* and *Lithocarpus litseifolius* require a semi-shaded environment for their maximum quercetin accumulation. Due to these variations, there is a need to know at which level of light intensity will the selected plants accumulate high amount of quercetin.

Some authors hypothesised that lower light intensity favours the accumulation of monohydroxy B-ring flavonoid while high light intensity favours the accumulation of dihydroxy B-ring flavonoid. Quercetin falls under the latter group, but in some plants, it is not produced under high light intensity. In addition, the selected plants are abandoned, common tropical plants. There is no available study on the photoprotective role of quercetin to the plants. Moreover, there is no available information on the activity of antioxidant enzymes and oxidative stress of

the plants. This makes it necessary to test the hypothesis on the selected plants, and determine whether high light intensity will favour the accumulation of quercetin.

Some secondary metabolites like salicylic acid can affect plant photosynthesis, where a high concentration of the acid can decrease photosynthesis. A previous hypothesis on how quercetin may decrease plant photosynthesis by inhibiting the activity of ATPase and electron transport rate indicates that flavonoids can also affect photosynthesis. Quercetin can inhibit electron transport chain of *Croton ciliatoglandulifer*. This provides the need to know whether quercetin accumulation can affect the plants photosynthesis.

When the light intensity is high, plants tend to accumulate high amount of secondary metabolites. This makes it necessary to know whether the production of high amount of secondary metabolites under high light intensity is due to high photosynthesis or due to the production of photoprotective secondary metabolites for protection against the excess light.

Lastly, the selected plants have different genetic makeup and grow in different light conditions (sun-exposed, semi-shaded and shaded environments). This provides the need to know how varying light intensity can influence the photosynthesis, pigment, and total flavonoid contents of the plants growing in their natural environment.

1.3 Aim and objectives of the study

This research aimed at identifying the photoprotective role of quercetin to selected light-adapted and shade-tolerant plant species. This was achieved by following specific objectives of the study, which are:

- i. To analyse how varying light intensity affects some gas exchange characteristics, pigment and production of total flavonoid content (TFC) of sun-exposed, semi-shaded and shaded plant species.
- ii. To evaluate the photoprotective role of quercetin to the study plants.
- iii. To correlate how the production of quercetin can affect the photosynthesis of the selected light adapted and shade tolerant plants

1.4 Scope of the study

This research focused on identifying the photoprotective role of quercetin to selected light-adapted and shade-tolerant plant species. *Mikania micrantha*, *Clidemia hirta* and *Tetracera sarmentosa* were selected for the study because the research is aimed at identifying the photoprotective role of quercetin to plants growing in their natural environment. Besides, the studied plants were found growing in sun-exposed, semi-shaded, and shaded regions of the mountain. Other reasons that justify selection of the plants include their medicinal application and their ability to grow in about 60 % of the forest. The initial step was achieved by obtaining approval from Perbadanan Taman Negara Johor (PTNJ). The plants were then selected and identified. The gas exchange was measured using the LI-6400 portable photosynthesis system. The total photosynthetic pigment contents, anthocyanin, total flavonoid, antioxidant enzymes, malondialdehyde, total soluble sugar, and soluble protein content of the plants were determined spectrophotometrically. Quercetin was quantified using high-performance liquid chromatography (HPLC). Finally, the results obtained were analysed statistically using SPSS statistical software.

1.5 Significance of the study

This research points out the benefit of producing quercetin to the plants. The findings of the study provide knowledge on the photoprotective role of quercetin to plants growing in their natural environment. This is necessary because most of the informations available on the photoprotective role of flavonoids were performed under a controlled environment. It also provides knowledge on the necessary measures required for improving the physiology of the studied plants by suggesting the best light condition for the selected plant's maximum photosynthesis. Moreover, it helps in providing information on how the studied plants survive in a constantly changing environment (that is how shaded plants tolerate lower light intensity and how sun exposed plants adapt to high light intensity), in addition to the knowledge on how plants use secondary metabolites instead of antioxidant enzymes for protection from reactive oxygen species generated due to excess light. The research provides answers to the tested hypothesis (how the activity of antioxidant enzymes is

related to quercetin biosynthesis, how light intensity affects monohydroxy B-ring and dihydroxy B-ring flavonoids, and how quercetin accumulation may affect photosynthesis). The findings identified the plants with high CO₂ absorption capacity which can be used for bioremediation and rehabilitation applications. Furthermore, the findings can be used by phytochemists when considering massive extraction of flavonoids.



REFERENCES

- Agati, G., Azzarello, E., Pollastri, S., & Tattini, M. (2012). Flavonoids as antioxidants in plants: Location and functional significance. *Plant Science*, *196*, 67–76.
- Agati, G., Biricolti, S., Guidi, L., Ferrini, F., Fini, A., & Tattini, M. (2011). The biosynthesis of flavonoids is enhanced similarly by UV radiation and root zone salinity in *L. vulgare* leaves. *Journal of Plant Physiology*, *168*(3), 204–212.
- Agati, G., Brunetti, C., Di Ferdinando, M., Ferrini, F., Pollastri, S., & Tattini, M. (2013). Functional roles of flavonoids in photoprotection: New evidence, lessons from the past. *Plant Physiology and Biochemistry*, *72*, 35–45.
- Agati, G., Galardi, C., Gravano, E., Romani, A., & Tattini, M. (2002). Flavonoid distribution in tissues of *Phillyrea latifolia* L. leaves as estimated by microspectrofluorometry and multispectral fluorescence microimaging. *Photochemistry and Photobiology*, *76*(3), 350–60. Retrieved from
- Agati, G., Matteini, P., Goti, A., & Tattini, M. (2007). Chloroplast-located flavonoids can scavenge singlet oxygen. *New Phytologist*, *174*(1), 77–89.
- Aigner, S., Holzinger, A., Karsten, U., & Kranner, I. (2017). The freshwater red alga *Batrachospermum turfosum* (Florideophyceae) can acclimate to a wide range of light and temperature conditions. *European Journal of Phycology*, *52*(2), 238–249.
- Akhtar, T. A., Lees, H. A., Lampi, M. A., Enstone, Daryl, Brain, R. A., & Greenberg, B. M. (2010). Photosynthetic redox imbalance influences flavonoid biosynthesis in *Lemna gibba*. *Plant, Cell and Environment*, *33*(7), 1205–1219.
- Alabri, Z.K., Hussain, J., Mabood, F., Ur, N., Ali, L. (2018). Fluorescence spectroscopy- Partial Least Square Regression method for the quantification of Quercetin in *Euphorbia masirahensis*. *Measurement*, *121*, 355–359.

- Aleric, K., & Kirkman, L. (2005). Growth and photosynthetic responses of the federally endangered shrub, *Lindera melissifolia* (Lauraceae), to varied light environments. *American Journal of Botany*, 92(4), 682–689.
- Alexieva, V., Sergiev, I., Mapelli, S., & Karanov, E. (2001). The effect of drought and ultraviolet radiation on growth and stress markers in pea and wheat. *Plant, Cell and Environmen*, 24, 1337–1344.
- Ali, K., Maltese, F., Figueiredo, A., Rex, M., Fortes, A. M., Zyprian, E., & Choi, Y. H. (2012). Alterations in grapevine leaf metabolism upon inoculation with *Plasmopara viticola* in different time-points. *Plant Science*, 191–192, 100–107.
- Amalesh, S., Das, G., & Das, K. S. (2011). Roles of flavonoids in Plants. *Int J Pharm Sci Tech*, 6(1), 12–35.
- Andersson, B., & Aro, E. M. (1997). Proteolytic activities and proteases of plant chloroplasts. *Physiologia Plantarum*, 100(4), 780–793.
- Anelise, W., Daniele, C. de O., Janine, de M., Karin, G., & Clarice, M. B. R. (2007). Validation of UV Spectrophotometric and HPLC Methods for Quantitative Determination of Atenolol in Pharmaceutical Preparations. *Latin American Journal of Pharmacy*, 26(5), 765–770.
- Ang, L. F., Yam, M. F., Tan, Y., Fung, T., Kiang, P. K., & Darwin, Y. (2014). HPLC Method for Simultaneous Quantitative Detection of Quercetin and Curcuminoids in Traditional Chinese Medicines. *Journal of Pharmacopuncture*, 17(4), 36–49.
- Anttonen, M. J., Hoppula, K. I., Nestby, R., Verheul, M. J., & Karjalainen, R. O. (2006). Influence of Fertilization, Mulch Color, Early Forcing, Fruit Order, Planting Date, Shading, Growing Environment, and Genotype on the Contents of Selected Phenolics in Strawberry (*Fragaria ananassa* Duch.) Fruits. *Journal of Agricultural and Food Chemistry*, 54(7), 2614–2620.
- Arakawa, O. (1993). Effect of Ultraviolet Light on Anthocyanin Synthesis in Light-Colored Sweet Cherry, cv. Sato Nishiki. *Engei Gakkai Zasshi*, 62(3), 543–546.
- Arena, M. E., Postemsky, P. D., & Curvetto, N. R. (2017). Scientia Horticulturae Changes in the phenolic compounds and antioxidant capacity of *Berberis microphylla* G. Forst. berries in relation to light intensity and fertilization. *Scientia Horticulturae*, 218, 63–71.

- Assefa, A. D., & Keum, Y. S. (2017). Effect of extraction solvent and various drying methods on polyphenol content and antioxidant activities of yuzu (*Citrus junos* Sieb ex Tanaka). *Journal of Food Measurement and Characterization*, 11(2), 576–585.
- Awad, M. A., Wagenmakers, P. S., & De Jager, A. (2001). Effects of light on flavonoid and Chlorogenic acid levels in the skin of “Jonagold” apples. *Scientia Horticulturae*, 88(4), 289–298.
- Azuma, A., Yakushiji, H., Koshita, Y., & Kobayashi, S. (2012). Flavonoid biosynthesis-related genes in grape skin are differentially regulated by temperature and light conditions. *Planta*, 236(4), 1067–1080.
- Baranović, G., & Šegota, S. (2017). Infrared spectroscopy of flavones and flavonols. Reexamination of the hydroxyl and carbonyl vibrations in relation to the interactions of flavonoids with membrane lipids. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 1–43.
- Barnes, C., Tibbitts, T., Sager, J., Deitzer, G., Bubenheim, D., Koerner, G., & Bugbee, B. (1993). Accuracy of quantum sensors measuring yield photon flux and photosynthetic photon flux. *HortScience*, 28(12), 1197–1200.
- Barnes, P. W., Flint, S. D., & Caldwell, M. M. (1987). Photosynthesis Damage and Protective Pigments in Plants from A Latitudinal Arctic/Alpine Gradient Exposed to Supplemental UV-B Radiation in the Field. *Arctic and Alpine Research*, 19(1), 21–27.
- Basahi, J. M., Ismail, I. M., & Hassan, I. A. (2014). Effects of Enhanced UV-B Radiation and Drought Stress on Photosynthetic Performance of Lettuce (*Lactuca sativa* L. Romaine) Plants. *Annual Research & Review in Biology*, 4(11), 1739–1756.
- Becker, C. (2014). Impact of radiation, temperature and growth stage on the concentration of flavonoid glycosides and caffeic acid derivatives in red leaf lettuce. University of Berlin.
- Beggs, C. J., Kuhn, K., Bocker, R., & Wellmann, E. (1987). Phytochrome-induced flavonoid biosynthesis in mustard (*Sinapis alba* L.) cotyledons. Enzymic control and differential regulation of anthocyanin and quercetin formation. *Planta*, 172(1), 121–126.
- Belgio, E., Trsková, E., Kotabová, E., Ewe, D., Prášil, O., & Kaňa, R. (2017). High light acclimation of *Chromera velia* points to photoprotective NPQ.

Photosynthesis Research, 135(1–3), 263–274.

- Bergquist, S. Å., Gertsson, U. E., Nordmark, L. Y., & Olsson, M. E. (2007). Effects of shade nettings, sowing time and storage on baby spinach flavonoids. *Journal of the Science of Food and Agriculture*, 87(13), 2464–2471.
- Bermúdez-cardona, M. B., Américo, J., Filho, W., & Rodrigues, F. Á. (2015). Leaf Gas Exchange and chlorophyll a Fluorescence in Maize Leaves Infected with *Stenocarpella macrospora*. *Biochemistry and Cell Biology*, 105(1), 26–34.
- Bertamini, M., Muthuchelian, K., & Nedunchezian, N. (2006). Shade effect alters leaf pigments and photosynthetic responses in Norway spruce (*Picea abies* L.) grown under field conditions. *Photosynthetica*, 44(2), 227–234.
- Bian, Z., Cheng, R., Wang, Y., Yang, Q., & Lu, C. (2018). Effect of green light on nitrate reduction and edible quality of hydroponically grown lettuce (*Lactuca sativa* L) under short-term continuous light from red and blue light-emitting diodes. *Environmental and Experimental Botany*, 153(2017), 63–71.
- Bienert, G. P., Schjoerring, J. K., & Jahn, T. P. (2006). Membrane transport of hydrogen peroxide. *Biochimica et Biophysica Acta (BBA) - Biomembranes*, 1758(8), 994–1003.
- Binggeli, P. (2005). *Clidemia hirta* (L.) D. Don. In *Crop Protection Compendium* (pp. 1–12).
- Biosciences, L. C. (2002). Using the LI-6400 portable photosynthesis system. Li-Cor Biosciences Inc.: Lincoln, NE.
- Borlongan, I. A. G., Gerung, G. S., Nishihara, G. N., & Terada, R. (2016). Light and temperature effects on photosynthetic activity of *Eucheuma denticulatum* and *Kappaphycus alvarezii* (brown and green color morphotypes) from Sulawesi Utara, Indonesia. *Phycological Research*, 65(1), 69–79.
- Borlongan, I. A., Nishihara, G. N., Shimada, S., & Terada, R. (2017). Effects of temperature and PAR on the photosynthesis of *Kappaphycus* sp. (*Solieriaceae Rhodophyta*) from Okinawa, Japan, at the northern limit of native *Kappaphycus* distribution in the western Pacific. *Phycologia*, 56(4), 444–453.
- Brainard, D. C., Bellinder, R. R., & DiTommaso, A. (2005). Effects of canopy shade on the morphology, phenology, and seed characteristics of Powell amaranth (*Amaranthus powellii*). *Weed Science*, 53(2), 175–186.
- Briskin, D. P., & Gawienowski, M. C. (2001). Differential effects of light and

nitrogen on production of hypericins and leaf glands in *Hypericum perforatum*. *Plant Physiology and Biochemistry*, 39(12), 1075–1081.

- Brodowska, K. M. (2017). Natural flavonoids: classification, potential role, and application of flavonoid analogues. *European Journal of Biological Research*, 7(2), 108–123.
- Brugnoli, E., & Bjorkman, O. (1992). Growth of cotton under continuous salinity stress: influence on allocation pattern, stomatal and non-stomatal components of photosynthesis and dissipation of excess light energy. *Planta*, 187(3), 335–347.
- Brunetti, C., Di Ferdinando, M., Fini, A., Pollastri, S., & Tattini, M. (2013). Flavonoids as antioxidants and developmental regulators: Relative significance in plants and humans. *International Journal of Molecular Sciences*, 14(2), 3540–3555.
- Bunce, J. A. (2016). Light dependence of carboxylation capacity for C3 photosynthesis models. *Photosynthetica*, 54(4), 484–490.
- CABI (2019). Invasive Species Compendium. Wallingford, UK: CAB International. www.cabi.org/isc.
- Cambie, R. C., & Ash, J. (1994). *Fijian Medicinal Plants*. CSIRO Publishing
- Carvalho, I. S., Cavaco, T., Carvalho, L. M., & Duque, P. (2010). Effect of photoperiod on flavonoid pathway activity in sweet potato (*Ipomoea batatas* (L.) Lam.) leaves. *Food Chemistry*, 118(2), 384–390.
- Cen, Y. P., & Borman, J. F. (1990). The Response of Bean Plants to UV-B Radiation Under Different Irradiances of Background Visible Light. *Journal of Experimental Botany*, 41(11), 1489–1495.
- Céspedes, C. L., Achnine, L., Lotina-Hennsen, B., Salazar, J. R., Gómez-Garibay, F., & Calderón, J. S. (2001). Inhibition of photophosphorylation and electron transport by flavonoids and biflavonoids from endemic *Tephrosia* spp. of Mexico. *Pesticide Biochemistry and Physiology*, 69(2), 63–76.
- Chai, T., & Wong, F. (2012). Antioxidant properties of aqueous extracts of *Selaginella willdenowii*. *Journal Of Medicinal Plants*, 6(7), 1289–1296.
- Chalker-Scott, L. (1999). Environmental significance of anthocyanins in plant stress responses. *Photochemistry and Photobiology*, 70(1), 1–9.
- Chan, E. W. C., Lim, Y. Y., Wong, L. F., Lianto, F. S., Wong, S. K., Lim, K. K., & Lim, T. Y. (2008). Antioxidant and tyrosinase inhibition properties of leaves

- and rhizomes of ginger species. *Food Chemistry*, 109(3), 477–483.
- Chang, C. C., Yang, M. H., & Wen, H. M. (2002). Estimation of Total Flavonoid Content in Propolis by Two Complementary Colorimetric Methods. *Journal of Food and Drug Analysis*, 10(3), 178–182.
- Chen, B., Zou, D., Zhu, M., & Yang, Y. (2017). Effects of CO₂ levels and light intensities on growth and amino acid contents in red seaweed *Gracilaria lemaneiformis*. *Aquaculture Research*, 48(6), 2683–2690.
- Chen, C., Luo, X., Jin, G., Cheng, Z., & Pan, X. (2017). Shading effect on survival, growth, and contents of secondary metabolites in micropropagated *Anoectochilus* plantlets. *Brazilian Journal of Botany*, 79(2), 1–9.
- Chen, Y. E., Zhang, C. M., Su, Y. Q., Ma, J., Zhang, Z. W., Yuan, M., & Yuan, S. (2016). Responses of photosystem II and antioxidative systems to high light and high temperature co-stress in wheat. *Environmental and Experimental Botany*, 135, 45–55.
- Cheng, A. X., Yu, M., Wang, G. G., Wu, T., Zhang, C., Journal, S., & October, N. (2013). Growth, morphology and biomass allocation in response to light gradient in five subtropical evergreen broadleaved tree seedlings. *Journal of Tropical Forest Science*, 25(4), 537–546.
- Cheng, G. X., Li, R. J., Wang, M., Huang, L. J., Khan, A., Ali, M., & Gong, Z. H. (2018). Variation in leaf color and combine effect of pigments on physiology and resistance to whitefly of pepper (*Capsicum annuum* L.). *Scientia Horticulturae*, 229(June 2017), 215–225.
- Cheng, S. J. (2016). *Shedding Light on Photosynthesis : The Impacts of Atmospheric Conditions and Plant Canopy Structure on Ecosystem Carbon Uptake*. University of Michigan.
- Chuyong, G. B., & Acidri, T. (2017). Light and moisture levels affect growth and physiological parameters differently in *Faidherbia albida* (Delile) A. Chev. seedlings. *Acta Physiologiae Plantarum*, 39(5), 1–6.
- Cockshull, K. E., Graves, C. J., & Cave, C. R. J. (1992). The influence of shading on yield of glasshouse tomatoes. *Journal of Horticultural Science*, 67(1), 11–24.
- Colombo, S. L., Pairoba, C. F., & Andreo, C. S. (1996). Inhibitory Effect of Shikimic Acid on PEP Carboxylase Activity. *Plant and Cell Physiology*, 37(6), 870–872.
- Cortell, J. M., & Kennedy, J. A. (2006). Effect of shading on accumulation of

- flavonoid compounds in (*Vitis vinifera* L.) Pinot noir fruit and extraction in a model system. *Journal of Agricultural and Food Chemistry*, 54(22), 8510–8520.
- Crozier, A. (2018). Micronutrient Information Center, Linus Pauling Institute, Oregon State University, Corvallis.
- Crupi, P., Pichierri, A., Basile, T., & Antonacci, D. (2013). Postharvest stilbenes and flavonoids enrichment of table grape cv Redglobe (*Vitis vinifera* L.) as affected by interactive UV-C exposure and storage conditions. *Food Chemistry*, 141(2), 802–808.
- Cullen, J. (2006). Practical Plant Identification. Cambridge: Cambridge university press.
- Dai, Y., Shen, Z., Liu, Y., Wang, L., Hannaway, D., & Lu, H. (2009). Effects of shade treatments on the photosynthetic capacity, chlorophyll fluorescence, and chlorophyll content of *Tetrastigma hemsleyanum* Diels et Gilg. *Environmental and Experimental Botany*, 65(2–3), 177–182.
- D'Andrea, G. (2015). Quercetin: a flavonol with multifaceted therapeutic applications. *Fitoterapia*, 106, 256–271.
- Dar, F.A., Jain, K., Jain, B., Modak, M. (2017). Preliminary Phytochemical Analysis and Characterization of Flavonoid Moiety from *Vitex negundo* Leaves Origin in Madhya Pradesh State by HPLC Study. *UK Journal of Pharmaceuticals and Biosciences*. 5, 60–65.
- Darko, E., Heydarizadeh, P., Schoefs, B., & Sabzalian, M. R. (2014). Photosynthesis under artificial light: the shift in primary and secondary metabolism. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 369(1640), 20130243–20130243.
- Dawiyah, R. Y. A., Yunus, A., Samanhudi, Y., & Widiyastuti, W. (2018). Shading and vermicompost effect on growth and flavonoid content of *Tapak Liman* (*Elephantopus scaber* L). In *IOP Conference Series: Earth and Environmental Science* (pp. 1–10).
- Day, M. D., Kawi, A., Kurika, K., Dewhurst, C. F., Waisale, S., Saul-Maora, J., & Senaratne, K. A. D. (2012). *Mikania micrantha* Kunth (Asteraceae) (Mile-a-Minute): Its Distribution and Physical and Socioeconomic Impacts in Papua New Guinea. *Pacific Science*, 66(2), 213–223.
- De Azevedo, C. H., & Rodriguez-Amaya, D. B. (2005). Carotenoid composition of

- kale as influenced by maturity, season and minimal processing. *Journal of the Science of Food and Agriculture*, 85(4), 591–597.
- De Nardo, T., Ian, C. S., Halim, Y., Francis, D., & Saona, L. E. R. (2015). Rapid and Simultaneous Determination of Lycopene and β -Carotene Contents in Tomato Juice by Infrared Spectroscopy. *Journal of Agricultural and Food Chemistry*, 57, 1105–1112.
- Debski, H., Wiczowski, W., Szawara-Nowak, D., Baczek, N., Szwed, M., & Horbowicz, M. (2017). Enhanced Light Intensity Increases Flavonol and Anthocyanin Concentrations but Reduces Flavone Levels in the Cotyledons of Common Buckwheat Seedlings. *Cereal Research Communications*, 45(2), 225–233.
- Del Valle, J. C., Buide, M. L., Casimiro-Soriguer, I., Whittall, J. B., & Narbona, E. (2015). On flavonoid accumulation in different plant parts: variation patterns among individuals and populations in the shore campion (*Silene littorea*). *Frontiers in Plant Science*, 6(939), 1–13.
- Demmig-Adams, B., Adams III, W. W., Barker, D. H., Logan, B. A., Bowling, D. R., & Verhoeven, A. S. (1996). Using chlorophyll fluorescence to assess the fraction of absorbed light allocated to thermal dissipation of excess excitation. *Physiologia Plantarum*.
- Deng, B., Shang, X., Fang, S., Li, Q., Fu, X., & Su, J. (2012). Integrated effects of light intensity and fertilization on growth and flavonoid accumulation in *Cyclocarya paliurus*. *Journal of Agricultural and Food Chemistry*, 60(25), 6286–6292.
- DeWalt, S. J., Denslow, J. S., & Hamrick, J. L. (2004). Biomass allocation, growth, and photosynthesis of genotypes from native and introduced ranges of the tropical shrub *Clidemia hirta*. *Oecologia*, 138(4), 521–531.
- Díez, M. C., Moreno, F., & Gantiva, E. (2017). Effects of light intensity on the morphology and CAM photosynthesis of *Vanilla planifolia* Andrews. *Revista Facultad Nacional de Agronomía*, 70(1), 8023.
- Distelbarth, H., Nagale, T., & Heyer, A. G. (2013). Responses of antioxidant enzymes to cold and high light are not correlated to freezing tolerance in natural accessions of *Arabidopsis thaliana*. *Plant Biology*, 1, 1–9.
- Dong, Y.-H., Beuning, L., Davies, K., Mitra, D., Morris, B., & Kootstra, A. (1998). Expression of pigmentation genes and photo-regulation of anthocyanin

- biosynthesis in developing Royal Gala apple flowers. *Australian Journal of Plant Physiology*, 25(2), 245.
- Dou, H., Niu, G., Gu, M., & Masabni, J. (2017). Effects of Light Quality on Growth and Phytonutrient Accumulation of Herbs under Controlled Environments. *Horticulturae*, 3(2), 36.
- Downey, M. O., Harvey, J. S., & Robinson, S. P. (2004). The effect of bunch shading on berry development and flavonoid accumulation in Shiraz grapes. *Australian Journal of Grape and Wine Research*, 10, 55–73.
- Edreva, A. (2005). The importance of non-photosynthetic pigments and cinnamic acid derivatives in photoprotection. *Agriculture, Ecosystems and Environment*, 106(2–3 SPEC. ISS.), 135–146.
- Ethier, G. J., & Livingston, N. J. (2004). On the need to incorporate sensitivity to CO₂ transfer conductance into the Farquhar-von Caemmerer-Berry leaf photosynthesis model. *Plant, Cell and Environment*, 27(2), 137–153.
- Ewa, B. U., Honda, C., Bessho, H., Kondo, S., Wada, M., Kobayashi, S., & Moriguchi, T. (2006). Expression analysis of anthocyanin biosynthetic genes in apple skin : Effect of UV-B and temperature. *Plant Science*, 170, 571–578.
- Félix, L. F., Rafael, C.-'Alvarez¹, & Iv'an, G. (2003). Relations between electron transport rates determined by pulse amplitude modulated chlorophyll fluorescence and oxygen evolution in macroalgae under different light conditions. *Photosynthesis Research*, 75, 259–275.
- Fan, X. X., Xu, Z. G., Liu, X. Y., Tang, C. M., Wang, L. W., & Han, X. lin. (2013). Effects of light intensity on the growth and leaf development of young tomato plants grown under a combination of red and blue light. *Scientia Horticulturae*, 153, 50–55.
- Fan, Y., Wang, Y., Tan, R., & Zhang, Z. (1998). Seasonal and sexual variety of *Ginkgo* flavonol glycosides in the leaves of *Ginkgo biloba* L. *Europe PMC*, 23(5), 267–269.
- Fanglian, H. (2011). Bradford Protein Assay. *Bio-Protocol*, 1(6), 4–5.
- Farquhar, G. D., von Caemmerer, S., & Berry, J. A. (1980). A biochemical model of photosynthetic CO₂ assimilation in leaves of C₃ species. *Planta*, 149(1), 78–90.
- Favory, J. J., Stec, A., Gruber, H., Rizzini, L., Oravec, A., Funk, M., & Ulm, R. (2009). Interaction of COP1 and UVR8 regulates UV-B-induced

- photomorphogenesis and stress acclimation in *Arabidopsis*. *The EMBO Journal*, 28(5), 591–601.
- Feng, F., Li, M., Ma, F., & Cheng, L. (2013). Phenylpropanoid metabolites and expression of key genes involved in anthocyanin biosynthesis in the shaded peel of apple fruit in response to sun exposure. *Plant Physiology and Biochemistry*, 69, 54–61.
- Feng, S., Wang, Y., Yang, S., Xu, Y., & Chen, X. (2010). Anthocyanin biosynthesis in pears is regulated by a R2R3-MYB transcription factor PyMYB10. *Planta*, 232(1), 245–255.
- Ferdinando, F. B., Adam, T., Stephen, F., Subhalakshmi, N., Ramaswamy, N., Lynne, A. S., & Jayant, K. (2010). Enzymatic Synthesis and Characterization of PolyQuercetin. *Journal of Macromolecular Science, Part A*, 47, 1191–1196.
- Ferdinando, M. Di, Brunetti, C., & Fini, A. (2012). Flavonoids as Antioxidants in Plants Under Abiotic Stresses. In P. Ahmad & M. N. V. Prasad (Eds.), *Abiotic Stress Responses in Plants: Metabolism, Productivity and Sustainability* (pp. 159–179). Springer.
- Fern, K., & Fern, A. (2014). Contact - Useful Tropical Plants. Retrieved September 28, 2018, from <http://tropical.theferns.info/contact.php>
- Fernando, W., Ratnasooriya, W., & Deraniyagala, S. (2009). Antinociceptive activity of aqueous leaf extract of *Tetracera sarmentosa* L. in rats. *Pharmacognosy Research*, 1(6), 381–386.
- Ferreres, F., Figueiredo, R., Bettencourt, S., Carqueijeiro, I., Oliveira, J., Gil-Izquierdo, A., & Sottomayor, M. (2011). Identification of phenolic compounds in isolated vacuoles of the medicinal plant *Catharanthus roseus* and their interaction with vacuolar class III peroxidase: An H₂O₂ affair? *Journal of Experimental Botany*, 62(8), 2841–2854.
- Ferreira, F., Lorena, M., Rius, S., & Casati, P. (2012). Flavonoids: biosynthesis, biological functions, and biotechnological applications. *Frontiers in Plant Science*, 3, 222.
- Fini, A., Brunetti, C., Ferdinando, M. Di, Ferrini, F., & Tattini, M. (2011). Stress-induced flavonoid biosynthesis and the antioxidant machinery of plants. *Plant Signaling and Behavior*, 6(5), 709–711.
- Fonseca, G. M., Rushing, J. W., Rajapakse, N. C., Thomas, R. L., & Riley, M. B.

- (2006). *Potential Implications of Medicinal Plant Production in Controlled Environments: The case of Feverfew (Tanacetum parthenium)*. *HortScience* (Vol. 41).
- Fonseca, J. M., Rushing, J. W., Rajapakse, N. C., Thomas, R. L., & Riley, M. B. (2006). Potential implications of medicinal plant production in controlled environments: The case of feverfew (*Tanacetum parthenium*). *HortScience*, *41*(3), 531–535.
- Fu, Y., Li, H. Y., Yu, J., Liu, H., Cao, Z. Y., Manukovsky, N. S., & Liu, H. (2017). Interaction effects of light intensity and nitrogen concentration on growth, photosynthetic characteristics and quality of lettuce (*Lactuca sativa* L. Var. youmaicai). *Scientia Horticulturae*, *214*, 51–57.
- Fujita, A., Goto-Yamamoto, N., Aramaki, I., & Hashizume, K. (2006). Organ-specific transcription of putative flavonol synthase genes of grapevine and effects of plant hormones and shading on flavonol biosynthesis in grape berry skins. *Bioscience, Biotechnology, and Biochemistry*, *70*(3), 632–8. Retrieved from
- Gao, K., Li, P., Watanabe, T., & Walter Helbling, E. (2008). Combined effects of ultraviolet radiation and temperature on morphology, photosynthesis, and DNA of *Arthrospira* (Spirulina) *Platensis* (Cyanophyta). *Journal of Phycology*, *44*(3), 777–786.
- Gardeli, A., & Pappas, C. (2017). Estimation of antioxidant activity of different mixed herbal infusions using attenuated total reflectance Fourier transform infrared spectroscopy and chemometrics, (January). <https://doi.org/10.9755/ejfa.2016-11-1670>.
- Gardner, K. H., & Correa, F. (2012). How plants see the invisible. *Plant Science*, *335*(6075), 1451–1452.
- Gauthier, P. P., Battle, M. O., Griffin, K. L., & Bender, M. L. (2018). Measurement of gross photosynthesis, respiration in the light, and mesophyll conductance using H₂¹⁸O labeling. *Plant Physiology*, (7), 1–49.
- Gavin, N. M., & Durako, M. J. (2012). Localization and antioxidant capacity of flavonoids in *Halophila johnsonii* in response to experimental light and salinity variation. *Journal of Experimental Marine Biology and Ecology*, *416–417*(April), 32–40.
- Gazolla, A. P., Maria, F., Marangoni, L., Curado, J., Nascimento, R. C., Claudio, L.,

- & Silva, F. G. (2017). The influence of light quality on phenolic acid and biflavonoid production in *Anacardium othonianum* Rizz . seedlings grown in vitro. *Australian Journal of Crop Science*, 11(05), 528–534.
- GBIF Secretariat. (2018). *Mikania micrantha* Kunth. Retrieved February 11, 2018, from <https://www.gbif.org/species/5398421>
- Ghale, B. (2013). Morphological trait difference, growth and ecophysiological performance of *Mikania micrantha* grown under contrasting light and nutrient regimes. Norwegian University of life science.
- Ghasemzadeh, A., & Ghasemzadeh, N. (2011). Flavonoids and phenolic acids : Role and biochemical activity in plants and human. *Journal of Medicinal Plants Research*, 5(31), 6697–6703.
- Ghasemzadeh, A., Jaafar, H. Z. E., & Rahmat, A. (2010). Synthesis of phenolics and flavonoids in ginger (*Zingiber officinale* Roscoe) and their effects on photosynthesis rate. *International Journal of Molecular Sciences*, 11(11), 4539–4555.
- Ghasemzadeh, A., Jaafar, H. Z. E., Rahmat, A., Wahab, P. E. M., & Halim, M. R. A. (2010). Effect of different light intensities on total phenolics and flavonoids synthesis and anti-oxidant activities in young ginger varieties (*Zingiber officinale* Roscoe). *International Journal of Molecular Sciences*, 11(10), 3885–3897.
- Gianmaria, M., Gianpaolo, G., & Stiina, K. (2018). The Effect of Light Spectrum on the Morphology and Cannabinoid Content of *Cannabis sativa* L. *Medical Cannabis and Cannabinoids*, 1(6), 19–27.
- Gonçalves, A. A., & Rocha, M. de O. C. (2017). Safety and Quality of Antimicrobial Packaging Applied to Seafood. *MOJ Food Processing & Technology*, 4(1), 1–7.
- Gould, K. S. (2004). Nature's Swiss Army Knife : The Diverse Protective Roles of Anthocyanins in Leaves. *Journal of Biomedicine and Biotechnology*, 5, 314–320.
- Graham, T. L. (1998). Flavonoid and flavonol glycoside metabolism in *Arabidopsis*. *Plant Physiology and Biochemistry*, 36(1–2), 135–144.
- Gregoriou, K., Pontikis, K., & Vemmos, S. (2007). Effects of reduced irradiance on leaf morphology, photosynthetic capacity, and fruit yield in olive (*Olea europaea* L). *Photosynthetica*, 45(2), 172–181.

- Gressel, J. (1979). Blue light photoreception. *Photochemistry and Photobiology*, 30(6), 749–754.
- Griffin, J. J., Ranney, T. G., & Pharr, D. M. (2004). Photosynthesis, chlorophyll fluorescence, and carbohydrate content of *Illicium taxa* grown under varied irradiance. *Journal of the American Society for Horticultural Science*, 129(1), 46–53.
- Grubby, C. (2014). Growing 101: The Basics of Plant Lighting. Retrieved December 19, 2018, from <https://www.maximumyield.com/growing-101-the-basics-of-plant-lighting/2/1246>.
- Gu, J., Zhou, Z., Li, Z., Chen, Y., Wang, Z., Zhang, H., & Yang, J. (2017). Photosynthetic Properties and Potentials for Improvement of Photosynthesis in Pale Green Leaf Rice under High Light Conditions. *Frontiers in Plant Science*, 8(June), 1–14.
- Gyu, T., Kim, N. D., & Hyun, S. (2017). Effects of light sources on major flavonoids and antioxidant activity in common buckwheat sprouts. *Food Science and Biotechnology*, 1–8.
- Hagen, S. F., Borge, G. I. A., Bengtsson, G. B., Bilger, W., Berge, A., Haffner, K., & Solhaug, K. A. (2007). Phenolic contents and other health and sensory related properties of apple fruit (*Malus domestica* Borkh., cv. Aroma): Effect of postharvest UV-B irradiation. *Postharvest Biology and Technology*, 45(1), 1–10.
- Hectors, K., van Oevelen, S., Guisez, Y., Prinsen, E., & Jansen, M. A. K. (2012). The phytohormone auxin is a component of the regulatory system that controls UV-mediated accumulation of flavonoids and UV-induced morphogenesis. *Physiologia Plantarum*, 145(4), 594–603.
- Heijde, M., & Ulm, R. (2012). UV-B photoreceptor-mediated signalling in plants. *Trends in Plant Science*, 17(4), 230–237. Retrieved from <https://www.sciencedirect.com/science/article/pii/S1360138512000088>
- Hemm, M. R., Rider, S. D., Ogas, J., Murry, D. J., & Chapple, C. (2004). Light induces phenylpropanoid metabolism in *Arabidopsis* roots. *The Plant Journal*, 38(5), 765–778.
- Heraut-Bron, V., Robin, C., Varlet-Grancher, C., Afif, D., & Guckert, A. (1999). Light quality (red:far-red ratio): does it affect photosynthetic activity, net CO₂ assimilation, and morphology of young white clover leaves? *Canadian*

Journal of Botany, 77(10), 1425–1431.

- Hogewoning, S. W., Trouwborst, G., Maljaars, H., Poorter, H., van Ieperen, W., & Harbinson, J. (2010). Blue light dose-responses of leaf photosynthesis, morphology, and chemical composition of *Cucumis sativus* grown under different combinations of red and blue light. *Journal of Experimental Botany*, 61(11), 3107–3117.
- Hollósy, F. (2002). Effects of ultraviolet radiation on plant cells. *Micron*, 33(2), 179–197.
- Hoque, E., & Remus, G. (1996). Reflective Light Properties of Tissue Layers in Beech (*Fagus sylvatica* L.) Leaves. *Photochemistry and Photobiology*, 63(4), 498–506.
- Hu, Y., Pan, Z. J., Liao, W., Li, J., Gruget, P., & Kitts, D. D. (2016). Determination of Antioxidant Capacity and Phenolic Content of Chocolate by Attenuated Total Reflectance-Fourier Transformed-Infrared Spectroscopy. *Food Chemistry*, 1–28.
- Hughes, M. N. (2009). *The photoprotective role of anthocyanin pigments in leaf tissues*. Wake Forest University.
- Iannotti, M. (2018). What Does Full Sun or Partial Shade Mean? Retrieved February 13, 2019, from <https://www.thespruce.com/what-is-full-sun-partial-shade-1402372>
- Ibrahim, M. H., & Jaafar, H. Z. E. (2012). Primary, Secondary Metabolites, H₂O₂, Malondialdehyde and Photosynthetic Responses of *Orthosiphon stamineus* Benth. to Different Irradiance Levels. *Molecules*, 17, 1159–1176.
- Ibrahim, M. H., Jaafar, H. Z. E., Karimi, E., & Ghasemzadeh, A. (2014). Allocation of secondary metabolites, photosynthetic capacity, and antioxidant activity of Kacip Fatimah (*Labisia pumila* Benth) in response to CO₂ and light intensity. *The Scientific World Journal*.
- Ibrahim, M. H., Jaafar, H. Z. E., Rahmat, A., & Rahman, Z. A. (2011). The relationship between phenolics and flavonoids production with total non structural carbohydrate and photosynthetic rate in *Labisia pumila* Benth. under high CO₂ and nitrogen fertilization. *Molecules*, 16(1), 162–174.
- Irshad, M., Debnath, B., Mitra, S., Arafat, Y., Li, M., Sun, Y., & Qiu, D. (2018). Accumulation of anthocyanin in callus cultures of red-pod okra (*Abelmoschus esculentus* (L.) Hongjiao) in response to light and nitrogen levels. *Plant*

Cell, Tissue and Organ Culture (PCTOC), 1–11.

- Ishak, A. H., Shafie, N. H., & Bahari, H. (2018). From Weed to Medicinal Plant : Antioxidant Capacities and Phytochemicals of Various Extracts of *Mikania micrantha*. *International Journal of Agriculture & Biology*, 20(3), 561–568.
- Ishida, A., Toma, T., & Marjenah. (1999). Leaf gas exchange and chlorophyll fluorescence in relation to leaf angle, azimuth, and canopy position in the tropical pioneer tree, *Macaranga conifera*. *Tree Physiology*, 19(2), 117–124.
- Ishikita, H., Saenger, W., & Biesiadka, J. (2006). How photosynthetic reaction centers control oxidation power in chlorophyll pairs P680, P700, and P870. *National Acad Sciences*, 103(26), 9855–9860.
- Islam, M. A., Tarkowská, D., Clarke, J. L., Blystad, D. R., Gislerød, H. R., Torre, S., & Olsen, J. E. (2014). Impact of end-of-day red and far-red light on plant morphology and hormone physiology of poinsettia. *Scientia Horticulturae*, 174(1), 77–86.
- Jaakola, L., & Hohtola, A. (2010). Effect of latitude on flavonoid biosynthesis in plants. *Plant, Cell and Environment*, 33(8), 1239–1247.
- Jaakola, L., Määttä-Riihinen, K., Kärenlampi, S., & Hohtola, A. (2004). Activation of flavonoid biosynthesis by solar radiation in bilberry (*Vaccinium myrtillus* L.) leaves. *Planta*, 218(5), 721–728.
- Jaakola, L., Poole, M., Jones, M. O., Kämäräinen-Karppinen, T., Koskimäki, J. J., Hohtola, A., & Seymour, G. B. (2010). A SQUAMOSA MADS Box Gene Involved in the Regulation of Anthocyanin Accumulation in Bilberry Fruits. *Plant Physiology*, 153, 1619–1629.
- Jacobs, M., & Rubery, P. H. (1988). Naturally Occurring Auxin Transport Regulators. *Science*, 241(4863), 346–349.
- Jang, Y., Mun, B., Seo, T., Lee, J., Oh, S., & Chun, C. (2013). Effects of light quality and intensity on the carbon dioxide exchange rate, growth, and morphogenesis of grafted pepper transplants during healing and acclimatization. *Korean Journal of Horticultural Science and Technology*, 31(1), 14–23.
- Jansen, M. A., van den Noort, R. E., Tan, M. Y., Prinsen, E., Lagrimini, L. M., & Thorneley, R. N. (2001). Phenol-oxidizing peroxidases contribute to the protection of plants from ultraviolet radiation stress. *Plant Physiology*, 126(3), 1012–23.

- Jeong, S. T., Goto-Yamamoto, N., Kobayashi, S., & Esaka, M. (2004). Effects of plant hormones and shading on the accumulation of anthocyanins and the expression of anthocyanin biosynthetic genes in grape berry skins. *Plant Science*, 167, 247–252.
- Jespersen, E., Brix, H., & Sorrell, B. K. (2017). Acclimation to light and avoidance of photoinhibition in *Typha latifolia* is associated with high photosynthetic capacity and xanthophyll pigment content. *Functional Plant Biology*, 44(8), 774–784.
- Jia, H.-J., Araki, A., & Okamoto, G. (2005). Influence of fruit bagging on aroma volatiles and skin coloration of ‘Hakuho’ peach (*Prunus persica* Batsch). *Postharvest Biology and Technology*, 35(1), 61–68.
- Jifon, J. L., & Syvertsen, J. P. (2003). Moderate shade can increase net gas exchange and reduce photoinhibition in citrus leaves. *Tree Physiology*, 23(2), 119–127.
- Juergeniemk, G., Boje, K., Huewel, S., Lohmann, C., Galla, H. J., & Nahrstedt, A. (2003). In Vitro studies indicate that miquelianin (quercetin 3-O-β-D-glucuronopyranoside) is able to reach the CNS from the small intestine. *Planta medica*, 69(11), 1013-1017.
- Jyothilakshmi, M., Jyothis, M., & Latha, M. S. (2015). Antidermatophytic Activity of *Mikania micrantha* Kunth: An Invasive Weed. *Pharmacognosy Research*, 7(1), S20-5.
- Kadomura-Ishikawa, Y., Miyawaki, K., Noji, S., & Takahashi, A. (2013). Phototropin 2 is involved in blue light-induced anthocyanin accumulation in *Fragaria ananassa* fruits. *Journal of Plant Research*, 126(6), 847–857.
- Kakani, V. G., Reddy, K. R., Zhao, D., & Mohammed, A. R. (2003). Effects of ultraviolet-B radiation on cotton (*Gossypium hirsutum* L.) morphology and anatomy. *Annals of Botany*, 91(7), 817–826.
- Kalaji, H. M., Schansker, G., Brestic, M., Bussotti, F., Calatayud, A., Ferroni, L., & Ferroni, L. (2017). Frequently asked questions about chlorophyll fluorescence , the sequel. *Photosynth Res*, 132, 13–66.
- Karimi, E., Jaafar, H. Z. E., Ghasemzadeh, A., & Ibrahim, M. H. (2013). Light intensity effects on production and antioxidant activity of flavonoids and phenolic compounds in leaves, stems and roots of three varieties of *Labisia pumila* benth. *Australian Journal of Crop Science*, 7(7), 1016–1023.
- Kato, M. C., Hikosaka, K., Hirotsu, N., Makino, A., & Hirose, T. (2003). The excess

light energy that is neither utilized in photosynthesis nor dissipated by photoprotective mechanisms determines the rate of photoinactivation in photosystem II. *Plant and Cell Physiology*, 44(3), 318–325.

- Kesarkar, S., Bhandage, A., Deshmukh, S., Shevkar, K., & Abhyankar, M. (2009). Flavonoids : An Overview. *Journal of Pharmacy Research*, 2(6), 1148–1154.
- Khan, S. R., Rose, R., Haase, D. L., & Sabin, T. E. (2000). Effects of shade on morphology, chlorophyll concentration, and chlorophyll fluorescence of four Pacific Northwest conifer species. *New Forests*, 19(2), 171–186.
- Khan, W., Prithiviraj, B., & Smith, D. L. (2003). Photosynthetic responses of corn and soybean to foliar application of salicylates. *Journal of Plant Physiology*, 160(5), 485–492.
- Khodary, S. E. A. (2004). Effect of Salicylic Acid on the Growth, Photosynthesis and Carbohydrate Metabolism in Salt Stressed Maize Plants. *International Journal of Agriculture & Biology*, 6(1), 1–4.
- Khoo, H. E., Azlan, A., Tang, S. T., & Lim, S. M. (2017). Anthocyanidins and anthocyanins: colored pigments as food, pharmaceutical ingredients, and the potential health benefits. *Food & Nutrition Research*, 61(1), 1361779.
- Kim, D. O., Jeong, S. W., & Lee, C. Y. (2003). Antioxidant capacity of phenolic phytochemicals from various cultivars of plums. *Food Chemistry*, 81(3), 321–326.
- King-Díaz, B., Granados-Pineda, J., Bah, M., Rivero-Cruz, J. F., & Lotina-Hennsen, B. (2015). Mexican propolis flavonoids affect photosynthesis and seedling growth. *Journal of Photochemistry and Photobiology B: Biology*, 151, 213–220.
- Kitajima, K., & Hogan, K. P. (2003). Increases of chlorophyll a/b ratios during acclimation of tropical woody seedlings to nitrogen limitation and high light. *Plant, Cell and Environment*, 26(6), 857–865.
- Kobayashi, S., Goto-Yamamoto, N., & Hirochika, H. (2004). Retrotransposon-induced mutations in grape skin color. *Science (New York, N.Y.)*, 304(5673), 982.
- Koes, R., Verweij, W., & Quattrocchio, F. (2005). Flavonoids: a colorful model for the regulation and evolution of biochemical pathways. *Trends in Plant Science*, 10(5), 236–242. <https://doi.org/10.1016/J.TPLANTS.2005.03.002>
- Kokalj Ladan, M., Straus, J., Tavčar Benković, E., & Kreft, S. (2017). FT-IR-based

- method for rutin, quercetin and quercitrin quantification in different buckwheat (*Fagopyrum*) species. *Scientific Reports*, 7(1), 7226.
- Kolb, C. A., Kaser, M. A., Kopecky, J., & Zotz, G. (2001). Effects of natural intensities of visible and ultraviolet radiation on. *Plant Physiology*, 127(3), 863–875.
- Koller, D. (1990). Light-driven leaf movements. *Plant, Cell and Environment*, 13(7), 615–632.
- Kong, J. M., Chia, L. S., Goh, N. K., Chia, T. F., & Brouillard, R. (2003). Analysis and biological activities of anthocyanins. *Phytochemistry*, 64(5), 923–933.
- Koyama, K., Ikeda, H., Poudel, P. R., & Goto-Yamamoto, N. (2012). Light quality affects flavonoid biosynthesis in young berries of Cabernet Sauvignon grape. *Phytochemistry*, 78, 54–64.
- Kumar, S., & Pandey, A. K. (2013). Chemistry and biological activities of Flavonoids: An Overview. *The Scientific World Journal*, 16(12), 1–17.
- Kumari, R., Singh, S., & Agrawal, S. B. (2009). Effects of supplemental ultraviolet-b radiation on growth and physiology of *Acorus calamus* L. (sweet flag). *Acta Biologica Cracoviensis Series Botanica*, 51(2), 19–27.
- Kurata, H., Matsumura, S., & Furusaki, S. (1997). Light irradiation causes physiological and metabolic changes for purine alkaloid production by a *Coffea arabica* cell suspension culture. *Plant Science*, 123(1–2), 197–203.
- Lambers, H., Chapin, F. S., & Pons, T. L. (2008). *Plant Physiological Ecology* (2nd ed.). Springer New York.
- Lariguet, P., & Dunand, C. (2005). Plant Photoreceptors: Phylogenetic Overview. *Journal of Molecular Evolution*, 61(4), 559–569.
- Lätti, A. K., Riihinen, K. R., & Kainulainen, P. S. (2008). Analysis of Anthocyanin Variation in Wild Populations of Bilberry (*Vaccinium myrtillus* L.) in Finland. *Journal of Agricultural and Food Chemistry*, 56(1), 190–196.
- Lee, J. (2005). Determination of Total Monomeric Anthocyanin Pigment Content of Fruit Juices, Beverages, Natural Colorants, and Wines by the pH Differential Method: Collaborative Study. *Journal Of AOAC International*, 88(5), 1269–1278.
- Leng, P., Shuchai, S., Tianhua, W., Xiangning, J., & Shasheng, W. (2002). Effects of light intensity and light quality on photosynthesis, flavonol glycoside and terpene lactone contents of *Ginkgo biloba* L. seedlings. *Journal of Plant*

- Resources and Environment*, 11(1), 1–4. Retrieved from
- Lesne, A. (2008). Robustness: confronting lessons from physics and biology. *Biological Reviews*, 83(4), 509–32.
- Lev-yadun, S., & Gould, K. S. (2009). Role of Anthocyanins in Plant Defence. In Anthocyanins (pp. 21–48). Springer New York.
- Li, A., Li, S., Wu, X., Zhang, J., He, A., Zhao, G., & Yang, X. (2016). Effect of Light Intensity on Leaf Photosynthetic Characteristics and Accumulation of Flavonoids in *Lithocarpus litseifolius* (Hance) Chun. (Fagaceae). *Open Journal of Forestry*, 06(05), 445–459. <https://doi.org/10.4236/ojf.2016.65034>
- Li, C. X., Chang, S. X., Khalil-Ur-Rehman, M., Xu, Z. G., & Tao, J. M. (2017). Effect of irradiating the leaf abaxial surface with supplemental light-emitting diode lights on grape photosynthesis. *Australian Journal of Grape and Wine Research*, 23(1), 58–65.
- Li, H., Jiang, D., Wollenweber, B., Dai, T., & Cao, W. (2010). Effects of shading on morphology, physiology and grain yield of winter wheat. *European Journal of Agronomy*, 33(4), 267–275.
- Li, J., Li, G., Wang, H., & Wang Deng, X. (2011). Phytochrome Signaling Mechanisms. *The Arabidopsis Book*, 9, e0148.
- Li, Q., Deng, M., Xiong, Y., Coombes, A., & Zhao, W. (2014). Morphological and photosynthetic response to high and low irradiance of *Aeschynanthus longicaulis*. *Scientific World Journal*, 2014, 1–9.
- Li, Q., & Kubota, C. (2009). Effects of supplemental light quality on growth and phytochemicals of baby leaf lettuce. *Environmental and Experimental Botany*, 67, 59–64.
- Li, Y., Xin, G., Wei, M., Shi, Q., Yang, F., & Wang, X. (2017). Carbohydrate accumulation and sucrose metabolism responses in tomato seedling leaves when subjected to different light qualities. *Scientia Horticulturae*, 225(7), 490–497.
- Liang, N., Lu, X., Hu, Y., & Kitts, D. (2016). Application of Attenuated Total Reflectance-Fourier Transformed Infrared (ATR-FTIR) Spectroscopy to Determine Chlorogenic acid Isomer Profile and Antioxidant Capacity of Coffee Beans Antioxidant Capacity of Coffee Beans. *Journal of Agricultural and Food Chemistry*, 1–38.
- Lichtenthaler, H. K., & Buschmann, C. (2001). chlorophylls and Carotenoids :

- Measurement and Characterization by UV-VIS. In *Current Protocols in Food Analytical Chemistry* (pp. 1–8).
- Lima, C. C., Lemos, R. P. L., & Conserva, L. M. (2014). Dilleniaceae family: an overview of its ethnomedicinal uses, biological and phytochemical profile. *Journal of Pharmacognosy and Phytochemistry*, 3(2), 181–204.
- Lingakumar, K., & Kulandaivelu, G. (1993). Changes induced by ultraviolet-B radiation in vegetative growth, foliar characteristics and photosynthetic activities in *Vigna unguiculata*. *Functional Plant Biology*, 20(3), 299–308.
- Liu, C., Guo, C., Wang, Y., & Ouyang, F. (2002). Effect of light irradiation on hairy root growth and artemisinin biosynthesis of *Artemisia annua* L. *Process Biochemistry*, 38(4), 581–585.
- Liu, H., Fu, Y., Hu, D., Yu, J., & Liu, H. (2018). Effect of green, yellow and purple radiation on biomass, photosynthesis, morphology and soluble sugar content of leafy lettuce via spectral wavebands “knock out.” *Scientia Horticulturae*, 236(2), 10–17.
- Liu, L., Gitz, D. C., & McClure, J. W. (1995). Effects of UV- B on flavonoids, ferulic acid, growth and photosynthesis in barley primary leaves. *Physiologia Plantarum*, 93(4), 725–733.
- Liu, S. L., Ma, M. D., Pan, Y. Z., Wei, L. L., He, C. X., & Yang, K. M. (2013). Effects of light regimes on photosynthetic characteristics and antioxidant system in seedlings of two alder species. *Chinese Journal of Plant Ecology*, 36(10), 1062–1074.
- Liu, Y., Fang, S., Yang, W., Shang, X., & Fu, X. (2018). Light quality affects flavonoid production and related gene expression in *Cyclocarya paliurus*. *Journal of Photochemistry & Photobiology, B: Biology*, 179(11), 66–73.
- Liu, Y., Qian, C., Ding, S., Shang, X., Yang, W., & Fang, S. (2016). Effect of light regime and provenance on leaf characteristics, growth and flavonoid accumulation in *Cyclocarya paliurus* (Batal) Iljinskaja coppices. *Botanical Studies*, 57(1), 1–13.
- Lobo, F. de A., de Barros, M. P., Dalmagro, H. J., Dalmolin, Â. C., Pereira, W. E., de Souza, É. C., & Rodríguez Ortíz, C. E. (2013). Fitting net photosynthetic light-response curves with Microsoft Excel - a critical look at the models. *Photosynthetica*, 51(3), 445–456.
- Lombard, K., Peffley, E. B., Lombard, K. A., Geoffriau, E., & Peffley, E. (2002).

- Flavonoid quantification in onion by spectrophotometric and high performance liquid Chromatography analysis. *HortScience*, 37(4), 682–685
- Lopez, T., Corbin, C., Falguieres, A., Doussot, J., Montguillon, J., Hagège, D., & Lainé, É. (2016). Secondary metabolite accumulation, antibacterial and antioxidant properties of in vitro propagated *Clidemia hirta* L. extracts are influenced by the basal culture medium. *Comptes Rendus Chimie*, 19(9), 1071–1076.
- Løvdal, T., Olsen, K. M., Slimestad, R., Verheul, M., & Lillo, C. (2010). Synergetic effects of nitrogen depletion, temperature, and light on the content of phenolic compounds and gene expression in leaves of tomato. *Phytochemistry*, 71(5–6), 605–613.
- Lu, T., Meng, Z., Zhang, G., Qi, M., Sun, Z., Liu, Y., & Li, T. (2017). Sub-high Temperature and High Light Intensity Induced Irreversible Inhibition on Photosynthesis System of Tomato Plant (*Solanum lycopersicum* L.). *Frontiers in Plant Science*, 08(March), 1–16.
- Lu, X., Ross, C. F., Powers, J. R., & Rasco, B. A. (2011a). Determination of Quercetins in Onion (*Allium cepa*) Using Infrared Spectroscopy. *Journal of Agricultural and Food Chemistry*, 59, 6376–6382.
- Lu, X., Ross, C. F., Powers, J. R., Aston, D. E., & Rasco, B. A. (2011b). Determination of Total Phenolic Content and Antioxidant Activity of Garlic (*Allium sativum*) and Elephant Garlic (*Allium ampeloprasum*) by Attenuated Total Reflectance À Fourier Transformed Infrared Spectroscopy. *Journal of Agricultural and Food Chemistry*, 59, 5215–5221.
- Ma, X., Song, L., Yu, W., Hu, Y., Liu, Y., Wu, J., & Ying, Y. (2015). Growth, physiological, and biochemical responses of *Camptotheca acuminata* seedlings to different light environments. *Frontiers in Plant Science*, 6, 321.
- Ma, Z., Li, S., & Zhang, M. (2010). Light Intensity Affects Growth, Photosynthetic Capability, and Total Flavonoid Accumulation of *Anoectochilus* Plants Light Intensity Affects Growth, Photosynthetic Capability, and Total Flavonoid Accumulation of *Anoectochilus* Plants. *HortScience*, 45(6), 863–867.
- Maciej, H., Maria, K., Dorota, N., & Anna, K. (2001). Infrared spectrum analysis of some flavonoids. *Acta Poloniae Pharmaceutica-Drug Research*, 58(6), 415–420.
- Manetas, Y. (2006). Why some leaves are anthocyanic and why most anthocyanic

leaves are red? *Flora: Morphology, Distribution, Functional Ecology of Plants*, 201(3), 163–177.

Manivannan, A., Soundararajan, P., Park, Y. G., Wei, H., Kim, S. H., & Jeong, B. R. (2017). Blue and red light-emitting diodes improve the growth and physiology of in vitro-grown carnations ‘Green Beauty’ and ‘Purple Beauty.’ *Horticulture Environment and Biotechnology*, 58(1), 12–20.

Marshall, B., & Biscoe, P. V. (1980). A Model for C₃ Leaves Describing the Dependence of Net Photosynthesis on Irradiance. *Journal of Experimental Botany*, 31(1), 29–39.

Martín, J., Navas, M. J., Moreno, M. J., & Asuero, G. A. (2017). Anthocyanin Pigments: Importance, Sample Anthocyanin Pigments: Preparation and Extraction. In *Phenolic Compounds - Natural Sources, Importance and Applications* (pp. 117–152). InTech.

Martínez Pastur, G., Lencinas, M. V., Peri, P. L., & Arena, M. (2007). Photosynthetic plasticity of *Nothofagus pumilio* seedlings to light intensity and soil moisture. *Forest Ecology and Management*, 243(2–3), 274–282.

Martínez-Lüscher, J., Torres, N., Hilbert, G., Richard, T., Sánchez-Díaz, M., Delrot, S., & Gomès, E. (2014). Ultraviolet-B radiation modifies the quantitative and qualitative profile of flavonoids and amino acids in grape berries. *Phytochemistry*, 102, 106–114.

Marques, G. S., Leão, W. F., Lyra, M. A. M., Peixoto, M. S., Monteiro, R. P. M., Rolim, L. A., ... Soares, L. A. L. (2013). Comparative evaluation of UV/VIS and HPLC analytical methodologies applied for quantification of flavonoids from leaves of *Bauhinia forficata*. *Brazilian Journal of Pharmacognosy*, 23(1), 51–57.

Matus, J. T., Loyola, R., Vega, A., Peña-Neira, A., Bordeu, E., Arce-Johnson, P., & Alcalde, J. A. (2009). Post-veraison sunlight exposure induces MYB-mediated transcriptional regulation of anthocyanin and flavonol synthesis in berry skins of *Vitis vinifera*. *Journal of Experimental Botany*, 60(3), 853–867.

Maxwell, K., & Johnson, G. N. (2000). chlorophyll fluorescence — a practical guide. *Journal of Experimental Botany*, 51(345), 659–668.

McCree, K. J. (1971). The action spectrum, absorptance and quantum yield of photosynthesis in crop plants. *Agricultural Meteorology*, 9, 191–216.

- Meng, X., Xing, T., & Wang, X. (2004). The role of light in the regulation of anthocyanin accumulation in *Gerbera hybrida*. *Plant Growth Regulation*, *44*, 243–250.
- Meyer, S., Cerovic, Z. G., Goulas, Y., Montpied, P., Demotes-Mainard, S., Bidel, L. P. R., & Dreyer, E. (2006). Relationships between optically assessed polyphenols and chlorophyll contents, and leaf mass per area ratio in woody plants: A signature of the carbon-nitrogen balance within leaves? *Plant, Cell and Environment*, *29*(7), 1338–1348.
- Michelina, C., Ferdinando, P., Flavia, B., Simona, P., Sabina, M., Paola, N., & Severina, P. (2015). Silica/quercetin sol-gel hybrids as antioxidant dental implant materials. *Science and Technology of Advanced Materials*, *16*, 1–11.
- Mierziak, J., Kostyn, K., & Kulma, A. (2014). Flavonoids as important molecules of plant interactions with the environment. *Molecules*, *19*(10), 16240–16265.
- Migné, A., Delebecq, G., Davoult, D., Spilmont, N., Menu, D., Janquin, M., & Menu, D. (2018). Photosynthesis of *Laminaria digitata* over the immersion and emersion alternation of spring tides under sunny and hot weather (No. hal-01827565).
- Mishra, A., Sharma, A. K., Kumar, S., Saxena, A. K., & Pandey, A. K. (2013). *Bauhinia variegata* leaf extracts exhibit considerable antibacterial, antioxidant, and anticancer activities. *BioMed Research International*, *2013*, 1.
- Miyamoto, K. (1997). Renewable biological systems for alternative sustainable energy production. Japan: Food & Agriculture Org.
- Mizuno, T., Amaki, W., & Watanabe, H. (2011). Effects of monochromatic light irradiation by led on the growth and anthocyanin contents in leaves of cabbage seedlings. *Acta Horticulturae*, (907), 179–184.
- Moe, R., Morgan, L., & Grindal, G. (2002). Growth and plant morphology of *Cucumis sativus* and *Fuchsia hybrid* are influenced by light quality during the photoperiod and by diurnal temperature alternations. *Acta Horticulturae*, *580*, 229–234.
- Moghaddam, S. S., Ibrahim, R., Damalas, C. A., & Noorhosseini, S. A. (2017). Effects of Gamma Stress and Carbon Dioxide on Eight Bioactive Flavonoids and Photosynthetic Efficiency in *Centella asiatica*. *Journal of Plant Growth Regulation*, *36*(4), 957–969.

- Mohd, N. H. W. S. W., Ahmad, F., & Heng Yen, K. (2015). Chemical constituents from *Piper caninum* and antibacterial activity. *Journal of Applied Pharmaceutical Science*, 5(06), 20-025.
- Momayyezi, M., & Guy, R. D. (2017). Blue light differentially represses mesophyll conductance in high vs low latitude genotypes of *Populus trichocarpa* Torr. *Journal of Plant Physiology*, 213, 122–128.
- Morales-Flores, F., Olivares-Palomares, K. S., Aguilar-Laurents, M. I., Rivero-Cruz, J. F., Lotina-Hennsen, B., & King-Díaz, B. (2015). Flavonoids Affect the Light Reaction of Photosynthesis in Vitro and in Vivo as Well as the Growth of Plants. *Journal of Agricultural and Food Chemistry*, 63(37), 8106–8115.
- Mori, K., Goto-Yamamoto, N., Kitayama, M., & Hashizume, K. (2007). Loss of anthocyanins in red-wine grape under high temperature. *Journal of Experimental Botany*, 58(8), 1935–1945.
- Mori, K., Sugaya, S., & Gemma, H. (2005). Decreased anthocyanin biosynthesis in grape berries grown under elevated night temperature condition. *Scientia Horticulturae*, 105(3), 319–330.
- Mosaleeyanon, K., Zobayed, S. M. A., Afreen, F., & Kozai, T. (2005). Relationships between net photosynthetic rate and secondary metabolite contents in St. John's wort. *Plant Science*, 169(3), 523–531.
- Moutinho-Pereira, J. M., Bacelar, E. A., Goncalves, B., Ferreira, H. F., & Correia, C. M. (2010). Effects of Open-Top Chambers on physiological and yield attributes of field grown grapevines. *Acta Physiologiae Plantarum*, 32, 395–403.
- Müller, V., Albert, A., Barbro Winkler, J., Lankes, C., Noga, G., & Hunsche, M. (2013). Ecologically relevant UV-B dose combined with high PAR intensity distinctly affect plant growth and accumulation of secondary metabolites in leaves of *Centella asiatica* L. Urban. *Journal of Photochemistry and Photobiology B: Biology*, 127, 161–169.
- Müller, V., Lankes, C., Schmitz-Eiberger, M., Noga, G., & Hunsche, M. (2013). Estimation of flavonoid and centelloside accumulation in leaves of *Centella asiatica* L. Urban by multiparametric fluorescence measurements. *Environmental and Experimental Botany*, 93, 27–34.
- Muneer, S., Kim, E. J., Park, J. S., & Lee, J. H. (2014). Influence of green, red and blue light emitting diodes on multiprotein complex proteins and

- photosynthetic activity under different light intensities in lettuce leaves (*Lactuca sativa* L.). *International Journal of Molecular Sciences*, 15(3), 4657–4670.
- Murchie, E. H., & Horton, P. (1997). Acclimation of photosynthesis to irradiance and spectral quality in British plant species: chlorophyll content, photosynthetic capacity and habitat preference. *Plant, Cell and Environment*, 20(4), 438–448.
- Nam, T. G., Jin, Y., Seok, L., & Eom, H. (2018). Flavonoid accumulation in common buckwheat (*Fagopyrum esculentum*) sprout tissues in response to light. *Horticulture, Environment, and Biotechnology*, 1–9.
- Narbona, E., Buide, M. L., Casimiro-Soriguer, I., & Valle, J. C. Del. (2014). Polimorfismos de color floral: causas e implicaciones evolutivas. *Revista Ecosistemas*, 23(3), 36–47.
- Nawkar, G., Maibam, P., Park, J., Sahi, V., Lee, S., Kang, C., Kang, C. H. (2013). UV-Induced Cell Death in Plants. *International Journal of Molecular Sciences*, 14(1), 1608–1628.
- Neill, S. O. (2002). *The functional role of anthocyanins in leaves*. PhD Thesis. University of Auckland.
- Ngoc, T., Aung, A., Naing, H., & Kil, C. (2016). Influences of different light sources and light / dark cycles on anthocyanin accumulation and plant growth in Petunia. *Journal of Plant Biotechnology*, 43, 119–124.
- Nguyen, H. T., & Goycoolea, F. M. (2017). Chitosan/Cyclodextrin/TPP Nanoparticles Loaded with Quercetin as Novel Bacterial Quorum Sensing Inhibitors. *Molecules*, 22(1975), 1–23.
- Nicotra, A. B., Chazdon, R. L., & Schlichting, C. D. (1997). Patterns of genotypic variation and phenotypic plasticity of light response in two tropical Piper (Piperaceae) species. *American Journal of Botany*, 84(11), 1542–1552.
- Niglas, A., Papp, K., Sękiewicz, M., & Sellin, A. (2017). Short-term effects of light quality on leaf gas exchange and hydraulic properties of silver birch (*Betula pendula*). *Tree Physiology*, 37(9), 1218–1228.
- Nishiyama, Y., Suleyman, I. A., & Murata, N. (2006). A new paradigm for the action of reactive oxygen species in the photoinhibition of photosystem II. *Biochimica et Biophysica Acta (BBA)-Bioenergetics*, 1757(7), 742–749.
- Niu, S. S., Xu, C. J., Zhang, W. S., Zhang, B., Li, X., Lin-Wang, K., & Chen, K. S.

- (2010). Coordinated regulation of anthocyanin biosynthesis in Chinese bayberry (*Myrica rubra*) fruit by a R2R3 MYB transcription factor. *Planta*, 231(4), 887–899.
- Oliveira, R. N., Mancini, M. C., Cabral, F., Oliveira, S. De, Passos, T. M., & Quilty, B. (2016). FTIR analysis and quantification of phenols and flavonoids of five commercially available plants extracts used in wound healing Análise por FTIR e quantificação de fenóis e flavonóides de cinco produtos naturais disponíveis comercialmente utilizados no tr. *Revista Materia*, 21(3), 767–779.
- Olle, M., & Viršilė, A. (2013). The effects of light-emitting diode lighting on greenhouse plant growth and quality. *Agricultural and Food Science*, 22, 223–234.
- Oshita, K., Suzuki, T., & Kawono, T. (2018). Possible Roles of Near-infrared Light on the Photosynthesis in *Synechocystis* sp. PCC6803 under Solar Simulating Artificial Light. *Environment Control in Biology*, 56(1), 17–21.
- Osmond, C. B. (1994). What is photoinhibition? Some insights from comparisons of shade and sun plants. *Photoinhibition of photosynthesis: from molecular mechanisms to the field* (pp. 1–24).
- Özer, H. (2017). Effects of shading and organic fertilizers on tomato yield and quality. *Pak Journal of Botany*, 49(5), 1849–1855.
- Pacheco, F. V., Alvarenga, I. C. A., Junior, P. M. R., Pinto, J. E. B. P., Avelar, R. de P., & Alvarenga, A. A. (2014). Growth and production of secondary compounds in monkey-pepper (*Piper aduncum* L.) leaves cultivated under altered ambient light. *Australian Journal of Crop Science*, 8(11), 1510–1516.
- Pala, U. C., & Toklucu, K. A. (2011). Effect of UV-C light on anthocyanin content and other quality parameters of pomegranate juice. *Journal of Food Composition and Analysis*, 24, 790–795.
- Palapol, Y., Ketsa, S., Lin-Wang, K., Ferguson, I. B., & Allan, A. C. (2009). A MYB transcription factor regulates anthocyanin biosynthesis in mangosteen (*Garcinia mangostana* L.) fruit during ripening. *Planta*, 229(6), 1323–1334.
- Pan, J., & Guo, B. (2016). Effects of light intensity on the growth, photosynthetic characteristics, and flavonoid content of *Epimedium pseudowushanense* B.L.Guo. *Molecules*, 21(11), 1–12.
- Panche, A. N., Diwan, A. D., & Chandra, S. R. (2016). Flavonoids: An overview. *Journal of Nutritional Science*, 5(e47), 1–15.

- Park, H. R., Daun, Y., Park, J. K., & Bark, K. M. (2013). Spectroscopic properties of flavonoids in various aqueous-organic solvent mixtures. *Bulletin of Korean Chemical Society*, *34*(1), 211–220.
- Pauly, G., & Moretti, C. (2003). Use of plant extracts with an anti-radical-type action. US Grant. Retrieved from
- Pedroso, R. C. N., Branquinho, N. A. A., Hara, A. C. B. A. M., Costa, A. C., Silva, F. G., Pimenta, L. P., & Januario, A. H. (2017). Impact of light quality on flavonoid production and growth of *Hyptis marrubioides* seedlings cultivated in vitro. *Brazilian Journal of Pharmacognosy*, *27*(4), 466–470.
- Pego, J. V., Kortstee, A. J., Huijser, C., & Smeekens, S. C. (2000). Photosynthesis, sugars and the regulation of gene expression. *Journal of Experimental Botany*, *51*, 407–16. Retrieved from
- Peltzer, D., & Polle, A. (2001). Diurnal fluctuations of antioxidative systems in leaves of field-grown beech trees (*Fagus sylvatica*): Responses to light and temperature. *Physiologia Plantarum*, *111*(2), 158–164.
- Pereira, G. E., Gaudillere, J. P., Pieri, P., Hilbert, G., Maucourt, M., Deborde, C., & Rolin, D. (2006). Microclimate Influence on Mineral and Metabolic Profiles of Grape Berries. *Journal of Agricultural and Food Chemistry*, *54*(18), 6765–6775.
- Pérez-López, U., Sgherri, C., Miranda-Apodaca, J., Micaelli, F., Lacuesta, M., Mena-Petite, A., & Muñoz-Rueda, A. (2018). Concentration of phenolic compounds is increased in lettuce grown under high light intensity and elevated CO₂. *Plant Physiology and Biochemistry*, *123*, 233–241.
- Peri, P. L., Lucas, R. J., & Moot, D. J. (2007). Dry matter production, morphology and nutritive value of *Dactylis glomerata* growing under different light regimes. *Agroforestry Systems*, *70*(1), 63–79.
- Peters, H. A. (2001). *Clidemia hirta* Invasion at the Pasoh Forest Reserve: An Unexpected Plant Invasion in an Undisturbed Tropical Forest¹. *Biotropica*, *33*(1), 60–68.
- Philpoteducation. (2018). Essential ideas. Retrieved October 6, 2018, from https://www.philpoteducation.com/pluginfile.php/1136/mod_book/chapter/1131/2.1.9b.jpg
- Pietrini, F., Iannelli, M. A., & Massacci, A. (2002). Anthocyanin accumulation in the illuminated surface of maize leaves enhances protection from photo-

- inhibitory risks at low temperature, without further limitation to photosynthesis. *Plant, Cell and Environment*, 25(10), 1251–1259.
- Pintér, J., Kósa, E., Hadi, G., Hegyi, Z., Spitzkó, T., Tóth, Z., & Marton, L. C. (2007). Effect of increased UV-B radiation on the anthocyanin content of maize (*Zea mays* L.) leaves. *Acta Agronomica Hungarica*, 55(1), 7–17.
- Proestos, C., & Komaitis, M. (2013). Analysis of Naturally Occurring Phenolic Compounds in Aromatic Plants by RP-HPLC Coupled to Diode Array Detector (DAD) and GC-MS after Silylation. *Foods*, 2, 90–99.
- Pudasaini, A., & Zoltowski, B. D. (2013). Zeitelupe senses blue-light fluence to mediate circadian timing in *Arabidopsis thaliana*. *Biochemistry*, 52(40), 7150–7158.
- Putri, D. P., Widyastuti, Y., Dewi, W. S., & Yunus, A. (2018). The effect of shade and vermicompost application on yield and flavonoid levels of Tempuyung (*Sonchus arvensis*). In *IOP Conf. Series: Earth and Environmental Science* (pp. 1–12).
- Qin, Z., Mao, D. J., Quan, G. M., Zhang, J., Xie, J. F., & DiTommaso, A. (2012). Physiological and morphological responses of invasive *Ambrosia artemisiifolia* (common ragweed) to different irradiances. *Botany*, 90(12), 1284–1294.
- Rabara, R. C., Behrman, G., Timbol, T., & Rushton, P. J. (2017). Effect of Spectral Quality of Monochromatic LED Lights on the Growth of Artichoke Seedlings. *Frontiers in Plant Science*, 8(2), 1–9.
- Rabinowicz, P. D., Braun, E. L., Wolfe, A. D., Bowen, B., & Grotewold, E. (1999). Maize R2R3 Myb Genes: Sequence Analysis Reveals Amplification in the Higher Plants. *Genetics*, 153(1), 427–444.
- Rajni, G., 2000. Photosynthesis and the Z-scheme. University of Illinois at Urbana-Champaign.
- Ramakrishna, A., & Ravishankar, G. A. (2011). Influence of abiotic stress signals on secondary metabolites in plants. *Plant Signaling and Behavior*, 6(11), 1720–1731.
- Ramawat, K. G., Dass, S., & Mathur, M. (2009). The Chemical Diversity of Bioactive Molecules and Therapeutic Potential of Medicinal Plants. In *Herbal Drugs: Ethnomedicine to Modern Medicine* (pp. 7–32). Berlin, Heidelberg: Springer Berlin Heidelberg.

- Ravaglia, D., Espley, R. V., Henry-Kirk, R. A., Andreotti, C., Ziosi, V., Hellens, R. P., & Allan, A. C. (2013). Transcriptional regulation of flavonoid biosynthesis in nectarine (*Prunus persica*) by a set of R2R3 MYB transcription factors. *BMC Plant Biology*, *13*(1), 68. <https://doi.org/10.1186/1471-2229-13-68>
- Raven, P., Evert, R., & Eichhorn, S. (2005). *Photosynthesis, Light, and Life. Biology of Plants* (Vol. 7).
- Ravidran, N., & Nirmal, K. (2005). *Babu. Ginger: the genus Zingiber*.
- Re, G. A., Piluzzaa, G., Sanna, F., Molinub, M. G., & Sulasa, L. (2018). Polyphenolic composition and antioxidant capacity of legume based swards are affected by light intensity in a Mediterranean agroforestry system Running. *Jsfa*, 1–25.
- Rehana, S., Ahmed, F., Zeba, N., Husna, A., & Hossain, F. (2018). Effect of sunlight and artificial light on micropropagation of potato (*Solanum tuberosum* L) plantlets. *Archives of Agriculture and Environmental Science*, *3*(2), 151–156.
- Ruiz, J. M., Rivero, R. M., López-Cantarero, I., & Romero, L. (2003). Role of Ca²⁺ in the metabolism of phenolic compounds in tobacco leaves (*Nicotiana tabacum* L.). *Plant Growth Regulation*, *41*(2), 173–177.
- Sakhonwasee, S., Tummachai, K., & Nimnoy, N. (2017). Influences of LED Light Quality and Intensity on Stomatal Behavior of Three Petunia Cultivars Grown in a Semi-closed System. *Environment Control in Biology*, *55*(2), 93–103.
- Sakihama, Y., & Yamasaki, H. (2002). Lipid Peroxidation Induced by Phenolics in Conjunction with Aluminum Ions. *Biologia Plantarum*, *45*(2), 249–254.
- Salvatierra, A., Pimentel, P., Moya-Leon, M. A., Caligari, P. D. S., & Herrera, R. (2010). Comparison of transcriptional profiles of flavonoid genes and anthocyanin contents during fruit development of two botanical forms of *Fragaria chiloensis* ssp. *chiloensis*. *Phytochemistry*, *71*(16), 1839–1847.
- Särkkä, L. E., Jokinen, K., Ottosen, C. O., & Kaukoranta, T. (2017). Effects of HPS and LED lighting on cucumber leaf photosynthesis, light quality penetration and temperature in the canopy, plant morphology and yield. *Agricultural and Food Science*, *26*(2), 101–109.
- Senthilkumar, S. R., Thirumal, S., Arulmozhi, K. T., & Mythili, N. (2018). FT-IR analysis and correlation studies on the antioxidant activity, total phenolics and

total flavonoids of Indian commercial teas (*Camellia sinensis* L .) - A novel approach FT-IR IR analysis and correlation studies on the antioxidant activity, total p. *International Research Journal of Biological Sciences*, 6(3), 1–7.

Sevik, H., Cetin, M., Kapucu, O., Aricak, B., Canturk, U. (2017). Effects of light on morphologic and stomatal characteristics of Turkish fir needles (*Abies nordmanniana* subsp. *Bornmulleriana* Mattf.). *Fresenius Environmental Bulletin*, 26(10), 6579–6587.

Shao, Q., Wang, H., Guo, H., Zhou, A., Huang, Y., Sun, Y., & Li, M. (2014). Effects of shade treatments on photosynthetic characteristics, chloroplast ultrastructure, and physiology of *Anoectochilus roxburghii*. *PLoS ONE*, 9(2), 1.

Sharkey, T. D. (2016). What gas exchange data can tell us about photosynthesis. *Plant Cell and Environment*, 39(6), 1161–1163.

Sharkey, T. D., Bernacchi, C. J., Farquhar, G. D., & Singsaas, E. L. (2007). Fitting photosynthetic carbon dioxide response curves for C3 leaves. *Plant, Cell and Environment*, 30(9), 1035–1040.

Shen, H., Hong, L., Ye, W., Cao, H., & Wang, Z. (2007). The influence of the holoparasitic plant *Cuscuta campestris* on the growth and photosynthesis of its host *Mikania micrantha*. *Journal of Experimental Botany*, 58(11), 2929–2937.

Shivashankara, K., Mithila, J., & Maiti, S. (2000). Effect of different light intensities on growth and yield of betel vine (*Piper betle* L.). *Journal of Plantation Crops*, 28(3), 196–200.

Shourie, A., Tomar, P., Srivastava, D., & Chauhan, R. (2014). Enhanced Biosynthesis of Quercetin Occurs as A Photoprotective Measure in *Lycopersicon esculentum* Mill . under Acute UV-B Exposure. *Brazilian Archives of Biology and Technology*, 57(3), 317–325.

Shui-Yuan, C., Xu, F., & Wang, Y. (2009). Advances in the study of flavonoids in *Ginkgo biloba* leaves. *Journal of Medicinal Plants Research*, 3(13), 1248–1252.

Silva, S. T., Bertolucci, S. K. V., da Cunha, S. H. B., Lazzarini, L. E. S., Tavares, M. C., & Pinto, J. E. B. P. (2017). Effect of light and natural ventilation systems on the growth parameters and carvacrol content in the in vitro cultures of *Plectranthus amboinicus* (Lour.) Spreng. *Plant Cell, Tissue and Organ*

Culture, 129(3), 501–510.

- Singh, A. S., Jones, A. M. P., Shukla, M. R., & Saxena, P. K. (2017). High light intensity stress as the limiting factor in micropropagation of sugar maple (*Acer saccharum* Marsh.). *Plant Cell, Tissue and Organ Culture (PCTOC)*, 129(2), 209–221.
- Singh, G. (2016). *Plant Systematics, Third Edition*. (G. Singh, Ed.). Science Publishers.
- Slattery, R. A., VanLoocke, A., Bernacchi, C. J., Zhu, X. G., & Ort, D. R. (2017). Photosynthesis, Light Use Efficiency, and Yield of Reduced-chlorophyll Soybean Mutants in Field Conditions. *Frontiers in Plant Science*, 8(4), 1–19.
- Smith, J., Burritt, D. J., & Bannister, P. (2000). Shoot Dry Weight, chlorophyll and UV-B-absorbing Compounds as Indicators of a Plant's Sensitivity to UV-B Radiation. *Annals of Botany*, 86(6), 1057–1063.
- Sousa Paiva, É. A., Dos Santos Isaias, R. M., Aguiar Vale, F. H., & De Senna Queiroz, C. G. (2003). The influence of light intensity on anatomical structure and pigment contents of *Tradescantia pallida* (Rose) Hunt. cv. purpurea boom (commelinaceae) leaves. *Brazilian Archives of Biology and Technology*, 46(4), 617–624.
- Sozmen, A. B., Erhan, C., Sozmen, E. Y., & Ovez, B. (2018). Author's Accepted Manuscript. *Biocatalysis and Agricultural Biotechnology*, 1–27.
- Sumanta, N., Haque, C. I., Nishika, J., & Suprakash, R. (2014). Spectrophotometric Analysis of chlorophylls and Carotenoids from Commonly Grown Fern Species by Using Various Extracting Solvents. *Research Journal of Chemical Sciences*, 4(9), 63–69.
- Stanton, K. M., Weeks, S. S., Dana, M. N., & Mickelbart, M. V. (2010). Light exposure and shade effects on growth, flowering, and leaf morphology of *Spiraea alba* du roi and *Spiraea tomentosa* L. *HortScience*, 45(12), 1912–1916.
- Steindal, A. L. H., Mølmann, J., Bengtsson, G. B., & Johansen, T. J. (2013). Influence of Day Length and Temperature on the Content of Health-Related Compounds in Broccoli (*Brassica oleracea* L. var. *italica*). *Journal of Agricultural and Food Chemistry*, 61(45), 10779–10786.
- Stutte, G. W., & Edney, S. (2009). Photoregulation of Bioprotectant Content of Red Leaf Lettuce with Light-emitting Diodes. *HortScience*, 44(1), 79–82.

- Su, W. H., Zhang, G. F., Li, X. H., Gu, F. X., & Shi, B. L. (2006). Effect of light intensity and light quality on growth and total flavonoid accumulation of *Erigeron breviscapus*. *Chinese Traditional and Herbal Drugs*.
- Sun, Y., Qian, M., Wu, R., Niu, Q., Teng, Y., & Zhang, D. (2014). Postharvest pigmentation in red Chinese sand pears (*Pyrus pyrifolia* Nakai) in response to optimum light and temperature. *Postharvest Biology and Technology*, *91*, 64–71.
- Szopa, A., Starzec, A., & Ekiert, H. (2018). The importance of monochromatic lights in the production of phenolic acids and flavonoids in shoot cultures of *Aronia melanocarpa*, *Aronia arbutifolia* and *Aronia prunifolia*. *Journal of Photochemistry & Photobiology, B: Biology*, 1–23.
- Szymańska, R., Ślesak, I., Orzechowska, A., & Kruk, J. (2017). Physiological and biochemical responses to high light and temperature stress in plants. *Environmental and Experimental Botany*, *139*, 165–177.
- Takos, A. M., Jaffe, F. W., Jacob, S. R., Bogs, J., Robinson, S. P., & Walker, A. R. (2006). Light-Induced Expression of a MYB Gene Regulates Anthocyanin Biosynthesis in Red Apples. *Plant Physiology*, *142*(3), 1216–1232.
- Tang, H., Hu, Y. Y., Yu, W.-W., Song, L. L., & Wu, J. S. (2015). Growth, photosynthetic and physiological responses of *Torreya grandis* seedlings to varied light environments. *Trees*, *29*(4), 1011–1022.
- Tattini, M., Galardi, C., Pinelli, P., Massai, R., Remorini, D., & Agati, G. (2004). Differential accumulation of flavonoids and hydroxycinnamates in leaves of *Ligustrum vulgare* under excess light and drought stress. *New Phytologist*, *163*(3), 547–561.
- Tattini, M., Guidi, L., Morassi-Bonzi, L., Pinelli, P., Remorini, D., Degl’Innocenti, E., & Agati, G. (2005). On the role of flavonoids in the integrated mechanisms of response of *Ligustrum vulgare* and *Phillyrea latifolia* to high solar radiation. *New Phytologist*, *167*(2), 457–470.
- Taulavuori, K., Pyysalo, A., & Taulavuori, E. (2018). Responses of phenolic acid and flavonoid synthesis to blue and blue-violet light depends on plant species. *Environmental and Experimental Botany*, 1–18.
- Tegelberg, R. (2002). Impact of elevated ultraviolet-B radiation on three northern deciduous woody plants. University of Joensuu.
- Teramura, A. H., & Sullivan, J. H. (1994). Effects of UV-B radiation on

- photosynthesis and growth of terrestrial plants. *Photosynthesis Research*, 39(3), 463–473.
- Thrane, J. E., Kyle, M., Striebel, M., Haande, S., Grung, M., Rohrlack, T., & Andersen, T. (2015). Spectrophotometric analysis of pigments: A critical assessment of a high-throughput method for analysis of algal pigment mixtures by spectral deconvolution. *PLoS ONE*, 10(9), 1–24.
- Thwe, A. A., Vercambre, G., Gautier, H., Gay, F., & Phattaralerphong, J. (2014). Response of photosynthesis and chlorophyll fluorescence to acute ozone stress in tomato (*Solanum lycopersicum* Mill.). *Photosynthetica*, 52(1), 105–116.
- Tian-gen, C., Chang-peng, X., Ming-nan, Q., Hong-long, Z., Qing-feng, S., & Xing-guang, Z. (2017). Evaluation of Protocols for Measuring Leaf Photosynthetic Properties of Field-Grown Rice. *Rice Science*, 24(1), 1–9.
- Tianzi, C., & Baolong, Z. (2016). Measurements of Proline and Malondialdehyde Content and Antioxidant Enzyme Activities in Leaves of Drought Stressed Cotton. *Bio-Protocol*, 6(17), 1–14.
- Tinoco-Ojanguren, C., & Pearcy, R. W. (1992). Dynamic stomatal behavior and its role in carbon gain during lightflecks of a gap phase and an understory *Piper* species acclimated to high and low light. *Oecologia*, 92(2), 222–228.
- Tourism Malaysia (2014). Gunung Ledang. Retrieved 25 May 2014.
- Treutter, D. (2005). Significance of flavonoids in plant resistance and enhancement of their biosynthesis. *Plant Biology*, 7(6), 581–591.
- Treutter, D. (2006). Significance of flavonoids in plant resistance : a review, 147–157.
- Triantaphylides, C., Krischke, M., Hoerberichts, F. A., Ksas, B., Gresser, G., Havaux, M., & Mueller, M. J. (2008). Singlet Oxygen Is the Major Reactive Oxygen Species Involved in Photooxidative Damage to Plants. *Plant Physiology*, 148(2), 960–968.
- Triantaphylidès, C., & Havaux, M. (2009). Singlet oxygen in plants: production, detoxification and signaling. *Trends in Plant Science*, 14(4), 219–228.
- Trojak, M., & Skowron, E. (2017). Role of anthocyanins in high-light stress response. *World Scientific News*, 81(2), 150–168.
- Tsurunaga, Y., Takahashi, T., Katsube, T., Kudo, A., Kuramitsu, O., Ishiwata, M., & Matsumoto, S. (2013). Effects of UV-B irradiation on the levels of

- anthocyanin, rutin and radical scavenging activity of buckwheat sprouts. *Food Chemistry*, 141(1), 552–556.
- Tu, J., Zhang, Z., Cui, C., Yang, M., Li, Y., & Zhang, Y. (2017). Fast Separation and Determination of Flavonoids in Honey Samples by Capillary Zone Electrophoresis. *Kem. Ind*, 66(3–4), 129–134.
- Turturica, M., Oancea, A. M., Rapeanu, G., & Bahrim, G. (2015). Anthocyanins: naturally occurring fruit pigments with functional properties mihaela. *The Annals of the University Dunărea de Jos of Galați Fascicle VI – Food Technology*, 39(1), 9–24.
- Uddin, M. M., Islam, M. A., Hosen, M. T., Alam, M. S., Alam, M. N., Faruk, M., & Uddin, S. B. (2017). Estimation of in vivo neuropharmacological and in vitro antioxidant effects of *Tetracera sarmentosa*. *Cogent Biology*, 3(1), 1–10.
- Uddin, M. M. N., Kabir, M. S. H., Hasan, M., Al Mahmud, Z., Alam Bhuiya, N. M. M., Ahmed, F., & Alam, M. S. (2018). Assessment of the antioxidant, thrombolytic, analgesic, anti-inflammatory, antidepressant and anxiolytic activities of leaf extracts and fractions of *Tetracera sarmentosa* (L.) Vahl. *Journal of Basic and Clinical Physiology and Pharmacology*, 29(1), 81–93.
- Uleberg, E., Rohloff, J., Jaakola, L., Tröst, K., Junttila, O., Häggman, H., & Martinussen, I. (2012). Effects of Temperature and Photoperiod on Yield and Chemical Composition of Northern and Southern Clones of Bilberry (*Vaccinium myrtillus* L.). *Journal of Agricultural and Food Chemistry*, 60(42), 10406–10414.
- Violet-Chabrand, S., Matthews, J. S. A., Simkin, A. J., Raines, C. A., & Lawson, T. (2017). Importance of Fluctuations in Light on Plant Photosynthetic Acclimation. *Plant Physiology*, 173(4), 2163–2179.
- Vimolmangkang, S., Zheng, D., Han, Y., Khan, M. A., Soria-Guerra, R. E., & Korban, S. S. (2014). Transcriptome analysis of the exocarp of apple fruit identifies light-induced genes involved in red color pigmentation. *Gene*, 534(1), 78–87.
- Wada, M., Kagawa, T., & Sato, Y. (2003). Chloroplast movement. *Annual Review of Plant Biology*, 54(1), 455–468.
- Wan Nurhayati, W. H., Norli Arlizan, T., & Nurdiana, S. (2013). Effect of *Mikania micrantha* leaf extract on the level of blood glucose and hepatic glycogen in the normal and alloxan-induced diabetic rats. *Npajj*, 9(10), 398–402.

- Wang, C. L., Guo, Q. S., Zhu, Z. B., & Cheng, B. X. (2017). Physiological characteristics, dry matter, and active component accumulation patterns of *Changium smyrnioides* in response to a light intensity gradient. *Pharmaceutical Biology*, 55(1), 581–589.
- Wang, C. Y., Chen, C. T., & Wang, S. Y. (2009). Changes of flavonoid content and antioxidant capacity in blueberries after illumination with UV-C. *Food Chemistry*, 117(3), 426–431.
- Wang, H., Gui, M., Tian, X., Xin, X., Wang, T., & Li, J. (2017). Effects of UV-B on vitamin C, phenolics, flavonoids and their related enzyme activities in mung bean sprouts (*Vigna radiata*). *International Journal of Food Science and Technology*, 52(3), 827–833.
- Wang, J., Lu, W., Tong, Y., & Yang, Q. (2016). Leaf Morphology, Photosynthetic Performance, chlorophyll Fluorescence, Stomatal Development of Lettuce (*Lactuca sativa* L.) Exposed to Different Ratios of Red Light to Blue Light. *Frontiers in Plant Science*, 7(250), 1–10.
- Wang, M. L., Jiang, Y. S., Wei, J. Q., Wei, X., Qi, X. X., Jiang, S. Y., & Wang, Z. M. (2007). Effects of irradiance on growth, photosynthetic characteristics, and artemisinin content of *Artemisia annua* L. *Photosynthetica*, 46(1), 17–20.
- Wang, S. Y., Chen, C. T., & Wang, C. Y. (2009). The influence of light and maturity on fruit quality and flavonoid content of red raspberries. *Food Chemistry*, 112(3), 676–684.
- Wang, Y., Zhou, B., Sun, M., Li, Y., & Kawabata, S. (2012). UV-A light induces anthocyanin biosynthesis in a manner distinct from synergistic blue + UV-B light and UV-A/blue light responses in different parts of the hypocotyls in turnip seedlings. *Plant and Cell Physiology*, 53(8), 1470–1480.
- Wilson, K. E., Thompson, J. E., Huner, N. P. A., & Greenberg, B. M. (2001). Effects of Ultraviolet-A Exposure on Ultraviolet-B-induced Accumulation of Specific Flavonoids in *Brassica napus*. *Photochemistry and Photobiology*, 73(6), 678–684.
- Winkel-Shirley, B. (2002). Biosynthesis of flavonoids and effects of stress. *Current Opinion in Plant Biology*, 5(3), 218–223.
- Winter, K., & Virgo, A. (1998). Elevated CO₂ enhances growth in the rain forest understory plant, *Piper cordulatum*, at extremely low light intensities. *Flora Jena*, 193(3), 323–326.

- Wu, G., Liu, H., Hua, L., Luo, Q., Lin, Y., He, P., & Ye, Q. (2018). Differential Responses of Stomata and Photosynthesis to Elevated Temperature in Two Co-occurring Subtropical Forest Tree Species. *Frontiers in Plant Science*, 9(April), 1–8.
- Wu, H. C., & Lin, C. C. (2012). Red Light-emitting Diode Light Irradiation Improves Root and Leaf Formation in Difficult-to-propagate *Protea cynaroides* L. Plantlets In Vitro. *HortScience* (Vol. 47). American Society for Horticultural Science. Retrieved from <http://hortsci.ashspublications.org/content/47/10/1490.short>
- Wulandari, L., Retnaningtyas, Y., & Lukman, H. (2016). Analysis of Flavonoid in Medicinal Plant Extract Using Infrared Spectroscopy and Chemometrics. *Journal of Analytical Methods in Chemistry*, 2016, 1–6.
- XiaoYing, L., ShiRong, G., ZhiGang, X., XueLei, J., & Tezuka, T. (2011). Regulation of chloroplast ultrastructure, cross-section anatomy of leaves, and morphology of stomata of cherry tomato by different light irradiations of light-emitting diodes. *HortScience*, 46(2), 217–221.
- Xie, B. D., & Wang, H. T. (2006). Effects of light spectrum and photoperiod on contents of flavonoid and terpene in leaves of *Ginkgo biloba* L. *Journal of Nanjing Forestry University*, 30, 51–54.
- Xu, Y., Wang, G., Cao, F., Zhu, C., Wang, G., & El-Kassaby, Y. A. (2014). Light intensity affects the growth and flavonol biosynthesis of Ginkgo (*Ginkgo biloba* L.). *New Forests*, 45(6), 765–776.
- Xue, W., Li, X. Y., Zhu, J. T., Lin, L. S., & Wang, Y. J. (2011). Effects of shading on leaf morphology and response characteristics of photosynthesis in *Alhagi sparsifolia*. *Chinese Journal of Plant Ecology*, 35(1), 82–90.
- Yabuzaki, J. (2017). Carotenoids Database: structures, chemical fingerprints and distribution among organisms. *Database*, 2017(1).
- Yamane, T., Jeong, S. T., Goto-Yamamoto, N., Koshita, Y., & Kobayashi, S. (2006). Effects of Temperature on Anthocyanin Biosynthesis in Grape Berry Skins. *American Journal of Enology and Viticulture*, 57, 54–59.
- Yang, F., Song, L., Wang, H., Wang, J., Xu, Z., & Xing, N. (2015). Quercetin in prostate cancer: Chemotherapeutic and chemopreventive effects, mechanisms and clinical application potential. *Oncology reports*, 33(6), 2659–2668.
- Yang, W., Liu, Y., Fang, S., Ding, H., Zhou, M., & Shang, X. (2017). Variation in

- growth, photosynthesis and water-soluble polysaccharide of *Cyclocarya paliurus* under different light regimes. *IForest*, 10(2), 468–474.
- Yao, X. yang, Liu, X. ying, Xu, Z. gang, & Jiao, X. lei. (2017). Effects of light intensity on leaf microstructure and growth of rape seedlings cultivated under a combination of red and blue LEDs. *Journal of Integrative Agriculture*, 16(1), 97–105.
- Ye, S., Shao, Q., Xu, M., Li, S., Wu, M., Tan, X., & Su, L. (2017). Effects of Light Quality on Morphology, Enzyme Activities, and Bioactive Compound Contents in *Anoectochilus roxburghii*. *Frontiers in Plant Science*, 8(5), 1–7.
- Yu, L. Y. (2015). Evaluation of the anti-oxidant and anti-aging properties of extracts from four types of plants. Universiti Sains Malaysia.
- Yu, W., Liu, Y., Song, L., Jacobs, D. F., Du, X., Ying, Y., & Wu, J. (2016). Effect of Differential Light Quality on Morphology, Photosynthesis, and Antioxidant Enzyme Activity in *Camptotheca acuminata* Seedlings. *Journal of Plant Growth Regulation*, 36(1), 148–160.
- Yuan, M., Jia, X., Ding, C., Zeng, H., Du, L., Yuan, S., & Liu, J. (2015). Effect of fluorescence light on phenolic compounds and antioxidant activities of soybeans (*Glycine max* L. Merrill) during germination. *Food Science and Biotechnology*, 24(5), 1859–1865.
- Zandi, K., Teoh, B. T., Sam, S. S., Wong, P. F., Mustafa, M., & AbuBakar, S. (2011). Antiviral activity of four types of bioflavonoid against dengue virus type-2. *Virology Journal*, 8(1), 560.
- Zhang, L. L., & Wen, D. Z. (2009). Responses of photosynthetic parameters of *Mikania micrantha* and *Chromolaena odorata* to contrasting irradiance and soil moisture. *Journal of Plant Resources*, 122, 69–79.
- Zhang, L. Y., Ye, W. H., Cao, H. L., & Feng, H. L. (2004). *Mikania micrantha* H.B.K. in China - an overview. *Weed Research*, 44(1), 42–49.
- Zhang, Q., Liu, M., & Ruan, J. (2017). Metabolomics analysis reveals the metabolic and functional roles of flavonoids in light-sensitive tea leaves. *BMC Plant Biology*, 17(1), 1–10.
- Zhang, X., C Allan, A., Yi, Q., Chen, L., Li, K., Shu, Q., & Su, J. (2011). Differential Gene Expression Analysis of Yunnan Red Pear, *Pyrus Pyrifolia*, During Fruit Skin Coloration. *Plant Molecular Biology Reporter*, 29(2), 305–314.

- Zhang, Z., & Rongfeng, H. (2013). Analysis of Malondialdehyde, chlorophyll Proline, Soluble Sugar, and Glutathione Content in *Arabidopsis* seedling. *Bio-Protocol*, 3(14), 1–8.
- Zhao, D., Maoyin, L., Jianmin, X., & Zhe, T. (1999). Effects of light on cell growth and flavonoids biosynthesis in callus cultures of *Saussurea medusa* Maxim. *Acta phytophysiologica Sinica* (Vol. 25). Shanghai ke xue ji shu chu ban she. Retrieved from <https://europepmc.org/abstract/cba/321524>
- Zhen, S., & van Iersel, M. W. (2017). Far-red light is needed for efficient photochemistry and photosynthesis. *Journal of Plant Physiology*, 209, 115–122.
- Zheng, L., & Van Labeke, M.-C. (2017). Long-Term Effects of Red- and Blue-Light Emitting Diodes on Leaf Anatomy and Photosynthetic Efficiency of Three Ornamental Pot Plants. *Frontiers in Plant Science*, 8(5), 1–12.
- ZhiMin, Y., ShaoJian, Z., AiTong, H., YouFei, Z., & JingYi, Y. (2000). Response of cucumber plants to increased UV-B radiation under water stress. *Journal of Environmental Sciences*, 12(2), 236–240.
- Zhong, J., Seki, T., Kinoshita, S., & Yoshida, T. (1991). Effect of light irradiation on anthocyanin production by suspended culture of *Perilla frutescens*. *Biotechnology and Bioengineering*, 38(6), 653–658.
- Zhou, R., Su, W. H., Zhang, G. F., Zhang, Y. N., & Guo, X. R. (2016). Relationship between flavonoids and photoprotection in shade-developed *Erigeron breviscapus* transferred to sunlight. *Photosynthetica*, 54(2), 201–209.
- Zhou, Y., & Singh, B. R. (2004). Effect of Light on Anthocyanin Levels in Submerged, Harvested Cranberry Fruit. *Journal of Biomedicine and Biotechnology*, 5, 259–263.
- Zhu, H., Li, X., Zhai, W., Liu, Y., Gao, Q., Liu, J., & Zhu, Y. (2017). Effects of low light on photosynthetic properties, antioxidant enzyme activity, and anthocyanin accumulation in purple pak-choi (*Brassica campestris* ssp. *Chinensis* Makino). *PLoS ONE*, 12(6), 1–17.
- Zhu, W., Jia, Q., Wang, Y., Zhang, Y., & Xia, M. (2012). The anthocyanin cyanidin-3-O- β -glucoside, a flavonoid, increases hepatic glutathione synthesis and protects hepatocytes against reactive oxygen species during hyperglycemia: Involvement of a cAMP–PKA-dependent signaling pathway. *Free Radical Biology and Medicine*, 52(2), 314–327.

- Zlatev, Z. S., J. C. Lidon, F., & Kaimakanova, M. (2012). Plant physiological responses to UV-B radiation. *Emirates Journal of Food and Agriculture*, 24(6), 481–501.
- Zoratti, L., Karppinen, K., Luengo, E., HÅggman, H., & Jaakola, L. (2014). Light-controlled flavonoid biosynthesis in fruits. *Frontiers in Plant Science*, 5(October), 1–17. <https://doi.org/10.3389/fpls.2014.00534>
- Zoratti, L., Karppinen, K., Luengo Escobar, A., HÅggman, H., & Jaakola, L. (2014). Light-controlled flavonoid biosynthesis in fruits. *Frontiers in Plant Science*, 5, 534.

