

Pervasiveness of biological impacts of artificial light at night

Kevin J. Gaston¹, Simone Ackermann, Jonathan Bennie, Daniel T.C. Cox, Benjamin B. Phillips, Alejandro Sánchez de Miguel and Dirk Sanders

Environment & Sustainability Institute, University of Exeter, Penryn, Cornwall, TR10 9FE, U.K.

¹E-mail: k.j.gaston@exeter.ac.uk

Synopsis Artificial light at night (ALAN) and its associated biological impacts have regularly been characterised as predominantly urban issues. Although far from trivial, this would imply that these impacts only affect ecosystems that are already heavily modified by humans and are relatively limited in their spatial extent, at least as compared with some key anthropogenic pressures on the environment that attract much more scientific and public attention, such as climate change or plastic pollution. However, there are a number of reasons to believe that ALAN and its impacts are more pervasive, and therefore need to be viewed from a broader geographic perspective rather than an essentially urban one. Here we address, in turn, 11 key issues when considering the degree of spatial pervasiveness of the biological impacts of ALAN. First, the global extent of ALAN is likely itself commonly underestimated, as a consequence of limitations of available remote sensing data sources and how these are processed. Second and third, more isolated (rural) and mobile (e.g., vehicle headlight) sources of ALAN may have both very widespread and important biological influences. Fourth and fifth, the occurrence and impacts of ALAN in marine systems and other remote settings, need much greater consideration. Sixth, seventh and eighth, there is growing evidence for important biological impacts of ALAN at low light levels, from skyglow, and over long distances (because of the altitudes from which it may be viewed by some organisms), all of which would increase the areas over which impacts are occurring. Ninth and tenth, ALAN may exert indirect biological effects that may further expand these areas, because it has a landscape ecology (modifying movement and dispersal and so hence with effects beyond the direct extent of ALAN), and because ALAN interacts with other anthropogenic pressures on the

environment. Finally, ALAN is not stable, but increasing rapidly in global extent, and shifting towards wavelengths of light that often have greater biological impacts.

Introduction

There has long been recognition that the introduction of artificial light into the nighttime environment has biological impacts. Early observations particularly highlighted the delayed retention of leaves on trees close to, and the attraction of insects and birds toward, outdoor light sources (e.g., Allen 1880; Gastman 1886; Matzke 1936; Schroeder 1945; Cochran and Graber 1958; Verheijen 1960). Nonetheless, it is only recently that artificial light at night (ALAN) has been regarded as a significant anthropogenic environmental pressure (Longcore and Rich 2004; Rich and Longcore 2006; Hölker et al. 2010; Gaston et al. 2014; Gaston 2018; Owens et al. 2020). This development has particularly been spurred by two things. First, satellite observations of the Earth at night have highlighted the widespread geographic occurrence of direct ALAN emissions, detected as the vertically emitted or reflected component (Sullivan 1989; Román et al. 2018; Levin et al. 2020), and this has been further emphasized by modelling of the extent of skyglow (indirect ALAN; artificial brightening of the night sky that results predominantly from upwardly emitted artificial light being scattered in the atmosphere by water, dust, and gas molecules; Cinzano et al. 2001; Falchi et al. 2016).

Second, a rapidly growing body of studies has documented many ways in which the disruption of natural light regimes by ALAN has biological impacts. These include at the levels of the individual (e.g., physiology, behaviour; Da Silva et al. 2015; Brüning et al. 2018; Grubisic et al. 2019), the population or species (e.g., abundance, distribution, reproduction, mortality, dispersal; Dominoni et al. 2013; Gaston and Bennie 2014; Stone et al. 2012; Davies et al. 2017; Rodríguez et al. 2017), the community (e.g., species composition and richness, trophic structure; e.g., Sanders and Gaston 2018), through to the ecosystem (e.g., pollination, seed dispersal; Lewanzik and Voigt 2014; Knop et al. 2017). Impacts have been documented in all environmental realms (marine, freshwater, terrestrial), in a wide array of habitat types, and across microbes, plants, fungi and animals (Gaston et al. 2013; Bennie et al. 2016; Sanders et al. 2021). The mechanisms by which these effects occur are increasingly well understood (Gaston et al. 2013; Dominoni 2015; Falcon et al. 2020).

This said, how pervasive or systemic the biological impact of ALAN actually is, and in this sense how comparable to other anthropogenic pressures on the environment, remains rather unclear. On the one hand, ALAN continues to be characterised widely as essentially an urban issue, and to be discussed most commonly in the context of towns and cities. Whilst its effects would then be tightly linked to where most people occur, this would actually suggest a relatively limited spatial extent compared to many other anthropogenic pressures (e.g., climate change, ocean acidification, plastic pollution). Estimates of global urban coverage are highly variable - in large part dependent on definitions of urban land cover, and the spatial resolution and accuracy of the data used - but consensus seems to be that this is less than 1% of the land surface (e.g., Potere et al. 2009; Schneider et al. 2009; Li et al. 2017). Some national and regional urban coverages can be several-fold larger, but are almost invariably still quite limited (excepting some small highly developed nations and regions; Zhou et al. 2015). Of course, these zones of influence are markedly increased by the inclusion of skyglow, but this has recently been estimated still 'only' to occur over less than a quarter of global land area (Falchi et al. 2016).

On the other hand, natural light regimes are the strongest and most predictable environmental fluctuations that organisms typically experience and play a fundamental role in biology (Bradshaw and Holzapfel 2010; Gaston et al. 2017). They can be exquisitely sensitive to variation in these regimes, through diel, lunar and seasonal cycles. Thus, for example, (i) at low latitudes where annual variation in day length is less than an hour, some plant species are still using such changes as biological cues (e.g., Borchert and Rivera 2001; Rivera and Borchert 2001); (ii) in the high Arctic polar night zooplankton undertake diel vertical migration response to diel variation in solar radiance despite the sun never rising above the horizon (Ludvigsen et al. 2018); and (iii) nocturnal insects can see colour, fly, navigate, manoeuvre at fine scales and land at known targets at very low nighttime light levels (Sponberg et al. 2015; Warrant 2017). This would suggest that altering natural light regimes even mildly could have potent biological effects.

In this paper we highlight a series of factors that suggest that the biological effects of ALAN may indeed be spatially much more pervasive than is often understood. We focus foremost on factors that have in the main received limited research attention, often because this is challenging to undertake. We consider, first estimations of the extent of ALAN. We then turn to a series of ways in which ALAN may be more widespread than an 'urban' characterisation would imply. We address, in turn, isolated sources, mobile sources, marine systems, remote sources, light levels, skyglow, elevation, landscape ecology, interactions with other factors. Finally, we consider what future changes might portend. We end with some concluding remarks. Throughout, our intent is not to provide a comprehensive systematic review, but rather to explore and link key issues and provide some examples.

Estimations of the extent of ALAN

Remote sensing data, especially those derived from satellite-borne sensors, have played a pivotal role in documenting the extent, composition and dynamics of ALAN (Levin et al. 2020). Indeed, although other platforms are becoming available (Colomb et al. 2003; Li et al. 2019; Zhang et al. 2019), most understanding of these issues has arisen from data obtained from the Defense Meteorological Satellite Program Operational Linescan System (DMSP-OLS) and, more recently, the Visible Infrared Imaging Radiation Suite (VIIRS) on the Suomi National Polar-Orbiting Partnership satellite. Whilst they have been invaluable, these sources have important limitations. In determining the extent of direct emissions of ALAN these include that:

- (i) the spatial resolution of data acquisition remains quite coarse (DMSP-OLS - 2.7 km; VIIRS - 740 m), making them better suited to detecting emissions from collections of, rather than individual, lights;
- (ii) variation in the angle of data acquisition means that emissions from some sources of artificial lighting are more apparent than others, with emissions close to the nadir, and those from well shielded and more horizontally projected sources, being less detectable (combining data obtained at multiple different times and thus different angles can reduce this issue);
- (iii) emissions that are shielded by natural vegetation (e.g., under tree canopies, as may occur in some rural, typically tropical, communities) will not be well detected;

- (iv) data are consistently acquired at particular local times of day when this may not necessarily be at its peak and which means that the inevitable diel variation is not captured (DMSP-OLS - 20.00; VIIRS - 01.30);
- (v) sensors are panchromatic, and insensitive to emissions in the blue part of the spectrum, which are particularly associated with the broad 'white' light-emitting diode (LED) technology that is increasingly being used in outdoor lighting (Bierman 2012). This can result in substantial underestimation (e.g., 50%) of the overall intensity of ALAN emissions;
- (vi) sensors are sensitive to emissions in the infrared, and thus to those from volcanoes and fires, as well as high pressure sodium (HPS) lamps; and
- (vii) some ALAN emissions will be removed when processing data to reduce contributions of lighting from non-ALAN sources, such as airglow, the aurora and fires. This is particularly true of ALAN emissions that appear to vary on short timescales, either because they genuinely do so (e.g., because of limited or erratic availability of power, or because they are mobile), or because they appear to do so as a consequence of variation in the angles of data acquisition at different times.

Many of these limitations are greatly exacerbated by the need to obtain satellite data in the absence of substantial cloud cover, conditions that occur infrequently in some regions and times of year. Many of the limitations have variously been overcome with the use of other sensor platforms (e.g., balloons, drones, manned aircraft, the International Space Station), but these have thus far been extremely limited in the ground/sea coverage that has been achieved and in the frequency of repeat data acquisition for the same areas (Levin et al. 2020).

Direct light can be observed at an almost indefinite distance from the source (subject to the curvature of the earth and atmospheric scattering), but shading, geometric dilution and the attenuation of light in the atmosphere mean that direct illuminance declines with distance from the source and can vary by orders of magnitude over short distances. Any estimate of the global area affected by ALAN emissions is therefore necessarily dependent on the spatial resolution used to measure it and the intensity threshold used.

Processing data from VIIRS to estimate the global coverage of both direct ALAN emissions and skyglow, at a spatial resolution of 1.61 km x 2.12 km, ALAN extends over 49.5% of the land surface between 59°N and 55°S (Fig. 1; this is where confounding effects of albedo, airglow, the aurora and permanent snow and ice on imagery are relatively small). By way of comparison, when sampled at the same resolution a recent attempt to map ambient human densities (averaged over 24 hours) reveals that between the same latitudinal limits people and ALAN co-occur over 39.4% of the global land surface, but that people occur over 63% of that surface (Fig. 2). Whilst there are doubtless areas in which people are present but producing little or no ALAN emissions outside of buildings, the combination of these estimates suggests that the extent of ALAN may indeed be markedly underestimated by satellite data in isolation due to low detection thresholds. Such satellite data, and models based upon them, have frequently been used to assess the extent of the biological impacts of ALAN (e.g., Bennie et al. 2014a; Duffy et al. 2015; Gaston et al. 2015; Correa-Cano et al. 2018; Koen et al. 2018; Garrett et al. 2020). These may thus prove to be significant underestimates.

Isolated sources

Much of the difference between the extent of ALAN as estimated from nighttime satellite data and as inferred from the distribution of people will arise from the occurrence of lighting at low densities outside of towns and cities, and away from foci of industrial activity (e.g., airports, ports, power stations, mining sites) and major transport arteries (e.g., motorways). Indeed, in understanding the biological impacts of ALAN it is likely to be important to distinguish between two situations. The first is that in which these emissions arise from multiple sources and essentially contribute to most local organisms living in a much altered 'light environment'; this is linked to various attempts to determine what contribution different kinds of sources (from streetlights, housing, businesses, advertisement hoardings etc) make to this environment (Kuechly et al. 2012; Kyba et al. 2020). The second situation is that in which the effect of ALAN on organisms is driven by their responses to individual light sources. As evidenced by a recent formal meta-analysis (Sanders et al. 2021), to date studies of the biological impacts of ALAN have tended to focus foremost on the former circumstance. This has typically involved observations in the vicinity of streetlights or subjecting organisms to experimental treatments intended to simulate such light conditions. This is, in effect, also a focus on the

impacts of downward emissions from light sources, and that is often how these are narratively and figuratively predominantly conceptualised.

Considering the isolated light sources that predominate in more rural areas focusses attention on an issue that is of less importance in urban areas, namely how far emissions are spatially propagated. For true diffuse point sources, the illuminance of a receiving plane decreases with the square of the distance from the light source (the inverse square law):

$$E = k(I/r^2)$$

where E is illuminance, I is the luminous intensity of the source, r is the distance and k a constant (dependent on the units used; Schreuder 2010). Hence, illuminance at or close to the horizontal from isolated inadequately shielded streetlights and other such sources will tend to decline quite quickly. When a source is small in relation to the distance this equation also works as an approximation for non-point sources, and large sources can usefully be treated as multiple small ones. For 'bundled' sources, where emissions are shaped into a parallel beam by a mirror or lens, the inverse square law underestimates the level of illuminance at a given distance with, all else being equal, the luminous intensity remaining roughly constant with distance. Indeed, for such sources, aside from intervening obstacles (vegetation, buildings, topography), the distance over which they can be detected is principally determined by the curvature of the earth. Most external lighting sources (such as street lights) fall closer to the inverse-square law than a focused, bundled source, but some spotlights, floodlights, vehicle headlights and some modern LED lighting fixtures emit more parallel beams of light and so luminance will decline with distance at a much slower rate.

The distances over which different organisms respond to isolated artificial light sources remain surprisingly poorly understood. Most attention in this regard has been paid to insects, particularly in connection with understanding the areas over which light traps are effectively sampling. For moths, estimates of this distance vary greatly but suggest that attraction is limited - within generally a few metres (Baker and Sadovy 1978);

generally < 30 m for 50% of individuals (Beck and Linsenmair 2006); generally < 10 m, but up to 80 m (Truxa and Fiedler 2012); generally < 10 m, but up to 50 m (van Grunsven et al. 2014); or up to at least 50 m (Plaut 1971). For midges, this distance is generally 2-4 m (Venter et al. 2012), but up to a maximum of 15.5 m (Kirkeby et al. 2013). This suggests that for the majority of individuals these distances are low, although given that in the main these studies are based on mark-recapture over short periods, the potential that some individuals may travel much greater distances remains.

The other group for which there is some information on distances over which individuals respond to artificial light sources is seabirds (a group which can experience high mortality as a result of such attraction; Rodríguez et al. 2017). In the main this information is indirect, being based on the separation between colonies and the sources of lighting (e.g., Imber 1975; Rodrigues et al. 2012; Rodríguez et al. 2014; Syopsz et al. 2018). However, one study using GPS data-loggers to track the maiden and second flights of Cory's shearwater *Calonectris diomedea* fledglings from nest-burrows found that these were grounded by artificial lights at distances up to 16 km (Rodríguez et al. 2015a). Another found that fledglings of three species of petrels were grounded by artificial lights at a mean distance of ~5 km and up to >20 km from their colonies (Rodríguez et al. 2015b).

Similar kinds of measurements do not seem to have been conducted for other groups of organisms, and thus the range of attraction of artificial light sources is essentially unknown. Moreover, equivalent experiments testing repulsion effects of isolated lights have not been done.

In addition to attraction or repulsion, isolated artificial lights in landscapes could have other effects, including through visual confusion or distraction. For example, fireflies and glow worms may find it harder to detect light signals made by other individuals (e.g., Owens and Lewis 2018; Desouhant et al. 2019; Lewis et al. 2020). Attraction of isolated lights can also have second-order effects, for example through attraction of predators (e.g., spiders, frogs, bats) to aggregations of prey that have been directly attracted (Canário et al. 2012; Minnaar et al. 2015; Rodríguez et al. 2021).

Mobile sources

Attention to ALAN predominantly focuses on static sources of emissions (e.g., streetlights, building lights). Many mobile, and therefore often temporally sporadic, sources (e.g., from road vehicles, rail vehicles, shipping) will not be detected by remote sensing because they are projected predominantly in a horizontal plane, and indeed, if they are detected may be removed in the processing of data to reduce the influence of fires (natural and human-caused) and gas flares.

Nonetheless, mobile sources may both be extensive and disproportionately contribute to ALAN beyond urban areas (where the vast majority of emissions will be from static sources, and where the majority of streetlights occur). The global road network alone is estimated to extend over 36 million km (CIA 2020), and in 2015 there were an estimated 947 million passenger cars and 335 million commercial vehicles in use (OICA 2020); in the UK, one of the only regions for which data are accessible, 16-48% of traffic is on the road outside of daylight hours, depending on the time of year (Department for Transport 2016). In many regions, road coverage is sufficiently dense that most land may be exposed to light emissions from vehicle headlights. In particular, headlights introduce artificial light into areas that do not experience streetlights, especially in rural areas (including protected areas) that may be otherwise buffered from many other anthropogenic impacts on the environment. Vehicle headlights produce a focused beam (a 'bundled' source; see above) that is projected horizontally and travels further and at higher intensities than light emissions from streetlighting. For example, modelling suggests that half of land in Great Britain is less than 216 m from a road, and that around 70% of land may be exposed to vehicle headlight emissions, whilst only a small portion of this is exposed to emissions from streetlighting (Fig. 3; Phillips et al. 2021).

Most consideration of the environmental impacts of vehicle headlights has focussed on the dazzling of vertebrates, especially mammals, and the resultant potential for causing collisions with vehicles (e.g., Outen 2002). However, particularly importantly, light emissions from vehicle headlights are commonly experienced locally as irregularly timed pulses as vehicles pass, rather than as continuous lighting. In laboratory contexts,

light pulses, even when infrequent and very short-lived, are capable of major perturbations to the circadian rhythms of organisms, to their visual systems and to their behaviour (for review see Gaston and Holt 2018). The rapid and unpredictable fluctuations in light levels experienced by roadside organisms are likely also to lead to greater problems with physiological and behavioural habituation.

Alongside road vehicles, the other major source of mobile ALAN emissions is shipping. There were estimated to be 3.7 million marine fishing vessels in 2015 (Rousseau et al. 2019), and 98,000 vessels in the global merchant fleet in 2020 (UNCTAD 2020). These vessels can use lights for generalist (e.g., anti-collision, deck operations) or more specialist (e.g., fishing, anchor handling, diving) activities, although much is emitted simply as a consequence of a lack of routine use of blinds or other forms of blackout. Attention to the biological impacts of ALAN from shipping has focused foremost on the potential for bird strikes, particularly in regions where seabirds are abundant (Dick and Donaldson 1978; Black 2005; Merkel and Johansen 2011). However, there may be others (see below).

Marine systems

The vast majority of research into the extent of ALAN and into its biological impacts has concerned terrestrial and freshwater systems (113/126 studies in the meta-analysis of Sanders et al. 2021). Nonetheless, ALAN is widely experienced by marine systems, and seems likely to have similarly major biological implications. Davies et al. (2014) estimated from satellite-derived data that 22% of the world's coastlines (excluding Antarctica) were exposed to ALAN, although this is likely to be a marked underestimate for reasons already discussed. Obviously much of this coastal lighting can carry far out to sea because light paths are typically unhindered, but more locally may influence much of the extent of natural coastal ecosystems including that of, fast disappearing, tidal flats (Murray et al. 2019). Other forms of largely static ALAN emissions in marine systems arise from repeated or long-term mooring of vessels in the same localities, from offshore oil and gas platforms, and increasingly from wind turbine arrays. Shipping as a source of mobile emissions has already been mentioned, with the global coverage of its routes over even a single year being extremely widespread (e.g., Halpern et al. 2008). However, particularly significant is the use of lights, for deck operations, or

explicitly as lures, by fishing vessels, with studies having demonstrated large aggregations of such lighting in some parts of the oceans (e.g., <https://earthobservatory.nasa.gov/features/Malvinas>).

The biological impacts of ALAN in marine systems seem likely to be at least as diverse as those occurring terrestrially, with evidence already for effects on timing of coral spawning (Ayalon et al. 2020); invertebrate settlement (Davies et al. 2015); behaviour of pelagic organisms (Berge et al. 2020); turtle nesting and orientation (Thums et al. 2016; Silva et al. 2017; Hu et al. 2018; Vandersteen et al. 2020); and seabird grounding and mortality (Wiese et al. 2001; Le Corre et al. 2002; Jones and Francis 2003; Rodríguez et al. 2014; Syposz et al. 2018). Impacts on diel vertical migration of zooplankton could be especially profound, given its importance for global carbon cycling.

Remote sources

Whilst not distinct from the considerations mentioned thus far, the sometimes remote occurrence of ALAN emissions is worth emphasis. Such emissions occur at remote industrial sites (e.g., mines, sawmills, oil rigs), on oceanic islands, at isolated desert and forest settlements, tourist lodges and research sites, and at Arctic and Antarctic bases (Fig. 4). In the context of some other environmental pressures (e.g., plastic pollution), such occurrences often attract much scientific and media attention. Ironically, of course, demonstrating those occurrences for other pressures often itself entails the introduction - if only temporarily - of ALAN.

The impacts of its introduction, even temporarily, into “light-naïve” areas that have previously not experienced ALAN may be much greater than those in regions where ALAN has been present within the wider landscape for some time. This is suggested by historical accounts where gas or electric light was introduced for the first time. In the early 19th Century, the entomologist Edward Doubleday (1837), for example, reported trapping moths in rural New York State by removing the glass window of a room lit by a single lamp – he cited reports that in summer hundreds of moths would be swept from the floor of the room in the morning. Early reports from lighthouses reported mass mortality of migrating birds through attraction followed by collision and/or exhaustion - 10,000 birds were reported killed in one season at Gatteville lighthouse in France, and

3,200 in two nights at Belle-Ile (AP 1917). More recently, gas flaring on oil production facilities in the Arctic has been shown to attract large numbers of birds (Day et al. 2015).

Remote sources of ALAN may also have disproportionate biological impacts if, as seems likely, they are often bright and poorly or unshielded, resulting in emissions above the horizontal. This may occur because of perceived needs in the absence of widespread lighting sources, because of weaker regulations, or weaker enforcement of regulations.

Light levels

Implicit to much discussion about its biological impacts is that these will be increased by greater levels of ALAN. This may indeed often be the case, with the suppression of melatonin production, for example, showing a clear positive dose response relationship (Grubisic et al. 2019). Given that higher levels of ALAN tend often (although far from exclusively) to be associated with urban areas, this would encourage belief that these are where the biological impacts are likely to be most severe (subject to adaptation and habituation). However, a recent meta-analysis, addressing a wide variety of impacts, found no evidence for a systematic increase in effect sizes with levels of ALAN (Sanders et al. 2021). Moreover, it has been shown that the complexities of cascading effects through food webs can mean that lower intensities have greater impacts than higher ones (Sanders et al. 2018). If these results generalize, then the fact that ALAN is considerably more widely spatially distributed at lower levels than at higher ones becomes much more significant in interpreting the extent of its impacts.

Skyglow

Thus far we have focussed principally on the occurrence and biological impact of direct emissions of ALAN. In terms of the extent of ALAN then skyglow is also an important consideration. As well as being directly associated with urban areas it has been found to be detectable over distances of hundreds of kilometers from these sources (Luginbuhl et al. 2014), with its reach being greater, through amplification, on cloudy nights (Kyba et al. 2011; Jechow et al. 2017). This means that skyglow extends into many areas identified as

important for biodiversity and into many areas protected for biological conservation (Garrett et al. 2020).

It is widely presumed that skyglow has biological impacts, both by influencing levels of organismal activity and by obscuring natural diel, lunar and seasonal cycles. This is reinforced by the aforementioned evidence for impacts of ALAN at low intensities. This said, studies of these biological impacts, which can have significant practical challenges, have thus far been scarce, and limited to experimental demonstrations of the influence of skyglow on the movements of organisms in aquatic systems (Moore et al. 2000; Torres et al. 2020).

Skyglow tends predominantly to be measured and modelled at the zenith, but is at its brightest on the horizon (for examples see Jechow et al. 2017). This could serve to extend its spatial influence much further, if animals used the increased brightness of horizons either to improve their detection of prey or of predators, or for orientation (see Limpus and Kamrowski 2013). It would be surprising if some species at least did not exploit these opportunities, although it seems likely to be hard to demonstrate in the field.

Elevation

The biological impacts of ALAN tend foremost to be thought of, and measured (including for skyglow), in terms of the emissions that are likely to be sensed or experienced by organisms at ground level. However, flying animals may (literally) have very different perspectives, and be influenced by ALAN on very different spatial scales. Ignoring atmospheric refraction and any intervening obstacles (vegetation, buildings, topography),

$$d \approx 3.57\sqrt{h}$$

where d is distance to the horizon (in km), and h is height above sea level (in m).

For example, whilst nocturnally migrating birds commonly fly at relatively low altitudes (a few hundred metres), they may regularly do so up to 6000-7000 m above sea level, and sometimes even higher (Liechti

and Schaller 1999; Bruderer et al. 2018; Sjöberg et al. 2018). This equates to direct emissions of ALAN potentially being visible over distances of 10s to 100s of kilometres. There is evidence that this can result in these birds being attracted to ALAN sources from long distances, which may result in course adjustments through to landfall and stopover in lit areas (Bowlin et al. 2015; La Sorte et al. 2017; McLaren et al. 2018; but see Cabrera-Cruz et al. 2020).

Landscape ecology

Aside from consideration of possible influences on migratory movements, most observations and studies of the biological impacts of ALAN have concerned local scales. This ignores the fact that ALAN could have landscape scale effects, and the extent to which ALAN influences the perceived connectivity or fragmentation of habitats, and can act as a barrier or as a facilitator of movements. By modifying or restricting patterns of movement, dispersal and migration at a landscape scale, there is potential for ALAN to have population-level effects even in landscapes where only a small proportion of the area is exposed to ALAN and for species where the duration of direct exposure (as a proportion of the life of an individual) is relatively brief. It is well known that road networks can have profound effects on the movement of animals and thus on the viability of populations (van der Ree et al. 2015), despite the actual area of land directly affected by roads being much smaller than that unaffected. Similarly, networks and patches of light may have landscape effects as barriers and population sinks. Modifying the ability of species to move through landscapes could alter metapopulation dynamics, gene flow between populations and foraging opportunities for individuals. Modelling and some field studies have shown that these effects could be very significant (e.g., Beier 1995; Bennie et al. 2014b; Laforge et al. 2019).

Interactions with other factors

In common with most discussion of the biological impacts of ALAN, thus far we have treated this as an environmental pressure in isolation from others. However, virtually no such pressure acts entirely independently. Key therefore in understanding the biological impacts of ALAN is to consider whether it is likely to exacerbate other forms of pollution, particularly those that act unequally across the diel cycle and

for which there is evidence that the impacts could be greater at night, such as global warming asymmetry (Cox et al. 2020b) and ocean acidification (Price et al. 2012).

Future changes

We have focused throughout this piece on the current situation. However, just as important is how ALAN, and its biological impacts, will change in the future. In terms of raw infrastructure, studies have variously projected that globally urban areas (the fastest growing land use) will increase by a factor of 1.8-5.9 by 2100 (Gao and O'Neill 2020), the global road network by 1.65, and road vehicle numbers by 2.0 by 2050 (Dulac 2013). Furthermore, the vast majority of this growth (e.g., 90% of growth in road use) will come from regions outside of Europe and North America (Dulac 2013; Gao and O'Neill 2020), where existing ALAN is lower, so such growth is likely to result in a disproportionate increase in the extent of ALAN. This will serve directly to exacerbate global biological impacts of ALAN, including through many of the mechanisms highlighted in this paper.

It also seems likely that global biodiversity will continue to decline, with suggestions that even substantive steps are unlikely quickly to reverse this trend (e.g., Leclère et al. 2020). This will mean that any negative biological impacts of ALAN are likely to become proportionally yet more significant.

These effects could potentially be somewhat offset if there were strong evolutionary responses to the biological impacts of ALAN. As yet, evidence of such responses is rather limited (but see Altermatt and Ebert 2016). This might be anticipated given that the reliance on natural light cues for the timings of many components of biological activity is evolutionarily deep-rooted. However, the selection pressure exerted by ALAN may be strong (particularly where it leads to large effects on organismal fitness through reduced reproduction or increased mortality), and thus nonetheless encourage evolutionary responses. Whether the paucity of evidence for these responses is simply a consequence of a paucity of studies remains an open question.

In conclusion

There are regions of the earth that continue to experience natural light regimes. And there are more extensive areas over which the changes to those regimes caused by ALAN are, whilst detectable, unlikely to have substantive biological impacts. But it is also clear that given the diversity of sources and ways in which ALAN is emitted, and the diversity of ways in which it impacts biological systems, the proportion of the earth over which ALAN occurs at levels at which it is likely to have such impacts is marked. It is certainly vastly greater than the areal coverage by cities and towns in which context it is usually discussed and more comparable to that over which many other key anthropogenic environmental pressures have impacts. Particularly given the rapidity with which the global extent of ALAN is growing, this highlights the importance of studying its biological impacts in non-urban settings and for the purpose of understanding its impacts in those settings.

Acknowledgements

We are grateful to A. Rodríguez for kind assistance, and two anonymous reviewers for their helpful comments.

Funding

This work was supported in part by Natural Environment Research Council grant NE/P01156X/1.

Data availability

No new data are used in this manuscript.

References

- Allen JA. 1880. Destruction of birds by light-houses. *Bull Nutt Ornithol Club* 5:131-138.
- Altermatt F, Ebert D. 2016. Reduced flight-to-light behaviour of moth populations exposed to long-term urban light pollution. *Biol Lett* 12:20160111.
- “A.P.” 1917. Les oiseaux et les phares. *Le Chenil et l’Echo de Elevage* 51:602-603.

- Ayalon I, Rosenberg Y, Benichou JIC, Campos CLD, Sayco SLG, Nada MAL, Baquiran JIP, Ligson CA, Avisar D, Conaco C, Kuechly HU, Kyba CCM, Cabaitan PC, Levy O. 2020. Coral gametogenesis collapse under artificial light pollution. *Curr Biol* 31:1-7.
- Baker RR, Sadovy Y. 1978. The distance and nature of the light-trap response of moths. *Nature* 276:818-821.
- Beck J, Linsenmair KE. 2006. Feasibility of light-trapping in community research on moths: Attraction radius of light, completeness of samples, nightly flight times and seasonality of southeast-asian hawkmoths (Lepidoptera: Sphingidae). *J Res Lepid* 39:18-37.
- Beier P. 1995. Dispersal of juvenile cougars in fragmented habitat. *J Wildl Manage* 59:228-237.
- Bennie J, Davies TW, Cruse D, Gaston KJ. 2016. Ecological effects of artificial light at night on wild plants. *J Ecol* 104:611-620.
- Bennie J, Davies TW, Inger R, Gaston KJ. 2014b. Mapping artificial lightscapes for ecological studies. *Methods Ecol Evol* 5:534-540.
- Bennie J, Duffy JP, Inger R, Gaston KJ. 2014a. Biogeography of time partitioning in mammals. *Proc Natl Acad Sci USA* 111:13727-13732.
- Berge J, Geoffroy M, Daase M, Cottier F, Priou P, Cohen JH, Johnsen G, McKee D, Kostakis I, Renaud PE, Vogedes D, Anderson P, Last KS, Gauthier S. 2020. Artificial light during the polar night disrupts Arctic fish and zooplankton behaviour down to 200m depth. *Commun Biol* 3:102.
- Bierman A. 2012. Will switching to led outdoor lighting increase sky glow? *Light Res Technol* 44:449-458.
- Black A. 2005. Light induced seabird mortality on vessels operating in the Southern Ocean: incidents and mitigation measures. *Antarc Sci* 17:67-68.
- Borchert R, Rivera G. 2001. Photoperiodic control of seasonal development and dormancy in tropical stem-succulent trees. *Tree Physiol* 21:213-221.
- Bowlin MS, Enstrom DA, Murphy BJ, Plaza E, Jurich P, Cochran J. 2015. Unexplained altitude changes in a migrating thrush: long-flight altitude data from radio-telemetry. *Auk* 132:808-816.
- Bradshaw WE, Holzapfel CM. 2010. Light, time, and the physiology of biotic response to rapid climate change in animals. *Annu Rev Physiol* 72:147-166.

- Bruderer B, Peter D, Korner-Nievergelt F. 2018. Vertical distribution of bird migration between the Baltic Sea and the Sahara. *J Ornithol* 159:315-336.
- Brüning A, Kloas W, Preuer T, Hölker F. 2018. Influence of artificially induced light pollution on the hormone system of two common fish species, perch and roach, in a rural habitat. *Conserv Physiol* 6:368.
- Cabrera-Cruz SA, Cohen EB, Smolinsky JA, Buler JJ. 2020. Artificial light at night is related to broad-scale stopover distributions of nocturnally migrating landbirds along the Yucatan peninsula, Mexico. *Remote Sens* 12:395.
- Canário F, Leitão AH, Tomé R. 2012. Predation attempts by short-eared and long-eared owls on migrating songbirds attracted to artificial lights. *J Raptor Res* 46:232-234.
- [CIA] Central Intelligence Agency. 2020. The World Fact Book. <https://www.cia.gov/library/publications/the-world-factbook/index.html>.
- Cinzano P, Falchi F, Elvidge CD. 2001. The first World Atlas of the artificial night sky brightness. *Mon Not R Astron Soc* 328:689-707.
- Cochran WW, Graber RR. 1958. Attraction of nocturnal migrants by lights on a television tower. *Wilson Bull* 70:378-380.
- Colomb R, Alonso C, Nollmann I. 2003. SAC-C mission and the international am constellation for earth observation. *Acta Astron* 52:995-1005.
- Correa-Cano ME, Goettsch B, Duffy JP, Bennie J, Inger R, Gaston KJ. 2018. Erosion of natural darkness in the geographic ranges of cacti. *Sci Rep* 8:4347.
- Cox DTC, Maclean IMD, Gardner AS, Gaston KJ. 2020b. Global variation in diurnal asymmetry in temperature, cloud cover, specific humidity and precipitation and its association with Leaf Area Index. *Glob Change Biol* 26:7099-7111.
- Cox DTC, Sánchez de Miguel A, Dzurjak SA, Bennie J, Gaston KJ. 2020a. National scale spatial variation in artificial light at night. *Remote Sens* 12:1591.
- Da Silva A, Valcu M, Kempnaers B. 2015. Light pollution alters the phenology of dawn and dusk singing in common European songbirds. *Phil Trans R Soc B* 370:20140126.

- Davies TW, Bennie J, Cruse D, Blumgart D, Inger R, Gaston KJ. 2017. Multiple night-time light-emitting diode lighting strategies impact grassland invertebrate assemblages. *Global Change Biol* 23:2641-2648.
- Davies TW, Coleman M, Griffith KM, Jenkins SR. 2015. Night-time lighting alters the composition of marine epifaunal communities. *Biol Lett* 11:20150080.
- Davies TW, Duffy J, Bennie J, Gaston KJ. 2014. The nature, extent, and ecological implications of marine light pollution. *Front Ecol Environ* 12:347-355.
- Day RH, Rose JR, Prichard AK, Streever B. 2015. Effects of gas flaring on the behaviour of night-migrating birds at an artificial oil-production island, Arctic Alaska. *Arctic* 68:367-379.
- Department for Transport. 2016. Road traffic statistics (TRA) - statistical data set, <https://www.gov.uk/government/statistical-data-sets/road-traffic-statistics-tra>
- Desouhant E, Gomes E, Mondy N, Amat I. 2019. Mechanistic, ecological, and evolutionary consequences of artificial light at night for insects: review and prospective. *Entom Exp Appl* 167:37-58.
- Dick MH, Donaldson W. 1978. Fishing vessel endangered by crested auklet landings. *Condor* 80:235-236.
- Dominoni D, Quetting M, Partecke J. 2013. Artificial light at night advances avian reproductive physiology. *Proc R Soc B* 280:20123017.
- Dominoni DM. 2015. The effects of light pollution on biological rhythms of birds: an integrated, mechanistic perspective. *J Ornithol* 156 (Suppl 1):S409-S418.
- Doubleday E. 1837. Communications on the natural history of North America. *Entomol Mag* 5:21-34.
- Duffy JP, Bennie J, Durán AP, Gaston KJ. 2015. Mammalian ranges are experiencing erosion of natural darkness. *Sci Rep* 5:12042.
- Dulac J. 2013. Global land transport infrastructure requirements: estimating road and railway infrastructure capacity and costs to 2050. <https://webstore.iea.org/global-land-transport-infrastructure-requirements>.
- Falchi F, Cinzano P, Duriscoe D, Kyba CCM, Elvidge CD, Baugh K, Portnov BA, Rybnikova NA, Furgoni R. 2016. The new world atlas of artificial night sky brightness. *Sci Adv* 2:e1600377.
- Falcon J, Torriglia A, Attia D, Viénot F, Gronfier C, Behar-Cohen F, Martinsons C, Hicks D. 2020. Exposure to artificial light at night and the consequences for flora, fauna, and ecosystems. *Front Neurosci* 14:321.

- Gao J, O'Neill BC. 2020. Mapping global urban land for the 21st century with data-driven simulations and Shared Socioeconomic Pathways. *Nature Comm* 11:2302.
- Garrett JK, Donald PF, Gaston KJ. 2020. Skyglow extends into world's Key Biodiversity Areas. *Anim Conserv* 23:153-159.
- Gastman EA. 1886. Birds killed by electric light towers at Decatur, Ill. *Am Nat* 20:981.
- Gaston KJ. 2018. Lighting up the nighttime. *Science* 362:744-746.
- Gaston KJ, Bennie J. 2014. Demographic effects of artificial nighttime lighting on animal populations. *Environ Rev* 22:323-330.
- Gaston KJ, Bennie J, Davies TW, Hopkins J. 2013. The ecological impacts of nighttime light pollution: A mechanistic appraisal. *Biol Rev* 88:912-927.
- Gaston KJ, Davies TW, Nedelec SL, Holt LA. 2017. Impacts of artificial light at night on biological timings. *Annu Rev Ecol Evol Syst* 48:49-68.
- Gaston KJ, Duffy JP, Bennie J. 2015. Quantifying the erosion of natural darkness in the global protected area system. *Conserv Biol* 29:1132-1141.
- Gaston KJ, Duffy JP, Gaston S, Bennie J, Davies TW. 2014. Human alteration of natural light cycles: causes and ecological consequences. *Oecologia* 176:917-931.
- Gaston KJ, Holt LA. 2018. Nature, extent and ecological implications of night-time light from road vehicles. *J Appl Ecol* 55:2296-2307.
- Grubisic M, Haim A, Bhusal P, Dominoni DM, Gabriel KMA, Jechow A, Kupprat F, Lerner A, Marchant P, Riley W, Stebelova K, van Grunsven RHA, Zeman M, Zubidat AE, Hölker F. 2019. Light pollution, circadian photoreception, and melatonin in vertebrates. *Sustainability* 11:6400.
- Halpern BS, Walbridge S, Selkoe KA, Kappel CV, Micheli F, D'Agrosa C, Bruno JF, Casey KS, Ebert C, Fox HE, Fujita R, Heinemann D, Lenihan HS, Madin EMP, Perry MT, Selig ER, Spalding M, Steneck R, Watson R. 2008. A global map of human impact on marine ecosystems. *Science* 319:948-952.
- Hölker F, Moss R, Griefahn B, Kloas W, Voigt CC, Henckel D, Hänel A, Kappeler PM, Völker S, Schwöpe A, Franke S, Uhrandt D, Fischer J, Klenke B, Wolter C, Tockner K. 2010. The dark side of light: a transdisciplinary research agenda for light pollution policy. *Ecol Soc* 15:13.

- Hu Z, Hu H, Huang Y. 2018. Association between nighttime artificial light pollution and sea turtle nest density along Florida coast: A geospatial study using VIIRS remote sensing data. *Environ Poll* 239:30-42.
- Imber MJ. 1975. Behaviour of petrels in relation to the moon and artificial lights. *Notornis* 22:302-306.
- Jechow A, Kolláth Z, Ribas SJ, Spoelstra H, Hölker F, Kyba CCM. 2017. Imaging and mapping the impact of clouds on skyglow with all-sky photometry. *Sci Rep* 7:6741.
- Jones J, Francis CM. 2003. The effects of light characteristics on avian mortality at lighthouses. *J Avian Biol* 34:328-333.
- Kirkeby C, Græsbøll K, Stockmarr AS, Christiansen LE, Bødker R. 2013. The range of attraction for light traps catching *Culicoides* biting midges (Diptera: Ceratopogonidae). *Parasites Vectors* 6:67.
- Knop E, Zoller L, Ryser R, Gerpe C, Hörler M, Fontaine C. 2017. Artificial light at night as a new threat to pollination. *Nature* 548:206-209.
- Koen EL, Minnaar C, Roever CL, Boyles JG. 2018. Emerging threat of the 21st century lightscape to global biodiversity. *Glob Change Biol* 24:2315-2324.
- Kuechly HU, Kyba CCM, Ruhtz T, Lindemann C, Wolter C, Fischer J, Hölker F. 2012. Aerial survey and spatial analysis of sources of light pollution in Berlin, Germany. *Remote Sens Environ* 126:39-50.
- Kyba CCM, Ruby A, Kuechly HU, Kinzey B, Miller N, Sanders J, Barentine J, Kleinodt R, Espey B. 2020. Direct measurement of the contribution of street lighting to satellite observations of nighttime light emissions from urban areas. *Lighting Res Technol* 104, online early.
- Kyba CCM, Ruhtz T, Fischer J, Hölker F. 2011. Cloud coverage acts as an amplifier for ecological light pollution in urban ecosystems. *PLoS One* 6:e17307.
- Laforge A, Pauwels J, Faure B, Bas Y, Kerbiriou C, Fonderflick J, Besnard A. 2019. Reducing light pollution improves connectivity for bats in urban landscapes. *Landsc Ecol* 34:793-809.
- La Sorte FA, Fink D, Buler JJ, Farnsworth A, Cabrera-Cruz SA. 2017. Seasonal associations with urban light pollution for nocturnally migrating bird populations. *Global Change Biol* 23:4609-4619.
- Le Corre M, Ollivier A, Ribes S, Jouventin P. 2002. Light-induced mortality of petrels: a 4-year study from Réunion Island (Indian Ocean). *Biol Conserv* 105:93-102.

- Leclère D, Obersteiner M, Barrett M, Butchart SM et al. 2020. Bending the curve of terrestrial biodiversity needs an integrated strategy. *Nature* 585:551-556.
- Levin N, Kyba CCM, Zhang Q, Sánchez de Miguel A, Román MO, Li X, Portnov BA, Molthan AL, Jechow A, Miller SD, Wang Z, Shrestha RM, Elvidge CD. 2020. Remote sensing of night lights: A review and an outlook for the future. *Remote Sens Environ* 237:111443.
- Lewanzik D, Voigt CC. 2014. Artificial light puts ecosystem services of frugivorous bats at risk. *J Appl Ecol* 51:388-394.
- Lewis SM, Wong CH, Owens ACS, Fallon C, Jepsen S, Thancharoen A, Wu C, De Cock R, Novák M, López-Palafox T, Khoo V, Reed JM. 2020. A global perspective on firefly extinction threats. *BioScience* 70:157-167.
- Li X, Chen G, Liu X, Liang X, Wang S, Chen Y, Pei F, Xu X. 2017. A new global land-use and land-cover change product at a 1-km resolution for 2010 to 2100 based on human–environment interactions. *Ann Am Assoc Geogr* 107:1040-1059.
- Li X, Liu Z, Chen X, Sun J. 2019. Assessing the ability of Luojia 1-01 imagery to detect feeble nighttime lights. *Sensors* 19:3708.
- Liechti F, Schaller E. 1999. The use of low-level jets by migrating birds. *Naturwissenschaften* 86:549-551.
- Limpus C, Kamrowski RL. 2013. Ocean-finding in marine turtles: the importance of low horizon elevation as an orientation cue. *Behaviour* 150:863-893.
- Longcore T, Rich C. 2004. Ecological light pollution. *Front Ecol Environ* 2:191-198.
- Ludvigsen M, Berge J, Geoffroy M, Cohen JH, De La Torre PR, Nornes SM, Singh H, Sørensen AJ, Daase M, Johnsen G. 2018. Use of an Autonomous Surface Vehicle reveals small-scale diel vertical migrations of zooplankton and susceptibility to light pollution under low solar irradiance. *Sci Adv* 4:eaap9887.
- Luginbuhl CB, Boley PA, Davis DR. 2014. The impact of light source spectral power distribution on skyglow. *J Quant Spectrosc Rad Transf* 139:21-26.
- Matzke EB. 1936. The effect of street lights in delaying leaf-fall in certain trees. *Am J Bot* 23:446-452.
- Mclaren JD, Buler JJ, Schreckengost T, Smolinsky JA, Boone M, van Loon, EE, Dawson DK, Walters EL. 2018. Artificial light at night confounds broad-scale habitat use by migrating birds. *Ecol Lett* 21:356-364.

- Merkel FR, Johansen KL. 2011. Light-induced bird strikes on vessels in Southwest Greenland. *Mar Poll Bull* 62:2330-2336.
- Minnaar C, Boyles JG, Minnaar IA, Sole CL, McKechnie AE. 2014. Stacking the odds: light pollution may shift the balance in an ancient predator-prey arms race. *J Appl Ecol* 52:522-531.
- Moore MV, Pierce SM, Walsh HM, Kvalvik SK, Lim JD. 2000. Urban light pollution alters the diel vertical migration of *Daphnia*. *Verh Int Ver Limnol* 27:779-782.
- Murray NJ, Phinn SR, DeWitt M, Ferrari R, Johnston R, Lyons MB, Clinton N, Thau D, Fuller RA. 2019. The global distribution and trajectory of tidal flats. *Nature* 565:222-225.
- OICA 2020. International Organization of Motor Vehicle Manufacturers, Motorization rate 2015 – worldwide, <http://www.oica.net/category/vehicles-in-use/>
- Outen AR. 2002. The ecological effects of road lighting. In B Sherwood, D Cutler, J Burton (eds), *Wildlife and roads: the ecological impact*, pp. 133-155. Imperial College Press, London.
- Owens ACS, Cochard P, Durrant J, Farnworth B, Perkin EK, Seymoure B. 2020. Light pollution is a driver of insect declines. *Biol Conserv* 241:108259.
- Owens ACS, Lewis SM. 2018. The impact of artificial light at night on nocturnal insects: A review and synthesis. *Ecol Evol* 8:11337-11358.
- Phillips BB, Bullock JM, Osborne JL, Gaston KJ. 2021. Spatial extent of road pollution: a national analysis. *STOTEN*, 773:145589.
- Plaut H. 1971 Distance of attraction of moths of *Spodoptera littoralis* to BL radiation, and recapture of moths released at different distances of an ESA blacklight standard trap. *J Econ Entom* 64:1402-1404.
- Potere D, Schneider A, Angel S, Civco DL. 2009. Mapping urban areas on a global scale: which of the eight maps now available is more accurate? *Int J Remote Sens* 30:6531-6558.
- Price NN, Martz TR, Brainard RE, Smith JE. 2012. Diel Variability in seawater pH relates to calcification and benthic community structure on coral reefs. *PLoS One* 7:e43843.
- Rich C, Longcore T (eds). 2006. *Ecological consequences of artificial night lighting*. Island Press, Washington.
- Rivera G, Borchert R. 2001. Induction of flowering in tropical trees by a 30-min reduction in photoperiod: evidence from field observations and herbarium specimens. *Tree Physiol* 21:201-212.

- Rodrigues P, Aubrecht C, Gil A, Longcore T, Elvidge C. 2012. Remote sensing to map influence of light pollution on Cory's shearwater in São Miguel Island, Azores Archipelago. *Eur J Wildl Res* 58:147-155.
- Rodríguez A, Burgan G, Dann P, Jessop R, Negro JJ, Chiaradia A. 2014. Fatal attraction of short-tailed shearwaters to artificial lights. *PLoS One* 9:e110114.
- Rodríguez A, García D, Rodríguez B, Cardona E, Parpal L, Pons P. 2015b. Artificial lights and seabirds: is light pollution a threat for the threatened Balearic petrels? *J Ornithol* 156:893-902.
- Rodríguez A, Holmes ND, Ryan PG, Wilson K-J, Faulquier L, Murillo Y, Raine AF, Penniman JF, Neves V, Rodríguez B, Negro JJ, Chiaradia A, Dann P, Anderson T, Metzger B, Shirai M, Deppe L, Wheeler J, Hodum P, Gouveia C, Carmo V, Carreira GP, Delgado-Alburquerque L, Guerra-Correa C, Couzi F-X, Travers M, Corre ML. 2017. Seabird mortality induced by land-based artificial lights. *Conserv Biol* 31:986-1001.
- Rodríguez A, Orozco-Valor PM, Sarasola JH. 2021. Artificial light at night as a driver of urban colonization by an avian predator. *Landsc Ecol* 36:17-27.
- Rodríguez A, Rodríguez B, Negro JJ. 2015a. GPS tracking for mapping seabird mortality induced by light pollution. *Sci Rep* 5:10670.
- Román MO, Wang Z, Sun Q, Kalb V, Miller SD, Molthan A, Schultz L, Bell J, Stokes EC, Pandey B, Seto KC, Hall D, Oda T, Wolfe RE, Lin G, Golpayegani N, Devadiga S, Davidson C, Sarkar S, Praderas C, Schmaltz J, Boller R, Stevens J, González OMR, Padilla E, Alonso J, Detrés Y, Armstrong R, Miranda I, Conte Y, Marrero N, MacManus K, Esch T, Masuoka EJ. 2018. NASA's Black Marble nighttime lights product suite. *Remote Sens Environ* 210:113-143.
- Rousseau Y, Watson RA, Blanchard JL, Fulton EA. 2019. Evolution of global marine fishing fleets and the response of fished resources. *Proc Nat Acad Sci USA* 116:12238-12243.
- Sanders D, Frago E, Kehoe R, Patterson C, Gaston KJ. 2021. A meta-analysis of biological impacts of artificial light at night. *Nat Ecol Evol* 5:74-81.
- Sanders D, Gaston KJ. 2018. How ecological communities respond to artificial light at night. *J Exp Zool A* 329:394-400.
- Sanders D, Kehoe R, Cruse D, van Veen FJF, Gaston KJ. 2018. Low levels of artificial light at night change food web dynamics. *Curr Biol* 28:2474-2478.

- Schneider A, Friedl MA, Potere D. 2009. A new map of global urban extent from MODIS satellite data. *Environ Res Lett* 4:044033.
- Schreuder D. 2010. *Outdoor lighting: physics, vision and perception*. Springer.
- Schroeder CA. 1945. Tree foliation affected by street lights. *Arborists News* 10:1-3.
- Silva E, Marco A, da Graça J, Pérez H, Abella E, Patino-Martinez J, Martins S, Almeida C. 2017. Light pollution affects nesting behavior of loggerhead turtles and predation risk of nests and hatchlings. *J Photochem Photobiol B* 173:240-249.
- Sjöberg S, Pedersen L, Malmiga G, Alerstam T, Hansson B, Hasselquist D, Thorup K, Tøttrup AP, Andersson A, Bäckman J. 2018. Barometer logging reveals new dimensions of individual songbird migration. *J Avian Biol* 49:e01821.
- Sponberg S, Dyhr JP, Hall RW, Daniel TL. 2015. Luminance-dependent visual processing enables moth flight in low light. *Science* 348:1245-1248.
- Stone EL, Jones G, Harris S. 2012. Conserving energy at a cost to biodiversity? Impacts of led lighting on bats. *Glob Change Biol* 18:2458-2465.
- Sullivan III WT. 1989. A 10 km resolution image of the entire night-time Earth based on cloud-free satellite photographs in the 400–1100 nm band. *Int J Remote Sens* 10:1-5.
- Syposz M, Gonçalves F, Carty M, Hoppitt W, Manco F. 2018. Factors Influencing Manx Shearwater grounding on the west coast of Scotland. *Ibis* 160:846-854.
- Thums M, Whiting SD, Reisser J, Pendoley KL, Pattiaratchi CB, Proietti M, Hetzel Y, Fisher R, Meekan MG. 2016. Artificial light on water attracts turtle hatchlings during their near shore transit. *R Soc Open Sci* 3:160142.
- Torres D, Tidau S, Jenkins S, Davies T. 2020. Artificial skyglow disrupts celestial migration at night. *Curr Biol* 30:R696-R697.
- Truxa C, Fiedler K. 2012. Attraction to light - from how far do moths (Lepidoptera) return to weak artificial sources of light? *Eur J Entomol* 109:77-84.
- UNCTAD. 2020. *Handbook of Statistics 2020*. <https://unctadstat.unctad.org/EN/>
- van der Ree R, Smith DJ, Grilo C. (eds) 2015. *Handbook of Road Ecology*. Wiley Blackwell, Oxford.

- van Grunsven RHA, Lham D, van Geffen KG, Veenendaal EM. 2014. Range of attraction of a 6-w moth light trap. *Entomol Exp Applic* 152:87-90.
- Vandersteen J, Kark S, Sorrell K, Levin N. 2020. Quantifying the impact of light pollution on sea turtle nesting using ground-based imagery. *Remote Sens* 12:1785.
- Venter G, Majatladi D, Labuschagne K, Boikanyo S, Morey L. 2012. The attraction range of the Onderstepoort 220V light trap for *Culicoides* biting midges as determined under South African field conditions. *Vet Parasitol* 190:222-229.
- Verheijen FJ. 1960. The mechanisms of the trapping effect of artificial light sources upon animals. *Arch Neerl Zool* 13:1-107.
- Warrant EJ. 2017. The remarkable visual capacities of nocturnal insects: vision at the limits with small eyes and tiny brains. *Phil Trans R Soc B* 372:20160063.
- Wiese FK, Montevecchi WA, Davoren GK, Huettmann F, Diamond AW, Linke J. 2001. Seabirds at risk around offshore oil platforms in the North-west Atlantic. *Mar Poll Bull* 42:1285-1290.
- Zhang G, Guo X, Li D, Jiang B. 2019. Evaluating the potential of LJ1-01 nighttime light data for modeling socio-economic parameters. *Sensors* 19:1465.
- Zhou Y, Smith SJ, Zhao K, Imhoff M, Thomson A, Bond-Lamberty B, Asrar GR, Zhang X, He C, Elvidge CD. 2015. A global map of urban extent from nightlights. *Environ Res Lett* 10:054011.

Figure legends

Fig. 1 Spatial variation in artificial light at night over the global land surface. The layer gives an estimated combined extent of direct emissions and skyglow between 59°N and 55°S at 1.61 * 1.21 km resolution in Behrmann equal area projection (see Cox et al. [2020a] for a full description of methods). In brief, the layer gives VIIRS day/night band (DNB) values corrected for albedo and skyglow. To avoid the confounding effects of the aurora and permanent snow and ice, we excluded pixels above 59°N and 55°S. Once albedo had been removed, DNB airglow was detected over the majority of the land surface. Airglow varies with latitude and therefore to provide the best possible mask while minimising the obscuring of true direct ALAN emissions, for each 200 km latitudinal band we calculated the median DNB value and converted all values below the median to 0 (mean across latitudinal bands 0.098; min 0.015, max 0.202). To incorporate skyglow we used values for artificial brightness from the New World Atlas of Artificial Sky Brightness (Falchi et al. 2016). Following Falchi et al. (2016) we set artificial brightness values below 0.00174 mcd/m² to 0, because these were considered indistinguishable from a pristine night sky. To map the extent of both forms of ALAN, values of skyglow were added to values of direct ALAN emissions before converting all values greater than 0 to 1. and thereby creating a binary layer of where ALAN is present (white) or absent (black).

Fig. 2 Spatial variation in human population density. The layer was created by adjusting the raw LandScan 2016 population count layer for area and projecting to Behrmann equal area (EASE-Grid 2.0: EPSG:6933; Brodzik et al. 2012) to give population density per km² at a resolution of 1.61 * 2.12 km². The colour banding is log₁₀ at intervals of 0.25.

Fig. 3 Estimated variation across Great Britain in ALAN from (a) streetlights and (b) vehicle headlights in terms of maximum exposure, i.e. light level when a vehicle passes, and (c) average exposure, accounting for traffic volumes (based on Phillips et al. 2021). Maps (a) and (c) are modelled using a simple inverse square decay function (ignoring topography), with respect to distance from roads, and varied event frequency based on the type of roads.

Fig. 4 Four examples of ALAN found in remote places in the world. (A) island of Tahiti, a popular tourist destination in the Pacific Ocean, with most ALAN arising from the coastline because the island interior is mostly uninhabited (17.6509° S, 149.4260° W); (B) central island of Santa Cruz in the Galapagos archipelago, home to the Galapagos National Park, with the principal source of ALAN coming from the town of Puerto Ayora (0.6144° S, 90.3451° W); (C) remote volcanic archaeological site, Easter Island, with much of the island being protected as a world heritage site and the ALAN being produced by the main town, Hanga Roa (27.1127° S, 109.3497° W); and (D) small villages producing ALAN on the track of the Amazon river (bottom track) in South America and the Japura River (top track), while being surrounded by protected and conserved land (2°14'52.5"S 66°39'19.6"W). ALAN data from VIIRS 2019 composite.