

**Texas School Survey of Drug and Alcohol Use
Methodology Report
2014**

For the Texas Department of State Health Services

Prepared by

Miner P. Marchbanks III, Ph.D.

James A. Dyer, Ph.D.

Allison Seibert, M.Ed.

Shannon Peairson, M.S.

Public Policy Research Institute

Texas A&M University

College Station, TX 77843-4476

Table of Contents

INTRODUCTION.....	1
SURVEY INSTRUMENT	2
SURVEY CONTENT	3
SURVEY SAMPLE	4
SELECTION OF CAMPUSES.....	4
SOLICITING CAMPUS PARTICIPATION	5
SELECTION OF CLASSROOMS WITHIN CAMPUSES.....	6
SURVEY ADMINISTRATION PROCEDURES	7
DATA ENTRY AND ANALYSES	8
EXAGGERATED RESPONSES	8
UNREPORTED GRADE LEVELS	8
QUALITY CONTROL MEASURES	9
WEIGHTS, STANDARD ERRORS, AND CONFIDENCE INTERVALS	10
CONCLUSION	11
APPENDIX A - SURVEY INSTRUMENTS	A-1
APPENDIX B - PARTICIPATING CAMPUSES	B-1
APPENDIX C - CLASSROOM SELECTION GUIDELINES	C-1
APPENDIX D - SCHOOL SURVEY ADMINISTRATION	D-1
APPENDIX E - COMPUTATION OF SAMPLE WEIGHTS	E-1
APPENDIX F - STANDARD ERROR COMPUTATION.....	F-1
APPENDIX G - STANDARD ERROR & CONFIDENCE INTERVAL TABLES.....	G-1

Introduction

The Public Policy Research Institute (PPRI), in conjunction with the Texas Department of State Health Services (DSHS) conducted the fourteenth statewide survey of drug and alcohol use among Texas public school students in the spring of 2014. Originally implemented in 1988 as a component of a larger survey assessing substance use among the state's general population, the school survey has since become an ongoing, independent project. District surveys are offered every year with a statewide survey conducted every two years. The 2014 effort provides follow-up data reflecting changes over the past twenty-six years in grades six through twelve.

The *Texas School Survey* project has two primary objectives. First, it serves to inform state and local policy-makers about the extent and nature of youth substance use in Texas public schools. Second, the statewide survey provides a standard of comparison for districts conducting local assessments of drug and alcohol use.

The purpose of this document is to describe the methodology used to administer the *2014 Texas School Survey of Drug and Alcohol Use*. Following a brief introduction to the survey instrument itself, attention is focused on sample selection and survey administration procedures. Next, methods for data processing and quality control are described and the report concludes with a review of standard error estimates.

Survey Instrument

Since its inception in 1988, the *Texas School Survey of Drug and Alcohol Use* has received slight revisions throughout the years for purposes of incorporating new drugs or other relevant topics to measure alcohol and drug use trends among youth. However, major revisions rarely occur and typically result from the need to reduce survey administration time in schools. The last major revision occurred in 1998 where similar questions assessing substance use prevalence rates were grouped together in a matrix format to reduce survey administration time. Again, in the fall of 2012, PPRI and DSHS proposed several changes to improve the quality of survey results while maintaining compatibility with previous survey data.

In previous survey administration years, two versions of the Texas School Survey of Drug and Alcohol Use were administered. The elementary version of the survey was a simplified three-page instrument created for students in grades four through six. Elementary students were asked about only four types of substances including tobacco (cigarettes, snuff, and chewing tobacco), alcohol (beer, wine, wine coolers, and liquor), inhalants, and marijuana. The secondary instrument was a six-page questionnaire designed for students in grades seven through twelve. Secondary students were asked about the same substances as elementary students, as well as a broader range of illicit drugs including powdered cocaine, crack, hallucinogens, steroids, ecstasy, Rohypnol, DXM, methamphetamine, and heroin. Other sets of questions on both the elementary and secondary instruments were designed to assess behavioral correlates of substance use and students' perceptions of support available to help them cope with substance-related problems. Rather than maintain these two separate forms, PPRI developed one survey form for grades 6 through 12. The decision to migrate to one form was made based on increased concerns from parents and school districts about surveying students as young as 4th and 5th grade as well as the need to streamline administration procedures for schools. Therefore, starting in 2014, the survey design involved surveying students in grades 6 through 12 using one survey form.

During the fall of 2012, PPRI and DSHS revised the *Texas School Survey of Drug and Alcohol Use* secondary survey instrument with the purpose of increasing the accuracy of responses, preventing respondent fatigue, and to include responses from students in grade 6. The list of revisions for the 2014 administration included:

- (1) Removing questions that assessed length of attendance in the school district, receipt of an allowance, parents' education level, use of uppers or downers, method of marijuana use, and incidents of getting into trouble with police or teachers because of alcohol and/or drug use;
- (2) Revising the list of extracurricular activities (deleted drama, speech, or art clubs; combined student government with student newspaper and yearbook; and combined academic club or societies with academic competition groups);
- (3) Addition of synthetic marijuana (spice, K2) and methamphetamine (speed, crystal meth, ice, or crank) to major drug lists;
- (4) Removing hallucinogens from major drug lists except core prevalence questions (*How recently have you used...*);
- (5) Combining cigarettes and smokeless tobacco categories into 'tobacco', and beer, wine coolers, wine, and liquor into 'alcohol' on major drug lists;
- (6) Including 'not prescribed to you' as qualifier for prescription drug use; and

(7) Combining science class with social studies class for sources of information.

It should be noted that the removal of uppers and downers and addition of synthetic marijuana and methamphetamine to the major drug lists has resulted in changes to the categorization of 'any illicit drug use' in survey data and reports. Therefore, comparing changes in prevalence rates overtime for 'any illicit drug use' should be interpreted with caution.

Survey Content

To assess possible differences in student responses between the 2012 survey form and the proposed 2014 version, PPRI conducted a pilot study during the spring of 2013 in two public school districts. All students in grades 7-12 were provided either the 2012 secondary version or the proposed 2014 version of the survey. Students in grades 6 received either the 2012 elementary version or the proposed 2014 version of the survey. The surveys were administered using the standard *Texas School Survey* protocols. Very few significant differences were found between the 2012 and 2014 survey forms. Many significant findings from bivariate analyses were no longer significant once student demographic, academic, and personal characteristics were controlled for in the statistical analyses. When examining the analyses for grades 7-12, the number of statistically significant results falls within a normal range given the overall analyses completed.

Over time, PPRI has administered a paper-pencil version of the *Texas School Survey of Drug and Alcohol Use*. Since 2012, schools have had the option of choosing between a web-based version of the survey or the traditional paper-pencil version. The web survey is programmed in LimeSurvey, a widely used software application for internet surveys, and hosted on a secure server at Texas A&M University. The paper-pencil questionnaire is in a format that can be scanned optically, similar to that used for standardized testing. Both versions are designed for anonymous self-administration by students with the aid of a staff member to pass out the survey or web survey tokens, read a common set of instructions, monitor the class during survey administration, and collect the instruments or web survey tokens after the survey is completed. The survey instrument is included in Appendix A.

Survey Sample

In recent years, recruiting school districts has become increasingly challenging where greater numbers of school districts, particularly large school districts, refuse to participate in the survey. Decreases in participation rates can significantly impact the ability to generalize survey results to the Texas youth population. To address these concerns and attempt to improve participation rates, PPRI implemented a new sampling strategy for the 2014 statewide administration.

Selection of Campuses

The sample of students for the 2014 survey was designed to be a stratified, weighted, random sample of all public school students between the sixth and twelfth grades in the state. In order to make administration practical, students were selected using a multi-stage stratified sampling procedure. This involved sampling schools, and then classrooms within schools.

Schools were the primary sampling units (PSU) and were sampled with probability of selection proportionate to size¹. This means that a school with 200 students is twice as likely to be sampled as a school with 100 students. The probabilities of selection were increased for strata that required larger samples given reporting needs. For instance, border schools are oversampled to provide sufficient numbers of schools in the border area to allow for separate Border/Non-Border reports. Other oversampled areas included schools in rural areas, middle school or junior high campuses, and schools in smaller geographic areas needed for the regional reports. For each grade level in a sample school, eight classrooms were randomly selected. In schools with fewer than eight classrooms in a grade, all classrooms were surveyed. All students in a sampled classroom were asked to participate in the survey.

Initially 352 schools were sampled. This number was selected to provide an estimated sample size to report with statistical confidence on the drug and alcohol related behavior of youth in the public schools statewide. It was large enough to report separately for subgroups defined by characteristics such as gender and ethnicity. It also provided enough oversampling to compensate for the expectation that some schools would not participate. Later, DSHS approached PPRI about creating regional reports defined by the eleven designated Public Health Regions. Because some of the regions are rural in nature, few schools from these regions fell into the original state sample; additional schools were sampled in several regions—a total of 128 additional schools. As the recruiting process progressed, the reluctance of many schools to participate became apparent. In order to compensate for this, the research team added an additional 119 campuses to the sample. The original sample and the additional samples were all randomly selected with probability of selection proportionate to the student population as described above. In total, 599 campuses were sampled to be recruited. Again, the sample size was determined so that after accounting for nonparticipation, it would be large enough to provide reliable estimates for subgroups defined by variables such as gender, ethnicity, and economically disadvantaged within each grade level.

As detailed in the section “Weights, Standard Errors, and Confidence Intervals” below, the sample of completed surveys was weighted for the probability of selection as well as the

¹ For discussion of weighting with probability proportionate to size see William Cochran, *Sampling Techniques*, Third Edition. New York: John Wiley and Sons, pp 250-258.

populations of the strata of grades and regions. The weighted sample can be considered a representative sample of public school students in Texas between the 6th and 12th grades.

As described above, this sampling methodology is a change from what the Texas School Survey has used in the past. Previously the PSUs were the districts and not the campuses. The representativeness of the sample is dependent on selected school districts agreeing to participate in the survey, thus district refusals can significantly impact the ability to generalize results to the Texas youth population. The current sampling procedure is more similar to that used in the Youth Risk Behavior Survey (YRBS) and other surveys of students in high schools.

The new survey form, sampling design, and recruitment strategies should make the data more representative than data collected in the past. It should be noted that one must interpret apparent trends in the data with caution.²

Soliciting Campus Participation

In preparation for recruiting schools, PPRI survey coordinators, Shannon Pearison and Allison Seibert, reviewed the list of sampled schools to determine whether PPRI had a contact at the district level from previous survey administration years. If PPRI worked with the school district in the past, they were contacted directly to explain the changes in sample design. In schools where no previous contacts had been established, survey coordinators initiated contact with the school principal along with a district testing coordinator, health specialist, or other individual that seemed like they would be involved with student health/substance use surveys.

PPRI started calling each randomly selected school in September after the initial letters were sent to discuss survey administration procedures and answer any questions. PPRI completed a minimum of three calls to each school. For non-responsive campuses, PPRI contacted the superintendent or other district staff to request approval for conducting the survey at a particular campus. Schools interested in participating in the 2014 survey were instructed to complete a basic participation form via the project's website (<http://texasschoolsurvey.org>). One hundred and six of the original 599 selected campuses participated in the study. Most campuses declined due to the lack of time and resources involved in survey administration.

Recruitment methods for the state survey were similar to past survey administration years where school contacts were sent a packet of information including a recruitment letter, letter of support from DSHS, copy of the survey instrument, and frequently asked question guide. When sending the initial recruitment letters, PPRI offered a \$500 monetary incentive to the state sampled campuses as an incentive to increase participation rates and reduce the burden of survey administration on schools. The purpose of payment was to help schools defray costs associated with implementing this survey, such as costs associated with the distribution of parental notification forms and other administrative costs. In the past, the primary incentive used by PPRI to garner school participation was the district level report. Due to the removal of Safe and Drug Free Schools and Communities (SDFSC) funding, which required district level reporting of drug and alcohol use, the value of this incentive has waned over time. Therefore, PPRI did not actively promote local level reports as incentives for participation.

District level reports were offered to both districts with state sampled campuses as well as districts with no state sampled campuses for an additional cost. When districts requested this, additional data were collected for district level reports. Those with state sampled campuses received the additional surveys and district level report at a discounted rate. Additional data

² Fulton, Janet E. and Susan A. Carlson. 2012. "Surveillance of Physical Activity" in Ainsworth, Barbara E. and Caroline A. Macera, eds. *Physical Health and Public Practice*. Boca Raton, Florida: CRC Press, p 231.

collected from non-state campuses to produce the district level reports were not included in the state data analysis.

Selection of Classrooms within Campuses

The selection of classrooms was performed by campus personnel based on a set of guidelines provided by PPRI (illustrated in Appendix C). Members of campus staff were asked to make a list by grade (according to teacher’s last name or some other convenient method) of all classes held during a selected class period or subject area. PPRI randomly selected classes from this list until the target numbers per grade level were reached.

TABLE 1. Number of Surveys Included in State Sample

	Total Non-Blank Surveys	Usable Surveys	Number rejected	Percent rejected
Total	40,007	38,413	1,594*	4.0%

*Surveys were rejected because the responses indicated exaggeration or the survey could not be matched to a sampled school and grade.

TABLE 2. Survey Distribution by Grade

Grade	Number of Usable Surveys	Percentage
Grade 6	4950	12.9%
Grade 7	5753	15.0%
Grade 8	5570	14.5%
Grade 9	6340	16.5%
Grade 10	5914	15.4%
Grade 11	5427	14.1%
Grade 12	4459	11.6%
Total	38,413	100%

Survey Administration Procedures

Each participating school had the ability to administer either an online version of the survey via a secure website, or the traditional paper-pencil administration. Schools selected which method of administration they preferred when signing up for the survey. Out of the 106 participating state schools, 31 administered the online version of the survey and 75 administered the paper-pencil version.

An Information Sheet regarding the survey was provided for required dissemination to parents/guardians a minimum of two weeks prior to survey administration. The Information Sheet explained the voluntary nature of the survey and that students may choose to opt out of participation at any time. It also provided a mechanism for parents to let the school know if they did not want their child to participate. This form also emphasized that participation is completely anonymous.

Relevant personnel in the selected campuses were provided with complete instructions and materials necessary to administer the survey (see Appendix D). Classrooms were selected randomly by PPRI based on information from an Excel spreadsheet from the district. Teachers in selected classrooms were given a script to read so that all students would receive a standardized set of instructions. The scripts were provided to the campuses at least a week before survey administration in order to allow teachers sufficient time to familiarize themselves with the survey procedures. Teachers were also asked to complete a Classroom Identification Form that provided data on the number of students that are currently enrolled in the class. This information was useful for computing error estimates.

Survey administration for the paper-pencil version followed standard *Texas School Survey* procedures. Students received a computer-scannable questionnaire booklet to record their answers to survey questions by filling in corresponding ovals with a number two pencil. Students were instructed to place the questionnaire in a large manila envelope once the survey was completed. Teachers were instructed to place any blank or unused surveys back in the manila envelope to be returned to PPRI.

Survey administration for the web version followed the standard *Texas School Survey* procedures with slight modifications due to the variation in mode. Students completed the web survey either in their regular classroom using portable laptop computers, or by taking the survey in the school computer lab. Instead of receiving paper copies of the survey instrument, students in sampled classrooms were provided with unique, randomly generated six digit alpha-numeric codes, referred to as survey tokens. These survey tokens could only be used once and provided access to the online survey via a secure website. To ensure only sampled classrooms completed the online survey, all unused and used survey tokens were collected by the survey administrator when surveys were completed. Additionally, PPRI limited access to the online survey during the district's scheduled survey administration period.

The envelopes containing either the survey tokens or survey instruments from all participating classrooms were collected and returned to PPRI.

Data Entry and Analyses

As noted earlier, the format of the paper-pencil survey instruments enabled them to be scanned optically. Upon receipt at PPRI, the paper-pencil surveys were logged in, coded and scanned by staff or trained personnel.

Exaggerated Responses

Because the *Texas School Survey* data is based entirely upon respondents' description of their own behavior, it is inevitable that some students will under- or over-report their use of drugs or alcohol, and to the extent possible PPRI attempted to identify and eliminate data from those respondents. Two checks were incorporated into the data analysis program to identify exaggerators. First, students were asked about their use of a false drug called "Damaxil". Data from students claiming to have used this substance were considered suspect and dropped from the analyses.

Second, checks were run to identify any students claiming extremely high levels of drug and alcohol use. Students were defined as exaggerators if they reported using three or more drugs (other than tobacco, alcohol, inhalants, or steroids) daily and if they reported using four or more drugs (other than tobacco, alcohol, or steroids) eleven or more times in the past month. As in those cases where students reported using "Damaxil", data from students reporting exaggerated use were also dropped from the analyses.

Unreported Grade Levels

When students failed to report their grade level, it was impossible to determine unequivocally in which grade these students' data should be analyzed. When a grade level was missing, an estimate of the grade was made based on the students' age and the data were retained. Table 3 identifies the range of students' ages and the corresponding grade levels that were assigned. If both grade and age were missing, the data were dropped from the analyses.

TABLE 3. Age-Based Grade Assignments.

Age	Grade
11	6 th Grade
12	7 th Grade
13	8 th Grade
14	9 th Grade
15	10 th Grade
16	11 th Grade
17 or older	12 th Grade

Quality Control Measures

To ensure the quality of the statewide survey data, a number of internal checks were put into place to guide survey processing. First, a quality control analyst oversaw the implementation of all pre- and post-analysis quality control procedures. As the following paragraphs describe, many aspects of PPRI's plan for quality control were embedded in automated procedures. However, there is no replacement for human oversight. The quality control analyst monitored and tracked the processing of each campus' surveys from the initial mailing through the production of the final state report. Responsibilities included ensuring that surveys were properly coded and scanned and checking for anomalies in the final table of results.

In addition to the safeguards resulting from careful project oversight, there were also a number of procedural checks against error. For example, there was a possibility, however remote, that after the bindings of a set of survey instruments were cut, the instruments could be dropped or otherwise placed out of order. If this occurred, it is conceivable that some pages of data could have been read into the incorrect computer record. To resolve this problem, each instrument used in the 2014 survey was printed with a five-digit "litho-code" number. With this coding process, every page of a given instrument is printed with the same scannable number, but a unique number is assigned to every instrument. By using the litho-code, when each page of an instrument is scanned it will automatically be read into the correct computer record. In this way, even if the pages from different instruments were shuffled together and read randomly, all data derived from the same instrument would automatically be read to the same data record. Additional quality control measures were conducted on approximately ten percent of the surveys processed and included counting the number of surveys in a stack after it was read by the optical scanner to make sure the number of surveys inserted corresponded to the number of surveys read by the scanner and comparing the physical survey responses to the output generated by the scanner software.

Programming checks were also incorporated into the data analysis program by cross-analysis. That is, the same data was run in several different ways using existing programs, and program outputs were then compared for consistency. Confidence is high that these quality control features will ensure valid and reliable survey findings.

Weights, Standard Errors, and Confidence Intervals

Weights were applied to each case based on the strata (unique subgroups discussed below), district, and campus. The weights were applied so that the aggregation of students in each campus, district, and strata reflected their proportions in the actual district, campus, and strata populations. The formula use to determine these weights are presented in Appendix E. The strata used in weighting included grade, region, and ethnicity.

Even when random sampling is performed, responses do not perfectly match the population. This can occur for a variety of reasons including random oversampling of one strata or non-response. Weighting allows for the data that are collected to be adjusted to better approximate the true population paramaters by reducing the weight given to those “oversampled” and increasing the weight given to those “undersampled.”

Standard errors and confidence intervals were estimated for each grade and within the major subgroups. A more detailed discussion of this approach to error estimation and examples of computing it with commonly used statistical analysis software are found in Appendix F. The table of standard errors and confidence intervals for past year and lifetime use of substances by grades are presented in Appendix G.

Conclusion

As discussed earlier, the sample design was changed in response to the problem of non-participation from school districts. For a variety of reasons, willingness to participate has been declining. The results were not as good as we had hoped. Schools have indicated a reluctance to give up class time due to perceived pressures regarding required standardized testing. Further, many individuals have reported that they are “surveyed-out” due to the increased requests by researchers to collect data in their schools. Many have made the decision to not allow further surveys.

Moving forward, exploring new options to improve participation is crucial. We have already made the survey available online, offered flexible implementation dates, provided monetary incentives and targeted the schools directly. Still, there are likely other avenues that would pay dividends. PPRI is exploring the option of compensating the survey administrators for their time. Implementing the survey is not a trivial task for the school administrator; perhaps a token of appreciation would help. There is precedence in other states for a state mandate for schools to participate in the survey. This option is outside of our control; however, it would likely lead to a dramatic increase in participation. Finally, PPRI is in ongoing communication with individuals across the state including school personnel as well as DSHS staff to explore avenues to increase participation. While the new approach did not yield high response rates, the research team is confident that in the absence of the changes the response rate would have been lower.

The *Texas School Survey* has become a valuable policy tool for both state and local educators and policy-makers. The survey, administered every two years, provides timely and relevant information about current drug and alcohol use patterns among young people enrolled in Texas’ public schools. Various regional breakdowns including border, non-border and regional analyses provide the ability to compare various diverse areas of Texas with the state as a whole. These results can yield important information on the unique needs of different regions in Texas, thus informing policy makers for purposes of program design and resource allocation for substance abuse prevention among youth in Texas. Furthermore, longitudinal analysis can provide insight into changes in drug and alcohol prevalence over time. As was noted in the introduction, every state survey culminates in a Texas Department of State Health Services publication providing an overview of findings to date.

Appendix A
Survey Instrument

1. Are you:

Male Female

2. What grade are you in?

6 7 8 9 10 11 12

3. Do you live with both of your parents?

Yes No

4. How old are you?

10 or younger 11 12 13 14 15 16 17 18 19+

5. Are you:

White African American Mexican American Asian American Native American Other

6. On average what grades do you get? (DARKEN ONE BUBBLE ONLY)

Mostly A's Mostly B's Mostly C's Mostly D's Mostly F's

7. Do you have a job?

Yes No

8. During the current school year, do you qualify for a free or reduced price school lunch?

Yes
 No
 Don't Know

9. Do you regularly participate in any of the following extra-curricular activities? (DARKEN ONE BUBBLE FOR EACH LINE: a-i.)

	Yes	No	
a.	<input type="radio"/>	<input type="radio"/>	School athletics (UIL, Varsity, Junior Varsity)
b.	<input type="radio"/>	<input type="radio"/>	School band/orchestra
c.	<input type="radio"/>	<input type="radio"/>	School choir
d.	<input type="radio"/>	<input type="radio"/>	Drill team or cheerleading squad
e.	<input type="radio"/>	<input type="radio"/>	Student government, newspaper, or yearbook
f.	<input type="radio"/>	<input type="radio"/>	Academic clubs, societies, or competition groups (UIL, math, science, foreign language, etc.)
g.	<input type="radio"/>	<input type="radio"/>	Other school clubs or student groups
h.	<input type="radio"/>	<input type="radio"/>	Athletic teams OUTSIDE of school
i.	<input type="radio"/>	<input type="radio"/>	Other clubs or groups OUTSIDE of school

10. How safe do you feel when you are: (DARKEN ONE BUBBLE FOR EACH LINE: a-c.)

	Very Safe	Somewhat Safe	Not Very Safe	Not Safe At All	Don't Know
a. in your home	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. out in your neighborhood	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. at school	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

11. How many of your FRIENDS would you say: (DARKEN ONE BUBBLE FOR EACH LINE: a-e.)

	None	A Few	Some	Most	All
a. Feel close to their parents?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Sometimes carry weapons like a knife or gun?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Care about making good grades?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Belong to a gang or are interested in becoming a gang member?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. Wish they could drop out of school?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

12. Do one or both of your parents usually attend school-sponsored open houses or PTA meetings? Yes No

Below is a list of things some people sniff to get high. They are called **INHALANTS**.

13. How recently, if ever, have you sniffed, huffed, or inhaled the following INHALANTS for "kicks" or to get "high":
(DARKEN ONE BUBBLE FOR EACH LINE: a-h.)

	Never heard of/ Never used it	Used at least once in the Past Month	Used at least once Since School Began in the Fall	Used at least once In Your Lifetime
a. Spray paint to get high?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Whiteout, correction fluid, magic markers to get high?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Computer dusting sprays to get high?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Helium, butane, propane, whippets (nitrous oxide), refrigerants/Freon gas to get high?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. Glue to get high?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
f. Toluene/tolly, paint thinner, other solvents to get high?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
g. Gasoline, octane booster, carburetor cleaner to get high?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
h. Other aerosols/sprays (deodorant spray, Pam, hair spray, room spray, etc.) to get high?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

14. How recently, if ever, have you used the following TOBACCO, ALCOHOL, and INHALANT products even one time:
(DARKEN ONE BUBBLE FOR EACH LINE: a-g.)

	Never heard of/ Never used it	Used at least once in the Past Month	Used at least once Since School Began in the Fall	Used at least once In Your Lifetime
a. Cigarettes?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Smokeless Tobacco?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Beer?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Wine Coolers (hard lemonade, hard sodas, etc.)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. Wine?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
f. Liquor (vodka, rum, whiskey, etc.)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
g. Inhalants (whiteout, spray paint, glue, gas, etc.) to get high?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

15. How recently, if ever, have you used the following DRUGS even one time:
(DARKEN ONE BUBBLE FOR EACH LINE: a-k.)

	Never heard of/ Never used it	Used at least once in the Past Month	Used at least once Since School Began in the Fall	Used at least once In Your Lifetime
a. Marijuana?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Cocaine (not crack)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Crack?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Damaxil?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. Hallucinogens (LSD, PCP, Mushrooms, etc.)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
f. Rohypnol (roches, roofies, etc.)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
g. Steroids?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
h. Ecstasy?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
i. Heroin (black tar, cheese, chiva, brown heroin)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
j. Methamphetamine (speed, crystal meth, ice, or crank)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
k. Synthetic Marijuana (spice, K2)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

16. How recently, if ever, have you taken the following PRESCRIPTION DRUGS not prescribed to you or only for the experience or feeling it caused even one time:
(DARKEN ONE BUBBLE FOR EACH LINE: a-e.)

	Never heard of/ Never used it	Used at least once in the Past Month	Used at least once Since School Began in the Fall	Used at least once In Your Lifetime
a. Codeine cough syrup (Lean, Nods, AC/DC)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. OxyContin, Percodan, Percocet, or Oxycodone?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Vicodin, Lortab, Lorcet, or Hydrocodone?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Valium or Diazepam?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. Xanax or Alprazolam?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

SERIAL #



PLEASE DO NOT WRITE IN THIS AREA

17. How recently, if ever, have you taken the following OVER-THE-COUNTER DRUGS only for the experience or feeling it caused even one time:

a. DXM, Triple C's, Skittles, or Coricidin?

Never heard of/Never used it	Used at least once in the Past Month	Used at least once Since School Began in the Fall	Used at least once In Your Lifetime
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

18. IN THE PAST 30 DAYS, how many times, if any, have you used: (DARKEN ONE BUBBLE FOR EACH LINE: a-f.)

	Never heard of/ Not used it	1 - 2 Times	3 - 10 Times	11+ Times
a. Tobacco?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Alcohol?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Inhalants (whiteout, spray paint, glue, gas, etc.) to get high?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Marijuana?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. Cocaine (not crack)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
f. Crack?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

19. About how many of your close friends use: (DARKEN ONE BUBBLE FOR EACH LINE: a-d.)

	Never heard of/ None	A few	Some	Most	All
a. Tobacco?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Alcohol?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Inhalants (whiteout, spray paint, glue, gas, etc.)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Marijuana?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

20. If you wanted some, how difficult would it be to get: (DARKEN ONE BUBBLE FOR EACH LINE: a-l.)

	Never heard of it	Impossible	Very difficult	Somewhat difficult	Somewhat easy	Very easy
a. Tobacco?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Alcohol?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Inhalants (whiteout, spray paint, glue, gas, etc.)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Marijuana?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. Cocaine (not crack)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
f. Crack?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
g. Damaxil?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
h. Steroids?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
i. Ecstasy?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
j. Heroin (black tar, cheese, chiva, brown heroin)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
k. Methamphetamine (speed, crystal meth, ice, or crank)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
l. Synthetic Marijuana (spice, K2)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

21. How old were you when you first used: (DARKEN ONE BUBBLE FOR EACH LINE: a-l.)

	Never used it	8 or younger	9	10	11	12	13	14	15	16	17	18 or older
a. Tobacco?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Alcohol?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Inhalants (whiteout, spray paint, glue, gas, etc.)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Marijuana?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. Cocaine (not crack)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
f. Crack?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
g. Damaxil?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
h. Steroids?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
i. Ecstasy?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
j. Heroin (black tar, cheese, chiva, brown heroin)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
k. Methamphetamine (speed, crystal meth, ice, or crank)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
l. Synthetic Marijuana (spice, K2)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

SERIAL #



PLEASE DO NOT WRITE IN THIS AREA

22. How often do you normally use:
(DARKEN ONE BUBBLE FOR EACH LINE: a-l.)

	Never used it	Every day	Several times a week	Several times a month	About once a month	About once a year	Less than once a year
a. Tobacco?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Alcohol?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Inhalants (whiteout, spray paint, glue, gas, etc.)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Marijuana?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. Cocaine (not crack)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
f. Crack?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
g. Damaxil?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
h. Steroids?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
i. Ecstasy?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
j. Heroin (black tar, cheese, chiva, brown heroin)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
k. Methamphetamine (speed, crystal meth, ice, or crank)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
l. Synthetic Marijuana (spice, K2)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

23. When you drink alcoholic beverages, how many drinks do you usually have AT ONE TIME, on average:
(DARKEN ONE BUBBLE FOR EACH LINE: a-d.)

	Never drink this beverage	12 or more drinks	9 - 11 drinks	5 - 8 drinks	3 - 4 drinks	2 drinks	1 drink	Less than 1 drink
a. Beer	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Wine Coolers (hard lemonade, hard sodas, etc.)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Wine	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Liquor (vodka, rum, whiskey, etc.)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

24. IN THE PAST 30 DAYS, on how many days have you had 5 or more drinks of alcohol AT ONE TIME?

	Never had 5 or more drinks at one time	None	1 day	2 days	3 to 5 days	6 to 9 days	10 or more days
a. Alcohol	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

25. How often do you get alcoholic beverages from the following sources?
(DARKEN ONE BUBBLE FOR EACH LINE: a-e.)

	Do Not Drink	Never	Seldom	Most of the time	Always
a. At home	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. From friends	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. From a store	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. At parties	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. Other source	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

26. How dangerous do you think it is for kids your age to use:
(DARKEN ONE BUBBLE FOR EACH LINE: a-l.)

	Very Dangerous	Somewhat Dangerous	Not Very Dangerous	Not Dangerous At All	Don't Know
a. Tobacco?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Alcohol?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Inhalants (whiteout, spray paint, glue, gas, etc.)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Marijuana?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. Cocaine (not crack)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
f. Crack?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
g. Ecstasy?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
h. Steroids?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
i. Heroin (black tar, cheese, chiva, brown heroin)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
j. Methamphetamine (speed, crystal meth, ice, or crank)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
k. Synthetic Marijuana (spice, K2)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
l. Hydrocodone (Vicodin, Lortab, or Lorcet) not prescribed to you?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

27. SINCE SCHOOL BEGAN IN THE FALL, on how many DAYS have you . . .
(DARKEN ONE BUBBLE FOR EACH LINE: a-e.)

	None	1-3 days	4-9 days	10+ days
a. missed a whole day of school because you "skipped" or "cut"?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. missed a whole day of school because you were ill?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. missed a whole day of school for some other reason?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. been sent by a teacher to someone like the Principal, Dean, or Guidance Counselor because of your conduct or attitude?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. had someone from your home be called to school because of your conduct or attitude?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

28. DURING THE PAST 12 MONTHS, how many times (if any) have you...
(DARKEN ONE BUBBLE FOR EACH LINE: a-f.)

	None	1-3 times	4-9 times	10+ times
a. gotten into difficulties of any kind with your friends because of your drinking?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. driven a car when you've had a good bit to drink?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. been criticized by someone you were dating because of drinking?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. gotten into difficulties of any kind with your friends because of your drug use?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. driven a car when you've felt high from drugs?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
f. been criticized by someone you were dating because of drug use?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

29. SINCE SCHOOL BEGAN IN THE FALL, on how many DAYS (if any) have you attended at least one class while "high," "drunk" or "stoned" on . . .
(DARKEN ONE BUBBLE FOR EACH LINE: a-d.)

	None	1-3 days	4-9 days	10+ days
a. beer, wine coolers, wine, or hard liquor?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. marijuana?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. inhalants?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. some other drug(s)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

30. Alcohol or other drugs are sometimes used at teenage parties. Thinking of the parties you attended this school year . . .
(DARKEN ONE BUBBLE FOR EACH LINE: a-b.)

	Never	Seldom	Half the time	Most of the time	Always	Don't Know	Didn't attend any parties
a. how often was alcohol used?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. how often were marijuana and/or other drugs used?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

31. SINCE SCHOOL BEGAN IN THE FALL, have you sought help, other than from family or friends, for problems in any way connected with your use (if any) of alcohol, marijuana, or other drugs?

	Yes	No
	<input type="radio"/>	<input type="radio"/>

32. If you had a drug or alcohol problem and needed help, who would you go to?
(DARKEN ONE BUBBLE FOR EACH LINE: a-h.)

	Yes	No
a. A counselor or program in school?	<input type="radio"/>	<input type="radio"/>
b. Another adult in school (such as nurse, teacher, or coach)?	<input type="radio"/>	<input type="radio"/>
c. A counselor or program outside of school?	<input type="radio"/>	<input type="radio"/>
d. Your parents?	<input type="radio"/>	<input type="radio"/>
e. A medical doctor?	<input type="radio"/>	<input type="radio"/>
f. Your friends?	<input type="radio"/>	<input type="radio"/>
g. Another adult (such as relative, clergyman, or other family friend)?	<input type="radio"/>	<input type="radio"/>
h. I wouldn't go to anyone.	<input type="radio"/>	<input type="radio"/>

33. SINCE SCHOOL BEGAN IN THE FALL, have you gotten any information on drugs or alcohol from the following SCHOOL sources?
(DARKEN ONE BUBBLE FOR EACH LINE: a-g.)

	Yes	No
a. A school health class?	<input type="radio"/>	<input type="radio"/>
b. An assembly program at school?	<input type="radio"/>	<input type="radio"/>
c. A school guidance counselor?	<input type="radio"/>	<input type="radio"/>
d. A science or social study class?	<input type="radio"/>	<input type="radio"/>
e. A student group or club meeting at school?	<input type="radio"/>	<input type="radio"/>
f. An invited school guest?	<input type="radio"/>	<input type="radio"/>
g. Another source at school?	<input type="radio"/>	<input type="radio"/>

34. How do your parents feel about kids your age. . . (DARKEN ONE BUBBLE FOR EACH LINE: a-c.)

	Strongly Disapprove	Mildly Disapprove	Neither Approve nor Disapprove	Mildly Approve	Strongly Approve	Don't Know
a. using tobacco?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. drinking alcohol?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. using marijuana?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

35. IN THE PAST SCHOOL YEAR, have you been tested for drug use by someone at school? Yes No

36. Do you think drug testing should be required for all student athletes? Yes No Don't Know

SUPPLEMENTAL SURVEY QUESTIONS

S-1 A B C D E F G H I J	S-2 A B C D E F G H I J	S-3 A B C D E F G H I J	S-4 A B C D E F G H I J	S-5 A B C D E F G H I J	S-6 A B C D E F G H I J	S-7 A B C D E F G H I J	S-8 A B C D E F G H I J	S-9 A B C D E F G H I J	S-10 A B C D E F G H I J
S-11 A B C D E F G H I J	S-12 A B C D E F G H I J	S-13 A B C D E F G H I J	S-14 A B C D E F G H I J	S-15 A B C D E F G H I J	S-16 A B C D E F G H I J	S-17 A B C D E F G H I J	S-18 A B C D E F G H I J	S-19 A B C D E F G H I J	S-20 A B C D E F G H I J
S-21 A B C D E F G H I J	S-22 A B C D E F G H I J	S-23 A B C D E F G H I J	S-24 A B C D E F G H I J	S-25 A B C D E F G H I J	S-26 A B C D E F G H I J	S-27 A B C D E F G H I J	S-28 A B C D E F G H I J	S-29 A B C D E F G H I J	S-30 A B C D E F G H I J

THANK YOU FOR BEING PART OF THIS
IMPORTANT PROJECT



PLEASE DO NOT WRITE IN THIS AREA

SERIAL #

Appendix B
Participating Campuses

DISTRICT	CAMPUS
ALICE ISD	ALICE H S
GRAND PRAIRIE ISD	ANDREW JACKSON MIDDLE
APPLE SPRINGS ISD	APPLE SPRINGS H S
ATHENS ISD	ATHENS H S
AVERY ISD	AVERY MIDDLE SCHOOL
BELTON ISD	BELTON H S
BLOOMING GROVE ISD	BLOOMING GROVE H S
BLOOMINGTON ISD	BLOOMINGTON HS
BOERNE ISD	BOERNE SAMUEL V CHAMPION HS
BOWIE ISD	BOWIE H S
BRONTE ISD	BRONTE HS
BROOKELAND ISD	BROOKELAND HS
BROWNFIELD ISD	BROWNFIELD MIDDLE
CAMERON ISD	CAMERON YOE HIGH SCHOOL
LEANDER ISD	CANYON RIDGE MIDDLE SCHOOL
CENTRAL HEIGHTS ISD	CENTRAL HEIGHTS H S
BROWNSBORO ISD	CHANDLER INTERMEDIATE
CHILDRESS ISD	CHILDRESS J H
COMANCHE ISD	COMANCHE H S
CONNALLY ISD	CONNALLY HIGH SCHOOL
CYPRESS-FAIRBANKS ISD	COOK MIDDLE SCHOOL
COPPERAS COVE ISD	COPPERAS COVE H S
COTTON CENTER ISD	COTTON CENTER SCHOOL
CYPRESS-FAIRBANKS ISD	CYPRESS WOODS HS
DAWSON ISD	DAWSON SCHOOL
YSLETA ISD	DEL VALLE HS
HIDALGO ISD	DIAZ J H
DONNA ISD	DORA M SAUCEDA MIDDLE
WYLIE ISD	DR AL DRAPER INTERMED
SPRING ISD	DUEITT MIDDLE
ECTOR COUNTY ISD	ECTOR J H
EDNA ISD	EDNA JUNIOR HIGH SCHOOL
GILMER ISD	GILMER INT
ALDINE ISD	GRANTHAM ACADEMY
RIO GRANDE CITY CISD	GRULLA MIDDLE
BROWNSVILLE ISD	HANNA H S
HARLINGEN CISD	HARLINGEN H S
HAWKINS ISD	HAWKINS MIDDLE
HEREFORD ISD	HEREFORD H S
IOLA ISD	IOLA H S

JACKSBORO ISD	JACKSBORO MIDDLE
CYPRESS-FAIRBANKS ISD	JERSEY VILLAGE HIGH SCHOOL
KARNES CITY ISD	KARNES CITY H S
KERMIT ISD	KERMIT H S
KLEIN ISD	KLEIN FOREST H S
LEANDER ISD	KNOX WILEY MIDDLE SCHOOL
SAN ANGELO ISD	LEE MIDDLE SCHOOL
LOCKHART ISD	LOCKHART J H
FRISCO ISD	LONE STAR H S
LAKE WORTH ISD	LUCYLE COLLINS MIDDLE SCHOOL
LUMBERTON ISD	LUMBERTON INT
LUMBERTON ISD	LUMBERTON MIDDLE
MAY ISD	MAY EL
PORT ARTHUR ISD	MEMORIAL 9TH GRADE ACADEMY AT AUST
MOUNT ENTERPRISE ISD	MT ENTERPRISE H S
NEW CANEY ISD	NEW CANEY H S
ALIEF ISD	OWENS INT
PASADENA ISD	PASADENA HIGH SCHOOL
POTEET ISD	POTEET H S
PRAIRILAND ISD	PRAIRILAND H S
QUITMAN ISD	QUITMAN JH
KILLEEN ISD	ROBERT M SHOEMAKER H S
AZLE ISD	SANTO FORTE JUNIOR HIGH SCHOOL
SHARYLAND ISD	SHARYLAND NORTH J H
CALHOUN COUNTY ISD	TRAVIS MIDDLE
UNITED ISD	UNITED H S
VERNON ISD	VERNON H S
WESLACO ISD	WESLACO H S
WHITNEY ISD	WHITNEY H S
YSLETA ISD	YSLETA MS
DUMAS ISD	DUMAS J H
CROSBYTON CISD	CROSBYTON H S
DE LEON ISD	PERKINS MS
LAMAR CISD	READING JH
HITCHCOCK ISD	HITCHCOCK H S
ROCHELLE ISD	ROCHELLE SCHOOL
CROCKETT ISD	CROCKETT J H
LAREDO ISD	NIXON H S
LA FERIA ISD	NOEMI DOMINGUEZ EL
SAN BENITO CISD	MILLER JORDAN MIDDLE
MINEOLA ISD	MINEOLA H S
MINEOLA ISD	MINEOLA MIDDLE
GALENA PARK ISD	NORTH SHORE SENIOR HIGH SCHOOL

GALENA PARK ISD	GALENA PARK HS
PAMPA ISD	Pampa HS
PALESTINE ISD	Palestine HS
PALESTINE ISD	STORY INT
GRAPEVINE-COLLEYVILLE ISD	Colleyville Middle School
CORSICANA ISD	Corsicana HS
EAGLE MT-SAGINAW ISD	Wayside Middle School
EAGLE MT-SAGINAW ISD	Ed Wilkie Middle School
EAGLE MT-SAGINAW ISD	Boswell High School
POINT ISABEL ISD	Port Isabel JH
KELLER ISD	INDIAN SPRINGS MIDDLE
KELLER ISD	KELLER MIDDLE
KELLER ISD	TRINITY SPRINGS MIDDLE
NIXON-SMILEY CISD	NIXON-SMILEY HS
NIXON-SMILEY CISD	NIXON-SMILEY MS
PATTON SPRINGS ISD	PATTON SPRINGS SCHOOL
GRAPE CREEK ISD	GRAPE CREEK H S
GRAPE CREEK ISD	GRAPE CREEK MIDDLE
KRESS ISD	KRESS HS
EUSTACE ISD	Eustace HS
WELLS ISD	Wells HS
LA JOYA ISD	La Joya HS
LA JOYA ISD	Juan De Dios Salinas MS

Appendix C
Classroom Selection Guidelines

Texas School Survey of Drug and Alcohol Use

Sampling Guide for Survey Administration

For the Texas School Survey to be administered in your school/district, PPRI will need to randomly select classes for participation in the study. The random selection process will allow us to generate survey results that are representative of the student population within your school/district. Detailed instructions for completing this master list are noted below.

If you are surveying students at an **individual school only, you will need to provide a master list of classes for the specific grade levels contained at your school (only grades 6-12 are eligible to participate).

**If you are surveying students at the district level, you will need to provide a master list of classes at each district campus with grades 6-12.

Instructions:

The Survey Coordinator will receive this guide (*Sampling Guide for Survey Administration*) and a Microsoft Excel spreadsheet titled *Sampling Template*. Please use this guide (*Sampling Guide for Survey Administration*) when completing the *Sampling Template*.

- For schools conducting the survey via paper pencil, choose *either* a **class period** (i.e. 1st, 3rd) in which all students are present or a **subject** (i.e. English, Science) in which all students must be enrolled;
- For schools conducting the survey online, choose only a subject (i.e. English, Science) in which all students must be enrolled).

The Excel spreadsheet titled *Sampling Template*, has column and row titles to indicate where to insert the necessary information. Using the Excel spreadsheet, please complete the following steps:

1. List all teachers for that class period or that subject in the first column;
2. List the class period, regardless of whether class period or subject was selected.
3. List the number of students per grade level in each of those classes.

Please submit the Excel spreadsheet electronically to: txschoolsurvey@ppri.tamu.edu. PPRI will randomly pick entire classes until we reach the target numbers per grade level for the sample. Then, we'll send you the Master List of those classes randomly selected to be surveyed in order for the Parental Notification Letters to be sent home to the parents of students in the selected classes.

Appendix D
School Survey Administration
Survey Coordinator's Responsibilities
Campus Coordinators' Responsibilities
Teachers' Guide

Texas School Survey of Drug and Alcohol Use
Survey Coordinator's Responsibilities- Paper-Pencil Administration

1. Distribution of the Survey Administration Guide

The *Survey Administration Guide* enclosed in this mailing is designed to instruct the individuals who will administer the *Texas School Survey* at each campus. Each individual involved in the survey administration should receive a copy. The guides should be sent to the campuses in the time to be distributed one week prior to the scheduled survey administration date. This will allow time for everyone involved to read through the information and resolve any questions.

2. Distribution of Survey Instruments and Classroom Envelopes

Enclosed is a set of envelopes and survey instruments. There should be approximately one envelope per classroom and a sufficient number of survey instrument forms as requested by the Survey Coordinator. If pre-packaged materials were requested, classroom envelopes will contain 30 blank survey instruments and a copy of the *Survey Administration Guide*. If classrooms were randomly selected by PPRI for participation, all survey materials will be pre-packaged according to campus name, teacher name, and classroom enrollment numbers. The envelopes will be used to package the blank survey forms sent to each class and seal the completed instruments after survey administration (see Campus Coordinators' Responsibilities).

At least one week prior to the scheduled survey date, Survey Coordinators should forward the appropriate number of survey instruments and envelopes to each campus. Each campus will need as many envelopes as there are classes participating. In case of mass administration, approximately one envelope per 24 students should be used.

3. Packaging the Completed Surveys for Delivery to PPRI

Completed survey instruments will be sealed in envelopes and collected from each classroom. As the Survey Coordinator, you should clearly designate to each Campus Coordinator where the surveys should be sent from there. All completed surveys should be collected at a central location. Completed surveys should remain in the sealed envelopes, and the envelopes from each campus should be grouped together and **CLEARLY LABELED** with the campus name. **BLANK SURVEYS SHOULD BE** returned to PPRI along with the classroom envelopes. All survey materials should be boxed, and the boxes should be labeled sequentially (i.e., "one of three, two of three," etc.). Materials should then be returned to PPRI in a single mailing. Ship to:

Texas School Survey
PPRI / Texas A&M University
3833 Texas Avenue South, Suite 130
Bryan, TX 77802

Texas School Survey of Drug and Alcohol Use
Survey Coordinator's Responsibilities- Online Administration

1. Distribution of the Online Survey Administration Guide

The *Online Survey Administration Guide* enclosed in this mailing is designed to instruct the individuals who will administer the *Texas School Survey* at each campus. Each individual involved in the survey administration should receive a copy.

The guides should be sent to the campuses in the time to be distributed one week prior to the scheduled survey administration date. This will allow time for everyone involved to read through the information and resolve any questions.

2. Distribution of Online Survey Tokens and Classroom Envelopes

Enclosed is a set of envelopes and online survey tokens. These survey tokens are unique numbers printed on a small piece of paper that each student will need to log in to the online survey. There should be approximately one envelope per classroom and a sufficient number of online survey tokens as requested by the Survey Coordinator. If pre-packaged materials were requested, classroom envelopes will contain 30 online survey tokens and a copy of the *Online Survey Administration Guide*. If classrooms were randomly selected by PPRI for participation, all survey materials will be pre-packaged according to campus name, teacher name, and classroom enrollment numbers. The envelopes will be used to package the online survey tokens sent to each class and seal the online survey tokens after survey administration (see Campus Coordinators' Responsibilities).

At least one week prior to the scheduled survey date, Survey Coordinators should forward the appropriate number of survey tokens, classroom envelopes, and *Online Survey Administration Guides* to each campus. Each campus will need as many envelopes as there are classes participating.

3. Packaging the Survey Materials for Delivery to PPRI

The online survey tokens will be sealed in envelopes and collected from each classroom. As the Survey Coordinator, you should clearly designate to each Campus Coordinator where the classroom envelopes should be sent from there. All classroom envelopes should be collected at a central location. The classroom envelopes from each campus should be grouped together and **CLEARLY LABELED** with the campus name. **UNUSED ONLINE SURVEY TOKENS SHOULD BE** returned to PPRI along with the classroom envelopes.

All survey materials should be boxed, and the boxes should be labeled sequentially (i.e., "one of three, two of three," etc.). Materials should then be returned to PPRI in a single mailing. Ship to:

Texas School Survey
PPRI / Texas A&M University
3833 Texas Avenue South, Suite 130
Bryan, TX 77802

*Texas School Survey of Drug and Alcohol Use
Campus Coordinators' Responsibilities- Paper-Pencil Administration*

1. Distribution of the Guide to Survey Administration

The *Survey Administration Guide* is designed to instruct the individuals who will administer the *Texas School Survey* at each campus. Each individual involved in the survey administration should receive a copy. All survey packets should have a copy of the *Survey Administration Guide and Procedure* enclosed and additional copies have been provided to the Survey Coordinator. The Guide should be distributed to all survey administrators (i.e. teachers) approximately one week prior to the scheduled administration date so that everyone will have adequate time to read through the information and resolve any questions.

2. Distribution of Survey Instruments to Classes

The Survey Coordinator will send to each participating campus the appropriate number of survey instruments and enough envelopes to provide one to each class being surveyed. If the materials are not already pre-packaged in the classroom envelopes by campus and teacher name, Campus Coordinators should follow these steps:

- a) Count out a set of surveys for the appropriate number of students in each participating class;
- b) Place each set of surveys in a classroom envelope; and
- c) Label each classroom envelope to indicate which class it is along with **the campus name**. Do **not** seal the envelopes;
- d) Ensure each classroom envelope contains a copy of the *Survey Administration Guide and Procedure*.

On the day the survey is to be given, envelopes filled with the correct number of survey instruments and a copy of the *Survey Administration Guide and Procedure* should be distributed to each survey administrator. The individuals administering the survey should have reviewed the *Survey Administration Guide and Procedure* (described above) prior to this point and should already be aware of their responsibilities.

3. Collecting Completed Survey Instruments

After the survey has been administered in each class, teachers will be instructed to seal the completed survey forms in the envelopes. Unused blank forms should be kept together **OUTSIDE THE ENVELOPE**. It is recommended that teachers not leave the room with the sealed envelopes. Rather, someone who did not participate in the actual survey administration should be sent around to collect the envelopes.

Once envelopes have been collected from every classroom, they should be forwarded to the location designated by the Survey Coordinator where they will be collected and returned to PPRI in a **single mailing**.

Thank you very much for your assistance with this important research project.

*Texas School Survey of Drug and Alcohol Use
Campus Coordinators' Responsibilities- Online Administration*

1. Distribution of the Guide to Online Survey Administration

The *Online Survey Administration Guide* is designed to instruct the individuals who will administer the *Texas School Survey* at each campus. Each individual involved in the survey administration should receive a copy. All survey packets should have a copy of the *Online Survey Administration Guide* enclosed and additional copies have been provided to the Survey Coordinator. The Guide should be available for all survey administrators (i.e. teachers) to review approximately one week prior to the scheduled administration date so that everyone will have adequate time to read through the information and resolve any questions.

2. Distribution of Online Survey Tokens to Classes

The Survey Coordinator will send to each participating campus the appropriate number of online survey tokens and enough classroom envelopes to provide one to each class being surveyed. These survey tokens are unique numbers printed on a small piece of paper that each student will need to log in to the online survey. If the materials are not already pre-packaged in the classroom envelopes by campus and teacher name, Campus Coordinators should follow these steps:

- a) Count out a set of online survey tokens for the appropriate number of students in each participating class;
- b) Place each set of online survey tokens in a classroom envelope;
- c) Label each classroom envelope to indicate which class it is along with **the campus name**. Do **not** seal the envelopes.
- d) Ensure each classroom envelope contains a copy of the *Online Survey Administration Guide and Procedure*

On the day the survey is to be given, envelopes filled with the online survey tokens and a copy of the *Online Survey Administration Guide and Procedure* should be distributed to each survey administrator. The individuals administering the survey should have reviewed the *Online Survey Administration Guide and Procedure* (described above) prior to this point and should already be aware of their responsibilities.

3. Collecting Survey Materials

After the survey has been administered in each class, teachers will be instructed to seal all used and unused survey tokens in the envelopes. It is recommended that teachers not leave the room with the sealed envelopes. Rather, someone who did not participate in the actual survey administration should be sent around to collect the envelopes.

Once envelopes have been collected from every classroom, they should be forwarded to the location designated by the Survey Coordinator where they will be collected and returned to PPRI in a **single mailing**.

Thank you very much for your assistance with this important research project.

Texas School Survey of Drug and Alcohol Use
Survey Administration Guide- Paper-Pencil Administration

Your school has elected to participate in the *Texas School Survey of Drug and Alcohol Use*. The survey is sponsored and designed by the Texas Department of State Health Services (DSHS) and is facilitated by the Public Policy Research Institute (PPRI) at Texas A&M University. The school needs your assistance in administering the survey. PPRI has prepared this Survey Administration Guide to provide information and instructions for conducting the survey project.

- Your classroom envelope should contain blank surveys and an attached Classroom Identification Form to be filled out during survey administration. You will find this Classroom Identification Form attached as a label to your classroom envelope. It is very important that you write the **campus name** (if it is not already written on the label) and **number of students enrolled in your class** on this label.
- Please read all instructions carefully before administering the survey. It is important that these procedures are followed so that administration of the survey is uniform throughout the school.
- Your attitude towards this survey is extremely important. If students sense the importance of this information, they will take the survey more seriously. Please stress to the students that their responses are **entirely anonymous**.
- In addition, your physical position in the classroom may influence the way students answer questions. While students are taking the survey, do not walk around the room. Please sit or stand in a part of the room away from students.
- Try to keep students as quiet as possible throughout survey administration. If students are allowed to talk about the contents of the survey or their own experiences, peer influence could cause students to either inflate or under-report their own drug and alcohol habits.
- The survey should take approximately 30-45 minutes to administer.
- To maintain student confidentiality, **do not view completed survey instruments**.
- Students must use a #2 pencil. Be prepared to provide pencils to students who may not have one. Surveys completed in pen or marker will not be processed.

Thank you very much for your assistance with this important research project.

Texas School Survey of Drug and Alcohol Use
Survey Administration Procedure- Paper-Pencil Administration

1.) Before handing out surveys, please tell students the following:

“This survey is for the Public Policy Research Institute at Texas A&M University. Survey results will be used for evaluating and planning substance abuse programs for students in grades 6 through 12. No student is required to participate in this survey. This survey will ask questions about your experiences with drugs and alcohol. Do not turn the pages or begin answering questions until we have gone over the survey instructions.”

2.) Make sure all students have a #2 pencil to complete the survey. Hand out the surveys and please tell students the following:

I cannot answer questions about specific substances or tell you what questions mean. If you are not familiar with a substance listed on the survey, mark the category “Never heard of it.”

3.) Go over the instructions listed on the front of the survey booklet. Be sure to read aloud the Introduction, Directions, and Example Question.

4.) Please emphasize the following points to the students:

- This survey is completely voluntary.
- No one will know how individual students answer questions.
- If you do not feel comfortable answering a question or feel you cannot answer a question honestly, leave it blank.
- A drink means a can or bottle of beer or wine cooler, a 4 ounce glass of wine, a shot of liquor, or a mixed drink.

5.) If students refuse to take the survey, have them work quietly at their seat or send them to an area designated by your campus coordinator.

6.) Collect all completed surveys and place in envelope. Please do not view completed surveys instruments. **Fill out the attached Classroom Identification Form** which is attached as a label to the front of your classroom envelope. It is very important that the **campus name** and **number of students enrolled in your class** is written on this label. PLEASE DO NOT PUT UNUSED SURVEYS IN THE ENVELOPE. Seal the envelope.

7.) Return classroom envelope and unused surveys to your campus coordinator.

Thank you very much for your assistance with this important research project.

Texas School Survey of Drug and Alcohol Use
Survey Administration Guide- Online Administration

Your school has elected to participate in the *Texas School Survey of Drug and Alcohol Use*. The survey is sponsored and designed by the Texas Department of State Health Services (DSHS) and is facilitated by the Public Policy Research Institute (PPRI) at Texas A&M University. The school needs your assistance in administering the survey. PPRI has prepared this Online Survey Administration Guide to provide information and instructions for conducting the survey project.

- Your classroom envelope should contain a small piece of paper for each student with a number, which is an online survey token. The survey tokens are unique numbers that each student will need to log in to the online survey.
- You will also find a label on the front of your classroom envelope titled ‘Classroom Identification Form’. This form will need to be filled out during survey administration. It is very important that you write the **campus name** (if it is not already written on the label) and **number of students enrolled in your class** on this label.
- Please read all instructions carefully before administering the survey. It is important that these procedures are followed so that administration of the survey is uniform throughout the school.
- Your attitude towards this survey is extremely important. If students sense the importance of this information, they will take the survey more seriously. Please stress to the students that their responses are **entirely anonymous**. The tokens are completely random and there is no way to know which student responded to a particular survey.
- In addition, your physical position in the classroom may influence the way students answer questions. While students are taking the survey, do not walk around the room. Please sit or stand in a part of the room away from students.
- Try to keep students as quiet as possible throughout survey administration. If students are allowed to talk about the contents of the survey or their own experiences, peer influence could cause students to either inflate or under-report their own drug and alcohol habits.
- The survey should take approximately 30-45 minutes to administer.

Thank you very much for your assistance with this important research project.

*Texas School Survey of Drug and Alcohol Use
Survey Administration Procedure- Online Administration*

- 1.) Have all participating students enter the following address into their web browser:
<http://txss.tamu.edu>
- 2.) **Before handing out the online survey tokens, please tell students the following:** “This survey is for the Public Policy Research Institute at Texas A&M University. Survey results will be used for evaluating and planning substance abuse programs for students in grades 6 through 12. No student is required to participate in this survey. This survey will ask questions about your experiences with drugs and alcohol. Please do not begin answering questions until we have gone over the survey instructions.”
- 3.) Hand out the online survey tokens and please tell students the following:
 - This is a survey that measures tobacco, drug, and alcohol use and other information relating to tobacco, alcohol, inhalant, and drug use;
 - This survey is being given to as many as 250,000 students in the state of Texas;
 - There is no way for anyone to know how you respond to the questions;
 - You can skip any questions or not take the survey at all if you do not want to. There is no penalty if you skip questions or leave the survey blank;
 - On the survey, a drink means a can or bottle of beer or wine cooler, a 4 ounce glass of wine, a shot of liquor, or a mixed drink;
 - I cannot answer questions about specific substances or tell you what questions mean;
 - If you are not familiar with a substance listed, mark the category “Never heard of it.”
 - **At the end of the survey, please make sure you click the ‘Submit’ button**

If you feel you need to talk with someone about problems with tobacco, inhalants, alcohol, or drugs, please call, toll free, 1-877-9-NO-DRUG (1-877-966-3784) for immediate and confidential help, 24 hours a day, seven days a week. This study has been reviewed and approved by the Institutional Review Board - Human Subjects in Research, Texas A&M University. For research-related questions regarding subjects' rights, the Institutional Review Board may be contacted at (979) 458-4067 or irb@tamu.edu.

- 6.) Tell the students to type the number on their paper in the token box and click ‘ENTER’. The students will then need to read the instructions and click ‘NEXT’ to proceed to survey questions.
- 7.) If students refuse to take the survey, have them work quietly at their seat or send them to an area designated by your campus coordinator.
- 8.) Collect all papers with the survey tokens and place in envelope. **Please fill out the attached Classroom Identification Form** which is attached as a label to the front of your classroom envelope. It is very important that the **campus name and number of students enrolled in your class** is written on this label. Seal the envelope and return classroom envelope to your campus coordinator.

Thank you very much for your assistance with this important research project.

Appendix E
Computation of Sample Weights

Weighting the Texas School Survey

- The weights for the Texas School Survey of Substance Abuse are calculated in the following manner: State Sample
 - Campus Weight
 - Probability of each school being sampled
 - A =Total number of “adjusted” students in school (over/under-sampled)³
 - B =total “adjusted” population in region
 - D =total number of schools sampled
 - School’s probability $X=(AxD)/B$
 - Raw weight
 - $H= 1/X$
 - Sample weight adjustment
 - A =Total number of “adjusted” students in school (over/under-sampled)
 - J = Total number of actual students in school
 - Adjustment $K=J/A$
 - Preliminary Weight
 - Preliminary Weight $P= K*H$
 - Population Weight Adjustment
 - L =Total population (In Region)
 - Q = Total population (In Region) after applying Weight (P) for participating schools
 - Population Weight Adjustment $R=Q/L$
 - Campus Weight
 - Campus Weight $Z = P/R$
- Student Weight
 - Probability of each Student sampled in school
 - G =Number of students in grade
 - C = Number of students sampled in grade
 - Student Weight $Y=G/C$
 - Weight
 - Weight $W=Y*Z$
- Post Hoc Weights
 - Adjustment for grade distribution
 - U = State population in grade
 - V = State sample/grade population after applying weight W
 - Grade Weight Adjustment $GW=W*(U/V)$
 - Adjustment for race/ethnicity
 - S =Region Race/ethnicity population
 - T = Region Race/ethnicity population after applying weight GW
 - Final Weight $rweight=GW*(S/T)$

³ “(over/under-sampled)” refers to the increased probability attached to students in certain schools such as border schools and junior high campuses as discussed in the “Selection of Campuses” section of the report.

Appendix F

Standard Error Computation

Standard Error Computation for the Texas School Survey of Substance Use

The standard errors of the percentages were estimated using the SURVEYFREQ procedure in the Statistical Analysis System (SAS) software package.¹ The variance estimation techniques used in the program are specifically designed to address the complexity of the sample design involving multistage, stratified, cluster samples. A Taylor series estimation of the variance is used to determine the standard error of the proportion of each response. An example of the coding for the program is listed below. A subgroup is defined by a variable identifying separate parts of the sample for which the errors are to be separately estimated. When the error is to be estimated for the entire sample, the subgrouping is omitted. The variable grade is the strata and the cluster is the variable school. The weight is the final sample weight assigned to each observation which adjusts for both the different sample rates involved in the design and the post hoc weights for ethnicity. The var is the variable for which the error is estimated. In the estimates presented here the variables are dichotomous, in the category or not. This is appropriate since the categories of a prevalence category are not mutually exclusive. For example, the category of “lifetime use” includes all those who responded use in the “past month”, “school year”, as well as “lifetime”.

Detailed description of the technique, references, and use of the program can be found in the SAS on-line documentation (SAS/STAT(R) 9.22 User's Guide. (n.d.). Retrieved March 26, 2015, from

http://support.sas.com/documentation/cdl/en/statug/63347/HTML/default/viewer.htm#surveyfreq_toc.htm). The program can be used to produce error estimates, tests of significance, and analysis for other research questions using this data.

Example code:

```
proc surveyfreq data=name;
    by subgrouping (if any);
    weight weight;
    strata grade;
    cluster school;
    tables var;
```

¹ Estimates of the percentages in this report may differ from the Final Report slightly due to rounding differences between SAS and Stata, the software utilized to produce the Final Report.

Appendix G

Standard Error & Confidence Interval Tables

Lifetime Use of Selected Substances - Overall				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	19.9	18.3	21.6	0.85
CIGARETTES	17.7	16.1	19.2	0.8
SMOKELESS TOBACCO	8	7.2	8.7	0.39
ALCOHOL	46.5	44.1	48.9	1.24
BEER	35.1	32.9	37.3	1.11
WINE COOLER	28.6	26.6	30.7	1.05
WINE	28.9	27.2	30.6	0.86
LIQUOR	30.7	28.3	33.1	1.23
ANY ILLICIT DRUG	21.3	19.4	23.1	0.93
INHALANTS	12.3	11.5	13.1	0.42
MARIJUANA	20.4	18.5	22.2	0.94
COCAINE OR CRACK	2	1.8	2.3	0.14
MARIJUANA ONLY	12.2	11	13.4	0.61
COCAINE	1.8	1.5	2	0.13
CRACK	0.5	0.4	0.6	0.06
HALLUCINOGEN	2.2	1.9	2.6	0.18
ROHYPNOL	0.8	0.6	0.9	0.08
STEROIDS	1	0.8	1.1	0.07
ECSTASY	2.3	2	2.7	0.19
HEROIN	0.4	0.3	0.5	0.04
METHAMPHETAMINE	0.6	0.5	0.8	0.07
SYNTHETIC MARIJ	5.7	5.1	6.4	0.31
CODEINE	10	9.3	10.8	0.39
OXYCONTIN ETC	2.2	1.9	2.6	0.17
VICODIN ETC	4.5	3.8	5.2	0.36
VALIUM ETC	0.9	0.8	1.1	0.08
XANAX ETC	2.8	2.4	3.2	0.21
DXM, ETC.	3.3	2.9	3.7	0.2

Past Month Use of Selected Substances - Overall				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	7.4	6.6	8.2	0.4
CIGARETTES	5.8	5.1	6.5	0.34
SMOKELESS TOBACCO	3.3	2.9	3.7	0.21
ALCOHOL	19.1	17.6	20.6	0.76
BEER	12.1	10.9	13.4	0.61
WINE COOLER	10.8	9.8	11.9	0.52
WINE	8.2	7.4	8.9	0.36
LIQUOR	11.8	10.6	12.9	0.58
ANY ILLICIT DRUG	8.6	7.7	9.5	0.46
INHALANTS	4.2	3.7	4.6	0.23
MARIJUANA	7.9	7	8.8	0.44
COCAINE OR CRACK	0.6	0.5	0.8	0.06
MARIJUANA ONLY	3.7	3.3	4.2	0.22
COCAINE	0.5	0.4	0.6	0.06
CRACK	0.2	0.1	0.2	0.03
HALLUCINOGEN	0.7	0.5	0.8	0.07
ROHYPNOL	0.3	0.2	0.4	0.04
STEROIDS	0.3	0.3	0.4	0.04
ECSTASY	0.6	0.4	0.8	0.1
HEROIN	0.2	0.1	0.2	0.03
METHAMPHETAMINE	0.2	0.1	0.3	0.03
SYNTHETIC MARIJ	1.5	1.3	1.8	0.12
CODEINE	4.8	4.3	5.2	0.21
OXYCONTIN ETC	1.2	1	1.4	0.1
VICODIN ETC	2.2	1.8	2.5	0.17
VALIUM ETC	0.5	0.4	0.6	0.05
XANAX ETC	1.4	1.2	1.6	0.11
DXM, ETC.	1.5	1.2	1.7	0.12

Lifetime Use of Selected Substances - Overall African American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	14.4	12.1	16.8	1.19
CIGARETTES	12.7	10.5	15	1.15
SMOKELESS TOBACCO	4.3	3.1	5.5	0.6
ALCOHOL	46.2	42.4	50.1	1.96
BEER	26.3	23.4	29.3	1.5
WINE COOLER	31	27.2	34.7	1.9
WINE	29.8	27.2	32.4	1.32
LIQUOR	30.4	26.6	34.3	1.96
ANY ILLICIT DRUG	23.4	19.8	27	1.82
INHALANTS	10.2	8.7	11.6	0.74
MARIJUANA	22.7	19.1	26.3	1.81
COCAINE OR CRACK	0.8	0.4	1.2	0.19
MARIJUANA ONLY	17.1	14.4	19.9	1.39
COCAINE	0.7	0.3	1.1	0.19
CRACK	0.2	0.1	0.4	0.08
HALLUCINOGEN	1.4	0.7	2	0.32
ROHYPNOL	0.3	0.1	0.5	0.1
STEROIDS	1.3	0.8	1.8	0.25
ECSTASY	1.6	1	2.2	0.3
HEROIN	0.3	0.1	0.6	0.13
METHAMPHETAMINE	0.9	0.2	1.7	0.36
SYNTHETIC MARIJ	4	2.8	5.2	0.61

Past Month Use of Selected Substances - Overall African American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	4.1	2.9	5.3	0.62
CIGARETTES	3.3	2.3	4.3	0.5
SMOKELESS TOBACCO	1.8	1	2.6	0.41
ALCOHOL	17.1	14.6	19.5	1.23
BEER	7.2	5.7	8.8	0.8
WINE COOLER	10.3	8.1	12.6	1.15
WINE	7.8	6.3	9.3	0.74
LIQUOR	11.1	9.1	13.1	1.02
ANY ILLICIT DRUG	9	7.3	10.6	0.86
INHALANTS	3.5	2.3	4.6	0.57
MARIJUANA	8.1	6.5	9.7	0.82
COCAINE OR CRACK	0.3	0.1	0.6	0.12
MARIJUANA ONLY	5.2	4.2	6.3	0.54
COCAINE	0.3	0.1	0.5	0.11
CRACK	0.1	0	0.3	0.07
HALLUCINOGEN	0.5	0.2	0.7	0.14
ROHYPNOL	0.1	0	0.2	0.06
STEROIDS	0.5	0.1	0.8	0.16
ECSTASY	0.6	0.2	1	0.2
HEROIN	0.2	0	0.4	0.09
METHAMPHETAMINE	0.1	0	0.2	0.06
SYNTHETIC MARIJ	1.2	0.5	1.8	0.35

CODEINE	13.1	10.9	15.2	1.07
OXYCONTIN ETC	2.1	1.3	2.9	0.4
VICODIN ETC	4.3	2.9	5.8	0.73
VALIUM ETC	0.6	0.2	0.9	0.16
XANAX ETC	2.2	1.5	2.8	0.33
DXM, ETC.	2.7	1.8	3.6	0.46

CODEINE	5.9	4.7	7	0.58
OXYCONTIN ETC	1.1	0.6	1.6	0.25
VICODIN ETC	1.9	1.1	2.6	0.38
VALIUM ETC	0.4	0.1	0.7	0.14
XANAX ETC	1.1	0.7	1.6	0.23
DXM, ETC.	1.1	0.5	1.6	0.27

Lifetime Use of Selected Substances - Overall Mexican American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	21.8	20.3	23.3	0.76
CIGARETTES	19.8	18.3	21.2	0.74
SMOKELESS TOBACCO	8	7.2	8.7	0.39
ALCOHOL	49.2	47.1	51.3	1.07
BEER	39.1	37.1	41.2	1.04
WINE COOLER	29.6	27.7	31.6	0.98
WINE	28.6	27.1	30.2	0.79
LIQUOR	32.2	29.9	34.4	1.13
ANY ILLICIT DRUG	24.3	22.6	26.1	0.89
INHALANTS	14.3	13.4	15.2	0.48
MARIJUANA	23.2	21.5	25	0.89
COCAINE OR CRACK	3.1	2.7	3.5	0.21
MARIJUANA ONLY	13.3	12.2	14.5	0.61
COCAINE	2.7	2.3	3.1	0.21
CRACK	0.8	0.6	1	0.11
HALLUCINOGEN	2.1	1.8	2.5	0.18
ROHYPNOL	1.2	1	1.5	0.12
STEROIDS	0.9	0.7	1.1	0.1
ECSTASY	2.8	2.2	3.3	0.27
HEROIN	0.5	0.4	0.6	0.07
METHAMPHETAMINE	0.7	0.5	0.9	0.09
SYNTHETIC MARIJ	6.9	6.2	7.5	0.34
CODEINE	10.6	9.7	11.5	0.47
OXYCONTIN ETC	2	1.7	2.4	0.18
VICODIN ETC	3.9	3.3	4.5	0.31
VALIUM ETC	1.1	0.8	1.3	0.12
XANAX ETC	3	2.4	3.6	0.29
DXM, ETC.	3.8	3.3	4.3	0.24

Past Month Use of Selected Substances - Overall Mexican American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	7.9	7.1	8.6	0.39
CIGARETTES	6.4	5.7	7.1	0.36
SMOKELESS TOBACCO	3.1	2.7	3.6	0.23
ALCOHOL	21.3	19.9	22.7	0.72
BEER	14.3	13.1	15.5	0.63
WINE COOLER	12.1	11.1	13.2	0.54
WINE	9	8.3	9.8	0.37
LIQUOR	12.5	11.3	13.7	0.6
ANY ILLICIT DRUG	10.1	9.2	11.1	0.5
INHALANTS	5.3	4.8	5.9	0.29
MARIJUANA	9.2	8.2	10.1	0.49
COCAINE OR CRACK	1	0.8	1.2	0.1
MARIJUANA ONLY	4.3	3.7	4.8	0.26
COCAINE	0.8	0.6	1	0.1
CRACK	0.3	0.1	0.4	0.06
HALLUCINOGEN	0.8	0.5	1	0.12
ROHYPNOL	0.5	0.3	0.6	0.07
STEROIDS	0.3	0.2	0.4	0.05
ECSTASY	0.8	0.5	1.1	0.15
HEROIN	0.2	0.1	0.3	0.04
METHAMPHETAMINE	0.3	0.2	0.4	0.06
SYNTHETIC MARIJ	2.2	1.8	2.5	0.17
CODEINE	5	4.4	5.5	0.27
OXYCONTIN ETC	0.9	0.7	1.1	0.1
VICODIN ETC	1.9	1.5	2.2	0.17
VALIUM ETC	0.5	0.4	0.7	0.08
XANAX ETC	1.4	1.2	1.7	0.14
DXM, ETC.	1.8	1.5	2.1	0.15

Lifetime Use of Selected Substances - Overall White				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	20.5	17.9	23.2	1.34
CIGARETTES	17.5	15.2	19.9	1.18
SMOKELESS TOBACCO	9.8	8.4	11.2	0.71
ALCOHOL	44.3	40.5	48.1	1.94
BEER	34.2	30.8	37.5	1.71
WINE COOLER	28	24.8	31.2	1.63
WINE	29.3	26.5	32.2	1.44
LIQUOR	30.1	26.4	33.9	1.89
ANY ILLICIT DRUG	17.2	14.8	19.6	1.22
INHALANTS	10.5	9.4	11.7	0.57
MARIJUANA	16.6	14.2	18.9	1.21
COCAINE OR CRACK	1.2	0.9	1.6	0.17
MARIJUANA ONLY	9.7	8.2	11.1	0.74
COCAINE	1.1	0.8	1.4	0.16
CRACK	0.3	0.2	0.4	0.07
HALLUCINOGEN	2.6	2.1	3.2	0.28
ROHYPNOL	0.4	0.2	0.6	0.09

Past Month Use of Selected Substances - Overall White				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	8.4	7.2	9.7	0.65
CIGARETTES	6.1	5.1	7.1	0.51
SMOKELESS TOBACCO	4.2	3.5	4.9	0.37
ALCOHOL	17.4	15.2	19.6	1.12
BEER	11.6	9.8	13.4	0.92
WINE COOLER	9.9	8.4	11.5	0.78
WINE	7.2	6.2	8.3	0.54
LIQUOR	11.6	9.8	13.5	0.93
ANY ILLICIT DRUG	6.7	5.5	7.8	0.58
INHALANTS	2.9	2.3	3.4	0.27
MARIJUANA	6.2	5.1	7.3	0.57
COCAINE OR CRACK	0.4	0.2	0.5	0.09
MARIJUANA ONLY	2.7	2.1	3.2	0.27
COCAINE	0.3	0.1	0.4	0.07
CRACK	0.1	0	0.2	0.05
HALLUCINOGEN	0.6	0.4	0.8	0.09
ROHYPNOL	0.1	0.1	0.2	0.04

STEROIDS	1	0.8	1.3	0.13
ECSTASY	2	1.5	2.5	0.23
HEROIN	0.3	0.2	0.4	0.06
METHAMPHETAMINE	0.5	0.4	0.7	0.07
SYNTHETIC MARIJ	4.9	4	5.8	0.45
CODEINE	8.4	7.4	9.4	0.51
OXYCONTIN ETC	2.6	2.1	3.2	0.27
VICODIN ETC	5.6	4.5	6.7	0.56
VALIUM ETC	1	0.7	1.3	0.15
XANAX ETC	2.9	2.3	3.4	0.29
DXM, ETC.	2.7	2.2	3.2	0.27

STEROIDS	0.3	0.2	0.5	0.08
ECSTASY	0.5	0.3	0.6	0.1
HEROIN	0.1	0	0.2	0.04
METHAMPHETAMINE	0.2	0.1	0.2	0.04
SYNTHETIC MARIJ	0.9	0.6	1.1	0.12
CODEINE	4.2	3.5	4.8	0.32
OXYCONTIN ETC	1.6	1.2	2	0.19
VICODIN ETC	2.7	2.2	3.3	0.28
VALIUM ETC	0.6	0.4	0.8	0.1
XANAX ETC	1.5	1.2	1.9	0.18
DXM, ETC.	1.2	0.9	1.5	0.14

Lifetime Use of Selected Substances - Overall Female				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	18.5	16.8	20.2	0.85
CIGARETTES	17.2	15.6	18.8	0.82
SMOKELESS TOBACCO	5.4	4.7	6.1	0.35
ALCOHOL	48.4	45.7	51.1	1.37
BEER	34.2	31.9	36.4	1.14
WINE COOLER	32.8	30.5	35.2	1.22
WINE	31.6	29.6	33.6	1.01
LIQUOR	32.5	29.9	35	1.31
ANY ILLICIT DRUG	20.4	18.6	22.2	0.93
INHALANTS	13.3	12.3	14.4	0.53
MARIJUANA	19.5	17.6	21.3	0.93
COCAINE OR CRACK	1.6	1.3	1.9	0.16
MARIJUANA ONLY	12.6	11.3	13.9	0.65
COCAINE	1.3	1	1.5	0.13
CRACK	0.5	0.4	0.7	0.08
HALLUCINOGEN	1.5	1.2	1.7	0.13
ROHYPNOL	0.8	0.6	1	0.09
STEROIDS	0.8	0.6	1	0.09
ECSTASY	2.1	1.7	2.6	0.22
HEROIN	0.4	0.2	0.5	0.06
METHAMPHETAMINE	0.6	0.4	0.8	0.11
SYNTHETIC MARIJ	5	4.4	5.6	0.3
CODEINE	9.3	8.5	10.1	0.41
OXYCONTIN ETC	2.1	1.7	2.5	0.19
VICODIN ETC	4.6	3.8	5.4	0.41
VALIUM ETC	0.9	0.7	1.1	0.09
XANAX ETC	2.9	2.4	3.4	0.25
DXM, ETC.	3.2	2.8	3.7	0.23

Past Month Use of Selected Substances - Overall Female				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	6.5	5.7	7.2	0.39
CIGARETTES	5.6	4.9	6.3	0.35
SMOKELESS TOBACCO	2.1	1.7	2.5	0.2
ALCOHOL	19.9	18.4	21.4	0.78
BEER	11.7	10.5	12.8	0.59
WINE COOLER	12.9	11.7	14.1	0.61
WINE	9	8.2	9.9	0.44
LIQUOR	12.5	11.3	13.7	0.62
ANY ILLICIT DRUG	8.1	7.2	9.1	0.5
INHALANTS	4.3	3.7	4.8	0.28
MARIJUANA	7.6	6.7	8.6	0.48
COCAINE OR CRACK	0.4	0.3	0.6	0.06
MARIJUANA ONLY	3.8	3.3	4.4	0.29
COCAINE	0.4	0.3	0.5	0.06
CRACK	0.1	0.1	0.2	0.03
HALLUCINOGEN	0.5	0.4	0.6	0.06
ROHYPNOL	0.3	0.2	0.4	0.05
STEROIDS	0.2	0.1	0.3	0.05
ECSTASY	0.6	0.3	0.8	0.15
HEROIN	0.1	0.1	0.2	0.03
METHAMPHETAMINE	0.2	0.1	0.3	0.05
SYNTHETIC MARIJ	1.3	1.1	1.6	0.14
CODEINE	4.4	4	4.9	0.25
OXYCONTIN ETC	1.1	0.9	1.4	0.13
VICODIN ETC	2.2	1.8	2.6	0.2
VALIUM ETC	0.5	0.4	0.7	0.07
XANAX ETC	1.5	1.2	1.8	0.14
DXM, ETC.	1.5	1.1	1.8	0.16

Lifetime Use of Selected Substances - Overall Male				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	21.3	19.4	23.2	0.96
CIGARETTES	18.1	16.3	19.8	0.88
SMOKELESS TOBACCO	10.5	9.3	11.6	0.57
ALCOHOL	44.6	42.2	47	1.21
BEER	36	33.6	38.3	1.2
WINE COOLER	24.4	22.6	26.3	0.95
WINE	26.4	24.7	28	0.83
LIQUOR	28.9	26.5	31.3	1.22
ANY ILLICIT DRUG	22.1	20.1	24.1	1.01
INHALANTS	11.3	10.4	12.2	0.45
MARIJUANA	21.2	19.2	23.2	1.02
COCAINE OR CRACK	2.5	2	2.9	0.21

Past Month Use of Selected Substances - Overall Male				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	8.3	7.4	9.3	0.48
CIGARETTES	6	5.2	6.8	0.4
SMOKELESS TOBACCO	4.4	3.8	5	0.3
ALCOHOL	18.2	16.6	19.8	0.81
BEER	12.6	11.2	14	0.73
WINE COOLER	8.8	7.8	9.7	0.5
WINE	7.3	6.5	8.1	0.39
LIQUOR	11	9.8	12.2	0.62
ANY ILLICIT DRUG	9.1	8.1	10.1	0.52
INHALANTS	4.1	3.5	4.6	0.27
MARIJUANA	8.1	7.1	9.1	0.51
COCAINE OR CRACK	0.8	0.6	1	0.11

MARIJUANA ONLY	11.9	10.6	13.1	0.62
COCAINE	2.2	1.8	2.7	0.21
CRACK	0.5	0.3	0.7	0.09
HALLUCINOGEN	3	2.4	3.5	0.28
ROHYPNOL	0.8	0.6	1	0.11
STEROIDS	1.2	0.9	1.4	0.12
ECSTASY	2.5	2.1	2.9	0.21
HEROIN	0.4	0.3	0.6	0.07
METHAMPHETAMINE	0.7	0.5	0.9	0.08
SYNTHETIC MARIJ	6.5	5.7	7.3	0.39
CODEINE	10.8	9.8	11.7	0.48
OXYCONTIN ETC	2.4	2	2.8	0.21
VICODIN ETC	4.4	3.7	5.2	0.38
VALIUM ETC	1	0.8	1.2	0.11
XANAX ETC	2.7	2.2	3.1	0.21
DXM, ETC.	3.4	2.9	3.9	0.24

MARIJUANA ONLY	3.6	3.1	4.1	0.27
COCAINE	0.7	0.5	0.9	0.1
CRACK	0.2	0.1	0.3	0.06
HALLUCINOGEN	0.9	0.7	1.1	0.12
ROHYPNOL	0.3	0.2	0.4	0.06
STEROIDS	0.4	0.3	0.6	0.07
ECSTASY	0.7	0.5	0.9	0.1
HEROIN	0.2	0.1	0.3	0.05
METHAMPHETAMINE	0.2	0.1	0.3	0.04
SYNTHETIC MARIJ	1.7	1.4	2	0.16
CODEINE	5.1	4.6	5.6	0.26
OXYCONTIN ETC	1.2	1	1.5	0.13
VICODIN ETC	2	1.7	2.4	0.19
VALIUM ETC	0.5	0.4	0.6	0.07
XANAX ETC	1.3	1	1.5	0.13
DXM, ETC.	1.5	1.2	1.7	0.13

Lifetime Use of Selected Substances - Overall Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	20.4	18	22.9	1.22
CIGARETTES	19.3	16.9	21.7	1.19
SMOKELESS TOBACCO	6.1	5.2	7	0.47
ALCOHOL	44.9	41.3	48.4	1.77
BEER	34.7	31.2	38.2	1.76
WINE COOLER	26.5	23.2	29.8	1.63
WINE	25.4	23	27.8	1.2
LIQUOR	27.9	24.6	31.1	1.61
ANY ILLICIT DRUG	23.1	20	26.3	1.59
INHALANTS	14.3	12.9	15.7	0.7
MARIJUANA	22	18.9	25.2	1.56
COCAINE OR CRACK	3.1	2.4	3.7	0.31
MARIJUANA ONLY	11.6	9.8	13.4	0.9
COCAINE	2.8	2.2	3.3	0.28
CRACK	0.6	0.4	0.8	0.11
HALLUCINOGEN	2	1.5	2.5	0.23
ROHYPNOL	1.4	1	1.8	0.21
STEROIDS	0.9	0.6	1.1	0.12
ECSTASY	3.9	2.7	5.1	0.59
HEROIN	0.5	0.3	0.6	0.09
METHAMPHETAMINE	0.6	0.3	1	0.16
SYNTHETIC MARIJ	7.6	6.4	8.9	0.62
CODEINE	8.8	7.4	10.2	0.71
OXYCONTIN ETC	1.7	1.2	2.1	0.21
VICODIN ETC	3.1	2	4.1	0.51
VALIUM ETC	1	0.6	1.3	0.17
XANAX ETC	3	2.2	3.8	0.41
DXM, ETC.	3	2.2	3.8	0.41

Past Month Use of Selected Substances - Overall Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	7.6	6.5	8.8	0.59
CIGARETTES	6.8	5.8	7.9	0.53
SMOKELESS TOBACCO	2.3	1.8	2.8	0.27
ALCOHOL	19.7	17.5	22	1.13
BEER	13.1	11.3	14.9	0.88
WINE COOLER	10.9	9.2	12.7	0.89
WINE	8.3	7.4	9.3	0.47
LIQUOR	10.9	9.4	12.4	0.76
ANY ILLICIT DRUG	10.1	8.3	11.8	0.86
INHALANTS	5.7	5.1	6.3	0.31
MARIJUANA	8.9	7.4	10.4	0.76
COCAINE OR CRACK	1.1	0.8	1.4	0.16
MARIJUANA ONLY	3.5	2.9	4.1	0.3
COCAINE	0.9	0.7	1.2	0.15
CRACK	0.3	0.2	0.4	0.07
HALLUCINOGEN	0.7	0.5	1	0.11
ROHYPNOL	0.5	0.3	0.7	0.11
STEROIDS	0.4	0.3	0.6	0.09
ECSTASY	1.3	0.8	1.9	0.27
HEROIN	0.1	0	0.2	0.05
METHAMPHETAMINE	0.2	0.1	0.4	0.08
SYNTHETIC MARIJ	2.6	2	3.1	0.28
CODEINE	3.7	3	4.3	0.33
OXYCONTIN ETC	0.7	0.5	0.9	0.11
VICODIN ETC	1.4	0.9	2	0.27
VALIUM ETC	0.5	0.3	0.7	0.1
XANAX ETC	1.5	1	2	0.25
DXM, ETC.	1.4	0.9	1.8	0.23

Lifetime Use of Selected Substances - Overall Non-Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	19.8	17.9	21.8	0.97
CIGARETTES	17.4	15.6	19.2	0.9
SMOKELESS TOBACCO	8.3	7.4	9.2	0.46
ALCOHOL	46.8	44	49.6	1.43
BEER	35.1	32.6	37.6	1.27
WINE COOLER	29	26.6	31.4	1.21
WINE	29.5	27.6	31.5	0.99

Past Month Use of Selected Substances - Overall Non-Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	7.4	6.5	8.3	0.45
CIGARETTES	5.6	4.9	6.4	0.38
SMOKELESS TOBACCO	3.4	3	3.9	0.24
ALCOHOL	19	17.3	20.7	0.86
BEER	12	10.6	13.4	0.7
WINE COOLER	10.8	9.7	12	0.59
WINE	8.1	7.3	8.9	0.41

LIQUOR	31.2	28.4	34	1.42
ANY ILLICIT DRUG	20.9	18.9	23	1.05
INHALANTS	11.9	11	12.9	0.46
MARIJUANA	20.1	18	22.2	1.06
COCAINE OR CRACK	1.9	1.6	2.2	0.15
MARIJUANA ONLY	12.3	11	13.7	0.69
COCAINE	1.6	1.3	1.9	0.14
CRACK	0.5	0.4	0.6	0.07
HALLUCINOGEN	2.3	1.9	2.7	0.2
ROHYPNOL	0.7	0.5	0.8	0.08
STEROIDS	1	0.8	1.2	0.08
ECSTASY	2.1	1.7	2.4	0.19
HEROIN	0.4	0.3	0.5	0.05
METHAMPHETAMINE	0.6	0.5	0.8	0.08
SYNTHETIC MARIJ	5.4	4.8	6.1	0.34
CODEINE	10.2	9.4	11.1	0.44
OXYCONTIN ETC	2.3	1.9	2.7	0.19
VICODIN ETC	4.8	3.9	5.6	0.42
VALIUM ETC	0.9	0.8	1.1	0.09
XANAX ETC	2.7	2.3	3.2	0.23
DXM, ETC.	3.3	2.9	3.8	0.22

LIQUOR	11.9	10.6	13.2	0.67
ANY ILLICIT DRUG	8.4	7.4	9.4	0.51
INHALANTS	3.9	3.4	4.4	0.25
MARIJUANA	7.7	6.7	8.7	0.5
COCAINE OR CRACK	0.6	0.4	0.7	0.07
MARIJUANA ONLY	3.8	3.3	4.3	0.25
COCAINE	0.5	0.3	0.6	0.06
CRACK	0.2	0.1	0.2	0.03
HALLUCINOGEN	0.7	0.5	0.8	0.08
ROHYPNOL	0.3	0.2	0.4	0.05
STEROIDS	0.3	0.2	0.4	0.05
ECSTASY	0.5	0.3	0.7	0.1
HEROIN	0.2	0.1	0.2	0.03
METHAMPHETAMINE	0.2	0.1	0.3	0.03
SYNTHETIC MARIJ	1.4	1.1	1.6	0.13
CODEINE	4.9	4.5	5.4	0.24
OXYCONTIN ETC	1.2	1	1.5	0.11
VICODIN ETC	2.3	1.9	2.7	0.2
VALIUM ETC	0.5	0.4	0.6	0.06
XANAX ETC	1.4	1.2	1.6	0.12
DXM, ETC.	1.5	1.2	1.7	0.13

Lifetime Use of Selected Substances - Grade 6				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	5.8	4.3	7.4	0.76
CIGARETTES	4.7	3.2	6.3	0.76
SMOKELESS TOBACCO	1.8	1.2	2.5	0.31
ALCOHOL	23.1	19.9	26.3	1.6
BEER	15.4	12.6	18.2	1.41
WINE COOLER	9.7	8.1	11.3	0.8
WINE	13.5	11.2	15.7	1.11
LIQUOR	7.2	5.9	8.6	0.66
ANY ILLICIT DRUG	4.5	2.7	6.3	0.91
INHALANTS	12	10.1	13.8	0.93
MARIJUANA	3.6	1.6	5.5	0.96
COCAINE OR CRACK	1	0.2	1.8	0.4
MARIJUANA ONLY	1.9	0.9	2.9	0.49
COCAINE	0.6	0	1.3	0.35
CRACK	0.6	0.1	1	0.24
HALLUCINOGEN	0.2	0	0.3	0.08
ROHYPNOL	0.3	0	0.5	0.1
STEROIDS	1	0.7	1.3	0.15
ECSTASY	0.1	0	0.2	0.05
HEROIN	0.5	0.2	0.7	0.11
METHAMPHETAMINE	0.4	0.1	0.7	0.13
SYNTHETIC MARIJ	1.3	0.3	2.3	0.51
CODEINE	5.6	5	6.3	0.32
OXYCONTIN ETC	0.6	0.4	0.8	0.11
VICODIN ETC	0.6	0.4	0.8	0.11
VALIUM ETC	0.1	0	0.2	0.05
XANAX ETC	0.4	0.2	0.5	0.09
DXM, ETC.	1.9	1.3	2.6	0.33

Past Month Use of Selected Substances - Grade 6				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	1.6	1.1	2.2	0.26
CIGARETTES	1.3	0.8	1.7	0.23
SMOKELESS TOBACCO	0.6	0.1	1	0.22
ALCOHOL	6.7	5.1	8.3	0.8
BEER	2.9	1.9	3.9	0.48
WINE COOLER	3.3	2.5	4.2	0.44
WINE	3	2.2	3.7	0.38
LIQUOR	1.7	1.1	2.3	0.32
ANY ILLICIT DRUG	1.4	0.7	2	0.3
INHALANTS	5.6	4.4	6.8	0.61
MARIJUANA	0.9	0.3	1.5	0.3
COCAINE OR CRACK	0.3	0	0.5	0.13
MARIJUANA ONLY	0.4	0	0.9	0.21
COCAINE	0	0	0.1	0.02
CRACK	0.2	0	0.5	0.13
HALLUCINOGEN	0.1	0	0.1	0.04
ROHYPNOL	0.2	0	0.3	0.08
STEROIDS	0.1	0	0.3	0.06
ECSTASY	0	0	0.1	0.02
HEROIN	0.2	0	0.4	0.08
METHAMPHETAMINE	0.1	0	0.2	0.05
SYNTHETIC MARIJ	0.3	0	0.6	0.15
CODEINE	2.8	2.3	3.2	0.23
OXYCONTIN ETC	0.2	0	0.3	0.07
VICODIN ETC	0.2	0	0.3	0.08
VALIUM ETC	0.1	0	0.1	0.04
XANAX ETC	0.1	0	0.2	0.05
DXM, ETC.	0.6	0.4	0.8	0.08

Lifetime Use of Selected Substances - Grade 6 African American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	8.1	4.1	12	1.95
CIGARETTES	7	1.8	12.2	2.56

Past Month Use of Selected Substances - Grade 6 African American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	1.4	0	2.9	0.76
CIGARETTES	1.3	0	2.9	0.76

SMOKELESS TOBACCO	1.2	0	2.9	0.8
ALCOHOL	23.8	18.9	28.7	2.42
BEER	13.3	7.7	18.9	2.74
WINE COOLER	9.7	6.2	13.1	1.68
WINE	15.6	12.6	18.6	1.47
LIQUOR	8.3	6.6	10	0.82
ANY ILLICIT DRUG	5.9	0.7	11.1	2.56
INHALANTS	11.7	8.9	14.5	1.37
MARIJUANA	5.4	0.1	10.7	2.61
COCAINE OR CRACK	0.7	0	1.7	0.49
MARIJUANA ONLY	3.8	0.3	7.3	1.72
COCAINE	0.6	0	1.5	0.46
CRACK	0.1	0	0.4	0.12
HALLUCINOGEN	0	*	*	*
ROHYPNOL	0	*	*	*
STEROIDS	0.1	0	0.4	0.12
ECSTASY	0	*	*	*
HEROIN	0	*	*	*
METHAMPHETAMINE	1.6	0	3.7	1.02
SYNTHETIC MARIJ	1.3	0	3.3	0.96
CODEINE	7.5	4.5	10.5	1.49
OXYCONTIN ETC	0.3	0	0.7	0.22
VICODIN ETC	0.2	0	0.6	0.2
VALIUM ETC	0.1	0	0.3	0.12
XANAX ETC	0.1	0	0.4	0.13
DXM, ETC.	2	0.3	3.6	0.82

SMOKELESS TOBACCO	0	0	0.1	0.04
ALCOHOL	8.5	5.8	11.3	1.37
BEER	3	0.9	5.1	1.03
WINE COOLER	3.4	0.5	6.3	1.43
WINE	4.4	0.8	8.1	1.79
LIQUOR	3.6	2.4	4.7	0.57
ANY ILLICIT DRUG	1.7	0	3.7	0.97
INHALANTS	5.9	0.7	11.1	2.57
MARIJUANA	1.6	0	3.6	0.98
COCAINE OR CRACK	0.1	0	0.3	0.11
MARIJUANA ONLY	1.5	0	3.5	0.96
COCAINE	0	*	*	*
CRACK	0.1	0	0.3	0.11
HALLUCINOGEN	0	*	*	*
ROHYPNOL	0	*	*	*
STEROIDS	0.1	0	0.3	0.11
ECSTASY	0	*	*	*
HEROIN	0	*	*	*
METHAMPHETAMINE	0	*	*	*
SYNTHETIC MARIJ	0	0	0.1	0.02
CODEINE	5	2.7	7.2	1.09
OXYCONTIN ETC	0.1	0	0.2	0.08
VICODIN ETC	0	*	*	*
VALIUM ETC	0.1	0	0.3	0.12
XANAX ETC	0	0	0.1	0.04
DXM, ETC.	0.9	0	2	0.55

Lifetime Use of Selected Substances - Grade 6 Mexican American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	7.4	6.1	8.6	0.61
CIGARETTES	5.9	4.7	7.2	0.62
SMOKELESS TOBACCO	2.5	1.5	3.5	0.5
ALCOHOL	26.9	23.8	29.9	1.51
BEER	18	15.1	20.8	1.42
WINE COOLER	12.5	9.9	15.1	1.28
WINE	15.2	12.2	18.3	1.51
LIQUOR	9.1	6.7	11.6	1.22
ANY ILLICIT DRUG	5.9	4.2	7.5	0.8
INHALANTS	14.4	12.5	16.2	0.91
MARIJUANA	4.9	2.9	6.9	0.98
COCAINE OR CRACK	1.4	0.2	2.6	0.61
MARIJUANA ONLY	2.2	1.4	3.1	0.43
COCAINE	0.7	0	1.7	0.48
CRACK	1	0	2	0.49
HALLUCINOGEN	0.1	0	0.3	0.07
ROHYPNOL	0.4	0.1	0.7	0.17
STEROIDS	1.2	0.6	1.8	0.31
ECSTASY	0.2	0	0.3	0.08
HEROIN	0.6	0	1.3	0.31
METHAMPHETAMINE	0.2	0	0.3	0.08
SYNTHETIC MARIJ	1.8	0.4	3.3	0.71
CODEINE	4.6	3.5	5.8	0.57
OXYCONTIN ETC	0.9	0.3	1.4	0.28
VICODIN ETC	0.7	0.3	1	0.17
VALIUM ETC	0.1	0	0.3	0.08
XANAX ETC	0.5	0.2	0.9	0.16
DXM, ETC.	2.1	1.4	2.7	0.33

Past Month Use of Selected Substances - Grade 6 Mexican American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	2.7	1.9	3.6	0.43
CIGARETTES	2	1.2	2.8	0.41
SMOKELESS TOBACCO	1.1	0.2	1.9	0.42
ALCOHOL	8.6	6.8	10.4	0.89
BEER	4.3	3.2	5.4	0.56
WINE COOLER	4.5	3.2	5.8	0.66
WINE	3.5	2.5	4.5	0.48
LIQUOR	2.1	1.2	3	0.45
ANY ILLICIT DRUG	2.1	1.3	2.8	0.37
INHALANTS	6.9	5.4	8.3	0.71
MARIJUANA	1.4	0.8	2	0.29
COCAINE OR CRACK	0.5	0	1.1	0.31
MARIJUANA ONLY	0.5	0.2	0.8	0.16
COCAINE	0	0	0.1	0.02
CRACK	0.4	0	1.1	0.31
HALLUCINOGEN	0.1	0	0.2	0.07
ROHYPNOL	0.3	0	0.6	0.16
STEROIDS	0.2	0	0.3	0.07
ECSTASY	0	0	0.1	0.02
HEROIN	0.3	0	0.6	0.16
METHAMPHETAMINE	0.1	0	0.1	0.04
SYNTHETIC MARIJ	0.6	0	1.2	0.3
CODEINE	1.7	1.1	2.4	0.32
OXYCONTIN ETC	0.1	0	0.2	0.07
VICODIN ETC	0.2	0	0.3	0.09
VALIUM ETC	0.1	0	0.3	0.07
XANAX ETC	0.2	0	0.5	0.11
DXM, ETC.	0.6	0.3	0.9	0.16

Lifetime Use of Selected Substances - Grade 6 White				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	3.6	1.5	5.6	1.02
CIGARETTES	2.8	1.1	4.5	0.85
SMOKELESS TOBACCO	1.3	0.5	2.2	0.43
ALCOHOL	18.6	13.9	23.2	2.29
BEER	13.6	9.2	18.1	2.21
WINE COOLER	6.5	4.1	8.8	1.17
WINE	10.2	7.3	13.1	1.44
LIQUOR	5.4	3.3	7.4	1
ANY ILLICIT DRUG	2.6	1	4.1	0.77
INHALANTS	9.7	6.4	12.9	1.61
MARIJUANA	1.5	0.3	2.7	0.58
COCAINE OR CRACK	0.6	0	1.5	0.46
MARIJUANA ONLY	1	0	2	0.52
COCAINE	0.5	0	1.4	0.46
CRACK	0.1	0	0.2	0.07
HALLUCINOGEN	0.3	0	0.6	0.17
ROHYPNOL	0.2	0	0.5	0.15
STEROIDS	1	0.5	1.5	0.25
ECSTASY	0.1	0	0.2	0.07
HEROIN	0.4	0.1	0.7	0.14
METHAMPHETAMINE	0.4	0.1	0.7	0.16
SYNTHETIC MARIJ	0.4	0	0.9	0.23
CODEINE	6.8	5.6	8.1	0.64
OXYCONTIN ETC	0.6	0.1	1.1	0.26
VICODIN ETC	0.8	0.3	1.3	0.27
VALIUM ETC	0.1	0	0.3	0.07
XANAX ETC	0.3	0	0.6	0.15
DXM, ETC.	2	0.4	3.6	0.79

Past Month Use of Selected Substances - Grade 6 White				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	0.5	0	1	0.25
CIGARETTES	0.3	0	0.7	0.19
SMOKELESS TOBACCO	0.2	0	0.4	0.11
ALCOHOL	3.8	2.7	4.8	0.52
BEER	1.2	0.5	1.9	0.35
WINE COOLER	2	1.1	2.8	0.42
WINE	1.8	1.2	2.4	0.28
LIQUOR	0.7	0.1	1.3	0.3
ANY ILLICIT DRUG	0.5	0.2	0.9	0.17
INHALANTS	4.2	2.6	5.8	0.8
MARIJUANA	0.2	0	0.4	0.1
COCAINE OR CRACK	0.1	0	0.2	0.07
MARIJUANA ONLY	0	0	0.1	0.04
COCAINE	0	0	0.1	0.04
CRACK	0.1	0	0.2	0.07
HALLUCINOGEN	0.1	0	0.2	0.06
ROHYPNOL	0	0	0.1	0.03
STEROIDS	0.1	0	0.3	0.1
ECSTASY	0	0	0.1	0.02
HEROIN	0.2	0	0.5	0.14
METHAMPHETAMINE	0.2	0	0.5	0.13
SYNTHETIC MARIJ	0.1	0	0.2	0.06
CODEINE	3.6	2.4	4.8	0.6
OXYCONTIN ETC	0.4	0	0.8	0.21
VICODIN ETC	0.4	0	0.8	0.2
VALIUM ETC	0	*	*	*
XANAX ETC	0.1	0	0.2	0.05
DXM, ETC.	0.6	0.2	1	0.19

Lifetime Use of Selected Substances - Grade 6 Female				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	4.4	2.9	5.9	0.74
CIGARETTES	3.9	2.5	5.3	0.71
SMOKELESS TOBACCO	1.1	0.5	1.8	0.31
ALCOHOL	20.8	16.9	24.6	1.92
BEER	12.6	9.8	15.4	1.38
WINE COOLER	9.4	7.7	11.1	0.84
WINE	12.3	10.1	14.5	1.09
LIQUOR	6.2	4.6	7.8	0.81
ANY ILLICIT DRUG	3.4	2.2	4.5	0.59
INHALANTS	11.4	9.5	13.3	0.97
MARIJUANA	2.6	1.1	4	0.72
COCAINE OR CRACK	0.5	0.1	0.9	0.22
MARIJUANA ONLY	1.4	0.5	2.3	0.47
COCAINE	0.2	0	0.6	0.17
CRACK	0.3	0	0.6	0.16
HALLUCINOGEN	0	0	0.1	0.02
ROHYPNOL	0.1	0	0.2	0.07
STEROIDS	0.8	0.3	1.4	0.27
ECSTASY	0.1	0	0.3	0.07
HEROIN	0.5	0	1	0.27
METHAMPHETAMINE	0.2	0	0.4	0.1
SYNTHETIC MARIJ	0.8	0.2	1.5	0.32
CODEINE	6.4	5.3	7.5	0.55
OXYCONTIN ETC	0.7	0.2	1.3	0.27

Past Month Use of Selected Substances - Grade 6 Female				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	1.4	0.7	2.2	0.36
CIGARETTES	1.2	0.6	1.9	0.33
SMOKELESS TOBACCO	0.5	0.1	0.8	0.17
ALCOHOL	6.1	4	8.2	1.05
BEER	2.5	1.3	3.6	0.56
WINE COOLER	3.7	2.3	5.2	0.72
WINE	2.5	1.4	3.6	0.55
LIQUOR	1.1	0.5	1.8	0.34
ANY ILLICIT DRUG	1.3	0.2	2.4	0.54
INHALANTS	5.2	4.1	6.2	0.53
MARIJUANA	1.1	0	2.2	0.56
COCAINE OR CRACK	0.1	0	0.1	0.04
MARIJUANA ONLY	0.6	0	1.4	0.4
COCAINE	0	0	0.1	0.03
CRACK	0	0	0.1	0.03
HALLUCINOGEN	0	0	0	0.01
ROHYPNOL	0	0	0.1	0.02
STEROIDS	0.2	0	0.4	0.09
ECSTASY	0	0	0	0.01
HEROIN	0.1	0	0.3	0.1
METHAMPHETAMINE	0.1	0	0.3	0.1
SYNTHETIC MARIJ	0.3	0	0.6	0.17
CODEINE	2.9	2	3.8	0.44
OXYCONTIN ETC	0.2	0	0.5	0.12

VICODIN ETC	0.7	0.2	1.2	0.23
VALIUM ETC	0.1	0	0.3	0.07
XANAX ETC	0.4	0.1	0.7	0.13
DXM, ETC.	2.1	0.9	3.3	0.59

VICODIN ETC	0.2	0	0.5	0.13
VALIUM ETC	0.1	0	0.2	0.07
XANAX ETC	0.2	0	0.5	0.11
DXM, ETC.	0.5	0.2	0.7	0.14

Lifetime Use of Selected Substances - Grade 6 Male				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	7.2	5.1	9.2	1.04
CIGARETTES	5.5	3.6	7.4	0.97
SMOKELESS TOBACCO	2.5	1.6	3.5	0.49
ALCOHOL	25.3	22.3	28.4	1.53
BEER	17.9	14.7	21.2	1.61
WINE COOLER	10.1	8.1	12.1	0.99
WINE	14.5	11.8	17.2	1.34
LIQUOR	8.2	6.1	10.3	1.05
ANY ILLICIT DRUG	5.5	2.9	8.1	1.29
INHALANTS	12.5	9.9	15.2	1.32
MARIJUANA	4.5	2	6.9	1.23
COCAINE OR CRACK	1.4	0.1	2.8	0.68
MARIJUANA ONLY	2.3	1.2	3.5	0.57
COCAINE	1	0	2.4	0.68
CRACK	0.8	0	1.5	0.36
HALLUCINOGEN	0.3	0	0.6	0.16
ROHYPNOL	0.4	0	0.8	0.19
STEROIDS	1.1	0.8	1.5	0.17
ECSTASY	0.1	0	0.2	0.05
HEROIN	0.4	0.1	0.8	0.18
METHAMPHETAMINE	0.6	0.1	1.1	0.26
SYNTHETIC MARIJ	1.7	0.3	3.1	0.7
CODEINE	5	3.9	6.1	0.54
OXYCONTIN ETC	0.5	0.2	0.9	0.16
VICODIN ETC	0.5	0.2	0.8	0.17
VALIUM ETC	0.1	0	0.2	0.05
XANAX ETC	0.3	0	0.6	0.15
DXM, ETC.	1.8	1.2	2.5	0.32

Past Month Use of Selected Substances - Grade 6 Male				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	1.8	1.2	2.4	0.3
CIGARETTES	1.2	0.6	1.9	0.32
SMOKELESS TOBACCO	0.6	0	1.2	0.29
ALCOHOL	7.3	5.8	8.8	0.74
BEER	3.2	2.2	4.2	0.49
WINE COOLER	3	1.9	4.1	0.53
WINE	3.5	2.6	4.4	0.46
LIQUOR	2.2	1.4	2.9	0.36
ANY ILLICIT DRUG	1.3	0.8	1.8	0.25
INHALANTS	6	4	8	1
MARIJUANA	0.7	0.2	1.2	0.23
COCAINE OR CRACK	0.4	0	0.9	0.25
MARIJUANA ONLY	0.3	0	0.6	0.13
COCAINE	0	0	0	0.01
CRACK	0.4	0	0.9	0.25
HALLUCINOGEN	0.1	0	0.3	0.07
ROHYPNOL	0.3	0	0.6	0.15
STEROIDS	0.1	0	0.2	0.05
ECSTASY	0	0	0.1	0.02
HEROIN	0.3	0	0.6	0.15
METHAMPHETAMINE	0	0	0.1	0.04
SYNTHETIC MARIJ	0.4	0.1	0.7	0.15
CODEINE	2.6	1.7	3.5	0.46
OXYCONTIN ETC	0.1	0	0.3	0.08
VICODIN ETC	0.2	0	0.3	0.09
VALIUM ETC	0	0	0.1	0.03
XANAX ETC	0	0	0.1	0.03
DXM, ETC.	0.8	0.4	1.1	0.16

Lifetime Use of Selected Substances - Grade 6 Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	6.1	3.2	9	1.22
CIGARETTES	5.2	2.2	8.2	1.26
SMOKELESS TOBACCO	1.6	0.6	2.6	0.42
ALCOHOL	22.2	19.8	24.6	1.02
BEER	13	9	17.1	1.71
WINE COOLER	8.7	7.7	9.7	0.43
WINE	10.2	8.5	11.9	0.73
LIQUOR	5.3	4.1	6.5	0.51
ANY ILLICIT DRUG	4.6	2.8	6.4	0.77
INHALANTS	15	12.2	17.8	1.18
MARIJUANA	4	2.4	5.6	0.69
COCAINE OR CRACK	0.3	0	0.7	0.21
MARIJUANA ONLY	1.8	0.7	2.9	0.47
COCAINE	0.1	0	0.4	0.13
CRACK	0.1	0	0.4	0.1
HALLUCINOGEN	0.2	0	0.6	0.15
ROHYPNOL	0	0	0.2	0.05
STEROIDS	1	0.4	1.5	0.23
ECSTASY	0.2	0	0.6	0.16

Past Month Use of Selected Substances - Grade 6 Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	2	0.2	3.8	0.76
CIGARETTES	1.6	0	3.3	0.73
SMOKELESS TOBACCO	0.6	0.1	1.1	0.2
ALCOHOL	8.2	5.2	11.2	1.28
BEER	4	1.3	6.6	1.13
WINE COOLER	3.3	2.2	4.4	0.47
WINE	3.3	1.7	5	0.69
LIQUOR	0.8	0.1	1.5	0.29
ANY ILLICIT DRUG	1.6	0.5	2.7	0.46
INHALANTS	7.6	6.1	9.1	0.64
MARIJUANA	1.3	0.5	2	0.3
COCAINE OR CRACK	0.2	0	0.5	0.15
MARIJUANA ONLY	0.4	0.1	0.8	0.15
COCAINE	0.1	0	0.2	0.06
CRACK	0.1	0	0.4	0.1
HALLUCINOGEN	0.1	0	0.4	0.1
ROHYPNOL	0	0	0.2	0.05
STEROIDS	0.4	0	0.9	0.19
ECSTASY	0	0	0.2	0.05

HEROIN	0.2	0	0.4	0.11
METHAMPHETAMINE	0.5	0.1	0.9	0.16
SYNTHETIC MARIJ	1.5	0	3.2	0.7
CODEINE	3.1	1.7	4.5	0.59
OXYCONTIN ETC	0.6	0.2	1	0.17
VICODIN ETC	0.9	0	1.8	0.37
VALIUM ETC	0.2	0	0.6	0.14
XANAX ETC	0.7	0.5	1	0.11
DXM, ETC.	2.1	1.1	3.2	0.45

HEROIN	0.2	0	0.4	0.1
METHAMPHETAMINE	0.2	0	0.5	0.14
SYNTHETIC MARIJ	0.4	0	0.9	0.2
CODEINE	1.5	0.3	2.7	0.51
OXYCONTIN ETC	0.3	0	0.7	0.15
VICODIN ETC	0.1	0	0.3	0.09
VALIUM ETC	0.1	0	0.3	0.09
XANAX ETC	0.2	0	0.4	0.11
DXM, ETC.	1	0.4	1.6	0.25

Lifetime Use of Selected Substances - Grade 6 Non-Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	5.8	4	7.5	0.87
CIGARETTES	4.7	2.9	6.4	0.88
SMOKELESS TOBACCO	1.9	1.2	2.6	0.35
ALCOHOL	23.3	19.5	27.1	1.91
BEER	15.8	12.5	19.2	1.66
WINE COOLER	9.9	8.1	11.8	0.94
WINE	14.1	11.5	16.6	1.28
LIQUOR	7.6	6.1	9.2	0.77
ANY ILLICIT DRUG	4.4	2.3	6.6	1.07
INHALANTS	11.4	9.3	13.5	1.06
MARIJUANA	3.5	1.2	5.8	1.14
COCAINE OR CRACK	1.1	0.2	2.1	0.46
MARIJUANA ONLY	1.9	0.7	3	0.58
COCAINE	0.7	0	1.5	0.41
CRACK	0.6	0.1	1.2	0.27
HALLUCINOGEN	0.1	0	0.3	0.1
ROHYPNOL	0.3	0	0.5	0.12
STEROIDS	1	0.6	1.3	0.17
ECSTASY	0.1	0	0.2	0.05
HEROIN	0.5	0.3	0.8	0.12
METHAMPHETAMINE	0.4	0.1	0.7	0.16
SYNTHETIC MARIJ	1.2	0	2.4	0.59
CODEINE	6.1	5.5	6.8	0.33
OXYCONTIN ETC	0.6	0.4	0.9	0.12
VICODIN ETC	0.5	0.3	0.7	0.1
VALIUM ETC	0.1	0	0.2	0.05
XANAX ETC	0.3	0.1	0.5	0.1
DXM, ETC.	1.9	1.1	2.7	0.38

Past Month Use of Selected Substances - Grade 6 Non-Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	1.6	1	2.1	0.27
CIGARETTES	1.2	0.7	1.7	0.23
SMOKELESS TOBACCO	0.5	0	1.1	0.26
ALCOHOL	6.4	4.6	8.3	0.91
BEER	2.7	1.6	3.7	0.53
WINE COOLER	3.3	2.3	4.4	0.51
WINE	2.9	2	3.8	0.44
LIQUOR	1.9	1.1	2.6	0.38
ANY ILLICIT DRUG	1.3	0.6	2	0.35
INHALANTS	5.2	3.8	6.7	0.71
MARIJUANA	0.9	0.2	1.6	0.36
COCAINE OR CRACK	0.3	0	0.6	0.15
MARIJUANA ONLY	0.4	0	1	0.25
COCAINE	0	0	0	0.01
CRACK	0.3	0	0.6	0.15
HALLUCINOGEN	0.1	0	0.1	0.04
ROHYPNOL	0.2	0	0.4	0.09
STEROIDS	0.1	0	0.2	0.06
ECSTASY	0	0	0	0.02
HEROIN	0.2	0	0.4	0.09
METHAMPHETAMINE	0.1	0	0.2	0.06
SYNTHETIC MARIJ	0.3	0	0.7	0.17
CODEINE	3	2.5	3.5	0.24
OXYCONTIN ETC	0.2	0	0.3	0.08
VICODIN ETC	0.2	0	0.4	0.09
VALIUM ETC	0.1	0	0.1	0.04
XANAX ETC	0.1	0	0.3	0.06
DXM, ETC.	0.5	0.4	0.7	0.08

Lifetime Use of Selected Substances - Grade 7				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	10	7.7	12.3	1.14
CIGARETTES	8.8	6.6	11	1.12
SMOKELESS TOBACCO	3.2	2.4	3.9	0.38
ALCOHOL	30.9	26.5	35.2	2.18
BEER	21.7	18.1	25.4	1.81
WINE COOLER	14.4	11.6	17.2	1.4
WINE	18	14.6	21.4	1.7
LIQUOR	12	9.7	14.4	1.19
ANY ILLICIT DRUG	8	6	10	1
INHALANTS	14.2	12	16.5	1.12
MARIJUANA	6.9	4.9	9	1.01
COCAINE OR CRACK	0.8	0.5	1.2	0.19
MARIJUANA ONLY	4	2.8	5.3	0.62
COCAINE	0.5	0.2	0.9	0.16

Past Month Use of Selected Substances - Grade 7				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	2.6	1.9	3.2	0.34
CIGARETTES	1.9	1.3	2.4	0.28
SMOKELESS TOBACCO	1	0.5	1.4	0.23
ALCOHOL	9.7	7.7	11.6	0.97
BEER	5.4	4.1	6.8	0.69
WINE COOLER	4.6	3.5	5.7	0.57
WINE	4.5	3.4	5.6	0.54
LIQUOR	3.5	2.5	4.5	0.5
ANY ILLICIT DRUG	3.4	2.4	4.4	0.52
INHALANTS	5.6	4.4	6.9	0.63
MARIJUANA	2.8	1.8	3.8	0.52
COCAINE OR CRACK	0.4	0.2	0.6	0.1
MARIJUANA ONLY	1.4	0.7	2.1	0.36
COCAINE	0.3	0.1	0.5	0.1

CRACK	0.4	0.2	0.6	0.09
HALLUCINOGEN	0.6	0.3	0.8	0.12
ROHYPNOL	0.5	0.2	0.7	0.14
STEROIDS	0.6	0.4	0.9	0.14
ECSTASY	0.6	0.2	1	0.2
HEROIN	0.7	0.5	0.9	0.11
METHAMPHETAMINE	0.5	0.3	0.7	0.09
SYNTHETIC MARIJ	2.2	1.5	3	0.37
CODEINE	5.6	4.6	6.6	0.5
OXYCONTIN ETC	0.5	0.3	0.8	0.11
VICODIN ETC	0.9	0.6	1.2	0.17
VALIUM ETC	0.3	0.1	0.4	0.07
XANAX ETC	0.5	0.3	0.7	0.1
DXM, ETC.	3.2	2.3	4.1	0.44

CRACK	0.1	0	0.2	0.05
HALLUCINOGEN	0.2	0	0.3	0.06
ROHYPNOL	0.1	0	0.2	0.05
STEROIDS	0.2	0	0.4	0.08
ECSTASY	0.1	0	0.2	0.04
HEROIN	0.4	0.2	0.6	0.08
METHAMPHETAMINE	0.2	0.1	0.4	0.07
SYNTHETIC MARIJ	0.8	0.4	1.3	0.21
CODEINE	2.7	2	3.4	0.34
OXYCONTIN ETC	0.3	0.1	0.5	0.09
VICODIN ETC	0.4	0.2	0.6	0.1
VALIUM ETC	0.1	0	0.2	0.05
XANAX ETC	0.3	0.1	0.4	0.07
DXM, ETC.	1.3	0.6	2	0.34

Lifetime Use of Selected Substances - Grade 7 African American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	10.2	6.1	14.2	2.01
CIGARETTES	8.4	5.9	10.9	1.23
SMOKELESS TOBACCO	3.6	0	8	2.15
ALCOHOL	38.2	30.8	45.6	3.67
BEER	17.9	12	23.8	2.92
WINE COOLER	22.3	15.7	28.9	3.27
WINE	25	17.3	32.7	3.81
LIQUOR	15.2	10.2	20.3	2.49
ANY ILLICIT DRUG	11.1	7.5	14.7	1.78
INHALANTS	13.4	8.6	18.2	2.37
MARIJUANA	9	5	13.1	2.01
COCAINE OR CRACK	0.3	0	0.8	0.26
MARIJUANA ONLY	6.3	2.7	9.9	1.78
COCAINE	0.2	0	0.7	0.22
CRACK	0.1	0	0.4	0.12
HALLUCINOGEN	0.4	0	1.1	0.33
ROHYPNOL	0	*	*	*
STEROIDS	0.7	0	1.6	0.43
ECSTASY	0.6	0	1.6	0.47
HEROIN	1	0	2.2	0.63
METHAMPHETAMINE	1	0	2.2	0.63
SYNTHETIC MARIJ	2.7	0	5.4	1.34
CODEINE	4.9	2.1	7.8	1.42
OXYCONTIN ETC	0.6	0	1.4	0.38
VICODIN ETC	0.2	0	0.5	0.18
VALIUM ETC	0.2	0	0.5	0.18
XANAX ETC	0.2	0	0.5	0.18
DXM, ETC.	1.8	0.3	3.3	0.74

Past Month Use of Selected Substances - Grade 7 African American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	1.5	0	3.1	0.79
CIGARETTES	0.8	0	1.8	0.51
SMOKELESS TOBACCO	0.8	0	1.9	0.57
ALCOHOL	13	8.1	18	2.45
BEER	4.1	0.6	7.5	1.7
WINE COOLER	8.4	5.8	11.1	1.3
WINE	7.8	3.4	12.3	2.2
LIQUOR	4.2	0.9	7.6	1.65
ANY ILLICIT DRUG	5.5	3.5	7.4	0.96
INHALANTS	5.5	1.1	9.8	2.14
MARIJUANA	4.3	2.2	6.5	1.06
COCAINE OR CRACK	0.2	0	0.7	0.22
MARIJUANA ONLY	2.6	0.2	4.9	1.16
COCAINE	0.2	0	0.7	0.22
CRACK	0	*	*	*
HALLUCINOGEN	0	*	*	*
ROHYPNOL	0	*	*	*
STEROIDS	0.5	0	1.3	0.41
ECSTASY	0	*	*	*
HEROIN	1	0	2.2	0.63
METHAMPHETAMINE	0.6	0	1.7	0.54
SYNTHETIC MARIJ	1.5	0	3.4	0.93
CODEINE	3.6	1.2	6	1.19
OXYCONTIN ETC	0.5	0	1.2	0.36
VICODIN ETC	0	*	*	*
VALIUM ETC	0	*	*	*
XANAX ETC	0.2	0	0.5	0.18
DXM, ETC.	0.6	0	1.4	0.4

Lifetime Use of Selected Substances - Grade 7 Mexican American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	13.1	10.9	15.4	1.12
CIGARETTES	11.7	9.5	13.8	1.08
SMOKELESS TOBACCO	4	2.8	5.1	0.57
ALCOHOL	35.8	31.6	40	2.1
BEER	27	22.9	31.2	2.06
WINE COOLER	16.3	13.2	19.4	1.54
WINE	19.3	16.1	22.6	1.62
LIQUOR	15	12.7	17.2	1.14
ANY ILLICIT DRUG	11.2	8.7	13.7	1.24

Past Month Use of Selected Substances - Grade 7 Mexican American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	3.4	2.4	4.5	0.52
CIGARETTES	2.5	1.7	3.4	0.43
SMOKELESS TOBACCO	1.3	0.6	2	0.34
ALCOHOL	12.7	10.4	14.9	1.12
BEER	7.4	5.8	9.1	0.82
WINE COOLER	5.8	4.2	7.3	0.78
WINE	5.7	4.2	7.3	0.76
LIQUOR	4.9	3.7	6.2	0.61
ANY ILLICIT DRUG	4.5	3.1	6	0.72

INHALANTS	18.4	15.4	21.4	1.48
MARIJUANA	10.1	7.6	12.7	1.27
COCAINE OR CRACK	1.4	0.8	2	0.29
MARIJUANA ONLY	5.6	4.1	7.1	0.75
COCAINE	0.9	0.4	1.4	0.25
CRACK	0.7	0.4	1	0.15
HALLUCINOGEN	0.7	0.3	1.1	0.2
ROHYPNOL	0.8	0.4	1.2	0.21
STEROIDS	0.6	0.3	1	0.18
ECSTASY	0.9	0.2	1.6	0.35
HEROIN	0.8	0.5	1.2	0.17
METHAMPHETAMINE	0.6	0.3	0.9	0.15
SYNTHETIC MARIJ	3.4	2.4	4.4	0.49
CODEINE	6.5	5.3	7.8	0.64
OXYCONTIN ETC	0.7	0.2	1.1	0.2
VICODIN ETC	1.1	0.7	1.5	0.21
VALIUM ETC	0.3	0.1	0.5	0.1
XANAX ETC	0.7	0.4	1	0.14
DXM, ETC.	4.6	3.5	5.8	0.57

INHALANTS	7.6	5.8	9.4	0.89
MARIJUANA	3.9	2.4	5.3	0.7
COCAINE OR CRACK	0.6	0.3	1	0.18
MARIJUANA ONLY	1.8	1.1	2.5	0.35
COCAINE	0.4	0.1	0.8	0.18
CRACK	0.2	0	0.4	0.1
HALLUCINOGEN	0.3	0.1	0.4	0.1
ROHYPNOL	0.1	0	0.3	0.09
STEROIDS	0.2	0	0.4	0.08
ECSTASY	0.1	0	0.2	0.06
HEROIN	0.4	0.2	0.6	0.11
METHAMPHETAMINE	0.3	0	0.5	0.12
SYNTHETIC MARIJ	1.3	0.6	2	0.36
CODEINE	3	2.2	3.8	0.41
OXYCONTIN ETC	0.4	0.1	0.7	0.15
VICODIN ETC	0.6	0.4	0.8	0.1
VALIUM ETC	0.2	0	0.4	0.09
XANAX ETC	0.4	0.1	0.6	0.11
DXM, ETC.	2	1.1	3	0.47

Lifetime Use of Selected Substances - Grade 7 White				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	6.5	3.6	9.3	1.42
CIGARETTES	5.6	2.9	8.3	1.36
SMOKELESS TOBACCO	2.2	1.1	3.3	0.56
ALCOHOL	23	18.1	28	2.47
BEER	16	12.4	19.6	1.8
WINE COOLER	10.2	7.1	13.4	1.57
WINE	14.7	10.3	19.2	2.22
LIQUOR	7.4	4.9	9.9	1.25
ANY ILLICIT DRUG	3.1	1.8	4.5	0.66
INHALANTS	8.8	6.8	10.9	1.01
MARIJUANA	2.5	1.3	3.7	0.61
COCAINE OR CRACK	0.3	0.1	0.4	0.09
MARIJUANA ONLY	1.6	0.7	2.5	0.46
COCAINE	0.2	0	0.3	0.08
CRACK	0.2	0	0.3	0.07
HALLUCINOGEN	0.4	0.1	0.7	0.16
ROHYPNOL	0.2	0	0.3	0.07
STEROIDS	0.7	0.3	1.1	0.18
ECSTASY	0.1	0	0.3	0.06
HEROIN	0.3	0.1	0.5	0.11
METHAMPHETAMINE	0.3	0.1	0.5	0.12
SYNTHETIC MARIJ	0.7	0.2	1.2	0.25
CODEINE	4.8	3.4	6.3	0.7
OXYCONTIN ETC	0.5	0.1	0.8	0.18
VICODIN ETC	0.9	0.4	1.3	0.21
VALIUM ETC	0.2	0	0.4	0.09
XANAX ETC	0.5	0.2	0.8	0.15
DXM, ETC.	1.5	0.5	2.5	0.47

Past Month Use of Selected Substances - Grade 7 White				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	1.9	0.9	2.9	0.48
CIGARETTES	1.4	0.6	2.2	0.39
SMOKELESS TOBACCO	0.7	0.1	1.4	0.33
ALCOHOL	4.8	3.5	6.2	0.67
BEER	2.9	1.5	4.3	0.7
WINE COOLER	2.2	1.3	3	0.44
WINE	2	1.3	2.8	0.37
LIQUOR	1.5	0.6	2.3	0.42
ANY ILLICIT DRUG	1.4	0.7	2.2	0.39
INHALANTS	2.7	1.5	3.9	0.58
MARIJUANA	1.1	0.4	1.9	0.36
COCAINE OR CRACK	0.1	0	0.3	0.07
MARIJUANA ONLY	0.7	0	1.4	0.34
COCAINE	0.1	0	0.2	0.06
CRACK	0.1	0	0.1	0.04
HALLUCINOGEN	0.1	0	0.4	0.12
ROHYPNOL	0	0	0.1	0.03
STEROIDS	0.1	0	0.3	0.07
ECSTASY	0	0	0.1	0.03
HEROIN	0.2	0	0.4	0.09
METHAMPHETAMINE	0.1	0	0.2	0.06
SYNTHETIC MARIJ	0.2	0	0.3	0.09
CODEINE	2	1.2	2.8	0.4
OXYCONTIN ETC	0.2	0	0.3	0.08
VICODIN ETC	0.4	0.1	0.7	0.16
VALIUM ETC	0.1	0	0.2	0.06
XANAX ETC	0.2	0	0.5	0.11
DXM, ETC.	0.4	0.1	0.8	0.18

Lifetime Use of Selected Substances - Grade 7 Female				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	9.8	7.2	12.3	1.28
CIGARETTES	8.8	6.2	11.3	1.29
SMOKELESS TOBACCO	2.7	1.8	3.5	0.42
ALCOHOL	32.2	27.4	37	2.39

Past Month Use of Selected Substances - Grade 7 Female				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	2.5	1.6	3.3	0.42
CIGARETTES	1.8	1.1	2.5	0.36
SMOKELESS TOBACCO	0.8	0.5	1.2	0.19
ALCOHOL	10.2	7.8	12.6	1.21

BEER	21	17.1	25	1.96
WINE COOLER	15.6	12.3	18.8	1.62
WINE	19.4	15.5	23.2	1.91
LIQUOR	13.4	10.4	16.4	1.51
ANY ILLICIT DRUG	7.6	5.5	9.6	1.03
INHALANTS	16.3	13	19.6	1.64
MARIJUANA	6.1	4.1	8.1	1
COCAINE OR CRACK	0.9	0.4	1.5	0.29
MARIJUANA ONLY	3.8	2.4	5.1	0.68
COCAINE	0.6	0.2	1	0.22
CRACK	0.5	0.1	0.9	0.18
HALLUCINOGEN	0.6	0.2	1	0.2
ROHYPNOL	0.5	0.2	0.9	0.18
STEROIDS	0.7	0.4	1.1	0.18
ECSTASY	0.5	0	0.9	0.2
HEROIN	0.8	0.4	1.2	0.21
METHAMPHETAMINE	0.4	0.2	0.7	0.13
SYNTHETIC MARIJ	1.9	1	2.9	0.46
CODEINE	6.7	5.3	8.2	0.73
OXYCONTIN ETC	0.7	0.3	1.1	0.19
VICODIN ETC	1.3	0.8	1.8	0.25
VALIUM ETC	0.5	0.2	0.7	0.14
XANAX ETC	0.7	0.4	1.1	0.18
DXM, ETC.	3.4	2.3	4.5	0.56

BEER	5.6	3.8	7.3	0.86
WINE COOLER	5.3	3.6	7	0.83
WINE	4.9	3.5	6.4	0.74
LIQUOR	3.9	2.4	5.3	0.7
ANY ILLICIT DRUG	3	1.8	4.1	0.57
INHALANTS	6.2	4.4	8	0.89
MARIJUANA	2.3	1.3	3.3	0.51
COCAINE OR CRACK	0.4	0.1	0.7	0.15
MARIJUANA ONLY	1.1	0.5	1.7	0.29
COCAINE	0.3	0	0.7	0.15
CRACK	0.1	0	0.2	0.05
HALLUCINOGEN	0.2	0	0.4	0.1
ROHYPNOL	0.1	0	0.3	0.08
STEROIDS	0.2	0	0.3	0.06
ECSTASY	0.1	0	0.2	0.05
HEROIN	0.4	0.1	0.7	0.16
METHAMPHETAMINE	0.2	0	0.4	0.09
SYNTHETIC MARIJ	0.6	0	1.3	0.31
CODEINE	2.9	2	3.8	0.44
OXYCONTIN ETC	0.5	0.2	0.8	0.17
VICODIN ETC	0.7	0.4	1	0.16
VALIUM ETC	0.3	0	0.5	0.1
XANAX ETC	0.4	0.1	0.6	0.12
DXM, ETC.	1.4	0.3	2.4	0.52

Lifetime Use of Selected Substances - Grade 7 Male				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	10.3	7.7	12.9	1.27
CIGARETTES	8.8	6.3	11.3	1.23
SMOKELESS TOBACCO	3.6	2.4	4.8	0.59
ALCOHOL	29.6	24.9	34.3	2.35
BEER	22.4	18.3	26.5	2.04
WINE COOLER	13.1	9.9	16.3	1.6
WINE	16.7	13	20.4	1.86
LIQUOR	10.7	8.3	13	1.17
ANY ILLICIT DRUG	8.4	6.2	10.5	1.07
INHALANTS	12.1	10.1	14	0.97
MARIJUANA	7.7	5.5	10	1.12
COCAINE OR CRACK	0.7	0.3	1.1	0.2
MARIJUANA ONLY	4.3	3	5.7	0.68
COCAINE	0.5	0.1	0.8	0.17
CRACK	0.3	0.1	0.4	0.09
HALLUCINOGEN	0.6	0.3	0.9	0.14
ROHYPNOL	0.4	0.2	0.7	0.14
STEROIDS	0.6	0.2	1	0.19
ECSTASY	0.7	0.3	1.2	0.23
HEROIN	0.5	0.2	0.8	0.15
METHAMPHETAMINE	0.5	0.2	0.8	0.14
SYNTHETIC MARIJ	2.5	1.7	3.3	0.38
CODEINE	4.5	3.6	5.5	0.48
OXYCONTIN ETC	0.4	0	0.7	0.16
VICODIN ETC	0.5	0.2	0.9	0.18
VALIUM ETC	0.1	0	0.1	0.03
XANAX ETC	0.3	0.1	0.5	0.11
DXM, ETC.	2.9	2	3.8	0.44

Past Month Use of Selected Substances - Grade 7 Male				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	2.6	1.7	3.5	0.44
CIGARETTES	1.8	1.2	2.5	0.32
SMOKELESS TOBACCO	1.1	0.4	1.8	0.35
ALCOHOL	9.2	7.4	11	0.91
BEER	5.2	3.7	6.7	0.75
WINE COOLER	3.9	2.9	4.9	0.5
WINE	4	2.8	5.2	0.59
LIQUOR	3.1	2.1	4.1	0.49
ANY ILLICIT DRUG	3.8	2.4	5.2	0.69
INHALANTS	5	3.8	6.3	0.63
MARIJUANA	3.3	1.8	4.7	0.72
COCAINE OR CRACK	0.3	0.1	0.6	0.11
MARIJUANA ONLY	1.7	0.6	2.8	0.56
COCAINE	0.2	0	0.4	0.09
CRACK	0.1	0	0.3	0.07
HALLUCINOGEN	0.2	0	0.3	0.07
ROHYPNOL	0.1	0	0.1	0.04
STEROIDS	0.2	0	0.5	0.15
ECSTASY	0.1	0	0.2	0.05
HEROIN	0.3	0.1	0.6	0.12
METHAMPHETAMINE	0.2	0	0.4	0.1
SYNTHETIC MARIJ	1	0.5	1.5	0.25
CODEINE	2.4	1.7	3.2	0.37
OXYCONTIN ETC	0.1	0	0.2	0.07
VICODIN ETC	0.2	0	0.4	0.09
VALIUM ETC	0	0	0.1	0.02
XANAX ETC	0.1	0	0.2	0.06
DXM, ETC.	1.3	0.8	1.8	0.25

Lifetime Use of Selected Substances - Grade 7 Border			
Category	Prevalence	Boundaries	Standard

Past Month Use of Selected Substances - Grade 7 Border			
Category	Prevalence	Boundaries	Standard

		Lower	Upper	Error
TOBACCO	11.5	8	15	1.51
CIGARETTES	10.6	7.5	13.8	1.37
SMOKELESS TOBACCO	3.1	1.7	4.5	0.59
ALCOHOL	28.5	21.9	35.1	2.85
BEER	20.8	15.7	25.9	2.22
WINE COOLER	13.2	8	18.3	2.23
WINE	13.5	9.5	17.6	1.75
LIQUOR	12.1	8.1	16.1	1.74
ANY ILLICIT DRUG	9.7	4.6	14.8	2.23
INHALANTS	17.3	12.1	22.5	2.27
MARIJUANA	8.4	3.6	13.3	2.09
COCAINE OR CRACK	1.6	0.1	3.1	0.66
MARIJUANA ONLY	4.5	1.4	7.6	1.34
COCAINE	1.4	0	2.7	0.59
CRACK	0.6	0	1.2	0.28
HALLUCINOGEN	0.8	0.1	1.5	0.3
ROHYPNOL	0.6	0	1.3	0.31
STEROIDS	0.9	0.5	1.4	0.19
ECSTASY	0.8	0.1	1.5	0.3
HEROIN	0.4	0	0.9	0.24
METHAMPHETAMINE	0.6	0	1.3	0.31
SYNTHETIC MARIJ	3.7	2.2	5.2	0.65
CODEINE	4.2	2.5	5.9	0.74
OXYCONTIN ETC	0.6	0.1	1	0.21
VICODIN ETC	1.1	0.7	1.5	0.18
VALIUM ETC	0.3	0	0.6	0.12
XANAX ETC	0.6	0.1	1.1	0.2
DXM, ETC.	3.1	1.6	4.6	0.63

		Lower	Upper	Error
TOBACCO	3.5	1.8	5.2	0.74
CIGARETTES	2.9	1.5	4.3	0.59
SMOKELESS TOBACCO	1	0.4	1.6	0.25
ALCOHOL	9.2	7.1	11.4	0.94
BEER	5.6	3.5	7.7	0.93
WINE COOLER	4	2.1	5.8	0.8
WINE	3.9	2.8	5	0.47
LIQUOR	3.8	2.8	4.9	0.47
ANY ILLICIT DRUG	4.5	1.8	7.1	1.16
INHALANTS	7.5	4.7	10.2	1.18
MARIJUANA	3.4	0.8	6	1.12
COCAINE OR CRACK	0.8	0	1.8	0.43
MARIJUANA ONLY	1.6	0	3.3	0.73
COCAINE	0.7	0	1.7	0.43
CRACK	0.2	0	0.6	0.16
HALLUCINOGEN	0.4	0.1	0.8	0.15
ROHYPNOL	0.1	0	0.3	0.08
STEROIDS	0.6	0.2	1	0.17
ECSTASY	0.3	0	0.6	0.12
HEROIN	0.1	0	0.3	0.09
METHAMPHETAMINE	0.5	0	1.2	0.31
SYNTHETIC MARIJ	1.5	0.5	2.4	0.41
CODEINE	1.7	0.6	2.8	0.47
OXYCONTIN ETC	0.2	0	0.5	0.12
VICODIN ETC	0.4	0.2	0.7	0.1
VALIUM ETC	0.1	0	0.4	0.11
XANAX ETC	0.4	0	0.9	0.21
DXM, ETC.	1.1	0.2	2	0.4

Lifetime Use of Selected Substances - Grade 7 Non-Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	9.8	7.2	12.4	1.31
CIGARETTES	8.5	5.9	11	1.27
SMOKELESS TOBACCO	3.2	2.3	4.1	0.44
ALCOHOL	31.3	26.2	36.4	2.52
BEER	21.9	17.7	26.1	2.1
WINE COOLER	14.6	11.3	17.8	1.61
WINE	18.8	14.9	22.7	1.93
LIQUOR	12	9.3	14.8	1.37
ANY ILLICIT DRUG	7.7	5.4	9.9	1.12
INHALANTS	13.7	11.1	16.2	1.24
MARIJUANA	6.7	4.4	9	1.14
COCAINE OR CRACK	0.7	0.3	1.1	0.19
MARIJUANA ONLY	4	2.5	5.4	0.7
COCAINE	0.4	0.1	0.7	0.15
CRACK	0.4	0.2	0.6	0.1
HALLUCINOGEN	0.5	0.3	0.8	0.13
ROHYPNOL	0.5	0.1	0.8	0.15
STEROIDS	0.6	0.3	0.9	0.15
ECSTASY	0.6	0.1	1	0.22
HEROIN	0.7	0.5	1	0.12
METHAMPHETAMINE	0.5	0.3	0.7	0.09
SYNTHETIC MARIJ	2	1.2	2.8	0.4
CODEINE	5.9	4.7	7	0.57
OXYCONTIN ETC	0.5	0.3	0.8	0.12
VICODIN ETC	0.9	0.5	1.3	0.19
VALIUM ETC	0.2	0.1	0.4	0.08

Past Month Use of Selected Substances - Grade 7 Non-Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	2.4	1.6	3.1	0.37
CIGARETTES	1.7	1.1	2.3	0.29
SMOKELESS TOBACCO	1	0.4	1.5	0.27
ALCOHOL	9.8	7.5	12.1	1.14
BEER	5.4	3.8	7	0.79
WINE COOLER	4.7	3.4	6	0.65
WINE	4.6	3.3	5.8	0.63
LIQUOR	3.4	2.3	4.6	0.58
ANY ILLICIT DRUG	3.2	2	4.4	0.58
INHALANTS	5.3	3.9	6.7	0.71
MARIJUANA	2.7	1.5	3.8	0.58
COCAINE OR CRACK	0.3	0.1	0.5	0.1
MARIJUANA ONLY	1.4	0.5	2.2	0.4
COCAINE	0.2	0	0.4	0.09
CRACK	0.1	0	0.2	0.05
HALLUCINOGEN	0.1	0	0.3	0.07
ROHYPNOL	0.1	0	0.2	0.06
STEROIDS	0.1	0	0.3	0.09
ECSTASY	0	0	0.1	0.03
HEROIN	0.5	0.3	0.7	0.09
METHAMPHETAMINE	0.2	0.1	0.3	0.07
SYNTHETIC MARIJ	0.7	0.3	1.2	0.23
CODEINE	2.8	2.1	3.6	0.39
OXYCONTIN ETC	0.3	0.1	0.5	0.1
VICODIN ETC	0.4	0.2	0.7	0.11
VALIUM ETC	0.1	0	0.3	0.06

XANAX ETC	0.5	0.3	0.7	0.12
DXM, ETC.	3.2	2.2	4.2	0.51

XANAX ETC	0.3	0.1	0.4	0.08
DXM, ETC.	1.4	0.6	2.2	0.4

Lifetime Use of Selected Substances - Grade 8				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	15	12.4	17.7	1.31
CIGARETTES	13.5	11.1	15.9	1.19
SMOKELESS TOBACCO	5.6	4.3	7	0.67
ALCOHOL	40.3	36.4	44.2	1.93
BEER	29.3	26.1	32.6	1.62
WINE COOLER	21	18	24	1.49
WINE	23.1	20.7	25.5	1.19
LIQUOR	22	18.6	25.3	1.66
ANY ILLICIT DRUG	14.4	11.4	17.5	1.53
INHALANTS	14.8	12.9	16.7	0.95
MARIJUANA	13.2	10.4	16	1.4
COCAINE OR CRACK	1.8	1.1	2.5	0.36
MARIJUANA ONLY	7.6	5.9	9.3	0.84
COCAINE	1.3	0.7	1.9	0.29
CRACK	0.9	0.6	1.3	0.19
HALLUCINOGEN	1.2	0.8	1.5	0.17
ROHYPNOL	0.8	0.3	1.2	0.2
STEROIDS	1.1	0.8	1.5	0.19
ECSTASY	1.5	0.9	2	0.28
HEROIN	0.3	0.2	0.5	0.08
METHAMPHETAMINE	0.7	0.4	1	0.16
SYNTHETIC MARIJ	4.3	3.2	5.5	0.57
CODEINE	7.5	6	9	0.75
OXYCONTIN ETC	1.1	0.7	1.5	0.2
VICODIN ETC	2.1	1.2	3.1	0.48
VALIUM ETC	0.8	0.5	1.1	0.15
XANAX ETC	1.5	1	2	0.25
DXM, ETC.	3.7	2.8	4.6	0.47

Past Month Use of Selected Substances - Grade 8				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	5.1	4	6.2	0.54
CIGARETTES	4	3.1	4.9	0.44
SMOKELESS TOBACCO	2	1.4	2.7	0.33
ALCOHOL	13.2	10.8	15.5	1.17
BEER	7.7	6	9.5	0.88
WINE COOLER	6.2	4.8	7.6	0.69
WINE	4.9	3.8	6.1	0.57
LIQUOR	6.8	5.4	8.2	0.7
ANY ILLICIT DRUG	5.3	4	6.5	0.63
INHALANTS	4.4	3.2	5.6	0.6
MARIJUANA	4.6	3.4	5.7	0.57
COCAINE OR CRACK	0.5	0.3	0.7	0.12
MARIJUANA ONLY	1.9	1.3	2.4	0.26
COCAINE	0.3	0.1	0.4	0.08
CRACK	0.3	0.1	0.5	0.11
HALLUCINOGEN	0.4	0.2	0.6	0.1
ROHYPNOL	0.2	0.1	0.3	0.07
STEROIDS	0.4	0.2	0.6	0.1
ECSTASY	0.4	0.2	0.7	0.14
HEROIN	0.1	0	0.2	0.04
METHAMPHETAMINE	0.1	0	0.2	0.05
SYNTHETIC MARIJ	1.6	1	2.2	0.28
CODEINE	3.6	2.9	4.4	0.39
OXYCONTIN ETC	0.5	0.2	0.7	0.12
VICODIN ETC	1.1	0.5	1.6	0.26
VALIUM ETC	0.4	0.2	0.6	0.1
XANAX ETC	0.6	0.3	0.9	0.13
DXM, ETC.	1.9	1.3	2.5	0.29

Lifetime Use of Selected Substances - Grade 8 African American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	9.6	4.1	15.2	2.72
CIGARETTES	9	3.6	14.4	2.66
SMOKELESS TOBACCO	1.9	0.2	3.7	0.87
ALCOHOL	39.1	30.4	47.8	4.3
BEER	20.3	13.7	26.9	3.26
WINE COOLER	19.8	12.7	26.9	3.48
WINE	21.2	13.5	28.8	3.75
LIQUOR	18.5	9.2	27.9	4.62
ANY ILLICIT DRUG	12	8.3	15.6	1.8
INHALANTS	11.2	7	15.3	2.05
MARIJUANA	10.4	6.6	14.2	1.86
COCAINE OR CRACK	0.2	0	0.5	0.14
MARIJUANA ONLY	8.5	4.3	12.8	2.1
COCAINE	0	*	*	*
CRACK	0.2	0	0.5	0.14
HALLUCINOGEN	0	*	*	*
ROHYPNOL	0	*	*	*
STEROIDS	2.8	0.4	5.2	1.16
ECSTASY	0.3	0	0.8	0.22
HEROIN	0	*	*	*
METHAMPHETAMINE	0.6	0	1.7	0.51

Past Month Use of Selected Substances - Grade 8 African American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	1.4	0	2.9	0.71
CIGARETTES	1.3	0	2.6	0.67
SMOKELESS TOBACCO	0.5	0	1.2	0.33
ALCOHOL	9.2	6.8	11.7	1.19
BEER	3.3	1.1	5.4	1.06
WINE COOLER	3.4	0.9	5.9	1.22
WINE	4.1	2.3	5.8	0.87
LIQUOR	4.3	1.4	7.2	1.44
ANY ILLICIT DRUG	2.3	1	3.7	0.66
INHALANTS	2.9	1.1	4.7	0.9
MARIJUANA	1.9	0.7	3.2	0.61
COCAINE OR CRACK	0	*	*	*
MARIJUANA ONLY	1.8	0.5	3	0.6
COCAINE	0	*	*	*
CRACK	0	*	*	*
HALLUCINOGEN	0	*	*	*
ROHYPNOL	0	*	*	*
STEROIDS	0.9	0	2.2	0.64
ECSTASY	0.1	0	0.3	0.08
HEROIN	0	*	*	*
METHAMPHETAMINE	0	*	*	*

SYNTHETIC MARIJ	2.4	0	5	1.24
CODEINE	9.4	3.5	15.3	2.91
OXYCONTIN ETC	0.1	0	0.2	0.07
VICODIN ETC	1.7	0	4.3	1.27
VALIUM ETC	0	*	*	*
XANAX ETC	0.2	0	0.5	0.15
DXM, ETC.	5.2	1.5	8.9	1.81

SYNTHETIC MARIJ	0.5	0	1.2	0.36
CODEINE	4.5	0.9	8	1.74
OXYCONTIN ETC	0	*	*	*
VICODIN ETC	1.4	0	4	1.26
VALIUM ETC	0	*	*	*
XANAX ETC	0	*	*	*
DXM, ETC.	3.7	1	6.4	1.33

Lifetime Use of Selected Substances - Grade 8 Mexican American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	18.5	16.8	20.2	0.86
CIGARETTES	16.9	15	18.8	0.94
SMOKELESS TOBACCO	6.6	5.5	7.6	0.53
ALCOHOL	45	41.9	48.1	1.55
BEER	35	32.2	37.7	1.36
WINE COOLER	24.7	21.4	28.1	1.66
WINE	25.5	22.9	28	1.27
LIQUOR	26.2	22.6	29.8	1.79
ANY ILLICIT DRUG	19.6	16.8	22.4	1.39
INHALANTS	18.5	16.4	20.6	1.04
MARIJUANA	18	15.3	20.7	1.35
COCAINE OR CRACK	3.2	2.2	4.2	0.51
MARIJUANA ONLY	9.4	7.8	11.1	0.83
COCAINE	2.4	1.5	3.3	0.46
CRACK	1.5	1	2.1	0.25
HALLUCINOGEN	1.7	1.2	2.2	0.25
ROHYPNOL	1.4	0.7	2.1	0.34
STEROIDS	0.9	0.5	1.3	0.2
ECSTASY	2.2	1.5	2.9	0.34
HEROIN	0.5	0.2	0.8	0.14
METHAMPHETAMINE	1	0.5	1.6	0.28
SYNTHETIC MARIJ	6.3	4.9	7.7	0.69
CODEINE	9.2	7.3	11.1	0.97
OXYCONTIN ETC	1.2	0.7	1.6	0.23
VICODIN ETC	2.6	1.2	3.9	0.67
VALIUM ETC	1.2	0.8	1.7	0.22
XANAX ETC	2	1.1	2.8	0.42
DXM, ETC.	4.4	3	5.7	0.66

Past Month Use of Selected Substances - Grade 8 Mexican American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	6.6	5.5	7.7	0.56
CIGARETTES	5.3	4.2	6.3	0.53
SMOKELESS TOBACCO	2.5	1.8	3.3	0.37
ALCOHOL	16.7	14.1	19.3	1.28
BEER	10.3	8.3	12.3	0.99
WINE COOLER	8.5	6.6	10.3	0.93
WINE	6.5	5.4	7.6	0.55
LIQUOR	8.7	6.9	10.5	0.89
ANY ILLICIT DRUG	7.7	6.2	9.2	0.75
INHALANTS	6.2	4.8	7.6	0.7
MARIJUANA	6.7	5.2	8.2	0.74
COCAINE OR CRACK	0.8	0.5	1.1	0.15
MARIJUANA ONLY	2.4	1.8	3.1	0.34
COCAINE	0.6	0.3	0.9	0.15
CRACK	0.4	0.2	0.7	0.13
HALLUCINOGEN	0.8	0.5	1.1	0.15
ROHYPNOL	0.4	0.1	0.7	0.13
STEROIDS	0.3	0	0.6	0.15
ECSTASY	0.6	0.2	1	0.19
HEROIN	0.2	0	0.3	0.08
METHAMPHETAMINE	0.2	0	0.4	0.09
SYNTHETIC MARIJ	2.4	1.6	3.1	0.37
CODEINE	4.6	3.5	5.7	0.55
OXYCONTIN ETC	0.6	0.3	0.9	0.15
VICODIN ETC	1.3	0.6	2	0.37
VALIUM ETC	0.6	0.3	1	0.17
XANAX ETC	0.8	0.3	1.3	0.23
DXM, ETC.	2.1	1.2	3	0.44

Lifetime Use of Selected Substances - Grade 8 White				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	12.6	8.5	16.7	2.04
CIGARETTES	10.9	7.3	14.4	1.77
SMOKELESS TOBACCO	5.4	3.1	7.7	1.14
ALCOHOL	35.6	29	42.1	3.26
BEER	24.7	19.7	29.6	2.46
WINE COOLER	17.1	12.8	21.4	2.13
WINE	21.2	17.6	24.9	1.81
LIQUOR	18	12.6	23.4	2.68
ANY ILLICIT DRUG	8.7	5.6	11.8	1.53
INHALANTS	11.5	8.8	14.3	1.35
MARIJUANA	8	5.2	10.8	1.39
COCAINE OR CRACK	0.6	0	1.2	0.31
MARIJUANA ONLY	5.5	3.1	7.8	1.18
COCAINE	0.3	0	0.6	0.17
CRACK	0.5	0	1.1	0.3
HALLUCINOGEN	0.7	0.2	1.3	0.27

Past Month Use of Selected Substances - Grade 8 White				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	4	2.4	5.6	0.79
CIGARETTES	2.9	1.6	4.1	0.62
SMOKELESS TOBACCO	1.7	0.7	2.6	0.47
ALCOHOL	9.4	7.1	11.8	1.17
BEER	5.7	3.8	7.6	0.95
WINE COOLER	3.7	2.6	4.9	0.58
WINE	2.8	1.7	3.9	0.54
LIQUOR	5.1	3.3	6.9	0.88
ANY ILLICIT DRUG	2.7	1.5	4	0.62
INHALANTS	2.3	1.4	3.2	0.44
MARIJUANA	2.4	1.3	3.4	0.52
COCAINE OR CRACK	0.3	0	0.9	0.29
MARIJUANA ONLY	1.3	0.6	2	0.34
COCAINE	0	*	*	*
CRACK	0.3	0	0.9	0.29
HALLUCINOGEN	0	*	*	*

ROHYPNOL	0.1	0	0.3	0.07
STEROIDS	1.1	0.6	1.5	0.21
ECSTASY	0.7	0.1	1.2	0.29
HEROIN	0.3	0.1	0.5	0.11
METHAMPHETAMINE	0.4	0	0.8	0.19
SYNTHETIC MARIJ	1.9	0.7	3.1	0.6
CODEINE	5.1	3.6	6.5	0.71
OXYCONTIN ETC	1	0.4	1.7	0.32
VICODIN ETC	1.7	0.6	2.9	0.56
VALIUM ETC	0.5	0.1	0.9	0.2
XANAX ETC	1.1	0.5	1.7	0.28
DXM, ETC.	2.2	1.1	3.3	0.53

ROHYPNOL	0	*	*	*
STEROIDS	0.3	0.1	0.6	0.13
ECSTASY	0.2	0	0.5	0.15
HEROIN	0	0	0.1	0.03
METHAMPHETAMINE	0	0	0.1	0.03
SYNTHETIC MARIJ	0.5	0.1	0.8	0.16
CODEINE	2.4	1.6	3.2	0.39
OXYCONTIN ETC	0.5	0.2	0.9	0.18
VICODIN ETC	0.8	0.2	1.3	0.26
VALIUM ETC	0.2	0	0.4	0.1
XANAX ETC	0.5	0.2	0.8	0.15
DXM, ETC.	1.1	0.5	1.7	0.29

Lifetime Use of Selected Substances - Grade 8 Female				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	14.4	11.2	17.6	1.59
CIGARETTES	13.4	10.3	16.4	1.53
SMOKELESS TOBACCO	4.1	3	5.3	0.59
ALCOHOL	42.3	36.9	47.8	2.72
BEER	29.8	24.6	34.9	2.56
WINE COOLER	24.2	19.7	28.7	2.24
WINE	26.2	22.6	29.8	1.8
LIQUOR	23.9	19	28.7	2.4
ANY ILLICIT DRUG	14.2	10.8	17.7	1.72
INHALANTS	17.1	14	20.2	1.55
MARIJUANA	12.8	9.8	15.9	1.52
COCAINE OR CRACK	2.5	1.3	3.7	0.61
MARIJUANA ONLY	7.2	5.3	9.1	0.95
COCAINE	1.7	0.8	2.7	0.47
CRACK	1.4	0.7	2.1	0.35
HALLUCINOGEN	1.1	0.6	1.6	0.26
ROHYPNOL	0.8	0.4	1.2	0.21
STEROIDS	1.2	0.8	1.7	0.24
ECSTASY	1.1	0.6	1.5	0.24
HEROIN	0.4	0.2	0.6	0.1
METHAMPHETAMINE	0.6	0.2	1.1	0.23
SYNTHETIC MARIJ	4.1	2.7	5.5	0.71
CODEINE	7.6	5.8	9.4	0.89
OXYCONTIN ETC	0.9	0.5	1.3	0.2
VICODIN ETC	2.3	1.2	3.4	0.55
VALIUM ETC	0.9	0.5	1.3	0.21
XANAX ETC	1.7	1.2	2.3	0.29
DXM, ETC.	3.6	2.5	4.7	0.55

Past Month Use of Selected Substances - Grade 8 Female				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	5.1	3.9	6.3	0.59
CIGARETTES	4.4	3.3	5.5	0.55
SMOKELESS TOBACCO	1.4	0.8	2	0.28
ALCOHOL	14.4	11.2	17.7	1.62
BEER	8.4	5.9	10.9	1.25
WINE COOLER	7.1	5.3	8.9	0.88
WINE	5.5	4	7.1	0.75
LIQUOR	8.3	6.4	10.2	0.95
ANY ILLICIT DRUG	5.2	3.6	6.8	0.79
INHALANTS	5	3.4	6.6	0.81
MARIJUANA	4.6	3.1	6	0.72
COCAINE OR CRACK	0.6	0.3	0.9	0.16
MARIJUANA ONLY	1.6	1	2.3	0.32
COCAINE	0.4	0.2	0.7	0.14
CRACK	0.4	0.1	0.6	0.12
HALLUCINOGEN	0.4	0.1	0.7	0.15
ROHYPNOL	0.2	0	0.4	0.09
STEROIDS	0.4	0	0.7	0.16
ECSTASY	0.2	0	0.3	0.08
HEROIN	0	0	0.1	0.03
METHAMPHETAMINE	0.1	0	0.2	0.08
SYNTHETIC MARIJ	1.4	0.9	1.9	0.25
CODEINE	3.5	2.4	4.6	0.56
OXYCONTIN ETC	0.4	0.1	0.6	0.12
VICODIN ETC	1.2	0.6	1.7	0.27
VALIUM ETC	0.4	0.1	0.6	0.13
XANAX ETC	0.6	0.3	1	0.18
DXM, ETC.	1.8	1	2.5	0.37

Lifetime Use of Selected Substances - Grade 8 Male				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	15.6	12.5	18.7	1.53
CIGARETTES	13.6	10.9	16.2	1.32
SMOKELESS TOBACCO	7.1	5	9.2	1.03
ALCOHOL	38.6	34.7	42.5	1.94
BEER	29.1	26	32.1	1.54
WINE COOLER	17.8	15.1	20.6	1.37
WINE	20.4	17.6	23.2	1.38
LIQUOR	20.2	17.5	22.9	1.36
ANY ILLICIT DRUG	14.7	11.6	17.8	1.57
INHALANTS	12.7	11.1	14.2	0.79
MARIJUANA	13.5	10.6	16.4	1.46

Past Month Use of Selected Substances - Grade 8 Male				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	5.1	3.7	6.5	0.7
CIGARETTES	3.5	2.4	4.7	0.55
SMOKELESS TOBACCO	2.6	1.6	3.5	0.47
ALCOHOL	11.8	9.7	13.8	1.04
BEER	7.2	5.4	9.1	0.93
WINE COOLER	5.2	3.8	6.6	0.69
WINE	4.3	3.2	5.5	0.57
LIQUOR	5.2	3.7	6.7	0.74
ANY ILLICIT DRUG	5.4	4	6.8	0.71
INHALANTS	3.8	2.6	4.9	0.57
MARIJUANA	4.6	3.3	5.8	0.62

COCAINE OR CRACK	1.1	0.6	1.7	0.27
MARIJUANA ONLY	8	6.2	9.9	0.94
COCAINE	0.9	0.4	1.3	0.22
CRACK	0.5	0.1	0.9	0.2
HALLUCINOGEN	1.2	0.7	1.8	0.27
ROHYPNOL	0.7	0.2	1.3	0.27
STEROIDS	1	0.4	1.5	0.28
ECSTASY	1.9	0.9	2.8	0.49
HEROIN	0.3	0.1	0.5	0.1
METHAMPHETAMINE	0.8	0.4	1.2	0.21
SYNTHETIC MARIJ	4.5	3.3	5.7	0.6
CODEINE	7.5	5.9	9.2	0.83
OXYCONTIN ETC	1.4	0.7	2	0.31
VICODIN ETC	2.1	1	3.1	0.51
VALIUM ETC	0.7	0.2	1.2	0.23
XANAX ETC	1.2	0.5	1.9	0.34
DXM, ETC.	3.9	2.7	5	0.59

COCAINE OR CRACK	0.4	0	0.8	0.21
MARIJUANA ONLY	2.1	1.4	2.8	0.37
COCAINE	0.2	0	0.4	0.1
CRACK	0.3	0	0.7	0.2
HALLUCINOGEN	0.4	0.1	0.7	0.16
ROHYPNOL	0.2	0	0.4	0.1
STEROIDS	0.3	0	0.6	0.15
ECSTASY	0.7	0.2	1.2	0.27
HEROIN	0.1	0	0.3	0.07
METHAMPHETAMINE	0.1	0	0.3	0.07
SYNTHETIC MARIJ	1.8	1	2.6	0.4
CODEINE	3.9	3	4.8	0.44
OXYCONTIN ETC	0.6	0.2	1	0.19
VICODIN ETC	1	0.3	1.7	0.33
VALIUM ETC	0.4	0	0.7	0.16
XANAX ETC	0.5	0.2	0.9	0.18
DXM, ETC.	2	1.2	2.8	0.41

Lifetime Use of Selected Substances - Grade 8 Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	18.9	14.1	23.8	2.11
CIGARETTES	18	13.1	23	2.16
SMOKELESS TOBACCO	6.1	3.4	8.7	1.16
ALCOHOL	43.9	36.5	51.3	3.22
BEER	33.6	26.8	40.3	2.92
WINE COOLER	21	15.1	26.9	2.56
WINE	23.2	17.7	28.8	2.42
LIQUOR	21	15.9	26.1	2.22
ANY ILLICIT DRUG	19.1	13	25.2	2.7
INHALANTS	18	13.9	22.1	1.81
MARIJUANA	17.4	11.3	23.5	2.7
COCAINE OR CRACK	2.5	1.2	3.7	0.57
MARIJUANA ONLY	9.1	6.2	12	1.29
COCAINE	1.7	0.6	2.8	0.48
CRACK	1.1	0.5	1.7	0.28
HALLUCINOGEN	1.4	0.3	2.5	0.49
ROHYPNOL	1.5	0.4	2.7	0.49
STEROIDS	0.5	0	1.4	0.4
ECSTASY	2.5	0.3	4.6	0.95
HEROIN	0.7	0.3	1.1	0.17
METHAMPHETAMINE	0.6	0.1	1	0.21
SYNTHETIC MARIJ	6.9	4.3	9.5	1.14
CODEINE	6.3	4.6	8	0.76
OXYCONTIN ETC	0.9	0	2.1	0.5
VICODIN ETC	0.7	0.1	1.4	0.29
VALIUM ETC	0.8	0.2	1.3	0.24
XANAX ETC	1.2	0.4	2.1	0.38
DXM, ETC.	2.4	1.2	3.5	0.5

Past Month Use of Selected Substances - Grade 8 Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	6.7	4.6	8.7	0.88
CIGARETTES	5.8	3.9	7.6	0.81
SMOKELESS TOBACCO	2.3	1.3	3.4	0.46
ALCOHOL	16.1	12.6	19.6	1.5
BEER	10.9	7.9	13.8	1.28
WINE COOLER	7.1	3.7	10.6	1.48
WINE	6.5	4.2	8.8	0.99
LIQUOR	7.1	4.9	9.2	0.93
ANY ILLICIT DRUG	8	5.3	10.7	1.2
INHALANTS	7.6	5.7	9.5	0.84
MARIJUANA	6.7	4.5	8.9	0.97
COCAINE OR CRACK	0.8	0.3	1.3	0.22
MARIJUANA ONLY	2.9	1.6	4.2	0.57
COCAINE	0.5	0	1	0.23
CRACK	0.4	0.1	0.8	0.16
HALLUCINOGEN	0.8	0.1	1.4	0.3
ROHYPNOL	0.6	0	1.2	0.25
STEROIDS	0	*	*	*
ECSTASY	1.4	0	2.8	0.64
HEROIN	0.3	0	0.7	0.19
METHAMPHETAMINE	0.2	0	0.4	0.11
SYNTHETIC MARIJ	3.3	1.8	4.8	0.65
CODEINE	2.4	1.5	3.2	0.37
OXYCONTIN ETC	0.3	0	0.6	0.14
VICODIN ETC	0.4	0	0.8	0.18
VALIUM ETC	0.3	0	0.6	0.14
XANAX ETC	0.5	0.1	0.9	0.18
DXM, ETC.	0.8	0	1.7	0.39

Lifetime Use of Selected Substances - Grade 8 Non-Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	14.4	11.5	17.2	1.41
CIGARETTES	12.7	10.2	15.3	1.24
SMOKELESS TOBACCO	5.6	4	7.1	0.76
ALCOHOL	39.7	35.3	44.1	2.17
BEER	28.6	25.1	32.2	1.77
WINE COOLER	21	17.6	24.4	1.7

Past Month Use of Selected Substances - Grade 8 Non-Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	4.8	3.6	6	0.59
CIGARETTES	3.7	2.7	4.6	0.47
SMOKELESS TOBACCO	2	1.2	2.7	0.37
ALCOHOL	12.6	10	15.3	1.32
BEER	7.2	5.3	9.1	0.95
WINE COOLER	6.1	4.5	7.6	0.76

WINE	23.1	20.4	25.8	1.34
LIQUOR	22.1	18.3	26	1.92
ANY ILLICIT DRUG	13.6	10.2	17	1.7
INHALANTS	14.3	12.2	16.3	1.03
MARIJUANA	12.4	9.3	15.5	1.54
COCAINE OR CRACK	1.7	0.8	2.5	0.41
MARIJUANA ONLY	7.3	5.4	9.2	0.96
COCAINE	1.2	0.5	1.9	0.33
CRACK	0.9	0.5	1.4	0.22
HALLUCINOGEN	1.1	0.7	1.5	0.18
ROHYPNOL	0.6	0.2	1.1	0.23
STEROIDS	1.2	0.8	1.6	0.21
ECSTASY	1.3	0.7	1.8	0.27
HEROIN	0.3	0.1	0.5	0.09
METHAMPHETAMINE	0.7	0.4	1.1	0.19
SYNTHETIC MARIJ	3.9	2.7	5	0.59
CODEINE	7.7	5.9	9.4	0.87
OXYCONTIN ETC	1.1	0.7	1.6	0.22
VICODIN ETC	2.4	1.3	3.5	0.56
VALIUM ETC	0.8	0.5	1.1	0.17
XANAX ETC	1.5	0.9	2.1	0.29
DXM, ETC.	3.9	2.9	5	0.52

WINE	4.6	3.3	5.9	0.64
LIQUOR	6.8	5.1	8.4	0.81
ANY ILLICIT DRUG	4.8	3.5	6.1	0.67
INHALANTS	3.9	2.6	5.1	0.63
MARIJUANA	4.2	3	5.4	0.62
COCAINE OR CRACK	0.4	0.2	0.7	0.13
MARIJUANA ONLY	1.7	1.1	2.2	0.27
COCAINE	0.3	0.1	0.4	0.08
CRACK	0.3	0.1	0.5	0.12
HALLUCINOGEN	0.3	0.1	0.5	0.1
ROHYPNOL	0.1	0	0.3	0.07
STEROIDS	0.4	0.2	0.7	0.12
ECSTASY	0.3	0.1	0.5	0.09
HEROIN	0	0	0.1	0.04
METHAMPHETAMINE	0.1	0	0.2	0.05
SYNTHETIC MARIJ	1.3	0.8	1.8	0.26
CODEINE	3.9	3	4.8	0.45
OXYCONTIN ETC	0.5	0.2	0.8	0.14
VICODIN ETC	1.2	0.6	1.8	0.31
VALIUM ETC	0.4	0.1	0.6	0.12
XANAX ETC	0.6	0.3	0.9	0.15
DXM, ETC.	2.1	1.4	2.7	0.32

Lifetime Use of Selected Substances - Grade 9				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	21	16.3	25.6	2.33
CIGARETTES	18.7	14.2	23.2	2.26
SMOKELESS TOBACCO	8.3	6.3	10.3	1.02
ALCOHOL	50.2	43.1	57.2	3.54
BEER	37.6	31	44.1	3.26
WINE COOLER	30.9	24.8	37	3.05
WINE	32	27.6	36.3	2.17
LIQUOR	33.5	27.3	39.7	3.11
ANY ILLICIT DRUG	22.3	17.5	27.1	2.4
INHALANTS	12.9	10.9	15	1.05
MARIJUANA	21.6	16.8	26.3	2.37
COCAINE OR CRACK	1.5	1.1	1.9	0.2
MARIJUANA ONLY	13.1	10.2	16	1.45
COCAINE	1.4	1	1.8	0.19
CRACK	0.3	0.1	0.4	0.07
HALLUCINOGEN	2.3	1.4	3.1	0.42
ROHYPNOL	1.1	0.5	1.6	0.28
STEROIDS	0.8	0.4	1.2	0.21
ECSTASY	2.5	1.8	3.2	0.35
HEROIN	0.3	0.1	0.5	0.08
METHAMPHETAMINE	0.5	0.2	0.7	0.11
SYNTHETIC MARIJ	6.4	4.7	8.1	0.85
CODEINE	10.4	7.8	13	1.3
OXYCONTIN ETC	1.9	1.3	2.4	0.27
VICODIN ETC	3.8	2.7	4.8	0.52
VALIUM ETC	0.8	0.5	1.2	0.2
XANAX ETC	2.9	2.1	3.6	0.38
DXM, ETC.	3.1	2.1	4.1	0.52

Past Month Use of Selected Substances - Grade 9				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	6.7	4.8	8.6	0.95
CIGARETTES	4.9	3.3	6.6	0.82
SMOKELESS TOBACCO	3.1	2.3	3.9	0.42
ALCOHOL	19.9	15.7	24.1	2.12
BEER	11.9	9	14.7	1.42
WINE COOLER	11	8.2	13.8	1.43
WINE	8.9	6.9	10.9	1
LIQUOR	12.3	9.5	15.1	1.41
ANY ILLICIT DRUG	9.1	7	11.1	1.04
INHALANTS	4	2.9	5.1	0.55
MARIJUANA	8.1	6.2	9.9	0.94
COCAINE OR CRACK	0.6	0.3	0.9	0.15
MARIJUANA ONLY	4	3	5	0.5
COCAINE	0.6	0.3	0.9	0.15
CRACK	0.2	0	0.3	0.07
HALLUCINOGEN	0.7	0.4	1	0.17
ROHYPNOL	0.4	0.2	0.7	0.12
STEROIDS	0.1	0	0.3	0.06
ECSTASY	0.7	0.4	1.1	0.17
HEROIN	0	0	0.1	0.03
METHAMPHETAMINE	0.1	0	0.2	0.05
SYNTHETIC MARIJ	2.1	1.3	2.8	0.37
CODEINE	5.2	3.7	6.6	0.73
OXYCONTIN ETC	1.1	0.7	1.5	0.2
VICODIN ETC	1.9	1.3	2.4	0.27
VALIUM ETC	0.5	0.2	0.8	0.13
XANAX ETC	1.5	1	1.9	0.22
DXM, ETC.	1.3	0.7	2	0.31

Lifetime Use of Selected Substances - Grade 9 African American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	14.1	7.7	20.5	3.18

Past Month Use of Selected Substances - Grade 9 African American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	3.6	1	6.2	1.28

CIGARETTES	11.7	5.6	17.8	3.03
SMOKELESS TOBACCO	4.7	2.6	6.9	1.07
ALCOHOL	49.6	39.6	59.5	4.93
BEER	26.5	19.1	34	3.68
WINE COOLER	33.8	22.3	45.3	5.72
WINE	31.5	27.5	35.5	1.97
LIQUOR	31	21.6	40.4	4.65
ANY ILLICIT DRUG	22.9	13.9	31.9	4.45
INHALANTS	12.2	9.4	15.1	1.42
MARIJUANA	22.5	13.8	31.2	4.3
COCAINE OR CRACK	0.6	0	1.3	0.37
MARIJUANA ONLY	17.4	11.5	23.3	2.94
COCAINE	0.5	0	1.1	0.34
CRACK	0.6	0	1.3	0.37
HALLUCINOGEN	1	0	2.1	0.53
ROHYPNOL	1	0	2	0.49
STEROIDS	0.9	0.1	1.7	0.39
ECSTASY	1.5	0	3.1	0.79
HEROIN	0.4	0	1.1	0.34
METHAMPHETAMINE	0.4	0	0.9	0.23
SYNTHETIC MARIJ	3.3	0.8	5.7	1.23
CODEINE	13.2	6.7	19.7	3.21
OXYCONTIN ETC	1.3	0	2.8	0.72
VICODIN ETC	3.4	1.8	4.9	0.77
VALIUM ETC	0.4	0	1.1	0.33
XANAX ETC	2	0.8	3.2	0.59
DXM, ETC.	2	0.4	3.6	0.8

CIGARETTES	2.5	0.8	4.1	0.83
SMOKELESS TOBACCO	1.7	0.2	3.3	0.78
ALCOHOL	17.2	10.8	23.6	3.16
BEER	5.1	2.8	7.4	1.12
WINE COOLER	9.5	4.2	14.8	2.61
WINE	6.7	4	9.4	1.34
LIQUOR	11	6.6	15.4	2.18
ANY ILLICIT DRUG	8.5	4.6	12.4	1.94
INHALANTS	3.9	2.4	5.4	0.76
MARIJUANA	7.6	4.2	11.1	1.72
COCAINE OR CRACK	0.5	0	1.2	0.36
MARIJUANA ONLY	5.5	3.1	7.9	1.18
COCAINE	0.4	0	1	0.33
CRACK	0.5	0	1.2	0.36
HALLUCINOGEN	0.2	0	0.5	0.16
ROHYPNOL	0.3	0	0.6	0.18
STEROIDS	0.2	0	0.5	0.14
ECSTASY	0.8	0	2.1	0.67
HEROIN	0.1	0	0.3	0.08
METHAMPHETAMINE	0.1	0	0.3	0.09
SYNTHETIC MARIJ	0.5	0	1.1	0.29
CODEINE	7	3.1	10.9	1.92
OXYCONTIN ETC	0.7	0	1.6	0.41
VICODIN ETC	1.3	0.3	2.4	0.52
VALIUM ETC	0.3	0	1	0.33
XANAX ETC	1.6	0.4	2.9	0.61
DXM, ETC.	0.8	0	1.6	0.39

Lifetime Use of Selected Substances - Grade 9 Mexican American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	22.4	19.7	25.2	1.36
CIGARETTES	21.1	18.7	23.5	1.22
SMOKELESS TOBACCO	7.4	5.7	9.1	0.84
ALCOHOL	52.6	48.3	57	2.17
BEER	43.1	38.8	47.3	2.13
WINE COOLER	31.9	28.1	35.7	1.9
WINE	32.1	29	35.2	1.56
LIQUOR	35.4	31.7	39.1	1.83
ANY ILLICIT DRUG	26.5	22.7	30.3	1.9
INHALANTS	13.4	11.5	15.3	0.95
MARIJUANA	25.6	22	29.2	1.8
COCAINE OR CRACK	2.4	1.6	3.1	0.37
MARIJUANA ONLY	15.3	12.6	18	1.35
COCAINE	2.2	1.5	2.9	0.34
CRACK	0.2	0	0.4	0.1
HALLUCINOGEN	2	1.1	2.9	0.44
ROHYPNOL	1.4	0.7	2	0.32
STEROIDS	0.8	0.3	1.3	0.25
ECSTASY	2.5	1.5	3.5	0.5
HEROIN	0.3	0	0.6	0.15
METHAMPHETAMINE	0.5	0.2	0.8	0.15
SYNTHETIC MARIJ	8.2	6.8	9.5	0.68
CODEINE	12	9.7	14.3	1.13
OXYCONTIN ETC	1.9	1.1	2.6	0.37
VICODIN ETC	3.5	2.2	4.8	0.65
VALIUM ETC	0.9	0.3	1.6	0.33
XANAX ETC	3.2	2	4.4	0.6
DXM, ETC.	3.7	2.4	5	0.64

Past Month Use of Selected Substances - Grade 9 Mexican American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	6.5	4.9	8.1	0.81
CIGARETTES	5.4	4	6.8	0.71
SMOKELESS TOBACCO	2.4	1.6	3.3	0.42
ALCOHOL	21.8	18.8	24.9	1.51
BEER	14.7	12.1	17.3	1.32
WINE COOLER	12.1	10.4	13.8	0.84
WINE	10.2	8.6	11.8	0.82
LIQUOR	12.5	10.1	14.9	1.18
ANY ILLICIT DRUG	10.8	8.5	13.1	1.16
INHALANTS	4.4	3	5.8	0.69
MARIJUANA	9.7	7.7	11.6	0.96
COCAINE OR CRACK	1	0.4	1.5	0.27
MARIJUANA ONLY	4.5	3.2	5.7	0.63
COCAINE	0.9	0.4	1.4	0.26
CRACK	0.2	0	0.4	0.09
HALLUCINOGEN	0.8	0.3	1.4	0.26
ROHYPNOL	0.7	0.3	1.1	0.21
STEROIDS	0.1	0	0.3	0.08
ECSTASY	0.7	0.2	1.1	0.23
HEROIN	0.1	0	0.2	0.05
METHAMPHETAMINE	0.2	0	0.3	0.08
SYNTHETIC MARIJ	3.2	2.2	4.3	0.53
CODEINE	5.8	4.5	7.1	0.66
OXYCONTIN ETC	1	0.5	1.5	0.26
VICODIN ETC	2.1	1.3	2.8	0.39
VALIUM ETC	0.6	0.2	1.1	0.23
XANAX ETC	1.8	1.1	2.5	0.35
DXM, ETC.	1.6	0.9	2.3	0.37

Lifetime Use of Selected Substances - Grade 9 White				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	22.9	15.8	30	3.57
CIGARETTES	19.3	12.4	26.3	3.48
SMOKELESS TOBACCO	11.8	8.3	15.2	1.73
ALCOHOL	48.9	38.4	59.3	5.22
BEER	36.9	27.5	46.3	4.71
WINE COOLER	30.3	21.6	39	4.36
WINE	32.7	25	40.4	3.86
LIQUOR	33.7	24.3	43.2	4.72
ANY ILLICIT DRUG	17.4	13.1	21.7	2.15
INHALANTS	12.8	8.7	16.8	2.01
MARIJUANA	16.8	12.3	21.2	2.22
COCAINE OR CRACK	0.8	0.4	1.3	0.22
MARIJUANA ONLY	9.1	6.4	11.8	1.35
COCAINE	0.8	0.4	1.3	0.22
CRACK	0.1	0	0.4	0.1
HALLUCINOGEN	3.1	1.5	4.7	0.79
ROHYPNOL	0.6	0	1.5	0.43
STEROIDS	0.9	0.3	1.5	0.29
ECSTASY	2.7	2	3.3	0.34
HEROIN	0.2	0	0.5	0.13
METHAMPHETAMINE	0.4	0.1	0.8	0.16
SYNTHETIC MARIJ	5.7	3.4	8	1.15
CODEINE	6.8	4	9.5	1.37
OXYCONTIN ETC	2.2	1.2	3.3	0.52
VICODIN ETC	4.5	2.7	6.4	0.91
VALIUM ETC	1	0.3	1.8	0.38
XANAX ETC	2.9	1.4	4.3	0.73
DXM, ETC.	2.5	1.2	3.7	0.62

Past Month Use of Selected Substances - Grade 9 White				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	8.4	5.6	11.3	1.44
CIGARETTES	5.3	2.9	7.8	1.22
SMOKELESS TOBACCO	4.8	3.5	6	0.64
ALCOHOL	19.3	13.6	25	2.83
BEER	11.6	7.7	15.5	1.95
WINE COOLER	10.7	6.4	14.9	2.12
WINE	8.1	5	11.3	1.58
LIQUOR	13.4	9	17.7	2.17
ANY ILLICIT DRUG	7.4	5.4	9.3	0.98
INHALANTS	3.5	1.7	5.2	0.86
MARIJUANA	6.7	4.6	8.8	1.05
COCAINE OR CRACK	0.3	0.1	0.5	0.11
MARIJUANA ONLY	3.2	2.1	4.4	0.58
COCAINE	0.3	0.1	0.5	0.11
CRACK	0	*	*	*
HALLUCINOGEN	0.7	0.2	1.1	0.23
ROHYPNOL	0.1	0	0.1	0.04
STEROIDS	0.1	0	0.3	0.08
ECSTASY	0.7	0.2	1.1	0.23
HEROIN	0	0	0	0.01
METHAMPHETAMINE	0.1	0	0.3	0.1
SYNTHETIC MARIJ	1.2	0.5	1.9	0.34
CODEINE	3.4	1.8	5	0.8
OXYCONTIN ETC	1.4	0.9	2	0.28
VICODIN ETC	2	1.2	2.7	0.37
VALIUM ETC	0.4	0.2	0.7	0.12
XANAX ETC	1.2	0.4	1.9	0.37
DXM, ETC.	1.2	0.3	2.1	0.44

Lifetime Use of Selected Substances - Grade 9 Female				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	19.2	15.4	23	1.91
CIGARETTES	18.1	14.5	21.6	1.79
SMOKELESS TOBACCO	5.5	4.1	6.9	0.71
ALCOHOL	52.7	46.5	58.9	3.1
BEER	37.4	31.7	43	2.81
WINE COOLER	36.8	30.4	43.3	3.23
WINE	34.8	30.4	39.3	2.22
LIQUOR	36.2	30.5	42	2.88
ANY ILLICIT DRUG	21.1	16.9	25.3	2.09
INHALANTS	14.5	11.5	17.5	1.5
MARIJUANA	20.5	16.5	24.4	1.99
COCAINE OR CRACK	1.2	0.5	1.8	0.33
MARIJUANA ONLY	13.2	10.7	15.8	1.26
COCAINE	1.1	0.5	1.7	0.3
CRACK	0.3	0	0.5	0.12
HALLUCINOGEN	1.6	0.8	2.3	0.37
ROHYPNOL	1.1	0.6	1.6	0.27
STEROIDS	0.8	0.3	1.3	0.25
ECSTASY	2.2	1.5	2.9	0.37
HEROIN	0.3	0	0.6	0.14
METHAMPHETAMINE	0.3	0.1	0.6	0.12
SYNTHETIC MARIJ	5.8	4.3	7.4	0.78
CODEINE	10.4	7.9	12.8	1.22

Past Month Use of Selected Substances - Grade 9 Female				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	5.4	3.5	7.4	0.97
CIGARETTES	4.8	3.1	6.5	0.86
SMOKELESS TOBACCO	1.7	0.9	2.4	0.37
ALCOHOL	21.4	17.6	25.2	1.9
BEER	12.1	9.7	14.5	1.22
WINE COOLER	13.7	10.6	16.9	1.57
WINE	10.3	7.9	12.7	1.19
LIQUOR	13.8	11	16.6	1.41
ANY ILLICIT DRUG	8.9	6.9	11	1.03
INHALANTS	4.3	2.6	6	0.85
MARIJUANA	8.2	6.5	9.8	0.84
COCAINE OR CRACK	0.7	0.2	1.1	0.23
MARIJUANA ONLY	4.1	3.1	5.2	0.54
COCAINE	0.6	0.2	1	0.21
CRACK	0.2	0	0.4	0.12
HALLUCINOGEN	0.7	0.3	1	0.19
ROHYPNOL	0.6	0.3	1	0.18
STEROIDS	0.1	0	0.2	0.06
ECSTASY	0.8	0.3	1.3	0.24
HEROIN	0.1	0	0.1	0.04
METHAMPHETAMINE	0.1	0	0.2	0.06
SYNTHETIC MARIJ	2.1	1.5	2.8	0.33
CODEINE	5.2	3.6	6.8	0.81

OXYCONTIN ETC	1.7	1.1	2.3	0.3
VICODIN ETC	3.7	2.4	5.1	0.68
VALIUM ETC	0.7	0.2	1.2	0.25
XANAX ETC	3.3	2.3	4.3	0.5
DXM, ETC.	3.2	2.1	4.4	0.57

OXYCONTIN ETC	0.7	0.4	1.1	0.17
VICODIN ETC	1.9	1.1	2.7	0.39
VALIUM ETC	0.5	0.2	0.9	0.19
XANAX ETC	1.9	1.2	2.5	0.33
DXM, ETC.	1.5	0.7	2.4	0.43

Lifetime Use of Selected Substances - Grade 9 Male				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	22.8	16.9	28.6	2.92
CIGARETTES	19.4	13.8	25	2.81
SMOKELESS TOBACCO	11	7.7	14.3	1.64
ALCOHOL	47.6	39.5	55.8	4.09
BEER	37.8	30.2	45.5	3.83
WINE COOLER	25.3	19.9	30.8	2.75
WINE	29	24.4	33.7	2.34
LIQUOR	30.8	23.8	37.9	3.52
ANY ILLICIT DRUG	23.3	17.8	28.9	2.79
INHALANTS	11.3	9.6	13	0.85
MARIJUANA	22.5	16.8	28.1	2.83
COCAINE OR CRACK	1.8	1.3	2.3	0.25
MARIJUANA ONLY	12.9	9.4	16.4	1.75
COCAINE	1.7	1.2	2.2	0.25
CRACK	0.2	0.1	0.4	0.09
HALLUCINOGEN	2.9	1.7	4.2	0.64
ROHYPNOL	1.1	0.3	1.8	0.36
STEROIDS	0.8	0.4	1.3	0.24
ECSTASY	2.6	1.7	3.5	0.45
HEROIN	0.2	0	0.4	0.11
METHAMPHETAMINE	0.5	0.2	0.8	0.15
SYNTHETIC MARIJ	6.9	4.9	9	1.04
CODEINE	10.3	7.3	13.3	1.5
OXYCONTIN ETC	2.1	1.3	2.9	0.39
VICODIN ETC	3.7	2.8	4.7	0.47
VALIUM ETC	1	0.5	1.5	0.25
XANAX ETC	2.3	1.7	3	0.34
DXM, ETC.	2.9	1.7	4.1	0.61

Past Month Use of Selected Substances - Grade 9 Male				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	7.9	5.7	10	1.08
CIGARETTES	5	3.3	6.8	0.88
SMOKELESS TOBACCO	4.5	3.2	5.8	0.67
ALCOHOL	18.4	13.5	23.3	2.47
BEER	11.6	8	15.3	1.82
WINE COOLER	8.4	5.7	11.1	1.34
WINE	7.4	5.3	9.6	1.06
LIQUOR	10.8	7.5	14	1.62
ANY ILLICIT DRUG	9	6.7	11.4	1.18
INHALANTS	3.6	2.7	4.5	0.44
MARIJUANA	7.9	5.6	10.2	1.14
COCAINE OR CRACK	0.6	0.3	1	0.17
MARIJUANA ONLY	3.8	2.5	5.2	0.68
COCAINE	0.6	0.2	0.9	0.17
CRACK	0.1	0	0.2	0.06
HALLUCINOGEN	0.8	0.3	1.2	0.24
ROHYPNOL	0.2	0	0.5	0.12
STEROIDS	0.2	0	0.4	0.1
ECSTASY	0.7	0.3	1.1	0.18
HEROIN	0	0	0.1	0.03
METHAMPHETAMINE	0.1	0	0.3	0.06
SYNTHETIC MARIJ	2	1	2.9	0.5
CODEINE	5.1	3.4	6.7	0.81
OXYCONTIN ETC	1.4	0.8	2	0.29
VICODIN ETC	1.8	1.3	2.3	0.24
VALIUM ETC	0.5	0.2	0.8	0.14
XANAX ETC	1.1	0.8	1.5	0.18
DXM, ETC.	1.2	0.5	1.8	0.32

Lifetime Use of Selected Substances - Grade 9 Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	19	14	24	2.21
CIGARETTES	18.1	13.5	22.7	2.03
SMOKELESS TOBACCO	5.7	3	8.3	1.18
ALCOHOL	43.2	34.3	52.1	3.95
BEER	33.5	25.4	41.6	3.57
WINE COOLER	24.8	18.7	30.9	2.7
WINE	26.3	20.9	31.7	2.41
LIQUOR	27.6	21.5	33.6	2.65
ANY ILLICIT DRUG	25	20.1	30	2.19
INHALANTS	10.8	5.9	15.7	2.16
MARIJUANA	24.3	19.2	29.3	2.23
COCAINE OR CRACK	2.7	1	4.4	0.75
MARIJUANA ONLY	13.1	8.3	18	2.14
COCAINE	2.5	1.1	3.9	0.62
CRACK	0.4	0	1	0.27
HALLUCINOGEN	2.2	0.4	4	0.8
ROHYPNOL	1.9	0.7	3.1	0.53
STEROIDS	0.5	0	0.9	0.2

Past Month Use of Selected Substances - Grade 9 Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	6.7	4.2	9.2	1.12
CIGARETTES	6.2	3.7	8.6	1.08
SMOKELESS TOBACCO	2.2	1	3.4	0.53
ALCOHOL	18.5	14.2	22.7	1.88
BEER	12.5	8.4	16.5	1.79
WINE COOLER	9.7	7.4	12	1.03
WINE	9.4	7.2	11.7	1.01
LIQUOR	10.8	8.8	12.8	0.89
ANY ILLICIT DRUG	11.1	7.1	15.1	1.77
INHALANTS	4.5	2.7	6.3	0.8
MARIJUANA	9.9	6.7	13.1	1.4
COCAINE OR CRACK	1.8	0.3	3.4	0.7
MARIJUANA ONLY	3.1	2	4.3	0.52
COCAINE	1.7	0.2	3.1	0.63
CRACK	0.4	0	1	0.27
HALLUCINOGEN	0.9	0.2	1.6	0.3
ROHYPNOL	1.1	0	2.2	0.5
STEROIDS	0.3	0	0.7	0.17

ECSTASY	3.8	1.3	6.2	1.08
HEROIN	0.2	0	0.5	0.16
METHAMPHETAMINE	0.5	0	1.2	0.29
SYNTHETIC MARIJ	8.7	5.1	12.2	1.57
CODEINE	9	6.1	11.9	1.28
OXYCONTIN ETC	1	0.2	1.7	0.33
VICODIN ETC	2	0.4	3.6	0.72
VALIUM ETC	0.4	0	0.9	0.2
XANAX ETC	3.7	1.5	5.9	0.98
DXM, ETC.	3.6	1.4	5.7	0.96

ECSTASY	1.3	0	3	0.74
HEROIN	0.2	0	0.5	0.16
METHAMPHETAMINE	0.2	0	0.5	0.14
SYNTHETIC MARIJ	4.1	1.5	6.7	1.17
CODEINE	3.6	1.6	5.6	0.87
OXYCONTIN ETC	0.6	0	1.3	0.29
VICODIN ETC	1.2	0.2	2.1	0.41
VALIUM ETC	0.2	0	0.6	0.15
XANAX ETC	1.8	0.7	2.9	0.49
DXM, ETC.	1.9	0	3.8	0.86

Lifetime Use of Selected Substances - Grade 9 Non-Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	21.3	15.9	26.7	2.7
CIGARETTES	18.8	13.6	24	2.59
SMOKELESS TOBACCO	8.7	6.3	11.1	1.2
ALCOHOL	51.2	42.9	59.5	4.16
BEER	38.2	30.6	45.8	3.8
WINE COOLER	31.9	24.6	39.1	3.6
WINE	32.8	27.7	37.9	2.55
LIQUOR	34.4	27.1	41.8	3.66
ANY ILLICIT DRUG	21.9	16.5	27.3	2.68
INHALANTS	13.3	10.9	15.7	1.2
MARIJUANA	21.1	15.8	26.4	2.65
COCAINE OR CRACK	1.3	1	1.7	0.18
MARIJUANA ONLY	13.1	9.8	16.4	1.64
COCAINE	1.2	0.9	1.6	0.18
CRACK	0.2	0.1	0.4	0.07
HALLUCINOGEN	2.3	1.3	3.2	0.47
ROHYPNOL	1	0.4	1.5	0.29
STEROIDS	0.9	0.4	1.4	0.24
ECSTASY	2.3	1.6	3	0.35
HEROIN	0.3	0.1	0.5	0.09
METHAMPHETAMINE	0.5	0.2	0.7	0.11
SYNTHETIC MARIJ	6.1	4.3	7.9	0.89
CODEINE	10.6	7.6	13.6	1.5
OXYCONTIN ETC	2	1.4	2.7	0.31
VICODIN ETC	4	2.8	5.2	0.61
VALIUM ETC	0.9	0.5	1.4	0.23
XANAX ETC	2.7	1.9	3.5	0.4
DXM, ETC.	3	1.9	4.2	0.57

Past Month Use of Selected Substances - Grade 9 Non-Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	6.7	4.5	8.8	1.08
CIGARETTES	4.7	2.9	6.5	0.9
SMOKELESS TOBACCO	3.2	2.3	4.2	0.49
ALCOHOL	20.1	15.2	25	2.45
BEER	11.8	8.6	15	1.6
WINE COOLER	11.2	7.9	14.5	1.66
WINE	8.8	6.6	11.1	1.13
LIQUOR	12.5	9.2	15.8	1.64
ANY ILLICIT DRUG	8.7	6.5	11	1.13
INHALANTS	3.9	2.7	5.2	0.62
MARIJUANA	7.8	5.7	9.8	1.02
COCAINE OR CRACK	0.5	0.2	0.7	0.11
MARIJUANA ONLY	4.2	3	5.3	0.58
COCAINE	0.4	0.2	0.6	0.11
CRACK	0.1	0	0.3	0.06
HALLUCINOGEN	0.7	0.3	1.1	0.19
ROHYPNOL	0.3	0.1	0.5	0.11
STEROIDS	0.1	0	0.3	0.06
ECSTASY	0.6	0.3	1	0.16
HEROIN	0	0	0.1	0.02
METHAMPHETAMINE	0.1	0	0.2	0.05
SYNTHETIC MARIJ	1.8	1.1	2.4	0.33
CODEINE	5.4	3.7	7.1	0.85
OXYCONTIN ETC	1.1	0.7	1.6	0.23
VICODIN ETC	2	1.4	2.6	0.31
VALIUM ETC	0.5	0.3	0.8	0.15
XANAX ETC	1.4	0.9	1.9	0.24
DXM, ETC.	1.3	0.6	1.9	0.32

Lifetime Use of Selected Substances - Grade 10				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	25.8	21.3	30.3	2.25
CIGARETTES	22.9	18.4	27.4	2.25
SMOKELESS TOBACCO	10.6	8.5	12.7	1.06
ALCOHOL	58.5	54	63	2.25
BEER	45.1	40.2	49.9	2.43
WINE COOLER	39.5	35.4	43.7	2.07
WINE	37.1	33.6	40.5	1.73
LIQUOR	43.8	39.6	47.9	2.08
ANY ILLICIT DRUG	29.8	25.9	33.7	1.93
INHALANTS	12.1	9.5	14.7	1.31
MARIJUANA	29	25.2	32.7	1.86
COCAINE OR CRACK	2.3	1.5	3.1	0.4
MARIJUANA ONLY	18	15.5	20.4	1.22

Past Month Use of Selected Substances - Grade 10				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	9.6	7.5	11.6	1.03
CIGARETTES	7.1	5.2	9.1	0.98
SMOKELESS TOBACCO	4.5	3.5	5.6	0.54
ALCOHOL	25.5	22.4	28.6	1.55
BEER	15.9	12.4	19.3	1.73
WINE COOLER	15.4	13	17.7	1.19
WINE	10.9	9.4	12.5	0.78
LIQUOR	16.6	14.9	18.3	0.85
ANY ILLICIT DRUG	11.5	9.7	13.4	0.94
INHALANTS	3.5	2.5	4.4	0.47
MARIJUANA	10.7	9.1	12.4	0.83
COCAINE OR CRACK	0.4	0.2	0.6	0.11
MARIJUANA ONLY	5.2	4.3	6.1	0.45

COCAINE	2.1	1.3	2.8	0.36
CRACK	0.6	0.3	1	0.18
HALLUCINOGEN	2.9	2.4	3.5	0.28
ROHYPNOL	1	0.5	1.5	0.26
STEROIDS	1.1	0.8	1.4	0.17
ECSTASY	3.1	2	4.1	0.53
HEROIN	0.4	0.2	0.6	0.1
METHAMPHETAMINE	0.7	0.3	1.1	0.2
SYNTHETIC MARIJ	8.4	6.2	10.5	1.06
CODEINE	13.1	11.3	14.9	0.91
OXYCONTIN ETC	3.3	2.6	3.9	0.32
VICODIN ETC	6.6	4.4	8.7	1.08
VALIUM ETC	1.2	0.8	1.6	0.19
XANAX ETC	3.7	2.4	4.9	0.61
DXM, ETC.	3.9	2.6	5.2	0.66

COCAINE	0.4	0.2	0.6	0.1
CRACK	0.1	0	0.2	0.05
HALLUCINOGEN	0.8	0.6	1	0.11
ROHYPNOL	0.5	0.1	0.9	0.18
STEROIDS	0.5	0.3	0.8	0.12
ECSTASY	0.7	0.3	1.2	0.23
HEROIN	0.1	0	0.3	0.06
METHAMPHETAMINE	0.3	0.1	0.5	0.09
SYNTHETIC MARIJ	2.2	1.6	2.7	0.29
CODEINE	5.9	4.8	7	0.54
OXYCONTIN ETC	1.9	1.5	2.3	0.2
VICODIN ETC	3.2	2.2	4.1	0.47
VALIUM ETC	0.7	0.4	1	0.16
XANAX ETC	1.9	1.2	2.6	0.35
DXM, ETC.	1.7	1	2.4	0.34

Lifetime Use of Selected Substances - Grade 10 African American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	15	8.6	21.4	3.19
CIGARETTES	13.3	7.7	18.9	2.76
SMOKELESS TOBACCO	4.9	1.5	8.3	1.68
ALCOHOL	56.2	50	62.4	3.07
BEER	34	30	38.1	1.99
WINE COOLER	39	33.2	44.8	2.88
WINE	35.7	31.5	39.8	2.05
LIQUOR	41.6	36.1	47.1	2.72
ANY ILLICIT DRUG	28.3	19.4	37.1	4.39
INHALANTS	9.9	6.8	12.9	1.5
MARIJUANA	28.1	19.4	36.7	4.29
COCAINE OR CRACK	1	0	2.4	0.66
MARIJUANA ONLY	21.4	16.5	26.4	2.45
COCAINE	1	0	2.4	0.66
CRACK	0.2	0	0.6	0.2
HALLUCINOGEN	1.8	0	3.6	0.89
ROHYPNOL	0.2	0	0.5	0.14
STEROIDS	2.2	0.9	3.6	0.67
ECSTASY	2.1	0.4	3.8	0.86
HEROIN	0.1	0	0.2	0.06
METHAMPHETAMINE	0.1	0	0.3	0.09
SYNTHETIC MARIJ	4.7	1.3	8.1	1.69
CODEINE	16.5	13.4	19.6	1.54
OXYCONTIN ETC	2.1	0.2	4	0.92
VICODIN ETC	3.5	0.4	6.5	1.5
VALIUM ETC	0.3	0	0.6	0.17
XANAX ETC	2	0.2	3.7	0.86
DXM, ETC.	2.3	0.6	4.1	0.87

Past Month Use of Selected Substances- Grade 10 African American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	4.6	1.3	7.9	1.64
CIGARETTES	3.3	0.9	5.8	1.22
SMOKELESS TOBACCO	2.4	0.3	4.5	1.04
ALCOHOL	20.1	13.3	27	3.41
BEER	7.5	2.6	12.4	2.43
WINE COOLER	11.8	4.7	19	3.55
WINE	9.5	4.8	14.1	2.31
LIQUOR	15.3	11.1	19.5	2.08
ANY ILLICIT DRUG	10.2	6.3	14.1	1.94
INHALANTS	2.5	0.7	4.3	0.91
MARIJUANA	9.4	5.7	13.1	1.84
COCAINE OR CRACK	0.1	0	0.2	0.06
MARIJUANA ONLY	6.1	3.5	8.6	1.25
COCAINE	0.1	0	0.2	0.06
CRACK	0	*	*	*
HALLUCINOGEN	0.6	0	1.2	0.31
ROHYPNOL	0	*	*	*
STEROIDS	0.7	0	1.8	0.53
ECSTASY	1	0	2.3	0.64
HEROIN	0	*	*	*
METHAMPHETAMINE	0	*	*	*
SYNTHETIC MARIJ	1.1	0.2	2.1	0.47
CODEINE	6.5	3.5	9.5	1.5
OXYCONTIN ETC	1.2	0	2.5	0.67
VICODIN ETC	1.1	0	2.2	0.58
VALIUM ETC	0.2	0	0.5	0.13
XANAX ETC	0.2	0	0.5	0.16
DXM, ETC.	0.6	0	1.3	0.36

Lifetime Use of Selected Substances - Grade 10 Mexican American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	26.8	23.5	30.1	1.65
CIGARETTES	24.1	20.5	27.6	1.77
SMOKELESS TOBACCO	10.9	8.6	13.2	1.15
ALCOHOL	60.1	56.2	64.1	1.95
BEER	47.9	43.7	52.1	2.1
WINE COOLER	39.7	36.4	43.1	1.65
WINE	35.4	32.1	38.6	1.63
LIQUOR	44.1	40.6	47.7	1.77

Past Month Use of Selected Substances - Grade 10 Mexican American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	10	7.9	12	1.03
CIGARETTES	7.4	5.5	9.3	0.94
SMOKELESS TOBACCO	4.8	3.2	6.3	0.76
ALCOHOL	27.8	25.2	30.4	1.29
BEER	17.7	15	20.3	1.32
WINE COOLER	16.5	14.6	18.3	0.92
WINE	11.4	10	12.7	0.68
LIQUOR	16.7	14.7	18.7	1

ANY ILLICIT DRUG	33	30.1	35.8	1.42
INHALANTS	11.8	9.6	13.9	1.08
MARIJUANA	32	29.1	34.8	1.43
COCAINE OR CRACK	3.5	2.5	4.5	0.48
MARIJUANA ONLY	19.5	16.3	22.8	1.62
COCAINE	3	2.1	4	0.48
CRACK	1	0.4	1.6	0.31
HALLUCINOGEN	2.7	2.1	3.4	0.33
ROHYPNOL	1.6	0.8	2.5	0.41
STEROIDS	0.9	0.4	1.4	0.24
ECSTASY	3.6	2.2	5.1	0.72
HEROIN	0.5	0.2	0.8	0.15
METHAMPHETAMINE	1	0.3	1.6	0.32
SYNTHETIC MARIJ	9.2	7.2	11.2	1
CODEINE	12.4	9.8	14.9	1.26
OXYCONTIN ETC	2.5	1.5	3.5	0.5
VICODIN ETC	5.1	3.3	6.9	0.89
VALIUM ETC	1.6	0.9	2.3	0.34
XANAX ETC	3.4	2.2	4.6	0.6
DXM, ETC.	3.8	2.4	5.3	0.72

ANY ILLICIT DRUG	13.6	11.6	15.6	1
INHALANTS	3.7	2.7	4.6	0.48
MARIJUANA	12.5	10.5	14.5	1
COCAINE OR CRACK	0.7	0.4	1	0.17
MARIJUANA ONLY	6.3	4.7	7.8	0.78
COCAINE	0.6	0.3	0.9	0.15
CRACK	0.2	0	0.4	0.08
HALLUCINOGEN	0.8	0.4	1.2	0.18
ROHYPNOL	0.8	0.3	1.4	0.26
STEROIDS	0.6	0.3	0.8	0.15
ECSTASY	1	0.3	1.8	0.37
HEROIN	0.2	0	0.4	0.1
METHAMPHETAMINE	0.4	0.1	0.7	0.15
SYNTHETIC MARIJ	3	1.9	4	0.51
CODEINE	5.3	3.9	6.6	0.67
OXYCONTIN ETC	1.1	0.5	1.7	0.3
VICODIN ETC	2.4	1.4	3.4	0.49
VALIUM ETC	0.8	0.2	1.4	0.29
XANAX ETC	1.9	1.1	2.6	0.39
DXM, ETC.	2	1.2	2.8	0.4

Lifetime Use of Selected Substances - Grade 10 White				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	29.9	24.1	35.8	2.92
CIGARETTES	26.2	20.2	32.2	2.98
SMOKELESS TOBACCO	13.2	10.4	16.1	1.4
ALCOHOL	58.3	52	64.6	3.14
BEER	47.4	41.1	53.7	3.14
WINE COOLER	41.8	35.7	47.9	3.05
WINE	40.1	35.2	45	2.45
LIQUOR	45.3	39.2	51.3	3
ANY ILLICIT DRUG	26.8	21.9	31.7	2.46
INHALANTS	13.1	8.9	17.3	2.09
MARIJUANA	26.1	21.4	30.9	2.38
COCAINE OR CRACK	1.4	0.5	2.4	0.46
MARIJUANA ONLY	15.3	12	18.7	1.66
COCAINE	1.4	0.5	2.3	0.46
CRACK	0.4	0	0.8	0.22
HALLUCINOGEN	3.7	2.7	4.7	0.51
ROHYPNOL	0.6	0.1	1.2	0.27
STEROIDS	1	0.3	1.7	0.34
ECSTASY	2.5	1.3	3.8	0.62
HEROIN	0.3	0	0.6	0.14
METHAMPHETAMINE	0.5	0.1	0.8	0.19
SYNTHETIC MARIJ	8.6	5.7	11.5	1.45
CODEINE	12.7	9.7	15.8	1.53
OXYCONTIN ETC	4.5	3.2	5.9	0.68
VICODIN ETC	9.5	6.1	12.9	1.71
VALIUM ETC	1.2	0.6	1.9	0.3
XANAX ETC	4.7	3.1	6.3	0.79
DXM, ETC.	3.9	2.6	5.1	0.64

Past Month Use of Selected Substances - Grade 10 White				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	12.1	8.9	15.2	1.57
CIGARETTES	8.8	5.7	11.8	1.52
SMOKELESS TOBACCO	5.8	4.1	7.4	0.83
ALCOHOL	25.4	21.6	29.2	1.89
BEER	17.7	12.9	22.5	2.41
WINE COOLER	16.7	13.1	20.2	1.77
WINE	11.4	9.1	13.7	1.15
LIQUOR	17.7	14.1	21.2	1.78
ANY ILLICIT DRUG	9.5	6.9	12	1.28
INHALANTS	4	2.1	5.8	0.93
MARIJUANA	9	6.5	11.6	1.27
COCAINE OR CRACK	0.2	0	0.4	0.11
MARIJUANA ONLY	3.7	2.3	5.1	0.7
COCAINE	0.2	0	0.4	0.11
CRACK	0	0	0.1	0.02
HALLUCINOGEN	1	0.6	1.4	0.21
ROHYPNOL	0.4	0	0.9	0.25
STEROIDS	0.5	0	0.9	0.2
ECSTASY	0.4	0	0.7	0.17
HEROIN	0.2	0	0.4	0.12
METHAMPHETAMINE	0.2	0	0.5	0.14
SYNTHETIC MARIJ	1.8	0.9	2.7	0.45
CODEINE	6.4	4.7	8.2	0.87
OXYCONTIN ETC	3	1.7	4.3	0.65
VICODIN ETC	4.9	3.3	6.5	0.82
VALIUM ETC	0.8	0.2	1.4	0.3
XANAX ETC	2.7	1.8	3.6	0.46
DXM, ETC.	1.6	0.8	2.4	0.41

Lifetime Use of Selected Substances - Grade 10 Female				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	25.7	21.4	30	2.16
CIGARETTES	24.1	19.5	28.7	2.3
SMOKELESS TOBACCO	7.1	5.6	8.6	0.75

Past Month Use of Selected Substances - Grade 10 Female				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	9.3	7.5	11.1	0.91
CIGARETTES	7.9	6.1	9.7	0.9
SMOKELESS TOBACCO	3	2.3	3.8	0.37

ALCOHOL	62.1	56.5	67.8	2.84
BEER	45.9	41	50.8	2.45
WINE COOLER	46.5	42.1	51	2.2
WINE	41.8	38.1	45.5	1.86
LIQUOR	47.6	43.2	51.9	2.17
ANY ILLICIT DRUG	30.8	27.1	34.6	1.86
INHALANTS	13.5	11	16.1	1.29
MARIJUANA	30.2	26.6	33.8	1.8
COCAINE OR CRACK	1.7	0.9	2.6	0.43
MARIJUANA ONLY	19.7	16.9	22.4	1.37
COCAINE	1.4	0.7	2.1	0.37
CRACK	0.6	0.1	1	0.22
HALLUCINOGEN	2.5	1.6	3.5	0.47
ROHYPNOL	1	0.4	1.7	0.33
STEROIDS	0.6	0.2	1	0.19
ECSTASY	3.1	2	4.2	0.56
HEROIN	0.3	0.1	0.6	0.14
METHAMPHETAMINE	0.8	0.3	1.3	0.25
SYNTHETIC MARIJ	8.2	6.3	10.2	0.96
CODEINE	11.7	8.9	14.4	1.37
OXYCONTIN ETC	3.5	2.6	4.4	0.46
VICODIN ETC	7.3	5	9.6	1.13
VALIUM ETC	1.3	0.8	1.7	0.22
XANAX ETC	4	2.7	5.4	0.65
DXM, ETC.	4.3	2.9	5.8	0.72

ALCOHOL	28.6	25.8	31.4	1.38
BEER	16.6	13.1	20.2	1.77
WINE COOLER	18.9	16.7	21.1	1.08
WINE	12.8	11.7	13.8	0.53
LIQUOR	19.6	17.6	21.5	0.97
ANY ILLICIT DRUG	12	10.3	13.8	0.89
INHALANTS	4	2.7	5.3	0.66
MARIJUANA	11.6	9.9	13.3	0.84
COCAINE OR CRACK	0.3	0.1	0.5	0.1
MARIJUANA ONLY	5.8	4.3	7.4	0.76
COCAINE	0.2	0	0.3	0.08
CRACK	0.1	0	0.2	0.06
HALLUCINOGEN	0.6	0.3	0.9	0.15
ROHYPNOL	0.4	0	0.7	0.16
STEROIDS	0.2	0	0.5	0.11
ECSTASY	0.5	0.2	0.9	0.19
HEROIN	0.1	0	0.2	0.05
METHAMPHETAMINE	0.4	0.1	0.7	0.16
SYNTHETIC MARIJ	2.2	1.4	3	0.4
CODEINE	5.5	4.1	7	0.72
OXYCONTIN ETC	2.3	1.5	3.2	0.43
VICODIN ETC	3.9	2.5	5.2	0.69
VALIUM ETC	0.9	0.4	1.3	0.21
XANAX ETC	2.5	1.5	3.5	0.51
DXM, ETC.	1.8	1.1	2.6	0.37

Lifetime Use of Selected Substances - Grade 10 Male				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	25.7	20.3	31	2.68
CIGARETTES	21.6	16.6	26.6	2.51
SMOKELESS TOBACCO	14.3	11	17.7	1.69
ALCOHOL	54.7	50.5	58.9	2.1
BEER	44.2	38.4	49.9	2.86
WINE COOLER	32.2	27.8	36.7	2.21
WINE	32.2	28.5	36	1.87
LIQUOR	39.7	34.6	44.8	2.54
ANY ILLICIT DRUG	28.7	24.2	33.2	2.26
INHALANTS	10.7	7.5	14	1.63
MARIJUANA	27.7	23.3	32.1	2.2
COCAINE OR CRACK	2.8	1.5	4	0.63
MARIJUANA ONLY	16.3	13.6	19	1.35
COCAINE	2.6	1.4	3.8	0.6
CRACK	0.7	0.2	1.2	0.25
HALLUCINOGEN	3.4	2.6	4.1	0.38
ROHYPNOL	1	0.3	1.7	0.36
STEROIDS	1.7	1	2.3	0.33
ECSTASY	3.1	1.9	4.3	0.61
HEROIN	0.5	0.2	0.7	0.14
METHAMPHETAMINE	0.6	0.2	1.1	0.24
SYNTHETIC MARIJ	8.6	5.7	11.4	1.42
CODEINE	14.6	13	16.2	0.81
OXYCONTIN ETC	3.1	2.3	3.9	0.39
VICODIN ETC	5.9	3.2	8.5	1.32
VALIUM ETC	1.2	0.6	1.8	0.29
XANAX ETC	3.3	1.8	4.8	0.73
DXM, ETC.	3.5	2	4.9	0.71

Past Month Use of Selected Substances - Grade 10 Male				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	9.7	7	12.4	1.35
CIGARETTES	6.3	3.9	8.7	1.21
SMOKELESS TOBACCO	6	4.2	7.7	0.87
ALCOHOL	22.2	18.4	26.1	1.93
BEER	15.1	11.4	18.7	1.84
WINE COOLER	11.8	8.8	14.8	1.51
WINE	9	6.5	11.5	1.25
LIQUOR	13.6	11.5	15.7	1.05
ANY ILLICIT DRUG	11.1	8.6	13.5	1.22
INHALANTS	3	1.9	4.1	0.54
MARIJUANA	9.9	7.9	12	1.02
COCAINE OR CRACK	0.6	0.2	1	0.19
MARIJUANA ONLY	4.6	3.6	5.6	0.48
COCAINE	0.6	0.2	0.9	0.18
CRACK	0.1	0	0.3	0.06
HALLUCINOGEN	1	0.5	1.5	0.23
ROHYPNOL	0.7	0.1	1.3	0.32
STEROIDS	0.8	0.4	1.3	0.22
ECSTASY	1	0.4	1.6	0.31
HEROIN	0.2	0	0.4	0.11
METHAMPHETAMINE	0.2	0	0.3	0.09
SYNTHETIC MARIJ	2.2	1.2	3.1	0.47
CODEINE	6.3	4.9	7.6	0.69
OXYCONTIN ETC	1.4	0.8	2.1	0.31
VICODIN ETC	2.5	1.5	3.5	0.51
VALIUM ETC	0.6	0.2	1	0.22
XANAX ETC	1.3	0.7	1.9	0.3
DXM, ETC.	1.6	0.8	2.4	0.41

Lifetime Use of Selected Substances - Grade 10 Border

Past Month Use of Selected Substances - Grade 10 Border

Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	26.1	22.1	30	1.7
CIGARETTES	25.4	21.8	29	1.57
SMOKELESS TOBACCO	7.2	5.2	9.2	0.87
ALCOHOL	57.8	52.8	62.8	2.16
BEER	45.5	40.3	50.7	2.27
WINE COOLER	38.3	32.8	43.7	2.36
WINE	33.2	28.2	38.3	2.18
LIQUOR	41.4	36.5	46.2	2.1
ANY ILLICIT DRUG	32.5	27.2	37.9	2.32
INHALANTS	14.8	12.7	16.9	0.91
MARIJUANA	31.8	26.7	36.9	2.21
COCAINE OR CRACK	4.3	2.1	6.5	0.94
MARIJUANA ONLY	15.9	12.2	19.6	1.59
COCAINE	3.7	2	5.4	0.74
CRACK	1.1	0.2	2	0.4
HALLUCINOGEN	3.8	2.6	5.1	0.55
ROHYPNOL	2	0.6	3.5	0.63
STEROIDS	1.1	0.5	1.7	0.27
ECSTASY	6.6	3.2	10	1.49
HEROIN	0.9	0.4	1.4	0.24
METHAMPHETAMINE	1.3	0	2.9	0.7
SYNTHETIC MARIJ	12	8.7	15.4	1.47
CODEINE	12.2	9.3	15.2	1.28
OXYCONTIN ETC	2.1	0.8	3.4	0.56
VICODIN ETC	4.6	0.9	8.2	1.58
VALIUM ETC	1.6	0.8	2.4	0.34
XANAX ETC	3.5	2.3	4.6	0.5
DXM, ETC.	5	0.9	9	1.76

Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	9.2	6.2	12.2	1.31
CIGARETTES	8.1	5.7	10.5	1.04
SMOKELESS TOBACCO	2.8	1.4	4.3	0.63
ALCOHOL	26.3	22	30.7	1.9
BEER	16.3	12.9	19.7	1.48
WINE COOLER	15.6	12.8	18.3	1.17
WINE	11.1	8.3	13.8	1.2
LIQUOR	14.4	10.9	18	1.54
ANY ILLICIT DRUG	14.1	9.3	19	2.08
INHALANTS	5	3.9	6.1	0.47
MARIJUANA	12.7	8.1	17.3	1.98
COCAINE OR CRACK	1.1	0.5	1.6	0.24
MARIJUANA ONLY	4.8	3.7	5.8	0.44
COCAINE	0.7	0.3	1.1	0.16
CRACK	0.4	0	0.8	0.17
HALLUCINOGEN	1.3	0.7	1.9	0.26
ROHYPNOL	1	0.3	1.6	0.27
STEROIDS	0.7	0.3	1.2	0.19
ECSTASY	2.6	0.8	4.3	0.76
HEROIN	0.1	0	0.5	0.14
METHAMPHETAMINE	0.6	0	1.3	0.29
SYNTHETIC MARIJ	3.1	1.6	4.5	0.63
CODEINE	4.2	3.2	5.3	0.45
OXYCONTIN ETC	0.7	0.1	1.2	0.24
VICODIN ETC	2.2	0.3	4.1	0.82
VALIUM ETC	0.8	0.3	1.3	0.22
XANAX ETC	1.1	0.3	2	0.38
DXM, ETC.	2.2	0.7	3.7	0.67

Lifetime Use of Selected Substances - Grade 10 Non-Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	25.7	20.5	30.9	2.59
CIGARETTES	22.5	17.4	27.7	2.55
SMOKELESS TOBACCO	11.2	8.6	13.7	1.26
ALCOHOL	58.6	53.3	63.8	2.61
BEER	45	39.4	50.6	2.78
WINE COOLER	39.7	34.9	44.5	2.4
WINE	37.7	33.6	41.7	2.01
LIQUOR	44.2	39.3	49	2.43
ANY ILLICIT DRUG	29.4	25	33.7	2.16
INHALANTS	11.6	8.7	14.6	1.45
MARIJUANA	28.5	24.3	32.7	2.08
COCAINE OR CRACK	2	1.3	2.8	0.37
MARIJUANA ONLY	18.3	15.5	21.1	1.4
COCAINE	1.8	1.1	2.5	0.36
CRACK	0.6	0.2	1	0.19
HALLUCINOGEN	2.8	2.2	3.4	0.31
ROHYPNOL	0.8	0.3	1.4	0.27
STEROIDS	1.1	0.7	1.5	0.19
ECSTASY	2.5	1.6	3.4	0.43
HEROIN	0.3	0.1	0.5	0.1
METHAMPHETAMINE	0.6	0.3	1	0.18
SYNTHETIC MARIJ	7.8	5.5	10	1.11
CODEINE	13.2	11.1	15.3	1.05
OXYCONTIN ETC	3.5	2.7	4.2	0.37
VICODIN ETC	6.9	4.4	9.4	1.24

Past Month Use of Selected Substances - Grade 10 Non-Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	9.6	7.2	12	1.19
CIGARETTES	7	4.8	9.2	1.1
SMOKELESS TOBACCO	4.8	3.5	6.1	0.64
ALCOHOL	25.4	21.8	28.9	1.76
BEER	15.8	11.8	19.8	1.98
WINE COOLER	15.3	12.6	18.1	1.36
WINE	10.9	9.1	12.7	0.89
LIQUOR	17	15	19	0.99
ANY ILLICIT DRUG	11.1	9.2	13.1	0.98
INHALANTS	3.2	2.2	4.2	0.51
MARIJUANA	10.4	8.7	12.2	0.87
COCAINE OR CRACK	0.3	0.1	0.5	0.1
MARIJUANA ONLY	5.3	4.2	6.4	0.53
COCAINE	0.3	0.1	0.5	0.1
CRACK	0.1	0	0.1	0.04
HALLUCINOGEN	0.7	0.5	0.9	0.11
ROHYPNOL	0.4	0	0.8	0.2
STEROIDS	0.5	0.2	0.7	0.13
ECSTASY	0.5	0.2	0.7	0.14
HEROIN	0.1	0	0.3	0.06
METHAMPHETAMINE	0.2	0.1	0.4	0.09
SYNTHETIC MARIJ	2	1.4	2.6	0.3
CODEINE	6.2	4.9	7.5	0.64
OXYCONTIN ETC	2.1	1.6	2.5	0.22
VICODIN ETC	3.3	2.3	4.4	0.54

VALIUM ETC	1.2	0.7	1.6	0.21
XANAX ETC	3.7	2.3	5.1	0.71
DXM, ETC.	3.7	2.3	5.1	0.69

VALIUM ETC	0.7	0.3	1.1	0.18
XANAX ETC	2	1.2	2.9	0.41
DXM, ETC.	1.6	0.9	2.4	0.37

Lifetime Use of Selected Substances - Grade 11				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	32.2	28.7	35.6	1.74
CIGARETTES	28.7	25	32.4	1.85
SMOKELESS TOBACCO	13.1	11.1	15.2	1.02
ALCOHOL	64	58.6	69.5	2.72
BEER	50.5	45	56	2.76
WINE COOLER	44.8	39.5	50.1	2.65
WINE	40	36.4	43.6	1.8
LIQUOR	50.6	46.5	54.7	2.06
ANY ILLICIT DRUG	36.4	32.5	40.4	1.96
INHALANTS	10	8.6	11.5	0.71
MARIJUANA	35.5	31.1	39.8	2.17
COCAINE OR CRACK	3.5	2.5	4.6	0.53
MARIJUANA ONLY	20.9	18.7	23.1	1.1
COCAINE	3.3	2.3	4.3	0.5
CRACK	0.6	0.3	0.9	0.15
HALLUCINOGEN	4.6	3.6	5.6	0.5
ROHYPNOL	1	0.5	1.5	0.25
STEROIDS	1.1	0.7	1.5	0.2
ECSTASY	4.5	3.3	5.8	0.65
HEROIN	0.2	0.1	0.4	0.09
METHAMPHETAMINE	0.9	0.4	1.4	0.26
SYNTHETIC MARIJ	9.3	6.9	11.7	1.19
CODEINE	14.7	12.6	16.8	1.05
OXYCONTIN ETC	4.7	3.6	5.8	0.55
VICODIN ETC	9.6	7.7	11.6	0.98
VALIUM ETC	1.8	1.4	2.3	0.24
XANAX ETC	6.3	5.1	7.4	0.59
DXM, ETC.	4.1	3.3	5	0.43

Past Month Use of Selected Substances - Grade 11				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	13.1	11.5	14.7	0.79
CIGARETTES	10.6	8.9	12.3	0.85
SMOKELESS TOBACCO	5.4	4.6	6.2	0.4
ALCOHOL	30.1	25.6	34.5	2.22
BEER	21.1	17	25.2	2.03
WINE COOLER	18.7	15.2	22.3	1.77
WINE	12.7	10.9	14.5	0.92
LIQUOR	21.3	18.3	24.4	1.53
ANY ILLICIT DRUG	16.3	13.8	18.7	1.24
INHALANTS	3.2	2.5	3.9	0.35
MARIJUANA	15.2	12.6	17.8	1.32
COCAINE OR CRACK	1.3	0.7	1.8	0.27
MARIJUANA ONLY	7.2	6	8.4	0.6
COCAINE	1.2	0.7	1.8	0.26
CRACK	0.1	0	0.2	0.05
HALLUCINOGEN	1.5	1	2.1	0.28
ROHYPNOL	0.4	0.1	0.6	0.13
STEROIDS	0.6	0.3	0.8	0.12
ECSTASY	1.6	0.5	2.7	0.54
HEROIN	0.1	0	0.1	0.03
METHAMPHETAMINE	0.3	0	0.6	0.15
SYNTHETIC MARIJ	1.9	1.1	2.6	0.37
CODEINE	7.2	5.7	8.6	0.73
OXYCONTIN ETC	2.5	1.7	3.2	0.37
VICODIN ETC	4.2	3.3	5	0.41
VALIUM ETC	0.9	0.5	1.3	0.18
XANAX ETC	3	2.2	3.8	0.41
DXM, ETC.	1.9	1.3	2.5	0.28

Lifetime Use of Selected Substances - Grade 11 African American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	22.1	19	25.2	1.55
CIGARETTES	18.7	14.4	23.1	2.14
SMOKELESS TOBACCO	7.1	4.9	9.2	1.06
ALCOHOL	54.9	44.7	65	5.03
BEER	31.5	23.4	39.6	4.02
WINE COOLER	44.9	33.4	56.4	5.7
WINE	36.5	28	45.1	4.24
LIQUOR	44.8	38.3	51.4	3.25
ANY ILLICIT DRUG	37.6	29.7	45.5	3.92
INHALANTS	6.9	3.8	10	1.53
MARIJUANA	36.7	28	45.4	4.32
COCAINE OR CRACK	0.6	0	1.2	0.32
MARIJUANA ONLY	27.5	22.5	32.4	2.46
COCAINE	0.5	0	1	0.27
CRACK	0.1	0	0.3	0.1
HALLUCINOGEN	2.4	0.8	3.9	0.76
ROHYPNOL	0.4	0	0.9	0.24
STEROIDS	0.8	0	2.1	0.61
ECSTASY	3.4	0.6	6.3	1.42
HEROIN	0.5	0	1.6	0.54

Past Month Use of Selected Substances - Grade 11 African American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	7.7	4.8	10.7	1.45
CIGARETTES	5.8	3.3	8.3	1.24
SMOKELESS TOBACCO	3.2	1.6	4.8	0.78
ALCOHOL	20.8	13.4	28.2	3.69
BEER	10.5	5.9	15.2	2.33
WINE COOLER	15.5	10	20.9	2.7
WINE	9.9	6.1	13.7	1.89
LIQUOR	14.3	10	18.5	2.09
ANY ILLICIT DRUG	16.5	11.2	21.8	2.61
INHALANTS	2.3	1.1	3.5	0.59
MARIJUANA	14.6	9.3	19.9	2.62
COCAINE OR CRACK	0.6	0	1.2	0.32
MARIJUANA ONLY	10.3	7.7	12.8	1.28
COCAINE	0.5	0	1	0.27
CRACK	0.1	0	0.3	0.1
HALLUCINOGEN	1	0	2	0.51
ROHYPNOL	0.2	0	0.6	0.19
STEROIDS	0.5	0	1.6	0.54
ECSTASY	1.3	0	2.7	0.7
HEROIN	0	*	*	*

METHAMPHETAMINE	0.6	0	1.8	0.56
SYNTHETIC MARIJ	6.3	2	10.6	2.15
CODEINE	16.2	11.4	21	2.38
OXYCONTIN ETC	5.4	1.7	9.1	1.83
VICODIN ETC	9.9	6.5	13.2	1.66
VALIUM ETC	1.7	0	3.6	0.93
XANAX ETC	5	2.7	7.3	1.15
DXM, ETC.	3.1	0.6	5.6	1.24

METHAMPHETAMINE	0	*	*	*
SYNTHETIC MARIJ	1.4	0	2.9	0.74
CODEINE	8.3	5.1	11.6	1.6
OXYCONTIN ETC	3.4	0.5	6.3	1.44
VICODIN ETC	3.6	1.5	5.6	1.02
VALIUM ETC	1.5	0	3.3	0.89
XANAX ETC	2.7	0.4	4.9	1.13
DXM, ETC.	0.4	0	1	0.3

Lifetime Use of Selected Substances - Grade 11 Mexican American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	34.3	30.4	38.2	1.93
CIGARETTES	31	27.3	34.7	1.86
SMOKELESS TOBACCO	13	10.2	15.8	1.38
ALCOHOL	66.5	62.2	70.9	2.16
BEER	55.7	51.5	60	2.13
WINE COOLER	44.3	39.7	48.9	2.29
WINE	37.8	33.9	41.8	1.98
LIQUOR	52	48.3	55.7	1.83
ANY ILLICIT DRUG	40.3	36.8	43.8	1.74
INHALANTS	11.7	9.6	13.8	1.05
MARIJUANA	39	35	43	1.98
COCAINE OR CRACK	5.2	3.8	6.7	0.72
MARIJUANA ONLY	21.7	18.8	24.6	1.45
COCAINE	5.1	3.7	6.5	0.72
CRACK	0.7	0.2	1.1	0.21
HALLUCINOGEN	4.1	2.5	5.6	0.79
ROHYPNOL	1.6	0.8	2.4	0.41
STEROIDS	0.9	0.3	1.4	0.27
ECSTASY	5.3	3.7	6.8	0.79
HEROIN	0.1	0	0.2	0.05
METHAMPHETAMINE	0.8	0.2	1.5	0.32
SYNTHETIC MARIJ	10.9	8.5	13.3	1.2
CODEINE	16.4	13.7	19.1	1.36
OXYCONTIN ETC	4.3	3	5.7	0.67
VICODIN ETC	7.9	5.9	10	1.03
VALIUM ETC	1.9	1	2.9	0.5
XANAX ETC	7.4	5	9.9	1.23
DXM, ETC.	4.8	3.8	5.9	0.54

Past Month Use of Selected Substances -Grade 11 Mexican American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	13.5	11.7	15.3	0.88
CIGARETTES	11.6	9.8	13.3	0.85
SMOKELESS TOBACCO	5.3	3.9	6.7	0.72
ALCOHOL	33.3	29.4	37.2	1.95
BEER	24.9	21	28.8	1.94
WINE COOLER	20.5	17	24	1.72
WINE	14	11.3	16.6	1.32
LIQUOR	23.7	20.4	27	1.64
ANY ILLICIT DRUG	18.5	15.8	21.3	1.37
INHALANTS	4.4	2.9	5.9	0.76
MARIJUANA	17.2	14.1	20.2	1.53
COCAINE OR CRACK	1.9	1.1	2.8	0.42
MARIJUANA ONLY	8	6.4	9.5	0.77
COCAINE	1.9	1.1	2.8	0.42
CRACK	0.2	0	0.4	0.09
HALLUCINOGEN	1.5	0.2	2.8	0.65
ROHYPNOL	0.6	0	1.1	0.28
STEROIDS	0.3	0.1	0.6	0.13
ECSTASY	2	0.4	3.6	0.78
HEROIN	0	0	0.1	0.02
METHAMPHETAMINE	0.5	0	1.1	0.32
SYNTHETIC MARIJ	2.2	1.3	3.1	0.44
CODEINE	8.2	6.4	10	0.89
OXYCONTIN ETC	1.8	1	2.6	0.4
VICODIN ETC	3.1	1.9	4.2	0.59
VALIUM ETC	0.7	0.2	1.2	0.23
XANAX ETC	3.3	2.2	4.3	0.5
DXM, ETC.	2.5	1.6	3.3	0.43

Lifetime Use of Selected Substances - Grade 11 White				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	34.4	29.9	38.9	2.25
CIGARETTES	30.3	26.5	34.2	1.93
SMOKELESS TOBACCO	16.7	13.5	19.8	1.57
ALCOHOL	65.3	59.6	71.1	2.89
BEER	52.7	46.8	58.6	2.94
WINE COOLER	48.2	42	54.3	3.07
WINE	44.3	40.5	48.2	1.94
LIQUOR	52.4	47.2	57.6	2.59
ANY ILLICIT DRUG	31.6	27.7	35.4	1.91
INHALANTS	8.8	6.8	10.8	0.99
MARIJUANA	30.8	26.9	34.8	1.97
COCAINE OR CRACK	2.4	1	3.7	0.66
MARIJUANA ONLY	17.7	15.4	20.1	1.19
COCAINE	2.1	0.8	3.4	0.63
CRACK	0.5	0	1	0.24

Past Month Use of Selected Substances - Grade 11 White				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	15.2	12.4	18.1	1.43
CIGARETTES	11.6	9.4	13.8	1.09
SMOKELESS TOBACCO	6.8	4.8	8.9	1.04
ALCOHOL	29.8	26	33.6	1.91
BEER	21	17.5	24.6	1.77
WINE COOLER	19.1	15.2	22.9	1.92
WINE	11.9	10.6	13.2	0.65
LIQUOR	21.3	18.3	24.2	1.45
ANY ILLICIT DRUG	13.5	10.9	16	1.26
INHALANTS	1.8	0.8	2.8	0.5
MARIJUANA	13	10.5	15.5	1.25
COCAINE OR CRACK	0.6	0	1.3	0.36
MARIJUANA ONLY	5.2	4	6.3	0.58
COCAINE	0.6	0	1.3	0.36
CRACK	0	*	*	*

HALLUCINOGEN	5.7	4.5	6.8	0.56
ROHYPNOL	0.5	0.1	0.9	0.21
STEROIDS	1.5	0.6	2.4	0.46
ECSTASY	4.1	2.7	5.6	0.71
HEROIN	0.3	0	0.6	0.18
METHAMPHETAMINE	1	0.4	1.5	0.27
SYNTHETIC MARIJ	8.8	6	11.6	1.41
CODEINE	11.9	9.4	14.4	1.24
OXYCONTIN ETC	5.3	4.1	6.4	0.58
VICODIN ETC	12.3	10	14.7	1.18
VALIUM ETC	1.9	0.6	3.2	0.63
XANAX ETC	5.6	4.3	7	0.68
DXM, ETC.	3.6	2.2	5.1	0.73

HALLUCINOGEN	1.3	0.5	2.2	0.41
ROHYPNOL	0.2	0	0.5	0.14
STEROIDS	0.8	0.2	1.5	0.32
ECSTASY	1.2	0.4	2	0.39
HEROIN	0.1	0	0.3	0.09
METHAMPHETAMINE	0.2	0	0.5	0.14
SYNTHETIC MARIJ	1.5	0.5	2.4	0.46
CODEINE	5.6	3.3	7.9	1.13
OXYCONTIN ETC	3.3	2.1	4.5	0.59
VICODIN ETC	6	4.9	7.2	0.57
VALIUM ETC	1	0.4	1.6	0.31
XANAX ETC	2.7	1.6	3.8	0.55
DXM, ETC.	1.7	0.9	2.6	0.44

Lifetime Use of Selected Substances - Grade 11 Female				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	30.3	26.8	33.8	1.76
CIGARETTES	28.4	24.6	32.1	1.88
SMOKELESS TOBACCO	9.2	7	11.4	1.12
ALCOHOL	68.9	64	73.8	2.45
BEER	49	43.2	54.7	2.87
WINE COOLER	52.9	48.3	57.5	2.3
WINE	44.5	40.4	48.5	2.03
LIQUOR	53.5	50.2	56.8	1.64
ANY ILLICIT DRUG	34.7	30.4	39.1	2.17
INHALANTS	11	9.6	12.4	0.7
MARIJUANA	33.8	29.2	38.3	2.27
COCAINE OR CRACK	2.1	1.4	2.8	0.36
MARIJUANA ONLY	22.1	19.4	24.8	1.33
COCAINE	1.8	1.1	2.5	0.36
CRACK	0.5	0.2	0.8	0.16
HALLUCINOGEN	2.5	2	3.1	0.28
ROHYPNOL	1.1	0.5	1.7	0.3
STEROIDS	0.6	0.2	1.1	0.22
ECSTASY	4.6	2.5	6.6	1.01
HEROIN	0.1	0	0.2	0.07
METHAMPHETAMINE	0.8	0	1.8	0.47
SYNTHETIC MARIJ	7.4	5.4	9.4	1
CODEINE	11.7	9.6	13.9	1.07
OXYCONTIN ETC	4.6	3.1	6.1	0.75
VICODIN ETC	10.3	7.6	12.9	1.34
VALIUM ETC	1.7	1.1	2.3	0.28
XANAX ETC	6.2	4.5	8	0.87
DXM, ETC.	3.6	2.4	4.7	0.55

Past Month Use of Selected Substances - Grade 11 Female				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	11.1	9.4	12.8	0.86
CIGARETTES	9.7	7.9	11.5	0.91
SMOKELESS TOBACCO	3.5	2.7	4.3	0.41
ALCOHOL	31.5	27.3	35.7	2.12
BEER	20.1	16.5	23.8	1.81
WINE COOLER	23	19.3	26.6	1.82
WINE	13.9	12	15.9	0.97
LIQUOR	22	19.2	24.9	1.41
ANY ILLICIT DRUG	16.1	12.5	19.6	1.77
INHALANTS	2.8	2	3.5	0.38
MARIJUANA	15.3	11.9	18.8	1.74
COCAINE OR CRACK	0.7	0.3	1.1	0.22
MARIJUANA ONLY	8.4	6.3	10.4	1.03
COCAINE	0.7	0.2	1.1	0.21
CRACK	0.1	0	0.2	0.07
HALLUCINOGEN	1	0.5	1.5	0.24
ROHYPNOL	0.5	0.1	1	0.24
STEROIDS	0.5	0	0.9	0.21
ECSTASY	1.9	0.1	3.6	0.87
HEROIN	0	*	*	*
METHAMPHETAMINE	0.4	0	0.9	0.25
SYNTHETIC MARIJ	1.3	0.6	1.9	0.31
CODEINE	5.5	3.8	7.2	0.86
OXYCONTIN ETC	2.4	1.3	3.6	0.58
VICODIN ETC	4.1	2.9	5.3	0.59
VALIUM ETC	0.8	0.4	1.2	0.22
XANAX ETC	2.6	1.9	3.3	0.34
DXM, ETC.	2.1	1.1	3	0.47

Lifetime Use of Selected Substances - Grade 11 Male				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	33.9	29.4	38.4	2.23
CIGARETTES	28.9	24.1	33.7	2.4
SMOKELESS TOBACCO	17.1	14.3	19.8	1.38
ALCOHOL	59.4	53.2	65.5	3.06
BEER	52.4	46.7	58	2.83
WINE COOLER	36.8	30.9	42.7	2.96
WINE	35.6	31.8	39.4	1.9
LIQUOR	47.8	42.6	53	2.6
ANY ILLICIT DRUG	38.1	33.6	42.7	2.28
INHALANTS	9.2	7.4	11	0.9

Past Month Use of Selected Substances - Grade 11 Male				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	15.2	13.1	17.2	1.01
CIGARETTES	11.7	9.7	13.7	1
SMOKELESS TOBACCO	7.3	6	8.6	0.66
ALCOHOL	28.9	23.9	34	2.52
BEER	22.2	17.2	27.3	2.52
WINE COOLER	14.6	11	18.3	1.84
WINE	11.5	9.2	13.9	1.17
LIQUOR	20.7	16.8	24.6	1.96
ANY ILLICIT DRUG	16.4	13.8	19	1.28
INHALANTS	3.6	2.5	4.7	0.55

MARIJUANA	37.2	32.1	42.3	2.55
COCAINE OR CRACK	5.1	3.2	6.9	0.91
MARIJUANA ONLY	19.7	17	22.4	1.35
COCAINE	4.9	3.1	6.7	0.88
CRACK	0.7	0.2	1.2	0.27
HALLUCINOGEN	6.7	4.9	8.5	0.91
ROHYPNOL	0.8	0.3	1.4	0.26
STEROIDS	1.6	0.7	2.4	0.43
ECSTASY	4.5	3.5	5.6	0.53
HEROIN	0.4	0.1	0.7	0.15
METHAMPHETAMINE	0.9	0.4	1.5	0.27
SYNTHETIC MARIJ	11.3	8.3	14.4	1.52
CODEINE	17.7	15	20.3	1.32
OXYCONTIN ETC	4.8	3.5	6.1	0.65
VICODIN ETC	8.9	7.1	10.7	0.89
VALIUM ETC	2	1.3	2.7	0.35
XANAX ETC	6.3	4.8	7.7	0.72
DXM, ETC.	4.5	3.4	5.7	0.59

MARIJUANA	15	12.1	17.9	1.46
COCAINE OR CRACK	1.9	0.9	2.9	0.5
MARIJUANA ONLY	5.9	4.3	7.5	0.79
COCAINE	1.8	0.8	2.8	0.5
CRACK	0.1	0	0.2	0.06
HALLUCINOGEN	2.1	1	3.2	0.54
ROHYPNOL	0.2	0	0.4	0.1
STEROIDS	0.7	0.3	1.1	0.22
ECSTASY	1.3	0.5	2.1	0.4
HEROIN	0.1	0	0.3	0.07
METHAMPHETAMINE	0.3	0	0.6	0.18
SYNTHETIC MARIJ	2.4	1.4	3.5	0.54
CODEINE	8.9	7.3	10.5	0.79
OXYCONTIN ETC	2.4	1.6	3.3	0.43
VICODIN ETC	4.2	3.1	5.3	0.55
VALIUM ETC	1	0.5	1.6	0.29
XANAX ETC	3.3	1.9	4.7	0.69
DXM, ETC.	1.7	1.2	2.3	0.26

Lifetime Use of Selected Substances - Grade 11 Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	31.4	27.8	35	1.59
CIGARETTES	30.3	27.1	33.6	1.44
SMOKELESS TOBACCO	8.9	6.1	11.7	1.23
ALCOHOL	64	57.8	70.2	2.7
BEER	51	44.5	57.6	2.84
WINE COOLER	42.7	37.6	47.9	2.24
WINE	37.4	31.8	43	2.37
LIQUOR	49	44.5	53.4	1.92
ANY ILLICIT DRUG	38.3	32.9	43.7	2.34
INHALANTS	12.7	9.8	15.5	1.26
MARIJUANA	37	31.9	42.1	2.21
COCAINE OR CRACK	4.7	3.9	5.6	0.35
MARIJUANA ONLY	19.4	16.2	22.7	1.43
COCAINE	4.7	4	5.3	0.29
CRACK	0.8	0.3	1.3	0.22
HALLUCINOGEN	2.7	1.4	4	0.55
ROHYPNOL	2.6	0.5	4.7	0.9
STEROIDS	1.5	0.4	2.6	0.47
ECSTASY	6.4	3	9.9	1.49
HEROIN	0.6	0	1.4	0.35
METHAMPHETAMINE	1.1	0	2.3	0.5
SYNTHETIC MARIJ	12.4	10	14.8	1.04
CODEINE	14.5	10.7	18.4	1.71
OXYCONTIN ETC	3.6	1.8	5.3	0.75
VICODIN ETC	6.2	1.9	10.6	1.88
VALIUM ETC	2	0.1	3.9	0.83
XANAX ETC	7.5	4.4	10.7	1.37
DXM, ETC.	3.3	0.8	5.9	1.12

Past Month Use of Selected Substances - Grade 11 Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	13.3	11.4	15.2	0.84
CIGARETTES	12	10.3	13.7	0.76
SMOKELESS TOBACCO	3.6	2	5.1	0.67
ALCOHOL	33.5	27.7	39.2	2.48
BEER	23.7	19	28.5	2.05
WINE COOLER	20	15.7	24.2	1.84
WINE	13.1	10.6	15.5	1.03
LIQUOR	22.2	19.3	25.1	1.25
ANY ILLICIT DRUG	16.2	12.6	19.8	1.56
INHALANTS	3.9	2	5.8	0.84
MARIJUANA	14.2	11.6	16.8	1.14
COCAINE OR CRACK	1.1	0.3	1.8	0.33
MARIJUANA ONLY	5.6	4.8	6.4	0.36
COCAINE	1	0.3	1.6	0.28
CRACK	0.3	0	0.8	0.2
HALLUCINOGEN	0.8	0	1.6	0.34
ROHYPNOL	0.5	0	1.2	0.29
STEROIDS	0.8	0	1.6	0.35
ECSTASY	2	0	4	0.89
HEROIN	0	*	*	*
METHAMPHETAMINE	0.1	0	0.2	0.07
SYNTHETIC MARIJ	3.1	2.1	4	0.41
CODEINE	7	5	9	0.9
OXYCONTIN ETC	2	1.2	2.7	0.33
VICODIN ETC	3.1	0.8	5.3	0.97
VALIUM ETC	1.3	0.1	2.4	0.51
XANAX ETC	5.2	3.1	7.3	0.92
DXM, ETC.	1.9	0.4	3.3	0.65

Lifetime Use of Selected Substances - Grade 11 Non-Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	32.3	28.2	36.4	2.02
CIGARETTES	28.4	24.2	32.7	2.12
SMOKELESS TOBACCO	13.8	11.5	16.2	1.18
ALCOHOL	64	57.7	70.4	3.15
BEER	50.4	44.1	56.8	3.17

Past Month Use of Selected Substances - Grade 11 Non-Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	13.1	11.2	14.9	0.91
CIGARETTES	10.4	8.5	12.4	0.96
SMOKELESS TOBACCO	5.7	4.8	6.6	0.45
ALCOHOL	29.5	24.5	34.5	2.48
BEER	20.7	16.1	25.3	2.28

WINE COOLER	45.1	38.9	51.3	3.1
WINE	40.4	36.3	44.6	2.06
LIQUOR	50.8	46	55.7	2.41
ANY ILLICIT DRUG	36.1	31.7	40.6	2.22
INHALANTS	9.6	8.1	11.1	0.74
MARIJUANA	35.2	30.3	40.2	2.48
COCAINE OR CRACK	3.4	2.1	4.6	0.61
MARIJUANA ONLY	21.2	18.6	23.7	1.27
COCAINE	3.1	2	4.2	0.56
CRACK	0.6	0.2	0.9	0.17
HALLUCINOGEN	4.9	3.8	5.9	0.52
ROHYPNOL	0.7	0.3	1.2	0.22
STEROIDS	1	0.6	1.5	0.21
ECSTASY	4.2	2.9	5.6	0.69
HEROIN	0.2	0	0.3	0.08
METHAMPHETAMINE	0.8	0.3	1.4	0.29
SYNTHETIC MARIJ	8.8	6.2	11.4	1.3
CODEINE	14.8	12.4	17.2	1.2
OXYCONTIN ETC	4.9	3.6	6.1	0.63
VICODIN ETC	10.2	8	12.3	1.07
VALIUM ETC	1.8	1.3	2.3	0.25
XANAX ETC	6.1	4.8	7.4	0.65
DXM, ETC.	4.2	3.3	5.2	0.47

WINE COOLER	18.5	14.5	22.6	2.02
WINE	12.6	10.5	14.7	1.06
LIQUOR	21.2	17.7	24.7	1.76
ANY ILLICIT DRUG	16.3	13.4	19.1	1.42
INHALANTS	3	2.3	3.8	0.38
MARIJUANA	15.4	12.3	18.4	1.53
COCAINE OR CRACK	1.3	0.7	1.9	0.31
MARIJUANA ONLY	7.4	6	8.8	0.7
COCAINE	1.3	0.7	1.9	0.3
CRACK	0.1	0	0.1	0.03
HALLUCINOGEN	1.7	1.1	2.3	0.29
ROHYPNOL	0.4	0.1	0.7	0.15
STEROIDS	0.5	0.3	0.8	0.13
ECSTASY	1.5	0.3	2.8	0.61
HEROIN	0.1	0	0.2	0.04
METHAMPHETAMINE	0.4	0	0.7	0.17
SYNTHETIC MARIJ	1.7	0.9	2.5	0.39
CODEINE	7.2	5.5	8.9	0.84
OXYCONTIN ETC	2.5	1.7	3.4	0.44
VICODIN ETC	4.3	3.4	5.3	0.46
VALIUM ETC	0.9	0.5	1.2	0.19
XANAX ETC	2.6	1.9	3.4	0.37
DXM, ETC.	1.9	1.3	2.5	0.31

Lifetime Use of Selected Substances - Grade 12				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	34	28.9	39.2	2.6
CIGARETTES	30.3	25.1	35.5	2.61
SMOKELESS TOBACCO	15.2	12.7	17.7	1.26
ALCOHOL	64.3	58.9	69.8	2.72
BEER	51.4	45.6	57.2	2.91
WINE COOLER	45.5	40.9	50.1	2.32
WINE	42.9	40	45.8	1.44
LIQUOR	53.3	49.2	57.3	2.05
ANY ILLICIT DRUG	38.7	35.2	42.3	1.76
INHALANTS	9	7.2	10.8	0.91
MARIJUANA	38.1	34.8	41.4	1.66
COCAINE OR CRACK	3.9	2.7	5	0.57
MARIJUANA ONLY	23.5	21.7	25.3	0.89
COCAINE	3.7	2.6	4.9	0.56
CRACK	0.3	0	0.5	0.11
HALLUCINOGEN	4.7	3.9	5.6	0.42
ROHYPNOL	1	0.5	1.4	0.23
STEROIDS	1.2	0.8	1.7	0.22
ECSTASY	4.8	3.6	6.1	0.63
HEROIN	0.5	0.1	0.8	0.18
METHAMPHETAMINE	0.9	0.3	1.5	0.31
SYNTHETIC MARIJ	9.3	7.3	11.4	1.04
CODEINE	14.8	12.2	17.3	1.27
OXYCONTIN ETC	4.2	3	5.4	0.59
VICODIN ETC	9.7	7.8	11.6	0.93
VALIUM ETC	1.8	1.2	2.4	0.3
XANAX ETC	5.2	4	6.4	0.62
DXM, ETC.	3.3	2.2	4.4	0.55

Past Month Use of Selected Substances - Grade 12				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	15.8	12.9	18.7	1.46
CIGARETTES	13	10.3	15.8	1.39
SMOKELESS TOBACCO	7.4	5.4	9.5	1.01
ALCOHOL	32.7	28.5	36.9	2.08
BEER	23.7	19.5	27.9	2.1
WINE COOLER	19.3	17.1	21.6	1.11
WINE	13.9	12.3	15.4	0.78
LIQUOR	23.9	21.2	26.6	1.35
ANY ILLICIT DRUG	15.9	13.2	18.7	1.37
INHALANTS	2.6	1.5	3.6	0.51
MARIJUANA	15.2	12.7	17.7	1.24
COCAINE OR CRACK	1.2	0.7	1.7	0.26
MARIJUANA ONLY	7.1	5.7	8.6	0.73
COCAINE	1.1	0.7	1.6	0.23
CRACK	0.1	0	0.2	0.06
HALLUCINOGEN	1.3	0.9	1.8	0.24
ROHYPNOL	0.3	0.1	0.5	0.11
STEROIDS	0.5	0.2	0.8	0.17
ECSTASY	1.1	0.5	1.7	0.3
HEROIN	0.3	0	0.5	0.13
METHAMPHETAMINE	0.3	0.1	0.4	0.09
SYNTHETIC MARIJ	2.1	1	3.1	0.53
CODEINE	6.6	5.5	7.7	0.55
OXYCONTIN ETC	2.1	1.3	2.8	0.37
VICODIN ETC	5	3.8	6.2	0.6
VALIUM ETC	1.1	0.8	1.4	0.17
XANAX ETC	2.9	2.3	3.4	0.28
DXM, ETC.	1.6	0.9	2.3	0.34

Lifetime Use of Selected Substances - Grade 12 African American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	

Past Month Use of Selected Substances - Grade 12 African American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	

TOBACCO	20.9	12	29.7	4.38
CIGARETTES	20.2	11.5	28.8	4.25
SMOKELESS TOBACCO	6	0.7	11.4	2.62
ALCOHOL	57.4	44.8	70	6.19
BEER	37.2	32	42.4	2.55
WINE COOLER	42.5	33.7	51.3	4.33
WINE	39.2	31.5	46.9	3.79
LIQUOR	47.9	37.7	58	4.99
ANY ILLICIT DRUG	42.2	37.2	47.2	2.47
INHALANTS	6	3.1	8.8	1.42
MARIJUANA	42.2	37.2	47.2	2.46
COCAINE OR CRACK	1.9	0.7	3	0.55
MARIJUANA ONLY	31.2	26.3	36.2	2.44
COCAINE	1.9	0.7	3	0.56
CRACK	0	*	*	*
HALLUCINOGEN	3.5	1.1	5.9	1.19
ROHYPNOL	0.3	0	1	0.35
STEROIDS	1.9	0.7	3	0.58
ECSTASY	2.6	1.3	3.9	0.65
HEROIN	0.3	0	1	0.35
METHAMPHETAMINE	2.4	0	6.7	2.11
SYNTHETIC MARIJ	7.1	2.3	11.8	2.35
CODEINE	21	13.2	28.8	3.85
OXYCONTIN ETC	4.5	2.8	6.1	0.82
VICODIN ETC	10.7	7.2	14.1	1.7
VALIUM ETC	1.1	0	2.4	0.6
XANAX ETC	5.3	2.7	7.9	1.28
DXM, ETC.	3.2	0.2	6.3	1.52

TOBACCO	7.6	2.8	12.5	2.4
CIGARETTES	7.4	2.7	12.2	2.34
SMOKELESS TOBACCO	3.7	0	8.1	2.17
ALCOHOL	28	20.3	35.8	3.82
BEER	16.3	12.2	20.4	2.02
WINE COOLER	18.9	11.2	26.7	3.82
WINE	11.6	7	16.2	2.26
LIQUOR	22	15.1	29	3.41
ANY ILLICIT DRUG	16.5	10.1	22.9	3.17
INHALANTS	1.3	0	2.7	0.69
MARIJUANA	15.9	9.5	22.3	3.15
COCAINE OR CRACK	0.8	0	1.9	0.57
MARIJUANA ONLY	7.9	3.5	12.4	2.19
COCAINE	0.8	0	1.9	0.57
CRACK	0	*	*	*
HALLUCINOGEN	1.4	0	3	0.78
ROHYPNOL	0.3	0	1	0.35
STEROIDS	0.4	0	1.2	0.37
ECSTASY	0.6	0	1.6	0.48
HEROIN	0.3	0	1	0.35
METHAMPHETAMINE	0	*	*	*
SYNTHETIC MARIJ	3	0	7.4	2.14
CODEINE	5.3	2.9	7.7	1.19
OXYCONTIN ETC	1.9	1	2.9	0.46
VICODIN ETC	5.7	3.4	8	1.12
VALIUM ETC	0.7	0	1.4	0.37
XANAX ETC	2.7	1.7	3.6	0.48
DXM, ETC.	1	0	2.2	0.57

Lifetime Use of Selected Substances - Grade 12 Mexican American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	35.8	31.3	40.4	2.27
CIGARETTES	33.3	28.7	38	2.32
SMOKELESS TOBACCO	14.3	12.3	16.2	0.98
ALCOHOL	64.6	59.8	69.5	2.42
BEER	54.3	48.2	60.3	3.03
WINE COOLER	45	40.7	49.2	2.12
WINE	39.7	36.3	43.2	1.72
LIQUOR	52.7	48.6	56.9	2.07
ANY ILLICIT DRUG	41.3	37	45.5	2.13
INHALANTS	9.9	7.8	12	1.04
MARIJUANA	40.2	36.1	44.4	2.07
COCAINE OR CRACK	5.6	3.8	7.4	0.91
MARIJUANA ONLY	24	21.3	26.7	1.36
COCAINE	5.5	3.7	7.4	0.92
CRACK	0.2	0	0.5	0.15
HALLUCINOGEN	4.3	3.2	5.3	0.52
ROHYPNOL	1.3	0.7	2	0.32
STEROIDS	1	0.4	1.5	0.28
ECSTASY	6	3.5	8.5	1.25
HEROIN	0.6	0.1	1	0.21
METHAMPHETAMINE	0.7	0.2	1.2	0.24
SYNTHETIC MARIJ	9.7	7.6	11.7	1.02
CODEINE	15.1	11.9	18.3	1.59
OXYCONTIN ETC	3.5	1.9	5.2	0.81
VICODIN ETC	8.3	6.3	10.2	0.98
VALIUM ETC	1.6	0.8	2.3	0.37
XANAX ETC	4.8	3.1	6.6	0.87

Past Month Use of Selected Substances -Grade 12 Mexican American				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	15.8	13.3	18.3	1.24
CIGARETTES	13.7	11	16.5	1.37
SMOKELESS TOBACCO	6	4.5	7.4	0.73
ALCOHOL	34.1	29.7	38.5	2.19
BEER	26.1	22.1	30	1.97
WINE COOLER	21.4	18.3	24.4	1.52
WINE	14.2	12.3	16	0.92
LIQUOR	24	20.7	27.3	1.63
ANY ILLICIT DRUG	16.9	14.1	19.7	1.39
INHALANTS	3.4	2.3	4.6	0.58
MARIJUANA	16	13.4	18.7	1.32
COCAINE OR CRACK	1.6	1	2.3	0.33
MARIJUANA ONLY	8	6.3	9.6	0.83
COCAINE	1.6	1	2.3	0.33
CRACK	0.1	0	0.2	0.06
HALLUCINOGEN	1.3	0.7	2	0.33
ROHYPNOL	0.4	0	0.7	0.18
STEROIDS	0.4	0	0.8	0.22
ECSTASY	1.6	0.7	2.4	0.42
HEROIN	0.4	0	0.8	0.21
METHAMPHETAMINE	0.4	0	0.8	0.22
SYNTHETIC MARIJ	2.4	1.6	3.3	0.44
CODEINE	7.1	5.2	9	0.95
OXYCONTIN ETC	1.7	0.8	2.6	0.43
VICODIN ETC	4.5	3	6	0.75
VALIUM ETC	0.7	0.3	1.2	0.21
XANAX ETC	2.4	1.2	3.5	0.59

DXM, ETC.	3.1	1.9	4.3	0.6
-----------	-----	-----	-----	-----

DXM, ETC.	1.6	0.8	2.3	0.36
-----------	-----	-----	-----	------

Lifetime Use of Selected Substances - Grade 12 White				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	39.9	35.3	44.5	2.28
CIGARETTES	32.9	27.4	38.3	2.73
SMOKELESS TOBACCO	21.6	18.7	24.5	1.46
ALCOHOL	68.8	65.2	72.5	1.83
BEER	55.4	49.8	60.9	2.77
WINE COOLER	49.9	45.5	54.4	2.21
WINE	48.6	44.7	52.4	1.92
LIQUOR	58.5	55	62	1.76
ANY ILLICIT DRUG	36.6	31.8	41.3	2.35
INHALANTS	8.6	6.4	10.9	1.15
MARIJUANA	36.3	31.7	41	2.32
COCAINE OR CRACK	3.1	2	4.1	0.53
MARIJUANA ONLY	21.1	18.3	23.8	1.36
COCAINE	2.8	1.9	3.7	0.44
CRACK	0.4	0	0.8	0.2
HALLUCINOGEN	5.6	4.1	7.1	0.75
ROHYPNOL	0.8	0.2	1.3	0.26
STEROIDS	1.1	0.1	2.2	0.51
ECSTASY	4.6	3.3	5.9	0.65
HEROIN	0.4	0	1	0.3
METHAMPHETAMINE	0.7	0.3	1.1	0.21
SYNTHETIC MARIJ	10.1	7.6	12.6	1.26
CODEINE	12.2	9.8	14.7	1.23
OXYCONTIN ETC	5.2	3.8	6.6	0.69
VICODIN ETC	11.7	8.8	14.6	1.45
VALIUM ETC	2.5	1.5	3.5	0.49
XANAX ETC	6	4.2	7.7	0.88
DXM, ETC.	3.6	1.8	5.4	0.89

Past Month Use of Selected Substances - Grade 12 White				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	20.3	17.1	23.4	1.57
CIGARETTES	15	12.2	17.8	1.41
SMOKELESS TOBACCO	11.4	7.9	14.9	1.74
ALCOHOL	35	30.4	39.6	2.28
BEER	25.4	19.6	31.1	2.87
WINE COOLER	18.7	15.7	21.8	1.51
WINE	15.2	13.5	17	0.87
LIQUOR	26.6	22.3	30.8	2.12
ANY ILLICIT DRUG	14.1	10.8	17.4	1.65
INHALANTS	1.4	0.4	2.3	0.48
MARIJUANA	13.4	10.4	16.5	1.51
COCAINE OR CRACK	0.9	0.1	1.8	0.41
MARIJUANA ONLY	5.4	3.4	7.4	1
COCAINE	0.7	0.2	1.2	0.27
CRACK	0.3	0	0.6	0.16
HALLUCINOGEN	1.2	0.7	1.8	0.28
ROHYPNOL	0.3	0	0.6	0.15
STEROIDS	0.4	0	1.2	0.37
ECSTASY	0.8	0	1.7	0.43
HEROIN	0.1	0	0.3	0.11
METHAMPHETAMINE	0.3	0	0.6	0.18
SYNTHETIC MARIJ	1.3	0.4	2.2	0.46
CODEINE	6.7	5.3	8.1	0.68
OXYCONTIN ETC	2.8	1.8	3.8	0.5
VICODIN ETC	6.1	4.2	7.9	0.92
VALIUM ETC	1.8	1.1	2.5	0.37
XANAX ETC	4	3	5	0.5
DXM, ETC.	1.9	0.9	2.9	0.48

Lifetime Use of Selected Substances - Grade 12 Female				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	28.6	24.2	33.1	2.21
CIGARETTES	26.8	22.3	31.4	2.28
SMOKELESS TOBACCO	9.5	7.1	11.9	1.2
ALCOHOL	64.7	59.6	69.9	2.58
BEER	47.7	43.3	52.1	2.21
WINE COOLER	49.4	44.3	54.6	2.57
WINE	45.5	42.5	48.5	1.51
LIQUOR	52.6	48.8	56.4	1.9
ANY ILLICIT DRUG	34.7	31.6	37.8	1.54
INHALANTS	7.9	6.1	9.8	0.93
MARIJUANA	34.2	31.3	37	1.42
COCAINE OR CRACK	2.3	1.5	3.2	0.44
MARIJUANA ONLY	23.8	22.2	25.4	0.8
COCAINE	2.4	1.5	3.2	0.43
CRACK	0	0	0	0.02
HALLUCINOGEN	2.1	1.4	2.8	0.34
ROHYPNOL	0.9	0.4	1.4	0.25
STEROIDS	0.8	0.3	1.2	0.23
ECSTASY	3.8	2.5	5.2	0.67
HEROIN	0.1	0	0.2	0.05

Past Month Use of Selected Substances - Grade 12 Female				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	12.1	9.5	14.7	1.3
CIGARETTES	10.9	8.1	13.8	1.42
SMOKELESS TOBACCO	4.3	2.2	6.5	1.08
ALCOHOL	30.3	26.6	34	1.84
BEER	18.7	15.1	22.3	1.81
WINE COOLER	21.3	19	23.7	1.17
WINE	14.5	12.3	16.8	1.14
LIQUOR	21.4	18.8	24.1	1.31
ANY ILLICIT DRUG	11.8	9.3	14.2	1.22
INHALANTS	1.6	0.8	2.4	0.4
MARIJUANA	11.6	9.2	14	1.2
COCAINE OR CRACK	0.3	0.1	0.6	0.12
MARIJUANA ONLY	5.9	4.1	7.7	0.91
COCAINE	0.3	0.1	0.6	0.12
CRACK	0	0	0	0.02
HALLUCINOGEN	0.5	0.2	0.8	0.16
ROHYPNOL	0.2	0	0.3	0.08
STEROIDS	0.2	0	0.4	0.09
ECSTASY	0.5	0	1.1	0.28
HEROIN	0	0	0.1	0.03

METHAMPHETAMINE	0.9	0	2.1	0.59
SYNTHETIC MARIJ	7	5.1	8.9	0.95
CODEINE	11.1	8.3	13.9	1.42
OXYCONTIN ETC	3.2	2.1	4.2	0.51
VICODIN ETC	7.9	6.2	9.5	0.82
VALIUM ETC	1.4	0.9	1.9	0.25
XANAX ETC	4.4	2.8	6.1	0.82
DXM, ETC.	2.1	1.1	3.1	0.49

METHAMPHETAMINE	0.2	0	0.4	0.11
SYNTHETIC MARIJ	1.4	0.1	2.8	0.66
CODEINE	5.9	4.6	7.1	0.62
OXYCONTIN ETC	1.4	0.7	2	0.31
VICODIN ETC	4.2	3	5.5	0.61
VALIUM ETC	1	0.6	1.5	0.22
XANAX ETC	2.6	1.7	3.5	0.44
DXM, ETC.	1.2	0.4	2	0.4

Lifetime Use of Selected Substances - Grade 12 Male				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	39.5	33.2	45.7	3.11
CIGARETTES	33.8	27.5	40	3.11
SMOKELESS TOBACCO	21.1	16.9	25.4	2.11
ALCOHOL	63.8	57.5	70.1	3.14
BEER	54.7	47.1	62.3	3.8
WINE COOLER	41.5	36.2	46.7	2.63
WINE	40.4	35.8	45	2.28
LIQUOR	53.7	48.5	58.8	2.58
ANY ILLICIT DRUG	42.7	37.8	47.5	2.41
INHALANTS	9.9	7.3	12.4	1.27
MARIJUANA	41.9	37.2	46.6	2.36
COCAINE OR CRACK	5.4	3.7	7.2	0.87
MARIJUANA ONLY	23.1	20.4	25.8	1.36
COCAINE	5.2	3.5	6.9	0.85
CRACK	0.5	0.1	0.9	0.21
HALLUCINOGEN	7.3	5.8	8.8	0.76
ROHYPNOL	1	0.3	1.8	0.37
STERIODS	1.7	0.7	2.6	0.48
ECSTASY	5.7	3.9	7.5	0.89
HEROIN	0.9	0.2	1.6	0.35
METHAMPHETAMINE	0.9	0.4	1.5	0.28
SYNTHETIC MARIJ	11.8	9.1	14.6	1.37
CODEINE	18.6	15.8	21.5	1.42
OXYCONTIN ETC	5.3	3.8	6.8	0.75
VICODIN ETC	11.4	8.8	14.1	1.33
VALIUM ETC	2.3	1.4	3.2	0.45
XANAX ETC	6.1	4.6	7.5	0.73
DXM, ETC.	4.5	2.9	6.2	0.82

Past Month Use of Selected Substances - Grade 12 Male				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	19.4	15.5	23.4	1.98
CIGARETTES	15	11.5	18.6	1.78
SMOKELESS TOBACCO	10.6	7.7	13.6	1.47
ALCOHOL	35	29.8	40.2	2.6
BEER	28.5	23.3	33.7	2.6
WINE COOLER	17.3	14.2	20.3	1.51
WINE	13.1	10.7	15.5	1.18
LIQUOR	26.2	22.8	29.6	1.69
ANY ILLICIT DRUG	19.9	16	23.7	1.93
INHALANTS	3.4	1.9	4.9	0.75
MARIJUANA	18.6	15	22.2	1.8
COCAINE OR CRACK	2.1	1.2	3	0.45
MARIJUANA ONLY	8.3	6.3	10.2	0.97
COCAINE	1.9	1.1	2.7	0.4
CRACK	0.2	0	0.5	0.11
HALLUCINOGEN	2.2	1.3	3.1	0.44
ROHYPNOL	0.5	0.1	0.9	0.2
STERIODS	0.8	0.2	1.4	0.31
ECSTASY	1.7	0.8	2.6	0.46
HEROIN	0.5	0	1	0.25
METHAMPHETAMINE	0.3	0	0.7	0.18
SYNTHETIC MARIJ	2.7	1.6	3.9	0.57
CODEINE	7.5	5.9	9.1	0.8
OXYCONTIN ETC	2.8	1.8	3.8	0.49
VICODIN ETC	5.5	3.8	7.2	0.83
VALIUM ETC	1.2	0.5	1.8	0.32
XANAX ETC	3.2	2.3	4.2	0.47
DXM, ETC.	2.1	1.3	2.8	0.4

Lifetime Use of Selected Substances - Grade 12 Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	37.9	34.7	41.1	1.43
CIGARETTES	34.9	30.9	39	1.82
SMOKELESS TOBACCO	12.8	11.2	14.4	0.72
ALCOHOL	66.2	60.1	72.3	2.71
BEER	56.7	50.4	63.1	2.82
WINE COOLER	47.3	41.2	53.3	2.68
WINE	41.5	36.3	46.7	2.3
LIQUOR	51.2	42.6	59.8	3.81
ANY ILLICIT DRUG	42.5	35.3	49.8	3.25
INHALANTS	9.8	5.9	13.8	1.77
MARIJUANA	40.7	33	48.5	3.49
COCAINE OR CRACK	6.8	4.5	9.1	1.04
MARIJUANA ONLY	22.8	18.3	27.3	2.03

Past Month Use of Selected Substances - Grade 12 Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	15.4	13.6	17.3	0.84
CIGARETTES	14.6	12.7	16.4	0.84
SMOKELESS TOBACCO	4.5	2	7	1.12
ALCOHOL	33.5	29.6	37.3	1.69
BEER	24.7	20.9	28.4	1.65
WINE COOLER	22.2	17.3	27.2	2.18
WINE	13.9	11.8	15.9	0.9
LIQUOR	23.3	19.8	26.9	1.55
ANY ILLICIT DRUG	19.1	15	23.2	1.84
INHALANTS	2.3	1.2	3.4	0.5
MARIJUANA	18.2	14.4	22	1.71
COCAINE OR CRACK	2.4	1.6	3.2	0.35
MARIJUANA ONLY	7.7	5.2	10.2	1.11

COCAINE	6.8	4.5	9.2	1.04
CRACK	0.1	0	0.3	0.1
HALLUCINOGEN	3.7	1.8	5.5	0.82
ROHYPNOL	1.6	0.1	3.1	0.65
STEROIDS	0.6	0	1.4	0.37
ECSTASY	8.9	4	13.9	2.18
HEROIN	0.3	0	0.7	0.16
METHAMPHETAMINE	0	0	0.2	0.05
SYNTHETIC MARIJ	10.6	7.5	13.6	1.35
CODEINE	15.4	9	21.7	2.83
OXYCONTIN ETC	3.9	2.1	5.6	0.77
VICODIN ETC	7.8	3.8	11.8	1.75
VALIUM ETC	1.8	0.7	2.9	0.48
XANAX ETC	5.3	0.2	10.5	2.27
DXM, ETC.	1.6	0.1	3.2	0.69

COCAINE	2.4	1.6	3.2	0.35
CRACK	0.1	0	0.3	0.09
HALLUCINOGEN	1	0	2.1	0.47
ROHYPNOL	0.4	0	0.7	0.17
STEROIDS	0.4	0	1	0.28
ECSTASY	2.5	0.6	4.3	0.8
HEROIN	0.1	0	0.3	0.1
METHAMPHETAMINE	0	*	*	*
SYNTHETIC MARIJ	3.1	1.4	4.9	0.77
CODEINE	6.8	3.8	9.9	1.36
OXYCONTIN ETC	1.3	0	2.6	0.59
VICODIN ETC	3.5	1.3	5.7	0.98
VALIUM ETC	0.9	0.3	1.5	0.27
XANAX ETC	2.3	0	4.7	1.04
DXM, ETC.	0.8	0	1.7	0.41

Lifetime Use of Selected Substances - Grade 12 Non-Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	33.4	27.6	39.3	2.91
CIGARETTES	29.5	23.7	35.3	2.89
SMOKELESS TOBACCO	15.6	12.6	18.6	1.5
ALCOHOL	64	57.8	70.2	3.1
BEER	50.5	44	57	3.23
WINE COOLER	45.2	40	50.5	2.62
WINE	43.1	39.8	46.4	1.66
LIQUOR	53.6	48.9	58.3	2.33
ANY ILLICIT DRUG	38.1	34.3	41.9	1.9
INHALANTS	8.8	6.8	10.8	1
MARIJUANA	37.7	34.1	41.3	1.79
COCAINE OR CRACK	3.4	2.2	4.5	0.58
MARIJUANA ONLY	23.6	21.6	25.6	0.99
COCAINE	3.2	2.1	4.4	0.56
CRACK	0.3	0	0.5	0.12
HALLUCINOGEN	4.9	4	5.9	0.46
ROHYPNOL	0.9	0.4	1.4	0.24
STEROIDS	1.3	0.8	1.8	0.25
ECSTASY	4.2	3.1	5.2	0.52
HEROIN	0.5	0.1	0.9	0.21
METHAMPHETAMINE	1.1	0.4	1.8	0.35
SYNTHETIC MARIJ	9.1	6.8	11.5	1.17
CODEINE	14.7	11.8	17.5	1.4
OXYCONTIN ETC	4.3	2.9	5.6	0.68
VICODIN ETC	10	7.9	12.1	1.04
VALIUM ETC	1.8	1.2	2.5	0.34
XANAX ETC	5.2	3.9	6.4	0.63
DXM, ETC.	3.5	2.3	4.8	0.63

Past Month Use of Selected Substances - Grade 12 Non-Border				
Category	Prevalence	Boundaries		Standard Error
		Lower	Upper	
TOBACCO	15.9	12.5	19.3	1.7
CIGARETTES	12.8	9.6	15.9	1.58
SMOKELESS TOBACCO	7.9	5.5	10.3	1.19
ALCOHOL	32.6	27.8	37.4	2.4
BEER	23.6	18.7	28.4	2.42
WINE COOLER	18.9	16.5	21.2	1.17
WINE	13.9	12.1	15.7	0.9
LIQUOR	24	20.8	27.1	1.56
ANY ILLICIT DRUG	15.4	12.4	18.4	1.51
INHALANTS	2.6	1.4	3.8	0.59
MARIJUANA	14.7	12	17.4	1.37
COCAINE OR CRACK	1	0.5	1.5	0.27
MARIJUANA ONLY	7.1	5.4	8.7	0.83
COCAINE	0.9	0.4	1.4	0.23
CRACK	0.1	0	0.3	0.07
HALLUCINOGEN	1.4	0.8	1.9	0.27
ROHYPNOL	0.3	0.1	0.6	0.12
STEROIDS	0.5	0.1	0.9	0.19
ECSTASY	0.9	0.3	1.4	0.28
HEROIN	0.3	0	0.6	0.15
METHAMPHETAMINE	0.3	0.1	0.5	0.1
SYNTHETIC MARIJ	1.9	0.7	3.1	0.58
CODEINE	6.6	5.4	7.8	0.6
OXYCONTIN ETC	2.2	1.4	3	0.41
VICODIN ETC	5.2	3.9	6.6	0.67
VALIUM ETC	1.1	0.7	1.5	0.2
XANAX ETC	3	2.4	3.5	0.27
DXM, ETC.	1.7	1	2.5	0.39