

<https://helda.helsinki.fi>

Global Gene Expression Profiling of Body-Mass Index in Middle-Aged Danish Twins

Li, Weilong

2020-12-04

Li, W, Baumbach, J, Mohammadnejad, A, Lund, J, Larsen, M J, Hjelmborg, J V B, Mengel-From, J, Christensen, K, Christiansen, L & Tan, Q 2020, ' Global Gene Expression Profiling of Body-Mass Index in Middle-Aged Danish Twins ', Genetics and Genomics, vol. 1, no. 1, pp. 1-8 . <https://doi.org/10.31487/j.GG.2020.01.04>

<http://hdl.handle.net/10138/331717>

<https://doi.org/10.31487/j.GG.2020.01.04>

cc_by

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Available online at www.sciencerepository.org

Science Repository



Research Article

Global Gene Expression Profiling of Body-Mass Index in Middle-Aged Danish Twins

Weilong Li^{1,2}, Jan Baumbach^{3,4}, Afsaneh Mohammadnejad¹, Jesper Lund¹, Martin J. Larsen^{5,6}, Jacob v B. Hjelmborg¹, Jonas Mengel-From¹, Kaare Christensen^{1,6}, Lene Christiansen^{1,7} and Qihua Tan^{1,6*}

¹Epidemiology and Biostatistics, Department of Public Health, University of Southern Denmark, Denmark

²Population Research Unit, Faculty of Social Sciences, University of Helsinki, Helsinki, Finland

³Computational BioMedicine, Department of Mathematics and Computer Science, University of Southern Denmark, Denmark

⁴Chair of Experimental Bioinformatics, TUM School of Life Sciences, Technical University of Munich, Munich, Germany

⁵Department of Clinical Genetics, Odense University Hospital, Odense C, Denmark

⁶Unit of Human Genetics, Department of Clinical Research, University of Southern Denmark, Denmark

⁷Department of Clinical Immunology, Copenhagen University Hospital, Rigshospitalet, Copenhagen, Denmark

ARTICLE INFO

Article history:

Received: 27 October, 2020

Accepted: 6 November, 2020

Published: 4 December, 2020

Keywords:

BMI

gene expression

twins

generalized correlation

ABSTRACT

Objective: The body mass index (BMI) measured as weight in relation to height is an important monitor for obesity and diabetes, with individual variation under control by genetic and environmental factors. In transcriptome-wide association studies on BMI, the genetic contribution calls for controlling of genetic confounding that affects both BMI and gene expression. We performed a global gene expression profiling of BMI on peripheral blood cells using monozygotic twins for efficient handling of genetic make-ups.

Methods: We applied a generalized association method to genome-wide gene expression data on 229 pairs of monozygotic twins (age 56-80 years) for detecting diverse patterns of correlation between BMI and gene expression.

Results: We detected seven probes associated with BMI with $p < 1e-04$, among them two probes with $p < 1e-05$ ($p = 2.83e-06$ AAK1; $p = 7.83e-06$ LILRA3). In total, the analysis found 1579 probes with nominal $p < 0.05$. Biological pathway analysis of enriched pathways found 50 KEGG and 45 Reactome pathways ($FDR < 0.05$). The identified top functional pathways included immune function, JAK-STAT signalling, insulin signalling and regulation of energy metabolism.

Conclusion: This transcriptome-wide association study using monozygotic twins and generalized correlation identified differentially expressed genes and a broad spectrum of enriched biological pathways that may implicate metabolic health.

© 2020 Qihua Tan. Hosting by Science Repository. All rights reserved

Introduction

The body mass index (BMI) quantifies the amount of tissue mass including muscle, fat and bone in an individual. It is highly associated with cardiovascular disease and diabetes and has profound influences on life quality and mortality [1-3]. In clinical application, BMI is a simple and widely used metric for defining overweight ($25 \leq \text{BMI} < 30$, kg/m^2) and obesity ($\text{BMI} \geq 30$ kg/m^2), which are conditions tightly linked to the

metabolic syndrome (MetS). Many epidemiological and molecular studies have been conducted to find the genetic and non-genetic (environmental) mechanisms underlying individual BMI variation in order to identify potential causes of MetS and eventually to find strategies for mitigating its burden to public health. Multiple genetic variants have been reported to affect BMI in genome-wide association studies, albeit with the proportion of BMI variation accounted for remaining far from the overall genetic contribution estimated in twin

*Correspondence to: Professor Qihua Tan, M.D., Ph.D., Epidemiology and Biostatistics, Department of Public Health, Unit of Human Genetics, Department of Clinical Research, University of Southern Denmark, J.B. Winslows Vej 9 B, DK 5000, Odense C, Denmark; Tel: 004565503536; E-mail: qtan@health.sdu.dk

studies, although recent effort based on whole genome sequencing has largely increased the contribution by genomic sequence variations [4, 5]. Instead of focusing on the genetic polymorphisms which are static across life span, analysis of gene expression profiles can directly depict the dynamic activity of functional genes in regulating the variation of BMI, especially when controlling for genetics.

In transcriptome-wide association studies (TWAS), genetic variations can confound the relation between MetS related health conditions and gene expression (through functioning as cis- or trans-eQTLs). This is particularly crucial for BMI giving the high genetic contribution [4]. Compared to the ordinary case-control design using unrelated individuals, the use of twins has been proven a valuable approach in controlling genetic background, due to the shared genetic makeups in twins leveraging enriched statistical power [6, 7]. Moreover, current TWAS assumes linear relationship between gene expression and traits of interest, thus ignoring the diverse patterns of association in biology. By focusing on twins, we have investigated gene expression profiles in peripheral blood cells in association with BMI by introducing a generalized correlation method to capture different patterns (both linear and nonlinear) of association between gene expression and BMI while controlling for genetic confounding.

Methods

I Samples

This study is based on a twin cohort, the Middle Aged Danish Twins (MADT), from the Danish Twin Register [8]. There are in total 229 complete monozygotic twin pairs included in the analysis, 254 male and 204 female twins, with age at sampling ranging from 56 to 80 and BMI from 15.77 to 38.15. Anthropometric measures and whole blood samples were taken over the period from 2008 to 2011. Blood cell counts of 449 individuals were available and the blood cell counts of the other 9 individuals were imputed by the estimate CellCounts function in the R package minfi using DNA methylation data collected on the same blood samples [9, 10].

II Global Gene Expression Analysis

Whole blood samples were collected in PAXgene Blood RNA Tubes (PreAnalytiX GmbH, Hombrechtikon, Switzerland) and total RNA extracted using the PAXgene Blood miRNA kit (QIAGEN) according to the manufacturer's protocol. Concentration of the extracted RNA was determined using a NanoDrop spectrophotometer ND-8000 (NanoDrop Technologies), and the quality was assessed by the Agilent 2100 Bioanalyzer (Agilent Technologies).

Gene-expression analysis was performed using the Agilent SurePrint G3 Human GE 8×60K Microarray (Agilent Technologies), a dual-colour high-definition array containing 60K high quality probes of 60-mer. Sample labeling and array hybridization were carried out according to the 'Two-Colour Microarray-Based Gene Expression Analysis – Low Input Quick Amp Labeling'-protocol (Agilent Technologies). Samples were labeled with Cy5 and the reference consisting of a pool of 16 samples was labeled with Cy3. Hybridization, washing, scanning, and

quantification were performed according to the array manufacturer's recommendations.

III Data Pre-Processing

The raw intensity data was background-corrected using the NormExp method and was then within-array normalized by Loess normalization method and between-array normalized by quantile normalization [11]. The normalized values were used to calculate log₂-transformed Cy5/Cy3 ratios. Missing expression values were imputed by k-nearest neighbors averaging, and replicate probes were collapsed calculating the median. Data pre-processing was performed using the R packages limma [12]. All the probes on the Agilent SurePrint G3 array were re-annotated using GENCODE v.25 gene annotation database (Link).

IV Statistical Analysis

After normalization, we adjusted for covariates including age, sex, cell composition and first two PCs from the PCA (principal component analysis) on the gene expression data. We then applied a generalized measure of association, the generalized correlation coefficient (GCC), to investigate the association between intra-pair difference of BMI and intra-pair difference of expression to control the genetic and shared environmental effects, as proposed by Tan *et al.* [13]. GCC was computed using a ratio of maximum likelihoods for the marginal distribution and maximum weighted likelihoods for the joint distribution using the R package matie [14]. The mRNA probes with $p < 0.05$ were used for gene set enrichment analysis (GSEA) to detect biological pathways over-represented by the listed probes for functional interpretation [15].

Results

A total of 50599 probes were available on the microarray. We first removed house-keeping probes by calculating the coefficient of variation (CV) as standard deviation divided by mean of expression measurement for each probe. Probes with $CV < 0.1$ were filtered out leaving 37716 probes for subsequent analysis. We identified 2 probes with $p < 1e-05$ (A_33_P3289204 of *AAKI* gene, $p = 2.83e-06$; A_23_P79094 of *LILRA3* gene, $p = 7.83e-06$), 5 probes with $p < 1e-04$ (A_33_P3234809 of *PAX8* gene, $p = 4.37e-05$; A_32_P133767 of *C12orf42* gene, $p = 5.52e-05$; A_33_P3364060 of *HR* gene, $p = 6.11e-05$; A_33_P3213179 of *PPP1R3A* gene, $p = 8e-05$; A_23_P108280 of *CYP4F12* gene, $p = 9.87e-05$) and 1579 probes with $p < 0.05$ (Supplementary Table 1). Table 1 shows the top 20 probes ranked by p values ($p < 6.24e-04$). The Q-Q plot for probes with $p < 0.05$ is shown in (Figure 1). The figure does not display inflated statistical significance although the sample contains highly correlated monozygotic twin pairs. The Manhattan plot for probes with $p < 0.05$ are shown in (Figure 2).

Table 1: Top 20 probes from mRNA analysis. A is the association score or GCC from matie package.

Probe ID	Gene Symbol	p-value	A
A_33_P3289204	AAK1	2.83E-06	0.216
A_23_P79094	LILRA3	7.83E-06	0.207
A_33_P3234809	PAX8	4.37E-05	0.190
A_32_P133767	C12orf42	5.52E-05	0.187
A_33_P3364060	HR	6.11E-05	0.186
A_33_P3213179	PPP1R3A	8E-05	0.183
A_23_P108280	CYP4F12	9.87E-05	0.181
A_33_P3659808	PELP1	0.00014	0.177
A_23_P67702	ZNF85	0.00023	0.171
A_32_P9575	MRPL45	0.000236	0.171
A_23_P45851	HIAT1	0.000256	0.170
A_19_P00318014	SNHG11	0.000285	0.169
A_24_P395621	RUNDC1	0.000363	0.166
A_33_P3392250	FAM178A	0.000373	0.166
A_33_P3295328	CLEC6A	0.000381	0.165
A_23_P127948	ADM	0.00039	0.165
A_23_P131899	SDCBP2	0.000417456	0.164
A_23_P329924	HCAR2	0.000474119	0.163
A_33_P3351189	LOC100129603	0.000492092	0.162
A_23_P105973	SERPINA11	0.000623576	0.159

Table 2: KEGG pathways identified by GSEA over-representation analysis.

Gene Set Name	# Genes in Set (K)	Gene# Overlap (k)	in k/K	p-value	FDR q-value
KEGG_PATHWAYS_IN_CANCER	328	33	0.101	8.64E-09	1.61E-06
KEGG_JAK_STAT_SIGNALING_PATHWAY	155	17	0.110	1.08E-05	1.00E-03
KEGG_INSULIN_SIGNALING_PATHWAY	137	15	0.110	3.56E-05	1.76E-03
KEGG_DILATED_CARDIOMYOPATHY	92	12	0.130	3.78E-05	1.76E-03
KEGG_MAPK_SIGNALING_PATHWAY	267	22	0.082	5.72E-05	2.13E-03
KEGG_AXON_GUIDANCE	129	14	0.109	7.06E-05	2.19E-03
KEGG_HYPERTROPHIC_CARDIOMYOPATHY_HCM	85	11	0.129	8.49E-05	2.25E-03
KEGG_LEUKOCYTE_TRANSENDOTHELIAL_MIGRATION	118	13	0.110	1.09E-04	2.53E-03
KEGG_OLFACTORY_TRANSDUCTION	389	27	0.069	1.66E-04	3.44E-03
KEGG_NEUROACTIVE_LIGAND_RECEPTOR_INTERACTION	272	21	0.077	2.06E-04	3.84E-03
KEGG_CARDIAC_MUSCLE_CONTRACTION	80	10	0.125	2.36E-04	3.99E-03
KEGG_WNT_SIGNALING_PATHWAY	151	14	0.093	3.74E-04	5.14E-03
KEGG_TIGHT_JUNCTION	134	13	0.097	3.86E-04	5.14E-03
KEGG_PANCREATIC_CANCER	70	9	0.129	3.87E-04	5.14E-03
KEGG_NUCLEOTIDE_EXCISION_REPAIR	44	7	0.159	4.61E-04	5.56E-03
KEGG_LEISHMANIA_INFECTIO	72	9	0.125	4.78E-04	5.56E-03
KEGG_VIRAL_MYOCARDITIS	73	9	0.123	5.30E-04	5.80E-03
KEGG_T_CELL_RECEPTOR_SIGNALING_PATHWAY	108	11	0.102	7.05E-04	6.84E-03
KEGG_ARRHYTHMOGENIC_RIGHT_VENTRICULAR_CARDIOMYOPAT	76	9	0.118	7.15E-04	6.84E-03
HY_ARVC					
KEGG_GLYCOLYSIS_GLUONEOGENESIS	62	8	0.129	7.88E-04	6.84E-03
KEGG_INTESTINAL_IMMUNE_NETWORK_FOR_IGA_PRODUCTION	48	7	0.146	7.93E-04	6.84E-03
KEGG_FOCAL_ADHESION	201	16	0.080	8.09E-04	6.84E-03
KEGG_CELL_ADHESION_MOLECULES_CAMS	134	12	0.090	1.29E-03	1.04E-02
KEGG_P53_SIGNALING_PATHWAY	69	8	0.116	1.60E-03	1.11E-02
KEGG_PYRUVATE_METABOLISM	40	6	0.150	1.61E-03	1.11E-02
KEGG_MELANOGENESIS	102	10	0.098	1.62E-03	1.11E-02
KEGG_UBIQUITIN_MEDIATED_PROTEOLYSIS	138	12	0.087	1.66E-03	1.11E-02
KEGG_REGULATION_OF_ACTIN_CYTOSKELETON	216	16	0.074	1.72E-03	1.11E-02
KEGG_TGF_BETA_SIGNALING_PATHWAY	86	9	0.105	1.74E-03	1.11E-02

KEGG_BASAL_CELL_CARCINOMA	55	7	0.127	1.80E-03	1.12E-02
KEGG_ASTHMA	30	5	0.167	2.44E-03	1.45E-02
KEGG_LYSINE_DEGRADATION	44	6	0.136	2.65E-03	1.45E-02
KEGG_HUNTINGTONS_DISEASE	185	14	0.076	2.67E-03	1.45E-02
KEGG_ADHERENS_JUNCTION	75	8	0.107	2.73E-03	1.45E-02
KEGG_B_CELL_RECEPTOR_SIGNALING_PATHWAY	75	8	0.107	2.73E-03	1.45E-02
KEGG_ACUTE_MYELOID_LEUKEMIA	60	7	0.117	2.99E-03	1.54E-02
KEGG_CITRATE_CYCLE_TCA_CYCLE	32	5	0.1562	3.27E-03	1.65E-02
KEGG_COLORECTAL_CANCER	62	7	0.1129	3.60E-03	1.76E-02
KEGG_BASE_EXCISION_REPAIR	35	5	0.1429	4.87E-03	2.32E-02
KEGG_PURINE_METABOLISM	159	12	0.0755	5.30E-03	2.47E-02
KEGG_SMALL_CELL_LUNG_CANCER	84	8	0.0952	5.50E-03	2.49E-02
KEGG_PROXIMAL_TUBULE_BICARBONATE_RECLAMATION	23	4	0.1739	5.69E-03	2.52E-02
KEGG_ENDOMETRIAL_CANCER	52	6	0.1154	6.17E-03	2.67E-02
KEGG_METABOLISM_OF_XENOBIOTICS_BY_CYTOCHROME_P450	70	7	0.1	7.04E-03	2.98E-02
KEGG_NON_SMALL_CELL_LUNG_CANCER	54	6	0.1111	7.42E-03	3.07E-02
KEGG_DRUG_METABOLISM_CYTOCHROME_P450	72	7	0.0972	8.20E-03	3.27E-02
KEGG_CELL_CYCLE	128	10	0.0781	8.26E-03	3.27E-02
KEGG_GALACTOSE_METABOLISM	26	4	0.1538	8.91E-03	3.38E-02
KEGG_GLYCOSAMINOGLYCAN_BIOSYNTHESIS_HEPARAN_SULFATE	26	4	0.1538	8.91E-03	3.38E-02
KEGG_PHOSPHATIDYLINOSITOL_SIGNALING_SYSTEM	76	7	0.0921	1.09E-02	4.06E-02

Table 3: Reactome pathways identified by GSEA over-representation analysis.

Gene Set Name	# Genes	in# Genes	in k/K	p-value	FDR q-value
	Gene Set (K)	Overlap (k)			
REACTOME_IMMUNE_SYSTEM	933	72	0.077	9.41E-12	6.34E-09
REACTOME_SIGNALING_BY_GPCR	920	65	0.071	3.38E-09	1.14E-06
REACTOME_GPCR_DOWNSTREAM_SIGNALING	805	58	0.072	1.18E-08	2.66E-06
REACTOME_ADAPTIVE_IMMUNE_SYSTEM	539	43	0.080	5.58E-08	9.40E-06
REACTOME_HEMOSTASIS	466	37	0.079	5.16E-07	6.96E-05
REACTOME_DEVELOPMENTAL_BIOLOGY	396	32	0.081	2.04E-06	2.29E-04
REACTOME_METABOLISM_OF_LIPIDS_AND_LIPOPROTEINS	478	35	0.073	6.30E-06	5.73E-04
REACTOME_CELL_CYCLE	421	32	0.076	7.28E-06	5.73E-04
REACTOME_SIGNALLING_BY_NGF	217	21	0.097	7.65E-06	5.73E-04
REACTOME_MHC_CLASS_II_ANTIGEN_PRESENTATION	91	12	0.132	3.39E-05	2.28E-03
REACTOME_CELL_CYCLE_MITOTIC	325	25	0.077	5.74E-05	3.52E-03
REACTOME_GROWTH_HORMONE_RECEPTOR_SIGNALING	24	6	0.250	8.73E-05	4.90E-03
REACTOME_TRANSCRIPTIONAL_REGULATION_OF_WHITE_ADIPOCYTE_D DIFFERENTIATION	72	10	0.139	9.67E-05	5.01E-03
REACTOME_G_ALPHA1213_SIGNALLING_EVENTS	74	10	0.135	1.22E-04	5.89E-03
REACTOME_ANTIGEN_PROCESSING_UBIQUITINATION_PROTEASOME_D EGRADATION	212	18	0.085	1.80E-04	7.56E-03
REACTOME_TRANSMEMBRANE_TRANSPORT_OF_SMALL_MOLECULES	413	28	0.068	1.88E-04	7.56E-03
REACTOME_CLASS_I_MHC_MEDIATED_ANTIGEN_PROCESSING_PRESEN TATION	251	20	0.080	1.91E-04	7.56E-03
REACTOME_P75_NTR_RECEPTOR_MEDIATED_SIGNALLING	81	10	0.124	2.61E-04	9.79E-03
REACTOME_GPCR_LIGAND_BINDING	408	27	0.066	3.54E-04	1.26E-02
REACTOME_PHASE1_FUNCTIONALIZATION_OF_COMPOUNDS	70	9	0.129	3.87E-04	1.30E-02
REACTOME_OLFACTORY_SIGNALING_PATHWAY	328	23	0.070	4.23E-04	1.30E-02
REACTOME_METABOLISM_OF_CARBOHYDRATES	247	19	0.077	4.23E-04	1.30E-02
REACTOME_SIGNALING_BY_NOTCH	103	11	0.107	4.71E-04	1.38E-02
REACTOME_AXON_GUIDANCE	251	19	0.076	5.16E-04	1.45E-02
REACTOME_SIGNALING_BY_ILS	107	11	0.103	6.51E-04	1.76E-02
REACTOME_INNATE_IMMUNE_SYSTEM	279	20	0.072	7.41E-04	1.85E-02
REACTOME_NEURONAL_SYSTEM	279	20	0.072	7.41E-04	1.85E-02
REACTOME_METABOLISM_OF_AMINO_ACIDS_AND_DERIVATIVES	200	16	0.080	7.67E-04	1.85E-02
REACTOME_PLATELET_HOMEOSTASIS	78	9	0.115	8.64E-04	1.97E-02

REACTOME_COSTIMULATION_BY_THE_CD28_FAMILY	63	8	0.127	8.78E-04	1.97E-02
REACTOME_CYTOCHROME_P450_ARRANGED_BY_SUBSTRATE_TYPE	51	7	0.137	1.15E-03	2.49E-02
REACTOME_MEIOSIS	116	11	0.095	1.27E-03	2.69E-02
REACTOME_PLATELET_SENSITIZATION_BY_LDL	16	4	0.250	1.40E-03	2.77E-02
REACTOME_XENOBIOTICS	16	4	0.250	1.40E-03	2.77E-02
REACTOME_STRIATED_MUSCLE_CONTRACTION	27	5	0.185	1.50E-03	2.88E-02
REACTOME_NGF_SIGNALLING_VIA_TRKA_FROM_THE_PLASMA_MEMBRANE	137	12	0.088	1.56E-03	2.92E-02
REACTOME_GLUCOSE_METABOLISM	69	8	0.116	1.60E-03	2.92E-02
REACTOME_LIPOPROTEIN_METABOLISM	28	5	0.179	1.77E-03	3.15E-02
REACTOME_RNA_POL_I_RNA_POL_III_AND_MITOCHONDRIAL_TRANSCRIPTION	122	11	0.090	1.91E-03	3.31E-02
REACTOME_NRAGE_SIGNALS_DEATH_THROUGH_JNK	43	6	0.140	2.36E-03	3.97E-02
REACTOME_GENERIC_TRANSCRIPTION_PATHWAY	352	22	0.063	2.42E-03	3.99E-02
REACTOME_PRE_NOTCH_EXPRESSION_AND_PROCESSING	44	6	0.136	2.65E-03	4.26E-02
REACTOME_CELL_DEATH_SIGNALLING_VIA_NRAGE_NRF1_AND_NAIP1	60	7	0.117	2.99E-03	4.68E-02
REACTOME_DNA_REPAIR	112	10	0.089	3.24E-03	4.97E-02
REACTOME_ACTIVATION_OF_THE_API_FAMILY_OF_TRANSCRIPTION_FACTORS	10	3	0.300	3.32E-03	4.97E-02




Figure 1: Q-Q plots for single probe analysis ($p < 0.05$ only). The top probes (red) deviate upward from the p values of null hypothesis.

By submitting the genes in Supplementary (Table 1) to GSEA, we identified 50 KEGG (The Kyoto Encyclopedia of Genes and Genomes) pathways with $FDR < 0.05$ (Table 2). Among the top significant pathways are JAK/STAT signaling pathway ($FDR = 1e-03$), insulin signaling pathway ($FDR = 1.76e-03$), MAPK signaling pathway ($FDR = 2.13e-03$), axon guidance ($FDR = 2.19e-03$), etc. Considering the fact that KEGG focuses on intermediary metabolism, we further performed pathway analysis using the Reactome database containing higher-level pathways compiled based on a broad range of species [16]. Table 3 shows the 45 Reactome pathways identified with $FDR < 0.05$. Among them, nine are highly significant with $FDR < 1e-03$ ($p < 1e-05$) such as immune system ($FDR = 1.34e-09$), G protein-coupled receptor (GPCR) signaling pathway ($FDR = 1.14e-06$), hemostasis ($FDR = 6.96e-05$), developmental biology ($FDR = 2.29e-04$), signaling by nerve growth factor (NGF) ($FDR = 5.73e-04$), olfactory signaling ($FDR = 1.3e-02$) etc. Table 3 also contains multiple pathways directly implicated in metabolism including metabolism of lipids and lipoproteins ($FDR = 5.73e-04$), metabolism of carbohydrates ($FDR = 1.30e-02$), glucose metabolism ($FDR = 2.93e-02$) and lipoprotein metabolism ($FDR = 3.15e-02$).




Figure 2: Manhattan plot showing probe significance ($p < 0.05$ only) along each chromosome.

Discussion

The high genetic contribution to BMI variation as estimated in twins, and in family studies, suggests the need to study the relation between transcriptional activity and BMI while controlling for genetics, given the fact that genetic variations could affect both BMI and gene expression [4, 17-19]. Instead of using unrelated samples, focusing on genetically identical individuals (e.g. MZ twins) can control for genetic variations in the study samples. However, traditional statistical testing on dependent samples requires modeling the relatedness in sample clusters using, for example, the mixed effects modeling assigning sample correlation as random effect variables with increased model complexity and reduced power in statistical testing. The application of GCC on twin data provides a simple but efficient way for handling related samples in TWAS. Most importantly, generalized correlation is a non-parametric method by nature meaning that its assessment of correlation between gene expression and BMI was done without imposing any assumption such as the linear relationship in regression modeling. The latter is sensitive to outlier observations in model fitting. As a result, our reported gene expression markers detected by the model-free approach are biologically meaningful and significantly enriched in functional pathways closely implicated in metabolism, although, many other pathways e.g. related to blood cell functioning (immune biology and hemostasis) and neuro function were also enriched.

In (Figure 1), the top probes (red coloured) show upward deviation from p values under the null hypothesis suggesting their non-random association with BMI. In accordance, functional annotation and published literature confirm their biological relevance. In (Table 1), the most significant probe (A_33_P3289204, $p=2.83e-06$) is from *AAK1* gene. The gene has been shown as a positive regulator of the Notch pathway which is a novel regulator of metabolism [20, 21]. Both *AAK1* and Notch signaling are implicated in neurological impairments, which have been shown to relate to metabolic disorders [22-24]. The number 2 probe (A_23_P79094, $p=7.83e-06$) belongs to *LILRA3* for which a SNP (rs367070) has been reported to associate with HDL-C [25]. The human *LILRA* gene family has diverse functions characterized by regulation of inflammation and immune tolerance [26]. The *LILRA3* regulated immunity alteration could be involved in obesity, a condition characterized by chronic low-grade inflammation [27]. For the third gene in (Table 1), *PAX8*, differential DNA methylation was identified to associate with gestational famine exposure and metabolic traits [28]. Other interesting genes linked to top probes of (Table 1) include *PPP1R3A* whose SNP variation has been associated with risk of type 2 diabetes and *CYP4F12* which belongs to the cytochrome *P450 4 (CYP4)* family implicated in various biological functions including inflammation, cardiovascular health, and cancer [29, 30].

Biological pathway analysis was performed using KEGG and Reactome databases with both revealing significantly enriched functional pathways implicated in metabolic health together with other biological functions. For example, the top pathways in (Table 2) include pathways in cancer and insulin signaling. In the literature, development of insulin resistance and hyperinsulinemia has been shown as a clear link between adipose tissue expansion and etiology of diseases like obesity, type-2 diabetes and cancer [31]. Likewise, the implication of MAPK signaling in obesity-related immune paralysis and cancer has been reviewed very

recently [32]. Another top significant pathway, JAK/STAT signaling pathway in (Table 2) is a highly conserved functional pathway required for normal homeostasis which, when dysregulated, contributes to the development of obesity and diabetes [33].

Like the KEGG pathways (Table 2), the importance of immunity is also reflected in the top significant Reactome pathways shown in (Table 3) (immune system with FDR 6.34e-09 and adaptive immune system with FDR 9.40e-06). The phenomenon can be explained by the target tissue used in this study, i.e. whole blood, which comprises of immune cells. Moreover, results in (Table 3) emphasize high implication of GPCR signaling in BMI variation. This is highly interesting as the melanocortin-4 receptor (MC4R), a GPCR embedded in the membranes of nerve cells in the brain's appetite control center, has been shown to provide clues to obesity treatment [34]. Another study has found that pathway of GPCR was overrepresented and it is associated with pediatric obesity [35]. As a potential therapeutic target for intervention in cognitive deficits, the association of GPCR signaling pathway with BMI could reflect the intrinsic connection between cognition and obesity [36, 37]. Table 3 also contains significant pathways directly related to metabolism of lipids, lipoproteins and carbohydrates, etc. which are closely related to obesity development as well as pathways overlapping with (Table 2) including axon guidance reportedly to implicate in early-onset obesity [38], and olfactory signaling/transduction shown to regulate lipid metabolism through neuroendocrine signaling in *Caenorhabditis elegans* [38, 39]. Methylation of olfactory pathway genes has been associated with dietary intake and obesity features [40].

In summary, by introducing generalized correlation coefficient for assumption-free association analysis and using monozygotic twins to control genetic confounding, this transcriptome-wide association study on BMI using peripheral blood identified differentially expressed genes and their enriched biological pathways implicated in multiple biological functions including immune biology, hemostasis, neural function, cancer, and metabolism. Findings from this study merit replications using independent samples to verify expression markers and functional pathways for characterizing and determining non-genetic etiology of obesity and health conditions.

Statement of Ethics

The survey was approved by The Regional Scientific Ethical Committees for Southern Denmark (S-VF-19980072) and conducted in accordance with the Helsinki II declaration, with informed consent to participate in the survey obtained from all participants.

Funding

This study was jointly supported by the Lundbeck Foundation (grant number R170-2014-1353), the DFF research project 1 from the Danish Council for Independent Research, Medical Sciences (DFF-FSS) (grant number DFF – 6110-00114); the Novo Nordisk Foundation Medical and Natural Sciences Research Grant (grant number NNF13OC0007493). JB is grateful for funding from VILLUM Young Investigator grant nr. 13154 and funding from the BMBF center CLINSPECT-M.

Conflicts of Interest

None.

REFERENCES

1. De Gonzalez AB, Hartge P, Cerhan JR, Flint AJ, Hannan L et al. (2010) Body-mass index and mortality among 1.46 million white adults. *N Engl J Med* 363: 2211-2219. [[Crossref](#)]
2. NCD Risk Factor Collaboration (2016) Trends in adult body-mass index in 200 countries from 1975 to 2014: A pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *Lancet* 387: 1377-1396. [[Crossref](#)]
3. Gaziano TA, Opie LH (2009) Body-mass index and mortality. *Lancet* 374: 113-114. [[Crossref](#)]
4. Hjelmberg JVB, Fagnani C, Silventoinen K, McGue M, Korkeila M et al. (2008) Genetic influences on growth traits of BMI: A longitudinal study of adult twins. *Obesity* 16: 847-852. [[Crossref](#)]
5. Wainschtein P, Jain DP, Yengo L, Zheng Z, Shadyab AH et al. (2019) Recovery of trait heritability from whole genome sequence data. *BioRxiv*.
6. Li W, Christiansen L, Hjelmberg J, Baumbach J, Tan Q (2018) On the power of epigenome-wide association studies using a disease-discordant twin design. *Bioinformatics* 34: 4073-4078. [[Crossref](#)]
7. Tan Q (2013) Epigenetic epidemiology of complex diseases using twins. *Med Epigenet* 1: 46-51.
8. Gaist D, Bathum L, Skythhe A, Jensen TK, McGue M et al. (2000) Strength and anthropometric measures in identical and fraternal twins: No evidence of masculinization of females with male co-twins. *Epidemiology* 11: 340-343. [[Crossref](#)]
9. Aryee MJ, Jaffe AE, Corrada Bravo H, Ladd Acosta C, Feinberg AP et al. (2014) Minfi: A flexible and comprehensive Bioconductor package for the analysis of Infinium DNA methylation microarrays. *Bioinformatics* 30: 1363-1369. [[Crossref](#)]
10. Soerensen M, Li W, Debrabant B, Nygaard M, Mengel From J et al. (2019) Epigenome-wide exploratory study of monozygotic twins suggests differentially methylated regions to associate with hand grip strength. *Biogerontology* 20: 627-647. [[Crossref](#)]
11. Yang YH, Dudoit S, Luu P, Lin DM, Peng V et al. (2002) Normalization for cDNA microarray data: A robust composite method addressing single and multiple slide systematic variation. *Nucleic Acids Res* 30: e15. [[Crossref](#)]
12. Ritchie ME, Silver J, Oshlack A, Holmes M, Diyagama D et al. (2007) A comparison of background correction methods for two-colour microarrays. *Bioinformatics* 23: 2700-2707. [[Crossref](#)]
13. Tan Q, Christiansen L, Hjelmberg JVB, Christensen K (2015) Twin methodology in epigenetic studies. *J Exp Biol* 218: 134-139. [[Crossref](#)]
14. Murrell B, Murrell D, Murrell H (2016) Discovering general multidimensional associations. *PLoS One* 11: e0151551. [[Crossref](#)]
15. Subramanian A, Tomayo P, Mootha VK, Mukherjee S, Ebert BL et al. (2005) Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci USA* 102: 15545-15550. [[Crossref](#)]
16. Haw R, Hermjakob H, D'Eustachio P, Stein L (2011) Reactome pathway analysis to enrich biological discovery in proteomics data sets. *Proteomics* 11: 3598-3613. [[Crossref](#)]
17. Li S, Duan H, Pang Z, Zhang D, Duan H et al. (2013) Heritability of eleven metabolic phenotypes in Danish and Chinese twins: A cross-population comparison. *Obesity* 21: 1908-1914. [[Crossref](#)]
18. Skarić Jurić T, Ginsburg E, Kobylansky E, Malkin I, Naranić NS et al. (2003) Complex segregation analysis of body height, weight and BMI in pedigree data from Middle Dalmatia, Croatia. *Coll Antropol* 27: 135-149. [[Crossref](#)]
19. Lee KE, Klein BEK, Klein R (2003) Familial aggregation of components of the multiple metabolic syndrome in the Framingham Heart and Offspring Cohorts: Genetic Analysis Workshop Problem 1. *BMC Genet* 4: S94. [[Crossref](#)]
20. Gupta Rossi N, Ortica S, Meas Yedid V, Heuss S, Moretti J et al. (2011) The adaptor-associated kinase 1, AAK1, is a positive regulator of the notch pathway. *J Biol Chem* 286: 18720-18730. [[Crossref](#)]
21. Bi P, Kuang S (2015) Notch signaling as a novel regulator of metabolism. *Trends Endocrinol Metab* 26: 248-255. [[Crossref](#)]
22. Schubert KO, Föcking M, Prehn JHM, Cotter DR (2012) Hypothesis review: Are clathrin-mediated endocytosis and clathrin-dependent membrane and protein trafficking core pathophysiological processes in schizophrenia and bipolar disorder. *Mol Psychiatry* 17: 669-681. [[Crossref](#)]
23. Ables JL, Breunig JJ, Eisch AJ, Rakic P (2011) Not(ch) just development: Notch signalling in the adult brain. *Nat Rev Neurosci* 12: 269-283. [[Crossref](#)]
24. O'Brien PD, Hinder LM, Callaghan BC, Feldman EL (2017) Neurological consequences of obesity. *Lancet Neurol* 16: 465-477. [[Crossref](#)]
25. Stanley A, Ponde CK, Rajani RM, Ashavaid TF (2017) Association between genetic loci linked to HDL-C levels and Indian patients with CAD: a pilot study. *Heart Asia* 9: 9-13. [[Crossref](#)]
26. Hirayasu K, Arase H (2015) Functional and genetic diversity of leukocyte immunoglobulin-like receptor and implication for disease associations. *J Hum Genet* 60: 703-708. [[Crossref](#)]
27. Lee H, Lee IS, Choue R (2013) Obesity, Inflammation and Diet. *Pediatr Gastroenterol Hepatol Nutr* 16: 143-152. [[Crossref](#)]
28. Finer S, Iqbal MS, Lowe R, Ogunkolade BW, Pervin S et al. (2016) Is famine exposure during developmental life in rural Bangladesh associated with a metabolic and epigenetic signature in young adulthood? A historical cohort study. *BMJ Open* 6: e011768. [[Crossref](#)]
29. Sánchez Pozos K, Ortíz López MG, Peña Espinoza BI, Granados Silvestre MLA, Jiménez Jacinto V et al. (2018) Whole-exome sequencing in maya indigenous families: variant in PPP1R3A is associated with type 2 diabetes. *Mol Genet Genomics* 293: 1205-1216. [[Crossref](#)]
30. Jarrar YB, Lee SJ (2019) Molecular functionality of cytochrome P450 4 (CYP4) genetic polymorphisms and their clinical implications. *Int J Mol Sci* 20: 4274. [[Crossref](#)]
31. Poloz Y, Stambolic V (2015) Obesity and cancer, a case for insulin signaling. *Cell Death Dis* 6: e2037. [[Crossref](#)]
32. Donohoe F, Wilkinson M, Baxter M, Brennan DJ (2020) Mitogen-Activated Protein Kinase (MAPK) and Obesity-Related Cancer. *Int J Mol Sci* 21: 1241. [[Crossref](#)]
33. Gurzov EN, Stanley WJ, Pappas EG, Thomas HE, Gough DJ (2016) The JAK/STAT pathway in obesity and diabetes. *FEBS J* 283: 3002-3015. [[Crossref](#)]

34. Ghamari Langroudi M, Digby GJ, Sebag JA, Millhauser GL, Palomino R et al. (2015) G-protein-independent coupling of MC4R to Kir7.1 in hypothalamic neurons. *Nature* 520: 94-98. [[Crossref](#)]
35. Dorajoo R, Ong RTH, Sim X, Wang L, Liu W et al. (2017) The contribution of recently identified adult BMI risk loci to paediatric obesity in a Singaporean Chinese childhood dataset. *Pediatr Obes* 12: e46-e50. [[Crossref](#)]
36. Azam S, Haque ME, Jakaria M, Jo SH, Kim IS et al. (2020) G-Protein-Coupled Receptors in CNS: A Potential Therapeutic Target for Intervention in Neurodegenerative Disorders and Associated Cognitive Deficits. *Cells* 9: 506. [[Crossref](#)]
37. Smith E, Hay P, Campbell L, Trollor JN (2011) A review of the association between obesity and cognitive function across the lifespan: Implications for novel approaches to prevention and treatment. *Obes Rev* 12: 740-55. [[Crossref](#)]
38. Zeltser LM (2019) Axon Guidance Molecules Implicated in Early-Onset Obesity. *Trends Neurosci* 42: 439-440. [[Crossref](#)]
39. Mutlu AS, Gao SM, Zhang H, Wang MC (2020) Olfactory specificity regulates lipid metabolism through neuroendocrine signaling in *Caenorhabditis elegans*. *Nat Commun* 11: 1450. [[Crossref](#)]
40. Ramos Lopez O, Riezu Boj JI, Milagro FI, Zulet MA, Santos JL et al. (2019) Associations between olfactory pathway gene methylation marks, obesity features and dietary intakes. *Genes Nutr* 14: 11. [[Crossref](#)]