

# Involvement of the Argentine Precordillera terrane in the Famatinian mobile belt: U-Pb SHRIMP and metamorphic evidence from the Sierra de Pie de Palo

C. Casquet Departamento de Petrología y Geoquímica, Universidad Complutense, 28040 Madrid, Spain

E. Baldo Departamento de Geología, Universidad Nacional de Córdoba, Argentina

R.J. Pankhurst British Antarctic Survey, Cambridge CB3 0ET, United Kingdom

C.W. Rapela Centro de Investigaciones Geológicas, Universidad Nacional de La Plata, Calle 1 No. 644, La Plata, Argentina

C. Galindo Departamento de Petrología y Geoquímica, Universidad Complutense, 28040 Madrid, Spain

C.M. Fanning Research School of Earth Sciences, Australian National University, Canberra ACT 0200, Australia

J. Saavedra Consejo Superior de Investigaciones Científicas, Instituto de Agrobiología y Recursos Naturales, 37071 Salamanca, Spain

## ABSTRACT

New data suggest that the eastern margin of the Argentine Precordillera terrane comprises Grenvillian basement and a sedimentary cover derived from it that were together affected by Middle Ordovician deformation and metamorphism during accretion to the Gondwana margin. The basement first underwent low pressure/temperature (*P/T*) type metamorphism, reaching high-grade migmatitic conditions in places ( $686 \pm 40$  MPa,  $790 \pm 17$  °C), comparable to the Grenvillian  $M_2$  metamorphism of the supposed Laurentian counterpart of the terrane. The second metamorphism, recognized in the cover sequence, is of Famatinian age and took place under higher *P/T* conditions, following a clockwise *P-T* path (baric peak:  $1300 \pm 100$  Mpa,  $600 \pm 50$  °C). Low-U zircon overgrew detrital Grenvillian cores as pressure fell from its peak, and yields U-Pb SHRIMP ages of ca. 460 Ma. This is interpreted as the age of ductile thrusting coincident with early uplift; initial accretion to Gondwana must have occurred before this. The absence of late Neoproterozoic detrital zircons is consistent with a Laurentian origin of the Argentine Precordillera terrane.

**Keywords:** Argentina, Ordovician, continent collision, metamorphism, SHRIMP zircon dating.

## INTRODUCTION

The origin of the Argentine Precordillera terrane has kept those involved in Paleozoic paleogeography and tectonics intrigued for some time. It is widely accepted as an exotic terrane, derived from Laurentia and accreted to the southwestern margin of Gondwana between Early Ordovician and Early Devonian time (Dalla Salda et al., 1992; Astini et al., 1995; Dalziel, 1997; Rapela et al., 1998). Another suggestion (Aceñolaza and Toselli, 1999) is that the terrane was autochthonous to Gondwana, originating in the Neoproterozoic margin of South America–Africa–Antarctica, and that it reached its present location by prolonged strike-slip movement. We provide new data highly relevant to these problems.

In the Precordillera, there is a nonmetamorphosed to weakly metamorphosed Cambrian–Devonian sedimentary cover (e.g., Keller et al., 1998) underlain by crystalline basement of Grenville age, inferred from U-Pb ages of xenoliths in Miocene intrusions (Kay et al., 1996). Grenvillian ages have also been obtained from the metamorphic rocks of the Sierra de Pie de Palo, Western Sierras Pampeanas, east of the Precordillera (Fig. 1), considered to be the main easternmost exposure of the Argentine Precordillera terrane

basement. These rocks include a Neoproterozoic ophiolite sequence and orthogneisses (McDonough et al., 1993; Vujovich and Kay, 1998; Pankhurst and Rapela, 1998). Controls on the timing of the Paleozoic collision and tectonothermal history are sparse. A Middle Ordovician unconformity and change in style of sedimentation in the Precordillera are believed to indicate collision-related drowning of the former carbonate platform (Astini et al., 1995). Ramos et al. (1998) interpreted  $^{39}\text{Ar}$ – $^{40}\text{Ar}$  data from the Sierra de Pie de Palo as being consistent with recrystallization; a single amphibolite hornblende yielded a plateau age of  $464.3 \pm 0.3$  Ma. However, the nature and extent of associated metamorphism are unknown, as are its pressure-temperature (*P-T*) characteristics and its relationship to plutonism and deformation in the Early Ordovician Famatinian mobile belt of the Sierras Pampeanas (Pankhurst and Rapela, 1998).

Our U-Pb SHRIMP data and metamorphic evidence demonstrate that the easternmost exposures of the Argentine Precordillera terrane were involved in Famatinian orogenic events and underwent relatively high *P/T* type regional metamorphism. A metamorphosed sedimentary cover to the Grenvillian basement is recognized for the first time, consistent with a

Laurentian origin. The time of initial terrane collision and prograde Famatinian metamorphism is defined as slightly before 460 Ma.


## GEOLOGY OF SIERRA DE PIE DE PALO

The Sierra is an imbricate ductile thrust system with a top-to-the-west sense of relative movement. We distinguish a number of allochthonous tectonic units overlying epimetamorphic basement formed by the Cauce Group sedimentary rocks (Fig. 1). The lower unit consists of mafic and ultramafic rocks (for which a Grenvillian ophiolite origin was suggested by Vujovich and Kay, 1998), black schists, and quartzites. It is separated from the Cauce Group by the Pirquitas thrust, which underwent protracted activity from 464 Ma until ca. 396 Ma (Ramos et al., 1998). The upper units comprise monotonous migmatites, schists, and quartzites, topped by the Difunta Correa Sedimentary Sequence of marbles, Capelitic schists, and quartzites (Baldo et al., 1998). Amphibolites are widespread, particularly in the lower unit and Difunta Correa Sedimentary Sequence; most were probably dikes, although some are metasediments. Peraluminous and metaluminous orthogneisses are also common, as are mylonites, particularly at the main unit boundaries. At least two foliations are found in zones protected from ductile shearing; the second is usually a differentiated crenulation cleavage. Retrograde extensional shear zones, with a top-to-the-east sense of relative movement, are present in the eastern part of the Sierra. They are superimposed on the thrust zones and caused major thinning of the sedimentary sequence in the uppermost tectonic units. The thickness of the tectonic pile is at least 15 km.

## U-Pb SHRIMP GEOCHRONOLOGY

Sample SPP-1001 is a mylonitic Ca-pelitic schist from north of the mapped area (on the road to the top of the Sierra;  $31^{\circ}23'30''\text{S}$ ,  $68^{\circ}01'07''\text{W}$ ). Its mineral composition corresponds to similar rocks in the Difunta Correa

**Figure 1. Geological sketch map of southern Sierra de Pie de Palo. Inset shows location of Sierras Pampeanas (SP) and Argentine Precordillera (APC).**



Sedimentary Sequence: Qtz, Pl, Ms, Bt, Grt, Hbl, Ep, Ilm, Rt, and Zr (mineral abbreviations after Kretz, 1983). The separated zircons are subhedral and elongate, to 200  $\mu\text{m}$ . Cathodoluminescence imaging shows their composite nature; cores consisting of fragments and subrounded grains with oscillatory zoning characteristic of igneous crystallization are invariably overgrown by unzoned low-U rims, generally 10–60  $\mu\text{m}$  thick (Fig. 2). Discrete cores and rims were selected for ion microprobe U-Pb analysis on SHRIMP II at the Australian National University, using a beam with an  $\sim 25\text{-}\mu\text{m}$ -diameter spot (Compston et al., 1992). Data for 10 cores and 10 rims are plotted in Figure 2 (see Data Repository<sup>1</sup>). The cores have inferred crystallization ages of 1032–1224 Ma (apart from one with an age of ca. 2900 Ma). With a single exception, the rims have Th/U = 0.01–0.1, indicating a metamorphic origin, and plot close to 460 Ma; this is interpreted as the age of metamorphic overgrowth. Two rim analyses plot to the right of the main group and are thought to reflect subsequent Pb loss.


#### CONDITIONS OF METAMORPHISM

The *P-T* conditions of metamorphism were assessed from Grt microdomains using the computer program TWQ2.02 and the thermodynamic database BA96a.DAT (Berman,

1991). Two contrasting rocks are compared: a mylonitized Ca-pelitic schist (SPP-882) collected a few meters from sample SPP-1001, and a mylonitized migmatitic paragneiss (SPP-2057) from the bottom of one of the upper units (Fig. 1).

SPP-882 has a complex mineralogy (Qtz, Ms, Pg, Bt, Grt, Ky, St, Pl, Hbl, Ep, Rt, Ilm) with evidence of three superimposed foliations (Baldo et al., 1998). Garnet, Ms, Bt, Ky, St, Hbl, and Rt are prekinematic relative to the mylonitic foliation,  $S_{\text{myl}}$ , which is defined by a banded matrix of granoblastic Qtz lenses,

layers of Pl, Ep, and dynamically recrystallized Ms, Pg, and Bt. Garnet porphyroclasts enveloped by  $S_{\text{myl}}$  contain orientated inclusions that define a discordant sigmoidal internal  $S_1$  foliation. This suggests that Grt grew in microlithons during the development of a new external foliation ( $S_2$ ) that was obliterated by mylonitization, except in strain shadows where relics of deformed  $S_1$  are preserved. A differentiated crenulation cleavage is common in these rocks away from the shear zones. Minerals contained in garnet are Qtz, St, Mrg, Bt, Pg, Ep, Chl, Rt, Ilm, and Pl. Garnet shows



**Figure 2. Tera-Wasserburg U-Pb concordia plot for SHRIMP spot analyses of zircons from sample SPP-1001. Black ellipses are 68% confidence limits for analyses of crystal cores, uncorrected for common Pb; shaded ellipses are for low-U overgrowths (see cathodoluminescence image). Open ellipses show points shifted to concordia when corrected for common Pb, as estimated from measured  $^{204}\text{Pb}$  peaks, albeit with error magnification due to low Pb contents. Data for cores clearly indicate crystallization in interval 1000–1200 Ma, whereas rims formed at close to 460 Ma (excluding single cross-hatched ellipse, which seems to have undergone more recent Pb loss). There is no evidence for any event between these two ages.**

<sup>1</sup>GSA Data Repository item 2001084, U-Pb SHRIMP data for sample SPP-1001, and chemical analyses of minerals considered for *P-T* computations, is available on request from Documents Secretary, GSA, P.O. Box 9140, Boulder, CO 80301-9140, editing@geosociety.org, or at www.geosociety.org/pubs/ft2001.htm.




Figure 3. Pressure-temperature ( $P$ - $T$ ) plot of metamorphic conditions derived from samples from Sierra de Pie de Palo. White boxes: mylonitic Ca-pelitic schist SPP-882 (Baldo et al., 1998). Dark gray boxes: mylonitic migmatite SPP-2057. Light gray boxes:  $P$ - $T$  conditions for Grenvillian M2 metamorphism in Llano uplift, Texas (Carlson and Schwarze, 1997). Arrows:  $P$ - $T$  trajectories. Equilibrium curves are from Spear et al. (1999) and Spear (1993). Abbreviations after Kretz (1983).




Figure 4. Photomicrographs of mylonitized migmatite SPP-2057, showing (A) pre-mylonitic type II garnet and, (B) late mylonitic idiomorphic type III garnet. Respective chemical zoning is shown on right (C) (C is core; B is border). Abbreviations after Kretz (1983).

growth zoning with a continuous increase of  $X_{grs}$  ( $X$  = mole fraction) and  $X_{prp}$  relative to  $X_{alm}$  and  $X_{sp}$  toward the rim. Three  $P$ - $T$  points were determined in this rock (Fig. 3).  $M_1$  ( $650 \pm 50$  Mpa,  $550 \pm 50$  °C) corresponds to conditions during prograde growth, computed from core garnet and mineral inclusions.  $M_2$  ( $1300 \pm 100$  Mpa,  $600 \pm 50$  °C) corresponds to close to peak pressure conditions, computed from rim garnet and minerals protected from  $S_{myl}$  in a nearby strain shadow.  $M_3$  ( $900 \pm 130$  Mpa,  $570 \pm 5$  °C) represents mylonitization and is based on syntectonic matrix minerals.

SPP-2057 shows a penetrative mylonitic fabric that wraps around pre-tectonic Grt (type II), Pl, Ms, By, fibrolitic pseudomorphs after Ky blades, and Ilm. Sillimanite is also found included in Ms, defining an internal foliation discordant to  $S_{myl}$ . Scarce relicts of St (and Bt and type I Grt) are preserved armored in plagioclase porphyroclasts. Mylonitization was accompanied by dynamic recrystallization of micas along grain boundaries and of Qtz as ribbons; Rt formed at the expense of Ilm. Clusters of crystallites of an Al silicate (probably Ky) also formed during shearing, along grain boundaries of Pl, Grt, and mica, with a mean trend parallel to  $S_{myl}$ . Furthermore, there are very small ( $\sim 0.1$  mm) idiomorphic Grt grains that apparently grew across  $S_{myl}$  (type III, Fig. 4). These probably formed late during shearing and are chemically unlike the pre-tectonic garnets. Type I Grt is the poorest in Ca ( $alm_{79.2}prp_{15.2}grs_{3.7}sp_{1.65}$ ). Type II Grt porphyroclasts are relatively homogeneous and slightly richer in Ca ( $alm_{78-79.3}prp_{13-14.4}grs_{4.9-6}sp_{1.2-2.7}$ ), except at the edges, where an abrupt jump to type III compositions is found. Type III Grt is Ca rich and Mn poor and displays a strong zonation: increasing  $X_{Ca}$  and decreasing  $X_{Mg}$ ,  $X_{Fe}$ , and  $X_{Mn}$  toward the rim, where grossular content can be as high as 20 mol% (Fig. 4). Both pre-tectonic and recrystallized biotites are homogeneous, with Mg/(Fe + Mg) ratios of 0.54–0.48 and Ti contents of 0.32–0.38 atoms per formula unit (afu). However, relict Bt armored in Pl is richer in Ti (0.49 afu) and Mg (Mg/[Fe+Mg] = 0.6). Plagioclase shows normal zoning, from  $Ab_{70}$  (cores) to  $Ab_{77}$  (rims). Two  $P$ - $T$  points were determined (Fig. 3), one for the migmatitic stage ( $686 \pm 40$  Mpa,  $790 \pm 17$  °C) and one for the stage of type III garnet growth ( $1140 \pm 135$  Mpa,  $615 \pm 70$  °C). In the first case Sil, core Pl, and Ilm along with Bt and Grt armored in Pl, and Kfs (completely retrogressed to Ms) were considered; in the second, Ky, rim Pl, Ms, Bt, Rt, and rim type III Grt were used (see footnote 1).

## DISCUSSION

The Ca-metapelite contains detrital zircon derived by erosion of Grenvillian basement and thus represents a sedimentary cover se-

quence to the basement of the Argentine Precordillera terrane. This cover could be equivalent to the lower part of the Precordillera sedimentary sequence or much older, i.e., Neoproterozoic. The absence of derived zircons between ca. 1100 Ma and 460 Ma shows that it underwent no significant tectonothermal event until the Middle Ordovician. This age gap is a characteristic of the Grenville province of Laurentia: in Gondwana (including the Eastern Sierras Pampeanas) a succession of Neoproterozoic events is recorded in zircon inheritance patterns (Ireland et al., 1994; Sims et al., 1998; Pankhurst et al., 2001). Evidence for true Grenvillian basement in the Sierra de Pie de Palo includes conventional U-Pb zircon data from metabasites of the lower ophiolitic unit (McDonough et al., 1993) and a Rb-Sr isochron age for orthogneiss of unknown structural position (Pankhurst and Rapela, 1998). Thus an unconformity must exist, masked by Famatinian deformation.

Because no pre-Ordovician metamorphism is registered in the zircons of sample SPP-1001, and the three *P-T* points determined in the equivalent sample SPP-882 suggest a single clockwise *P-T* path, we infer that this path results from Famatinian metamorphism alone. Metamorphic conditions are estimated as between high *P-T* type (no blueschists or eclogites of Famatinian age have been found in the sequence) and medium *P-T* (Barrovian) type. We envisage that peak pressure conditions in these rocks fell during postcollisional uplift on the mylonitic shear zones (pro-shears). In contrast, migmatite SPP-2057 shows two contrasting metamorphic histories: an older event under relatively low *P-T* type conditions, and a younger (synmylonitic) event comparable to the close to peak conditions of sample SPP-882. The first might equate to the Grenvillian  $M_2$  metamorphism in the Llano uplift area of Texas (Carlson and Schwarze, 1997; Mosher, 1998), which is considered to be the Laurentian counterpart of the Argentine Precordillera terrane (e.g., Dickerson and Keller, 1998). Thus we envisage that the migmatitic gneisses and the equivalent schist-quartzite sequence, together with some orthogneisses, represent the true Grenvillian basement.

Zircon overgrowths indicate metamorphic reactions involving the release of Zr from minerals such as amphibole and garnet (Fraser et al., 1997). Both minerals occur in the pre-mylonitic assemblage of SPP-1001, but were consumed during mylonitization. Thus we suggest that 460 Ma is the age of ductile

thrusting in the Sierra de Pie de Palo. Because pressure had by then dropped from its peak (Fig. 3), this must be taken as a minimum for the age of underthrusting of the Argentine Precordillera terrane beneath post-Cambrian Gondwana, i.e., the initiation of collision. Premylonitic foliations and prograde metamorphism shown by the Difunta Correa Sedimentary Sequence probably developed earlier during underthrusting, consistent with an early Middle Ordovician age for the drowning of the Precordillera carbonate platform (Astini et al., 1995).

The Sierra de Pie de Palo thus represents part of the eastern margin of the Argentine Precordillera terrane that was thoroughly involved in the tectonothermal activity of the Famatinian mobile belt. Because this part of the belt was thrust westward along the Pirquitas thrust, the Precordillera does not show the effects of Famatinian events to any significant extent. Our results support a Laurentian origin of the Argentine Precordillera terrane and Early to Middle Ordovician collision with Gondwana.

#### ACKNOWLEDGMENTS

This work was supported by grant PB97-1246, Ministerio de Educación y Cultura, Spain, and grant PICT98-4189, Argentina. It is a contribution to International Geological Correlation Program project 436 (Pacific Gondwana Margin). We thank C.M. Powell and C. Van Staal for helpful reviews.

#### REFERENCES CITED

- Aceñolaza, F.G., and Toselli, A.J., 1999, Argentine Precordillera: Allochthonous or autochthonous Gondwanic? *Zentralblatt für Geologie und Paläontologie*, Teil I, Hefte 7/8, p. 743-756.
- Astini, R.A., Benedetto, J.L., and Vaccari, N.E., 1995, The early Paleozoic evolution of the Argentina Precordillera as a Laurentian rifted, drifted and collided terrane: A geodynamic model. *Geological Society of America Bulletin*, v. 107, p. 253-273.
- Baldo, E., Casquet, C., and Galindo, C., 1998, Datos preliminares sobre el metamorfismo de la Sierra de Pie de Palo, Sierras Pampeanas Occidentales (Argentina). *Geogaceta*, v. 24, p. 39-42.
- Berman, R.G., 1991, Thermobarometry using multi-equilibrium calculations: A new technique, with petrological applications. *Canadian Mineralogist*, v. 29, p. 833-855.
- Carlson, W., and Schwarze, E., 1997, Petrological significance of prograde homogenization of growth zoning in garnet: An example from the Llano uplift. *Journal of Metamorphic Geology*, v. 15, p. 631-644.
- Compston, W., Williams, I.S., Kirschvink, J.L., Zhang Zichao, and Ma Guogan, 1992, Zircon U-Pb ages for the Early Cambrian time-scale. *Geological Society of London Journal*, v. 149, p. 171-184.
- Dalla Salda, L., Cingolani, C.A., and Varela, R., 1992, Early Paleozoic belt of the Andes and southwestern South America: Result of Laurentia-Gondwana collision? *Geology*, v. 20, p. 617-620.
- Dalziel, I.W.D., 1997, Overview: Neoproterozoic-Paleozoic geography and tectonics: Review hypothesis, environmental speculation. *Geological Society of America Bulletin*, v. 109, p. 16-42.
- Dickerson, P.W., and Keller, M., 1998, The Argentine Precordillera: Its odyssey from the Laurentian Ouachita margin towards the Sierras Pampeanas of Gondwana,

- in* Pankhurst, R.J., and Rapela, C.W., eds., The proto-Andean margin of Gondwana: Geological Society [London] Special Publication 142, p. 85-105.
- Fraser, G., Ellis, D., and Eggins, S., 1997, Zirconium abundance in granulite-facies minerals, with implications for zircon geochronology in high-grade rocks. *Geology*, v. 25, p. 607-610.
- Ireland, T.R., Bradshaw, J.D., Muir, R.J., Weaver, S.D., and Adams, C.J., 1994, Zircon age distributions in granites, graywackes, and gneisses from the southwest Pacific Gondwana region: U.S. Geological Survey Circular 1107, p. 151.
- Kay, S.M., Orrell, S., and Abruzzi, J.M., 1996, Zircon and whole-rock Nd-Pb isotopic evidence for a Grenville age and Laurentian origin for the basement of the Precordillera in Argentina. *Journal of Geology*, v. 104, p. 637-648.
- Keller, M., Buggisch, W., and Lehnert, O., 1998, The stratigraphical record of the Argentina Precordillera and its plate-tectonic background, *in* Pankhurst, R.J., and Rapela, C.W., eds., The proto-Andean margin of Gondwana: Geological Society [London] Special Publication 142, p. 35-56.
- Kretz, R., 1983, Symbols for rock-forming minerals. *American Mineralogist*, v. 68, p. 277-279.
- McDonough, M.R., Ramos, V.A., Isachsen, C.E., Bowring, S.A., and Vujovich, G.I., 1993, Edades preliminares de circones del basamento de la Sierra de Pie de Palo, Sierras Pampeanas occidentales de San Juan: Sus implicancias para el supercontinente proterozoico de Rodinia: XII Congreso Geológico Argentino y II Congreso de Exploración de Hidrocarburos, Mendoza, Actas, v. 3, p. 340-342.
- Mosher, S., 1998, Tectonic evolution of the southern Laurentian Grenville orogenic belt. *Geological Society of America Bulletin*, v. 110, p. 1357-1375.
- Pankhurst, R.J., and Rapela, C.W., 1998, The proto-Andean margin of Gondwana: An introduction, *in* Pankhurst, R.J., and Rapela, C.W., eds., The proto-Andean margin of Gondwana: Geological Society [London] Special Publication 142, p. 1-9.
- Pankhurst, R.J., Rapela, C.W., and Fanning, C.M., 2001, Age and origin of coeval TTG, I- and S-type granites in the Famatinian belt of NW Argentina. *Royal Society of Edinburgh Transactions, Earth Sciences* (in press).
- Ramos, V., Dallmeyer, R.D., and Vujovich, G., 1998, Time constraints on the early Palaeozoic docking of the Precordillera, central Argentina, *in* Pankhurst, R.J., and Rapela, C.W., eds., The proto-Andean margin of Gondwana: Geological Society [London] Special Publication 142, p. 143-158.
- Rapela, C.W., Pankhurst, R.J., Casquet, C., Baldo, E., Saavedra, J., and Galindo, C., 1998, Early evolution of the proto-Andean margin of South America. *Geology*, v. 26, p. 707-710.
- Sims, J.P., Ireland, T.R., Camacho, A., Lyons, P., Pieters, P.E., Skirrow, R.G., Stuart-Smith, P.G., and Miró, R., 1998, U-Pb, Th-Pb and Ar-Ar geochronology from the southern Sierras Pampeanas, Argentina: Implications for the Palaeozoic tectonic development of the western Gondwana margin, *in* Pankhurst, R.J., and Rapela, C.W., eds., The proto-Andean margin of Gondwana: Geological Society [London] Special Publication 142, p. 259-281.
- Spear, F.S., 1993, Metamorphic phase equilibria and pressure-temperature-time paths. *Mineralogical Society of America Monograph*, 799 p.
- Spear, F.S., Kohn, M.J., and Cheney, J.T., 1999, *P-T* paths from anatectic pelites: Contributions to Mineralogy and Petrology, v. 134, p. 17-32.
- Vujovich, G.I., and Kay, S.M., 1998, A Laurentian? Grenville-age oceanic arc/back-arc terrane in the Sierra de Pie de Palo, Western Sierras Pampeanas, Argentina, *in* Pankhurst, R.J., and Rapela, C.W., eds., The proto-Andean margin of Gondwana: Geological Society [London] Special Publication 142, p. 159-179.

Manuscript received December 18, 2000  
 Revised manuscript received April 6, 2001  
 Manuscript accepted April 24, 2001

Printed in USA