


RESEARCH

Open Access



# Effect of birthweight measurement quality improvement on low birthweight prevalence in rural Ethiopia

Estifanos Baye<sup>1</sup>, Firehiwot Workneh Abate<sup>2</sup>, Michelle Eglovitch<sup>1</sup>, Fisseha Shiferie<sup>2</sup>, Ingrid E. Olson<sup>1</sup>, Tigest Shifraw<sup>2</sup>, Workagegnehu Tarekegn Kidane<sup>2</sup>, Kalkidan Yibeltal<sup>2</sup>, Sitota Tsegaye<sup>2</sup>, Mulatu Melese Derebe<sup>3</sup>, Sheila Isanaka<sup>4</sup>, Blair J. Wylie<sup>5</sup>, Rose L. Molina<sup>5</sup>, Grace J. Chan<sup>6,7</sup>, Amare Worku<sup>2</sup>, Luke C. Mullany<sup>8</sup>, Alemayehu Worku<sup>2</sup>, Yemane Berhane<sup>2</sup> and Anne C. C. Lee<sup>1\*</sup> 

## Abstract

**Background:** Low birthweight (LBW) (< 2500 g) is a significant determinant of infant morbidity and mortality worldwide. In low-income settings, the quality of birthweight data suffers from measurement and recording errors, inconsistent data reporting systems, and missing data from non-facility births. This paper describes birthweight data quality and the prevalence of LBW before and after implementation of a birthweight quality improvement (QI) initiative in Amhara region, Ethiopia.

**Methods:** A comparative pre-post study was performed in selected rural health facilities located in West Gojjam and South Gondar zones. At baseline, a retrospective review of delivery records from February to May 2018 was performed in 14 health centers to collect birthweight data. A birthweight QI initiative was introduced in August 2019, which included provision of high-quality digital infant weight scales (precision 5 g), routine calibration, training in birth weighing and data recording, and routine field supervision. After the QI implementation, birthweight data were prospectively collected from late August to early September 2019, and December 2019 to June 2020. Data quality, as measured by heaping (weights at exact multiples of 500 g) and rounding to the nearest 100 g, and the prevalence of LBW were calculated before and after QI implementation.

**Results:** We retrospectively reviewed 1383 delivery records before the QI implementation and prospectively measured 1371 newborn weights after QI implementation. Heaping was most frequently observed at 3000 g and declined from 26% pre-initiative to 6.7% post-initiative. Heaping at 2500 g decreased from 5.4% pre-QI to 2.2% post-QI. The percentage of rounding to the nearest 100 g was reduced from 100% pre-initiative to 36.5% post-initiative. Before the QI initiative, the prevalence of recognized LBW was 2.2% (95% confidence interval [CI]: 1.5–3.1) and after the QI initiative increased to 11.7% (95% CI: 10.1–13.5).

**Conclusions:** A QI intervention can improve the quality of birthweight measurements, and data measurement quality may substantially affect estimates of LBW prevalence.

**Keywords:** Low birthweight, Delivery registers, Birthweight heaping, Data quality improvement initiative, Ethiopia

\*Correspondence: alee6@bwh.harvard.edu

<sup>1</sup> Department of Pediatric Newborn Medicine, Brigham and Women's Hospital, Harvard Medical School, 75 Francis Street, Boston, MA 02115, USA

Full list of author information is available at the end of the article

## Background

Worldwide, approximately 15% of live births (21 million) were low birthweight (LBW) in 2015; 91% from low- and middle-income countries (LMICs) [1]. LBW, commonly



© The Author(s) 2021. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

caused by preterm birth and/or intrauterine growth restriction [2], leads to a range of short- and long-term health impacts, including respiratory distress and feeding intolerance, growth impairment, developmental delay, and higher risk of diabetes and cardiovascular diseases [3]. LBW newborns are twenty times more likely to die in the first year than normal weight newborns [4].

Though recent estimates have shown that some progress has been made in reducing the risk of LBW [1], intensified efforts are needed to meet the World Health Assembly global target of decreasing the proportion of infants with LBW by 30% (~14 million) by the end of 2025. Ethiopia set a goal to reduce the prevalence of LBW to 7% by 2025 from 10.8% in 2012 [5]. Accurately measuring newborn weight at birth is crucial to provide special care for LBW infants, monitor the burden of LBW in the population, evaluate access to interventions aimed to improve antenatal care, and planning appropriate actions to accelerate the reduction of neonatal morbidity and mortality [6]. In LMICs, however, the quality of birthweight data suffers from measurement and recording errors, inconsistent data reporting systems, and missing data from non-facility births [1]. Improving measurement, recording, and reporting of birthweight are therefore warranted to target interventions and track progress toward the global nutrition target [7]. Strengthening the existing routine health systems of LMICs has been recommended as an essential strategy to improve birthweight data quality [8]. The *Every Newborn* Action Plan endorsed by the World Health Assembly prioritizes measurement improvement, with a focus on strengthening routine facility-based data, to track the national 2030 milestones ( $\leq 12$  neonatal deaths and  $\leq 12$  stillbirths per 1000 live births) [9]. In LMICs, health system strengthening measures that include training of healthcare staff and supportive supervision showed to improve health facility data quality [10, 11]. The Ethiopian National Newborn and Child Survival Strategy (2015/16–2019/20) aims to strengthen the existing health information management system for improving the percentage of live births with a reported birthweight from 5.2% in 2013 to 95% by 2020 [12]. Notwithstanding, the recent global and national LBW estimates conducted by Blencowe and colleagues could not provide an estimated LBW prevalence from Ethiopia, due to lack of adequate birthweight data quality [1].

The Enhancing Nutrition and Antenatal Infection Treatment (ENAT) study is a pragmatic effectiveness study testing the impact of optimizing prenatal nutrition status and infection control on birth outcomes in rural Amhara (ISRCTN15116516). Prior to the study, we introduced a birthweight quality improvement (QI) initiative in all study sites with the objective of improving

birthweight data quality. Herein, we present the impact of QI on birthweight data quality measures and the prevalence of LBW before and after the implementation of the initiative.

## Methods

### Study design and population

A comparative pre-post study was conducted to determine the effect of QI on birthweight data quality and the proportion of newborns with LBW in rural Amhara, Ethiopia. Amhara region is subdivided into 13 administrative zones and 140 districts, and it has an estimated population size of over 21,000,000. Nearly 85% of the population live in rural areas. According to the 2019 Ethiopian Mini Demographic and Health Survey (EMDHS) report, in Amhara region, the percentage of institutional delivery and those receiving antenatal care from a skilled provider were 54.2% and 82.6%, respectively [13]. Approximately 23% of women of reproductive age had a body mass index of below  $18.5 \text{ kg/m}^2$  [14].

### Study sites and periods

#### Phase 1 (pre-QI)

A retrospective facility record review was performed in 14 health centers located in seven districts of three zones in Amhara, namely South Gondar, West Gojjam and North Wollo, from February to May 2018 (Additional file 1). The health centers were chosen as prospective sites for the parent ENAT study, and the populations were selected based on high rates of maternal malnutrition, risk of fetal growth restriction, and need for nutrition interventions. Geomorphological maps were used to identify drought and famine-prone districts, and local government and partners were consulted. Two health centers from each district were selected based on the current antenatal care volume, using data from the region's Health Management and Information System.

#### Phase 2 (post-QI)

After selection of the health centers for the parent ENAT study within the zones above (South Gondar, West Gojjam), birthweight QI was implemented in 12 study health centers and birthweight data were prospectively collected from late August to early September 2019, and December 2019 to June 2020. These study health facilities were selected in consultation with the Amhara Regional Health Bureau, Amhara Public Health Institute, and partners, with preference for health centers with higher ANC volume and access to transportation (Additional file 1). The phase 2 study was conducted with the objective of not only improving the quality of anthropometric measurements and recording, but also for testing study tools and procedures of the ENAT parent study.

**Birthweight measurement QI initiative package (Fig. 1)**

**Equipment**

A high-quality digital infant scale (ADE-M112600, Germany) (precision 5 g) was provided to each study health center for measuring the newborn weight at birth. Adequate weighing stations were set up within all health centers; scales were placed on a clean flat surface with the display clearly visible and calibrated every morning with the 1000 g and 2000 g standard weights. If the weight did not read between  $\pm 10$  g of the standardized weights, proper adjustments with the placement of scales were made until the correct measures were displayed.

**Practical training**

Health center staff ( $n = 1$  to 4 midwives per health center) were trained on how to accurately measure birthweight using digital weight scales. Methods to weigh the baby with clothing using the tare function were developed, based on methods proposed by the International Fetal and Newborn Growth Consortium for the 21st Century (INTERGROWTH-21<sup>st</sup>) [15]. Where weight measurement of a naked newborn was not possible, a scale was first tared with a blanket or cloth, and weight of the baby with the clothing or blanket was taken. Training on recording the full precision of birthweight values, without any rounding to the nearest number was given to all personnel involved in measuring anthropometry. We developed job aids demonstrating how to properly measure and document birth weight using step-by-step pictorial procedures and posted them on the delivery room walls (Additional file 2).

**Supportive supervision**

Ongoing field supervision was performed in all health centers by the field team (study physicians and field coordinators) throughout the post-QI phase. To ensure the

standard operating procedures were followed, the field team observed the midwives measuring and recording the newborns’ weight, checked facility birthweight records (labor/delivery registers) for heaping and rounding, and provided feedback when appropriate. In addition, a review meeting with all health center midwives and directors was held after 2 weeks of implementing the initiative (phase 2 study). Quality of birthweight data extracted from health center records, lessons learnt and ways to forward were discussed.

**Data collection and management**


Birthweight data were extracted from delivery registers by trained research staff using the Survey Solutions<sup>®</sup> electronic data collection software (version 20.10, World Bank, Washington DC, USA). Routine field supervision was conducted in all sites, and collected data were reviewed and cleaned by the data management team daily.


**Retrospective delivery register data extraction**

During phase 1, about 1400 health center delivery registers were retrospectively reviewed. All labor/delivery registers recorded from February to May 2018 ( $n \sim 100$  records from each health center) were included. Birthweight was measured using fully and semi-functional analog/spring weighing scales available in the health center (Fig. 2), with routine delivery room measuring practices and record keeping.

**Prospective birthweight data collection**

In phase 2, health center midwives measured the weight of newborns at birth ( $n \sim 100$  per health center) using precise digital infant weight scales (Fig. 2). In line with the routine practice, health center midwives weighed the newborn only once and documented on labor/delivery registers.





**Fig. 2** Birthweight scales used before and after initiative. **A)** Birthweight scale commonly used pre-QI. **B)** Birthweight scale used post-QI

**Data analyses**

Birthweight data quality was assessed for implausibility, heaping and rounding. The presence of implausible values defined as extreme or unlikely birthweight values, i.e., < 350 g or > 6000 g, was checked. Percentages of heaping exactly at multiples of 500 g (e.g., 1500 g, 2000 g, 2500 g, 3000 g, 3500 g, etc.) were calculated. Heaping index (HI) was also computed by dividing the number of exact weight values (e.g., 3000 g) by all weights within the adjacent 250 g brackets, excluding the exact values (e.g., 2750–2999 + 3001–3249) [16]. Proportions of birthweights rounded to the nearest 100 g increments were calculated. Histograms were constructed to visually inspect the birthweight distribution. To calculate the prevalence of LBW, the number of live births with a birthweight of less than 2500 g was divided by the total number of liveborn babies with reported birthweights. To examine the effect of birthweight heaping, prevalence of LBW was computed by including 50% of newborns who had exactly 2500 g as LBW [16, 17]. Data analysis was done using STATA v.15 (StataCorp LLC, College Station, TX, USA).

**Results**

**Quality of birthweight data**

Birthweight data from 1383 and 1371 live births were collected from delivery registers in phase 1 (pre-QI) and phase 2 (post-QI) studies, respectively. There were no missing or implausible birthweights recorded at any phase. A decrease in heaping (i.e., percentages and indices of weights exactly at multiplies of 500 g) was noted across the study phases. The proportion of heaping at 3000 g was highest at 26% (HI of 0.78) before QI and dropped to 6.7% (HI 0.16) post-implementation. For the critical 2500 g at which LBW is defined, the percentage


of heaping decreased by half: 5.4% pre-QI and 2.2% post-QI. About 10% of newborns were reported to have exactly 3000 g weight before the initiative, and this was reduced to 2% after the initiative. The heaping index was highest at 1.0 for 1500 g pre-QI (but only 2 observations were recorded between 1250 and 1749 g) and reduced to 0.06 post-QI (Table 1). The percentage of rounding to the nearest 100 g improved from 100% pre-initiative to 36.5% post-initiative (Fig. 3).

**Prevalence of LBW before and after QI implementation**

From health center labor/delivery registers, 2.2% (95% CI: 1.5–3.1) of newborns weighed less than 2500 g at birth

**Table 1** Birthweight measurement quality before and after QI initiative in rural health centers in Ethiopia

	Pre-QI (phase 1) N = 1383	Post-QI (phase 2) N = 1371
<b>Birthweights recorded</b>		
<i>Heaping (500 g increments)</i>	n (%)	n (%)
1500 g	2 (0.14%)	1 (0.07%)
2000 g	3 (0.22%)	11 (0.80%)
2500 g	75 (5.42%)	31 (2.26%)
3000 g	360 (26.00%)	92 (6.70%)
3500 g	127 (9.18%)	30 (2.19%)
<i>Heaping index (ratio)</i>		
1500 g	1.00	0.06
2000 g	0.27	0.24
2500 g	0.78	0.12
3000 g	0.78	0.16
3500 g	0.89	0.13
<i>Rounding</i>	n (%)	n (%)
Rounding to nearest 100 g	1383 (100.00%)	500 (36.50%)



pre-QI, and the percentage increased to 11.7% (95% CI: 10.1–13.5) post-QI. When half of newborns who weighed exactly 2500 g were included in the LBW calculation, the LBW rate increased to 4.9% (95% CI: 3.8–6.1) before initiative and 12.9% (95% CI: 11.1–14.7) after initiative. The proportion of newborns with 2000–2500 g also rose from 1.6% pre-QI to 9.3% post-QI (Table 2).

**Discussion**

Improving quality of birthweight data is essential for identifying infants at high risk for medical and neurodevelopmental problems, determining the burden of LBW, and monitoring access and usage of interventions aimed to enhance prenatal care and newborn survival. We observed a reduction in percentages of heaping and rounding and an increase in the proportion of LBW by about 10 percentage points after the implementation of birthweight QI initiative (provision of high-precision digital weight scales, practical training and supportive supervision).

We found that introduction of QI decreased birthweight heaping exactly at multiples of 500 g (e.g., 1500 g,

2000 g, 2500 g, 3000 g, 3500 g, etc.). Heaping, commonly observed when scales with low precision are used or continuous data are rounded, represents misclassification [18]. A recent study conducted in Bangladesh, Nepal and Tanzania showed significant heaping (19–67%) at 2500 g and 3000 g in health facilities [16]. The amount of heaping on 2500 g can have a significant effect on LBW estimation. Some newborns whose birthweights rounded to exactly 2500 g may have been lighter and should actually be classified in the LBW category, while some may have been heavier. The effect of this type of misclassification is critical in determining the proportion of LBW [18]. In this study, although pre- and post-initiative data appear symmetrical, the highest percentage of heaping observed at 3000 g (26%) pre-QI declined by three-quarters after the initiative. Heaping exactly at 2500 g also reduced from 5.4 pre-QI to 2.2% post-QI. Furthermore, 100% of data rounded to the nearest 100 g pre-initiative dropped by two-thirds post-initiative. In South Africa, a similar data improvement intervention, including training on data collection and providing feedback to healthcare staff with monthly data reviews and audits, increased the

**Table 2** Birthweight distribution before and after QI initiative in rural health centers in Ethiopia

Study design	Pre-QI (phase 1) Retrospective	Post-QI (phase 2) Prospective	Absolute difference
Birthweights recorded	N = 1383	N = 1371	
Birth weight	% (95% CI)	% (95% CI)	% (95% CI)
Mean (SD)	3083 (408)	2977 (463)	-106 (-138.6, -73.4)
< 2500 g	2.16 (1.5, 3.1)	11.74 (10.1, 13.5)	9.58 (7.7, 11.5)
≤ 2500 g	4.92 (3.8, 6.1)	12.91 (11.1, 14.7)	7.99 (5.9, 10.1)
2000–2500 g	1.59 (0.1, 2.2)	9.26 (7.7, 10.8)	7.67 (5.9, 9.4)
1500–2000 g	0.29 (0.01, 0.5)	1.82 (1.1, 2.3)	1.53 (0.8, 2.2)
< 1500 g	0.29 (0.01, 0.5)	0.66 (0.2, 1.1)	0.37 (-0.1, 0.9)

accuracy and completeness of health facility data [10]. Health system strengthening measures such as performance review feedback activities and enhanced supervision have also shown to increase ownership of data among healthcare workers in Rwanda [11]. The tendency to round birthweight may be due to the fact that there are no formal standards in recording birthweight within the health systems [19], low precision and common use of spring birthweight scales [20], and/or high workload of healthcare providers. Proper recording of birthweight data on health facility records, which can be used as a source of data for regional and national estimates, may improve the quality of birthweight data and decrease the need for statistical adjustments [1].

Based on data from pre-initiative phase, the percentage of newborns with LBW was about 2%. However, the prevalence increased to 12% after birthweight QI implementation, which was more consistent with a meta-analysis that included 4105 participants from nine observational studies showing 16% of newborns were LBW in Amhara region [21]. The 2016 Ethiopian Demographic and Health Survey report indicated a LBW prevalence of 22% in Amhara region, which was the highest nationally [14]. Recent studies performed at Debre Tabor hospital in South Gondar zone, and at Dangla Primary hospital in West Gojjam zone found a LBW prevalence of 10–12% among facility births [22, 23]. Estimation of the percentage of newborns with LBW is dependent on how newborns with a reported birthweight of exactly 2500 g are classified [19]. Data from LMICs showed an increase in the prevalence of LBW from 1.7% to 7.2% after re-allocating 50% of newborns with a birthweight of exactly 2500 g to the LBW category [16]. In our study, when we included half of newborns with exactly 2500 g as LBW in the calculation, the percentage increased by 2.7% pre-QI but declined to 1.2% after the initiative. In addition to recording errors, the change in estimated LBW rates due to differences in instrument precision is also possible [17]. A recent study reported lower heaping indices using digital scales compared to analog [24]. Compared to the traditional method focusing on control, audit and examination, the new approach of supportive supervision with a focus on strengthening routine health system, problem solving and training healthcare providers shows superior results in improving essential newborn care [25]. Our findings suggest that provision of digital weight scales together with periodic supervision may help to improve birthweight data quality and estimates of the true burden of LBW.

The present study has some limitations. Our findings may not be truly generalizable because women who give birth at home may differ in health and socioeconomic status from those who give birth in facilities, which may

lead to different rates of LBW. Since the main aim was to improve quality of birthweight data in study health centers prior to commencing the parent ENAT study, we did not conduct proper sample size calculation and collect birthweight data from all health centers or home births throughout the study phases. Stillbirths were not routinely measured or reported and hence not included in the analysis. Given that the trial was underway, we did not repeat observations after an interval of time from the initial QI initiative, hence we could not assess whether the improvements were sustained. Despite these limitations, we were able to assess birthweight within hours of birth, prior to significant weight loss, using high-precision digital weight scales at the health facilities.

## Conclusions

Implementation of a birthweight QI program improved data quality in rural Amhara with decreased percentages of heaping and rounding on health center records. These measurement improvements resulted in an increased prevalence of LBW from 2.2 to 11.7%. Availing precise digital weight scales, improving BW recording, and birth weighing guidelines along with supportive supervision may be essential to improve birthweight data quality in low-resource settings. Further implementation research is needed to assess the effectiveness and sustainability of the initiative on a larger-scale and estimate cost implications.

## Abbreviations

CI: Confidence intervals; EMDHS: Ethiopian Mini Demographic and Health Survey; ENAT Study: Enhancing Nutrition and Antenatal Infection Treatment Study; HI: Heaping Index; INTERGROWTH-21<sup>ST</sup>: International Fetal and Newborn Growth Consortium for the 21st Century; LBW: Low birthweight; LMICs: Low- and middle-income countries; QI: Quality improvement.

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12963-021-00265-0>.

**Additional file 1.** Health centers selected for Phase 1&2 studies (pre- and post-QI).

**Additional file 2.** Job aids posted on the delivery room walls.

## Acknowledgements

We would like to acknowledge the Amhara Regional Health Bureau, Amhara Public Health Institute and health center directors for their collaboration in implementing the study. We thank the ENAT field team (study nurses and data collectors) and health center midwives for their hard work and dedication to this study.

## Authors' contributions

EB participated in data acquisition, conducted data analyses, interpreted results and wrote the manuscript. FWA, ME, FS, IO, TS, WTK, KY, ST, MMD, AZ and AW participated in data acquisition and critical review of the manuscript. SI, BW, RM, GC, LM, AW and YB provided input on data acquisition and analysis and critical review of the manuscript. ACL conceived the research questions

and provided guidance on data acquisition and analysis and interpreting results, and critical review of the manuscript. All authors read and approved the final manuscript.

#### Funding

The ENAT study was funded by the Bill and Melinda Gates Foundation (OPP1184363).

#### Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

#### Declarations

##### Ethics approval and consent to participate

The study was reviewed by Institutional Review Boards of Mass General Brigham (USA), Addis Continental Institute of Public Health and Amhara Public Health Institute (Ethiopia), and by the National Research Ethics Review Committee (Ethiopia). Support letters from Amhara Public Health Institute, Zonal and District Health Bureaus and approval from health center directors were obtained prior to the study start. Data were extracted from health center records without personal identifiers.

##### Consent for publication

Not required.

##### Competing interests

The authors declare that they have no competing interests.

##### Author details

<sup>1</sup>Department of Pediatric Newborn Medicine, Brigham and Women's Hospital, Harvard Medical School, 75 Francis Street, Boston, MA 02115, USA. <sup>2</sup>Addis Continental Institute of Public Health, Addis Ababa, Ethiopia. <sup>3</sup>Amhara Public Health Institute, Bahir Dar, Ethiopia. <sup>4</sup>Departments of Nutrition and Global Health and Population, Harvard T.H. Chan School of Public Health, Boston, MA, USA. <sup>5</sup>Department of Obstetrics and Gynecology, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA, USA. <sup>6</sup>Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA, USA. <sup>7</sup>Department of Pediatrics, Harvard Medical School, Boston, MA, USA. <sup>8</sup>Department of International Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA.

Received: 23 June 2021 Accepted: 1 September 2021

Published online: 22 September 2021

#### References

- Blencowe H, Krusevec J, de Onis M, Black RE, An X, Stevens GA, Borghi E, Hayashi C, Estevez D, Cegolon L, et al. National, regional, and worldwide estimates of low birthweight in 2015, with trends from 2000: a systematic analysis. *Lancet Glob Health*. 2019;7:e849–60.
- Katz J, Lee AC, Kozuki N, Lawn JE, Cousens S, Blencowe H, Ezzati M, Bhutta ZA, Marchant T, Willey BA, et al. Mortality risk in preterm and small-for-gestational-age infants in low-income and middle-income countries: a pooled country analysis. *Lancet*. 2013;382:417–25.
- Zerbetto AB, Cortelo FM, Filho ÉBC. Association between gestational age and birth weight on the language development of Brazilian children: a systematic review. *J Pediatr (Rio J)*. 2015;91:326–32.
- UNICEF, WHO. Low birth weight: country, regional and global estimates. New York: UNICEF; 2004.
- Hailu I, Kidane A, Kovach T, Lemma F, Sethuraman K, Sommerfelt, Elisabeth A, Tekla T. A tool to support nutrition advocacy in Ethiopia: Ethiopia PROFILES 2012 estimates. Washington, DC, and Addis Ababa, Ethiopia: Food and Nutrition Technical Assistance III Project (FANTA)/FHI 360 and Federal Ministry of Health, Ethiopia; 2013.
- Chang KT, Mullany LC, Khatri SK, LeClerq SC, Munos MK, Katz J. Validation of maternal reports for low birthweight and preterm birth indicators in rural Nepal. *J Glob Health*. 2018;8:010604.
- Moxon SG, Ruysen H, Kerber KJ, Amouzou A, Fournier S, Grove J, Moran AC, Vaz LME, Blencowe H, Conroy N, et al. Count every newborn; a

- measurement improvement roadmap for coverage data. *BMC Pregnancy Childbirth*. 2015;15:S8.
- Doherty T, Kinney M. Low birthweight: will new estimates accelerate progress? *Lancet Glob Health*. 2019;7:e809–10.
- Moxon SG, Ruysen H, Kerber KJ, Amouzou A, Fournier S, Grove J, Moran AC, Vaz LM, Blencowe H, Conroy N, et al. Count every newborn; a measurement improvement roadmap for coverage data. *BMC Pregnancy Childbirth*. 2015;15(Suppl 2):S8.
- Mphatswe W, Mate KS, Bennett B, Ngidi H, Reddy J, Barker PM, Rollins N. Improving public health information: a data quality intervention in KwaZulu-Natal, South Africa. *Bull World Health Organ*. 2012;90:176–82.
- Wagenaar BH, Hirschhorn LR, Henley C, Gremu A, Sindano N, Chilengi R, Hingora A, Mboya D, Exavery A, Tani K, et al. Data-driven quality improvement in low-and middle-income country health systems: lessons from seven years of implementation experience across Mozambique, Rwanda, and Zambia. *BMC Health Serv Res*. 2017;17:830.
- Federal Ministry of Health, Maternal and Child Health Directorate. National Strategy for Newborn and Child Survival in Ethiopia: 2015/16–2019/20. Addis Ababa, Ethiopia. 2015.
- Ethiopian Public Health Institute (EPHI) [Ethiopia] and ICF. Ethiopia mini demographic and health survey 2019: key indicators. Rockville, MD: EPHI and ICF; 2019.
- Central statistical agency of Ethiopia and ICF. Ethiopia demographic and health survey 2016. Addis Ababa: CSA and ICF; 2016.
- Villar J, Cheikh Ismail L, Victora CG, Ohuma EO, Bertino E, Altman DG, Lambert A, Papageorgiou AT, Carvalho M, Jaffer YA, et al. International standards for newborn weight, length, and head circumference by gestational age and sex: the Newborn Cross-Sectional Study of the INTERGROWTH-21st Project. *Lancet*. 2014;384:857–68.
- Day LT, Gore-Langton GR, Rahman AE, Basnet O, Shabani J, Tahsina T, Poudel A, Shirima K, Ameen S, Ashish KC, et al. Labour and delivery ward register data availability, quality, and utility—Every Newborn-birth indicators research tracking in hospitals (EN-BIRTH) study baseline analysis in three countries. *BMC Health Serv Res*. 2020;20:737.
- Mullany LC, Darmstadt GL, Katz J, Khatri SK, Tielsch JM. Effect of instrument precision on estimation of low birth weight prevalence. *J Perinatol*. 2005;25:11–3.
- Blanc AK, Wardlaw T. Monitoring low birth weight: an evaluation of international estimates and an updated estimation procedure. *Bull World Health Organ*. 2005;83:178–85.
- Channon AA, Padmadas SS, McDonald JW. Measuring birth weight in developing countries: does the method of reporting in retrospective surveys matter? *Matern Child Health J*. 2011;15:12–8.
- Rekha C, Whelan RM, Reddy P, Reddy PS. Evaluation of adjustment methods used to determine prevalence of low birth-weight babies at a rural hospital in Andhra Pradesh. *India Indian J Public Health*. 2013;57:177–80.
- Endalamaw A, Engeda EH, Ekubagewargies DT, Belay GM, Tefera MA. Low birth weight and its associated factors in Ethiopia: a systematic review and meta-analysis. *Ital J Pediatr*. 2018;44:141.
- Talie A, Taddele M, Alemayehu M. Magnitude of low birth weight and associated factors among newborns delivered in Dangla Primary Hospital, Amhara Regional State, Northwest Ethiopia, 2017. *J Pregnancy*. 2019;2019:3587239.
- Mekie M, Taklual W. Magnitude of low birth weight and maternal risk factors among women who delivered in Debre Tabor Hospital, Amhara Region, Ethiopia: a facility based cross-sectional study. *Ital J Pediatr*. 2019;45:86.
- Kong S, Day LT, Zaman SB, Peven K, Salim N, Sunny AK, Shamba D, Rahman QS-u, Ashish KC, Ruysen H, et al. Birthweight: EN-BIRTH multi-country validation study. *BMC Pregnancy Childbirth*. 2021;21:240.
- Gupta R. Regular Appraisal of Program Implementation in District (RAPID): a supportive supervision approach to improve essential newborn care in Haryana India. *Int J Integr Care*. 2014;14:1. <https://doi.org/10.5334/ijic.1653>.

#### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

[onlineservice@springernature.com](mailto:onlineservice@springernature.com)