

Synthesis of the C1–C27 Fragment of Stambomycin D Validates Modular Polyketide Synthase-Based Stereochemical Assignments

Jieyan Lim,[‡] Venkaiah Chintalapudi,[‡] Haraldur G. Gudmundsson, Minh Tran, Alice Bernasconi, Araceli Blanco, Lijiang Song, Gregory L. Challis, and Edward A. Anderson*



Cite This: *Org. Lett.* 2021, 23, 7439–7444



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information


ABSTRACT: The stambomycins are a family of bioactive macrolides isolated from *Streptomyces ambofaciens*. Aside from two stereocenters installed through cytochrome P450 oxidations, their stereochemistry has been predicted by sequence analysis of the polyketide synthase. We report a synthesis of the C1–C27 fragment of stambomycin D, the spectroscopic data of which correlates well with that of the natural product, further validating predictive sequence analysis as a powerful tool for stereochemical assignment of complex polyketide natural products.



Stereochemical determination is a key element in natural product discovery, as bioactivity is often intrinsically linked to stereochemistry. It can also be one of the most challenging aspects, especially for polyketides where conformational flexibility and noncrystallinity render conclusive assignment challenging using NMR-based methods or X-ray crystallography.^{1–3} Computational approaches using NMR parameters are emerging as reliable tools^{4,5} but are unsuitable when individual stereoclusters are “insulated” by regions of flexible nonfunctionalized carbon chains, or by rigid (poly)alkene regions. NMR spectroscopy can equally be ambiguous for certain stereoclusters/conformations or complicated by overlapping signals in more complex natural products, rendering the extraction of coupling constants or nOes highly challenging and ultimately not definitive. This uncertainty provides a significant obstacle for synthesis and applications.⁶

Advances in bioinformatics have enabled the application of predictive sequence analysis of biosynthetic enzymes not only in the discovery of natural products but also in their structural and stereochemical determination.^{7–11} One example is the stambomycins, a family of 51-membered glycosylated macrolides discovered by Challis, Aigle, and co-workers in 2011 (Figure 1a).¹² These were identified as the metabolic products of a modular polyketide synthase (PKS) in *Streptomyces ambofaciens* through a genomics-driven approach involving rational genetic manipulation to induce transcription of the biosynthetic genes, which are poorly expressed in laboratory cultures. Four members of the family (A–D) were identified, differing at the C26 side chain, all of which showed potent antibacterial and antitumor activity. The planar structures and stereochemistry of the stambomycins were predicted via sequence analysis of the modular PKS responsible for their biosynthesis,^{12–14} with the exception of the C28 and C50 stereocenters, which are of non-PKS origin.¹⁵ Notably, the

a Genome sequence-predicted stereochemistry of the stambomycins



b Our target: C1–C27 fragment of the stambomycin D aglycon




Figure 1. (a) Stambomycins A–D and (b) the C1–C27 fragment and planned retrosynthesis.

Received: August 9, 2021

Published: September 8, 2021



stambomycins are one of the earliest structurally complex polyketides for which predictive sequence analysis was employed for stereochemical assignment, and remain one of the most elaborate examples to date.¹⁶


While the predicted planar structures of the stambomycins have been confirmed by NMR spectroscopy, their stereochemistry remains to be unequivocally confirmed. This inspired our interest in stambomycin D: a synthesis of this molecule would represent a powerful validation of sequence-based polyketide stereochemical assignment, and one that could offer a rapid and complementary approach to traditional NMR-based methods. Here, we report the synthesis of the C1–C27 aglycon fragment of stambomycin D and its comparison with the natural macrocycle. The excellent agreement between the synthetic and natural material supports the sequence-based stereochemical assignment in this region.

We envisioned that the northern and southern hemispheres of the stambomycin D aglycon would make ideal targets to establish a synthetic strategy for the entire molecule and allow a preliminary comparison of NMR data to support the predicted stereochemistry. To avoid the uncertainty of the C28 stereocenter, our initial target for the northern hemisphere consisted of the C1–C27 fragment **1** (Figure 1b). Retrosynthetically, **1** could be disconnected at the C11–C12 bond to reveal C1–C11 alkenyl iodide **2**, which could be coupled to a vinyl organometallic at C12, for example by Suzuki coupling. Disconnection at the C22–C23 bond reveals C13–C22 fragment **3** (in which the required boronic ester could be derived from manipulation of the ester group) and C23–C27 fragment **4**. Union of the latter two fragments could be achieved by asymmetric alkyne addition of **3** to **4**, followed by Hoveyda hydroboration/oxidation¹⁷ and reduction of the resulting propargylic alcohol to install the desired 1,3-*anti*-diol at C21/C23.

Synthesis of the C1–C11 fragment **2** (Scheme 1) commenced with an enantio- and diastereoselective Leighton crotylation¹⁸ of aldehyde **5** with *cis*-crotyltrichlorosilane **7** to give homoallylic alcohol **8** in 87% yield (89% *ee*, 15:1 *dr*). Cross metathesis of **8** with methyl acrylate afforded α,β -unsaturated ester **9** (90%), which was subjected to Evans–Prunet acetalization¹⁹ to obtain acetal **10** in 33% yield. Formation of this acetal appeared to be in an unfavorable equilibrium with the retro-Michael reaction, as the cyclization failed to reach completion even with extended reaction times; interestingly, the recovered starting material bore mainly a *Z*-alkene. This problem is attributed to the presence of the C8 (*R*)-methyl group, which must adopt an axial position in the six-membered cyclic acetal. Following a DIBALH reduction of the ester in **10**, a second Leighton crotylation was carried out on the resulting aldehyde, giving homoallylic alcohol **11** in 70% yield (10:1 *dr*). Protection of the alcohol as the PMB ether and subsequent oxidative cleavage of the terminal alkene afforded aldehyde **12**. A Mukaiyama aldol reaction of **12** with silyl ketene acetal **13** then gave the corresponding β -hydroxy ester (85%), which after oxidation of the alcohol, furnished β -keto ester **14** in 87% yield.

We expected that deprotection of the acetal under acidic conditions would also promote spontaneous cyclization of the resulting C7 hydroxyl group onto the C3 ketone to form the desired tetrahydropyran. This step proved unexpectedly challenging as the acetal was surprisingly robust; conditions that allowed for full conversion of the starting material also resulted in significant degradation and the formation of an

Scheme 1. Synthesis of C1–C11 Fragment **2**




unidentified side product which was difficult to separate from the product **15**. Various deprotection conditions were tested to achieve an optimal balance between conversion of the starting material and product formation, most of which involved different concentrations of aqueous HCl in MeOH/THF, as this acid was observed to give a relatively clean reaction. After fine-tuning the solvent ratio, temperature, and reaction time, it was found that use of 1.0 M aqueous HCl in MeOH/THF (1:1) at 35 °C for 22 h gave the desired tetrahydropyran **15** in 47% yield, with 46% recovered starting material. NOESY correlations and coupling constant analysis confirmed the relative stereochemistry of the various substituents on the 6-membered ring. Finally, PMB deprotection afforded the C1–C11 fragment **2** in 44% yield (12 steps from **5**).

Synthesis of the C13–C22 fragment **3** (Scheme 2a) began with 5-hexynal **16**. A Leighton crotylation¹⁸ was again employed to set the two adjacent stereocenters in homoallylic alcohol **17** (82%, 92% *ee*, > 20:1 *dr*). Adopting a similar strategy to that used for fragment **2**, alcohol **17** was protected as the PMB ether, with subsequent oxidative cleavage of the terminal alkene affording aldehyde **18**. A Mukaiyama aldol reaction of **18** with silyl ketene acetal **13** gave β -hydroxy ester **19** in 86% yield (5:1 *dr*); the stereochemistry of the alcohol was confirmed by Mosher ester analysis.²⁰ Finally, treatment of **19** with DDQ under anhydrous conditions resulted in the formation of the 1,3-PMP acetal, giving C13–C22 fragment **3** in 55% yield after removal of the minor diastereomer (6 steps from **16**).

Attention now turned to the construction of the C23–C27 aldehyde **4** (Scheme 2b). To install the hexyl-bearing stereocenter in this fragment, an enantioselective organocatalytic aldol reaction²¹ of octanal and formaldehyde was employed at the outset. This gave a lactol intermediate, which was subjected to a Wittig olefination to obtain enoate **21** in

Scheme 2. (a) Synthesis of C13–C22 Fragment 3 and (b) Synthesis of C23–C27 Fragment 4



67% yield (88% *ee*). Protection of the alcohol as the TBS ether (77%) and DIBALH reduction of the ester (82%) gave alcohol 22. Oxidation of alcohol 22 then afforded the C23–C27 aldehyde 4 in 93% yield.


With fragments 2–4 in hand, we proceeded to combine them toward the full C1–C27 fragment 1 (Scheme 3). First, a diastereoselective alkynylzinc addition²² of 3 to 4 afforded propargylic alcohol 23 in 62% yield (11:1 *dr*), with Mosher ester analysis confirming the stereochemistry of the alcohol. Hydroboration/oxidation of 23 employing a modification²³ of Hoveyda's conditions¹⁷ gave β -hydroxy ketone 24 in 80%

yield. Following an Evans–Saksena reduction²⁴ of the β -hydroxy ketone (>20:1 *dr*), the resulting 1,3-*anti*-diol was protected as the acetonide (25), which moreover served to confirm its stereochemistry through the Rychnovsky method.²⁵ A DIBALH reduction of the ester in 25 afforded the aldehyde, which was then alkynylated using the Ohira–Bestmann reagent. During alkynylation, it was observed that the PMP acetal was prone to ring-opening, presumably via enolization of the adjacent aldehyde under the mildly basic reaction conditions (K_2CO_3). This resulted in the formation of a side product which not only lowered the yield of the alkyne (26) but also led to problems with purification. It was eventually found that use of an excess of the Ohira–Bestmann reagent overcame this problem, enabling alkyne 26 to be obtained in 81% yield.

Following Zr-mediated hydroboration²⁶ of alkyne 26, the resulting vinylboronic ester 27 was coupled with C1–C11 fragment 2 via a Suzuki coupling. A variety of reaction conditions were screened, but use of Tl_2CO_3 ²⁷ was found to be essential for reaction success, giving the complete C1–C27 framework 28 in 53% yield. Deprotection of 28 proved nontrivial, as the C10–C13 1,3-diene was observed to be highly acid-sensitive and prone to degradation, potentially via acid-promoted cyclization of the C17 hydroxyl group. After much experimentation, we found that deprotection could be achieved using 0.1 M aqueous HCl in MeOH/THF without degradation of the diene. Lower acid concentrations of 0.05 and 0.02 M could also be used, although longer reaction times were required. Treatment of 28 with 0.1 M aqueous HCl in MeOH/THF for three hours at room temperature thus afforded C1–C27 fragment 1 in 41% yield.

Having obtained C1–C27 fragment 1, we were inspired to compare its NMR spectra with the corresponding NMR data of stambomycin D. To our delight, the data for 1 showed an excellent match with the reported¹² data for stambomycin D (Figure 2 and Supporting Information). Although slight discrepancies existed, this is not unexpected due to potential conformational differences between the acyclic fragment and

Scheme 3. Completion of C1–C27 Fragment 1






Figure 2. Comparison of (a) ^1H NMR and (b) ^{13}C NMR data of stambomycin D and C1–C27 fragment 1.

the cyclic macrolide. For example, the acyclic fragment contains a free hydroxyl at C5, whereas in the macrolide this oxygen atom is attached to the amino sugar mycaminose; in addition, the acyclic fragment is truncated at C27, as compared to the macrolide. These differences were therefore reflected in discrepancies in the ^{13}C NMR data of C5 and C27. An additional discrepancy was noted at the C19 protons; re-examination of the spectroscopic data for the natural product confirmed these signals should be reassigned. Overall, there is good agreement in the ^1H and ^{13}C NMR data between the C1–C27 fragment and stambomycin D, supporting the stereochemical assignment of this region of the natural product.

In summary, we have synthesized the C1–C27 “northern” fragment of the stambomycin D aglycon. Comparison of NMR data of this fragment with the reported data of stambomycin D showed good agreement between the two, providing preliminary proof of the accuracy of the sequence-based stereochemical assignment of the macrolide.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.1c02650>.

Experimental procedures and characterization data for novel compounds (PDF)

■ AUTHOR INFORMATION

Corresponding Author

Edward A. Anderson – Chemistry Research Laboratory, University of Oxford, Oxford OX1 3TA, U.K.; orcid.org/0000-0002-4149-0494; Email: edward.anderson@chem.ox.ac.uk

Authors

Jieyan Lim – Chemistry Research Laboratory, University of Oxford, Oxford OX1 3TA, U.K.

Venkaiah Chintalapudi – Chemistry Research Laboratory, University of Oxford, Oxford OX1 3TA, U.K.

Haraldur G. Gudmundsson – Chemistry Research Laboratory, University of Oxford, Oxford OX1 3TA, U.K.

Minh Tran – Chemistry Research Laboratory, University of Oxford, Oxford OX1 3TA, U.K.

Alice Bernasconi – Sezione Chimica Generale e Organica “A. Marchesini”, Università degli Studi di Milano, 20133 Milano, Italy

Araceli Blanco – Departamento de Química Orgánica, Facultad de Ciencias Químicas, Universidad de Salamanca, 37008 Salamanca, Spain

Lijiang Song – Department of Chemistry and Warwick Integrative Synthetic Biology Centre, University of Warwick, Coventry CV4 7AL, U.K.

Gregory L. Challis – Department of Chemistry and Warwick Integrative Synthetic Biology Centre, University of Warwick, Coventry CV4 7AL, U.K.; Department of Biochemistry and Molecular Biology and ARC Centre of Excellence for Innovations in Peptide and Protein Science, Biomedicine Discovery Institute, Monash University, Clayton, Victoria 3800, Australia; orcid.org/0000-0001-5976-3545

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/acs.orglett.1c02650>

Author Contributions

\ddagger J.L. and V.C. contributed equally to this work.

Notes

The authors declare the following competing financial interest(s): G.L.C. is a nonexecutive codirector of a Erebagen, Ltd. The other authors declare no competing interests.

ACKNOWLEDGMENTS

We thank the Agency for Science, Technology and Research (A*STAR) for a National Science Scholarship (J.L.), the Marie Skłodowska-Curie actions for an Individual Fellowship (V.C., GA No 702385), and the Leverhulme Trust for funding (H.G.G., RPG-2015-003). We are grateful to Jonathan Yong (University of Oxford) for his help in acquiring NMR spectra of **1**. E.A.A. thanks the EPSRC for additional support (EP/S013172/1).

REFERENCES

- (1) Matsumori, N.; Kaneno, D.; Murata, M.; Nakamura, H.; Tachibana, K. Stereochemical Determination of Acyclic Structures Based on Carbon-Proton Spin-Coupling Constants. A Method of Configuration Analysis for Natural Products. *J. Org. Chem.* **1999**, *64*, 866–876.
- (2) (a) Kobayashi, Y.; Lee, J.; Tezuka, K.; Kishi, Y. Toward Creation of a Universal NMR Database for the Stereochemical Assignment of Acyclic Compounds: The Case of Two Contiguous Propionate Units. *Org. Lett.* **1999**, *1*, 2177–2180. (b) Lee, J.; Kobayashi, Y.; Tezuka, K.; Kishi, Y. Toward Creation of a Universal NMR Database for the Stereochemical Assignment of Acyclic Compounds: Proof of Concept. *Org. Lett.* **1999**, *1*, 2181–2184. (c) Kobayashi, Y.; Hayashi, N.; Tan, C.-H.; Kishi, Y. Toward the Creation of NMR Databases in Chiral Solvents for Assignments of Relative and Absolute Stereochemistry: Proof of Concept. *Org. Lett.* **2001**, *3*, 2245–2248. (d) Fleury, E.; Lannou, M.-I.; Bistri, O.; Sautel, F.; Massiot, G.; Pancrazi, A.; Ardisson, J. Advances in the Universal NMR Database: Toward the Determination of the Relative Configurations of Large Polypropionates. *Eur. J. Org. Chem.* **2009**, *2009*, 4992–5001.
- (3) For reviews on NMR and related methods for stereochemical determination, see: (a) Menche, D. New methods for stereochemical determination of complex polyketides: configurational assignment of novel metabolites from myxobacteria. *Nat. Prod. Rep.* **2008**, *25*, 905–918. (b) Menna, M.; Imperatore, C.; Mangoni, A.; Della Sala, G.; Tagliatalata-Scafati, O. Challenges in the configuration assignment of natural products. A case-selective perspective. *Nat. Prod. Rep.* **2019**, *36*, 476–489.
- (4) (a) Smith, S. G.; Goodman, J. M. Assigning the Stereochemistry of Pairs of Diastereoisomers Using GIAO NMR Shift Calculation. *J. Org. Chem.* **2009**, *74*, 4597–4607. (b) Smith, S. G.; Goodman, J. M. Assigning Stereochemistry to Single Diastereoisomers by GIAO NMR Calculation: The DP4 Probability. *J. Am. Chem. Soc.* **2010**, *132*, 12946–1259. (c) Zanardi, M. M.; Sarotti, A. M. GIAO C–H COSY Simulations Merged with Artificial Neural Networks Pattern Recognition Analysis. Pushing the Structural Validation a Step Forward. *J. Org. Chem.* **2015**, *80*, 9371–9378. (d) Ermanis, K.; Parkes, K. E. B.; Agback, T.; Goodman, J. M. The optimal DFT approach in DP4 NMR structure analysis – pushing the limits of relative configuration elucidation. *Org. Biomol. Chem.* **2019**, *17*, 5886–5890.
- (5) (a) Marcarino, M. O.; Zanardi, M. M.; Cicetti, S.; Sarotti, A. M. NMR Calculations with Quantum Methods: Development of New Tools for Structural Elucidation and Beyond. *Acc. Chem. Res.* **2020**, *53*, 1922–1932. (b) Marcarino, M. O.; Cicetti, S.; Zanardi, M. M.; Sarotti, A. M. A critical review on the use of DP4+ in the structural elucidation of natural products: the good, the bad and the ugly. A practical guide. *Nat. Prod. Rep.* **2021**, DOI: 10.1039/D1NP00030F.
- (6) For examples, see: (a) Heinrich, M.; Murphy, J. J.; Ilg, M. K.; Letort, A.; Flasz, J.; Philipps, P.; Fürstner, A. Total Synthesis of Putative Chagosensine. *Angew. Chem., Int. Ed.* **2018**, *57*, 13575–13581. (b) Han, B. Y.; Lam, N. Y. S.; MacGregor, C. I.; Goodman, J. M.; Paterson, I. A synthesis-enabled relative stereochemical assignment of the C1–C28 region of hemicalide. *Chem. Commun.* **2018**, *54*, 3247–3250. (c) Paul, D.; Kundu, A.; Saha, S.; Goswami, R. K. Total synthesis: the structural confirmation of natural products. *Chem. Commun.* **2021**, *57*, 3307–3322. (d) Fuwa, H. Structure determination, correction, and disproof of marine macrolide natural products by chemical synthesis. *Org. Chem. Front.* **2021**, *8*, 3990–4023.
- (7) (a) Challis, G. L. Genome Mining for Novel Natural Product Discovery. *J. Med. Chem.* **2008**, *51*, 2618–2628. (b) Rutledge, P. J.; Challis, G. L. Discovery of microbial natural products by activation of silent biosynthetic gene clusters. *Nat. Rev. Microbiol.* **2015**, *13*, 509–523.
- (8) Bachmann, B. O.; Van Lanen, S. G.; Baltz, R. H. Microbial genome mining for accelerated natural products discovery: is a renaissance in the making? *J. Ind. Microbiol. Biotechnol.* **2014**, *41*, 175–184.
- (9) Ziemert, N.; Alanjary, M.; Weber, T. The evolution of genome mining in microbes – a review. *Nat. Prod. Rep.* **2016**, *33*, 988–1005.
- (10) Boddy, C. N. Bioinformatics tools for genome mining of polyketide and non-ribosomal peptides. *J. Ind. Microbiol. Biotechnol.* **2014**, *41*, 443–450.
- (11) Helfrich, E. J. N.; Reiter, S.; Piel, J. Recent advances in genome-based polyketide discovery. *Curr. Opin. Biotechnol.* **2014**, *29*, 107–115.
- (12) Laureti, L.; Song, L.; Huang, S.; Corre, C.; Leblond, P.; Challis, G. L.; Aigle, B. Identification of a bioactive 51-membered macrolide complex by activation of a silent polyketide synthase in *Streptomyces ambofaciens*. *Proc. Natl. Acad. Sci. U. S. A.* **2011**, *108*, 6258–6263.
- (13) Aigle, B.; Challis, G.; Laureti, L.; Song, L.; Leblond, P. Stambomycin and Derivatives, Their Production and Their Use as Drugs. US20120142622A1, June 7, 2012.
- (14) (a) Keatinge-Clay, A. T. Stereocontrol within polyketide assembly lines. *Nat. Prod. Rep.* **2016**, *33*, 141–149. (b) Keatinge-Clay, A. T. A Tylosin Ketoreductase Reveals How Chirality Is Determined in Polyketides. *Chem. Biol.* **2007**, *14*, 898–908. (c) Kwan, D. H.; Sun, Y.; Schulz, F.; Hong, H.; Popovic, B.; Sim-Stark, J. C. C.; Haydock, S. F.; Leadlay, P. F. Prediction and Manipulation of the Stereochemistry of Enoylreduction in Modular Polyketide Synthases. *Chem. Biol.* **2008**, *15*, 1231–1240.
- (15) Song, L.; Laureti, L.; Corre, C.; Leblond, P.; Aigle, B.; Challis, G. L. Cytochrome P450-mediated hydroxylation is required for polyketide macrolactonization in stambomycin biosynthesis. *J. Antibiot.* **2014**, *67*, 71–76.
- (16) For selected recent examples where bioinformatics was used in the stereochemical determination of polyketides, see: (a) Hu, Y.; Wang, M.; Wu, C.; Tan, Y.; Li, J.; Hao, X.; Duan, Y.; Guan, Y.; Shang, X.; Wang, Y.; Xiao, C.; Gan, M. Identification and Proposed Relative and Absolute Configurations of Niphimycins C–E from the Marine-Derived *Streptomyces* sp. IMB7–145 by Genomic Analysis. *J. Nat. Prod.* **2018**, *81*, 178–187. (b) Kim, M. C.; Machado, H.; Jang, K. H.; Trzoss, L.; Jensen, P. R.; Fenical, W. Integration of Genomic Data with NMR Analysis Enables Assignment of the Full Stereostructure of Neaumycin B, a Potent Inhibitor of Glioblastoma from a Marine-Derived *Micromonospora*. *J. Am. Chem. Soc.* **2018**, *140*, 10775–10784. (c) Pérez-Victoria, I.; Oves-Costales, D.; Lacret, R.; Martín, J.; Sánchez-Hidalgo, M.; Díaz, C.; Cautain, B.; Vicente, F.; Genilloud, O.; Reyes, F. Structure elucidation and biosynthetic gene cluster analysis of caniferolides A–D, new bioactive 36-membered macrolides from the marine-derived *Streptomyces caniferus* CA-271066. *Org. Biomol. Chem.* **2019**, *17*, 2954–2971. (d) Kim, M. C.; Winter, J. M.; Cullum, R.; Li, Z.; Fenical, W. Complementary Genomic, Bioinformatics, and Chemical Approaches Facilitate the Absolute Structure Assignment of Ionostatin, a Linear Polyketide from a Rare Marine-Derived Actinomycete. *ACS Chem. Biol.* **2020**, *15*, 2507–2515. (e) Low, Z. J.; Xiong, J.; Xie, Y.; Ma, G.-L.; Saw, H.; Thi Tran, H.; Wong, S. L.; Pang, L. M.; Fong, J.; Lu, P.; Hu, J.-F.; Liang, Y.; Miao, Y.; Liang, Z.-X. Discovery, biosynthesis and antifungal mechanism of the polyene-polyol meijiemycin. *Chem. Commun.* **2020**, *56*, 822–825.
- (17) Jang, H.; Zhugralin, A. R.; Lee, Y.; Hoveyda, A. H. Highly Selective Methods for Synthesis of Internal (α -) Vinylboronates through Efficient NHC–Cu-Catalyzed Hydroboration of Terminal Alkynes. Utility in Chemical Synthesis and Mechanistic Basis for Selectivity. *J. Am. Chem. Soc.* **2011**, *133*, 7859–7871.

(18) Suen, L. M.; Steigerwald, M. L.; Leighton, J. L. A new and more powerfully activating diamine for practical and scalable enantioselective aldehyde crotylsilylation reactions. *Chem. Sci.* **2013**, *4*, 2413–2417.

(19) Evans, D. A.; Gauchet-Prunet, J. A. Diastereoselective Synthesis of Protected Syn 1,3-Diols by Base-Catalyzed Intramolecular Conjugate Addition of Hemiacetal-Derived Alkoxide Nucleophiles. *J. Org. Chem.* **1993**, *58*, 2446–2453.

(20) (a) Dale, J. A.; Mosher, H. S. Nuclear Magnetic Resonance Enantiomer Reagents. Configurational Correlations via Nuclear Magnetic Resonance Chemical Shifts of Diastereomeric Mandelate, O-Methylmandelate, and α -Methoxy- α -trifluoromethylphenylacetate (MTPA) Esters. *J. Am. Chem. Soc.* **1973**, *95*, 512–519. (b) Ohtani, I.; Kusumi, T.; Kashman, Y.; Kakisawa, H. High-Field FT NMR Application of Mosher's Method. The Absolute Configurations of Marine Terpenoids. *J. Am. Chem. Soc.* **1991**, *113*, 4092–4096.

(21) Boeckman, R. K., Jr.; Biegasiewicz, K. F.; Tusch, D. J.; Miller, J. R. Organocatalytic Enantioselective α -Hydroxymethylation of Aldehydes: Mechanistic Aspects and Optimization. *J. Org. Chem.* **2015**, *80*, 4030–4045.

(22) Chen, W.; Tay, J.-H.; Ying, J.; Yu, X.-Q.; Pu, L. Catalytic Asymmetric Enyne Addition to Aldehydes and Rh(I)-Catalyzed Stereoselective Domino Pauson–Khand/[4 + 2] Cycloaddition. *J. Org. Chem.* **2013**, *78*, 2256–2265.

(23) Park, J. K.; Ondrusek, B. A.; McQuade, D. T. Regioselective Catalytic Hydroboration of Propargylic Species Using Cu(I)-NHC Complexes. *Org. Lett.* **2012**, *14*, 4790–4793.

(24) Evans, D. A.; Chapman, K. T.; Carreira, E. M. Directed Reduction of β -Hydroxy Ketones Employing Tetramethylammonium Triacetoxyborohydride. *J. Am. Chem. Soc.* **1988**, *110*, 3560–1578.

(25) Rychnovsky, S. D.; Skalitzky, D. J. Stereochemistry of Alternating Polyol Chains: ^{13}C NMR Analysis of 1,3-Diol Acetonides. *Tetrahedron Lett.* **1990**, *31*, 945–948.

(26) Wang, Y. D.; Kimball, G.; Prashad, A. S.; Wang, Y. Zr-Mediated hydroboration: stereoselective synthesis of vinyl boronic esters. *Tetrahedron Lett.* **2005**, *46*, 8777–8780.

(27) Uenishi, J.; Beau, J.-M.; Armstrong, R. W.; Kishi, Y. Dramatic Rate Enhancement of Suzuki Diene Synthesis: Its Application to Palytoxin Synthesis. *J. Am. Chem. Soc.* **1987**, *109*, 4756–4758.