



This is a repository copy of *Performance of the Hull Salford Cambridge Decision Rule (HSC DR) for early discharge of patients with findings on CT scan of the brain : a CENTER-TBI validation study.*

White Rose Research Online URL for this paper:  
<https://eprints.whiterose.ac.uk/179340/>

Version: Accepted Version

---

### Article:

Marincowitz, C. [orcid.org/0000-0003-3043-7564](https://orcid.org/0000-0003-3043-7564), Gravesteijn, B., Sheldon, T. et al. (2 more authors) (2021) Performance of the Hull Salford Cambridge Decision Rule (HSC DR) for early discharge of patients with findings on CT scan of the brain : a CENTER-TBI validation study. *Emergency Medicine Journal*. ISSN 1472-0205

<https://doi.org/10.1136/emered-2020-210975>

---

© 2021 Author(s) (or their employer(s)). Reuse of this manuscript version (excluding any databases, tables, diagrams, photographs and other images or illustrative material included where a another copyright owner is identified) is permitted strictly pursuant to the terms of the Creative Commons Attribution-NonCommercial 4.0 International (CC-BY-NC 4.0) <https://creativecommons.org/licenses/by-nc/4.0/>.

### Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial (CC BY-NC) licence. This licence allows you to remix, tweak, and build upon this work non-commercially, and any new works must also acknowledge the authors and be non-commercial. You don't have to license any derivative works on the same terms. More information and the full terms of the licence here:  
<https://creativecommons.org/licenses/>

### Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# Emergency Medicine Journal

## Performance of the Hull Salford Cambridge Decision Rule (HSC DR) for Early Discharge of patients with findings on CT brain scan: a CENTER-TBI validation study.

Journal:	<i>Emergency Medicine Journal</i>
Manuscript ID	emermed-2020-210975.R2
Article Type:	Original research
Date Submitted by the Author:	n/a
Complete List of Authors:	Marincowitz, Carl; The University of Sheffield, School of Health and Related Research (SchARR) Gravesteijn, Benjamin; Erasmus Medical Center, Department of Public Health Sheldon, Trevor; Queen Mary University of London Barts and The London School of Medicine and Dentistry, Institute of Population Health Sciences Steyerberg, Ewout; Leiden University Medical Center Department of Biomedical Data Sciences Lecky, Fiona; University of Sheffield and Salford Royal Hospital, School of Health and Related Research. Emergency Department, Salford Royal Hospital
Keywords:	Trauma, head, Trauma, research

SCHOLARONE™  
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

1  
2  
3 **Performance of the Hull Salford Cambridge Decision Rule (HSC DR) for Early Discharge of**  
4  
5 **patients with findings on CT brain scan: a CENTER-TBI validation study.**  
6  
7

8 Carl Marincowitz<sup>1</sup> NIHR Clinical Lecturer Emergency Medicine, MB BChir, PhD, MSc, BA (Hons),  
9 MRCEM  
10

11 B.Y. Gravesteijn<sup>2</sup> MSc, PhD Candidate  
12

13 Trevor A. Sheldon<sup>3</sup> Professor, MSc, MSc, DSc, FMedSci  
14

15 Ewout W. Steyerberg<sup>4</sup> Professor, MSc, PhD  
16

17 Fiona E. Lecky<sup>1,5</sup> Professor, Honorary Emergency Medicine Consultant, MB ChB, FRCS, DA, MSc, PhD,  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

FRCEM

1. **Corresponding Author.** Centre for Urgent and Emergency Care Research (CURE), Health Services  
Research School of Health and Related Research, University of Sheffield, Regent Court, 30 Regent  
Street, Sheffield, S1 4DA, UK, Fax: +44 (0)114 222 0749 Tel: (+44) (0)114 222 4345,

Email: [C.Marincowitz@Sheffield.ac.uk](mailto:C.Marincowitz@Sheffield.ac.uk)

2. Department of Public Health, Erasmus Medical Centre, P.O. Box 2040, 3000 CA, Rotterdam, The  
Netherlands. Email: [b.gravesteijn@erasmusmc.nl](mailto:b.gravesteijn@erasmusmc.nl)

3. Institute of Population Health Sciences, Barts and The London School of Medicine and Dentistry,  
Queen Mary University London, Yvonne Carter Building, 58 Turner Street, London E1 2AB, Email:  
[t.sheldon@qmul.ac.uk](mailto:t.sheldon@qmul.ac.uk)

4. Department of Biomedical Data Sciences, Leiden University Medical Center, Albinusdreef 2,  
2333 ZA Leiden, Tel: +31 71 526 9700, Email: [e.w.steyerberg@lumc.nl](mailto:e.w.steyerberg@lumc.nl)

5. Emergency Department, Salford Royal Hospital, Salford, UK. Email: [F.E.Lecky@Sheffield.ac.uk](mailto:F.E.Lecky@Sheffield.ac.uk)

## Abstract

### Background

There is international variation in hospital admission practices for patients with mild traumatic brain injury (TBI) and injuries on CT scan. Only a small proportion of patients require neurosurgical intervention, while many guidelines recommend routine admission of all patients. We aim to validate the Hull Salford Cambridge Decision Rule (HSC DR) and the Brain Injury Guideline (BIG) criteria to select low risk patients for discharge from the Emergency Department.

### Method

A cohort from 18 countries of GCS 13-15 patients with injuries on CT imaging was identified from the multi-centre CENTER-TBI study (conducted 2014 - 2017) for secondary analysis. A composite outcome measure encompassing need for ongoing hospital admission was used, including seizure activity, death, intubation, neurosurgical intervention, and neurological deterioration. We assessed the performance of our previously derived prognostic model, the HSC DR and the BIG criteria at predicting deterioration in this validation cohort.

### Results

Among 1047 patients meeting the inclusion criteria, 267 (26%) deteriorated. Our prognostic model achieved a C-statistic of 0.81 (95% CI, 0.78 to 0.84). The HSC DR achieved a sensitivity of 100% (95% CI: 97% to 100%) and specificity of only 4.7% (95% CI: 3.3% to 6.5%) for deterioration. Using the BIG criteria for discharge from the ED achieved a higher specificity (13.3%, 95% CI: 10.9% to 16.1%) and lower sensitivity (94.6%, 95% CI: 90.5 % to 97%), with

1  
2  
3 12/105 patients recommended for discharge subsequently deteriorating, compared to 0/34  
4  
5 with the HSC DR.  
6  
7

## 8 9 Conclusion

10  
11  
12 Our decision rule would have allowed 3.5% of patients to be discharged, none of whom  
13  
14 would have deteriorated. Use of the BIG criteria may result in too high a risk of  
15  
16 deterioration in a discharged patient to be used clinically. Further validation and  
17  
18 implementation studies are required to support use in clinical practice.  
19  
20  
21  
22

## 23 What is already known on this subject

24  
25  
26 NICE head injury guidelines state that following head injury, patients with “new, clinically  
27  
28 significant abnormalities on imaging” should be admitted for observation without defining  
29  
30 which injuries are clinically significant. We have previously empirically derived the first  
31  
32 prognostic model and decision rule (HSC-DR) to identify low risk patients with injuries on CT  
33  
34 who could be safely discharged from the ED.  
35  
36  
37  
38  
39

## 40 What this Study adds

41  
42  
43 We present the first validation study of our prognostic model and the HSC-DR. It shows that  
44  
45 application of the HSC-DR may allow a modest but safe reduction in inpatient admissions of  
46  
47 selected low risk patients with traumatic brain injuries identified by CT imaging.  
48  
49  
50

51 **Keywords:** Mild Traumatic Brain Injury; Prognostic Model; Clinical Decision Rule; Emergency  
52  
53 Department; Head Injury  
54  
55  
56  
57  
58  
59  
60

## Background

Over 2 million patients are admitted to hospital each year across Europe for traumatic brain injury (TBI; injury to the brain or alteration of brain function due to external force).<sup>1</sup> 95% of patients admitted to hospital and 36% of patients admitted to intensive care units with TBI have an initial Glasgow Coma Scale (GCS) of 13-15 and are defined as having mild injuries.<sup>2</sup> The management of mild TBI patients with injuries identified by CT imaging is controversial. Around 7% of initial GCS13-15 patients who present with head trauma have intra-cranial injuries or skull fractures identified on CT imaging but only around 1% of patients die or require neurosurgery.<sup>3</sup> Some studies advocate routine admission under specialist neurosurgical care and repeat CT imaging of all mild TBI patients with injuries identified on CT.<sup>4,5</sup> Some North American centres have adopted the consensus derived Brain Injury Guideline (BIG) criteria which advocates the discharge of selected patients from the ED (Supplementary Material 1).<sup>6</sup> In Europe there is variation in clinical practice with patients admitted under a range of specialties and with varying levels of intensity of inpatient care.<sup>2</sup> We recently developed the first empirically derived prognostic model and decision rule (the Hull Salford Cambridge Decision Rule (HSC DR)) predicting need for hospital admission in this population.<sup>7</sup> We compared the performance of the HSC DR and BIG criteria and found both had high sensitivity to clinical deterioration. The HSC DR maximised sensitivity at a cost of a specificity of 7% at the discharge threshold to ensure clinical safety, but implementation would have recommended fewer than one in ten TBI patients be discharged.<sup>7</sup> However, in the "COVID 19" era - where reducing hospital acquired infections is paramount, and in other resource constrained contexts, even small reductions in unnecessary hospital admissions are valuable. Application of this decision rule could – if externally validated – achieve this.<sup>7</sup>

1  
2  
3 The aims of this study were:  
4  
5

- 6 1. Externally validate and compare the performance of the HCS and BIG criteria  
7 decision rules, using an international dataset of patients attending Emergency  
8 Departments following traumatic brain injury.  
9
- 10 2. Evaluate the performance of the HCS and BIG criteria decision rules for mildly injured  
11 patients with TBI.  
12
- 13 3. Externally validate the empirically derived prediction model underpinning HSC-DR  
14 (recalibrating where required) using the CENTER TBI cohort.  
15  
16  
17  
18  
19  
20  
21  
22

## 23 **Methods**

### 24 *Study design*

25  
26  
27 An international dataset of patients with CT diagnosed TBI, was used to externally validate  
28 the two decision rules (BIG and HSC-DR) by comparing their sensitivity and specificity for  
29 predicting which patients required hospital admission for specific treatments.<sup>2,8</sup> The  
30 CENTER-TBI dataset was then used to recalibrate the HSC prediction model (which then  
31 feeds into the decision rule). The aim of the recalibration was to determine if the HSC  
32 decision rule performance could be improved using data from a more diverse population  
33 compared to the initial derivation dataset. We followed international guidelines (TRIPOD)  
34 for reporting of prognostic model validation.<sup>8</sup> The methods used to derive our prognostic  
35 model and the HSC-DR are available in the previously published protocol and derivation  
36 studies.<sup>7,9</sup>  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



### *Source of data*

Data for the core CENTER-TBI study were collected between December 2014 and 2017 at 63 centres across Europe and Israel and 4509 patients of all TBI severity were recruited, stratified by three strata of planned clinical management: ED only, admitted initially as a ward inpatient and admitted initially to intensive care. All patients were initially managed in the Emergency Department. Data were prospectively collected by trained research staff as detailed in the study protocol.<sup>10</sup> Follow up data were collected at 2-3 weeks, 3 months and 6 months with data collected on 83.4% of patients at 6-months.

### *Inclusion and exclusion criteria*

Patients aged 16 and over with an initial GCS 13-15 recorded in the ED and with either a skull fracture, intra-cranial haemorrhage or cerebral contusion identified on first CT scan - regardless of care pathway stratum were included, reflecting the population used in our derivation study.<sup>7</sup> Patients where initial GCS in the ED was unknown and patients where diffuse axonal injury was the sole injury identified on initial CT scan were excluded.

### *Outcome*

A composite outcome encompassing need for hospital admission was defined, matching the outcome in the model derivation study. This included: seizure as inpatient or at 2 week follow-up, death attributed to TBI within 30 days of first attendance, intubation recorded within 30 days of presentation, admission to ICU for any reason apart from close monitoring, neurosurgical intervention and recorded neurological deterioration (new deficit or drop in GCS of more than 1 point).

### Predictors

The original extended prediction model includes seven predictor variables for a composite outcome of deterioration encompassing need for hospital admission in this TBI population (Table 1).<sup>7</sup> The full prediction model is available in Supplementary Material 2. Six of these variables were used in our derivation study to form the simplified HSC DR which could be applied clinically to identify patients who could be safely discharged from the ED (Table 1 and Supplementary Material 2). The BIG criteria use 6 factors to risk stratify patient management (Supplementary Material 1). All factors in the prediction model and BIG criteria were available from data collected in CENTER-TBI.

**Table 1: Factors in extended prognostic model and HSC DR**

Factors in Extended model	HSC DR Discharge if	BIG Criteria Discharge after 6 hours if
Preinjury Anti-coagulation or anti-platelets	No	No
Initial GCS 13-15	GCS 15	13-15
First Neurological Examination	Normal	Normal
Number of Injuries on CT: 1-5 or Diffuse	1	
Injury severity on CT: Simple skull fracture Complex Skull Fracture Marshall IIa 1-2 bleeds < 5mm (total) Marshall IIb bleeds ≥ 5mm Marshall III/IV Marshall VI Brain stem/Cerebellar	Simple Skull fracture or 1-2 bleeds < 5mm total	Subdural ≤ 4mm Extradural ≤ 4mm 1 Intra-cerebral haemorrhage ≤ 4mm Trace Subarachnoid haemorrhage No skull fractures No Intra-ventricular haemorrhage
Injury Severity Score (body regions excluding head)	Up to 2 non-significant extra-cranial injuries(not requiring inpatient care, e.g closed fracture humerus)	
Intoxication		Not intoxicated
Hb	Not included in risk score	

### *Sample Size*

A minimum of between 100-200 events and 100-200 non-events per study sample has been recommended for validation studies of logistic regression models.<sup>11 12</sup> The validation cohort contained over 200 events and non-events.

### *Missing data*

To evaluate model performance, missing data were multiply imputed using the ICE STATA package on the assumption they were missing at random (fully described Supplementary Material 3).<sup>13</sup> Performance was averaged across imputed data sets.<sup>14 15</sup>

### *Decision Rule Performance*

All analysis was completed using STATA 16 (StataCorp. 2019. *Stata Statistical Software: Release 16*. College Station, TX: StataCorp LLC). Sensitivity, specificity of the HSC DR and of the BIG criteria to the composite outcome of deterioration were calculated in patients with complete data for either criteria. To be recommended for discharge all components of HSC DR or BIG criteria (Table 1) must be fulfilled. The proportion of patients recommended for discharge and accompanying risk of deterioration in a discharged patient (negative predictive value) were compared. In pre-specified exploratory subgroup analysis this was repeated in patients with less severe injuries as indicated by having a brain abbreviated injury score (AIS) or Marshall classification <3.<sup>16</sup> This represents patients without obvious midline shift or severe injuries on CT imaging and the population admitted for observation under ED care in the UK.

### *Model performance and recalibration*

1  
2  
3 Performance of the prediction model was assessed in the CENTER-TBI cohort using  
4  
5 measures of discrimination and calibration. Discrimination indicates how well the model  
6  
7 differentiates between patients who deteriorated and those who do not deteriorate and  
8  
9 was measured using the C-statistic (equivalent to the area under ROC curve).<sup>17</sup>  
10  
11

12  
13 Calibration measures how closely predictions made by the model match observed outcomes  
14  
15 (i.e. do predicted mean outcomes match observed mean outcomes).<sup>17</sup> Calibration was  
16  
17 assessed visually using a calibration plot and with estimates of the “calibration in the large”  
18  
19 (the ratio of expected versus observed numbers of events) and slope of the calibration plot  
20  
21 (the overall prognostic effects of predictors in the model). To account for differences  
22  
23 between the derivation and validation cohort and potential model over-fitting during  
24  
25 derivation, the intercept and coefficients of the prediction model were also re-estimated to  
26  
27 provide a re-calibrated model.  
28  
29  
30  
31  
32

### 33 *Clinical usefulness*

34  
35 Decision curve analysis was used to estimate the net benefit of using the prognostic model  
36  
37 to select patients for discharge from the ED.<sup>18 19</sup> Net benefit is estimated by the number of  
38  
39 true positives minus false positives multiplied by the clinical weight given to correct  
40  
41 classification across a range of probabilities of deterioration where discharge could be  
42  
43 considered.<sup>19</sup> The net benefit of using the prognostic model was compared visually in curves  
44  
45 using the BIG criteria’s single decision threshold and reference strategies of discharging no  
46  
47 or all patients.<sup>20</sup>  
48  
49  
50  
51  
52

### 53 **Ethics**

54  
55  
56  
57  
58  
59  
60

Ethics approval was obtained for each recruiting site, full details are available here

<https://www.center-tbi.eu/project/ethical-approval>. .

## **Patient and Public Involvement**

The Hull and East Yorkshire NHS Trust Trans-Humber Consumer Research Panel and Hull branch of the Headway charity helped inform developing the overall research aim of developing a predictive model to identify low risk patients with injuries on CT imaging who could be safely discharged from the ED.

## **Results**

### *Study population*

The cohort (n=1047) was mostly male, with over a third of patients aged over 65 and over 20% with either pre-injury anti-coagulant or anti-platelet use (Figure 1, Table 2). A total of 379 (36%) patients had data missing from at least one predictor variable value (mostly initial haemoglobin) used in the full prognostic model (Table 2). 12.1% patients had data missing in one or more predictor variable used in the HSC DR. Any clinical deterioration was noted among 267 patients (26%; 95% CI: 23% to 28%), including 212 patients (20%; 95% CI: 17% to 23%) who underwent neurosurgery, died, or were intubated and 25 patients had deaths attributable to TBI.

**Table 2: Characteristics of the study population (N=1047)**

Population Characteristic	Category	Mean (SD), min-max or N (%)	Missing data
Age	Years	54.8 (SD=19.7) 16-96	None
Age	≥65	384 (36.7%)	None
Sex	Male Female	688 (66%) 359 (34%)	None
GCS	15 14 13	677 (64.7%) 359 (24.7%) 111 (10.6%)	None
Stratum	ER Admission ICU	87 (8.3%) 587 (56%) 373 (35.6%)	None
Mechanism of Injury	High Velocity Trauma Blow to head/struck by object Ground level fall Fall from >1m or 5 stairs other	210 (20.1%) 183 (17.5%) 384 (36.7%) 218 (20.8%) 19 (1.8%)	33 (3.2%)
Intoxicated	Yes	242 (23.1%)	58 (5.5%)
Preinjury Anti-coagulation or anti-platelets	Anticoagulation use Antiplatelet use Both	72 (6.9%) 134(12.8%) 7 (0.7%)	12 (1.1%)
Abnormal First Neurological Examination	Yes	152 (14.5%)	71 (6.8%)
Haemoglobin	Grams/litre	135 (SD 19.9) 47-23.4	325 (31%)
Number of Injuries on CT	1 2 3 4 5 Multiple diffuse injury/>5	468 (44.7%) 243 (23.2%) 135 (12.9%) 81 (7.7%) 56 (5.4%) 64 (6.1%)	None
Injury severity on CT (Modified Marshall Classification described in detail Supplementary Material 2)	<b>1)</b> Simple Skull Fractures <b>2)</b> Complex Skull fractures <b>3)</b> 1-2 bleeds < 5mm (total) <b>4)</b> No or minimal mass effect <b>5)</b> Significant midline shift <b>6)</b> High/mixed-density lesion <b>7)</b> Cerebellar/Brain stem injury	19 (1.8%) 67 (6.4%) 426 (40.7%) 324 (31%) 29 (2.8%) 114 (10.9%) 68 (6.5%)	None
ISS	Body regions excluding head	17.3 (SD 20.6) 1-75 (range)	9 (0.9%)

### Decision Rule performance

The HCS DR achieved a sensitivity of 100% (95% CI: 98.8% to 100%), but very low specificity of 4.7% (95% CI: 3.3% to 6.5%) for the composite outcome of deterioration (Table 3). BIG 1 classification missed some events (sensitivity 94.6%, 95% CI: 90.5% to 97%), but had higher specificity (13.3%, 95% CI: 10.9% to 16.1%). Application of the HSC DR would have recommended discharge of only 3.5% of patients, compared to 11.4% patients recommended by the BIG criteria. However, patients recommended for discharge by the BIG criteria had a 11.4% (95% CI: 6.7% to 18.9%), risk of subsequent deterioration, compared to 0% (95% CI: 0% to 10.2%) with the HSC DR.

**Table 3: Performance of BIG and HSC Decision Rules \***

BIG Criteria Performance			
N=921	Deteriorated	Didn't deteriorate	
BIG1 (discharge from ED after 6 hours)	12	93	Sensitivity 94.6% (90.5-97%) Negative Predictive Value 88.6% (80.5 - 93.7%)
BIG 2/3 (admit)	210	606	Specificity 13.3% (10.9% - 16.1%) Positive Predictive Value 25.7% (22.8 - 28.9%)

HSC DR			
N=961	Deteriorated	Didn't Deteriorate	
Risk=0 (discharge)	0	34	Sensitivity 100% (98.8-100%) Negative Predictive Value 100% (87.4 - 100%)
Risk>0 (admit)	234	693	Specificity 4.7% (3.3-6.5%) Positive Predictive Value 25.2% (22.5 - 28.2%)

\*Full performance of the BIG are presented in Supplementary Material 4 and characteristics of patients recommended for discharge in Supplementary Material 5

### Sub-group analysis of less severely injured patients

One hundred and forty-six patients had AIS<3 and 800 patients had Marshall Classification <3 injuries. Use of the HSC DR would have facilitated discharge of 23% (34/146) of patients with brain AIS < 3, and 4.25% (34/800) of patients with Marshall Classification <3 injuries.

No patients selected for discharge by the HSC DR deteriorated (risk of deterioration 0%, 95% CI: 0% to 10.2%). Use of BIG criteria would have selected 26% (37/142) of patients with brain AIS < 3 injuries for discharge but with an 8.1% (95% CI: 2.8 % to 21.3%) risk of deterioration and 13.6% (105/770) of patients with Marshall classification < 3 injuries but with an 11.4% (95% CI: 6.7% to 18.9%) risk of deterioration (Table 4 and Supplementary Material 6).

**Table 4: Subgroup analysis AIS<3**

BIG 1			
N=142	Deteriorated	Didn't deteriorate	
BIG1 (discharge from ED after 6 hours)	3	34	Sensitivity 75% (42.8-93.3%) Negative Predictive Value 91.9% (77 – 97.9%)
BIG 2/3 (admit)	9	96	Specificity 26.2 (19-34.7%) Positive Predictive Value 8.6% (4.2 – 16.1%)

HSC DR			
N=146	Deteriorated	Didn't deteriorate	
Risk=0 (discharge)	0	34	Sensitivity 100% (69.99-100%) Negative Predictive Value 100% (87.4 - 100%)
Risk>0 (admit)	12	100	Specificity 25.4% (18.4-33.8%) Positive Predictive Value 10.7% (1075.9 - 18.313%)



1  
2  
3  
4  
5  
6 Twenty-seven patients were excluded from the cohort as the only injury identified on initial  
7  
8 CT imaging were diffuse axonal injury and therefore, they could not be assigned to a BIG  
9  
10 criterion. These injuries are equivalent to a Marshall score 4 severity and would be  
11  
12 recommend for admission by the HSC DR. Sensitivity analysis including these patients found  
13  
14 the HSC DR achieved a sensitivity (100% 95% CI: 98% to 100%) and specificity (4.5% 95% CI:  
15  
16 3.2% to 6.3%) to the composite outcome of deterioration.  
17  
18  
19

### 20 21 *Model Performance*

22  
23  
24 The original prognostic model achieved a C-statistic of 0.81 (95% CI, 0.78 to 0.84) in the  
25  
26 CENTER-TBI cohort (0.75 in the development cohort) and an estimated slope of the  
27  
28 calibration plot of 0.51 in the CENTER-TBI cohort (0.86 in the development cohort) (Figure  
29  
30 2i). The effect of re-calibration of both the intercept and coefficients is presented in Figure  
31  
32 2ii and the recalibrated model is presented in Supplementary Material 7. Measures of  
33  
34 calibration improved but the estimated C-statistic of the recalibrated model remained 0.81.  
35  
36  
37  
38  
39

### 40 *Clinical usefulness, analysis according to clinical tolerance for adverse outcomes*

41  
42  
43 Clinical usefulness depends on tolerance of risk of deterioration in those discharged without  
44  
45 observation. Figure 3 presents the decision curves and net benefit analysis for the selection  
46  
47 of patients either for a period of inpatient hospital observation or discharge directly from  
48  
49 the ED using the recalibrated prognostic model or BIG criteria in the CENTER-TBI cohort. Due  
50  
51 to the high risk of harm associated with discharging a patient who subsequently  
52  
53 deteriorates, the analysis was limited to those with a low predicted probability of  
54  
55 deterioration. Use of our recalibrated model showed potential benefit over an 'admit all'  
56  
57  
58  
59  
60

1  
2  
3 strategy if the threshold for the predicted probability of deterioration was over 2% (Figure  
4  
5  
6 3), which is potentially an acceptable clinical risk of deterioration in a discharged patient. If  
7  
8 2% is considered too high a risk to discharge a patient, given the harm associated with  
9  
10 deterioration in the community, then no net benefit over an “admit all” strategy was  
11  
12 demonstrated. The BIG criteria showed benefit over an ‘admit all’ strategy up to a threshold  
13  
14 for predicted probability of deterioration of around 12%.  
15  
16  
17

## 18 **Discussion**

### 21 *Summary*

22  
23  
24  
25 This study validated the performance of the BIG and HSC decision rules in a large  
26  
27 international dataset of patients with TBI, who had an overall deterioration prevalence of  
28  
29 26% (95%CI 23%, 28%). The BIG criteria achieved a sensitivity of 94.6% (95% CI: 90.5 % to  
30  
31 97%) and specificity of 13.3% (95% CI: 10.9% to 16.1%) and would have recommended  
32  
33 discharge of 11% of patients with an accompanying risk of subsequent deterioration of  
34  
35 11.4% (95% CI: 6.7 % to 18.9%). The HSC DR achieved a sensitivity of 100% (95% CI: 98% to  
36  
37 100%) and specificity of 4.7% (95% CI: 3.3% to 6.5%), comparable to that reported in the  
38  
39 development cohort (99.5% and 4.8% respectively). The HSC DR would have recommended  
40  
41 discharge of 3.5% of patients but with a subsequent risk of deterioration of 0% (95% CI: 0 %  
42  
43 to 10.2%). The prognostic model that underpins the HSC DR achieved a C-statistic of 0.81  
44  
45 and re-calibration improved accuracy of individual predicted risk of deterioration  
46  
47 (calibration).  
48  
49  
50  
51  
52  
53

54  
55 In the subgroup of patients with less severe injuries who are more likely to admitted under  
56  
57 non-specialist teams the BIG criteria recommended discharge of 26% of patients with brain  
58  
59  
60

1  
2  
3 AIS < 3 injuries for discharge but with an 8.1% (95% CI: 2.8 % to 21.3%) risk of deterioration.

4  
5 The HSC DR recommended discharge of 23% of patients of patient in this group with a risk  
6  
7  
8 of subsequent deterioration of 0% (95% CI: 0% to 10.2%).  
9

### 10 11 *Strengths*

12  
13  
14  
15 This study is the first external validation of the HSC-DR and, alongside our previous  
16  
17 development study, is the largest study to externally validate the BIG criteria and only study  
18  
19 to do so in a multi-centre European cohort of patients.<sup>4 21-23</sup> The CENTER-TBI study has good  
20  
21 prospective patient follow-up and so significant adverse outcomes in the community were  
22  
23 unlikely to have been missed. We have adhered to international guidelines for model  
24  
25 validation.<sup>8</sup> We explicitly addressed the potential clinical usefulness of the decision rule and  
26  
27 prognostic model according to a range of potential thresholds. This decision curve analysis  
28  
29 clarified that if quite low risks were already considered too high, e.g. corresponding to a  
30  
31 threshold of 1%, a treat all strategy would dominate. On the other hand, a less risk averse  
32  
33 clinical policy, such as accepting risks up to 10% as acceptable, would lead to greater value  
34  
35 of our rule or model (Fig 3).  
36  
37  
38  
39  
40  
41  
42

### 43 *Limitations*

44  
45  
46 Previous studies estimated that around 10% of initial GCS13-15 patients have skull fractures  
47  
48 or intra-cranial injuries identified on CT imaging, whilst in the CENTER-TBI study around 50%  
49  
50 of patients have injuries identified on imaging.<sup>3 24 25</sup> The CENTER-TBI population may be a  
51  
52 higher risk group than the clinical population assessed in the ED. There was a relatively high  
53  
54 proportion of missing data, especially for haemoglobin values. However, it is likely these  
55  
56 data were missing at random, i.e. only related to observed variables, and that imputation  
57  
58  
59  
60

1  
2  
3 methods we used are valid. Study recruitment for CENTER-TBI occurred at 2 sites  
4  
5  
6 (Cambridge and Salford) at which the case note review for derivation of our prognostic  
7  
8 model was conducted. These sites only contributed 6.9% of patients to the CENTER-TBI  
9  
10 validation cohort and exclusion of these patients did not materially affect our results  
11  
12 (Supplementary Material 8). Determining the significance of extra-cranial injuries in the  
13  
14 HSC-DR as derived from extra-cranial ISS score (including facial injuries) requires some  
15  
16 subjective clinical judgement.  
17  
18  
19  
20  
21  
22  
23  
24

#### 25 *Comparison to previous literature*

26  
27  
28 In the CENTER TBI cohort, 20% of patients underwent neurosurgery, died, or were intubated  
29  
30 compared to 13.1% in our development cohort and had a higher prevalence of deterioration  
31  
32 than reported in a previous systematic review.<sup>4</sup> This may reflect recruitment of more  
33  
34 severely injured patients to the CENTER-TBI study.  
35  
36  
37  
38

39 The BIG criteria for discharging patients from the ED achieved a lower sensitivity (94.6%)  
40  
41 and higher specificity (13.3%) than when applied to our development cohort (sensitivity  
42  
43 99.5% and specificity 4.8%). Application of the BIG criteria would have allowed 11.4% of  
44  
45 patients to be discharged from the ED which is similar to the 10% of patients estimated in  
46  
47 studies conducted where the BIG criteria was developed in the USA and 15% reported in an  
48  
49 external validation study.<sup>6 21 23</sup> The derivation and validation studies reported by the team  
50  
51 that developed the BIG criteria and available external validation studies report no adverse  
52  
53 outcomes in patients recommended for discharge by the BIG criteria.<sup>6 21-23 26</sup> In the CENTER-  
54  
55 TBI cohort, patients recommended for discharge had a 11.4% (95% CI: 6.7 % to 18.9%), risk  
56  
57  
58  
59  
60

1  
2  
3 of subsequently deteriorating. This may reflect the broader composite outcome measure  
4 used in our study and more comprehensive prospective follow-up of patients for  
5 deterioration. Some validation studies also modified the BIG criteria so that any patient with  
6 an initial GCS <15 was admitted to hospital.<sup>22</sup> The USA TBI population used for these studies  
7 also appears to be lower risk with a lower reported average age, anti-coagulant use and  
8 neurosurgical intervention rate.<sup>4,23</sup> The risk of deterioration when discharging a patient from  
9 the ED that is acceptable to patients and clinicians is subjective. When deriving the HSC-DR<sup>7</sup>  
10 we aimed to maximise sensitivity and aimed for a risk of a discharged patient deteriorating  
11 of around 1%, as this corresponds to other decision rules for discharging patients from the  
12 ED,<sup>25,27</sup> and may be a sufficiently low risk to consider routine discharge. However, significant  
13 variation in risk tolerance in clinicians and public representatives has been demonstrated,  
14 with some indicating that even a 1% risk of deterioration may be too high.<sup>28,29</sup> *Implications*  
15  
16 There is variation internationally in management and admission practices in this TBI  
17 population.<sup>4</sup> In the UK and other European countries guidelines recommend admission of all  
18 patients with TBI identified on CT imaging. This validation study shows a recalibrated version  
19 of our prognostic model could allow accurate prediction of risk of deterioration, and  
20 application of the HSC DR would have allowed a modest but safe reduction in hospital  
21 admissions for this group. The application of the BIG criteria would have discharged more  
22 patients but with a higher risk of subsequent deterioration in this European population,  
23 which may not be clinically acceptable. As indicated by our exploratory sub-group analysis,  
24 application of the HSC DR may be more beneficial when applied to lower risk populations  
25 more reflective of patients who attend the ED and are admitted for observation under  
26 Emergency Medicine or other non-neurosurgical specialities in the UK.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Our net benefit analysis using decision curves (Figure 3) showed use of our prognostic  
4 model may show benefit over an 'admit all' strategy if the threshold for the predicted  
5 probability of deterioration was over 2% and patients selected for discharge by the HSC DR  
6 had a 0% (95% CI: 0 % to 10.2%) risk of deterioration. This may be sufficiently low risk to use  
7 routinely. Research is needed to assess clinician and patient risk appetite in this population  
8 and assess the clinical impact of implementing the HSC DR where patient circumstances like  
9 intoxication or social circumstances may further affect whether a patient can be discharged.  
10  
11 Research to improve the accuracy of the prognostic model (e.g. through including  
12 biomarkers, other novel prognostic factors, or better classification of injury severity on CT  
13 imaging) is also needed. .

### 28 *Conclusion*

31 Use of the HSC DR would allow a modest but safe reduction in hospital admissions for mild  
32 TBI patients with injuries identified on CT. The BIG criteria appear to result in an  
33 unacceptably high risk of subsequent deterioration (one in ten) among discharged patients.  
34  
35 Future research should further validate our prognostic model and the HSC DR, consider safe  
36 implementation into clinical practice and assess whether inclusion of novel prognostic  
37 factors could improve the specificity of the model allowing more patients to be safely  
38 discharged.

### 50 **Author Disclosure Statement:**

51 No competing financial interests exist.

52  
53 Data used in preparation of this manuscript were obtained in the context of CENTER-TBI, a  
54 large collaborative project with the support of the European Union 7<sup>th</sup> Framework program  
55 (EC grant602150). Full list of participants and investigators are provided in Supplementary  
56  
57  
58  
59  
60

1  
2  
3 Material 9. Additional funding was obtained from the Hannelore Kohl Stiftung (Germany),  
4 from OneMind (USA) and from Integra LifeSciences Corporation (USA) (Grant Number Not  
5 Applicable/NA).  
6  
7

8  
9 FEL and EWS are supported by the European Union Framework 7 Collaborative European  
10 Neurotrauma Effectiveness Research in Traumatic Brain Injury ((EC grant 602150)). FEL is  
11 also supported by the NHS Trusts "Trauma Audit and Research Network -  
12 [www.tarn.ac.uk](http://www.tarn.ac.uk)"(Grant Number Not Applicable/NA).  
13  
14  
15

16  
17 CM is a National Institute for Health Research (NIHR) Clinical Lecturer in Emergency  
18 Medicine (Grant Number Not Applicable/NA). This publication presents independent  
19 research funded by the National Institute for Health Research and University of Sheffield.  
20  
21 The views expressed are those of the author(s) and not necessarily those of the University of  
22 Sheffield, the NHS, the NIHR or the Department of Health and Social Care  
23  
24  
25

#### 26 27 **Authors' contributions:**

28  
29 The idea for the study was conceived by CM, TAS and FEL. The analysis was completed by  
30 CM with specialist statistical advice from BYG and EWS and specialist clinical advice from  
31 FEL. All authors contributed to interpretation of results, read and approved the final  
32 manuscript.  
33  
34  
35  
36

#### 37 38 **Figures:**

39  
40  
41 Figure 1: STROBE flow diagram of selection of study population

42  
43 Figure 2: Slope of the calibration plot of original and re-calibrated prognostic model

44  
45 Figure 3: Decision Curve analysis  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## References

1. Majdan M, Plancikova D, Brazinova A, et al. Epidemiology of traumatic brain injuries in Europe: a cross-sectional analysis. *Lancet Public Health* 2016;1(2):e76-e83. doi: 10.1016/S2468-2667(16)30017-2 [published Online First: 2017/12/19]
2. Steyerberg EW, Wiegers E, Sewalt C, et al. Case-mix, care pathways, and outcomes in patients with traumatic brain injury in CENTER-TBI: a European prospective, multicentre, longitudinal, cohort study. *Lancet Neurol* 2019;18(10):923-34. doi: 10.1016/S1474-4422(19)30232-7 [published Online First: 2019/09/19]
3. Haydel MJ, Preston CA, Mills TJ, et al. Indications for computed tomography in patients with minor head injury. *N Engl J Med* 2000;343(2):100-5. doi: 10.1056/NEJM200007133430204
4. Marincowitz C, Lecky FE, Townend W, et al. The Risk of Deterioration in GCS13-15 Patients with Traumatic Brain Injury Identified by Computed Tomography Imaging: A Systematic Review and Meta-Analysis. *J Neurotrauma* 2018;35(5):703-18. doi: 10.1089/neu.2017.5259 [published Online First: 2018/01/13]
5. Thomas BW, Mejia VA, Maxwell RA, et al. Scheduled repeat CT scanning for traumatic brain injury remains important in assessing head injury progression. *J Am Coll Surg* 2010;210(5):824-30, 31-2. doi: 10.1016/j.jamcollsurg.2009.12.039
6. Joseph B, Friese RS, Sadoun M, et al. The BIG (brain injury guidelines) project: defining the management of traumatic brain injury by acute care surgeons. *J Trauma Acute Care Surg* 2014;76(4):965-9. doi: 10.1097/TA.0000000000000161
7. Marincowitz C, Lecky FE, Allgar V, et al. Development of a Clinical Decision Rule for the Early Safe Discharge of Patients with Mild Traumatic Brain Injury and Findings on Computed Tomography Brain Scan: A Retrospective Cohort Study. *J Neurotrauma* 2019 doi: 10.1089/neu.2019.6652 [published Online First: 2019/10/08]
8. Collins GS, Reitsma JB, Altman DG, et al. Transparent Reporting of a multivariable prediction model for Individual Prognosis Or Diagnosis (TRIPOD). *Ann Intern Med* 2015;162(10):735-6. doi: 10.7326/L15-5093-2 [published Online First: 2015/05/20]
9. Marincowitz C, Lecky FE, Townend W, et al. A protocol for the development of a prediction model in mild traumatic brain injury with CT scan abnormality: which patients are safe for discharge? *Diagnostic and Prognostic Research* 2018;2(1):6. doi: 10.1186/s41512-018-0027-4
10. Maas AI, Menon DK, Steyerberg EW, et al. Collaborative European NeuroTrauma Effectiveness Research in Traumatic Brain Injury (CENTER-TBI) A Prospective Longitudinal Observational Study. *Neurosurgery* 2014;76(1):67-80.
11. Vergouwe Y, Steyerberg EW, Eijkemans MJ, et al. Substantial effective sample sizes were required for external validation studies of predictive logistic regression models. *J Clin Epidemiol* 2005;58(5):475-83. doi: 10.1016/j.jclinepi.2004.06.017 [published Online First: 2005/04/23]
12. Collins GS, Ogundimu EO, Altman DG. Sample size considerations for the external validation of a multivariable prognostic model: a resampling study. *Stat Med* 2016;35(2):214-26. doi: 10.1002/sim.6787 [published Online First: 2015/11/11]
13. Royston P, White IR. Multiple imputation by chained equations (MICE): implementation in Stata. *J Stat Softw* 2011;45(4):1-20.
14. Sterne JA, White IR, Carlin JB, et al. Multiple imputation for missing data in epidemiological and clinical research: potential and pitfalls. *BMJ* 2009;338:b2393. doi: 10.1136/bmj.b2393 [published Online First: 2009/07/01]
15. Nguyen CD, Carlin JB, Lee KJ. Model checking in multiple imputation: an overview and case study. *Emerg Themes Epidemiol* 2017;14:8. doi: 10.1186/s12982-017-0062-6 [published Online First: 2017/08/31]



16. Association for the Advancement of Automotive Medicine. (2018). Abbreviated Injury Scale: 2015 Revision (6 ed.). Chicago, IL.
17. Steyerberg EW, Vickers AJ, Cook NR, et al. Assessing the performance of prediction models: a framework for traditional and novel measures. *Epidemiology* 2010;21(1):128-38. doi: 10.1097/EDE.0b013e3181c30fb2 [published Online First: 2009/12/17]
18. Vickers AJ, Elkin EB. Decision curve analysis: a novel method for evaluating prediction models. *Med Decis Making* 2006;26(6):565-74. doi: 10.1177/0272989X06295361 [published Online First: 2006/11/14]
19. Steyerberg EW, Vergouwe Y. Towards better clinical prediction models: seven steps for development and an ABCD for validation. *Eur Heart J* 2014;35(29):1925-31. doi: 10.1093/eurheartj/ehu207 [published Online First: 2014/06/06]
20. Vickers AJ, Van Calster B, Steyerberg EW. Net benefit approaches to the evaluation of prediction models, molecular markers, and diagnostic tests. *BMJ* 2016;352:i6. doi: 10.1136/bmj.i6 [published Online First: 2016/01/27]
21. Ross M, Pang PS, Raslan AM, et al. External retrospective validation of Brain Injury Guidelines criteria and modified guidelines for improved care value in the management of patients with low-risk neurotrauma. *Journal of neurosurgery* 2019;133(6):1880-85.
22. Capron GK, Voights MB, Moore III HR, et al. Not every trauma patient with a radiographic head injury requires transfer for neurosurgical evaluation: Application of the brain injury guidelines to patients transferred to a level 1 trauma center. *The American Journal of Surgery* 2017;214(6):1182-85.
23. Joseph B, Aziz H, Pandit V, et al. Prospective validation of the brain injury guidelines: Managing traumatic brain injury without neurosurgical consultation. *Journal of Trauma & Acute Care Surgery* 2014;77(6):984-88. doi: 10.1097/TA.0000000000000428
24. Ibanez J, Arikian F, Pedraza S, et al. Reliability of clinical guidelines in the detection of patients at risk following mild head injury: results of a prospective study. *Journal of Neurosurgery* 2004;100(5):825-34.
25. Stiell IG, Wells GA, Vandemheen K, et al. The Canadian CT Head Rule for patients with minor head injury. *Lancet* 2001;357(9266):1391-6.
26. Azim A, Jehan FS, Rhee P, et al. Big for small: Validating brain injury guidelines in pediatric traumatic brain injury. *J Trauma Acute Care Surg* 2017;83(6):1200-04. doi: 10.1097/TA.0000000000001611 [published Online First: 2017/06/08]
27. Battle C, Hutchings H, Lovett S, et al. Predicting outcomes after blunt chest wall trauma: development and external validation of a new prognostic model. *Crit Care* 2014;18(3):R98. doi: 10.1186/cc13873 [published Online First: 2014/06/03]
28. Than M, Herbert M, Flaws D, et al. What is an acceptable risk of major adverse cardiac event in chest pain patients soon after discharge from the Emergency Department?: A clinical survey. *International Journal of Cardiology* 2013;166(3):752-54. doi: 10.1016/j.ijcard.2012.09.171
29. O'Keefe ST. A cross-sectional study of doctors', managers' and public representatives' views regarding acceptable level of risk in discharges from the emergency department. *QJM: An International Journal of Medicine* 2014;108(7):533-38. doi: 10.1093/qjmed/hcu246




Figure 1: STROBE flow diagram of selection of study population

139x160mm (600 x 600 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60





Figure 2: Calibration slope of original and re-calibration prognostic model

176x280mm (600 x 600 DPI)

i) Use of prognostic model



ii) Use of the BIG criteria




Figure 3: Decision Curve analysis

209x233mm (600 x 600 DPI)

## Supplementary Material 1: The Brain Injury Guideline (BIG) criteria:

	BIG1 (Discharge from ED after 6 hours)	BIG2 (Non-specialist hospital admission)	BIG3* (Specialist hospital admission)
Neurological Examination	GCS13-15 Normal pupils No Focal Neurological deficit	GCS13-15 Normal pupils No Focal Neurological deficit	GCS<13 Or Abnormal pupils Or Focal Neurological deficit
Intoxicated	No	No/Yes	No/Yes
Anticoagulants or Anti-platelets	No	No	Yes
Skull Fracture	No	Non-displaced	Displaced
Intracranial Bleed	Subdural Haemorrhage <5mm Or Extradural Haemorrhage <5mm Or 1 Intraparenchymal Haemorrhage <5mm Or Trace Subarachnoid Haemorrhage	Subdural Haemorrhage 5-7mm Or Extradural Haemorrhage 5-7mm Or 1-2 Intraparenchymal Haemorrhages 5-7mm Or Localised Subarachnoid Haemorrhage	All other injuries
Intra-ventricular Haemorrhage	No	No	Yes

\*Patients must fulfil all the criteria of BIG1 or BIG2 to be categorised as such and are otherwise automatically in BIG3

## Supplementary Material 2: Risk Score

Factor	Coefficient	Risk Score Value
Preinjury Anti-coagulation or anti-platelets	0.3	1
GCS		
15	0 (Vs)	<b>GCS 15 0</b>
14	0.4	<b>GCS 14 1</b>
13	0.7	<b>GCS 13 2</b>
Normal first Neurological Examination	0.45	<b>Abnormal 1.5</b>
Number of Injuries on CT		
<b>1</b>	0 (Vs)	<b>1 0</b>
<b>2</b>	0.25	<b>2 1</b>
<b>3</b>	0.4	<b>3 1</b>
<b>4</b>	0.8	<b>4 3</b>
<b>5</b>	0.9	<b>5 3</b>
<b>Diffuse</b>	0.3	<b>Diffuse 1</b>
Injury severity on CT		
<b>1</b> simple skull fracture	0 (Vs)	<b>1 0</b>
<b>2</b> complex Skull Fracture	0.3	<b>2 1</b>
<b>3</b> Marshall IIa 1-2 bleeds < 5mm (total)	0.08	<b>3 0</b>
<b>4</b> Marshall IIb bleeds ≥ 5mm	0.7	<b>4 2</b>
<b>5</b> Marshall III/IV	1.7	<b>5 5</b>
<b>6</b> Marshall VI	2.7	<b>6 9</b>
<b>7</b> Brain stem/Cerebellar	1.7	<b>7 5</b>
ISS (body regions excluding head)	0.2	<b>Up to 2 non-significant extra-cranial injuries** 0</b> <b>Any significant extra-cranial injury or 3 or more injuries 2</b>
Hb	-0.01	Not included in risk score
Constant	-1.38	

\* Injuries exclude superficial lacerations and abrasions and a significant extra-cranial injury is defined as any injury requiring inpatient care

## Supplementary Material 3: Procedure for Multi-imputation of missing data

Missing data was assumed to be missing at random. Thirty-five imputed datasets were created on the basis of the fraction of missing information (around 35% of patients had missing data in at least one predictor variable in the extended prognostic model). The imputation model contained the composite outcome of deterioration, all predictive factors in the prognostic model, and additionally, age and sex. Model performance was averaged across imputed data sets.

Supplementary Material 4: Performance of BIG criteria across all 3 risk stratification categories

<b>BIG Criteria Performance</b>			
<b>BIG 1 (Discharge from ED after 6 hours)</b>			
N=921	<b>Deteriorated</b>	<b>Didn't deteriorate</b>	
N=105 Composite deterioration	12	93	
Neurosurg/Death/intubation	6	99	
<b>BIG 2 (non-specialist admission)</b>			
N=921	<b>Deteriorated</b>	<b>Didn't deteriorate</b>	
N=82 Composite deterioration	10	72	
Neurosurg/Death/intubation	8	74	
<b>BIG 3 (Neurosurgical Admission, repeat CT imaging)</b>			
N=921	<b>Deteriorated</b>	<b>Didn't deteriorate</b>	
N=734 Composite deterioration	200	534	
Neurosurg/Death/intubation	164	570	

## Supplementary Material 5: Characteristics of patients recommended for discharge

Population Characteristic	Category Mean (SD), min-max or N (%)	BIG 1 N=105	Recommended Discharge HSC DR N=34
Age	Years	52 (17.5) 17-84	48.4 (18.2) 25-80
Age	≥65	24 (22.9%)	6 (17.7%)
GCS	15	75 (71.4%)	34 (100%)
	14	24 (22.9%)	
	13	6 (5.7%)	
Intoxicated	Yes	0 (0%)	7 (20.6%)
Haemoglobin	Grams/litre	137 (SD 17.4) 8.3-16.3	143 (SD12.5) 12.7-15.6
Number of Injuries on CT	1	105 (100%)	34 (100%)
	2		
	3		
	4		
	5		
	Multiple diffuse injury/>5		
Injury severity on CT (Modified Marshall Classification described in detail supplementary Material )	1) Simple Skull Fractures	105 (100%)	1 (2.9%)
	3)1-2 bleeds < 5mm (total)		33 (97.1%)
ISS	Body regions excluding head	15.4 (12.1) 1-59	4.1 (2) 1-8



## Supplementary Material 6: Subgroup analysis Marshall Classification &lt;3

<b>HSC DR</b>			
<b>N=800</b>	<b>Deteriorated</b>	<b>Didn't deteriorate</b>	
Risk=0	0	34	Sensitivity 100% (96.6-100%)
Risk>0	137	629	Specificity 5.1% (3.6-7.2)

<b>BIG 1 (Discharge after 6 hours)</b>			
<b>N=770</b>	<b>Deteriorated</b>	<b>Didn't deteriorate</b>	
BIG1	12	93	Sensitivity 90.8% (84.2-95)
BIG 2/3	119	546	Specificity 14.6% (12-17.6)

## Supplementary Material 7: Recalibrated prognostic model

<b>Factor</b>	<b>Coefficient (optimism adjusted)</b>
Preinjury Anti-coagulation or anti-platelets	0.15
GCS	
15	0 (Vs)
14	0.2
13	0.36
Normal first Neurological Examination	0.23
Number of Injuries on CT	
<b>1</b>	0 (Vs)
<b>2</b>	0.13
<b>3</b>	0.2
<b>4</b>	0.41
<b>5</b>	0.46
<b>Diffuse</b>	0.15
Injury severity on CT*	
<b>1</b> simple skull fracture	0 (Vs)
<b>2</b> complex Skull Fracture	0.15
<b>3</b> 1-2 bleeds < 5mm	0.04
<b>4</b> Marshall II	0.36
<b>5</b> Marshall III/IV	0.87
<b>6</b> Marshall VI	1.38
<b>7</b> Brain stem/Cerebellar	0.87
ISS (body regions excluding head)	0.1
Hb	-0.005
Constant	-3.68

## Supplementary Material 8: Sensitivity analysis with 2 sites used in derivation study excluded

HSC DR			
N=893	Deteriorated	Didn't deteriorate	
Risk=0	0	31	Sensitivity 100% (98-100)
Risk>0	221	641	Specificity 4.6% (3.2-6.6%)

Confidential: For Review Only

## Supplementary Material 9: The CENTER-TBI participants and investigators:

Cecilia Åkerlund<sup>1</sup>, Krisztina Amrein<sup>2</sup>, Nada Andelic<sup>3</sup>, Lasse Andreassen<sup>4</sup>, Audny Anke<sup>5</sup>, Anna Antoni<sup>6</sup>, Gérard Audibert<sup>7</sup>, Philippe Azouvi<sup>8</sup>, Maria Luisa Azzolini<sup>9</sup>, Ronald Bartels<sup>10</sup>, Pál Barzó<sup>11</sup>, Romuald Beauvais<sup>12</sup>, Ronny Beer<sup>13</sup>, Bo-Michael Bellander<sup>14</sup>, Antonio Belli<sup>15</sup>, Habib Benali<sup>16</sup>, Maurizio Bernardino<sup>17</sup>, Luigi Beretta<sup>9</sup>, Morten Blaabjerg<sup>18</sup>, Peter Bragge<sup>19</sup>, Alexandra Brazinova<sup>20</sup>, Vibeke Brinck<sup>21</sup>, Joanne Brooker<sup>22</sup>, Camilla Brorsson<sup>23</sup>, Andras Buki<sup>24</sup>, Monika Bullinger<sup>25</sup>, Manuel Cabeleira<sup>26</sup>, Alessio Caccioppola<sup>27</sup>, Emiliana Calappi<sup>27</sup>, Maria Rosa Calvi<sup>9</sup>, Peter Cameron<sup>28</sup>, Guillermo Carbayo Lozano<sup>29</sup>, Marco Carbonara<sup>27</sup>, Simona Cavallo<sup>17</sup>, Giorgio Chevallard<sup>30</sup>, Arturo Chierogato<sup>30</sup>, Giuseppe Citerio<sup>31, 32</sup>, Hans Clusmann<sup>33</sup>, Mark Coburn<sup>34</sup>, Jonathan Coles<sup>35</sup>, Jamie D. Cooper<sup>36</sup>, Marta Correia<sup>37</sup>, Amra Čović<sup>38</sup>, Nicola Curry<sup>39</sup>, Endre Czeiter<sup>24</sup>, Marek Czosnyka<sup>26</sup>, Claire Dahyot-Fizelier<sup>40</sup>, Paul Dark<sup>41</sup>, Helen Dawes<sup>42</sup>, Véronique De Keyser<sup>43</sup>, Vincent Degos<sup>16</sup>, Francesco Della Corte<sup>44</sup>, Hugo den Boogert<sup>10</sup>, Bart Depreitere<sup>45</sup>, Đula Đilvesi<sup>46</sup>, Abhishek Dixit<sup>47</sup>, Emma Donoghue<sup>22</sup>, Jens Dreier<sup>48</sup>, Guy-Loup Dulière<sup>49</sup>, Ari Ercole<sup>47</sup>, Patrick Esser<sup>42</sup>, Erzsébet Ezer<sup>50</sup>, Martin Fabricius<sup>51</sup>, Valery L. Feigin<sup>52</sup>, Kelly Foks<sup>53</sup>, Shirin Frisvold<sup>54</sup>, Alex Furmanov<sup>55</sup>, Pablo Gagliardo<sup>56</sup>, Damien Galanaud<sup>16</sup>, Dashiell Gantner<sup>28</sup>, Guoyi Gao<sup>57</sup>, Pradeep George<sup>58</sup>, Alexandre Ghuysen<sup>59</sup>, Lelde Giga<sup>60</sup>, Ben Glocker<sup>61</sup>, Jagoš Golubovic<sup>46</sup>, Pedro A. Gomez<sup>62</sup>, Johannes Gratz<sup>63</sup>, Benjamin Gravesteijn<sup>64</sup>, Francesca Grossi<sup>44</sup>, Russell L. Gruen<sup>65</sup>, Deepak Gupta<sup>66</sup>, Juanita A. Haagsma<sup>64</sup>, Iain Haitsma<sup>67</sup>, Raimund Helbok<sup>13</sup>, Eirik Helseth<sup>68</sup>, Lindsay Horton<sup>69</sup>, Jilske Huijben<sup>64</sup>, Peter J. Hutchinson<sup>70</sup>, Bram Jacobs<sup>71</sup>, Stefan Jankowski<sup>72</sup>, Mike Jarrett<sup>21</sup>, Ji-yao Jiang<sup>58</sup>, Faye Johnson<sup>73</sup>, Kelly Jones<sup>52</sup>, Mladen Karan<sup>46</sup>, Angelos G. Kolias<sup>70</sup>, Erwin Kompanje<sup>74</sup>, Daniel Kondziella<sup>51</sup>, Evgenios Kornaropoulos<sup>47</sup>, Lars-Owe Koskinen<sup>75</sup>, Noémi Kovács<sup>76</sup>, Ana Kowark<sup>77</sup>, Alfonso Lagares<sup>62</sup>, Linda Lanyon<sup>58</sup>, Steven Laureys<sup>78</sup>, Fiona Lecky<sup>79, 80</sup>, Didier Ledoux<sup>78</sup>, Rolf Lefering<sup>81</sup>, Valerie Legrand<sup>82</sup>, Aurelie Lejeune<sup>83</sup>, Leon Levi<sup>84</sup>, Roger Lightfoot<sup>85</sup>, Hester Lingsma<sup>64</sup>, Andrew I.R. Maas<sup>43</sup>, Ana M. Castaño-León<sup>62</sup>, Marc Maegele<sup>86</sup>, Marek Majdan<sup>20</sup>, Alex Manara<sup>87</sup>, Geoffrey Manley<sup>88</sup>, Costanza Martino<sup>89</sup>, Hugues Maréchal<sup>49</sup>, Julia Mattern<sup>90</sup>, Catherine McMahon<sup>91</sup>, Béla Melegh<sup>92</sup>, David Menon<sup>47</sup>, Tomas Menovsky<sup>43</sup>, Ana Mikolic<sup>64</sup>, Benoit Misset<sup>78</sup>, Visakh Muraleedharan<sup>58</sup>, Lynnette Murray<sup>28</sup>, Ancuta Negru<sup>93</sup>, David Nelson<sup>1</sup>, Virginia Newcombe<sup>47</sup>, Daan Nieboer<sup>64</sup>, József Nyirádi<sup>2</sup>, Otesile Olubukola<sup>79</sup>, Matej Oresic<sup>94</sup>, Fabrizio Ortolano<sup>27</sup>, Aarno Palotie<sup>95, 96, 97</sup>, Paul M. Parizel<sup>98</sup>, Jean-François Payen<sup>99</sup>, Natascha Perera<sup>12</sup>, Vincent Perlbarg<sup>16</sup>, Paolo Persona<sup>100</sup>, Wilco Peul<sup>101</sup>, Anna Piippo-Karjalainen<sup>102</sup>, Matti Pirinen<sup>95</sup>, Horia Ples<sup>93</sup>, Suzanne Polinder<sup>64</sup>, Inigo Pomposo<sup>29</sup>, Jussi P. Posti<sup>103</sup>, Louis Puybasset<sup>104</sup>, Andreea Radoi<sup>105</sup>, Arminas Ragauskas<sup>106</sup>, Rahul Raj<sup>102</sup>, Malinka Rambadagalla<sup>107</sup>, Jonathan Rhodes<sup>108</sup>, Sylvia Richardson<sup>109</sup>, Sophie Richter<sup>47</sup>, Samuli Ripatti<sup>95</sup>, Saulius Rocka<sup>106</sup>, Cecilie Roe<sup>110</sup>, Olav Roise<sup>111, 112</sup>, Jonathan Rosand<sup>113</sup>, Jeffrey V. Rosenfeld<sup>114</sup>, Christina Rosenlund<sup>115</sup>, Guy Rosenthal<sup>55</sup>, Rolf Rossaint<sup>77</sup>, Sandra Rossi<sup>100</sup>, Daniel Rueckert<sup>61</sup>, Martin Rusnák<sup>116</sup>, Juan Sahuquillo<sup>105</sup>, Oliver Sakowitz<sup>90, 117</sup>, Renan Sanchez-Porras<sup>117</sup>, Janos Sandor<sup>118</sup>, Nadine Schäfer<sup>81</sup>, Silke Schmidt<sup>119</sup>, Herbert Schoechl<sup>120</sup>, Guus Schoonman<sup>121</sup>, Rico Frederik Schou<sup>122</sup>, Elisabeth Schwendenwein<sup>6</sup>, Charlie Sewalt<sup>64</sup>, Toril Skandsen<sup>123, 124</sup>, Peter Smielewski<sup>26</sup>, Abayomi Sorinola<sup>125</sup>, Emmanuel Stamatakis<sup>47</sup>, Simon Stanworth<sup>39</sup>, Robert Stevens<sup>126</sup>, William Stewart<sup>127</sup>, Ewout W. Steyerberg<sup>64, 128</sup>, Nino Stocchetti<sup>129</sup>, Nina Sundström<sup>130</sup>, Riikka Takala<sup>131</sup>, Viktória Tamás<sup>125</sup>, Tomas Tamosuitis<sup>132</sup>, Mark Steven Taylor<sup>20</sup>, Braden Te Ao<sup>52</sup>, Olli

1  
2  
3 Tenovuo<sup>103</sup>, Alice Theadom<sup>52</sup>, Matt Thomas<sup>87</sup>, Dick Tibboel<sup>133</sup>, Marjolein Timmers<sup>74</sup>,  
4 Christos Toliás<sup>134</sup>, Tony Trapani<sup>28</sup>, Cristina Maria Tudora<sup>93</sup>, Andreas Unterberg<sup>90</sup>, Peter  
5 Vajkoczy<sup>135</sup>, Shirley Vallance<sup>28</sup>, Egils Valeinis<sup>60</sup>, Zoltán Vámos<sup>50</sup>, Mathieu van der Jagt<sup>136</sup>,  
6 Gregory Van der Steen<sup>43</sup>, Joukje van der Naalt<sup>71</sup>, Jeroen T.J.M. van Dijck<sup>101</sup>,  
7 Thomas A. van Essen<sup>101</sup>, Wim Van Hecke<sup>137</sup>, Caroline van Heugten<sup>138</sup>,  
8 Dominique Van Praag<sup>139</sup>, Thijs Vande Vyvere<sup>137</sup>, Roel P. J. van Wijk<sup>101</sup>, Alessia Vargiolu<sup>32</sup>,  
9 Emmanuel Vega<sup>83</sup>, Kimberley Velt<sup>64</sup>, Jan Verheyden<sup>137</sup>, Paul M. Vespa<sup>140</sup>, Anne Vik<sup>123, 141</sup>,  
10 Rimantas Vilcinis<sup>132</sup>, Victor Volovici<sup>67</sup>, Nicole von Steinbüchel<sup>38</sup>, Daphne Voormolen<sup>64</sup>,  
11 Petar Vulekovic<sup>46</sup>, Kevin K.W. Wang<sup>142</sup>, Eveline Wieggers<sup>64</sup>, Guy Williams<sup>47</sup>, Lindsay Wilson<sup>69</sup>,  
12 Stefan Winzeck<sup>47</sup>, Stefan Wolf<sup>143</sup>, Zhihui Yang<sup>113</sup>, Peter Ylén<sup>144</sup>, Alexander Younsi<sup>90</sup>, Frederick  
13 A. Zeiler<sup>47,145</sup>, Veronika Zelinkova<sup>20</sup>, Agate Ziverte<sup>60</sup>, Tommaso Zoerle<sup>27</sup>  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1
- 2
- 3
- 4 <sup>1</sup> Department of Physiology and Pharmacology, Section of Perioperative Medicine and Intensive
- 5 Care, Karolinska Institutet, Stockholm, Sweden
- 6 <sup>2</sup> János Szentágothai Research Centre, University of Pécs, Pécs, Hungary
- 7 <sup>3</sup> Division of Surgery and Clinical Neuroscience, Department of Physical Medicine and
- 8 Rehabilitation, Oslo University Hospital and University of Oslo, Oslo, Norway
- 9 <sup>4</sup> Department of Neurosurgery, University Hospital Northern Norway, Tromsø, Norway
- 10 <sup>5</sup> Department of Physical Medicine and Rehabilitation, University Hospital Northern Norway,
- 11 Tromsø, Norway
- 12 <sup>6</sup> Trauma Surgery, Medical University Vienna, Vienna, Austria
- 13 <sup>7</sup> Department of Anesthesiology & Intensive Care, University Hospital Nancy, Nancy, France
- 14 <sup>8</sup> Raymond Poincaré hospital, Assistance Publique – Hôpitaux de Paris, Paris, France
- 15 <sup>9</sup> Department of Anesthesiology & Intensive Care, S Raffaele University Hospital, Milan, Italy
- 16 <sup>10</sup> Department of Neurosurgery, Radboud University Medical Center, Nijmegen, The
- 17 Netherlands
- 18 <sup>11</sup> Department of Neurosurgery, University of Szeged, Szeged, Hungary
- 19 <sup>12</sup> International Projects Management, ARTTIC, Munchen, Germany
- 20 <sup>13</sup> Department of Neurology, Neurological Intensive Care Unit, Medical University of
- 21 Innsbruck, Innsbruck, Austria
- 22 <sup>14</sup> Department of Neurosurgery & Anesthesia & intensive care medicine, Karolinska
- 23 University Hospital, Stockholm, Sweden
- 24 <sup>15</sup> NIHR Surgical Reconstruction and Microbiology Research Centre, Birmingham, UK
- 25 <sup>16</sup> Anesthésie-Réanimation, Assistance Publique – Hôpitaux de Paris, Paris, France
- 26 <sup>17</sup> Department of Anesthesia & ICU, AOU Città della Salute e della Scienza di Torino -
- 27 Orthopedic and Trauma Center, Torino, Italy
- 28 <sup>18</sup> Department of Neurology, Odense University Hospital, Odense, Denmark
- 29 <sup>19</sup> BehaviourWorks Australia, Monash Sustainability Institute, Monash University, Victoria,
- 30 Australia
- 31 <sup>20</sup> Department of Public Health, Faculty of Health Sciences and Social Work, Trnava
- 32 University, Trnava, Slovakia
- 33 <sup>21</sup> Quesgen Systems Inc., Burlingame, California, USA
- 34 <sup>22</sup> Australian & New Zealand Intensive Care Research Centre, Department of Epidemiology
- 35 and Preventive Medicine, School of Public Health and Preventive Medicine, Monash
- 36 University, Melbourne, Australia
- 37 <sup>23</sup> Department of Surgery and Perioperative Science, Umeå University, Umeå, Sweden
- 38 <sup>24</sup> Department of Neurosurgery, Medical School, University of Pécs, Hungary and
- 39 Neurotrauma Research Group, János Szentágothai Research Centre, University of Pécs,
- 40 Hungary
- 41 <sup>25</sup> Department of Medical Psychology, Universitätsklinikum Hamburg-Eppendorf, Hamburg,
- 42 Germany
- 43 <sup>26</sup> Brain Physics Lab, Division of Neurosurgery, Dept of Clinical Neurosciences, University of
- 44 Cambridge, Addenbrooke's Hospital, Cambridge, UK
- 45 <sup>27</sup> Neuro ICU, Fondazione IRCCS Cà Granda Ospedale Maggiore Policlinico, Milan, Italy
- 46 <sup>28</sup> ANZIC Research Centre, Monash University, Department of Epidemiology and Preventive
- 47 Medicine, Melbourne, Victoria, Australia
- 48 <sup>29</sup> Department of Neurosurgery, Hospital of Cruces, Bilbao, Spain
- 49 <sup>30</sup> NeuroIntensive Care, Niguarda Hospital, Milan, Italy
- 50 <sup>31</sup> School of Medicine and Surgery, Università Milano Bicocca, Milano, Italy
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- <sup>32</sup> NeuroIntensive Care, ASST di Monza, Monza, Italy
- <sup>33</sup> Department of Neurosurgery, Medical Faculty RWTH Aachen University, Aachen, Germany
- <sup>34</sup> Department of Anesthesiology and Intensive Care Medicine, University Hospital Bonn, Bonn, Germany
- <sup>35</sup> Department of Anesthesia & Neurointensive Care, Cambridge University Hospital NHS Foundation Trust, Cambridge, UK
- <sup>36</sup> School of Public Health & PM, Monash University and The Alfred Hospital, Melbourne, Victoria, Australia
- <sup>37</sup> Radiology/MRI department, MRC Cognition and Brain Sciences Unit, Cambridge, UK
- <sup>38</sup> Institute of Medical Psychology and Medical Sociology, Universitätsmedizin Göttingen, Göttingen, Germany
- <sup>39</sup> Oxford University Hospitals NHS Trust, Oxford, UK
- <sup>40</sup> Intensive Care Unit, CHU Poitiers, Poitiers, France
- <sup>41</sup> University of Manchester NIHR Biomedical Research Centre, Critical Care Directorate, Salford Royal Hospital NHS Foundation Trust, Salford, UK
- <sup>42</sup> Movement Science Group, Faculty of Health and Life Sciences, Oxford Brookes University, Oxford, UK
- <sup>43</sup> Department of Neurosurgery, Antwerp University Hospital and University of Antwerp, Edegem, Belgium
- <sup>44</sup> Department of Anesthesia & Intensive Care, Maggiore Della Carità Hospital, Novara, Italy
- <sup>45</sup> Department of Neurosurgery, University Hospitals Leuven, Leuven, Belgium
- <sup>46</sup> Department of Neurosurgery, Clinical centre of Vojvodina, Faculty of Medicine, University of Novi Sad, Novi Sad, Serbia
- <sup>47</sup> Division of Anaesthesia, University of Cambridge, Addenbrooke's Hospital, Cambridge, UK
- <sup>48</sup> Center for Stroke Research Berlin, Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Berlin, Germany
- <sup>49</sup> Intensive Care Unit, CHR Citadelle, Liège, Belgium
- <sup>50</sup> Department of Anaesthesiology and Intensive Therapy, University of Pécs, Pécs, Hungary
- <sup>51</sup> Departments of Neurology, Clinical Neurophysiology and Neuroanesthesiology, Region Hovedstaden Rigshospitalet, Copenhagen, Denmark
- <sup>52</sup> National Institute for Stroke and Applied Neurosciences, Faculty of Health and Environmental Studies, Auckland University of Technology, Auckland, New Zealand
- <sup>53</sup> Department of Neurology, Erasmus MC, Rotterdam, the Netherlands
- <sup>54</sup> Department of Anesthesiology and Intensive care, University Hospital Northern Norway, Tromsø, Norway
- <sup>55</sup> Department of Neurosurgery, Hadassah-hebrew University Medical center, Jerusalem, Israel
- <sup>56</sup> Fundación Instituto Valenciano de Neurorehabilitación (FIVAN), Valencia, Spain
- <sup>57</sup> Department of Neurosurgery, Shanghai Renji hospital, Shanghai Jiaotong University/school of medicine, Shanghai, China
- <sup>58</sup> Karolinska Institutet, INCF International Neuroinformatics Coordinating Facility, Stockholm, Sweden
- <sup>59</sup> Emergency Department, CHU, Liège, Belgium
- <sup>60</sup> Neurosurgery clinic, Pauls Stradins Clinical University Hospital, Riga, Latvia
- <sup>61</sup> Department of Computing, Imperial College London, London, UK
- <sup>62</sup> Department of Neurosurgery, Hospital Universitario 12 de Octubre, Madrid, Spain

- 1
- 2
- 3
- 4 63 Department of Anesthesia, Critical Care and Pain Medicine, Medical University of Vienna,  
5 Austria
- 6 64 Department of Public Health, Erasmus Medical Center-University Medical Center,  
7 Rotterdam, The Netherlands
- 8 65 College of Health and Medicine, Australian National University, Canberra, Australia
- 9 66 Department of Neurosurgery, Neurosciences Centre & JPN Apex trauma centre, All India  
10 Institute of Medical Sciences, New Delhi-110029, India
- 11 67 Department of Neurosurgery, Erasmus MC, Rotterdam, the Netherlands
- 12 68 Department of Neurosurgery, Oslo University Hospital, Oslo, Norway
- 13 69 Division of Psychology, University of Stirling, Stirling, UK
- 14 70 Division of Neurosurgery, Department of Clinical Neurosciences, Addenbrooke's Hospital  
15 & University of Cambridge, Cambridge, UK
- 16 71 Department of Neurology, University of Groningen, University Medical Center Groningen,  
17 Groningen, Netherlands
- 18 72 Neurointensive Care , Sheffield Teaching Hospitals NHS Foundation Trust, Sheffield, UK
- 19 73 Salford Royal Hospital NHS Foundation Trust Acute Research Delivery Team, Salford, UK
- 20 74 Department of Intensive Care and Department of Ethics and Philosophy of Medicine,  
21 Erasmus Medical Center, Rotterdam, The Netherlands
- 22 75 Department of Clinical Neuroscience, Neurosurgery, Umeå University, Umeå, Sweden
- 23 76 Hungarian Brain Research Program - Grant No. KTIA\_13\_NAP-A-II/8, University of Pécs,  
24 Pécs, Hungary
- 25 77 Department of Anaesthesiology, University Hospital of Aachen, Aachen, Germany
- 26 78 Cyclotron Research Center , University of Liège, Liège, Belgium
- 27 79 Centre for Urgent and Emergency Care Research (CURE), Health Services Research  
28 Section, School of Health and Related Research (SchARR), University of  
29 Sheffield, Sheffield, UK
- 30 80 Emergency Department, Salford Royal Hospital, Salford UK
- 31 81 Institute of Research in Operative Medicine (IFOM), Witten/Herdecke University, Cologne,  
32 Germany
- 33 82 VP Global Project Management CNS, ICON, Paris, France
- 34 83 Department of Anesthesiology-Intensive Care, Lille University Hospital, Lille, France
- 35 84 Department of Neurosurgery, Rambam Medical Center, Haifa, Israel
- 36 85 Department of Anesthesiology & Intensive Care, University Hospitals Southampton NHS  
37 Trust, Southampton, UK
- 38 86 Cologne-Merheim Medical Center (CMMC), Department of Traumatology, Orthopedic  
39 Surgery and Sportmedicine, Witten/Herdecke University, Cologne, Germany
- 40 87 Intensive Care Unit, Southmead Hospital, Bristol, Bristol, UK
- 41 88 Department of Neurological Surgery, University of California, San Francisco, California,  
42 USA
- 43 89 Department of Anesthesia & Intensive Care, M. Bufalini Hospital, Cesena, Italy
- 44 90 Department of Neurosurgery, University Hospital Heidelberg, Heidelberg, Germany
- 45 91 Department of Neurosurgery, The Walton centre NHS Foundation Trust, Liverpool, UK
- 46 92 Department of Medical Genetics, University of Pécs, Pécs, Hungary
- 47 93 Department of Neurosurgery, Emergency County Hospital Timisoara , Timisoara, Romania
- 48 94 School of Medical Sciences, Örebro University, Örebro, Sweden
- 49 95 Institute for Molecular Medicine Finland, University of Helsinki, Helsinki, Finland
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60



- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- <sup>96</sup> Analytic and Translational Genetics Unit, Department of Medicine; Psychiatric & Neurodevelopmental Genetics Unit, Department of Psychiatry; Department of Neurology, Massachusetts General Hospital, Boston, MA, USA
- <sup>97</sup> Program in Medical and Population Genetics; The Stanley Center for Psychiatric Research, The Broad Institute of MIT and Harvard, Cambridge, MA, USA
- <sup>98</sup> Department of Radiology, University of Antwerp, Edegem, Belgium
- <sup>99</sup> Department of Anesthesiology & Intensive Care, University Hospital of Grenoble, Grenoble, France
- <sup>100</sup> Department of Anesthesia & Intensive Care, Azienda Ospedaliera Università di Padova, Padova, Italy
- <sup>101</sup> Dept. of Neurosurgery, Leiden University Medical Center, Leiden, The Netherlands and Dept. of Neurosurgery, Medical Center Haaglanden, The Hague, The Netherlands
- <sup>102</sup> Department of Neurosurgery, Helsinki University Central Hospital
- <sup>103</sup> Division of Clinical Neurosciences, Department of Neurosurgery and Turku Brain Injury Centre, Turku University Hospital and University of Turku, Turku, Finland
- <sup>104</sup> Department of Anesthesiology and Critical Care, Pitié -Salpêtrière Teaching Hospital, Assistance Publique, Hôpitaux de Paris and University Pierre et Marie Curie, Paris, France
- <sup>105</sup> Neurotraumatology and Neurosurgery Research Unit (UNINN), Vall d'Hebron Research Institute, Barcelona, Spain
- <sup>106</sup> Department of Neurosurgery, Kaunas University of technology and Vilnius University, Vilnius, Lithuania
- <sup>107</sup> Department of Neurosurgery, Rezekne Hospital, Latvia
- <sup>108</sup> Department of Anaesthesia, Critical Care & Pain Medicine NHS Lothian & University of Edinburgh, Edinburgh, UK
- <sup>109</sup> Director, MRC Biostatistics Unit, Cambridge Institute of Public Health, Cambridge, UK
- <sup>110</sup> Department of Physical Medicine and Rehabilitation, Oslo University Hospital/University of Oslo, Oslo, Norway
- <sup>111</sup> Division of Orthopedics, Oslo University Hospital, Oslo, Norway
- <sup>112</sup> Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo, Norway
- <sup>113</sup> Broad Institute, Cambridge MA Harvard Medical School, Boston MA, Massachusetts General Hospital, Boston MA, USA
- <sup>114</sup> National Trauma Research Institute, The Alfred Hospital, Monash University, Melbourne, Victoria, Australia
- <sup>115</sup> Department of Neurosurgery, Odense University Hospital, Odense, Denmark
- <sup>116</sup> International Neurotrauma Research Organisation, Vienna, Austria
- <sup>117</sup> Klinik für Neurochirurgie, Klinikum Ludwigsburg, Ludwigsburg, Germany
- <sup>118</sup> Division of Biostatistics and Epidemiology, Department of Preventive Medicine, University of Debrecen, Debrecen, Hungary
- <sup>119</sup> Department Health and Prevention, University Greifswald, Greifswald, Germany
- <sup>120</sup> Department of Anaesthesiology and Intensive Care, AUVA Trauma Hospital, Salzburg, Austria
- <sup>121</sup> Department of Neurology, Elisabeth-TweeSteden Ziekenhuis, Tilburg, the Netherlands
- <sup>122</sup> Department of Neuroanesthesia and Neurointensive Care, Odense University Hospital, Odense, Denmark
- <sup>123</sup> Department of Neuromedicine and Movement Science, Norwegian University of Science and Technology, NTNU, Trondheim, Norway

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50
- 124 Department of Physical Medicine and Rehabilitation, St.Olavs Hospital, Trondheim University Hospital, Trondheim, Norway
- 125 Department of Neurosurgery, University of Pécs, Pécs, Hungary
- 126 Division of Neuroscience Critical Care, John Hopkins University School of Medicine, Baltimore, USA
- 127 Department of Neuropathology, Queen Elizabeth University Hospital and University of Glasgow, Glasgow, UK
- 128 Dept. of Department of Biomedical Data Sciences, Leiden University Medical Center, Leiden, The Netherlands
- 129 Department of Pathophysiology and Transplantation, Milan University, and Neuroscience ICU, Fondazione IRCCS Cà Granda Ospedale Maggiore Policlinico, Milano, Italy
- 130 Department of Radiation Sciences, Biomedical Engineering, Umeå University, Umeå, Sweden
- 131 Perioperative Services, Intensive Care Medicine and Pain Management, Turku University Hospital and University of Turku, Turku, Finland
- 132 Department of Neurosurgery, Kaunas University of Health Sciences, Kaunas, Lithuania
- 133 Intensive Care and Department of Pediatric Surgery, Erasmus Medical Center, Sophia Children's Hospital, Rotterdam, The Netherlands
- 134 Department of Neurosurgery, Kings college London, London, UK
- 135 Neurologie, Neurochirurgie und Psychiatrie, Charité – Universitätsmedizin Berlin, Berlin, Germany
- 136 Department of Intensive Care Adults, Erasmus MC– University Medical Center Rotterdam, Rotterdam, the Netherlands
- 137 icoMetrix NV, Leuven, Belgium
- 138 Movement Science Group, Faculty of Health and Life Sciences, Oxford Brookes University, Oxford, UK
- 139 Psychology Department, Antwerp University Hospital, Edegem, Belgium
- 140 Director of Neurocritical Care, University of California, Los Angeles, USA
- 141 Department of Neurosurgery, St.Olavs Hospital, Trondheim University Hospital, Trondheim, Norway
- 142 Department of Emergency Medicine, University of Florida, Gainesville, Florida, USA
- 143 Department of Neurosurgery, Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Berlin, Germany
- 144 VTT Technical Research Centre, Tampere, Finland
- 145 Section of Neurosurgery, Department of Surgery, Rady Faculty of Health Sciences, University of Manitoba, Winnipeg, MB, Canada

51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Åkerlund	Cecilia	cecilia.ai.akerlund@gmail.com
Amrein	Krisztina	tina.amrein84@gmail.com
Andelic	Nada	NADAND@ous-hf.no
Andreassen	Lasse	Lasse.Andreassen@unn.no
Anke	Audny	Audny.anke@unn.no
Antoni	Anna	anna.antoni@meduniwien.ac.at
Audibert	Gérard	g.audibert@chu-nancy.fr

1			
2			
3	Azouvi	Philippe	philippe.azouvi@rpc.aphp.fr
4	Azzolini	Maria Luisa	azzolini.marialuisa@hsr.it
5	Bartels	Ronald	Ronald.Bartels@radboudumc.nl
6	Barzó	Pál	pbarzo@gmail.com
7	Beauvais	Romuald	beauvais@arttic.eu
8	Beer	Ronny	ronny.beer@i-med.ac.at
9	Bellander	Bo-Michael	bo-michael.bellander@karolinska.se
10	Belli	Antonio	a.belli@bham.ac.uk
11	Benali	Habib	habib.benali@gmail.com
12	Berardino	Maurizio	maurizio_berardino@fastwebnet.it
13	Beretta	Luigi	beretta.luigi@hsr.it
14	Blaabjerg	Morten	morten.blaabjerg1@rsyd.dk
15	Bragge	Peter	peter.bragge@monash.edu
16	Brazinova	Alexandra	alexandra.brazinova@gmail.com
17	Brinck	Vibeke	vibeke.brinck@quesgen.com
18	Brooker	Joanne	Joanne.Brooker@monash.edu
19	Brorsson	Camilla	Camilla.Brorsson@umu.se
20	Buki	Andras	2saturn@gmail.com
21	Bullinger	Monika	bullinger@uke.de
22	Cabeleira	Manuel	mc916@cam.ac.uk
23	Caccioppola	Alessio	alessio.caccioppola@gmail.com
24	Calappi	Emiliana	calemy02@yahoo.it
25	Calvi	Maria Rosa	calvi.mariarosa@hsr.it
26	Cameron	Peter	peter.cameron@med.monash.edu.au
27	Carbayo Lozano	Guillermo	guillermobilbo@gmail.com
28	Carbonara	Marco	marco.carbonara@gmail.com
29	Castaña-León	Ana M.	ana.maria.castano.leon@gmail.com
30	Cavallo	Simona	cavallosimona1@gmail.com
31	Chevallard	Giorgio	giorgio.chevallard@ospedaleniguarda.it
32	Chierregato	Arturo	<a href="mailto:arturo.chierregato@ospedaleniguarda.it">arturo.chierregato@ospedaleniguarda.it</a>
33	Citerio	Giuseppe	giuseppe.citerio@unimib.it
34	Clusmann	Hans	hclusmann@ukaachen.de
35	Coburn	Mark Steven	mark.coburn@ukbonn.de
36	Coles	Jonathan	jpc44@wbic.cam.ac.uk
37	Cooper	Jamie D.	jamie.cooper@monash.edu
38	Correia	Marta	Marta.Correia@mrc-cbu.cam.ac.uk
39	Čović	Amra	amra.covic@med.uni-goettingen.de
40	Curry	Nicola	nicola.curry@ouh.nhs.uk
41	Czeiter	Endre	endre.czeiter@gmail.com
42	Czosnyka	Marek	mc141@medschl.cam.ac.uk
43	Dahyot-Fizelier	Claire	c.dahyot-fizelier@chu-poitiers.fr
44	Dark	Paul	paul.m.dark@manchester.ac.uk
45	Dawes	Helen	hdawes@brookes.ac.uk
46	De Keyser	Véronique	veronique.dekeyser@uza.be
47	Degos	Vincent	vincent.degoss@aphp.fr
48	Della Corte	Francesco	dellacorte.f@gmail.com
49			
50			
51			
52			
53			
54			
55			
56			
57			
58			
59			
60			

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

den Boogert	Hugo	Hugo.denBoogert@radboudumc.nl
Depreitere	Bart	bart.depreitere@uzleuven.be
Đilvesi	Đula	djuladjilvesi@gmail.com
Dixit	Abhishek	ad825@cam.ac.uk
Donoghue	Emma	emma.donoghue@monash.edu
Dreier	Jens	jens.dreier@charite.de
Dulière	Guy-Loup	glduliere@gmail.com
Ercole	Ari	ae105@cam.ac.uk
Esser	Patrick	pesser@brookes.ac.uk
Ezer	Erzsébet	ezererzsebet@yahoo.com
Fabricius	Martin	fabricius@dadlnet.dk
Feigin	Valery L.	valery.feigin@aut.ac.nz
Foks	Kelly	k.foks@erasmusmc.nl
Frisvold	Shirin	Shirin.Kordasti@unn.no
Furmanov	Alex	alexpuil@yahoo.com
Gagliardo	Pablo	pablog@fivan.org
Galanaud	Damien	galanaud@gmail.com
Gantner	Dashiell	dashiell.gantner@monash.edu
Gao	Guoyi	gao3@sina.com
George	Pradeep	george@incf.org
Ghuysen	Alexandre	A.Ghuysen@chu.ulg.ac.be
Giga	Lelde	lelde.giga@inbox.lv
Glocker	Ben	b.glocker@imperial.ac.uk
Golubović	Jagoš	jagosgolubovic@gmail.com
Gomez	Pedro A.	pagolopez@gmail.com
Gratz	Johannes	johannes.gratz@meduniwien.ac.at
Gravesteijn	Benjamin	b.gravesteijn@erasmusmc.nl
Grossi	Francesca	francesca.grossi@libero.it
Gruen	Russell L.	russell.gruen@anu.edu.au
Gupta	Deepak	drdeepakgupta@gmail.com
Haagsma	Juanita A.	j.haagsma@erasmusmc.nl
Haitsma	Iain	i.haitsma@erasmusmc.nl
Helbok	Raimund	Raimund.Helbok@tirol-kliniken.at
Helseth	Eirik	EHELSETH@ous-hf.no
Horton	Lindsay	lindsay.horton@stir.ac.uk
Huijben	Jilske	j.a.huijben@erasmusmc.nl
Hutchinson	Peter J.	pjah2@cam.ac.uk
Jacobs	Bram	b.jacobs@umcg.nl
Jankowski	Stefan	Stefan.Jankowski@sth.nhs.uk
Jarrett	Mike	mike.jarrett@quesgen.com
Jiang	Ji-yao	jiyaojiang@126.com
Johnson	Faye	faye.johnson@live.co.uk
Jones	Kelly	kejones@aut.ac.nz
Karan	Mladen	mladjokaran@gmail.com
Kolias	Angelos G.	angeloskolias@gmail.com
Kompanje	Erwin	erwinkompanje@me.com

1			
2			
3	Kondziella	Daniel	Daniel.Kondziella@regionh.dk
4	Kornaropoulos	Evgenios	ek481@cam.ac.uk
5	Koskinen	Lars-Owe	Lars-Owe.Koskinen@umu.se
6	Kovács	Noémi	kovacs.noemi@pte.hu
7	Lagares	Alfonso	algadoc@yahoo.com
8	Lanyon	Linda	lindal@incf.org
9	Laureys	Steven	steven.laureys@ulg.ac.be
10	Lecky	Fiona	f.e.lecky@sheffield.ac.uk
11	Ledoux	Didier	dledoux@chu.ulg.ac.be
12	Lefering	Rolf	Rolf.Lefering@uni-wh.de
13	Legrand	Valerie	Valerie.Legrand@iconplc.com
14	Lejeune	Aurelie	aurelie.lejeune@chru-lille.fr
15	Levi	Leon	llevi@rambam.health.gov.il
16	Lightfoot	Roger	Roger.Lightfoot@uhs.nhs.uk
17	Lingsma	Hester	h.lingsma@erasmusmc.nl
18	Maas	Andrew I.R.	andrew.maas@uza.be
19	Maegele	Marc	Marc.Maegele@t-online.de
20	Majdan	Marek	mmajdan@truni.sk
21	Manara	Alex	Alex.Manara@nbt.nhs.uk
22	Manley	Geoffrey	ManleyG@ucsf.edu
23	Maréchal	Hugues	Hugues.Marechal@chrcitadelle.be
24	Martino	Costanza	costmartino74@gmail.com
25	Mattern	Julia	Julia.Mattern@med.uni-heidelberg.de
26	McMahon	Catherine	Catherine.McMahon@thewaltoncentre.nhs.uk
27	Melegh	Béla	bela.melegh@aok.pte.hu
28	Menon	David	dkm13@cam.ac.uk
29	Menovsky	Tomas	tomas.menovsky@uza.be
30	Mikolic	Ana	a.mikolic@erasmusmc.nl
31	Misset	Benoit	Benoit.Misset@chuliege.be
32	Muraleedharan	Visakh	visakh@incf.org
33	Murray	Lynnette	lynnette.murray@monash.edu
34	Nair	Nandesh	nandesh.nair@uza.be
35	Negru	Ancuta	negruancu@gmail.com
36	Nelson	David	david.nelson@karolinska.se
37	Newcombe	Virginia	vfjn2@cam.ac.uk
38	Nieboer	Daan	d.nieboer@erasmusmc.nl
39	Nyirádi	József	nyiradi.jozsef@pte.hu
40	Oresic	Matej	matej.oresic@oru.se
41	Ortolano	Fabrizio	lupeda@gmail.com
42	Otesile	Olubukola	o.otesile@sheffield.ac.uk
43	Palotie	Aarno	aarno.palotie@helsinki.fi
44	Parizel	Paul M.	paul.parizel@uantwerpen.be
45	Payen	Jean-François	Jean-Francois.Payen@ujf-grenoble.fr
46	Perera	Natascha	perera@arttic.eu
47	Perlberg	Vincent	vincent.perlberg@gmail.com
48	Persona	Paolo	ppersona75@gmail.com
49			
50			
51			
52			
53			
54			
55			
56			
57			
58			
59			
60			

1			
2			
3	Peul	Wilco	W.C.Peul@lumc.nl
4	Piippo-Karjalainen	Anna	anna.piippo@hus.fi
5	Pirinen	Matti	matti.pirinen@helsinki.fi
6	Ples	Horia	horia.ples@neuromed.ro
7	Polinder	Suzanne	s.polinder@erasmusmc.nl
8	Pomposo	Inigo	inigo.pomposo@osakidetza.net
9	Posti	Jussi P.	jussi.posti@tyks.fi
10	Puybasset	Louis	louis.puybasset@aphp.fr
11	Rădoi	Andreea	aradoi@neurotrauma.net
12	Ragauskas	Arminas	telematics@ktu.lt
13	Raj	Rahul	rahul.raj@hus.fi
14	Rambadagalla	Malinka	malinka.rambadagalla@gmail.com
15	Rehorčíková	Veronika	rehorcikova@gmail.com
16	Rhodes	Jonathan	jrhodes1@staffmail.ed.ac.uk
17	Richardson	Sylvia	sylvia.richardson@mrc-bsu.cam.ac.uk
18	Richter	Sophie	sr773@cam.ac.uk
19	Ripatti	Samuli	samuli.ripatti@helsinki.fi
20	Rocka	Saulius	saulius.rocka@mf.vu.lt
21	Roe	Cecilie	e.c.t.roe@medisin.uio.no
22	Roise	Olav	olav.roise@medisin.uio.no
23	Rosand	Jonathan	jrosand@partners.org
24	Rosenfeld	Jeffrey	J.Rosenfeld@alfred.org.au
25	Rosenlund	Christina	chrisstenrose@gmail.com
26	Rosenthal	Guy	rosenthalg@hadassah.org.il
27	Rossaint	Rolf	RRossaint@ukaachen.de
28	Rossi	Sandra	sandrarossi0@gmail.com
29	Rueckert	Daniel	d.rueckert@imperial.ac.uk
30	Rusnák	Martin	mrusnak@igeh.org
31	Sahuquillo	Juan	sahuquillo@neurotrauma.net
32	Sakowitz	Oliver	oliver.sakowitz@gmail.com
33	Sanchez-Porras	Renan	renan_md@hotmail.com
34	Sandor	Janos	sandor.janos@sph.unideb.hu
35	Schäfer	Nadine	Nadine.Schaefer@uni-wh.de
36	Schmidt	Silke	silke.schmidt@uni-greifswald.de
37	Schoechl	Herbert	Herbert.Schoechl@auva.at
38	Schoonman	Guus	g.schoonman@tsz.nl
39	Schou	Rico Frederik	rico@mymedic.dk
40	Schwendenwein	Elisabeth	elisabeth.schwendenwein@meduniwien.ac.at
41	Sewalt	Charlie	c.sewalt@erasmusmc.nl
42	Skandsen	Toril	toril.skandsen@ntnu.no
43	Smielewski	Peter	ps10011@cam.ac.uk
44	Sorinola	Abayomi	sorinola_abayomi@hotmail.com
45	Stamatakis	Emmanuel	eas46@cam.ac.uk
46	Stanworth	Simon	simon.stanworth@nhsbt.nhs.uk
47	Kowark	Ana	akowark@ukaachen.de
48	Stevens	Robert	rstevens@jhmi.edu
49			
50			
51			
52			
53			
54			
55			
56			
57			
58			
59			
60			

1			
2			
3	Stewart	William	william.stewart@glasgow.ac.uk
4	Steyerberg	Ewout W.	e.steyerberg@erasmusmc.nl
5	Stocchetti	Nino	stocchet@policlinico.mi.it
6	Sundström	Nina	Nina.Sundstrom@vll.se
7	Takala	Riikka	riikka.takala@tyks.fi
8	Tamás	Viktória	tamas.viktoria@pte.hu
9	Tamosuitis	Tomas	tomas.tamosuitis@kaunoklinikos.lt
10	Taylor	Mark Steven	marktrnava@gmail.com
11	Te Ao	Braden	braden.teao@aut.ac.nz
12	Tenovuo	Olli	olli.tenovuo@tyks.fi
13	Theadom	Alice	alice.theadom@aut.ac.nz
14	Thomas	Matt	Matt.Thomas@nbt.nhs.uk
15	Tibboel	Dick	d.tibboel@erasmusmc.nl
16	Timmers	Marjolijn	mtimmers@hotmail.com
17	Tolias	Christos	christos.tolias@nhs.net
18	Trapani	Tony	tony.trapani@monash.edu
19	Tudora	Cristina Maria	cristina.tudora@neuromed.ro
20	Unterberg	Andreas	Andreas.Unterberg@med.uni-heidelberg.de
21	Vajkoczy	Peter	Peter.Vajkoczy@charite.de
22	Valeinis	Egils	Egils.Valeinis@latnet.lv
23	Vallance	Shirley	S.Vallance@alfred.org.au
24	Vámos	Zoltán	azozoka@gmail.com
25	Van der Jagt	Mathieu	m.vanderjagt@erasmusmc.nl
26	van der Naalt	Joukje	j.van.der.naalt@umcg.nl
27	Van der Steen	Gregory	gregory@webstone.be
28	van Dijck	Jeroen T.J.M.	<a href="mailto:j.van.dijck@haaglandenmc.nl">j.van.dijck@haaglandenmc.nl</a>
29	van Essen	Thomas A.	T.A.van_Essen@lumc.nl
30	Van Hecke	Wim	wim.vanhecke@icomatrix.com
31	van Heugten	Caroline	Caroline.vanheugten@maastrichtuniversity.nl
32	Van Praag	Dominique	dominique.vanpraag@uza.be
33	van Wijk	Roel	roel-van-wijk@ziggo.nl
34	Vande Vyvere	Thijs	thijs.vandevyvere@icomatrix.com
35	Vargiolu	Alessia	neurorianimazione@hsgerardo.org
36	Vega	Emmanuel	emmanuel.vega@chru-lille.fr
37	Velt	Kimberley	k.velt@erasmusmc.nl
38	Verheyden	Jan	jan.verheyden@icomatrix.com
39	Vespa	Paul M.	PVespa@mednet.ucla.edu
40	Vik	Anne	anne.vik@ntnu.no
41	Vilcinis	Rimantas	rimantas.vilcinis@kaunoklinikos.lt
42	Volovici	Victor	v.volovici@erasmusmc.nl
43	von Steinbüchel	Nicole	nvsteinbuechel@med.uni-goettingen.de
44	Voormolen	Daphne	d.voormolen@erasmusmc.nl
45	Vulekovic	Petar	pvulekovic@gmail.com
46	Wang	Kevin K.W.	kawangwang17@gmail.com
47	Wiegers	Eveline	e.wiegers@erasmusmc.nl
48	Williams	Guy	gbw1000@wbic.cam.ac.uk
49			
50			
51			
52			
53			
54			
55			
56			
57			
58			
59			
60			

1			
2			
3	Wilson	Lindsay	l.wilson@stir.ac.uk
4	Winzeck	Stefan	sw742@cam.ac.uk
5	Wolf	Stefan	stefan.wolf@charite.de
6	Yang	Zhihui	zhihuiyang@ufl.edu
7	Ylén	Peter	peter.ylen@vtt.fi
8	Younsi	Alexander	alexander.younsi@med.uni-heidelberg.de
9	Zeiler	Frederick A.	umzeiler@myumanitoba.ca
10	Ziverte	Agate	agate.ziverte@inbox.lv
11	Zoerle	Tommaso	tommaso.zoerle@policlinico.mi.it
12			
13			
14			
15			
16			
17			
18			
19			
20			
21			
22			
23			
24			
25			
26			
27			
28			
29			
30			
31			
32			
33			
34			
35			
36			
37			
38			
39			
40			
41			
42			
43			
44			
45			
46			
47			
48			
49			
50			
51			
52			
53			
54			
55			
56			
57			
58			
59			
60			

Confidential: For Review Only