



Botanical gardens as key resources and hazards for biosecurity

Mesfin Wondafrash^{1,2} · Michael J. Wingfield³ · John R. U. Wilson^{2,4} · Brett P. Hurley¹ · Bernard Slippers³ · Trudy Paap³

Received: 13 February 2021 / Revised: 29 March 2021 / Accepted: 5 April 2021
© The Author(s), under exclusive licence to Springer Nature B.V. 2021

Abstract

Biodiversity and economic losses resulting from invasive plant pests and pathogens are increasing globally. For these impacts and threats to be managed effectively, appropriate methods of surveillance, detection and identification are required. Botanical gardens provide a unique opportunity for biosecurity as they accommodate diverse collections of exotic and native plant species. These gardens are also often located close to high-risk sites of accidental invasions such as ports and urban areas. This, coupled with routine activities such as the movement of plants and plant material, and visits by millions of people each year, place botanical gardens at risk to the arrival and establishment of pests and pathogens. Consequently, botanical gardens can pose substantial biosecurity risks to the environment, by acting as bridgeheads for pest and pathogen invasions. Here we review the role of botanical gardens in biosecurity on a global scale. The role of botanical gardens has changed over time. Initially, they were established as physic gardens (gardens with medicinal plants), and their links with academic institutions led to their crucial role in the accumulation and dissemination of botanical knowledge. During the second half of the 20th century, botanical gardens developed a strong focus on plant conservation, and in recent years there has been a growing acknowledgement of their value in biosecurity research as sentinel sites to identify pest and pathogen risks (novel pest-host associations); for early detection and eradication of pests and pathogens; and for host range studies. We identify eight specific biosecurity hazards associated with botanical gardens and note potential management interventions and the opportunities these provide for improving biosecurity. We highlight the value of botanical gardens for biosecurity and plant health research in general, and the need for strategic thinking, resources, and capacity development to make them models for best practices in plant health.

Keywords Invasive pests · Invasive pathogens · Early detection · Early warning · Sentinel plants · Invasive alien species

Communicated by David Hawksworth.

This article belongs to the Topical Collection: Ex situ conservation.

✉ Mesfin Wondafrash
mesfin.gossa@fabi.up.ac.za

Extended author information available on the last page of the article

Introduction

Invasive alien pests and pathogens pose major threats to plant health globally, resulting in significant ecological and economic damage (Holmes et al. 2009; Aukema et al. 2011; Boyd et al. 2013; Lovett et al. 2016). Huge economic losses in the agricultural and forestry sector have been attributed to the damage caused by invasive alien pests, for example up to \$39 billion per year in the United States alone (Pimentel et al. 2005; Holmes et al. 2009; Pratt et al. 2017). Over the last few decades, there has been a rapid surge in the number of alien arthropod pests, microbial pathogens and parasitic nematodes (hereafter referred to as pests) of plants in many countries (Aukema et al. 2010; Liebhold et al. 2012; Santini et al. 2013; Wingfield et al. 2015; Hurley et al. 2016). This recent surge is commonly attributed to the increasing global network of trade, travel and, in particular, the movement of live plants (Hulme 2009; Roques et al. 2009; Santini et al. 2013).

Regulatory measures that are taken against alien pests include the listing of quarantine pests [as supported by Pest Risk Analyses (PRAs)] and plant passports/quarantine inspections i.e. visual inspections/lab analyses (Klapwijk et al. 2016). However, such interventions are severely hampered by a lack of knowledge on potential pests and novel pest-host associations (Eschen et al. 2019). Co-evolution between plant pests and their hosts, as well as pressures from competition, predation and parasitism, contribute to the complexity and stability of natural ecosystems. Outbreaks of native pests under natural environmental conditions, are therefore rare (Alpert 2006; Burdon and Thrall 2009). Consequently, many damaging invasive alien plant pests were unknown to science, or at least unknown to cause severe damage, prior to their arrival and establishment in a novel environment. Examples include Dutch elm disease, *Ophiostoma ulmi* sensu lato (Ophiostomatales: Ophiostomataceae) and chestnut blight, *Cryphonectria parasitica* (Diaporthales: Cryphonectriaceae) in Europe and America (Brasier 2000; Rigling and Prospero 2018); and the polyphagous shot hole borer (PSHB), *Euwallacea fornicatus* (Coleoptera: Curculionidae) and its fungal symbiont *Fusarium euwallaceae* (Hypocreales: Nectriaceae) in California, Israel and South Africa (Mendel et al. 2012; Eskalen et al. 2013; Paap et al. 2018). Once established in a novel environment, these and many other pests have caused devastating economic and environmental impacts.

Early detection of pest incursions after their arrival is essential for control and management interventions to achieve eradication or containment, and for impacts and risks to be managed cost-effectively (Mehta et al. 2007; Liebhold and Kean 2019). In Australia, it has been estimated that their biosecurity system (by detecting and responding to introductions of pests and thereby reducing future impacts) will result in a net national benefit of AUS\$314 billion (an equivalent of US\$210 billion) over the next 50 years (Dodd et al. 2020). However, the detection of pests at the early stages of an incursion is challenging (Klapwijk et al. 2016; Thakur et al. 2019; Paap et al. 2020). Detection is often impossible before signs and symptoms develop on the host plants, and visible impacts are observed in the recipient environment (Thakur et al. 2019). By this point in time, the pest may already be well established. In addition, the diagnostic processes required to identify the organism(s) responsible for the damage further add to the long lag time between arrival, detection and identification of alien species. This is further exacerbated by the limited resources available to monitor large areas, particularly in countries with minimal budgets for plant health surveillance.

One strategy to improve the detection rate of incursions and identification of pest risks is to monitor sentinel plants i.e., plants already present in the vicinity of high-risk sites or

in urban areas and used for regular inspection of pests (Britton et al. 2010; Eschen et al. 2019; Mansfield et al. 2019). A similar approach is to use sentinel plantings—plantings in the country of origin of the pests used to identify organisms in their native range that are likely to be harmful if introduced and establish elsewhere (Roques et al. 2015; Eschen et al. 2019). Sentinel plantings could be existing collections of plants in botanical gardens or plants planted for this purpose. Sentinel plants and sentinel plantings help to detect and identify pest risks effectively and are useful for early warning (Wylie et al. 2008; Sweeney et al. 2012; Paap et al., 2017; Morales-Rodríguez et al. 2019). An example of the potential role of sentinel plantings is the emerald ash borer, *Agilus planipennis* (Coleoptera: Buprestidae), one of the devastating invasive insect pests in the United States (Herms and McCullough 2014). The emerald ash borer was detected on North American ash species planted in the beetle's native range in China long before its introduction in North America (Liu et al. 2003).

Botanical gardens provide a unique resource of sentinel plants and sentinel plantings to identify recently introduced alien pests and novel pest-host associations. There are thousands of botanical gardens widely distributed across the world with diverse collections of native and exotic plants (Miller et al. 2015; Barham et al. 2016). The high diversity of plant species in botanical gardens and their proximity to high-risk sites of introduction makes them suitable sites for post-border surveillance and monitoring of alien pests. Staff in these gardens work amongst the plant collections on a daily basis. Therefore, raising awareness and building the capacity of garden staff in surveillance and detection of pests can provide increased opportunities for detection of pests. This is especially useful for countries with limited plant health surveillance and biosecurity budgets. On the other hand, botanical gardens pose various biosecurity hazards through their regular activities such as collection, cultivation, sharing and the sale of plant materials; the use of machinery, vehicles and equipment; outsourcing of organic material; and tourism. These activities could lead to the introduction, establishment and spread of invasive species (Coetzee et al. 2001; Dawson et al. 2008; Scalera et al. 2012; Harrower et al. 2018; Rigling and Prospero 2018).

In this review we consider the changing role of botanical gardens, identify the associated hazards and highlight the recent recognition of their value for biosecurity. We discuss their potential role as bridgeheads and conduits of pest invasions, but also their recent role as key resources for identifying pest risks and new pest-host associations, as well as their potential to aid in pest detection and eradication.

Changing attitudes to biosecurity in botanical gardens as their functions and roles evolve

The functions and the roles of botanical gardens have evolved significantly over time (Powledge 2011; Krishnan and Novy 2016). Initially (mid-sixteenth century), botanical gardens were established to study medicinal plants (Oldfield 2007). However, during the European expansion and exploration of the rest of the world (seventeenth to nineteenth century), botanical gardens such as the Royal Botanic Gardens, Kew in Britain and the Hortus Botanicus Leiden in the Netherlands were engaged in economic botany and cultivation of attractive plants (Brockway 1979). This led to the movement of plants at unprecedented levels and enhanced the collection and documentation of exotic and previously unknown plant specimens (Brockway 1979; Borsch and Lohne 2014; Krishnan and Novy 2016). During the second half of the twentieth century, botanical gardens around the world

developed a strong focus on the conservation of rare and threatened plant species and the sustainable use of biodiversity to mitigate the impacts of climate change, habitat loss, and many other factors (Maunder et al. 2000; Borsch and Lohne 2014). In the twenty-first century, the focus of botanical gardens has moved towards maintaining relevance to communities and meeting their needs, locally and globally. Along with their role in ex situ conservation of biodiversity, botanical gardens have increased their focus on impactful education, research, and ecological restoration (Krishnan and Novy 2016; Smith 2019). Furthermore, the expertise of botanical gardens is being used in in situ conservation of plant biodiversity and ecological restoration (Chen et al. 2009; Krishnan and Novy 2016; Chen and Sun 2018).

Despite the general evolving roles and functions of botanical gardens overtime, their role in relation to biosecurity issues has remained largely unappreciated until recently. During the early days of the establishment of botanical gardens (mid-sixteenth century), the concept of native and alien pests was not recognised. Yet, there was a significant level of economic exchange between Europe and the rest of the world, and the resultant exchange of goods could have enhanced the movement of pests (Walter 2012). Likewise, during the seventeenth to nineteenth century, biosecurity was not acknowledged nor considered important, despite the unprecedented level of movement of plants across the world for various purposes (Brockway 1979). It was only in 1878 that the first attempt to regulate the international movement of plants was made in response to the significant damage caused by the insect Grape Phylloxera, *Phylloxera vastatrix* (Hemiptera: Phylloxeridae) (now *Daktulosphaira vitifolia*), to the viticulture industry (Santini et al. 2018).

A growing awareness of the importance of biosecurity was observed among botanical gardens as their focus shifted from economic botany and they began to acknowledge their role in science and plant conservation. However, much of this focus was on plant invasions (Lowe et al. 2000; Heywood 2011; Hulme 2011; Sharrock et al. 2011), and little attention was given to the movement of harmful plant pests. In the United States, voluntary codes of conduct and ethics such as the ‘Chapel Hill Challenge’ and the ‘St Louis Declaration’ were launched for botanical gardens, arboreta and the horticultural industry in 1999 and 2002, respectively (Hulme 2011). These codes of conduct set recommendations for the removal of invasive plants from living collections and plant sales, and for the control of invasive plants in botanical gardens. They also set recommendations for undertaking risk assessments on new collections. As a result, botanical gardens had begun to take action to combat the risk of plant invasions associated with their conservation efforts. For example, the European Botanical Gardens Consortium established an initiative to identify potential invasive taxa from botanical collections in various gardens. The consortium also alerted collectors of a potential invasion threat posed by their plant collections. Botanical gardens in the United States have instituted similar initiatives for the identification of invasive horticultural species (Heywood 2011; Sharrock et al. 2011).

In the last two decades, scientists have recognised the role that botanical gardens can play in plant health and biosecurity and tried to align this role with the conservation role of gardens to minimise the introduction and spread of pests (Barham et al. 2016; Eschen et al. 2019). To this end, the International Plant Sentinel Network (IPSN) (<https://www.plantsentinel.org/>) was launched in 2013 with the aim of providing support and resources needed by botanical gardens and arboreta to conduct research on plant health (Barham et al. 2016). The IPSN is an initiative developed by Botanic Gardens Conservation International (BGCI), to facilitate collaboration among botanical gardens and arboreta, National Plant Protection Organizations (NPPOs) and plant health scientists. In this regard, botanical gardens such as the Royal Botanic Garden Edinburgh in the United Kingdom (UK) have taken

important steps to reduce opportunities for pests to develop in their ex situ conservation collections. They have made substantial efforts to develop protocols for quarantine and horticultural practices, and expanded visitor engagement and public education on plant pests. They have also implemented precautionary approaches for plant distribution in order to minimise the introduction of pests into the gardens and to halt their subsequent dispersal (Hayden 2020). The Royal Botanic Gardens Kew in the UK and the Botanic Gardens Biosecurity Network and Botanic Gardens Surveillance Network in Australia have formed important links with government plant health departments to enhance biosecurity through inspection and structured plant pest surveillance (<https://extensionaus.com.au/botanicgardensbiosecurity/structured-plant-pest-surveillance-by-botanic-gardens-staff/>, <https://www.kew.org/read-and-watch/behind-the-scenes-plant-quarantine-unit>).

Recently, botanical gardens have been used as sentinel sites for the detection of introduced alien pests (Paap et al. 2017; Eschen et al. 2019). Botanical gardens can also be viewed as *ex-patria* sentinel plantings or sentinel plantations, as proposed by Eschen et al. (2019). In this case, woody plants native to the importing country, which are growing in botanical gardens in the exporting country, can be used to monitor and identify pest risks. The identification of native pest–exotic host associations in the exporting country could provide valuable information of potential future invasion risks to the country of origin of the host plant (Britton et al. 2010; Barham et al. 2016; Paap et al. 2017; Eschen et al. 2019; Morales-Rodríguez et al. 2019).

The role of botanical gardens in biosecurity

The key elements and features that lend botanical gardens special importance in biosecurity, particularly in the study, detection and management of pests, are illustrated in Fig. 1 and the accompanying Table 1. Eight specific hazards were identified and are discussed, noting that the threats also provide opportunities for improved management and communication. In this section, we highlight the hazards and the opportunities botanical gardens present as bridgeheads and conduits for invasions; sentinel sites for the detection and eradication of pest incursions; important resources for research (focussing on the opportunities they provide to identify new pest-host association and thereby determine pest host range); and great opportunities for meaningful public engagement on biosecurity issues.

Botanical gardens as bridgeheads and conduits for invasions

Despite their clear role in biodiversity conservation, botanical gardens have historically acted as conduits for the introduction of invasive plants. For example, of the 34 plants listed by the IUCN as among 100 of the worst invasive species worldwide (Lowe et al. 2000), there is published evidence implicating botanical gardens as the most probable source of introduction for over half of these species (Hulme 2011).

There is no doubt that ex situ plant conservation plays a vital role in the conservation of plants, but it also poses high risk of pest introduction (Liebhold et al. 2012). Therefore, botanical gardens can also serve as pathways of introduction for invasive pests. It is possible for alien pests established in botanical gardens to spread further into the surrounding environment through various garden activities. Examples include invasion by chestnut blight, *C. parasitica* in the United States (Rigling and Prospero 2018) and the root rot fungus, *Armillaria mellea* (Agaricales: Physalacriaceae) in South Africa (Coetzee et al. 2001,




Fig. 1 Eight biosecurity hazards presented by botanical gardens and the opportunities they provide to improve management and communication. Details on the hazards, pathways of movement, real-world examples, management measures, and the corresponding opportunities are outlined in Table 1. The letters in the figures correspond to the letters in the table. The high diversity of native and exotic plant species in the gardens and their proximity to high risk sites such as ports and urban areas provides a unique opportunity for the detection and identification of pest risks. Materials going into the gardens such as seeds, tubers, cuttings, mulch, compost and soil could potentially transport and introduce pests to the gardens (a, d). On the other hand, materials leaving the gardens such as sold plants, prunings and dead plants can potentially transport pests established in the gardens to the external environment (b, h). Other activities of the gardens, including visits by local and international visitors (f), the use of machinery and equipment (e), and plant exchange between botanical gardens (c) may also serve as pathways of movement of pests to- and from the gardens. Additionally, pests may naturally disperse between managed estates of the gardens and the adjacent natural vegetation (g)

2003). In the United States, the first report of *C. parasitica* was on American chestnut in the Zoological Park of New York City in 1904 (Rigling and Prospero 2018). Subsequently, this pathogen spread to the surrounding environment and devastated native American chestnut trees. Similarly, *A. mellea* was initially detected in Company's Garden and Kirstenbosch National Botanical Garden in South Africa (Coetzee et al. 2001, 2003). It has recently been reported from the surrounding natural vegetation of Table Mountain National Park, a declared UNESCO world heritage site (Coetzee et al. 2018).

Plant donation from the Kirstenbosch National Botanical Garden in South Africa resulted in the accidental introduction of five sap-sucking hemipteran pests to London, the UK (Salisbury et al. 2011). Several pot-grown *Aloe* species together with a *Cheiridopsis glomerata* (Cryophyllales: Aizoaceae) were sent to London for the 2011 Royal Horticultural Society Chelsea Flower Show. At the end of the show, the plants were donated to the Royal Society of Chelsea. No pests were found on the plants despite assessments being made before and after the show. The donated plants were placed in the propagation department's quarantine facility. Within a month, it was clear that the plants were infested with several sap-sucking insects, which were later identified as five different insect species. Two

Table 1 Key biosecurity hazards and opportunities presented by botanical gardens, and potential management options

Hazard	Pathway(s)	Examples	Management and opportunities
a Plant materials are brought into the garden for cultivation and either the plants themselves or something associated with the plants or the soil material becomes invasive	<p>Escape: Botanical gardens & Zoos (publicGarden-Zoo/Aquaria)</p> <p>Contaminant: Contaminants of plants (contaminantOnPlants)</p> <p>Contaminant: Nursery material contaminant (contaminantNursery)</p> <p>Contaminant: Parasites of plants (parasiteOnPlant)</p> <p>Contaminant: Seed contaminant (seedContaminant)</p>	<p>In Cameroon, the trumpet tree (<i>Cecropia peltata</i>) was spread from a plantation in Limbe Botanical Garden into the forests of Mount Cameroon by fruit-eating bats and birds, where this tree outcompeted the native pioneer trees</p> <p>A genetically altered tropical alga (<i>Caulerpa taxifolia</i>) escaped from a public aquarium in Monaco into the Mediterranean Sea possibly with aquarium overflow (Scalera et al. 2012)</p> <p>At the Amani Botanical Garden (ABG), Tanzania, of 214 alien plant species found extant 38 had established and 16 had spread widely in the botanical garden (Dawson et al. 2008)</p> <p>The invasive root rot fungus, <i>Armillaria mellea</i> was introduced from Europe to the Company's Garden in Cape Town, South Africa, possibly on potted plants planted in the garden (Coetzee et al. 2001)</p> <p>In the United states, <i>Cryphonectria parasitica</i>, the causal agent of chestnut blight was first reported from the Zoological Park of New York City in 1904. The fungus was likely introduced via nursery stock imported from Asia (Rigling and Prospero 2018, Global Invasive Species Database)</p> <p>A container survey in New Zealand found eggs of brown marmorated stink bug (<i>Halyomorpha halys</i>) on leaves and foliage of imported plants (Harrower et al. 2018)</p> <p><i>Hymenocystus fraxineus</i>, the causal agent of Chalara ash dieback has been introduced to many countries through movement of diseased ash plants (Harrower et al. 2018)</p> <p>Two hundred cherry trees were sent in 1911 as a gift from the people of Japan to be planted in Washington DC. Upon their arrival found infested with crown gall disease and various insect pests (Liebhold and Griffin 2016)</p>	<p>Importers of alien species must comply with the phytosanitary regulations and demonstrate that the likelihood of escape is minimal, or the consequences of escape are not significant. It is also important to engage with various stakeholders including industries and the general public to create awareness about invasive species and the threats they pose</p> <p>Introduction of invasive species via contaminant pathway is closely linked to international trade. Thus, international regulations and phytosanitary measures/standards play very important role in reducing the problem. The exporting country should apply the sanitary and phytosanitary standards required while the importing country practices border controls and quarantine procedures. Harmonization of phytosanitary measures among countries is required, as is a stricter follow up on the enforcement of the enacted regulations and Acts</p> <p>The presence of invasive species that have escaped collections provide an opportunity to increase awareness of invasive plants in the region in a controlled setting</p> <p>Annual reports describing escape from plantings can be used to calculate the lag time between initial planting and earliest record of successful spread and thereby help to determine invasiveness of a species (e.g. Daehler 2009). This would enable an assessment of the risk of plant invasion from collections in botanical gardens</p> <p>Technological advancement provides immense opportunities to unravel cryptic species, divergent lineages and endophytic organisms which are difficult to detect and identify using classical approaches.</p> <p>Many pests are unknown prior to their arrival in a new environment. Identification of novel pest-host associations through sentinel plant research and early warning systems provide valuable information and aids the detection and identification process</p>

Table 1 (continued)

Hazard	Pathway(s)	Examples	Management and opportunities
b Plant material is propagated in the botanical garden (in a nursery) and sold to the general public, taken elsewhere, and that process leads to the spread of invasive plant species or organisms (contaminants) associated with the plants	Escape: Horticulture (horticulture) Escape: Ornamental (ornamental)NonHorticulture) Contaminant: Nursery material contaminant (contaminantNursery) Contaminant: Parasites of plants (parasitesOnPlants) Contaminant: Seed contaminant (seedContaminant)	Garden lupin (<i>Lupinus polyphyllos</i>) has been introduced to Europe, Australia and New Zealand for various purposes, including for ornamentation, soil stabilisation and cultivation Water hyacinth (<i>Pontederia crassipes</i>) has been introduced to different parts of the world as an ornamental plant <i>Phytophthora ramorum</i> , the causal agent of Ramorum disease in many plant species; appears to be transported with infested nursery plants and soil (Global Invasive Species Database)	The risk of spread of invasive species through plant sale can be minimised by focusing the sale on less risky plant species and following good phytosanitary practices within the nursery. Compliance with local regulations on plant movement is also equally important in reducing the risks Opportunities exist to create public awareness around invasive species on special occasions such as annual plant sale days. This can be done through oral presentations, posters and distribution of written information about invasive species, their pathways of movement, the impact and management measures
c Plant material is shared with other gardens or used for display purposes, and the process leads to the introduction and spread of invasive pest species	Contaminant: Nursery material (contaminantNursery) Contaminant: Parasite of plants (contaminantOnPlants) Contaminant: Seed contaminant (seedContaminant)	Plant donation from Kirstenbosch National Botanical Garden in South Africa to London for the 2011 Royal Horticultural Society Chelsea Flower Show resulted in the accidental introduction of five sap-sucking hemipteran pests to the UK. Fortunately, all the infested plants were destroyed before the spread of insects (Salisbury et al. 2011)	The donor should inspect plants for pests and issue a plant passport for the donated plants Upon arrival, the recipient should keep the plants in a quarantine facility and regularly inspect for pests and diseases. It is only after this process that the use of plants should be allowed
d Organic materials such as mulch, wood chips and soil can transport invasive species into the gardens	Contaminant: Habitat material (HabitatMaterial) Contaminant: Seed contaminant (seedContaminant)	Dispersal of the fungus <i>Phellinus noxius</i> , the causal agent of brown rot disease of trees is possible through transport of infested soil (Global Invasive Species Database) Common ragweed, <i>Ambrosia artemisiifolia</i> is common in construction sites suggesting its dispersal through transport of soil and gravel (Bullock et al. 2012) <i>Phytophthora cinnamomi</i> , an oomycete that causes root rot and dieback in various plant species, can be spread through soil and organic matter	Obtaining organic materials from trusted sources is very important. A trusted source should avoid making mulches and wood chips from infested plant parts. Fine chipping and composting is recommended to kill the contaminants. Allow mulch and wood chip to dry before use as fresh material would spread pests Soil solarisation is recommended to minimise the risk of introduction or spread of invasive species

Table 1 (continued)

Hazard	Pathway(s)	Examples	Management and opportunities
e Plant propagules and pests can be imported or exported or spread within the garden on vehicles, machinery or equipment	Stowaway: Machinery & equipment (machinery/Equipment)	<p>There are many examples of invasive plant species transported/dispersed by machinery, equipment and vehicles [CABI Invasive Species Compendium: Machinery and equipment (Pathway vector)]. E.g. Seeds of the alien invasive Common ragweed (<i>Ambrosia artemisiifolia</i> L.) have been transported in litter and soil by agricultural machinery from infested areas across Europe (Bullock et al. 2012)</p> <p>Plant pathogens such as <i>Phytophthora cinnamomi</i> and <i>Austrorhizium psidii</i>, the causal agent of myrtle rust spread by machinery, equipment and vehicles</p>	<p>Machinery, equipment and vehicles should be carefully washed/cleaned before entry or exit of the gardens</p> <p>It is equally important to limit the use of machinery and equipment to specific parts of the garden. If there is a need to use them in multiple sites, cleaning/washing is recommended between sites</p> <p>Limit the number of entry and exit points (gates) to reduce free movement of machinery, equipment and vehicles</p> <p>Display biosecurity signs at vehicle access points</p>
f Garden visitors accidentally either bring an invasive species into the garden or transport invasive species established in the garden to the external environment	Stowaway: People & luggage (people)	<p>65 fungal species were cultured from shoes of air travellers in a study conducted at Honolulu, the United States (Baker 1966)</p> <p>Spores of pathogenic fungi were detected from clothing of passengers in a study conducted at Wellington Airport in 1980 and 1982 (Sheridan 1989)</p> <p>The harlequin ladybird (<i>Harmonia axyridis</i>) has been reported arriving in new regions in suitcases</p> <p>Organic and inorganic debris and live insects were found on tents brought by air passengers at Auckland International Airport during December 1981 (Gadgil and Flint 1983)</p> <p>Plant pathogens such as <i>P. cinnamomi</i> and <i>A. psidii</i> can easily be transported via contaminated clothing, hats, footwear, equipment or vehicles</p>	<p>Creating public awareness about the threats and pathways of movement is important. The principle of “arrive clean, leave clean” should be encouraged and implemented. Demonstrate to the public and garden workers the value of ensuring all clothing, hats, footwear, tools, equipment, machinery and vehicles are free of weed seeds, mud, soil and organic matter before entering and exiting botanical gardens. This can be encouraged through signage, newsletters, social medias such as Twitter, Facebook and other media outlets, school teaching and citizen science</p> <p>The use of baths to clean shoes and air current to remove contaminants from hair, clothing and luggage of visitors would minimise the introduction and spread of invasive species</p>
g Invasive species naturally spread from the gardens to neighbouring areas or from neighbouring areas to the gardens	Natural dispersal (natural/Dispersal)	<p>The root rot fungus, <i>Armillaria mellea</i> which was initially reported from the Company’s Garden and subsequently from Kirstenbosch National Botanical Garden in South Africa, has gradually moved to the neighbouring Table Mountain National Park</p> <p><i>Cryphonectria parasitica</i>, the causal agent of chestnut blight, was initially reported on American chestnut in the Zoological Park of New York City and has subsequently spread to the surrounding environment and devastated native American chestnut trees</p>	<p>Botanical gardens are important sentinel plantings and sentinel sites. It is therefore important to conduct regular surveillance and monitoring to detect and identify invasive species spreading from the neighbouring environment to the botanical gardens</p> <p>Similarly, regular surveillance is recommended in the natural vegetation neighbouring botanical gardens to monitor the spread of invasive species established in botanical gardens and take the necessary management measures</p>

Table 1 (continued)

Hazard	Pathway(s)	Examples	Management and opportunities
h Invasive species can spread from botanical gardens to the surrounding environment via disposal/dumping of plant waste	Contaminant: Transportation of habitat material (transportationHabitatMaterial)	In Java, the invasive aquatic plant water hyacinth (<i>Pontederia crassipes</i>) became widely spread as a result of dumping of excessive weed from Bogor Botanical Garden into the Ciliwung River during the early twentieth century (Hulme 2011)	Plant waste disposal is one of the regular activities of botanical gardens. However, care should be taken not to dispose severely infected plants or plant parts in regular dumping sites as these will aid the spread of invasive species. Infested plants or plant parts should preferably be burnt in the gardens rather than disposing them out side of the gardens Disposal/dumping sites should be far from natural vegetation and high human traffic areas to minimise the risk of dispersal of invasive species

The hazards are linked to the relevant invasion pathway as they would operate for botanical gardens using terminology adopted by the Convention on Biological Diversity with proposed revisions by Harrower et al. (2018) and as recently included in the Darwin Core (dwc; pathway, see <https://dwc.tdwg.org/pw/#dwcpw> accessed 29 March 2021; Groom et al. 2019; the Darwin Core terms are given in brackets). Examples of how the hazards have led to invasions or potential invasions are given either specifically for botanical gardens, or for systems that would be expected to be analogous to botanical gardens, noting that in the latter case a different pathway might be involved from that specified here

of these species, the small mirid bug, *Aloea australis* (Hemiptera: Miridae) and the ice-plant scale, *Pulvinaria delottoi* (Hemiptera: Coccidae), were the first detections of these pests in Europe. Dried specimens of these pests were deposited in different insect reference collections and all the infested plants were destroyed. These examples demonstrate the need for quarantine and repeated inspections as during initial inspections, pests may have been at their immature stages and less visible.

Botanical gardens as sentinel sites for the detection and eradication of pest incursions

A recent move towards sentinel plant research has facilitated the detection and identification of emerging pest risks. Initiatives such as the IPSN, COST Action Global Warning (<https://www.cost.eu/actions/FP1401/#tabs{Name:overview}>), and the European Union Horizon 2020 HOMED (Holistic Management of Emerging Forest Pests and Diseases) project (<http://homed-project.eu/>), are coordinating sentinel plant research globally. These projects serve to highlight the many first reports of pests from botanical gardens and arboreta (Jock et al. 2000; Salisbury et al. 2011; Paap et al. 2018; Hulbert et al. 2019; Tchotet Tchoumi et al. 2019). In South Africa alone, 67 pest species (including fungi, oomycetes, insects and mites) were detected and identified from various botanical gardens over the past 23 years (1996–2019), 20 of which were first reports for the country (Wondafrash et al. in prep). In a recent review article, Mansfield et al. (2019) reported several cases of novel pest-host associations identified from sentinel plants, including plants grown in botanical gardens. These novel associations involved insects, fungi, bacteria and nematodes. Timely detection and identification of such pests aids eradication and containment of new incursions, thereby reducing the risks to natural vegetation and commercial and agricultural systems (Jock et al. 2000; Kenis et al. 2019).

Globally, thousands of eradication programmes have been implemented for alien forest insects since 1970. These include many cases of successes and failures (Liebhold and Kean 2019). Historical examples of eradication include many successful localised gypsy moth, *Lymantria dispar* (Lepidoptera: Erebididae) eradication programs in the United States and the Asian long-horn beetle, *Anoplophora glabripennis* from Chicago, the United States; Toronto, Canada; and Braunau, Austria (Liebhold and Kean 2019). Recently (April 2020), *E. fornicatus* was detected on plants growing in a greenhouse in the botanical gardens of Trauttmansdorff Castle in Italy and eradication is underway (<https://gd.eppo.int/reporting/article-6772>).

Eradication of plant pathogens once established in natural ecosystems presents a huge challenge (Paap et al. 2020). Yet, there are numerous examples of successful eradication of alien plant pathogens from controlled environments (Pluess et al. 2012). For example, inspection of known host plants of fire blight caused by the bacterium *Erwinia amylovora* (Enterobacterales: Erwiniaceae) in botanical gardens of Melbourne and Adelaide in Australia, resulted in its early detection in Royal Botanic Gardens, Melbourne (Jock et al. 2000). This is a pathogen endemic to North America and a causal agent of a serious disease of apple and pear trees and other rosaceous plants. Wide-ranging surveys and an intense host eradication program resulted in the removal of hundreds of trees, and protected Australia's pome fruit industry (Rodoni et al. 2002), valued at US\$560 million for the year 2014–15 (<https://apal.org.au/>). This demonstrates that the early recognition of new disease symptoms, coupled with rapid and accurate diagnostics, can lead to the successful

eradication of damaging organisms. Early detection and identification of pests is more easily achieved in botanical gardens than other sites.

Botanical gardens provide opportunities to determine pest host range

The diverse collection of exotic and native plant species in botanical gardens are valuable resources to investigate and determine host ranges of pests (Groenteman et al. 2015; Scott-Brown et al. 2018). Host range studies are not only valuable to the invaded region. Where exotic hosts are present, these studies can also inform other countries regarding possible future threats to their plant health, by contributing information on pests that are of regulatory interest (based on observed susceptibility in invaded ranges). These observations can be useful for early warning as they inform PRAs and assist with categorisation (quarantine status) of pests. Three examples of host range studies conducted in botanical gardens are presented below.

Example 1. *Euwallacea fornicatus*—*F. euwallaceae* complex

The host range of *E. fornicatus* and its fungal symbiont, *F. euwallaceae*, was studied at the Los Angeles Arboretum and the Huntington Botanical Gardens in its invasive range in California (Eskalen et al. 2013). Of the 335 tree species present in the gardens, 207 species (62%), from 58 plant families, showed signs and symptoms of *E. fornicatus* infestation. *Fusarium euwallaceae* was isolated from 54% of the *E. fornicatus* infested plant species. Trees infested by *E. fornicatus* and its fungal symbiont included native and agriculturally important species, and common street trees. This study showed the potential of the beetle and its fungal symbiont to establish in diverse plant communities in the United States and beyond. The techniques used in this study and the results have aided the study of the beetle and its fungal symbiont in its recent introduced range in South Africa.

Example 2. *Xylella fastidiosa* (Xanthomonadales: Xanthomonadaceae)

This is a xylem-dwelling bacterium known to cause disease in a variety of plant species. The symptoms range from leaf scorch, chlorosis or browning to stunted growth, branch die-back and death of infected plants (CABI 2020). It is vectored by the glassy-winged sharpshooter (GWSS), *Homalodisca vitripennis* (Hemiptera: Cicadellidae). The combination of the vector and the bacterium causes severe damage in plants of agricultural, ornamental and biodiversity importance (Pilkington et al. 2005). The host range of *X. fastidiosa* and its vector GWSS and the biocontrol potential of egg parasitoids against GWSS was studied on New Zealand plants in four botanical gardens and arboreta and public spaces in southern California (Groenteman et al. 2015). Signs of GWSS activity were observed on 26 of the 102 plants (25%) examined, while the bacterium was recovered at all the locations sampled and in 51% of the samples. This showed that several of New Zealand's indigenous plant species are susceptible to the bacterium.

Example 3. Red palm mite, *Raoiella indica* (Trombidiformes: Tenuipalpidae)

Since its detection in Martinique and St. Lucia in 2004, this polyphagous mite species has spread rapidly through the Neotropical region and has caused significant damage to a diverse range of plant species (Carrillo et al. 2012). The host range of *R. indica* was

investigated through periodic surveys in the Fairchild Tropical Botanic Garden in Florida, the United States and the Royal Botanic Gardens of Trinidad and Tobago. This has helped to produce an updated list of 91 reproductive host species of *R. indica*. This study also confirmed 27 new reproductive hosts, representing a 30% increase of the previously recorded host range (Carrillo et al. 2012).

Public engagement in botanical gardens

Opportunities exist for public engagement in plant protection via the gardens, leading to citizen science and better understanding of threats posed by pests, their pathways of movement and biosecurity measures. Botanical gardens are visited by many people on a daily basis. Biosecurity teams of the botanical gardens can use this opportunity to create awareness regarding pest and pathogen risks, pathways of movement and the recommended measures as per operational biosecurity guidelines. This could be achieved in various ways, including presentations, practical demonstrations of pests and pathogens in the gardens, group discussions and by posting biosecurity signage (see Hayden 2020).

Conclusions

Botanical gardens pose a variety of biosecurity threats—here we characterised eight specific threats (Fig. 1, Table 1). However, botanical gardens also provide great opportunities to study invasive pests, invasive plants and native pests that damage exotic plants. Botanical gardens are important sentinel sites for the detection and discovery of pest species, and are valuable resources for host range studies and the identification of novel pest-host associations. We believe some botanical gardens are already champions of biosecurity, and argue that others can expand their role from sentinel sites for biosecurity research to become models for best practices in plant health, in the global effort to limit the spread and impact of pests. This will require strategic thinking, resources, and capacity development. Yet, we strongly believe that such an approach would increase the value of the botanical gardens and the contribution that they make to biosecurity. Ex situ conservation is central to the role of many botanical gardens because of the increasing impacts of climate change, habitat loss, fragmentation, degradation, pollution and over-exploitation on plant biodiversity. This conservation role of botanical gardens needs to be aligned with biosecurity efforts in order to minimize the introduction of invasive pests to the gardens, and thus to halt their subsequent spread to the surrounding environment.

Acknowledgements The South African Department of Forestry, Fisheries and the Environment (DFFE) are thanked for funding noting that this publication does not necessarily represent the views or opinions of DFFE or its employees. The Forestry and Agricultural Biotechnology Institute (FABI) of the University of Pretoria is acknowledged for its infrastructural support. The horticulturalists at the various botanical gardens in South Africa deserve special thanks for their unreserved support on the project.

Authors contribution Conceptualization [TP, JR UW, MW], Formal analysis and investigation [TP, MW], Writing—original draft preparation [MW], Commenting the various versions of the manuscript and approval of the final version [MJW, JR UW, BPH, BS, TP].

Funding This work was supported by the South African Department of Forestry, Fisheries and the Environment (DFFE).

Availability of data and material Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.

Declarations

Conflict of interest The authors have no conflict of interest to declare that are relevant of the content of this article.

Consent for publication All the authors have given their consent for the submission and publication of this article.

References

- Alpert P (2006) The advantages and disadvantages of being introduced. *Biol Invasions* 8:1523–1534. <https://doi.org/10.1007/s10530-005-5844-z>
- Aukema JE, McCullough DG, Von Holle B, Liebhold AM, Britton K, Frankel SJ (2010) Historical accumulation of nonindigenous forest pests in the continental United States. *Bioscience* 60:886–897. <https://doi.org/10.1525/bio.2010.60.11.5>
- Aukema JE, Leung B, Kovacs K, Chivers C, Britton KO, Englin J, Frankel SJ, Haight RG, Holmes TP, Liebhold AM (2011) Economic impacts of non-native forest insects in the continental United States. *PLoS ONE* 6:e24587. <https://doi.org/10.1371/journal.pone.0024587>
- Baker EG (1966) Inadvertent distribution of fungi. *Can J Microbiol* 12(109):112. <https://doi.org/10.1139/m66-016>
- Barham E, Sharrock S, Lane C, Baker R (2016) The International Plant Sentinel Network: a tool for Regional and National Plant Protection Organizations. *Bulletin OEPP/EPPO Bulletin* 46:156–162
- Borsch T, Löhne C (2014) Botanic gardens for the future. Integrating research, conservation, environmental education and public recreation. *Ethiop J Biol Sci* 13:115–133
- Boyd IL, Freer-Smith PH, Gilligan CA, Godfray HCJ (2013) The consequence of tree pests and diseases for ecosystem services. *Science* 342:1235773
- Brasier CM (2000) Intercontinental spread and continuing evolution of the Dutch elm disease pathogens. In: Dunne CP (ed) *The elms: breeding, conservation and disease management*. Kluwer Academic Publishers, Boston, pp 61–72
- Britton K, White P, Kramer A, Hudler G (2010) A new approach to stopping the spread of invasive insects and pathogens: early detection and rapid response via a global network of sentinel plantings. *N Z J Sci* 40:109–114
- Brockway LH (1979) *Science and colonial expansion: the role of the British Royal Botanic Gardens*. Academic Press, New York
- Bullock J, Beale S, Chapman D, Schafer S, Roy D, Girardello M, Haynes T, Beal S, Wheeler B, Dickie I, Phang Z, Tinch R, Čivić K, Delbaere B, Jones-Walters L, Hilbert A, Schrauwen A, Prank M, Sofiev M, Niemelä S, Räisänen P, Lees B, Skinner M, Finch S, Brough C (2012) Assessing and controlling the spread and the effects of common ragweed in Europe (ENV.B.2/ETU/2010/0037). European Commission, Final report
- Burdon JJ, Thrall PH (2009) Coevolution of plants and their pathogens in natural habitats. *Science* 324:755–756. <https://doi.org/10.1126/science.1171663>
- CABI (2020) *Xylella fastidiosa* (Pierce's disease of grapevines). *Invasive Species Compendium*. <https://www.cabi.org/isc/datasheet/57195>. Accessed 20 Aug 2020
- Carrillo D, Amalin D, Hosein F, Roda A, Peña JE (2012) Host plant range of *Raoiella indica* (Acari: Tenuipalpidae) in areas of invasion of the New World. *Exp Appl Acarol* 57:271–289. <https://doi.org/10.1007/s10493-011-9487-8>
- Carter PCS (1989) Risk assessment and pest detection surveys for exotic pests and diseases which threaten commercial forestry in New Zealand. *N Z J For Sci* 19:353–374
- Chen G, Sun W (2018) The role of botanic gardens in scientific research, conservation, and citizen science. *Plant Divers* 40:181–188. <https://doi.org/10.1016/j.pld.2018.07.006>
- Chen J, Cannon CH, Hu H (2009) Tropical botanical gardens: at the in situ ecosystem management frontier. *Trends Plant Sci* 14:584–589. <https://doi.org/10.1016/j.tplants.2009.08.010>

- Coetzee MPA, Wingfield BD, Harrington TC, Steimel J, Coutinho TA, Wingfield MJ (2001) The root rot fungus *Armillaria mellea* introduced into South Africa by early Dutch settlers. *Mol Ecol* 10:387–396. <https://doi.org/10.1046/j.1365-294X.2001.01187.x>
- Coetzee MPA, Wingfield BD, Roux J, Crous JPW, Denman S, Wingfield MJ (2003) Discovery of two northern hemisphere *Armillaria* species on Proteaceae in South Africa. *Plant Pathol* 52:604–612. <https://doi.org/10.1046/j.1365-3059.2003.00879.x>
- Coetzee MPA, Musasira NY, Roux J, Roets F, van der Merwe NA, Wingfield MJ (2018) *Armillaria* root rot spreading into a natural woody ecosystem in South Africa. *Plant Pathol* 67:883–891. <https://doi.org/10.1111/ppa.12804>
- Daehler CC (2009) Short lag times for invasive tropical plants: evidence from experimental plantings in Hawai'i. *PLoS ONE* 4:e4462. <https://doi.org/10.1371/journal.pone.0004462>
- Dawson W, Mndolwa AS, Burslem D, Hulme PE (2008) Assessing the risks of plant invasions arising from collections in tropical botanical gardens. *Biodivers Conserv* 17:1979–1995. <https://doi.org/10.1007/s10531-008-9345-0>
- Dodd A, Stoeckl N, Baumgartner J, Kompas T (2020) Key result summary: Valuing Australia's biosecurity system. Centre of Excellence for Biosecurity Risk Analysis (CEBRA) Report
- Eschen R, Hanlon RO, Santini A, Vannini A, Roques A, Kirichenko N, Kenis M (2019) Safeguarding global plant health: the rise of sentinels. *J Pest Sci* 92:29–36. <https://doi.org/10.1007/s10340-018-1041-6>
- Eskalen A, Stouthamer R, Lynch SC, Twizeyimana M, Gonzalez A, Thibault T (2013) Host range of *Fusarium* dieback and its ambrosia beetle (Coleoptera: Scolytinae) vector in southern California. *Plant Dis* 97: 938–951
- Gadgil PD, Flint TN (1983) Assessment of the risk of introduction of exotic forest insects and diseases with imported tents. *N Z J For* 28:58–67
- Global Invasive Species Database. <http://www.iucngisd.org/gisd/>. Accessed 15 Nov 2020
- Groenteman R, Forgie SA, Hoddle MS, Ward DF, Goeke DF, Anand N (2015) Assessing invasion threats: novel insect-pathogen-natural enemy associations with native New Zealand plants in southern California. *Biol Invas* 17:1299–1305. <https://doi.org/10.1007/s10530-014-0804-0>
- Groom Q, Desmet P, Reyserhove L, Adriaens T, Oldoni D, Vanderhoeven S, Baskauf SJ, Chapman A, McGeoch M, Walls R, Wiczorek J, Wilson JR, Zermoglio PFF, Simpson A (2019) Improving Darwin Core for research and management of alien species. *Biodivers Inf Sci Stand* 3:e38084. <https://doi.org/10.3897/biss.3.38084>
- Harrower CA, Scalera R, Pagad S, Schönrogge K, Roy HE (2018) Guidance for interpretation of CBD categories on introduction pathways. Technical note prepared by IUCN for the European Commission
- Hayden K (2020) Botanic gardens and plant pathogens: a risk-based approach at the Royal Botanic Garden Edinburgh. *Sibbaldia. Int J Bot Garden Hortic* 18:127–139. <https://doi.org/10.24823/Sibbaldia.2020.293>
- Herms DA, McCullough DG (2014) Emerald ash borer invasion of North America: history, biology, ecology, impacts, and management. *Annu Rev Entomol* 59:13–30. <https://doi.org/10.1146/annurev-ento-011613-162051>
- Heywood VH (2011) The role of botanic gardens as resource and introduction centres in the face of global change. *Biodivers Conserv* 20:221–239. <https://doi.org/10.1007/s10531-010-9781-5>
- Holmes T, Aukema J, Von Holl B, Liebhold A, Sills E (2009) Economic impacts of invasive species in forests. *Ann NY Acad Sci* 1162:18–38
- Hulbert JM, Paap T, Burgess TI, Roets F, Wingfield MJ (2019) Botanical gardens provide valuable baseline *Phytophthora* diversity data. *Urban For and Urban Gree* 46:126461. <https://doi.org/10.1016/j.ufug.2019.126461>
- Hulme PE (2009) Trade, transport and trouble: managing invasive species pathways in an era of globalization. *J Appl Ecol* 46:10–18. <https://doi.org/10.1111/j.1365-2664.2008.01600.x>
- Hulme PE (2011) Addressing the threat to biodiversity from botanic gardens. *Trends Ecol Evol* 26:168–174. <https://doi.org/10.1016/j.tree.2011.01.005>
- Hurley BP, Garnas J, Wingfield MJ, Branco M, Richardson DM, Slippers B (2016) Increasing numbers and intercontinental spread of invasive insects on eucalypts. *Biol Invas* 18:921–933. <https://doi.org/10.1007/s10530-016-1081-x>
- Jock S, Rodoni B, Gillings M, Kim WS, Copes C, Merriman P, Geider K (2000) Screening of ornamental plants from the Botanic Gardens of Melbourne and Adelaide for the occurrence of *Erwinia amylovora*. *Australas Plant Path* 29:120–128. <https://doi.org/10.1071/AP00020>
- Kenis M, Hurley BP, Colombari F, Lawson S, Sun J, Wilcken C, Weeks R, Sathyapala S (2019) Guide to the classical biological control of insect pests in planted and natural forests. *FAO Forestry Paper* No. 182, Rome

- Klapwijk MJ, Hopkins AJM, Eriksson L, Pettersson M, Schroeder M, Lindelow A, Ronnberg J, Keskitalo ECH, Kenis M (2016) Reducing the risk of invasive forest pests and pathogens: Combining legislation, targeted management and public awareness. *Ambio* 45:223–234. <https://doi.org/10.1579/0044-7447-38.3.135>
- Krishnan S, Novy A (2016) The role of botanic gardens in the twenty-first century. *CAB Reviews* 11, No. 023
- Liebholt AM, Griffin RL (2016) The legacy of Charles Marlatt and efforts to limit plant pest invasions. *Am Entomol* 62:218–227. <https://doi.org/10.1093/ae/tmw072>
- Liebholt AM, Kean JM (2019) Eradication and containment of non-native forest insects: successes and failures. *J Pest Sci* 92:83–91. <https://doi.org/10.1007/s10340-018-1056-z>
- Liebholt A, Brockerhoff E, Garrett L, Parke J, Britton K (2012) Live plant imports: the major pathway for the forest insect and pathogen invasions of the US. *Front Ecol Environ* 10:135–143. <https://doi.org/10.1890/110198>
- Liu HP, Bauer LS, Gao R, Zhao T, Petrice TR, Haack RA (2003) Exploratory survey for the emerald ash borer, *Agilus planipennis* (Coleoptera: Buprestidae), and its natural enemies in China. *Great Lakes Entomol* 36:191–204
- Lovett GM, Weiss M, Liebholt AM, Holmes TP, Leung B, Lambert KF, Orwig DA, Campbell FT, Rosenthal J, McCullough DG, Wildova R, Ayres MP, Canham CD, Foster DR, Ladeau SL, Weldy T (2016) Nonnative forest insects and pathogens in the United States: Impacts and policy options. *Ecol Appl* 26: 1437–1455. <https://doi.org/10.1890/15-1176>.
- Lowe S, Browne M, Boudjelas S, De Poorter M (2000) 100 of the world's Worst Invasive Alien Species. A selection from the Global Invasive Species Database, The Invasive Species Specialist Group, a specialist group of the Species Survival Commission (SSC) of the World Conservation Union (IUCN), p 12
- Mansfield S, McNeill MR, Aalders LT, Bell NL, Kean JM, Barratt BIP, Boyd-Wilson K, Teulon DAJ (2019) The value of sentinel plants for risk assessment and surveillance to support biosecurity. *NeoBiota* 48:1–24. <https://doi.org/10.3897/neobiota.48.34205>
- Maunder M, Culham A, Alden B, Zizka G, Orliac C, Lobin W, Bordeu A, Ramirez JM, Glissmann-Gough S (2000) Conservation of the Toromiro tree: case study in the management of a plant extinct in the wild. *Conserv Biol* 14:1341–1350. <https://doi.org/10.1046/j.1523-1739.2000.98520.x>
- Mehta SV, Haight RG, Homans FR, Polasky S, Venette RC (2007) Optimal detection and control strategies for invasive species management. *Ecol Econ* 61:237–245. <https://doi.org/10.1016/j.ecolecon.2006.10.024>
- Mendel Z, Protosav A, Sharon M, Zveibil A, Ben Yahuda S, O'Donnell K, Rabaglia R, Wysoki M, Freeman S (2012) An Asian ambrosia beetle *Etw Wallacea fornicatus* and its novel symbiotic fungus *Fusarium* sp. pose a serious threat to Israeli avocado industry. *Phytoparasitica* 40:235–238. <https://doi.org/10.1007/s12600-012-0223-7>
- Miller A J, Novy A, Glover J, Kellogg EA, Maul JE, Raven P, Jackson PW (2015) Expanding the role of botanical gardens in the future of food. *Nat Plant*. Article Number 15078. <https://doi.org/https://doi.org/10.1038/nplants.2015.78>
- Morales-Rodríguez C, Anslan S, Auger-Rozenberg M-A, Augustin S, Baranchikov Y, Bellahirech A, Burokienė D, Čepukoit D, Čota E, Davydenko K, Doğmuş Lehtijärvi HT, Drenkhan R, Drenkhan T, Eschen R, Franić I, Glavendekić M, de Groot M, Kacprzyk M, Kenis M, Kirichenko N, Matsiakh I, Musolin DL, Nowakowska JA, O'Hanlon R, Prospero S, Roques A, Santini A, Talgø V, Tedersoo L, Uimari A, Vannini A, Witzell J, Woodward S, Zambounis A, Cleary M (2019) Forewarned is forearmed: harmonized approaches for early detection of potentially invasive pests and pathogens in sentinel plantings. *NeoBiota* 47:95–123. <https://doi.org/10.1016/j.biocontrol.2017.07.009>
- Oldfield S (2007) Great botanic gardens of the world. New Holland, London
- Paap T, Burgess T, Wingfield MJ (2017) Urban trees: bridge-heads for forest pest invasions and sentinels for early detection. *Biol Invasions* 19:3515–3526. <https://doi.org/10.1007/s10530-017-1595-x>
- Paap T, de Beer ZW, Migliorini D, Nel WJ, Wingfield MJ (2018) The polyphagous shot hole borer (PSHB) and its fungal symbiont *Fusarium euwallaceae*: a new invasion in South Africa. *Australas Plant Path* 47:231–237. <https://doi.org/10.1007/s13313-018-0545-0>
- Paap T, Wingfield MJ, Burgess TI, Hulbert JM, Santini A (2020) Harmonising the fields of invasion science and forest pathology. *NeoBiota* 62:301–3032. <https://doi.org/10.1016/j.tree.2004.07.021>
- Pilkington LJ, Irvin NA, Hoddle MS, Triapitsyn SV, Carey BG, Jones WA, Morgan DJW (2005) Introduced parasitic wasps could control glassy-winged sharpshooter. *Calif Agric* 59:223–228. <https://doi.org/10.3733/ca.v059n04p223>
- Pimentel D, Zuniga R, Morrison D (2005) Update on the environmental and economic costs associated with alien-invasive species in the United States. *Ecol Econ* 52:273–288. <https://doi.org/10.1016/j.ecolecon.2004.10.002>

- Pluess T, Jarosik V, Pysek P, Cannon R, Pergl J, Breukers A, Bacher S (2012) Which factors affect the success or failure of eradication campaigns against alien species? PLoS ONE 7(826):e48157. <https://doi.org/10.1371/journal.pone.0048157>
- Powledge F (2011) The evolving role of botanical gardens. Bioscience 61:743–749. <https://doi.org/10.1525/bio.2011.61.10.3>
- Pratt CF, Constantine KL, Murphy ST (2017) Economic impacts of invasive alien species on African smallholder livelihoods. Glob Food Sec 14:31–37. <https://doi.org/10.1016/j.gfs.2017.01.011>
- Rigling D, Prospero S (2018) Cryphonectria parasitica, the causal agent of chestnut blight: invasion history, population biology and disease control. Mol Plant Pathol 19:7–20. <https://doi.org/10.1111/mpp.12542>
- Rodoni B, Gardner R, Giles R, Cole M, Wimalajeewa S (2002) National surveys did not detect *Erwinia amylovora* on host plants in Australia. Acta Hort 590:39–45. <https://doi.org/10.17660/ActaHortic.2002.590.3>
- Roques A, Rabitsch W, Rasplus J-Y, Lopez-Vaamonde C, Nentwig W, Kenis M (2009) Alien terrestrial invertebrates of Europe. In: DAISIE, (ed) Handbook of alien species in Europe. Springer, Berlin, pp 63–79
- Roques A, Fan J, Courtial B, Zhang Y, Yart A, Auger-ozenberg M, Denux O, Kenis M, Baker R, Sun J (2015) Planting Sentinel European trees in Eastern Asia as a novel method to identify potential insect pest invaders. PLoS ONE. <https://doi.org/10.1371/journal.pone.0120864>
- Salisbury A, Malumphy C, Halstead AJ (2011) First incursion of *Aloea australis* (Hemiptera: Miridae) and *Pulvinaria delottoi* (Hemiptera: Coccidae) in Europe, and three other Hemipteran insects imported from South Africa. Br J Ent Nat Hist 24:217–220
- Santini A, Ghelardini L, De Pace C, Desprez-Loustau M-L, Capretti P, Chandelier A, Cech T, Chira D, Diamandis S, Gaitniekis T, Hantula J, Holdenrieder O, Jankovsky L, Jung T, Jurc D, Kirisits T, Kunca A, Lygis V, Malecka M, Marçais B, Schmitz S, Schumacher J, Solheim H, Solla A, Szabo I, Tsopeles P, Vannini A, Vetraino A, Webber J, Woodward S, Stenlid J (2013) Biogeographical patterns and determinants of invasion by forest pathogens in Europe. New Phytol 197:238–250. <https://doi.org/10.1111/j.1469-8137.2012.04364.x>
- Santini A, Liebhold A, Migliorini D, Woodward S (2018) Tracing the role of human civilization in the globalization of plant pathogens. ISME J 12:647–652. <https://doi.org/10.1098/rsos.150370>
- Scalera R, De Man D, Klausen B, Dickie L, Genovesi P (2012) European code of conduct on zoological gardens and aquaria and invasive alien species. Council of Europe, New edition November 2016
- Scott-Brown AS, Hodgetts J, Simmonds MSJ, Collins DW (2018) Potential role of botanic garden collections in predicting hosts at risk globally from invasive pests: a case study using *Scirtothrips dorsalis*. J Pest Sci 91:60–611. <https://doi.org/10.2307/2261425>
- Sharrock SL, Blackmore S, Cheney J, Dunn C, Husby C, Jebb M, Kennedy K, Kiehn M, Kramer A, Kueffer C, Raven P, Reichard S, Rejmanek M, Oldfield S, Wyse Jackson P (2011) The biodiversity benefits of botanic gardens. Trends Ecol Evol 26:433. <https://doi.org/10.1016/j.tree.2011.05.008>
- Sheridan JE (1989) Quarantine risks imposed by overseas passengers. N Z J For Sci 19(338):346
- Smith P (2019) The challenge for botanic garden science. Plants People Planet 1:38–43
- Sweeney J, Anderson RS, Webster RP, Nevill R (2012) First records of *Orchestes fagi* (L.) (Coleoptera: Curculionidae: Curculioninae) in North America, with a checklist of the North American Ramphini. Coleopt Bull 66:297–304. <https://doi.org/10.1649/072.066.0401>
- Tehotet Tchoumi JM, Coetzee MPA, Rajchenberg M, Roux J (2019) Taxonomy and species diversity of *Ganoderma* species in the Garden Route National Park of South Africa inferred from morphology and multilocus phylogenies. Mycologia 111:730–747. <https://doi.org/10.1080/00275514.2019.1635387>
- Thakur MP, van der Putten WH, Cobben MMP, van Kleunen M, Geisen S (2019) Microbial invasions in terrestrial ecosystems. Nat Rev Microbiol 17:621–631. <https://doi.org/10.1038/s41579-019-0236-z>
- Walter R (2012) Economic relations between Europe and the world: Dependence and Interdependence. European History Online (EGO), published by the Leibniz Institute of European History (IEG). <http://ieg-ego.eu/en/threads/europe-and-the-world/economic-relations>. Accessed 21 Aug 2020
- Wingfield MJ, Brockerhoff EG, Wingfield BD, Slippers B (2015) Planted forest health: the need for a global strategy. Science 349:832–836. <https://doi.org/10.1126/science.aac6674>
- Wylie FR, Griffiths M, King J (2008) Development of hazard site surveillance programs for forest invasive species: a case study from Brisbane, Australia. Austral For 71:229–235. <https://doi.org/10.1080/00049158.2008.10675040>

Authors and Affiliations

Mesfin Wondafrash^{1,2}  · Michael J. Wingfield³  · John R. U. Wilson^{2,4}  ·
Brett P. Hurley¹  · Bernard Slippers³  · Trudy Paap³ 

¹ Department of Zoology and Entomology, Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria 0002, South Africa

² South African National Biodiversity Institute, Kirstenbosch Research Centre, Cape Town, South Africa

³ Department of Biochemistry, Genetics and Microbiology, Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria 0002, South Africa

⁴ Centre for Invasion Biology, Department of Botany and Zoology, Stellenbosch University, Stellenbosch, South Africa