

Comparative assessment on rodent impacts and cultural perceptions of ecologically based rodent management in 3 Afro-Malagasy farming regions

Natasha L. CONSTANT,^{1,2} Lourens H. SWANEPOEL,³ Samual T. WILLIAMS,^{3,4,5}
Voahangy SOARIMALALA,^{6,7} Steven M. GOODMAN,^{6,8} Apia T. MASSAWE,⁹
Loth S. MULUNGU,⁹ Rhodes H. MAKUNDI,⁹ Mashaka E. MDANGI,¹⁰ Peter J. TAYLOR¹
and Steven R. BELMAIN¹¹

¹SARChI Chair on Biodiversity Value and Change and Core Team Member of the Centre for Invasion Biology, School of Mathematical and Natural Sciences, University of Venda, Thohoyandou, Johannesburg, South Africa, ²Sustainable Places Research Institute, Cardiff University, Cardiff, UK, ³Department of Zoology, School of Mathematical and Natural Sciences, University of Venda, Thohoyandou, South Africa, ⁴Department of Anthropology, Durham University, Durham, UK, ⁵Institute for Globally Distributed Open Research and Education (IGDORE), Johannesburg, South Africa, ⁶Association Vahatra, Antananarivo, Madagascar, ⁷Institut des Sciences et Techniques de l'Environnement Université de Fianarantsoa, Fianarantsoa, Madagascar, ⁸Field Museum of Natural History, Chicago, Illinois, USA, ⁹Pest Management Centre, Sokoine University of Agriculture, Morogoro, Tanzania, ¹⁰Ministry of Agriculture, Tanzania and ¹¹Natural Resources Institute University of Greenwich, Kent, UK

Abstract

Rodents generate negative consequences for smallholder farmers in Africa that directly impact household and livestock damage, food security, and public health. Ecologically Based Rodent Management (EBRM) seeks sustainable solutions for the mitigation of rodent damage through assessments of rodent population dynamics, agro-ecosystems, and socio-cultural contexts. We adopt a comparative approach across 3 rural Afro-Malagasy smallholder farming regions in South Africa, Tanzania, and Madagascar to assess the household impacts of rodent pests and current perceptions and preferences associated with several rodent control measures. We conducted focus group questionnaires and interviews in different study site locations. Rodents assert multiple impacts on Afro-Malagasy farmers demonstrating recurrent and emerging agricultural and household costs, and public health impacts. We identify a significant knowledge gap in educating communities about the application of different EBRM approaches in favor of acute poisons that are perceived to be more effective. Cultural issues and taboos also have a significant impact on the social acceptance of rodent hunting as well as biological control using

Correspondence: Natasha Louise Constant, SARChI Chair on Biodiversity Value and Change and Core Team Member of the Centre for Invasion Biology, School of Mathematical and Natural Sciences, University of Venda, Private Bag X5050, Thohoyandou 0950, South Africa.

Email: constantn@cardiff.ac.uk

indigenous predators. We advocate for an enhanced investigation of the socio-cultural beliefs associated with different rodent practices to understand the factors underlying social acceptance. A collaborative approach that integrates the perspectives of target communities to inform the design of EBRM initiatives according to the specific agro-ecosystem and socio-cultural context is necessary to ensure programmatic success.

Key words: Africa, beliefs, crop damage, ecologically based rodent management, farmer survey

INTRODUCTION

The African continent is home to 463 species of rodents, 77 of which damage crops and 12–20 are significant crop pests (Monadjem *et al.* 2015). For resource-poor farmers across Africa and Madagascar, rodents pose a significant threat to food security and agricultural productivity (Swanepoel *et al.* 2017b). Studies of Afro-Malagasy farming communities revealed that while estimated crop losses varied between cropping stages, the highest losses peaked during the seedling (46% median loss) and maturity stages (15% median loss) (Swanepoel *et al.* 2017b). However, during pest outbreaks, crop losses for maize can reach damage levels of 80–100% (Mwanjabe & Leirs 1997; Mwanjabe *et al.* 2002). Rodents damage clothes, blankets, and furniture; contaminate food stores; and attack small stock (Garba *et al.* 2014). Rodents also represent threats to public health as reservoirs of zoonotic diseases, including Lassa fever, plague, leptospirosis, and toxoplasmosis (Duchemin & Bitam 2017; Saez *et al.* 2018). Therefore, rodents assert multiple impacts on humans including food security and public health problems across national and global scales, yet these problems are rarely explored among Afro-Malagasy communities (Ziwa *et al.* 2013; Mead 2018).

Concerns of the long-term efficacy of chemical rodenticides due to resistance and secondary poisoning contributed to the development of Ecologically Based Rodent Management (EBRM) approaches (Singleton *et al.* 1999). This method aims to achieve rodent population suppression by the application of alternative forms of rodent control, using ecologically sound methods (Makundi & Massawe 2011). Mitigation of impact is achieved through assessments of rodent ecology and considering agro-ecological and socio-economic contexts (Massawe *et al.* 2012). Tools used by EBRM include habitat modification to reduce rodent key resources (Makundi *et al.* 1999), biological control through the use of indigenous predators to limit rodent numbers (Mahlaba *et al.* 2017; Williams *et al.* 2018), and assessments of rodent ecology to explore the appropriate timing of rodent control measures through preventive action (Makundi *et al.* 1999).

Explorations of EBRM by communities experiencing problems with rodent damage are often complicated by the cost-effectiveness and social acceptance (Ninh *et al.* 2016). These factors are important to develop culturally specific approaches for rodent management. Advocates of EBRM call for increased engagement of different ethnic groups to understand how cultural factors influence the ways farmers perceive and conduct rodent management (Palis *et al.* 2013). Knowledge, attitude, and perception studies have been an important component of EBRM across Southeast Asia (Brown *et al.* 2008; Ngaomei & Singh 2016), and are increasing in scope across Africa (Meheretu *et al.* 2010; Mulungu *et al.* 2015), but are largely lacking in South Africa and Madagascar. An exploration of people's knowledge and perceptions of rodent damage and management may provide indirect yet insightful information for promoting EBRM in different contexts.

We adopt a comparative approach herein, drawing on different case studies among different Afro-Malagasy communities to identify cross-cutting themes to assess: (i) the multiple impacts of rodent pests, (ii) current perceptions in rodent management, and (iii) preferences of different rodent management methods. We investigate these interactions focusing on 3 smallholder farming regions in South Africa, Tanzania, and Madagascar to compare key sites and countries, with different histories of rodent pest management. Tanzania has focused on applied rodent research and knowledge extension with respect to agricultural damage and rodent population outbreaks (Swanepoel *et al.* 2017b), while Madagascar has traditionally perceived rodents as a public health concern due to endemic plague (Andrianaivoarimanana *et al.* 2013) and South Africa has a more advanced private sector rodent control industry (SAPCA 2019). Each country is experiencing growing issues with rodent pests in smallholder farming communities, yet they have contrasting agro-ecosystems and environmental and social-cultural contexts. Each case example presents the setting of each study site, specific problem focus, methods, and key findings. We then discuss cross-cutting themes in relation to the impacts of rodent pests and




Figure 1 Map showing the locations of the study sites in South Africa, Tanzania, and Madagascar.

perceptions and preferences associated with different rodent management interventions. The analysis suggests several implications for promoting EBRM among Afro-Malagasy farming communities.

MATERIAL AND METHODS

The study site centers on Afro-Malagasy smallholder farming regions in South Africa, Tanzania, and Madagascar (Fig. 1).

Subsistence or semi-subsistence farming is practiced in all cases, typically in small plots of 1 ha or less. Differences between villages, districts, and countries in population size, land use types, physical geography, crops, storage methods, language, and rodent pests are summarized in Table 1.

Focus groups were conducted in the Vhembe/Mopani District of South Africa and the Handeni District of

Tanzania from April to May 2014. Focus groups consisted of a range of gender groups, farming communities, village leaders, and extension officers. A total of 18 focus groups (range: 6–13 informants) were carried out in South Africa and 5 focus groups in Tanzania (12 informants per focus group). Questions were posed to each group in the local language and responses were noted down and translated into English.

Firstly, questions focused on assessing rodent damage by asking farmers a series of opened ended questions (whether rodents are a problem in the community, how do they know if they are a problem, types of problems caused) followed by questions exploring the specificities of damage caused to household, livestock, human health, and crops. Questions addressing crop damage asked informants to estimate the total crop damage per annum for crop species at different stages of crop development (crop fields, harvest, post-harvest, and stores).

Table 1 Summary of site-specific information on each study site in South Africa, Tanzania, and Madagascar

District/Province	South Africa			Tanzania		Madagascar		Toamasina Province	
	Vhembe District	Mopani District	Handeni District	Fianarantsoa Province	Antananarivo Province	Antananarivo Province	Antananarivo Province	Antananarivo Province	Antananarivo Province
Target villages (population sizes)	Vyeboom (3579)	Ka-Ndengeza (5026)	Kwamisisi (5013), Magamba (3186), Kwachaga (3062), Suwa (6592), and Mazingara (6402)	Mahasoabe (34,000) and Kianjavato (5000)	Ambatolaona (5000) and Mahatsara (2000)	Ambatolaona (5000) and Mahatsara (2000)	Ambatolaona (5000) and Mahatsara (2000)	Ambatolaona (5000) and Mahatsara (2000)	Beforona (13,000)
Language/culture	Tshivenda	Xitsonga	Kiswahili is spoken by 54% (URT 2016), belonging to the Zigua ethnic group, the majority of which are Muslims	Principally Malagasy dialects and secondarily French are spoken. Ethnic groups include Betsileo, Tanala, Merina, Bezanozano, Vakinakarata, Betsimisaraka, Sakalava, and Antaisaka					
Elevation (m)	700	580	581	190–1100	600–2600	600–2600	600–2600	600–2600	0–1500
Rainfall (pa) (mm)	700–800		800–1500 (lower altitudes of Handeni), 400 (North Pare Mts)	1370	1317	1317	1317	1317	3246
Wet season	October–March		October–December and March–June	December–April	November–March	November–March	November–March	November–March	February–April
Dry season	May–August		July–September	July–December	April–October	April–October	April–October	April–October	September–November
Natural ecosystems	Granite Lowveld and Rocky Bushveld	Gravelotte	Semi deciduous Miombo Forest of the Zambesian Centre of endemism and the undifferentiated forests of the Zanzibar-Inhambane regional mosaic	Humid forest in Central Highlands dominated by <i>Weinmannia</i> (Cunoniaceae) and <i>Tambourissa</i> (Monimiaceae)	Middle-altitude dense rain forest with dominant tree genera, <i>Tambourissa</i> , <i>Weinmannia</i> , <i>Symphonia</i> , <i>Dalbergia</i> , and <i>Vernonia</i>	Middle-altitude dense rain forest with dominant tree genera, <i>Tambourissa</i> , <i>Weinmannia</i> , <i>Symphonia</i> , <i>Dalbergia</i> , and <i>Vernonia</i>	Middle-altitude dense rain forest with dominant tree genera, <i>Tambourissa</i> , <i>Weinmannia</i> , <i>Symphonia</i> , <i>Dalbergia</i> , and <i>Vernonia</i>	Middle-altitude dense rain forest with dominant tree genera, <i>Tambourissa</i> , <i>Weinmannia</i> , <i>Symphonia</i> , <i>Dalbergia</i> , and <i>Vernonia</i>	Coastal rainforest

(Continued)

Table 1 Continued

	South Africa	Tanzania	Madagascar	Toamasina Province
District/Province	Vhembe District	Mopani District	Handeni District	Antananarivo Province
Land use types	Cropping area, communal grazing land, and human settlement	Cropping area, human settlements, and degraded forest areas from agriculture and firewood and timber harvesting	Agricultural land and forested patches transformed from slash and burn practices and irrigated and rainfed upland rice cultivation with some dry crop fields	
Storage methods	Polypropylene bags in bedrooms or other secure areas of the house	Maize grains are stored in the roof of houses in polypropylene bags and used for cooking or selling	Rice either stored in granaries with difficult access for rodents and polypropylene bags stored in houses	
Main crops	Maize (<i>Zea mays</i>) and bambara nuts (<i>Vigna subterranea</i>) are dominant, but also pumpkins (<i>Curcubita</i> spp.), peanuts (<i>Arachis hypogaea</i>), and beans (<i>Phaseolus vulgaris</i>)	Maize, cassava (<i>Manihot esculenta</i>), beans, peanuts, banana (<i>Musa</i> spp.), sugar cane (<i>Saccharum</i> spp.), and different types of vegetables	Rice (<i>Oryza glaberrima</i>) from irrigated rice valleys and rain-fed uplands associated with slash-and-burn agriculture in areas of cleared forest or reclaimed secondary habitats. Other common food plants cultivated in these areas include cassava, peanuts, maize, sweet potato (<i>Ipomoea batatas</i>), and beans	Beforona: irrigated and upland rice
Main rodent pests	Rats (<i>Rattus rattus</i> , <i>R. norvegicus</i> , <i>R. tanezumii</i>), multimammate mice (<i>Mastomys coucha</i> , <i>M. natalensis</i>), and gerbils (<i>Gerbilliscus brantissii</i> , <i>G. leucogaster</i>)	Natal multimammate mouse (<i>M. natalensis</i>), African grass rat (<i>Arvicanthis neumanni</i>)	Rats (<i>R. rattus</i> , <i>R. norvegicus</i>) and house mouse (<i>Mus musculus</i>)	

Secondly, informants were asked to identify their level of agreement toward different statements by responding either “yes,” “no,” or “maybe.” The first series of statement explored perceptions of the effectiveness of different EBRM approaches, such as synchronizing the timing of crop planting, cleaning and improving hygiene on farms, and hunting and trapping rodents. The second series of statements explored perceptions of the effectiveness of individual and community management approaches. The third series of statements identified perceptions of the effectiveness of rodenticides such as poisons and anti-coagulants.

Thirdly, questions focused on preferences for the uptake of different types of rodent management, the perceived costs of different strategies, future management approaches, the application of biological control (using cats, birds of prey, snakes, and other reptiles), and perceptions toward the promotion of these approaches in their community. Each group was encouraged to discuss each of their responses, and in some cases represent collective answers. In certain cases, respondents often gave more than 1 answer to an individual question.

In Madagascar, 2 types of questionnaires were used during the interviews: (i) a household questionnaire discussed with people in their home setting and (ii) another for local women and men ranging in age from 18–60 years old as they worked in their agricultural fields. Individual questionnaires were used instead of focal groups because most people do not express personal ideas when in group settings. Data collection took place in local villages between June and July 2014 for Kianjavato and Mahasoabe and August and September 2015 for Ambatolaona, Mahatsara, and Beforona. Before the interviews took place, a review of the local fokontany (single or group of nearby villages) structure was conducted to determine the number of immediate villages and households. This list of households served as the basis for the selection of households and interviewees. Questionnaires focused on the same questions for the focus group interviews in South Africa and Tanzania. The original questionnaire was written in English and translated into French and investigators used the Malagasy language to conduct interviews.

Fieldwork was approved by the Research Ethics Committees of Sokoine University of Agriculture for activities in Tanzania, University of Venda for activities in South Africa, and Association Vahatra for activities in Madagascar. All qualitative data generated from focus groups and questionnaires were condensed as summary statistics based on the percentage of responses generated from informants.

RESULTS

Rodent impacts

Household damage

Rodents were reported as a problem by all respondents in the Vhembe/Mopani, Handeni District, and 98.2% of informants in eastern Madagascar. In the Vhembe/Mopani District, household damage was most commonly associated with damage to buildings (33.4%), food (30.0%), and personal possessions (28.3%) (Fig. S1a, Supporting Information). In the Handeni District, rodent household damage mainly targeted clothes (26.3%) and food storage (26.3%) (Fig. S1b, Supporting Information). In eastern Madagascar, reported rodent problems were most commonly associated with damage to buildings (28.4%), personal possessions (23.6%), clothes (22.9%), and crop damage (20.3%) (Fig. S1c, Supporting Information).

Crop damage

In the Vhembe/Mopani District, annual crop damage in fields was greatest for maize, peanuts, and bambara nuts (>40%) (Table 2). The Handeni District, experienced similar levels of annual crop damage (>40%) for maize, sunflowers (*Helianthus annuus*), and peanuts (Table 2). In the Handeni District, crop damage occurred throughout the planting, seedling, and maturity stage for maize (Table 2).

In eastern Madagascar, annual crop damage at harvest was high (>20%) for maize, rice, bambara nuts, peas, cassava, beans, and fruits (Table 2). In contrast, estimated crop damages at harvest (annual kg of crop damage per household) were variable, with high damages estimated for bambara nuts (only a single response: 150 kg), compared to rice (range: 1.2–60 kg) and beans (range: 6.3–12 kg) (Table 2).

In the Vhembe/Mopani District, post-harvest crop losses were higher than reports of damage in crop fields for maize and peanuts (>70%) and estimated crop losses in food stores were highest for maize and peanuts (>30%) (Table 2). In the Handeni District, estimated post-harvest crop damage were high for maize and sunflowers (>30%) (Table 2). In eastern Madagascar, post-harvest crop damage (annual kg of crop damage per household) was relatively low for maize (range: 3–20 kg) and cassava (range: 2–10 kg) (Table 2). Food damage in stores (annual kg of crop damage per household) was also low for bambara nuts (only a single response: 3 kg) and rice (range: 1.5–60 kg) (Table 2).

Table 2 Summary of estimates of crop damage by rodents for households in South Africa, Tanzania, and Madagascar per annum

Species	Crop damage in fields/% total crop damage per household	Crop damage at harvest/% total crop damage per household	Post-harvest crop damage/% total crop damage per household	Crop damage in stores/% total crop damage per household
<i>South Africa</i>				
Maize	Mean: 49% (range: 40–60%)	—	Single response: 80%	Mean: 49% (range: 20–70%)
Peanuts	Single response: 55%	—	Mean: 73% (range: 70–80%)	Mean: 46% (range 30–55%)
Bambara nuts	Mean: 47% (range: 40–50%)	—	—	—
Beans	—	—	Mean: 73% (range: 70–85%)	—
<i>Tanzania</i>				
Maize	Range: 50–100%	At planting (range: 50–100%), seedling (range: 10–20%), and maturity (range: 10–40%)	Stored grains (range: 30–60%)	Range: 30–50%
Sunflowers	Single response: 60%	At planting: single response 10%	Sunflowers whole seed: range: 30–80%	Range: 50–60%
Peanuts	Single response: 40%	At maturity: single response 20%	Single response: 40%	—
Cowpeas	—	At maturity: 20%	—	—
<i>Madagascar</i>				
Maize	—	Mean: 53% (range: 30–80%)	Mean: 11.5 kg (range: 3–20 kg)	—
Rice	—	Mean: 38% (range: 20–100%); mean: 13 kg (range 1.2–60 kg)	—	Mean: 13.3 kg (range: 1.5–60 kg)
Bambara nuts	—	Mean: 33% (range: 25–50%); one response: 150 kg	—	Single response: 3 kg
Peas	—	Single response: 33%	—	—
Cassava	—	Single response 25%	Mean: 8 kg (range 2–10 kg)	—
Beans	—	Single response: 75%; mean: 6.3 kg (range 6.3–12 kg)	—	—
Fruits	—	Single response: 33%	—	—

Livestock damage

In the Vhembe/Mopani District, 40.9% of participants reported no rodent problems relating to livestock rodent damage to livestock (Fig. 2a), although some evidence of livestock damage was mostly evidenced through feed

contamination (31.8%) and attacks on chickens/eggs (22.7%) (Fig. 2a).

In the Handeni District, rodent damage to livestock was corroborated mainly through attacks on chickens/eggs (80.0%) and calf wounds (20.0%) (Fig. 2b). In




Figure 2 Percentage of responses reporting livestock damage caused by rodents in the (a) Vhembe/Mopani District of South Africa ($n = 22$), (b) Handeni District of Tanzania ($n = 5$), and (c) eastern Madagascar ($n = 512$).




Figure 3 Percentage of responses reporting public health impacts caused by rodents in the (a) Vhembe/Mopani District of South Africa ($n = 17$), (b) Handeni District of Tanzania ($n = 12$), and (c) eastern Madagascar ($n = 512$).

eastern Madagascar, damage caused by rodents was commonly associated with attacks on chickens/eggs (65.4%), while 31.6% reported no livestock damage (Fig. 2c).

Public health

In the Vhembe/Mopani District, the majority of respondents reported no health impacts from rodents (94.1%), while 5.9% reported rodent bites (Fig. 3a).

In the Handeni District, rodent-related health problems were caused by rodent bites (41.7%), although some respondents reported knowledge of rodents spreading disease (8.3%), such as tetanus (8.3%), dysentery (8.3%), plague (8.3%), and unknown fevers (8.3%) (Fig. 3b).

In eastern Madagascar, 67.2% of informants reported no health impacts (Fig. 3c). However, some health impacts mentioned by informants were associated with unknown fevers (25.2%), fleas (2.3%), contaminated food (2.0%) and water (1.2%), coughs (0.4%), insomnia (0.4%), scabies (0.4%), and unspecified diseases (0.2%) (Fig. 3c).

Current perceptions in rodent management

EBRM Approaches

Over half of respondents across all case studies (Vhembe/Mopani District: 50%, Handeni District: 100%,

and eastern Madagascar: 55.5%) disagreed that synchronous planting of crops can reduce rodent populations (Fig. 4a).

The majority of farming communities in the Vhembe/Mopani District (55.6%) and eastern Madagascar (65.6%) supported the view that improving cleanliness on farms and surrounding areas can reduce rodent populations, while in the Handeni District, even numbers of people either supported (40%) or were against this view (40%) (Fig. 4a). In the Vhembe/Mopani District (72.2%) and eastern Madagascar (48.4%) of informants disagreed that hunting reduces the severity of observed rodent damage, while all informants in the Handeni District “maybe” agreed with this statement (Fig. 4a). Similarly, the majority of informants in the Vhembe/Mopani District (72.2%) and eastern Madagascar (61.0%) supported the view that trapping reduces the severity of rodent damage, while in the Handeni District, 40% agreed with this statement (Fig. 4a).

Individual and community rodent management

In eastern Madagascar, 61.2% of informants supported the view that individual rodent control is best to control their damage, while the majority of informants in the Vhembe/Mopani District (66.7%) and Handeni District (100%) disagreed with this statement (Fig. 4b). The majority of informants across the case studies agreed rodent management must be done continuously in order to be effective and community rodent control at any time during the cropping season is most effective (Fig. 4b). The majority of farmers in the Handeni District (100%) and eastern Madagascar (69.9%) supported the view that community rodent control at a specific stage of crop growth is most effective (Fig. 4b). In contrast, 58.8% of informants disagreed with this statement in the Vhembe/Mopani District (Fig. 4b).

Rodenticides

The majority of informants across the Afro-Malagasy study regions agreed that rodenticides reduce the severity of observed rodent damage (Fig. 4c). In the Vhembe/Mopani District, all informants agreed that acute poisons were “maybe” effective and safe to use, while in the Handeni District 100% disagreed with this statement (Fig. 4c). In the Vhembe/Mopani District, equal numbers of respondents supported (44.4%) and were against (44.4%) the view use that anticoagulants are an effective and safe poison to use, while 60% of informants in the Handeni District “maybe” agreed with this statement (Fig. 4c).

Preferences for rodent control

Uptake of rodent control measures

In the Vhembe/Handeni District, popular methods for rodent control in crop fields and households included acute poison (34.6%) and chronic poison (23.1%), which were all cited as effective and cheap management approaches (Fig. S5a and Table S1, Supporting Information).

In the Handeni District, acute poison (50.0%) and the non-steroidal anti-inflammatory drug indomethacin (30.0%) were popular forms of rodent control (Fig. 5b). Local support for acute poisons such as zinc phosphide was attributed to fast-acting results and applications were preferred in food stores and homesteads as opposed to crop fields (Table S1, Supporting Information). The economic costs of using acute poison and indomethacin were perceived as low, justifying their widespread use (Table S1, Supporting Information). In eastern Madagascar, snap traps (27.6%), granaries (17.4%), and cats (16.3%) were popular forms of rodent control (Fig. 5c) owing to their cost-effectiveness (Table S1, Supporting Information).

The majority of community members in the Vhembe/Mopani District stated they would continue to use snap traps (43.4%), anticoagulants (e.g. Rattex) (34.9%), followed by poison (e.g. Aldicarb) (17.4%) to manage rodent problems in the future because they were cheap and effective (Fig. S2a, Supporting Information). In the Handeni District, 83.3% and 16.7% of informants supported the continued future use of poisons and indomethacin, respectively (Fig. S2b, Supporting Information). In eastern Madagascar, cats (25.1%) and rodenticides (24.2%) were cited as the most popular forms of rodent control for continued use in the future (Fig. S2c, Supporting Information).

Biological control

The majority of informants across Afro-Malagasy farming communities believed that cats were effective in controlling rodent problems in their villages (Fig. 6).

In the Handeni District and eastern Madagascar, 83.3% and 61.0%, respectively, of respondents believed that owls, snakes, and other animals were effective in controlling rodent numbers (Fig. 6). However, in the Vhembe/Mopani District, 76.0% of respondents believed that owls, snakes, and other animals were ineffective (Fig. 6). The presence of snakes and owls was discouraged because of a fear of snakes and beliefs associated with owls such as witchcraft and bringing “evil to the village.” Informants who supported the use of cats to control rodents did not




Figure 4 Percentage of responses demonstrating current perceptions of (a) EBRM approaches, (b) individual and community rodent management, and (c) rodenticides across the Vhembe/Mopani District of South Africa ($n = 18$), Handeni District of Tanzania ($n = 5$), and eastern Madagascar ($n = 508$)




Figure 5 Percentage of responses supporting the application of different control methods for rodent management in the (a) Vhembe/Mopani District of South Africa ($n = 16$), (b) Handeni District of Tanzania ($n = 10$), and (c) eastern Madagascar ($n = 786$).




Figure 6 Percentage of responses supporting the effectiveness of cats in the Vhembe/Mopani District of South Africa ($n = 17$), Handeni District of Tanzania ($n = 12$), and eastern Madagascar ($n = 508$); the effectiveness of birds (owls), snakes, and other animals in the Vhembe/Mopani District of South Africa ($n = 17$), Handeni District of Tanzania ($n = 12$), and eastern Madagascar ($n = 498$); and the promotion of these approaches in their communities in the Vhembe/Mopani District of South Africa ($n = 16$), Handeni District of Tanzania ($n = 16$), and eastern Madagascar ($n = 447$).

tolerate the presence of other predators such as snakes and owls. Common informant statements included: “we will not tolerate other animals, other than cats” and “yes to cats, no to snakes and owls.”

In the Vhembe/Mopani District and eastern Madagascar, 100% and 85.5%, respectively, of community members were against the promotion of using indigenous

predators for rodent control, stating that the presence of these animals was ineffective against rodents (Fig. 6). In contrast, a high percentage of community members (62.5%) in the Handeni District supported the promotion of indigenous predators for rodent control (Fig. 6). Those against the use of biological control methods indicated that local beliefs and the perception of snakes as

dangerous animals prohibited them from supporting these strategies.

DISCUSSION

Rodent impacts

Rodent-induced problems are uniformly significant and widespread across Afro-Malagasy farming regions, causing household, crop, and livestock damage and public health threats.

Crop damage

In all countries, rodent damage to crops (pre- and post-harvest) was perceived to be of considerable importance. In terms of crop damage, our study revealed the importance of understanding the specificities of the agro-ecosystem and cropping systems to inform effective EBRM at local scales. For example, EBRM approaches should focus on cropping systems of high economic value such as maize (South Africa and Tanzania) and rice (eastern Madagascar) where socio-economic losses due to rodent damage are perceived to be especially high. Annual crop yields for smallholder maize farmers in the Limpopo Province of South Africa have been estimated at 320–1600 kg·ha⁻¹·year⁻¹ (Rootman & Stevens 2016). In light of these figures, our reported estimates of 80% losses for maize can constitute between 256–1280 kg·ha⁻¹·year⁻¹. In the Tanga region of Tanzania, crop yields for maize have been estimated at 604–1500 kg·ha⁻¹·year⁻¹ (Kaliba *et al.* 1998; URT 2007) therefore, reported losses can represent half (30–750 kg·ha⁻¹·year⁻¹) or a total loss of all crop yields. High crop damage for maize (>80%) has also been documented from field trials in Tanzania (Mwanjabe & Leirs 1997; Mwanjabe *et al.* 2002), confirming our results.

Conclusions based on estimates of crop damage at different stages (in the field, at harvest, post-harvest, and in stores) obtained from farmers in our study generally concur well with conclusions based on estimates from published field trials (Table S2, Supporting Information, based on the review of Swanepoel *et al.* 2017b). For example, greatest losses of maize due to rodents occur at the seedling stage (Myllymaki 1989; Mulungu *et al.* 2007), while losses of rice tend to be highest at the maturity stage of rice. EBRM should be focused on the most vulnerable stages for each crop type and region. Post-harvest estimated losses to rodents were high but variable across all countries, although actual losses reported in Madagascar (1.5–60 kg·ha⁻¹·year⁻¹) appeared

under-estimated compared to values published for smallholder rice farmers in eastern Madagascar (Hume 2006; 1500–3000 kg·ha⁻¹·year⁻¹). In all countries, polypropylene sacks are used to reduce storage maize and rice losses (Duplantier & Duchemin 2003; Leclerc-Madlala & Janowski 2004; Shabani *et al.* 2015).

Rodent-borne diseases

Rodent-borne zoonoses are common in Africa, including a number of communicable and infectious viruses, bacteria, helminthes, and protozoa (Meerburg *et al.* 2009). Compared to a low knowledge and perception of rodent-borne diseases in South Africa revealed by our study, an enhanced awareness of rodent borne diseases in Tanzania and Madagascar may be attributed to a history of public health and rodent control campaigns implemented and contemporary problems with endemic plague, which can cause public health emergencies in these countries (Duplantier & Rakotondravony 1999; Makundi *et al.* 2015). The seasonal epidemiology of plague in Madagascar has led to annual outbreaks for the last 30 years; however, plague has been a problem in Madagascar and Tanzania for over 100 years (Mead 2018). The symptoms of illness mentioned by communities would suggest that rickettsial and leptospiral infections are important health issues in Tanzania and Madagascar and several prevalence studies support these community reports (Key 1990; Crump *et al.* 2013).

Current perceptions in rodent management

The study explored local perceptions of several EBRM approaches as alternatives to rodenticides. In some cases, community perceptions reflected lack of uptake of research results; for example, while most communities did not agree, research has demonstrated that synchronized planting may limit rodent damage for cereals, such as maize (Makundi *et al.* 2010). Synchronizing planting may maximize the length of the fallow period causing rodent and populations to decrease because of a lack of food and shelter and a reduction in the duration of the breeding season (Leung *et al.* 1999). On the other hand, our surveys showed good awareness among all Afro-Malagasy communities sampled of the importance of sanitation programs for rodent control as confirmed by research undertaken in Africa (Makundi *et al.* 1999; Taylor *et al.* 2008). The promotion of other land use practices that may lead to the removal of rodent habitats may also be important. In Tanzania and Madagascar, where farmers practice slash and burn techniques after harvest or before planting maize

crops, this temporarily clears food sources and shelter for rodents and can act as important strategy for managing habitats that support high numbers of rodents (Duplantier & Rakotondravony 1999). In Madagascar, rodent abundance is lower in dry farming areas that provide food for rodents for short periods of time and the plots are often burned once per year (Duplantier & Rakotondravony 1999). Therefore, a combination of crop synchrony coupled with productive land use management and sanitation programs may be useful in all the case study regions to reduce the availability of food resources for rodents.

Although results were mixed, in Tanzania, informants generally endorsed the use of trapping to control rodents. Some countries, such as Madagascar, have a history of trapping campaigns implemented by plant protection technicians to enhance local awareness and action on the ground (Duplantier & Rakotondravony 1999). Trapping of rodents at the household level is appropriate in rural agricultural settings, particularly in our case study regions, where farmers lack other adequate resources for rodent control (Makundi *et al.* 1999; Taylor *et al.* 2008). Hunting has also been suggested as one of many approaches to tackle rodent populations, particularly in South-East Asia where rodents have been hunted through collective action by targeting rodent burrow systems or government bounty campaigns (Singleton *et al.* 2007). Hunting was perceived as ineffective in reducing the severity of observed rodent damage in South Africa and eastern Madagascar, while in Tanzania, all informants stated hunting “maybe” reduces rodent damage. The lack of support for hunting in some of the study regions may in part be explained by cultural beliefs and prohibitions that constrain social acceptance of the hunting of animals for food. During focus group discussions in Tanzania, those against hunting were largely Muslims who explained that hunting is usually not undertaken by local people because it is associated with food consumption and Islamic beliefs prohibit people from eating rodent meat. Similarly, in eastern Madagascar, the hunting of rodents for food is considered dirty, disgusting, and taboo; this conclusion is supported by data from the Northeast (Borgerson 2015).

Although individual rodent control was preferred among smallholder farmers in eastern Madagascar, our findings demonstrate a uniform support for community rodent control measures across the Afro-Malagasy farming regions that would provide a strong foundation for engaging such groups in collective action and cultivating a shared sense of responsibility for EBRM. Coordinated community methods for EBRM have been encouraged to increase the efficacy of rodent control compared to

individual approaches because labor efforts are shared across larger-scales (Palis *et al.* 2011). Some African cultures, such as the Vhavenda in the Vhembe District, have a strong history of community participation in agricultural activities; for example, clearing, weeding, and harvesting of crops were often completed in semi-communal labor groups (Stayt 2018). Coordinated community methods for rodent control can help reduce rodent population growth through breeding or immigration if done continuously (Belmain *et al.* 2015). Similarly, synchronizing control methods over relatively short periods of time at key periods, such as the onset of breeding, can also be effective, but this requires a detailed understanding of rodent population dynamics within particular agro-ecosystems (Brown & Khamphoukeo 2007).

Preferences for rodent control

Although all countries endorsed rodenticides, Tanzania and South Africa preferred acute rodenticides, while Madagascar preferred a wider variety of options, raising concerns of ethical application. The widespread application of rodenticides across Afro-Malagasy farming regions is linked to their historical use, private sector promotion, and government interventions, such as the free or subsidized distribution of rodenticides (Makundi *et al.* 1999). The preference for cheaper and faster-acting rodenticides which have negative environmental effects in terms of the risk of non-target species mortality poses a challenge for EBRM, requiring concerted educational programs across all countries.

Biological control of rodents using domestic and indigenous predators such as cats, dogs, raptors, and reptiles has garnered increasing research interest in Africa. Recent studies have demonstrated that the dual presence of cats and dogs in homesteads can reduce rodent activity, creating a “landscape of fear” among foraging rodents (Mahlaba *et al.* 2017). Some studies have explored the potential positive impacts of using nest boxes and perches to encourage owls and other avian predators into agricultural landscapes to reduce rodent numbers (Labuschagne *et al.* 2016). The feasibility of employing biological control measures for EBRM is dependent on the social acceptance of these species. Across the study regions, there was a consensus that domestic cats were an effective measure for controlling rodent pests and were socially acceptable owing to their familiarity as domestic animals. However, birds of prey and reptile predators were perceived as ineffective in South Africa and effective in Tanzania and Madagascar. Only rural farmers in Tanzania demonstrated a willingness to promote biological control

measures in their community compared to South Africa and Madagascar. Lack of support for the implementation of avian and reptilian predators for EBRM was linked to negative perceptions of snakes as dangerous because they provide ecosystem disservices, harmful to human health. Among the Vhavenda in South Africa, owls are believed to possess supernatural messages such as predicting sickness and through their association with witches' familiars (Mabogo 1990). Historically in all countries, witchcraft has been recognized by the state through law and through state practices that officialize witch-hunting (Ellis 2002; Kohnert 2003; Mesaki 2009). A belief in the magical properties of animals and occult practices that harness the harmful properties of magical creatures, such as witches using snakes to attack their victims (Bjerke 1969) have led to their persecution in Tanzania (Holmes *et al.* 2018). Among the Zigua of Tanzania, snakes are represented as metaphors for addressing changes in ways of life deemed negatively by the Zigua through discussions of the country's history of the slave trade and colonization (Walz 2016). In certain Malagasy cultural groups, certain animal species are entwined with local taboos, known locally as "*fady*." Negative taboos linked to snakes resulted in their frequent killing (Holmes *et al.* 2018). Our findings point to the importance of understanding historical and cultural beliefs that may influence social acceptance of EBRM approaches, particularly, those involving taboo animals.

CONCLUSION

Our study shows that a wide range of significant rodent impacts are universally experienced at household, agricultural, and public health levels across smallholder communities in South Africa, Tanzania, and Madagascar. The scale of these impacts, especially to farmers, and the vulnerable stages of major cropping systems revealed by our study is borne out by available field studies in Africa (Table S2, Supporting Information, summarized by Swanepoel *et al.* 2017b). When it comes to awareness and implementing of EBRM, uptake by communities may vary between countries. For example, government programs about frequent plague outbreaks have led to greater awareness of diseases risks of rodents in Madagascar and Tanzania compared to South Africa. Cultural issues and social taboos also have a significant impact, for example, on acceptance of hunting of rodents as well as biological control by encouraging species such as owls and snakes. We identify a large gap in educating communities about the use of more environmentally acceptable, second-generation rodenticides in favor of

fast-acting acute poisons that are perceived to be more effective.

In conclusion, the promotion of EBRM requires a more nuanced and targeted assessment of the cost-effectiveness, social acceptance, and ecological specificity when appraising rodent control measures in different contexts. Secondly, identifying factors that enable or constrain decisions for the uptake of different EBRM strategies is critical for developing culturally appropriate interventions to enhance tolerance among target communities. Researchers also need to understand local socio-cultural beliefs associated with different rodent practices that may enable or hinder EBRM uptake before project implementation. Thirdly, enhancing knowledge of the sustainability and appropriate application of different EBRM measures will require a multifaceted approach, where educational initiatives to address knowledge gaps are designed to be culturally sensitive in accordance with the context of the target community and coupled with scientific and local knowledge of rodent and predator population dynamics. Finally, implementing EBRM requires a shift toward participatory planning processes that integrate the perceptions and concerns of local people, practitioners, and policy makers to advocate for more sustainable and coordinated approaches.

ACKNOWLEDGMENTS

We thank all community participants involved in focus group discussions and questionnaires, and all university staff who provided field support across the 3 countries. This research was funded by 8 grants: (1) StopRats: European Union 9th European Development Fund grant from the African Caribbean and Pacific Science and Technology Programme (FED/2013/330-223); (2) African Union (EcoRodMan: AURGII/1/006/2016); (3) United Kingdom's Department for International Development (AgriTT/894); (4) SASOL Agricultural Trust; (5) South African National Research Foundation (LHS #88179); (6) South African National Research Foundation and Department of Science and Technology through the South African Research Chair in Biodiversity Value and Change hosted by the University of Venda and co-hosted by the Centre for Invasion Biology, Stellenbosch University (NC, PT #87311); (7) South African National Research Foundation (STW: UID 115040 and UID 107099); and (8) MEM also acknowledges the support of the Tanzania Commission for Science and Technology (COSTECH). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

REFERENCES

- Andrianaivoarimanana V, Kreppel K, Elissa N *et al.* (2013). Understanding the persistence of plague foci in Madagascar. *PLoS Neglected Tropical Diseases* **7**, e2382.
- Belmain SR, Htwe NM, Kamal NQ, Singleton GR (2015). Estimating rodent losses to stored rice as a means to assess efficacy of rodent management. *Wildlife Research* **42**, 132–42.
- Bjerke S (1969). The high god among the Zinza of north-western Tanzania. *Numen* **16**, 186–210.
- Borgerson C (2015). Illegal hunting on the Masoala Peninsula of Madagascar: Its extent, causes, and impact on lemurs and humans (PhD thesis). University of Massachusetts, Amherst, MA.
- Brown PR, Khamphoukeo K (2007). Farmers' knowledge, attitudes, and practices with respect to rodent management in the upland and lowland farming systems of the Lao People's Democratic Republic. *Integrative Zoology* **2**, 165–73.
- Brown PR, Yee N, Singleton GR *et al.* (2008). Farmers' knowledge, attitudes, and practices for rodent management in Myanmar. *International Journal of Pest Management* **54**, 69–76.
- Crump JA, Morrissey AB, Nicholson WL *et al.* (2013). Etiology of severe non-malaria febrile illness in northern Tanzania: a prospective cohort study. *PLoS Neglected Tropical Diseases* **7**, e232.
- Duchemin JB, Bitam L (2017). Les puces (Siphonaptera). In: Duvallat G, Fontenille D, Robert V, eds. *Entomologie Médicale et Vétérinaire* Quæ, IRD, Marseille and Versailles, France.
- Duplantier JM, Rakotondravony D (1999). The rodent problem in Madagascar: Agricultural pest and threat to human health. In: Singleton G, Hinds L, Leirs H, Zhang Z, eds. *Ecologically-Based Rodent Management*. ACIA, Canberra, Australia.
- Duplantier JM, Duchemin JB (2003). Introduced small mammals and their ectoparasites: A description of their colonisation and its consequences. In: Goodman SM, Benstead JP, eds. *The Natural History of Madagascar*. The University of Chicago Press, Chicago, IL.
- Ellis S (2002). Witch-hunting in central Madagascar 1828–1861. *Past & Present* **175**, 90–123.
- Garba M, Kane M, Gagare S *et al.* (2014). Local perception of rodent-associated problems in Sahelian urban areas: a survey in Niamey, Niger. *Urban Ecosystems* **17**, 573–84.
- Holmes G, Smith TA, Ward C (2018). Fantastic beasts and why to conserve them: animals, magic and biodiversity conservation. *Oryx* **52**, 231–9.
- Hume DW (2006). Swidden agriculture and conservation in eastern Madagascar: stakeholder perspectives and cultural belief systems. *Conservation and Society* **4**, 287.
- Kaliba AR, Verkuijl H, Mwangi W *et al.* (1998). *Adoption of maize production technologies in Eastern Tanzania*, International Maize and Wheat Improvement Center (CIMMYT), The United Republic of Tanzania and the southern African Centre for Cooperation in Agricultural Research (SACCAR), Mexico, DF.
- Key GE (1990). Control of the African striped ground squirrel, *Xerus erythropus*, in Kenya. Proceedings of the Fourteenth Vertebrate Pest Conference 1990; 6–8 Mar 1990, Sacramento, California. University of California, Davis, CA.
- Kohnert D (2003). Witchcraft and transnational social spaces: witchcraft violence, reconciliation and development in South Africa's transition process. *Journal of Modern African Studies* **41**, 217–45.
- Labuschagne L, Swanepoel LH, Taylor PJ, Belmain SR, Keith M (2016). Are avian predators effective biological control agents for rodent pest management in agricultural systems? *Biological Control* **101**, 94–102.
- Leclerc-Madlala S, Janowski M (2004). Perceptions and practices relevant to the transmission of plague, leptospirosis and toxoplasmosis. *NRI Report 2781* [Online]. Available from http://projects.nri.org/ratzooman/docs/mapate_final_anthropology_report.pdf://nri.org/projects/bandicoot/docs/Taylor_Integ_Zoo.pdf
- Leung LKP, Singleton GR, Rahmini S (1999). Ecologically-based population management of the rice-field rat in Indonesia. In: Singleton GR, Hinds LA, Leirs H, Zhang Z, eds. *Ecologically-based Management of Rodent Pests*. ACIAR, Canberra, Australia.
- Mabogo DEW (1990). The ethnobotany of the Vhavenda (Masters thesis). University of Pretoria, Pretoria, South Africa.
- Mahlaba TAM, Monadjem A, McCleery R, Belmain SR (2017). Domestic cats and dogs create a landscape of fear for pest rodents around rural homesteads. *PLoS ONE* **12**, e0171593.
- Makundi RH, Oguge NO, Mwanjabe PS (1999). Rodent pest management in East Africa—an ecological approach. In: Singleton G, Leirs H, Zhang Z,

- eds. *Ecologically-Based Rodent Management*. ACIAR, Canberra, Australia.
- Makundi RH, Ngowo MV, Massawe AW, Mshuza ME (2010). Community-based rodent pest management in Kilimanjaro region, Tanzania. Consultancy Report. Belgian Technical Cooperation and Ministry of Agriculture, Food Security and Cooperative, Dar es Salaam, Tanzania.
- Makundi RH, Massawe AW (2011). Ecologically based rodent management in Africa: potential and challenges. *Wildlife Research* **38**, 588–95.
- Makundi RH, Massawe AW, Borremans B, Laudisoit A, Katakweba A (2015). We are connected: flea host association networks in the plague outbreak focus in the Rift Valley, northern Tanzania. *Wildlife Research* **42**, 196–206.
- Massawe AW, Makundi RH, Mulungu LS, Katakweba A, Shayo TN (2012). Breeding dynamics of rodent species inhabiting farm–fallow mosaic fields in Central Tanzania. *African Zoology* **47**, 128–37.
- Mead PS (2018). Plague in Madagascar: A tragic opportunity for improving public health. *New England Journal of Medicine* **378**, 106–8.
- Meerburg BG, Singleton GR, Kijlstra A (2009). Rodent-borne diseases and their risks for public health. *Critical Reviews in Microbiology* **35**, 221–70.
- Meheretu Y, Kiros W, Seppe D, Dirk R, Rhodes M, Herwig L (2010). Farmers’ perspectives of rodent damage and management from the highlands of Tigray, Northern Ethiopia. *Crop Protection* **29**, 532–9.
- Mesaki S (2009). Witchcraft and the law in Tanzania. *International Journal of Sociology and Anthropology* **1**, 132–8.
- Monadjem A, Taylor PJ, Denys C, Cotterill FP (2015). *Rodents of Sub-Saharan Africa: A Biogeographic and Taxonomic Synthesis*. Walter de Gruyter, Berlin, Germany.
- Mulungu L, Mrosso F, Katakweba A *et al.* (2015). Farmer’s knowledge, attitude and practice on rodent management in lowland irrigated rice in Central-eastern Tanzania. *International Research Journal of Plant Sciences* **6**, 7–14.
- Mulungu LS, Makundi RH, Massawe AW, Leirs H (2007). Relationship between sampling intensity and precision for estimating damage to maize caused by rodents. *Integrative Zoology* **2**, 131–5.
- Mwanjabe PS, Leirs H (1997). An early warning system for IPM-based rodent control in smallholder farming systems in Tanzania. *Belgian Journal of Zoology* **127**, 49–58.
- Mwanjabe PS, Sirima FB, Lusingu J (2002). Crop losses due to outbreaks of *Mastomys natalensis* (Smith, 1834) Muridae, Rodentia, in the Lindi Region of Tanzania. *International Biodeterioration & Biodegradation* **49**, 133–7.
- Myllymaki A (1989). Denmark-Tanzania rodent control project—final report (unpublished). *Rodent Control Centre*. Morogoro, Tanzania.
- Ngaomei G, Singh EJ (2016). Farmers’ knowledge, attitudes and practices with respect to rodent management in the agricultural ecosystem of Tamenglong district, Manipur, North-East India. *Indian Journal of Applied Research* **2**, 536–40.
- Ninh HN, Palis FG, Aragon CT, Rejesus RM, Singleton GR (2016). Yield and income effects of ecologically-based rodent management in Mekong River Delta, Vietnam. *Asian Journal of Agriculture and Development* **13**, 55–74.
- Palis FG, Singleton GR, Brown PR, Huan NH, Umali C, Nga NTD (2011). Can humans outsmart rodents? Learning to work collectively and strategically. *Wildlife Research* **38**, 568–78.
- Palis FG, Sumalde ZM, Torres CS, Contreras AP, Datar FA (2013). *Impact Pathway Analysis of ACIAR’s Investment in Rodent Control in Vietnam, Lao PDR and Cambodia*. Australian Centre for International Agricultural Research, Canberra, Australia.
- Rootman G, Stevens JB (2016). Enhancing farmers’ organizational and experimentation capacities for soil fertility management in smallholder cropping systems in Vhembe district of Limpopo province in South Africa. *South African Journal of Agricultural Extension* **44**, 120–30.
- Saez AM, Haidara MC, Camara A *et al.* (2018). Rodent control to fight Lassa fever: Evaluation and lessons learned from a 4-year study in Upper Guinea. *PLoS Neglected Tropical Diseases* **12**, e0006829.
- Shabani I, Kimanya ME, Gichuhi PN, Bonsi C, Bovell-Benjamin AC (2015). Maize storage and consumption practices of farmers in Handeni District, Tanzania: Corollaries for mycotoxin contamination. *Open Journal of Preventive Medicine* **5**, 330.
- Singleton GR, Hinds LA, Leirs H, Zhang Z (1999). *Ecologically-Based Rodent Management*. ACIAR, Canberra, Australia.
- Singleton GR, Brown PR, Jacob J, Aplin KP (2007). Unwanted and unintended effects of culling: A case

- for ecologically-based rodent management. *Integrative Zoology* **2**, 247–59.
- South Africa Pest Control Association (SAPCA) (2019). About SAPCA: Objectives of the Organisation [Online]. Available from <http://sapca.org.za/index.php/about/>. Cited January 2020.
- Stayt HA (2018). *The Bavenda*. Routledge, Abingdon, UK.
- Swanepoel LH, Swanepoel CM, Brown PR *et al.* (2017a). Dataset: A systematic review of rodent pest research in Afro-Malagasy small-holder farming systems: Are we asking the right questions? *PLoS ONE* **12**, e0176621. <https://doi.org/10.13140/RG.2.2.24795.77609>.
- Swanepoel LH, Swanepoel CM, Brown PR *et al.* (2017b). A systematic review of rodent pest research in Afro-Malagasy small-holder farming systems: Are we asking the right questions? *PLoS ONE* **12**, e0174554.
- Taylor PJ, Arntzen L, Hayter M, Iles M, Frean J, Belmain S (2008). Understanding and managing sanitary risks due to rodent zoonoses in an African city: beyond the Boston Model. *Integrative Zoology* **3**, 38–50.
- United Republic of Tanzania (URT). (2007). National Sample Census of Agriculture: Volume Vd: Regional Report: Tanga Region [Cited Sep 2019]. Available from <http://www.fao.org/tempref/AG/Reserved/PPLPF/ftpOUT/GLiPHA/DATA/Queue/Working/tanzania/TANGA%20REGION%20REPORT.pdf>.
- Walz JR (2016). Zigua medicine, between mountains and ocean: People, performances, and objects in healing motion. *Histories of Medicine and Healing in the Indian Ocean World*. Springer, Berlin, Germany.
- Williams ST, Maree N, Taylor P, Belmain SR, Keith M, Swanepoel LH (2018). Predation by small mammalian carnivores in rural agro-ecosystems: An undervalued ecosystem service? *Ecosystem Services* **30**, 362–71.
- Ziwa MH, Matee MI, Hang'ombe BM, Lyamuya EF, Kilonzo BS (2013). Plague in Tanzania: An overview. *Tanzania Journal of Health Research* **15**, 252–8.

SUPPLEMENTARY MATERIALS

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Figure S1 Percentage of responses reporting household damage caused by rodents in the (a) Vhembe/Mopani District of South Africa ($n = 60$), (b) Handeni District of Tanzania ($n = 19$) and (c) eastern Madagascar ($n = 777$).

Figure S2 Percentage of responses supporting the future application of different control methods for rodent management in the (a) Vhembe/Mopani District of South Africa ($n = 23$), (b) Handeni District of Tanzania ($n = 6$) and (c) eastern Madagascar ($n = 714$).

Table S1 Respondents knowledge and perceptions of rodent management

Table S2 Summary of actual crop losses by rodents from field experiments across Africa from published studies. Data sourced from (Swanepoel *et al.* 2017)

Cite this article as:

Constant NL, Swanepoel LH, Williams ST *et al.* Comparative assessment on rodent impacts and cultural perceptions of ecologically based rodent management in 3 Afro-Malagasy farming regions. *Integrative Zoology*. 2020;**00**:1-17. <https://doi.org/10.1111/1749-4877.12447>