

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/331319724>

Factors affecting the success of artificial pack formation in an endangered, social carnivore: the African wild dog

Article in *Animal Conservation* · October 2019

DOI: 10.1111/acv.12490

CITATIONS

0

READS

280

22 authors, including:



Courtney Marneweck
University of Mpumalanga

12 PUBLICATIONS 25 CITATIONS

[SEE PROFILE](#)



Harriet T. Davies-Mostert
Endangered Wildlife Trust

62 PUBLICATIONS 885 CITATIONS

[SEE PROFILE](#)



Markus Gusset
Swiss Federal Office for Agriculture

76 PUBLICATIONS 1,491 CITATIONS

[SEE PROFILE](#)



David G. Marneweck
University of Pretoria

20 PUBLICATIONS 40 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:




Leopard status and habitat use drivers in a human-impacted region of southern Mozambique [View project](#)



Kruger carnivores [View project](#)

Factors affecting the success of artificial pack formation in an endangered, social carnivore: the African wild dog

C. Marneweck¹ , P. A. Becker², G. Beverley³, H. T. Davies-Mostert^{3,4}, C. du Plessis³, K. Forssman⁵, J. Graf⁶, M. Gusset⁷, M. Hofmeyr⁸, C. Kelly⁹, P. J. Kilian¹⁰, A. F. J. Marchal^{1,3}, D. G. Marneweck^{3,4}, K. Marnewick^{3,11}, C. Owen¹², G. Postiglione^{13,14}, R. Slotow¹², M. J. Somers^{4,15}, M. Szykman Gunther¹⁶, G. van Dyk¹⁷, B. Whittington-Jones¹⁸ & D. M. Parker^{1,19}

1 School of Biology and Environmental Sciences, University of Mpumalanga, Mbombela, South Africa

2 Washington Department of Fish and Wildlife, Olympia, WA, USA

3 Endangered Wildlife Trust, Johannesburg, South Africa

4 Eugène Marais Chair of Wildlife Management, Mammal Research Institute, University of Pretoria, Pretoria, South Africa

5 Department of Veterinary Tropical Diseases, University of Pretoria, Pretoria, South Africa

6 Association for Water & Rural Development, Hoedspruit, South Africa

7 Swiss Federal Office for Agriculture, Bern, Switzerland

8 Great Plains Conservation, Maun, Botswana

9 WildlifeACT, Cape Town, South Africa

10 Khamab Kalahari Reserve, Bray, South Africa

11 Department of Nature Conservation, Tshwane University of Technology, Pretoria, South Africa

12 School of Life Sciences, University of KwaZulu-Natal, Durban, South Africa

13 Department of Veterinary Medical Sciences, Bologna University, Bologna, Italy

14 Limpopo-Lipadi Private Game and Wilderness Reserve, Tsetsebjwe, Botswana

15 Centre for Invasion Biology, University of Pretoria, Pretoria, South Africa

16 Department of Wildlife, Humboldt State University, Arcata, CA, USA

17 Director of Wildlife, Tswalu Kalahari Reserve, Northern Cape, South Africa

18 Sharjah Environment and Protected Areas Authority, Sharjah, United Arab Emirates

19 Wildlife and Reserve Management Research Group, Department of Zoology and Entomology, Rhodes University, Grahamstown, South Africa

Keywords

group formation; *Lycaon pictus*; pre-release enclosure; separation; social integration; reintroduction; reproduction.

Correspondence

Courtney Marneweck, School of Biology and Environmental Sciences, University of Mpumalanga, Mbombela, South Africa.

Tel: +27 13 002 0223

Email: courtney.marneweck@gmail.com

Editor: Julie Young

Associate Editor: Vincenzo Penteriani

Received 09 July 2018; accepted 16 January 2019

doi:10.1111/acv.12490

Abstract

Social integration is an important factor when reintroducing group-living species, but examples of the formation of social groups before reintroduction are largely lacking. African wild dogs *Lycaon pictus* are endangered, and reintroductions have become a routine part of range expansion in South Africa. Wild dogs form packs that are essential to their reproduction and survival, and artificial pack formation is often required before reintroduction. We examined the proximate (i.e. social integration) and ultimate (i.e. reproduction) success of 43 artificial pack formation attempts in the South African managed metapopulation, between 1995 and 2018. The top (and dominant) model for proximate success included an interaction between total group size and an initial separation fence. Larger groups took longer to integrate, irrespective of initial separation, whereas smaller groups brought together immediately integrated faster than those that were initially separated. The top models for ultimate success included an interaction between the proportion of males and number of days spent in the pre-release enclosure, the total number of days in the enclosure and an interaction between the proportion of captive-sourced individuals and the total number of days in the enclosure. Ultimate success increased when packs spent less time in the enclosure, especially if those packs had a low proportion of males (i.e. female biased) or included >25% captive-sourced individuals. Neither the size of the artificially created pack nor the season in which the pack was released affected ultimate success. The success of social integration and reproductive success of artificially formed packs in this study was higher than for natural pack formations. We provide guidelines for optimizing future artificial pack formation in wild dogs for reintroduction success. Our results serve as an example of the practical importance of social behavior in successfully implementing conservation measures for group-living species.

Introduction

Animal translocations and reintroductions have become widespread and effective conservation tools (Seddon, Armstrong & Maloney, 2007). The main factors to consider when reintroducing animals include the selection of a suitable release site, a suitable founder population and removing the original cause of local extirpation (Fischer & Lindenmayer, 2000). However, when it comes to reintroducing group-living species, there is the additional consideration of social behavior (Somers & Gusset, 2009). Although the IUCN's Species Survival Commission Reintroduction Specialist Group provides universal standards and guidelines for reintroductions (IUCN/SSC, 1998, 2013; Seddon *et al.*, 2007), examples of reintroducing social groups are largely lacking. Furthermore, the importance of social structure is noted, but no advice is provided regarding the creation of viable social groups for reintroductions (Soorae, 2008, 2010, 2011, 2013). It is suggested that holding carnivores before reintroduction helps to break their homing tendencies, and allows acclimatization (Linnell *et al.*, 1997; Miller *et al.*, 1999; Briers-Louw, Verschueren & Leslie, 2019). But, for group-living carnivores, this is also an important time for social integration when artificial social groups are being formed (Gusset, Slotow & Somers, 2006). Despite understanding the importance of social behavior when reintroducing animals, guidelines are unavailable for managers implementing reintroductions of social species. Consequently, many decisions regarding group reintroductions are based on personal knowledge and experience, and the success of such an approach is variable.

African wild dogs *Lycaon pictus* were historically distributed throughout sub-Saharan Africa (Creel & Creel, 1998), but they have since disappeared from much of their historical range and, as a result, are listed as Endangered on the IUCN Red List of Threatened Species (Woodroffe & Sillero-Zubiri, 2012). Wild dogs display a complex social structure, forming packs with a nearly obligate cooperative breeding system (Creel & Creel, 2002). Thus, they rely on being part of a socially integrated pack for survival (Gusset *et al.*, 2006) and, in general, larger packs tend to be better at hunting, raising pups and avoiding threats (Courchamp & Macdonald, 2001; Creel & Creel, 2002; Buettner *et al.*, 2007; Rasmussen *et al.*, 2008; Davies *et al.*, 2016). Within South Africa, small and isolated populations are managed collectively as a metapopulation (Davies-Mostert, Mills & Macdonald, 2009). This involves the reintroduction of wild dogs through the translocation of a single-sex group (usually siblings) to a release site, joining them with an unrelated opposite-sex group and ultimately releasing them together as a newly formed pack (Gusset, 2010). This approach mimics natural dispersal processes (Creel & Creel, 2002) as, most often, the groups used in this study comprised individuals identified as dispersing and searching for mates. The managed metapopulation approach is required in South Africa, where no sufficiently large contiguous patches of suitable habitat remain aside from Kruger National Park, and it has been successful in increasing the wild dog population since its implementation in 1998 (Davies-Mostert, Mills & Macdonald, 2015).

Wild dogs disperse from their natal pack (primary dispersal) during their first year of sexual maturity (i.e. 2–3 years old (McNutt, 1996; Davies-Mostert *et al.*, 2015)) in search of unrelated opposite-sex dispersal groups with which to form a new pack. Although not always the case (Davies-Mostert *et al.*, 2015), males tend to disperse in larger groups than females (average of four individuals compared to an average of two for females (McNutt, 1996; Creel & Creel, 2002)) and, when dispersal groups meet, aggression can occur within the sexes due to an unestablished dominance hierarchy (Creel *et al.*, 1997a). The creation of social bonds during this time is crucial for the formation of a stable and breeding pack (McCreery, 2000). Furthermore, the associations between dispersing same-sex siblings are strong, and the presence of same-sex unrelated individuals may impede pack formation (de Villiers, Richardson & van Jaarsveld, 2003).

During artificial pack formation for reintroduction, wild dogs are kept in a pre-release enclosure to facilitate social integration of the two unrelated, opposite-sex groups, before release into the new area (Gusset *et al.*, 2006). Time spent together in the pre-release enclosure has a positive effect on wild dog survival post-release (Gusset *et al.*, 2010). Additionally, this pre-release enclosure time increases the social integration of wild dogs (Gusset *et al.*, 2008), and wild dogs kept in a pre-release enclosure are more likely to remain together after release (Gusset *et al.*, 2006). However, management records show that wild dogs are kept in different types of pre-release enclosures (1 vs. 2 compartments) and for vastly different lengths of time (WAG-SA, 1998–2018; KZN-WAG, 2004–2018). Pre-release resting group dynamics are an indicator of social integration, where a decreasing resting distance to a separating fence indicates successful pack formation when using a dual-compartment enclosure (Potgieter, O'Riain & Davies-Mostert, 2015). Pre-release enclosure guidelines for wild dogs are available (Potgieter *et al.*, 2012) but, due to resource constraints, many pre-release enclosures in South Africa comprise only one compartment (i.e. no separating fence), and two groups of wild dogs are often joined within the same enclosure from the start. Resource availability (e.g. pre-release enclosure) is, therefore, an important factor to consider when attempting to form a wild dog pack, especially when financial and human resources are limited. It is essential that we understand the dynamics of artificial pack formation under varying scenarios, to aim for optimal success of pack formation, and to reach conservation goals. Historically, captive-sourced wild dogs have also been used in reintroductions, but this was only successful when captive individuals were joined with wild-sourced individuals before release (Frantzen, Ferguson & de Villiers, 2001; Gusset *et al.*, 2010). Although wild dog reintroductions have been relatively successful in South Africa (Gusset *et al.*, 2010; Davies-Mostert *et al.*, 2015), and the importance of social integration is recognized (Gusset *et al.*, 2006), no formal assessment regarding the factors affecting artificial pack formation, and ultimate future reproductive success, has been undertaken.

To address this knowledge gap, we collated data on artificial pack formation attempts with wild dogs in South Africa

over the past 23 years (1995–2018), with the aim of investigating the factors affecting the proximate (i.e. social integration) and ultimate (i.e. reproduction) success of artificial pack formation. The relatedness of same-sex individuals affects social integration (de Villiers *et al.*, 2003; Gusset *et al.*, 2008). Therefore, we predicted that a male-biased sex ratio (to mimic natural dispersal (McNutt, 1996; Creel & Creel, 2002)) and a low proportion of unrelated same-sex individuals, would positively affect proximate success by decreasing the time taken for social integration. Because larger packs are generally more successful than smaller ones (Courchamp & Macdonald, 2001; Buettner *et al.*, 2007; Davies *et al.*, 2016), we also predicted that total group size would affect ultimate success, where larger groups would improve the likelihood of reproduction after release.

Materials and methods

Data collection

We collated data on the protocols followed, and the outcomes of artificial pack formations in wild dogs from the Wild Dog Managed Metapopulation Compendium (Potgieter *et al.*, 2012), and minutes of meetings of the Wild Dog Advisory Group of South Africa (WAG-SA, 1998–2018) and the KwaZulu-Natal Wild Dog Advisory Group (KZN-WAG, 2004–2018). Biannual or quarterly WAG meetings record data on the demographics, translocations and reintroductions of all wild dogs in South Africa. We extracted information recorded from individual reserve reports regarding artificial pack formations. In instances where details were missing, we made follow-up communications with individual managers, researchers and wildlife monitors. We were able to extract sufficient data for 43 separate artificial pack formation attempts over 23 years; that is, from the first attempted artificial pack formation recorded to the most recent (1995–2018; Supplementary material Table S1). Of these events, 37 resulted in the formation and release of a pack. For each successful artificial pack formation attempt, we recorded 10 factors (Table 1) based on current literature and our subsequent predictions. We defined age classes as pup (<1 year), yearling (1–2 years) and adult (>2 years old). If aggression occurred (the occurrence of ≥ 1 encounter causing physical injury with visible wounds), we classified it as either inter-(male-female/female-male) or intra-group (male-male/female-female). We calculated the proportion of unrelated same-sex individuals by the number of unrelated same-sex individuals divided by the total number of same-sex individuals. We defined the reproductive season as mating (February–April), denning (May–July) and non-denning (August–January) as per Van den Berghe *et al.* (2012). We considered social integration successful if a dominant pair formed and mixed-sex resting groups were observed within the pre-release enclosure, and counted the days to integration. A dominant pair is identifiable through mating behavior, over-marking and mate-guarding (Frame *et al.*, 1979; Creel & Creel, 2002). Spatial relationships mirror the strength of social bonds and, therefore, the social integration between the sexes (McCreery, 2000). Based on

Table 1 Factors investigated for their potential effect on the proximate and/or ultimate success (see text for details) of artificial pack formation in African wild dogs

Parameter	Parameter value	
	Mean \pm SE	Range
Total group size	6.72 \pm 0.47	2–16
Proportion of males	0.52 \pm 0.02	0.30–0.80
Proportion of adults	0.92 \pm 0.03	0.40–1.00
Proportion of unrelated males	0.07 \pm 0.03	0.00–0.80
Proportion of unrelated females	0.06 \pm 0.03	0.00–1.00
Proportion of captive-sourced animals	0.15 \pm 0.04	0.00–1.00
Total number of days in pre-enclosure	92.72 \pm 19.91	4–629
	Parameter description	
Initial separation	Separation ($n = 20$), no separation ($n = 23$)	
Aggression	Inter-group ($n = 9$), intra-group ($n = 2$), none ($n = 32$)	
Reproductive season at release	Mating ($n = 8$), denning ($n = 17$), non-denning ($n = 18$)	

McCreery (2000), we defined individuals resting within one adult wild dog length (approximately 90 cm) as resting together, and males and females observed in this manner over the inactive midday period during at least one scan sample, to be resting together. When considering events where wild dogs were initially separated, we used the metric of resting distance to the partitioning fence (Potgieter *et al.*, 2015). Only packs that were considered socially integrated were released and, finally, we considered reproduction successful if a female gave birth in the first breeding season after release.

Statistical analysis

Before analysis, we assessed collinearity between explanatory variables using variance inflation factors and Spearman rank correlation tests (Zar, 2010). For proximate success, we found multicollinearity between the proportion of adults and the proportion of unrelated males, and so removed the proportion of adults. We made this decision as Gusset *et al.* (2008) found no effect of age structure on artificial pack formation, and same-sex unrelated individuals are more likely to affect social integration (de Villiers *et al.*, 2003). For ultimate success, we also removed the proportion of adults, as well as separation fence and aggression, due to collinearity with the proportion of unrelated males.

To investigate the factors affecting the time taken to integrate socially (proximate success), we created 36 candidate generalized linear models with a Poisson distribution. We set the number of days until social integration within the enclosure as the response variable. We set the total group size (continuous), proportion of males (discrete), proportion of captive-sourced individuals (discrete), initial separation (categorical), aggression (categorical), proportion of unrelated males (discrete), proportion of unrelated females (discrete)

and reproductive season at time of joining (categorical), as explanatory variables. Additionally, as social integration is so complex (Gusset *et al.*, 2006), we also included all possible pairwise interactions as explanatory variables. For this analysis, we were only interested in groups that did socially integrate ($n = 37$), but also had to subset the data further to include only the events where the exact number of days taken to integrate socially was known ($n = 23$).

To investigate the factors affecting the ultimate success of artificial pack formation, we created a further 28 candidate generalized linear models with a Binomial distribution. We set denning in the first breeding season after release (yes/no) as the response variable. We then set the total group size, the proportion of males, the proportion of captive-sourced individuals, the proportion of unrelated males, the proportion of unrelated females, the reproductive season at the time of release and total number of days spent in the pre-release enclosure (continuous), as explanatory variables. We also included all possible pairwise interactions as explanatory variables. For this analysis, all packs that were released, and had the potential to breed, were included ($n = 29$; six packs that were released died (either entire pack or all of one sex) before their first breeding season and two packs were contracepted, and were thus not included).

To identify the best model(s), we used model selection based on Akaike's information criterion adjusted for small sample size (AIC_c), where models with ΔAIC_c values ≤ 2 are considered important (Burnham & Anderson, 2002) and then assessed model fit using R^2 . We performed all statistical analyses and created all figures in R (R Core Team, 2008) for Windows, using functions in the packages *lme4* (Bates *et al.*, 2014), *MuMIn* (Bartoń, 2013), *rsq* (Zhang, 2018) and *ggplot2* (Wickham, 2009).

Results

Proximate success

In general, the proximate success of artificial pack formation was high, where 37 of 40 (93%) groups socially integrated. The total group size, the presence of an initial separation fence, and the interaction between total group size and separation fence, affected the time to form artificial packs successfully (Table 2). When dealing with smaller group sizes, social integration occurred more quickly when there was no

initial separation fence; however, with larger group sizes, social integration occurred more slowly regardless of the presence of an initial separation fence (Fig. 1). Of the three events that did not result in social integration, it was the result of aggression, and one case of all females escaping from the pre-release enclosure before being united with the males.

Ultimate success

The ultimate success of artificial pack formation was also high, with 26 of 29 (90%) packs breeding in the first season post-release. Three models were highlighted as important (Table 3). First, the proportion of males, the total number of days in the pre-release enclosure, and the interaction between the two affected ultimate success. Female-biased packs (i.e. male proportion <0.5) had a higher probability of reproduction with decreased time in the enclosure (Fig. 2). Male-biased packs had a higher probability of reproduction with increased time in the enclosure; however, as the confidence interval surrounding male-biased packs is large, it is unlikely that this has a strong effect on reproductive success (Fig. 2, Table 3). Second, the total number of days in the enclosure affected ultimate success, where packs were more likely to reproduce with reduced time in the enclosure (Fig. 3). Third, the proportion of captive-sourced individuals, the total number of days in the enclosure and the interaction between the two affected ultimate success. Packs containing a higher proportion of captive-sourced individuals (i.e. >0.25) had a higher probability of reproduction with decreased time in the enclosure (Fig. 4). Packs containing no or a small proportion of captive-sourced individuals (i.e. <0.25) had a higher probability of reproduction with increased time in the enclosure. However, as the confidence interval is large, it is unlikely that this has a strong effect on reproductive success (Fig. 4, Table 3). Size of the newly formed packs and the season of release of the newly formed packs had little effect on the probability of reproduction in the first year post-release.

Discussion

Holding animals in an enclosure before release may increase reintroduction success (Fischer & Lindenmayer, 2000; Devineau *et al.*, 2011; Briers-Louw *et al.*, 2019). For carnivores, this allows animals to familiarize themselves with their new

Table 2 The five highest-ranking candidate models used to investigate factors affecting the proximate success of artificial pack formation in African wild dogs. The top model is indicated in bold, where $\Delta AIC_c \leq 2$. For a full table of all 36 candidate models, see supplementary material Table S2

Rank	Model	Coeff ^a	se ^a	ΔAIC_c	Weight	R^2
1	Group size + separation + groups size * separation	-0.21	0.05	0.00	>0.99	0.60
2	Prop unrelated males + separation + prop unrelated males * separation	-0.01	0.18	11.03	<0.01	0.50
3	Group size + season joined + group size * season joined	0.00	0.01	15.43	<0.01	0.60
4	Prop captive + separation + prop captive * separation	0.00	0.00	24.81	<0.01	0.38
5	Separation	2.85	0.52	29.43	<0.01	0.31

^aRefers to model interaction term only.




Figure 1 The number of days until social integration in artificially formed African wild dog packs based on total group size and the presence of an initial separation fence. The shaded region represents the 95% confidence interval.

environment and to break homing tendencies (Linnell *et al.*, 1997; Miller *et al.*, 1999). For group-living carnivores, it is also an important time for social integration in artificially formed groups (Hunter *et al.*, 2007; Gusset *et al.*, 2008). However, the optimal structure and integration of social groups used in the artificial formation of wild dog packs were not fully understood. Here, we defined two levels of success for artificial pack formation in wild dogs; namely pre-release social integration and post-release reproduction. With 37 of 40 events resulting in successful social integration, our results show that artificial pack formation in wild dogs has, in general, been very successful (93%). Gusset *et al.* (2009) calculated a natural pack formation probability of 64%, indicating that artificial pack formation in the managed metapopulation is greater than in free-roaming populations. Our results also show high ultimate success (90%), with 26 of 29 artificially created packs denning in the first breeding season after release, regardless of the season in which they were released. Gusset *et al.* (2009) calculated that the probability of a litter being produced by a newly formed pack (i.e. in the first breeding season after formation) was 33%, increasing to 66% for all subsequent years,

indicating that ultimate success of artificially formed packs in the present study was also very high.

Social integration has been highlighted as essential for wild dog reintroduction success (Gusset *et al.*, 2010), and wild dogs kept in a pre-release enclosure are more likely to remain together after release (Gusset *et al.*, 2006). Time spent in a pre-release enclosure increases social integration of wild dogs (Gusset *et al.*, 2008), and has also been found to facilitate the social integration of unrelated lions *Panthera leo* into viable social groups that go on to reproduce (Hunter *et al.*, 2007). Time spent together in a pre-release enclosure also has positive effects on wild dog survival upon release (Gusset *et al.*, 2010), and Gusset *et al.* (2006) suggested a pre-release enclosure period of up to 91 days for wild dogs (based on three case studies). Our results show that larger groups took longer to integrate than smaller ones, and therefore this factor should be considered when planning artificial pack formations. In addition, our analyses revealed that social integration was delayed when an initial separation fence was used with smaller groups but the presence of a separation fence did not affect larger groups. Gusset *et al.* (2006) report that there were insufficient data to conclude whether a separating fence facilitates social integration in wild dogs. Although it has been suggested that dispersal groups naturally come together over several days (Frame *et al.*, 1979), Hofmeyr (2001) suggested that opposite-sex groups should be introduced into the same enclosure immediately. Our results support the suggestion of joining opposite-sex groups immediately. It is suggested that the intrinsically rewarding physical properties of social behaviors, such as grooming and resting together, facilitates pair-bonding (Ágmo *et al.*, 2012). Such an approach would support the idea that groups should be placed together from the start to facilitate social bonding and the formation of a dominant pair. As the establishment of dominance hierarchies is extremely complex (Chase *et al.*, 2002), we propose that larger groups simply require more time to form a dominant pair and construct a hierarchy with more competing individuals.

With regard to ultimate success, packs were less likely to reproduce with increased time spent in the pre-release enclosure, suggesting there may be negative consequences of continuing to hold newly formed packs in captivity after signs of cohesive social integration are observed. Confinement-specific stressors, such as restricted movement, reduced retreat space and forced proximity to humans, can adversely affect animals in captivity (Morgan & Tromborg, 2007). Although temporary,

Table 3 The five highest-ranking candidate models used to investigate factors affecting the ultimate success of artificial pack formation in African wild dogs. The top models are indicated in bold, where $\Delta\text{AIC}_c \leq 2$. For a full table of all 28 candidate models, see supplementary material Table S3

Rank	Model	Coeff ^a	SE ^a	ΔAIC_c	Weight	R^2
1	Total days + prop males + total days * prop males	0.25	0.63	0.00	0.22	0.44
2	Total days	0.44	652.10	0.79	0.15	0.20
3	Prop captive + total days + prop captive * total days	-0.03	84.72	1.93	0.08	0.35
4	Season + total days + season * total days	-0.54	652.10	2.24	0.07	0.66
5	Prop males	-11.35	28.40	2.53	0.06	0.03

^aRefers to model interaction term only.




Figure 2 The proportion of artificially formed African wild dog packs reproducing based on the proportion of males and the total number of days in the enclosure pre-release. The shaded region represents the 95% confidence interval.




Figure 4 The proportion of artificially formed African wild dog packs reproducing based on the proportion of captive-sourced individuals and the total number of days in the enclosure pre-release. The shaded region represents the 95% confidence interval.




Figure 3 The proportion of artificially formed African wild dog packs reproducing based on the total number of days in the enclosure pre-release. The shaded region represents the 95% confidence interval.

the pre-release enclosure period is a form of captivity, where the aforementioned stressors exist. Prolonged confinement of wild animals has been shown to cause stress; for example, translocated wild eastern bettongs *Bettongia gaimardi* have higher faecal glucocorticoid metabolite concentrations when their release is delayed (Batson *et al.*, 2017). Ultimately, prolonged high levels of glucocorticoids (i.e. chronic stress) can be detrimental to reproduction (Pottinger, 1999). In contrast to the hypothesis that chronic stress causes reproductive suppression,

dominant wild dogs have higher basal corticosterone levels than subordinates (Creel *et al.*, 1997a; Creel, 2001), meaning that reproductive suppression in wild dogs is not driven by prolonged glucocorticoid elevation (i.e. chronic stress). However, this evidence relates only to free-roaming and stable packs. In addition to confinement stress, unfamiliar wild dogs brought together in a pre-release enclosure must also construct a dominance hierarchy. During this time, aggression can be severe (Creel *et al.*, 1997a) and individuals have limited space to avoid this aggression. Although capture and handling do not cause chronic stress in wild dogs (Creel, Creel & Monfort, 1997b), no physiological data exist regarding the impact of confinement on free-roaming wild dogs, especially groups of unfamiliar wild dogs brought together for artificial pack formation. We suggest that prolonged confinement, coupled with the joining of unknown groups, may cause chronic stress to such a level that it has a subsequent detrimental effect on reproduction. Nevertheless, such research is clearly needed for free-roaming wild dogs in confinement of different durations.

We also show that the pattern of decreased likelihood of reproduction with increased time in the enclosure is highlighted for female-biased packs. The hippocampus in females responds to stress differently to that in males, where females with high levels of oestrogen are especially sensitive (Shors, Chua & Falduto, 2001). Female tigers *Panthera tigris* are more sensitive to the stress of anthropogenic disturbance than males, resulting in higher faecal glucocorticoid metabolite concentrations (Bhattacharjee *et al.*, 2015). Also, captive female wild dogs have higher faecal cortisol metabolite concentrations than free-roaming females (Van der Weyde, Martin & Paris, 2016). If a pack contains a high proportion of females, then there are more females competing for dominance and aggression can be severe when a dominance hierarchy has not yet been established (Creel *et al.*, 1997a;

Creel, 2001). As such, the construction of a dominance hierarchy with more competing individuals may cause higher and prolonged levels of stress in these females. This, compounded with a long confinement period, may amount to chronic stress, which may have a subsequent detrimental effect on reproduction upon release. When packs contain a low proportion of females, there may be less stress in defining dominance amongst females and, thus, with confinement as the only stressor, it may be less likely to have such a detrimental impact on reproduction post-release. We suggest that confinement, coupled with female wild dogs potentially being more susceptible to chronic stress, can lead to female-biased packs being less successful post-release after a long confinement period.

Packs containing only captive-sourced individuals are not successful when it comes to reintroductions (Frantzen *et al.*, 2001). This was considered in subsequent reintroductions, where captive-sourced individuals were only used when joined with wild individuals when forming new packs (Gusset *et al.*, 2010). Some captive-sourced individuals were used in this study, and our results show that, for packs containing more than 25% captive-sourced individuals, their likelihood of reproduction significantly reduces with time spent in the enclosure. We suggest that the low ability of captive-sourced wild dogs to adapt to their environment (e.g. avoid lions, disease immunity) and contribute to the pack (e.g. communal hunting), coupled with the detrimental effect of prolonged confinement on the wild-sourced individuals with which they are joined, leads to packs being less likely to reproduce in their first season post-release.

Contrary to our expectations, total group size did not have a significant effect on ultimate success, at least not in the first breeding season following release. In general, larger packs are more successful hunters (Creel & Creel, 2002; Rasmussen *et al.*, 2008), have higher pup and juvenile survival (Buettner *et al.*, 2007; Davies *et al.*, 2016), select better den sites (Davies *et al.*, 2016), and are better at avoiding threats (Courchamp & Macdonald, 2001). Gusset & Macdonald (2010) reported a mean pack size for the managed metapopulation of 6.2 ± 3.4 (range = 2–17), similar to the present study (6.6 ± 0.5 , range = 2–16). This is lower than in larger, unmanaged populations (Creel, Mills & McNutt, 2004). However, metapopulation reserves tend to have low densities of competitors and high availability of suitable prey (Davies-Mostert *et al.*, 2009). Consequently, in line with previous research (Gusset *et al.*, 2008; Somers *et al.*, 2008; Gusset & Macdonald, 2010), our results suggest that pack size may not strongly affect the ultimate success of wild dog packs within the managed metapopulation, at least shortly after release.

The factors highlighted as important were not able to strongly predict the ultimate success of packs in this study (refer to model weights in Table 3). This is likely due to large variation in external variables at each reserve and each year, such as lion density, prey density and wild dog density. Unfortunately, these data were unavailable for our study, but our results support the notion that external factors are likely to be important when it comes to the ultimate success of reintroductions (i.e. reproduction). Another caveat to note is that

although the lineages of the wild dogs involved in our study were known, and relatedness was considered when choosing opposite-sex groups for artificial pack formation, genetic data were not available. We acknowledge the potential effect that genetic diversity and mate choice may have on both proximate and ultimate success, as wild dogs do show inbreeding avoidance (Becker *et al.*, 2012). However, recent genetic evidence pertaining to the last 10 years within the managed metapopulation suggests limited inbreeding and high levels of heterozygosity (Tensen *et al.* in press). We must also acknowledge the small samples sizes retained in our analyses, meaning we must be cautious in our conclusions.

As the first comprehensive study of artificial group formation in a social carnivore, our results provide a robust evaluation of current methods used for wild dog pack formation and provide practical guidelines for optimizing the formation of future packs. Currently, only ~600 wild dogs remain in South Africa, ~250 (42%) of which reside in the managed metapopulation (WAG-SA, 1998–2018). As such, optimizing actions and outcomes within the conservation strategy of the South African managed metapopulation is of high priority. In addition to providing important information for the reintroduction of wild dogs, our results show that it is possible to socially integrate unrelated individuals of social animals, which has applications for the reintroduction and restoration of many species. Within the Carnivora, this knowledge could be applied to artificial pack formation of endangered dholes *Cuon alpinus* and grey wolves *Canis lupus*. Outside of the Canidae, this could also be applied to creating coalitions of unrelated male cheetahs *Acinonyx jubatus* for reintroduction (Marnewick *et al.*, 2009; Boast *et al.*, 2018), as well as adding to the knowledge of creating viable lion prides (Slotow & Hunter, 2009). Our results can also provide a platform for the pair bonding or group formation of primates. In conclusion, our results highlight the necessity to incorporate species' social behavior into reintroductions, and the subsequent impact on conservation success that it can produce. Ultimately, these results can be used to positively impact wild dog populations, range expansion and recovery, as well as to stimulate thinking for the reintroduction of other threatened social species in fragmented landscapes, where dispersal corridors are becoming limited due to extensive anthropogenic pressure.

Implications for management

Reducing the number of pre-release enclosure days is vital at sites where resource restrictions mean that enclosures are multi-purpose, and the cost of provisioning wild dogs is high. For example, many protected areas in South Africa use pre-release enclosures for multiple species reintroductions, or for holding animals for game auctions, or temporary veterinary care. In addition to reducing resource costs, limiting the number of days in the pre-release enclosure also increases the probability of pack ultimate success. So, effort must be made to reduce the time packs spend in pre-release enclosures. Our results indicate that a separation fence is not necessary for artificial pack formation in wild dogs, as it delays

social integration of small groups and has no effect on large groups. A preliminary study has investigated the use of odour familiarity to increase social integration when two groups of unrelated, opposite-sex wild dogs are joined in one enclosure (Marchal *et al.* unpublished data). The method involves body rubbing of all individuals while still sedated, before waking together as one group, and preliminary results show rapid social cohesion, where eight out of nine events using this method resulted in rapid social integration (in one instance the females escaped from their enclosure before social integration could occur). We highlight the necessity for pre-release enclosure observations of inter-group social dynamics to assess social integration.

To optimize the process of artificial pack formation, we suggest that smaller groups be given at least 5 days to integrate socially if in one enclosure, and at least 10 days if in a double-compartmented enclosure. For large groups, we suggest that the wild dogs be given at least 14 days to integrate socially, regardless of enclosure type (see decision framework in Supplementary material Figure S1). In order to reduce the stress experience within the pre-release enclosure, we advise that, in accordance with WAG-SA pre-release enclosure guidelines (Potgieter *et al.*, 2012), stressful events (such as feeding days and enclosure checks) be kept regular, as predictable stressful events are less stressful than those occurring irregularly (Bassett & Buchanan-Smith, 2007). We also suggest that when captive-sourced individuals are to be used, they comprise <25% of the pack. Finally, we caution the long-term placement of packs in enclosures, as it is detrimental to their ultimate success.

Conflicts of Interest

The authors declare no conflicts of interest.

References

- Ågmo, A., Smith, A.S., Birnie, A.K. & French, J.A. (2012). Behavioral characteristics of pair bonding in the black tufted-ear marmoset (*Callithrix penicillata*). *Behaviour* **149**, 407–440.
- Bartón, K. (2013). *MuMIn: multi-model inference. R package version 1.9.13*. Vienna, Austria: The Comprehensive R Archive Network (CRAN).
- Bassett, L., Buchanan-Smith, H. M. (2007). Effects of predictability on the welfare of captive animals. *Appl. Anim. Behav. Sci.* **102**, 223–245.
- Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R.H.B., Singmann, H., Dai, B., Grothendieck, G. & Green, P. (2014). *Package 'lme4'*. Vienna: R Foundation for Statistical Computing.
- Batson, W.G., Gordon, I.J., Fletcher, D.B., Portas, T.J. & Manning, A.D. (2017). The effect of pre-release captivity on the stress physiology of a reintroduced population of wild eastern bettongs. *J. Zool.* **303**, 311–319.
- Becker, P.A., Miller, P.S., Gunther, M.S., Somers, M.J., Wildt, D.E. & Maldonado, J.E. (2012). Inbreeding avoidance influences the viability of reintroduced populations of African wild dogs (*Lycaon pictus*). *PLoS ONE* **7**, e37181.
- Bhattacharjee, S., Kumar, V., Chandrasekhar, M., Malviya, M., Ganswindt, A., Ramesh, K., Sankar, K. & Umaphathy, G. (2015). Glucocorticoid stress responses of reintroduced tigers in relation to anthropogenic disturbance in Sariska Tiger Reserve in India. *PLoS ONE* **10**, e0127626.
- Boast, L. K., Chelysheva, E. V., dervan Merwe, V., Schmidt-Kuntzel, A., Walker, E. H., Cilliers, D., Gusset, M. and Marker, L. (2018). Cheetah translocation and reintroduction programs: past, present, and future. In *Cheetahs: biology and conservation: 275–290*. Marker, L., Boast, L. K. and Schmidt-Kuntzel, A. (Eds). London, UK: Academic Press.
- Briers-Louw, W.D., Verschuuren, S. & Leslie, A.J. (2019). Big cats return to Majete Wildlife Reserve, Malawi: evaluating reintroduction success. *Afr. J. Wildl. Res.* **49**, 1–17.
- Buettner, U.K., Davies-Mostert, H.T., du Toit, J.T. & Mills, M.G.L. (2007). Factors affecting juvenile survival in African wild dogs (*Lycaon pictus*) in Kruger National Park, South Africa. *J. Zool.* **272**, 10–19.
- Burnham, K.P. & Anderson, D.R. (2002). *Model selection and multimodel inference: a practical information-theoretic approach*. 2nd edn. New York, NY: Springer Science & Business Media.
- Chase, I.D., Tovey, C., Spangler-Martin, D. & Manfredonia, M. (2002). Individual differences versus social dynamics in the formation of animal dominance hierarchies. *Proc. Natl Acad. Sci.* **99**, 5744–5749.
- Courchamp, F. & Macdonald, D.W. (2001). Crucial importance of pack size in the African wild dog *Lycaon pictus*. *Anim. Conserv.* **4**, 169–174.
- Creel, S. (2001). Social dominance and stress hormones. *Trends Ecol. Evol.* **16**, 491–497.
- Creel, S. & Creel, N.M. (1998). Six ecological factors that may limit African wild dogs, *Lycaon pictus*. *Anim. Conserv.* **1**, 1–9.
- Creel, S. & Creel, N.M. (2002). *The African wild dog: behavior, ecology, and conservation*. Princeton, NJ: Princeton University Press.
- Creel, S., Creel, N.M., Mills, M.G.L. & Monfort, S.L. (1997a). Rank and reproduction in cooperatively breeding African wild dogs: behavioral and endocrine correlates. *Behav. Ecol.* **8**, 298–306.
- Creel, S., Creel, N.M. & Monfort, S.L. (1997b). Radiocollaring and stress hormones in African wild dogs. *Conserv. Biol.* **11**, 544–548.
- Creel, S., Mills, M.G.L. & McNutt, J.W. (2004). Demography and population dynamics of African wild dogs in three critical populations. In *Biology and conservation of wild Canids: 337–350*. Macdonald, D.W. & Sillero-Zubiri, C. (Eds). Oxford, UK: Oxford University Press.

- Davies, A.B., Marneweck, D.G., Druce, D.J. & Asner, G.P. (2016). Den site selection, pack composition, and reproductive success in endangered African wild dogs. *Behav. Ecol.* **27**, 1869–1879.
- Davies-Mostert, H.T., Mills, M.G.L. & Macdonald, D.W. (2009). A critical assessment of South Africa's managed metapopulation recovery strategy for African wild dogs and its value as a template for large carnivore conservation elsewhere. In *Reintroduction of top-order predators*: 10–42. Hayward, M.W. & Somers, M.J. (Eds). Oxford, UK: Blackwell Publishing Ltd.
- Davies-Mostert, H.T., Mills, M.G.L. & Macdonald, D.W. (2015). The demography and dynamics of an expanding, managed African wild dog metapopulation. *Afr. J. Wildl. Res.* **45**, 258–273.
- Devineau, O., Shenk, T.M., Doherty, P.F. Jr, White, G.C. & Kahn, R.H. (2011). Assessing release protocols for Canada lynx reintroduction in Colorado. *J. Wildl. Manag.* **75**, 623–630.
- Fischer, J. & Lindenmayer, D.B. (2000). An assessment of the published results of animal relocations. *Biol. Cons.* **96**, 1–11.
- Frame, L.H., Malcolm, J.R., Frame, G.W. & Van Lawick, H. (1979). Social organization of African wild dogs (*Lycaon pictus*) on the Serengeti plains, Tanzania 1967–1978. *Zeitschrift für Tierpsychologie* **50**, 225–249.
- Frantzen, M.A.J., Ferguson, J.W.H. & de Villiers, M.S. (2001). The conservation role of captive African wild dogs (*Lycaon pictus*). *Biol. Cons.* **100**, 253–260.
- Gusset, M. (2010). The re-introduction of African wild dogs in South Africa. In *Global Re-introduction Perspectives: additional case-studies from around the globe*: 220–224. Soorae, P.S. (Ed). IUCN/SSC Re-introduction Specialist Group: Abu Dhabi, UAE.
- Gusset, M. & Macdonald, D.W. (2010). Group size effects in cooperatively breeding African wild dogs. *Anim. Behav.* **79**, 425–428.
- Gusset, M., Slotow, R. & Somers, M.J. (2006). Divided we fail: the importance of social integration for the re-introduction of endangered African wild dogs (*Lycaon pictus*). *J. Zool.* **270**, 502–511.
- Gusset, M., Ryan, S.J., Hofmeyr, M., Van Dyk, G., Davies-Mostert, H.T., Graf, J.A., Owen, C., Szykman, M., Macdonald, D.W., Monfort, S.L., Wildt, D.E., Maddock, A.H., Mills, M.G.L., Slotow, R. & Somers, M.J. (2008). Efforts going to the dogs? Evaluating attempts to re-introduce endangered wild dogs in South Africa. *J. Appl. Ecol.* **45**, 100–108.
- Gusset, M., Jakoby, O., Muller, M.S., Somers, M.J., Slotow, R. & Grimm, V. (2009). Dogs on the catwalk: modelling re-introduction and translocation of endangered wild dogs in South Africa. *Biol. Cons.* **142**, 2774–2781.
- Gusset, M., Stewart, G.B., Bowler, D.E. & Pullin, A.S. (2010). Wild dog reintroductions in South Africa: a systematic review and cross-validation of an endangered species recovery programme. *J. Nat. Conserv.* **18**, 230–234.
- Hofmeyr, M. (2001). Introduction of the African wild dog (*Lycaon pictus*) to game reserves. In *Proceedings of a symposium on the re-introduction of larger mammals to game reserves*. Faculty of Veterinary Sciences report, University of Pretoria, Pretoria, South Africa.
- Hunter, L.T.B., Pretorius, K., Carlisle, L.C., Rickelton, M., Walker, C., Slotow, R. & Skinner, J.D. (2007). Restoring lions *Panthera leo* to northern KwaZulu-Natal, South Africa: short-term biological and technical success but equivocal long-term conservation. *Oryx* **41**, 196–204.
- IUCN/SSC (1998). Guidelines for re-introductions. IUCN Species Survival Commission, Gland, Switzerland.
- IUCN/SSC (2013). *Guidelines for reintroductions and other conservation translocations*. Gland, Switzerland: IUCN Species Survival Commission.
- KZN-WAG. (2004–2018). *KwaZulu Natal wild dog advisory group*. Unpublished meeting minutes: KwaZulu-Natal, South Africa.
- Linnell, J.D.C., Aanes, R., Swenson, J.E., Odden, J. & Smith, M.E. (1997). Translocation of carnivores as a new method for managing problem animals: a review. *Biodivers. Conserv.* **6**, 1245–1257.
- Marnewick, K., Cilliers, D., Hayward, M. & Somers, M. (2009). Survival of cheetahs relocated from ranchlands to fenced protected areas. In *The re-introduction of top order predators*: 282–306. Hayward, M. & Somers, M. (Eds). Oxford, UK: Blackwell Publishing.
- McCreery, E.K. (2000). Spatial relationships as an indicator of successful pack formation in free-ranging African wild dogs. *Behaviour* **137**, 579–590.
- McNutt, J.W. (1996). Sex-biased dispersal in African wild dogs, *Lycaon pictus*. *Anim. Behav.* **52**, 1067–1077.
- Miller, B., Ralls, K., Reading, R.P., Scott, J.M. & Estes, J. (1999). Biological and technical considerations of carnivore translocation: a review. *Anim. Conserv.* **2**, 59–68.
- Morgan, K.N. & Tromborg, C.T. (2007). Sources of stress in captivity. *Appl. Anim. Behav. Sci.* **102**, 262–302.
- Potgieter, K.R., Whittington-Jones, B., Gusset, M., Mills, M.G.L. & Davies-Mostert, H.T. (2012). *The South African wild dog metapopulation compendium*. Unpublished report: Johannesburg, South Africa.
- Potgieter, K.R., O'Riain, M.J. & Davies-Mostert, H.T. (2015). Behavioural cues can be used to predict the outcome of artificial pack formation in African wild dogs (*Lycaon pictus*). *Afr. J. Wildl. Res.* **45**, 215–222.
- Pottinger, T. G. (1999). The impact of stress on animal reproductive activities. In *Stress physiology in animals*: 130–177. Baum, P. H. M. (Ed). Oxford, UK: Taylor & Francis.
- R Core Team (2008). *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Rasmussen, G.S., Gusset, M., Courchamp, F. & Macdonald, D.W. (2008). Achilles' heel of sociality revealed by energetic poverty trap in cursorial hunters. *Am. Nat.* **172**, 508–518.

- Seddon, P.J., Armstrong, D.P. & Maloney, R.F. (2007). Developing the science of reintroduction biology. *Conserv. Biol.* **21**, 303–312.
- Shors, T.J., Chua, C. & Falduto, J. (2001). Sex differences and opposite effects of stress on dendritic spine density in the male versus female hippocampus. *J. Neurosci.* **21**, 6292–6297.
- Slotow, R. and Hunter, L. T. B. (2009). Reintroduction decisions taken at the incorrect social scale devalue their conservation contribution: the African lion in South Africa. In *Reintroduction of top-order predators*. Hayward, M. W. and Somers, M. (Eds). Oxford, UK: Blackwell Publishing Ltd.
- Somers, M.J. & Gusset, M. (2009). The role of social behaviour in carnivore reintroductions. In *Reintroduction of top-order predators: 270–281*. Hayward, M.W. & Somers, M.J. (Eds). Oxford, UK: Blackwell Publishing Ltd.
- Somers, M.J., Graf, J.A., Szykman, M., Slotow, R. & Gusset, M. (2008). Dynamics of a small re-introduced population of wild dogs over 25 years: allee effects and the implications of sociality for endangered species' recovery. *Oecologia* **158**, 239–247.
- Soorae, P. S. (2008). Global re-introduction perspectives: re-introduction case-studies from around the globe. IUCN/SSC re-introduction specialist group: 2008. http://www.iucnsscrg.org/index.php?option=com_content&view=article&id=192&Itemid=587 (Accessed 06 November 2018).
- Soorae, P. S. (2010). Global re-introduction perspectives: Additional case-studies from around the globe. IUCN/SSC re-introduction specialist group 2010. http://www.iucnsscrg.org/index.php?option=com_content&view=article&id=192&Itemid=587 (Accessed 06 November 2018)
- Soorae, P. S. (2011). Global re-introduction perspectives: 2011. More case studies from around the globe. IUCN/SSC re-introduction specialist group and 2011. http://www.iucnsscrg.org/index.php?option=com_content&view=article&id=192&Itemid=587 (Accessed 06 November 2018)
- Soorae, P. S. (2013). Global re-introduction perspectives: 2013. Further case studies from around the globe. IUCN/SSC re-introduction specialist group: 2013. http://www.iucnsscrg.org/index.php?option=com_content&view=article&id=192&Itemid=587 (Accessed 06 November 2018).
- Van den Berghe, F., Paris, D.B., Van Soom, A., Rijsselaere, T., Van der Weyde, L., Bertschinger, H.J. & Paris, M.C. (2012). Reproduction in the endangered African wild dog: basic physiology, reproductive suppression and possible benefits of artificial insemination. *Anim. Reprod. Sci.* **133**, 1–9.
- Tensen, L., Van Vuuren, B., du Plessis, C. & Marneweck, D. G. (2019). African wild dogs: genetic viability of translocated populations across South Africa. *Biol. Conserv.* in press.
- Van der Weyde, L.K., Martin, G.B. & Paris, M.C. (2016). Monitoring stress in captive and free-ranging African wild dogs (*Lycaon pictus*) using faecal glucocorticoid metabolites. *Gen. Comp. Endocrinol.* **226**, 50–55.
- de Villiers, M.S., Richardson, P.R.K. & van Jaarsveld, A.S. (2003). Patterns of coalition formation and spatial association in a social carnivore, the African wild dog (*Lycaon pictus*). *J. Zool. (London)* **260**, 377–389.
- WAG-SA (1998–2018). *Wild dog advisory group of South Africa*. Unpublished meeting minutes: Johannesburg, South Africa.
- Wickham, H. (2009). *ggplot2: elegant graphics for data analysis*. New York, NY: Springer-Verlag.
- Woodroffe, R. & Sillero-Zubiri, C. (2012). *Lycaon pictus*. The IUCN red list of threatened species. e.T12436A16711116.
- Zar, J.H. (2010). *Biostatistical analysis*, 4th edn. New Jersey, USA: Prentice Hall.
- Zhang, D. (2018). *rsq: R-squared and related measures*. R package version 1.0.1. <https://cran.rproject.org/package=rsq>

Supporting information

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Table S1. Location and year of each attempted artificial pack formation in African wild dogs included in the study.

Table S2. The 36 candidate models used to investigate factors affecting proximate success of artificial pack formation in African wild dogs. The top model is indicated in bold, where $\Delta\text{AIC}_c \leq 2$.

Table S3. The 28 candidate models used to investigate factors affecting ultimate success of artificial pack formation in African wild dogs. The top models are indicated in bold, where $\Delta\text{AIC}_c \leq 2$.

Figure S1. Decision-making framework for optimizing the pre-release enclosure social integration process for African wild dog reintroductions.