

Convergent Evolution and Structural Adaptation to the Deep Ocean in the Protein-Folding Chaperonin CCT α

Alexandra A.-T. Weber ^{1,2,3,*}, Andrew F. Hugall¹, and Timothy D. O'Hara¹

¹Sciences, Museums Victoria, Melbourne, Victoria, Australia

²Centre de Bretagne, REM/EEP, Ifremer, Laboratoire Environnement Profond, Plouzané, France

³Zoological Institute, University of Basel, Switzerland

*Corresponding author: E-mail: aweber@museum.vic.gov.au.

Accepted: 5 August 2020

Data deposition: This project has been deposited at NCBI under the Bioproject accession PRJNA311384 and dryad packages: <https://doi.org/10.5061/dryad.db339> and <https://doi.org/10.5061/dryad.rb334>.

Abstract

The deep ocean is the largest biome on Earth and yet it is among the least studied environments of our planet. Life at great depths requires several specific adaptations; however, their molecular mechanisms remain understudied. We examined patterns of positive selection in 416 genes from four brittle star (Ophiuroidea) families displaying replicated events of deep-sea colonization (288 individuals from 216 species). We found consistent signatures of molecular convergence in functions related to protein biogenesis, including protein folding and translation. Five genes were recurrently positively selected, including chaperonin-containing TCP-1 subunit α (CCT α), which is essential for protein folding. Molecular convergence was detected at the functional and gene levels but not at the amino-acid level. Pressure-adapted proteins are expected to display higher stability to counteract the effects of denaturation. We thus examined *in silico* local protein stability of CCT α across the ophiuroid tree of life (967 individuals from 725 species) in a phylogenetically corrected context and found that deep-sea-adapted proteins display higher stability within and next to the substrate-binding region, which was confirmed by *in silico* global protein stability analyses. This suggests that CCT α displays not only structural but also functional adaptations to deep-water conditions. The CCT complex is involved in the folding of \sim 10% of newly synthesized proteins and has previously been categorized as a “cold-shock” protein in numerous eukaryotes. We thus propose that adaptation mechanisms to cold and deep-sea environments may be linked and highlight that efficient protein biogenesis, including protein folding and translation, is a key metabolic deep-sea adaptation.

Key words: Echinodermata, TCP-1, protein folding, positive selection, protein stability, pressure adaptation.

Significance

The deep ocean is a vast yet understudied ecosystem. Specifically, very little is known about how deep-sea animals are adapted to the inhospitable deep-sea conditions, such as freezing temperature and crushing pressure. We examined patterns of molecular evolution in >400 genes in hundreds of species from four brittle star families displaying independent events of deep-sea colonization. We found that functions related to the production of proteins were particularly important for deep-sea adaptation. Particularly, the gene CCT α , involved in protein folding, displayed functional and structural adaptations to life at great depths, highlighting its central role in adaptation to the deep ocean.

Introduction

The deep ocean (>200 m) covers approximately two-thirds of the global sea floor area, yet it is among the least studied

environments of our planet in terms of biodiversity, habitats, and ecosystem functioning (Ramirez-Llodra et al. 2010). It harbors specific environmental conditions such as high

© The Author(s) 2020. Published by Oxford University Press on behalf of the Society for Molecular Biology and Evolution.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

hydrostatic pressure (on average ~400 atm), low temperatures (0–4 °C), absence of light, and scarcity of food. Life at great depths requires multiple metabolic adaptations resulting in a physiological bottleneck (Gross and Jaenicke 1994) limiting the vertical distribution of species (Somero 1992; Brown and Thatje 2014). It has also been proposed that high pressure may drive speciation events (Brown and Thatje 2014; Gaither et al. 2016). Numerous cellular processes such as enzymatic processes, protein folding, assembly of multi-subunit proteins, and lipoprotein membranes are impacted by high pressure and low temperature (Jaenicke 1991; Somero 1992; Carney 2005; Pradillon and Gaill 2007). More specifically, cellular mechanisms of deep-sea adaptation have been recently reviewed (Yancey 2020) and include changes in membrane fluidity (high-pressure and low-temperature rigidify membranes), extrinsic adaptations (i.e., changes in the cellular milieu such as the expression of molecular chaperones or increased concentration of piezolytes, i.e., stabilizing solutes), and intrinsic adaptations (i.e., changes in protein sequences). An essential intrinsic adaptation of deep-sea proteins is their increased stability (i.e., more resistant to denaturation) compared with their shallow-water counterparts (Somero 1990, 1992, 2003; Gross and Jaenicke 1994; Siebenaller 2010). This has been shown in numerous studies, which mainly focused on a handful of proteins (such as lactate dehydrogenase) in deep-sea fishes (Siebenaller and Somero 1978, 1979; Dahlhoff and Somero 1991; Morita 2003, 2008; Wakai et al. 2014; Lemaire et al. 2018; Suka et al. 2019). Interestingly, patterns of protein adaptation to temperature show higher flexibility (i.e., decreased stability) with decreasing temperature (Fields et al. 2015; Saarman et al. 2017). As pressure and temperature strongly covary in the deep sea—temperature decreases as pressure increases—it therefore can be difficult to disentangle the respective combined or opposing effects of these factors on protein stability evolution.

Patterns of positive selection have been investigated to uncover genes underlying adaptation to specific environments, including the deep sea, in nonmodel species (Oliver et al. 2010; Kober and Pogson 2017; Sun et al. 2017; Weber et al. 2017; Zhang et al. 2017; Lan et al. 2018). Although valuable, these studies typically focus on a single or few shallow-deep transitions in a limited number of species and thus lack the comparative power to separate confounding effects. With almost 2,100 species, brittle stars (Ophiuroidea) are a large and ancient class of echinoderms (Stöhr et al. 2012, 2020). These diverse marine invertebrates have colonized every marine habitat, highlighting their strong adaptive abilities. Furthermore, their phylogeny is well resolved (O'Hara et al. 2014, 2017) and they represent a major component of the deep-sea fauna in terms of abundance and species diversity (Christodoulou et al. 2019; Christodoulou et al. 2020), making them important models for marine biodiversity and biogeography (Woolley et al. 2016; O'Hara et al. 2019). It is usually assumed that deep-sea organisms


colonized the deep sea from shallow waters (Brown and Thatje 2014); however, colonization from deep to shallow waters has also been reported (Brown and Thatje 2014; Bribiesca-Contreras et al. 2017). Four large independent ophiuroid families (Amphiuridae, Ophiodermatidae, Ophiomyxidae, and Ophiotrichidae) have a common ancestor from shallow water with extant species occurring in deep and shallow seas (Bribiesca-Contreras et al. 2017). Due to these repeated and independent deep-sea colonization events, these four brittle star families provide an ideal framework to test for convergent molecular evolution to the deep sea.

Materials and Methods


Phylogenomic Data Generation and Processing

The data used in this study are an extension (436 additional samples) of a previously published exon-capture phylogenomic datamatrix of 1,484 exons in 416 genes for 708 individual ophiuroid samples representative of the whole class Ophiuroidea (O'Hara et al. 2019). The full data set used here encompassed 1,144 individual ophiuroid samples accounting for 826 species. Details on specimen collection, environmental parameters, and list of species are available in [supplementary table S1, Supplementary Material](#) online. The set of 416 single-copy genes were first determined in a phylogenomic analysis to resolve the phylogeny of Ophiuroidea (O'Hara et al. 2014). These genes were rigorously screened to be one-to-one orthologs in 52 ophiuroid transcriptomes (representative of the entire class), in the sea urchin *Strongylocentrotus purpuratus*, the hemichordate *Saccoglossus kowalevskii*, and the fish *Danio rerio*. The subsequent exon-capture system laboratory, bioinformatic, and phylogenetic procedures are described in previous works (O'Hara et al. 2017; Hugall et al. 2016) and summarized in dryad packages <https://doi.org/10.5061/dryad.db339> and <https://doi.org/10.5061/dryad.rb334>. Briefly, base-calling used a minimum read coverage of five. Exon boundaries were initially based on the *S. purpuratus* and *Danio rerio* genomes and then revised using the exon-capture read mapping information. For all selection analyses, each codon immediately adjacent to exon boundaries was ignored. The primary data had IUPAC-coded heterozygous sites, which were then randomly resolved. However, these sites had little influence as both ambiguity-coded and randomly resolved data sets returned the same positive selection test results. A global phylogenetic tree of all 1,144 samples for 416 genes (273 kb sites) was generated via RAXML v.8.2.11 (Stamatakis 2014) using a codon position partition model. First, a fully resolved all-compatible consensus topology was generated from 200 RAXML fast bootstrap samples (the -f -d command), onto which branch lengths were then optimized using a codon position GTR- Γ model (the -f -e command). The tree was

A Phylogeny of Ophiuroidea



B 416 single-copy orthologs: ML reconstruction



C Candidate gene detection and analyses




Fig. 1—Workflow used in this study. (A) Schematic representation of the phylogeny of Ophiuroidea (redrawn from Christodoulou et al. [2019]). Four families (288 individuals from 216 species) with a shallow-water common ancestor and extant species in shallow (0–200 m) and deep (>200 m) environments are highlighted in different colors. The width of each triangle is proportional to the sampled number of species in each family. Ophiomyxidae (blue), Ophiodermatidae (green), Amphiuroidae (pink), and Ophiotrichidae (yellow). (B) For each family (number of species used in brackets) and each one of the 416 single-copy orthologs, maximum-likelihood (ML) reconstructions were performed. Trees are drawn for illustration only with the branches of shallow-water (0–200 m) species colored in black and deep-water (>200 m) species in blue. (C) For each resulting ML tree, deep species were labeled as foreground branches and four positive selection detection methods were used (BUSTED, aBSREL, MEME, and MNM). To detect and exclude candidate genes displaying relaxation of selection, that is, accumulation of substitutions not due to increased selection pressure, the method RELAX was used. The final set of candidate genes for each family encompassed genes positively selected in at least three methods and not displaying relaxation of selection. Convergent evolution was examined by overlapping candidate genes per family. For the most interesting candidate gene, amino-acid convergence and tertiary structure analyses were performed, and protein stability profiles and global protein stability were calculated.

then rooted according to O'Hara et al. (2017) defining the sister superorders Ophintegrida and Euryophiurida.

Four brittle star families were investigated that included species displaying independent events of deep-sea colonization from shallow water (Bribiesca-Contreras et al. 2017): Amphiuroidae (111 individuals from 95 species, depth range: –0.5 to –5,193 m; temperature range: –1.6 to 28.8 °C), Ophiodermatidae (60 individuals from 38 species, depth range: –0.5 to –1,668 m; temperature range: 2.6–28.3 °C), Ophiomyxidae (41 individuals from 29 species, depth range: –1.5 to –792 m; temperature range:

4.6–28.7 °C), and Ophiotrichidae (78 individuals from 62 species, depth range: –1 to –405 m; temperature range: 10.5–29.5 °C) (fig. 1A). Positive selection analyses were conducted separately per family. In total, 1,664 alignments were generated, representing each gene (416) in each family (4). In each alignment, a maximum of 30% missing data per sequence was allowed. Alignments that lacked deep species (>200 m) after filtering were not used. As all these four families belong to the superorder Ophintegrida, sequences of *Asteronyx loveni* belonging to the sister superorder Euryophiurida (Asteronychidae) were

Table 1

Common Positively Selected Candidate Genes in Three Families and Their Characteristics (Positively Selected in Three of Four Methods, Not Displaying Relaxation of Selection)

Gene Name	Description	Blast Reference Sequence	GO Terms: Biological Process	Positively Selected in
CCT α ^a	Chaperonin-containing TCP1 complex subunit α	XP_780270.1	<i>Protein folding</i>	Amphiuridae, Ophiidermatidae, Ophiomyxidae
PFD3	Prefoldin subunit 3	XP_797937.1	Macromolecular complex assembly; protein complex assembly; <i>protein folding</i>	Amphiuridae, Ophiidermatidae
tkt ^a	Transketolase 2 isoform X2	NP_1229589.1	Biological process	Amphiuridae, Ophiidermatidae
rpl34	Subunit ribosomal	XP_797232.1	Ribosome biogenesis; <i>translation</i>	Amphiuridae, Ophiomyxidae
rpl8	60S ribosomal L8	XP_796001.1	<i>Translation</i>	Amphiuridae, Ophiomyxidae

NOTE.—Common biological process annotation is shown in italics.

^aPositively selected in four of four methods, not displaying relaxation of selection.

used as outgroups. After filtering, 1,649 alignments were available for further analyses.

Phylogenetic Reconstruction and Positive Selection Analyses

For each of the 1,649 alignments, a maximum-likelihood (ML) phylogeny was reconstructed using RAXML v.8.2.11 with the following parameters: -x 12345 -# 100 -f a -m GTRGAMMA -p 12345 (fig. 1B). Deep (>200 m) species (tips) and monophyletic groups of deep species (nodes) were labeled as “Foreground” branches for positive selection analyses (Amphiuridae: 46 species; 11 independent events of deep-sea colonization; Ophiidermatidae: seven species; four independent events; Ophiomyxidae: ten species; four independent events; and Ophiotrichidae: four species; four independent events). Then, the package HyPhy was used to conduct several positive selection analyses (fig. 1C): 1) Branch-site Unrestricted Statistical Test for Episodic Diversification (BUSTED) (Murrell et al. 2015) to test for gene-wide positive selection (at least one site on at least one branch); 2) adaptive Branch-Site Random Effects Likelihood (aBSREL) (Smith et al. 2015) to detect specific branches evolving under episodic positive selection; 3) Mixed Effects Model of Evolution (MEME) (Murrell et al. 2012) to find sites evolving under episodic positive selection; 4) multinucleotide mutation (MNM) method; it has been shown recently that mutations at adjacent sites often occur as a result of the same mutational event (i.e., MNMs) and therefore may bias classical branch-site tests for positive selection (Venkat et al. 2018). The authors of that study developed a new model of positive selection detection incorporating MNMs, which we also used on deep lineages (>200 m). For each gene, *P* values were corrected for multiple testing using the Holm method (Holm 1979). The *P* value significance level used for all the positive selection detection methods was 0.05. Finally, we used 5) RELAX (Wertheim et al. 2015) to test for relaxation of selection and exclude potential candidate genes displaying relaxation of selection. For each of the four families, positively selected candidate genes of each

method were overlapped on a Venn diagram (fig. 1C and supplementary fig. S1, Supplementary Material online). To be considered as a candidate gene for positive selection in one family and to minimize the risk of false positives, a gene had to display a significant signal in at least three out of four methods including MEME and MNM (BUSTED, MEME, and MNM or MEME, MNM, and aBSREL) and not display relaxation of selection (RELAX). This set of candidates was used for functional annotation. Final sets of positively selected genes per family were then compared among each other to test for convergent evolution. To confirm that positive selection was detected only in deep-sea lineages, positive selection was also tested in shallow-water lineages for the five genes displaying convergent positive selection signatures (chaperonin-containing TCP-1 subunit α [CCT α], PFD3, tkt, rpl34, and rpl8; see Results, table 1, and supplementary table S3, Supplementary Material online). For each family, the same number of shallow-water species as was used for deep-water species was randomly labeled as “Foreground” in each gene tree and positive selection tests were performed for the five methods as described above.

Gene Ontology Annotations and Amino-Acid Convergence Analyses

To explore which functions may be involved in deep-sea adaptation, the representative sequence of each of the 416 genes was extracted from the sea urchin *S. purpuratus* genome and blasted against the nr database from NCBI using BLAST+. We used *S. purpuratus* as reference because sequence annotation for this species is of high quality (no high-quality brittle star reference genome is currently available) and to use a single complete representative sequence for each gene. The top 50 hits were extracted and loaded in BLAST2GO v.4.1. for annotation (Conesa et al. 2005). Mapping, annotation, and slim ontology (i.e., gene ontology [GO] subsets of broader categories) were performed with BLAST2GO using default parameters, except for the

annotation cutoff parameter, which was set to 45. GO categories were described using the level 3 of slim ontology.

CCT α , the only candidate gene displaying positive selection signal in three families (see Results) was further analyzed for signatures of convergent evolution. Specifically, amino-acid profiles were investigated for convergent shifts using PCOC (Rey et al. 2018). This method, which has been shown to display high sensitivity and specificity, detects convergent shifts in amino-acid preferences rather than convergent substitutions. The CCT α amino-acid alignment encompassing the four families and outgroups was used to generate a ML phylogeny as previously described but this time using the PROTGAMMAWAG protein model of sequence evolution (supplementary fig. S2, Supplementary Material online). For each family, positively selected branches resulting from aBSREL analyses were labeled as foreground branches (i.e., the branches with the convergent phenotype in the nucleotide topology) in four different scenarios: 1) Amphiuridae, Ophiidermatidae, and Ophiomyxidae; 2) Amphiuridae, Ophiidermatidae; 3) Amphiuridae, Ophiomyxidae; and 4) Ophiidermatidae, Ophiomyxidae. Detection of amino-acid convergence in these four scenarios was then performed using PCOC and a detection threshold of 0.9 (Rey et al. 2018).

Protein Structure Modeling, Protein Stability Profile, and Overall Protein Stability

To infer the position of positively selected mutations on CCT α , the corresponding amino-acid sequence of the individual *Amphiura constricta* MVF214041 was used to obtain the secondary and tertiary protein structures of this gene. This shallow-water species was chosen because its CCT α sequence had no missing data. The secondary structure was modeled using InterPro 72.0 web browser (<https://www.ebi.ac.uk/interpro>). The protein model was generated using the normal mode of the online Phyre² server (<http://www.sbg.bio.ic.ac.uk/~phyre2/>) (Kelley et al. 2015). The online server EzMol 1.22 was used for image visualization and production (<http://www.sbg.bio.ic.ac.uk/ezmol/>) (Reynolds et al. 2018). The online server NetSurfP-2.0 (<http://www.cbs.dtu.dk/services/NetSurfP/>) was used to predict amino-acid surface accessibility and secondary structures.

We then examined the protein stability profiles of CCT α across the whole ophiuroid class (967 sequences with <30% missing sites, representing 725 species) using eScape v2.1 (Gu and Hilser 2008, 2009). This algorithm calculates a per-site estimate of Gibbs free energy of stabilization based on a sliding window of 20 residues. More specifically, it models the contribution of each residue to the stability constant, a metric that represents the equilibrium of the natively folded and the multiple unfolded states of a protein (D'Aquino et al. 1996). Sites adapted to elevated pressure (or high temperature at atmospheric pressure) are expected to display stabilizing mutations (i.e., more negative ΔG values), whereas sites

adapted to low temperatures at atmospheric pressure are expected to display mutations increasing flexibility (i.e., decreasing stability, thus more positive ΔG values) (Fields et al. 2015; Saarman et al. 2017). For each site of the apical domain (codons 203–353), we calculated the average ΔG value for all 324 shallow-water species (424 individuals) (0–200 m) and 401 deep-water species (543 individuals) (>200 m) and their respective standard deviation. To test the difference between these average values in a phylogenetic context, we used phylogenetically corrected analysis of variance (ANOVA; R function `phylANOVA` of the `phytools` v.0.6-60 R package; 10,000 simulations). To correct for relatedness among species, we used the global RAXML phylogenetic tree pruned to the 967 tips. To investigate regions rather than individual codons, we contrasted shallow versus deep species along the whole gene, averaging ΔG values across ten residues and performing a phylogenetically corrected ANOVA as previously described.

Finally, we tested how the mutations at the positively selected sites, and the ones with local stability values significantly different between shallow and deep species in the phylogenetically corrected ANOVA, impacted the global protein stability. For each one of these sites, we identified “shallow-water” and “deep-water” mutations and used the shallow-water *Amphiura constricta* MVF214041 sequence as template to add these “shallow-water” and “deep-water” mutations one at the time. A model structure for each one of these new sequences was generated using the normal mode of the online Phyre² server (<http://www.sbg.bio.ic.ac.uk/~phyre2/>) (Kelley et al. 2015). The global stability of each model structure was then calculated using the “stability” function of the program FoldX with default values (Schymkowitz et al. 2005). For each site, the $\Delta\Delta G$ value between “deep-water” (mutant) and “shallow-water” (wild-type) mutations was calculated, where $\Delta\Delta G > 0.5$ kcal/mol indicates stabilizing mutations, $\Delta\Delta G < -0.5$ kcal/mol indicates destabilizing mutation, and -0.5 kcal/mol $< \Delta\Delta G < 0.5$ kcal/mol indicates neutral mutations (Khatun et al. 2004; Khan and Vihinen 2010).

Results

Five Genes Involved in Protein Biogenesis Are Recurrently Positively Selected in Deep-Sea Brittle Stars

We used 416 single-copy orthologs from 216 species (288 individuals) of four brittle star families (fig. 1A) to examine patterns of positive selection in deep-sea species (>200 m). For each gene of each family, we used four different positive selection methods and one method detecting relaxation of selection (fig. 1B and C). Given that our data set is not representative of the whole genome, we chose a stringent approach to focus on a few candidate genes displaying strong signals of positive selection. Therefore, we only kept candidate genes that had significant signatures of positive selection

in at least three methods and which also did not show relaxation of selection. We found 36 candidate genes in Amphiuroidae, nine in Ophiordermatidae, six in Ophiomyxidae, and none in Ophiotrichidae (supplementary table S2 and fig. S1, Supplementary Material online). Four genes involved in embryo development were positively selected in Amphiuroidae. Common functions among candidate genes from different families include protein folding, translation (i.e., ribosomal proteins), catabolic process, and cell proliferation (supplementary table S2, Supplementary Material online).

Five genes were positively selected in at least two families, which are notably involved in protein folding and in translation (table 1). Among these, CCT α was positively selected in all three families and significant in each one of the selection detection methods (table 1). To confirm that positive selection was detected only in shallow-deep transition, we performed positive selection analyses on these five common candidate genes but this time labeling shallow-water lineages as "Foreground." Most of the genes did not display signatures of positive selection in shallow-water environments, except PFD3 in Amphiuroidae and tkt in Ophiotrichidae, which were significant in the BUSTED, aBSREL, and MEME methods (supplementary table S3, Supplementary Material online). This suggests that these two genes are evolving faster not only in response to deep-sea conditions but also possibly in response to additional environmental conditions.

CCT α , the sole candidate gene detected as significantly positively selected in three families in the shallow-deep contrast, did not display signatures of positive selection in shallow-water environments (supplementary table S3, Supplementary Material online). It is a subunit of the octameric CCT complex, a cytosolic eukaryotic chaperonin having a central role in protein folding (fig. 2A–D) (Valpuesta et al. 2005; Bueno-Carrasco and Cuéllar 2018). The three other CCT subunits present in our data set, CCT δ , CCT ϵ , and CCT η , did not display consistent signatures of positive selection; rather, although positive selection was detected at the site level in all subunits, relaxation of selection was detected in all three subunits in Amphiuroidae (supplementary table S6, Supplementary Material online), indicating that the accumulation of mutations in these subunits is not clearly due to natural selection. Therefore, CCT α was the only one out of four subunits to show consistent signatures of positive selection across methods. The remaining four subunits of the CCT complex could not be tested as they were not present in our data set.

Interestingly, PFD3, a subunit of the hexameric cochaperone prefoldin interacting with CCT (Martín-Benito et al. 2002; Gestaut et al. 2019), was positively selected in two families (Amphiuroidae: MNM, MEME, and BUSTED; Ophiordermatidae: MNM, aBSREL, MEME, and BUSTED) (table 1; supplementary tables S2 and S6, Supplementary Material online; fig. 2D), although it was also positively

selected in shallow-water Amphiuroidae (supplementary table S3, Supplementary Material online). The two other prefoldin subunits present in our data set (PFD1 and PFD5) did not show a consistent signature of positive selection (supplementary table S4, Supplementary Material online), suggesting that subunits of this cochaperone can evolve relatively independently from each other. Finally, two ribosomal proteins (Rpl8 and Rpl34) were recurrently positively selected in Amphiuroidae and Ophiomyxidae, whereas the ribosomal protein S6 was positively selected in Ophiordermatidae (table 1 and supplementary table S2, Supplementary Material online).

CCT α and Deep-Sea Adaptation: Convergence at the Gene but Not the Amino-Acid Level

Although positive selection was detected at the gene level in three families, positive selection at the site level (MEME) was detected in all four families. The sites displaying positive selection in CCT α were not the same among the four families, except for site 26, which was shared between Amphiuroidae and Ophiotrichidae (fig. 2E and table 2 and supplementary table S5, Supplementary Material online). Three sites were found in the equatorial domain, that is, the ATP-binding region, two sites were found in the intermediate domain, and three sites were found in the apical domain, that is, the substrate-binding region (fig. 3) (Bueno-Carrasco and Cuéllar 2018). Furthermore, we found that seven out of eight positively selected sites increased the global protein stability, suggesting that these sites may be involved in structural adaptation of CCT α (table 2). This is corroborated by the fact that six out of eight mutations involve changes toward hydrophobic amino acids, which are known to increase protein stability (Kellis et al. 1988; Pace et al. 2011). Finally, convergent evolution at the site level was not detected when examining amino-acid profiles (PCOC posterior probabilities at all sites <0.9).

Energetic Landscapes Reveal Structural Adaptation within and next to the Substrate-Binding Region

Next, we calculated site-specific protein stability profiles of CCT α in 967 individuals from 725 species representative of the whole Ophiuroidea class, to test the hypothesis that deep-sea-adapted proteins are more stable than their shallow-water counterparts. For each site, we compared the average stability measure of 324 shallow-water species (0–200 m depth) versus 401 deep-water species (>200 m depth), where lower ΔG values correspond to higher stability (fig. 2F and supplementary fig. S3A, Supplementary Material online). We focused on the apical domain as it encompasses the substrate-binding region, whose position and structure are highly conserved across eukaryotes (Joachimciak et al. 2014). This region is composed of the proximal loop (PL), the apical hinge (AH) and Helix 11 (H11) (figs. 2A and 3). Although AH and H11 are almost invariant across all ophiuroids, the stability




FIG. 2—Structure and function of the CCT complex, selection analyses on CCT α , and comparison of local stability values from CCT α apical domain between shallow and deep species. (A) Model of tertiary structure of the CCT α subunit. Each subunit is composed of an apical domain (AD; green) containing the substrate-binding regions (PL: proximal loop; AH: apical hinge; H11: Helix 11), an intermediate domain (ID; blue) and an equatorial domain (ED; pink) containing the ATP-binding sites and where hydrolysis takes place. (B) Model of the top view of the CCT complex, encompassing eight paralogous subunits. (C) Quaternary structure model of the CCT complex encompassing a double ring of eight paralogous subunits. (D) Simplified model of prefoldin (PFD)-CCT interaction in the folding of newly synthesized actin or tubulin. (A–D) Adapted from Bueno-Carrasco and Cuéllar (2018). (E) Localization of the positively selected sites on the tertiary structure of CCT α in the four ophiuroid families investigated. (F) Average protein stability profiles and respective standard deviations (vertical bars) for each codon of the CCT α apical domain in 324 species (424 individuals) from shallow water (0–200 m; in black) and 401 species (543 individuals) from deep water (>200 m; in blue) representative of the whole ophiuroid class. A smaller (i.e., more negative) value of ΔG is indicative of substitutions increasing stability. The substrate-binding regions PL, AH, and H11 are highlighted as well as the positively selected sites.

Table 2Summary of the CCT α Positively Selected Sites in Each Family and Sites with Local Stability Values Significantly Different between Shallow and Deep Species

Site	Tertiary Structure	Detection Method	Amino-Acid		Amino-Acid	Amino-Acid	$\Delta\Delta G$ (kcal/mol)
			"Deep"	Amino-Acid Property			
26	ED	PS Amphiuroidae and PS Ophiotrichidae	A	Hydrophobic	S	Polar uncharged	34.7
175	ID	PS Amphiuroidae	Y	Hydrophobic	S	Polar uncharged	11.8
203 ^a	AD	phylANOVA	P	Special case	Q	Polar uncharged	-17.3
215 ^a	AD	phylANOVA	V	Hydrophobic	P	Special case	2.1
219	AD	PS Ophiotrichidae	I	Hydrophobic	T	Polar uncharged	3.7
263 ^a	AD	phylANOVA	H	Positively charged	Q	Polar uncharged	22.9
327 ^a	AD	phylANOVA	S, V	Polar uncharged, hydrophobic	A	Hydrophobic	11.9, 30.1
328 ^a	AD	phylANOVA	T	Polar uncharged	S	Polar uncharged	27.6
329	AD	PS Amphiuroidae	V	Hydrophobic	M	Hydrophobic	15.7
350	AD	PS Ophiidermatidae	I	Hydrophobic	K	Positively charged	15.7
375	ID	PS Amphiuroidae	V	Hydrophobic	I	Hydrophobic	12.3
462	ED	PS Ophiomyxidae	C	Special case	A	Hydrophobic	-3.8
465	ED	PS Ophiidermatidae	F	Hydrophobic	K	Positively charged	22.9

NOTE.—For each site, the "deep" and "shallow" amino acids are reported, as well as their impact on global protein stability: $\Delta\Delta G > 0.5$ kcal/mol: mutations increasing stability; $\Delta\Delta G < -0.5$ kcal/mol: mutations decreasing stability; and -0.5 kcal/mol $< \Delta\Delta G < 0.5$ kcal/mol: neutral mutations in terms of stability. ED, equatorial domain; ID, intermediate domain; AD, apical domain; PS, positively selected.

^aThe most common amino acids between shallow and deep species are reported.

measure was lower (i.e., more stable) in deep compared with shallow species within the PL and in two sites following the PL (codons 214–217), close to a positively selected site (codon 219) (fig. 2F and [supplementary fig. S3A, Supplementary Material](#) online). In contrast, three codons displayed significantly higher flexibility in deep compared with shallow species ([supplementary fig. S3A and B, Supplementary Material](#) online), suggesting that increased flexibility may play a role in deep-sea adaptation outside the ligand binding region, possibly related to low temperature. Nevertheless, when averaging ΔG values across ten codons, only the signal close to PL remained significant in the phylogenetically corrected ANOVA contrasting stability values of shallow and deep species ([supplementary fig. S4, Supplementary Material](#) online). This indicates that substitutions toward a more stable PL occurred independently in the ophiuroid tree of life. Finally, we examined how these mutations impacted the global (rather than local) stability value of the protein. We found that four out of five mutations increased the global protein stability ([table 2](#)), including the site located in the substrate-binding region (i.e., site 215 in the PL), confirming the results found with the local stability values.

Discussion

Protein Biogenesis Is Essential for Deep-Sea Adaptation

In this study, we examined patterns of positive selection in 416 genes from four brittle star families representing replicated instances of deep-sea colonization. We found that four genes involved in functions related to protein biogenesis including

protein folding and protein synthesis (i.e., ribosomal proteins) were convergently positively selected in three out of four families. Furthermore, an additional ribosomal protein was positively selected in Ophiidermatidae and two transcription factors were positively selected in Amphiuroidae. Previous studies have shown that several genes involved in genetic information processing such as protein folding, transcription, or translation (including ribosomal proteins) were positively selected in the hadal (>6,000 m depth) amphipod *Hirondella gigas* (Lan et al. 2017), in the bathyal (200–3,000 m depth) fish *Aldrovandia affinis* (Lan et al. 2018), and in the bathyal barnacle *Glyptelasma gigas* (Gan et al. 2020). Taken together, these and previous results indicate that adaptations in protein biogenesis are essential to deep-sea life.

At the individual family level, four genes related to embryo development were positively selected in Amphiuroidae. Interestingly, nine genes related to the same function (embryo development; GO:0009790) were detected as candidate genes for deep-sea adaptation between populations of the deep-sea fish *Coryphaenoides rupestris* living at different depths (Gaither et al. 2018). This suggests that there might be similar selection pressures on development at both young (i.e., species level) and old (i.e., family level) evolutionary time-scales for life at great depths. Finally, no candidate genes for deep-sea adaptation were found in Ophiotrichidae across different positive selection methods. This might be explained by the fact that this family is the youngest from a fossil and a phylogenetic point of view (O'Hara et al. 2017). In that sense, it started colonizing the deep sea later than the other families, and therefore there was possibly less time for natural selection




FIG. 3—Primary structure of CCT α . Ligand binding sites (PL, AH, and H11), ATP-binding sites, positively selected sites (method MEME), and sites displaying different local stability values between shallow and deep species (method phylANOVA) are highlighted. The sequence of the shallow-water Amphiuroidae *Amphiura constricta* is used as template.

to act. In support with this, the maximum sampling depth for Ophiotrichids was 405 m, whereas it was deeper for the three other families (Amphiuridae: 5,193 m; Ophiodermatidae: 1,668 m; and Ophiomyxidae: 792 m).

Independent Evolutionary Histories of Subunits from Macromolecular Complexes

CCT α and prefoldin 3 (PFD3), the two genes recurrently positively selected involved in protein folding, are subunits of octameric and hexameric complexes, respectively. The CCT complex is estimated to fold ~10% of newly synthesized proteins, including actin and tubulin, and is involved in numerous core cellular processes such as cytoskeleton formation, cell signaling, cell recognition, and protein degradation

(Bueno-Carrasco and Cuéllar 2018), whereas the prefoldin complex is known to assist CCT in actin folding (Martín-Benito et al. 2002). We tested four subunits of the octameric CCT complex, yet CCT α was the only one to show consistent signatures of positive selection across different methods. Indeed, the other tested subunits displayed relaxation of selection or no signatures of positive selection. This might be due to the different degrees of subunit specialization in CCT, as CCT α has intermediate binding properties (i.e., neither high ATP affinity nor high substrate affinity) compared with the other subunits (Bueno-Carrasco and Cuéllar 2018). Therefore, it is possible that CCT α can evolve more rapidly because it may experience less negative selection than the other subunits. In the same way, we tested three subunits of the hexameric prefoldin complex, and PFD3 was the only

subunit to be consistently positively selected. Although we acknowledge that we analyzed an existing data set and thus did not test all subunits of both macromolecular complexes (four out of eight subunits in CCT; three of six subunits in prefoldin), it appears surprising that individual subunits of macromolecular complexes can evolve relatively independently from each other.

We note that a previous study actually highlighted evolutionary independence of CCT subunits across old evolutionary timescales (i.e., among eukaryotes), showing that CCT α , CCT γ , and CCT ζ evolved under positive selection after duplication events which led to subfunctionalization in eukaryotes, most likely in response to folding increasingly complex cytosolic proteins (Fares and Wolfe 2003). Furthermore, it has been shown that the exact same mutation in the conserved ATP-binding region of all eight CCT subunits has dramatically different phenotypic outcomes in yeast, such as differential temperature sensitivity, differential growth in presence of actin and tubulin polymerization inhibitors, or differential over and underexpression of genes (Amit et al. 2010). This not only indicates that the functional activity of each subunit is not redundant but also possibly highlights “moonlighting” proteins (i.e., other functions of individual subunits that are carried out not as part of the full complex) (Jeffery 2009; Huberts and van der Klei 2010). In that sense, the functional coherence of several hetero-oligomeric protein complexes including CCT has been reviewed (Matalon et al. 2014). Interestingly, they found a lack of coherence among subunits at virtually all functional levels, including different levels of subunit abundance, differences in transcriptional regulation, differential subunit duplication or loss, and finally lack of coherence in genetic interaction. Explanations for this lack of coherence include enhancing complex assembly or subunit multifunctionality (Matalon et al. 2014). There is increasing evidence for the latter case, as many CCT monomer functions have been reported and recently reviewed (Vallin and Grantham 2019). Taken together, our results are consistent with the hypothesis that CCT α and PFD3 may be moonlighting proteins involved in deep-sea adaptation.

Convergent Evolution, Functional, and Structural Adaptations in CCT α

Our results suggest that convergent evolution was detected at the functional (i.e., protein folding, translation) and at the gene levels (i.e., CCT α , PFD3, tkt, rpl8, rpl34) in deep-sea adaptation. Yet, it was not detected at the amino-acid level in CCT α , as the majority of positively selected sites were different among families, except for one site shared between Amphiuroidae and Ophiotrichidae. This might be due to the old divergence time among the investigated brittle star families and the broad nature of the selection regime (i.e., depth and temperature). It has been shown that rates of molecular convergence decrease with time (Storz 2016) and the last

common ancestor of Amphiuroidae, Ophiidermatidae, and Ophiomyxidae is estimated to be ~250 Myr old (O'Hara et al. 2017). Furthermore, convergence at the amino-acid level is often less common than convergence at higher levels of biological hierarchy (e.g., gene, pathway, or species levels) (Tenaillon et al. 2012, 2016; Bolnick et al. 2018).

Our results indicate that CCT α evolved under positive selection during deep-sea colonization and speciation events. These mutations possibly reveal functional adaptations of CCT α , such as adaptations in ATP or substrate binding. Indeed, it has previously been reported that the shallow groove created by the conserved H11 and the flexible PL allows the binding of a variety of substrates (Yam et al. 2008; Joachimiak et al. 2014), suggesting that the detected substitutions in the PL and in adjacent amino acids allow efficient substrate binding in deep-sea species. Similarly, in a study on metabolic enzymes from 37 ctenophores, numerous sites associated with adaptation to depth, temperature, or both were located close to the substrate-binding region (Winnikoff et al. 2019). A major substrate of the CCT complex and its cochaperone prefoldin is actin, an essential cytoskeleton protein (Martín-Benito et al. 2002). Although it should be noted that CCT α is not directly involved in actin folding (actin binds the subunits CCT δ , CCT β , and CCT ϵ) (Llorca et al. 1999, 2000), examination of cytoskeletal proteins is of interest due to their essential role in cells. Unfortunately, as actin and tubulin were not included in our data set, no conclusions about cytoskeletal proteins of deep-sea brittle stars can yet be drawn. However, actin of deep-sea fishes has been extensively studied, showing that it displays structural adaptations to high-pressure which enhance stability (Morita 2003; Koyama and Aizawa 2008; Wakai et al. 2014; Yancey 2020). Furthermore, genome analysis of the hadal fish *Pseudoliparis swirei* revealed that a gene family related to cytoskeleton was expanded, and several genes from three GO categories linked to cytoskeleton were evolving under positive selection (Wang et al. 2019). Finally, 13 genes involved in cytoskeleton systems were positively selected in the deep-sea fish *Aldrovandia affinis* (Lan et al. 2018), suggesting that cytoskeletal proteins are pivotal for deep-sea adaptation.

Furthermore, we tested the hypothesis that CCT α is structurally adapted to deep-sea conditions, as a common feature of high-pressure-adapted proteins is to display increased stability (Somero 2003; Ohmae et al. 2013; Yancey and Siebenaller 2015). We investigated local and global stability in CCT α and found that four out of five sites with phylogenetically-corrected significantly different local stability showed increased global protein stability, suggesting that these mutations reflect structural deep-sea adaptations. Moreover, seven out of eight mutations at positively selected sites increased global protein stability and involved hydrophobic amino-acid substitutions, which are known to increase protein stability (Kellis et al. 1988; Pace et al. 2011). It has also been shown that several proteins from the abyssal fish *Coryphaenoides*

armatus displayed increased protein stability as an adaptation to hydrostatic pressure (Morita 2003, 2008; Brindley et al. 2008; Lemaire et al. 2018). Finally, several critical mutations in both the enzyme synthesizing trimethylamine oxide (a heavily studied stabilizing cosolute or piezolyte [Yancey 2020]) and in HSP90 have been proposed to enhance overall protein stability in the hadal fish *Pseudoliparis swirei* (Wang et al. 2019).

Role of Molecular Chaperones in Deep-Sea Adaptation

We have shown that over deep evolutionary timescales, CCT α displays recurrent signatures of accelerated evolution and structural adaptation in transition from shallow- to deep-sea habitats in brittle stars. This contrasts with previous findings on sea urchins where, CCT β —but not the other subunits—showed signatures of positive selection, yet not in the two deep-sea species included in the work (Kober and Pogson 2017). A functional equivalent of the eukaryotic chaperonin CCT in bacteria is the chaperonin complex GroEL/GroES (Gupta 1990; Mayhew et al. 1996). This chaperonin has been shown to be constitutively highly expressed in the deep-sea bacteria *Shewanella*, and that a major target of this chaperonin was a subunit of the 50S ribosome (Sato et al. 2015). Furthermore, convergent evolution in chaperones has previously been shown (Draceni and Pechmann 2019), and there is evidence that other groups of molecular chaperones such as HSP70 and HSP90 (Kim et al. 2013) are involved in deep-sea adaptation (Yancey 2020). For instance, the HSP70 gene family was expanded in a deep-sea mussel (Sun et al. 2017) and convergent amino-acid changes have been reported in four out of five copies of the HSP90 protein in a hadal fish (Wang et al. 2019). In our data set, two HSP90 genes were present (HSP90b1 and HSP90ab1), yet neither showed signatures of accelerated evolution. Nevertheless, only the examination of whole genomes will give a complete view of HSP family evolution in brittle stars, which should be explored in further studies.

Similarities between Deep-Sea and Cold Adaptation Mechanisms

There are numerous studies on the role of CCT at shorter evolutionary timescales, which reveal its role in cold-stress response. Notably, CCT has been characterized as a “cold-shock” protein in several eukaryotes due to the overexpression of the investigated subunits when organisms were exposed to cold stress (Somer et al. 2002; Kayukawa et al. 2005; Yin et al. 2011; He et al. 2017). Furthermore, CCT has been shown to display specific structural (Pucciarelli et al. 2006) and functional (Cuellar et al. 2014) adaptations to cold environment in Antarctic fish, in addition to being also overexpressed in Antarctic fish exposed to heat stress (Buckley and Somero 2009). Moreover, there is evidence for a link between

cold-stress response and high-pressure stress response in bacteria (Welch et al. 1993; Wemekamp-Kamphuis et al. 2002), although it has also been shown that heat-shock proteins are also expressed under high-pressure stress in bacteria (Aertsen et al. 2004) or constitutively in the deep-sea bacteria *Shewanella* (Sato et al. 2015). Additionally, transcriptome analyses revealed that cold-inducible protein families are expanded in a hadal amphipod (Lan et al. 2017), and several proteins involved in cold shock have been shown to evolve under positive selection in deep-sea amphipod and fish (Lan et al. 2018). Taken together, our findings support the hypothesis that cold-shock proteins play an important role in deep-sea adaptation (Brown and Thatje 2014), which was also proposed by others (Lan et al. 2018).

Limitations and Outlook

We acknowledge that our study lacks functional validation to demonstrate that the changes are truly adaptive (which would be experimentally demanding as CCT folds \sim 10% of newly synthesized proteins and we analyzed the sequences of more than 700 species), yet we minimized false inferences by applying stringent positive selection detection criteria. Furthermore, we used a proxy of functional validation by investigating *in silico* local and global protein stability in a data set with great comparative power, both in terms of phylogenetic and environmental diversity. Finally, direct experimental testing on deep-sea organisms remains technically challenging, so we made use of the power of molecular data to indirectly reveal new insights in deep-sea adaptation. Further studies should include whole genomes to obtain a more complete view of deep-sea adaptation hotspots and possible mechanisms, such as recent studies on deep-sea fishes (Gaither et al. 2018; Wang et al. 2019), hydrothermal vent mussel (Sun et al. 2017), or cold seep gastropod (Liu et al. 2020). Also, although we focused on intrinsic adaptations (i.e., at the genomic level), mechanisms of extrinsic adaptations (i.e., changes in the cellular milieu) through, for instance, osmolyte concentration (piezolytes such as trimethylamine oxide) or membrane phospholipid composition changes should not be overlooked (Somero 2003; Yancey and Siebenaller 2015; Yancey 2020). However, this is beyond the scope of this study. With increasing interests in deep-sea biodiversity, ecosystems, and resources in the last decades (Danovaro et al. 2014, 2017; Glover et al. 2018), these are exciting times for diving deeper into mechanisms of deep-sea adaptation.

Supplementary Material

Supplementary data are available at *Genome Biology and Evolution* online.

Acknowledgments

We are grateful to W. Salzburger for providing access to the HPC sciCORE cluster and to J. Sarrazin and S. Arnaud-Haond for comments on a previous version of this manuscript. We thank L. Bribiesca-Contreras for help with R analyses. We thank three anonymous reviewers and the editorial team for valuable comments on a previous version of this manuscript. Calculations were performed at sciCORE (<http://scicore.unibas.ch/>) scientific computing center at the University of Basel, Switzerland, and at DATARMOR (<http://www.ifremer.fr/pcdm>) scientific computing center at the Pôle de Calcul et de Données Marines (PCDM), Ifremer, Brest, France. This work was supported by an Endeavour Postdoctoral Fellowship awarded by the Australian Department of Education and Training (Grant No. 6534_2018 to A.A.-T.W.) and a Marie Skłodowska-Curie Global Fellowship awarded by the European Union's Horizon 2020 Research and Innovation Program (Grant No. 797326 to A.A.-T.W.). Open Access publication fees were covered with a Micro One World Grant awarded by the Marie Curie Alumni Association.

Literature Cited

- Aertsen A, et al. 2004. Heat shock protein-mediated resistance to high hydrostatic pressure in *Escherichia coli*. *Appl Environ Microbiol*. 70(5):2660–2666.
- Amit M, et al. 2010. Equivalent mutations in the eight subunits of the chaperonin CCT produce dramatically different cellular and gene expression phenotypes. *J Mol Biol*. 401(3):532–543.
- Bolnick DI, Barrett RD, Oke KB, Rennison DJ, Stuart YE. 2018. (Non) parallel evolution. *Annu Rev Ecol Syst*. 49(1):303–330.
- Bribiesca-Contreras G, Verbruggen H, Hugall AF, O'Hara TD. 2017. The importance of offshore origination revealed through ophiuroid phylogenomics. *Proc R Soc B* 284(1858):20170160.
- Brindley AA, et al. 2008. Enzyme sequence and its relationship to hyperbaric stability of artificial and natural fish lactate dehydrogenases. *PLoS One* 3(4):e2042.
- Brown A, Thatje S. 2014. Explaining bathymetric diversity patterns in marine benthic invertebrates and demersal fishes: physiological contributions to adaptation of life at depth. *Biol Rev*. 89(2):406–426.
- Buckley BA, Somero GN. 2009. cDNA microarray analysis reveals the capacity of the cold-adapted Antarctic fish *Trematomus bernacchii* to alter gene expression in response to heat stress. *Polar Biol*. 32(3):403–415.
- Bueno-Carrasco MT, Cuéllar J. 2018. Mechanism and function of the eukaryotic chaperonin CCT. *ELS John Wiley Sons Ltd Ed*. p. 1–9.
- Carney RS. 2005. Zonation of deep biota on continental margins. In: *Oceanography and marine biology*. Boca Raton (FL): Taylor & Francis Group. p. 221–288.
- Christodoulou M, O'Hara TD, Hugall AF, Arbizu PM. 2019. Dark Ophiuroid Biodiversity in a Prospective Abyssal Mine Field. *Curr Biol*. 29(22):3909–3912.e3.
- Christodoulou M, et al. 2020. Unexpected high abyssal ophiuroid diversity in polymetallic nodule fields of the northeast Pacific Ocean and implications for conservation. *Biogeosciences* 17(7):1845–1876.
- Conesa A, et al. 2005. Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics* 21(18):3674–3676.
- Cuellar J, et al. 2014. Assisted protein folding at low temperature: evolutionary adaptation of the Antarctic fish chaperonin CCT and its client proteins. *Biol Open* 3(4):261–270.
- Dahlhoff E, Somero GN. 1991. Pressure and temperature adaptation of cytosolic malate dehydrogenases of shallow and deep-living marine invertebrates: evidence for high body temperatures in hydrothermal vent animals. *J Exp Biol*. 159:473–487.
- Danovaro R, Corinaldesi C, Dell'Anno A, Snelgrove PV. 2017. The deep-sea under global change. *Curr Biol*. 27(11):R461–R465.
- Danovaro R, Snelgrove PV, Tyler P. 2014. Challenging the paradigms of deep-sea ecology. *Trends Ecol Evol*. 29(8):465–475.
- D'Aquino JA, et al. 1996. The magnitude of the backbone conformational entropy change in protein folding. *Proteins Struct Funct Bioinf*. 25:143–156.
- Draceni Y, Pechmann S. 2019. Pervasive convergent evolution and extreme phenotypes define chaperone requirements of protein homeostasis. *Proc Natl Acad Sci U S A*. 116(40):20009–20014.
- Fares MA, Wolfe KH. 2003. Positive selection and subfunctionalization of duplicated CCT chaperonin subunits. *Mol Biol Evol*. 20(10):1588–1597.
- Fields PA, Dong Y, Meng X, Somero GN. 2015. Adaptations of protein structure and function to temperature: there is more than one way to 'skin a cat'. *J Exp Biol*. 218(12):1801–1811.
- Gaither MR, et al. 2016. Depth as a driver of evolution in the deep sea: insights from grenadiers (Gadiformes: Macrouridae) of the genus *Coryphaenoides*. *Mol Phylogenet Evol*. 104:73–82.
- Gaither MR, et al. 2018. Genomics of habitat choice and adaptive evolution in a deep-sea fish. *Nat Ecol Evol*. 2(4):680–687.
- Gan Z, et al. 2020. Comparative transcriptomic analysis of deep-and shallow-water barnacle species (Cirripedia, Poecilasmatidae) provides insights into deep-sea adaptation of sessile crustaceans. *BMC Genomics*. 21(1):1–13.
- Gestaut D, et al. 2019. The chaperonin TRiC/CCT associates with prefoldin through a conserved electrostatic interface essential for cellular proteostasis. *Cell* 177(3):751–765.
- Glover AG, Wiklund H, Chen C, Dahlgren TG. 2018. Point of view: managing a sustainable deep-sea 'blue economy' requires knowledge of what actually lives there. *eLife* 7:e41319.
- Gross M, Jaenicke R. 1994. Proteins under pressure: the influence of high hydrostatic pressure on structure, function and assembly of proteins and protein complexes. *Eur J Biochem*. 221(2):617–630.
- Gu J, Hilser VJ. 2008. Predicting the energetics of conformational fluctuations in proteins from sequence: a strategy for profiling the proteome. *Structure* 16(11):1627–1637.
- Gu J, Hilser VJ. 2009. Sequence-based analysis of protein energy landscapes reveals nonuniform thermal adaptation within the proteome. *Mol Biol Evol*. 26(10):2217–2227.
- Gupta RS. 1990. Sequence and structural homology between a mouse T-complex protein TCP-1 and the 'chaperonin' family of bacterial (GroEL, 60–65 kDa heat shock antigen) and eukaryotic proteins. *Biochem Int*. 20(4):833–841.
- He Y, Wang L, Zhu W, Dong Z, Liu N. 2017. Effects of salinity on cold tolerance of Malaysian red tilapia. *Aquacult Int*. 25(2):777–792.
- Holm S. 1979. A simple sequentially rejective multiple test procedure. *Scand J Stat*. 6(2):65–70.
- Huberts DHEW, van der Klei IJ. 2010. Moonlighting proteins: an intriguing mode of multitasking. *Biochim Biophys Acta Mol Cell Res*. 1803(4):520–525.
- Hugall AF, O'Hara TD, Hunjan S, Nilsen R, Moussalli A. 2016. An exon-capture system for the entire class Ophiuroidea. *Mol Biol Evol*. 33(1):281–294.
- Jaenicke R. 1991. Protein stability and molecular adaptation to extreme conditions. In: *EJB reviews* 1991. Berlin, Germany: Springer. p. 291–304.

- Jeffery CJ. 2009. Moonlighting proteins—an update. *Mol Biosyst.* 5(4):345–350.
- Joachimiak LA, Walzthoeni T, Liu CW, Aebersold R, Frydman J. 2014. The structural basis of substrate recognition by the eukaryotic chaperonin TRiC/CCT. *Cell* 159(5):1042–1055.
- Kayukawa T, et al. 2005. Expression of mRNA for the t;chcomplex polypeptide-1, a subunit of chaperonin CCT, is upregulated in association with increased cold hardiness in *Delia antiqua*. *Cell Stress Chaperones* 10(3):204.
- Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJ. 2015. The Phyre2 web portal for protein modeling, prediction and analysis. *Nat Protoc.* 10(6):845–858.
- Kellis JT, Nyberg K, Säil D, Fersht AR. 1988. Contribution of hydrophobic interactions to protein stability. *Nature* 333(6175):784–786.
- Khan S, Vihinen M. 2010. Performance of protein stability predictors. *Hum Mutat.* 31(6):675–684.
- Khatun J, Khare SD, Dokholyan NV. 2004. Can contact potentials reliably predict stability of proteins? *J Mol Biol.* 336(5):1223–1238.
- Kim YE, Hipp MS, Bracher A, Hayer-Hartl M, Ulrich Hartl F. 2013. Molecular chaperone functions in protein folding and proteostasis. *Annu Rev Biochem.* 82(1):323–355.
- Kober KM, Pogson GH. 2017. Genome-wide signals of positive selection in stronglycentrotid sea urchins. *BMC Genomics.* 18(1):555.
- Koyama S, Aizawa M. 2008. Piezotolerance of the cytoskeletal structure in cultured deep-sea fish cells using DNA transfection and protein introduction techniques. *Cytotechnology* 56(1):19–26.
- Lan Y, et al. 2017. Molecular adaptation in the world's deepest-living animal: insights from transcriptome sequencing of the hadal amphipod *Hirondellea gigas*. *Mol Ecol.* 26(14):3732–3743.
- Lan Y, et al. 2018. De novo transcriptome assembly and positive selection analysis of an individual deep-sea fish. *BMC Genomics.* 19(1):394.
- Lemaire B, et al. 2018. Molecular adaptation to high pressure in cytochrome P450 1A and aryl hydrocarbon receptor systems of the deep-sea fish *Coryphaenoides armatus*. *Biochim Biophys Acta Proteins Proteomics* 1866(1):155–165.
- Liu R, et al. 2020. De novo genome assembly of limpet *Bathycyba lactea* (Gastropoda: Pectinodontidae), the first reference genome of a deep-sea gastropod endemic to cold seeps. *Genome Biol Evol.* 12(6):905–910.
- Llorca O, et al. 1999. Eukaryotic type II chaperonin CCT interacts with actin through specific subunits. *Nature* 402(6762):693–696.
- Llorca O, et al. 2000. Eukaryotic chaperonin CCT stabilizes actin and tubulin folding intermediates in open quasi-native conformations. *EMBO J.* 19(22):5971–5979.
- Martín-Benito J, et al. 2002. Structure of eukaryotic prefoldin and of its complexes with unfolded actin and the cytosolic chaperonin CCT. *EMBO J.* 21(23):6377–6386.
- Matalon O, Horovitz A, Levy ED. 2014. Different subunits belonging to the same protein complex often exhibit discordant expression levels and evolutionary properties. *Curr Opin Struct Biol.* 26:113–120.
- Mayhew M, et al. 1996. Protein folding in the central cavity of the GroEL–GroES chaperonin complex. *Nature* 379(6564):420–426.
- Morita T. 2003. Structure-based analysis of high pressure adaptation of α -actin. *J Biol Chem.* 278(30):28060–28066.
- Morita T. 2008. Comparative sequence analysis of myosin heavy chain proteins from congeneric shallow- and deep-living rattail fish (genus *Coryphaenoides*). *J Exp Biol.* 211(9):1362–1367.
- Murrell B, et al. 2012. Detecting individual sites subject to episodic diversifying selection. *PLoS Genet.* 8(7):e1002764.
- Murrell B, et al. 2015. Gene-wide identification of episodic selection. *Mol Biol Evol.* 32(5):1365–1371.
- O'Hara TD, Hugall AF, Thuy B, Moussalli A. 2014. Phylogenomic resolution of the class Ophiuroidea unlocks a global microfossil record. *Curr Biol.* 24(16):1874–1879.
- O'Hara TD, Hugall AF, Thuy B, Stöhr S, Martynov AV. 2017. Restructuring higher taxonomy using broad-scale phylogenomics: the living Ophiuroidea. *Mol Phylogenet Evol.* 107:415–430.
- O'Hara TD, Hugall AF, Woolley SN, Bribiesca-Contreras G, Bax NJ. 2019. Contrasting processes drive ophiuroid phylodiversity across shallow and deep seafloors. *Nature* 565(7741):636–639.
- Ohmae E, Miyashita Y, Kato C. 2013. Thermodynamic and functional characteristics of deep-sea enzymes revealed by pressure effects. *Extremophiles* 17(5):701–709.
- Oliver TA, et al. 2010. Whole-genome positive selection and habitat-driven evolution in a shallow and a deep-sea urchin. *Genome Biol Evol.* 2:800–814.
- Pace CN, et al. 2011. Contribution of hydrophobic interactions to protein stability. *J Mol Biol.* 408(3):514–528.
- Pradillon F, Gaill F. 2007. Adaptation to deep-sea hydrothermal vents: some molecular and developmental aspects. *J Mar Sci Technol.* 37–53.
- Pucciarelli S, Parker SK, Detrich HW, Melki R. 2006. Characterization of the cytoplasmic chaperonin containing TCP-1 from the Antarctic fish *Notothenia coriiceps*. *Extremophiles* 10(6):537–549.
- Ramirez-Llodra EZ, et al. 2010. Deep, diverse and definitely different: unique attributes of the world's largest ecosystem. *BioGeosciences* 7(9):2851–2899.
- Rey C, Guéguen L, Sémon M, Boussau B. 2018. Accurate detection of convergent amino-acid evolution with PCOC. *Mol Biol Evol.* 35(9):2296–2306.
- Reynolds CR, Islam SA, Sternberg MJ. 2018. EzMol: a web server wizard for the rapid visualization and image production of protein and nucleic acid structures. *J Mol Biol.* 430(15):2244–2248.
- Saarman NP, Kober KM, Simson WB, Pogson GH. 2017. Sequence-based analysis of thermal adaptation and protein energy landscapes in an invasive blue mussel (*Mytilus galloprovincialis*). *Genome Biol Evol.* 9(10):2739–2751.
- Sato H, Nakasone K, Yoshida T, Kato C, Maruyama T. 2015. Increases of heat shock proteins and their mRNAs at high hydrostatic pressure in a deep-sea piezophilic bacterium, *Shewanella violacea*. *Extremophiles* 19(4):751–762.
- Schymkowitz J, et al. 2005. The FoldX web server: an online force field. *Nucleic Acids Res.* 33(Web Server):W382–W388.
- Siebenaller J, Somero GN. 1978. Pressure-adaptive differences in lactate dehydrogenases of congeneric fishes living at different depths. *Science* 201(4352):255–257.
- Siebenaller JF. 2010. Effects of the deep-sea environment on invertebrates. In: *Comparative high press biology*. Enfield (NH): Science Publishers. p. 319–341.
- Siebenaller JF, Somero GN. 1979. Pressure-adaptive differences in the binding and catalytic properties of muscle-type (M 4) lactate dehydrogenases of shallow- and deep-living marine fishes. *J Comp Physiol B* 129(4):295–300.
- Smith MD, et al. 2015. Less is more: an adaptive branch-site random effects model for efficient detection of episodic diversifying selection. *Mol Biol Evol.* 32(5):1342–1353.
- Somer L, Shmulman O, Dror T, Hashmueli S, Kashi Y. 2002. The eukaryote chaperonin CCT is a cold shock protein in *Saccharomyces cerevisiae*. *Cell Stress Chaperones* 7(1):47.
- Somero GN. 1990. Life at low volume change: hydrostatic pressure as a selective factor in the aquatic environment. *Am Zool.* 30(1):123–135.
- Somero GN. 1992. Adaptations to high hydrostatic pressure. *Annu Rev Physiol.* 54(1):557–577.
- Somero GN. 2003. Protein adaptations to temperature and pressure: complementary roles of adaptive changes in amino acid sequence and internal milieu. *Comp Biochem Physiol B Biochem Mol Biol.* 136(4):577–591.
- Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30(9):1312–1313.

- Stöhr S, O'Hara TD, Thuy B. 2012. Global diversity of brittle stars (Echinodermata: Ophiuroidea). *PLoS One* 7(3):e31940.
- Stöhr S, O'Hara T, Thuy B, editor. 2020. World Ophiuroidea database. Available from: <http://www.marinespecies.org/ophiuroidea> (accessed August 19, 2020).
- Storz JF. 2016. Causes of molecular convergence and parallelism in protein evolution. *Nat Rev Genet.* 17(4):239–250.
- Suka A, et al. 2019. Stability of cytochromes *c'* from psychrophilic and piezophilic *Shewanella* species: implications for complex multiple adaptation to low temperature and high hydrostatic pressure. *Extremophiles* 23(2):239–248.
- Sun J, et al. 2017. Adaptation to deep-sea chemosynthetic environments as revealed by mussel genomes. *Nat Ecol Evol.* 1:0121.
- Tenaillon O, et al. 2012. The molecular diversity of adaptive convergence. *Science* 335(6067):457–461.
- Tenaillon O, et al. 2016. Tempo and mode of genome evolution in a 50,000-generation experiment. *Nature* 536(7615):165–170.
- Vallin J, Grantham J. 2019. The role of the molecular chaperone CCT in protein folding and mediation of cytoskeleton-associated processes: implications for cancer cell biology. *Cell Stress Chaperones* 24(1):17–27.
- Valpuesta JM, Carrascosa JL, Willison KR. 2005. Structure and function of the cytosolic chaperonin CCT. *Protein fold handbook*. Weinheim, Germany: Wiley-VCH. p. 725–755.
- Venkat A, Hahn MW, Thornton JW. 2018. Multinucleotide mutations cause false inferences of lineage-specific positive selection. *Nat Ecol Evol.* 2(8):1280–1288.
- Wakai N, Takemura K, Morita T, Kitao A. 2014. Mechanism of deep-sea fish α -actin pressure tolerance investigated by molecular dynamics simulations. *PLoS One* 9(1):e85852.
- Wang K, et al. 2019. Morphology and genome of a snailfish from the Mariana Trench provide insights into deep-sea adaptation. *Nat Ecol Evol.* 3(5):823–833.
- Weber AA-T, et al. 2017. Positive selection on sperm ion channels in a brooding brittle star: consequence of life-history traits evolution. *Mol Ecol.* 26(14):3744–3759.
- Welch TJ, Farewell A, Neidhardt FC, Bartlett DH. 1993. Stress response of *Escherichia coli* to elevated hydrostatic pressure. *J Bacteriol.* 175(22):7170–7177.
- Wemekamp-Kamphuis HH, Karatzas AK, Wouters JA, Abee T. 2002. Enhanced levels of cold shock proteins in *Listeria monocytogenes* LO28 upon exposure to low temperature and high hydrostatic pressure. *Appl Environ Microbiol.* 68(2):456–463.
- Wertheim JO, Murrell B, Smith MD, Kosakovsky Pond SL, Scheffler K. 2015. RELAX: detecting relaxed selection in a phylogenetic framework. *Mol Biol Evol.* 32(3):820–832.
- Winnikoff JR, Francis WR, Thuesen EV, Haddock SHD. 2019. Combing transcriptomes for secrets of deep-sea survival: environmental diversity drives patterns of protein evolution. *Integr Comp Biol.* 59(4):786–798.
- Woolley SN, et al. 2016. Deep-sea diversity patterns are shaped by energy availability. *Nature* 533(7603):393–410.
- Yam AY, et al. 2008. Defining the TRiC/CCT interactome links chaperonin function to stabilization of newly made proteins with complex topologies. *Nat Struct Mol Biol.* 15(12):1255–1262.
- Yancey PH. 2020. Cellular responses in marine animals to hydrostatic pressure. *J Exp Zool.* 333(6):398–420.
- Yancey PH, Siebenaller JF. 2015. Co-evolution of proteins and solutions: protein adaptation versus cytoprotective micromolecules and their roles in marine organisms. *J Exp Biol.* 218(12):1880–1896.
- Yin Q, et al. 2011. Molecular cloning of *Litopenaeus vannamei* TCP-1-eta gene and analysis on its relationship with cold tolerance. *Yi Chuan Hered.* 33(2):168–174.
- Zhang Y, et al. 2017. Adaptation and evolution of deep-sea scale worms (Annelida: Polynoidae): insights from transcriptome comparison with a shallow-water species. *Sci Rep.* 7:46205.

Associate editor: Marta Barluenga



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Weber, AA-T; Hugall, AF; O'Hara, TD

Title:

Convergent Evolution and Structural Adaptation to the Deep Ocean in the Protein-Folding Chaperonin CCT.

Date:

2020-11-03

Citation:

Weber, A. A. -T., Hugall, A. F. & O'Hara, T. D. (2020). Convergent Evolution and Structural Adaptation to the Deep Ocean in the Protein-Folding Chaperonin CCT.. *Genome Biol Evol*, 12 (11), pp.1929-1942. <https://doi.org/10.1093/gbe/evaa167>.

Persistent Link:

<http://hdl.handle.net/11343/277779>

File Description:

Published version

License:

CC BY-NC