

Review

Agroecological Strategies to Safeguard Insect Pollinators in Biodiversity Hotspots: Chile as a Case Study

Patricia A. Henríquez-Piskulich ¹, Constanza Schapheer ^{2,3}, Nicolas J. Vereecken ⁴ and Cristian Villagra ^{1,*}

¹ Instituto de Entomología, Universidad Metropolitana de Ciencias de la Educación, Av. José Pedro Alessandri 174, Santiago 7760197, Chile; patricia.andrea.henriquez@gmail.com

² Laboratorio de Sistemática y Evolución, Departamento de Silvicultura y Conservación de la Naturaleza, Universidad de Chile, Santiago 8320000, Chile; constanza.schapheer@ug.uchile.cl

³ Programa de Doctorado en Ciencias Silvoagropecuarias y Veterinarias, Campus Sur Universidad de Chile, Santa Rosa 11315, Santiago 8820808, Chile

⁴ Agroecology Lab, Interfaculty School of Bioengineering, Université Libre de Bruxelles (ULB), Boulevard du Triomphe CP 264/02, B-1050 Brussels, Belgium; nicolas.vereecken@ulb.ac.be

* Correspondence: cristian.villagra@umce.cl; Tel.: +56-999450246

Abstract: Industrial agriculture (IA) has been recognized among the main drivers of biodiversity loss, climate change, and native pollinator decline. Here we summarize the known negative effects of IA on pollinator biodiversity and illustrate these problems by considering the case of Chile, a “world biodiversity hotspot” (WBH) where food exports account for a considerable share of the economy in this country. Most of Chile’s WBH area is currently being replaced by IA at a fast pace, threatening local biodiversity. We present an agroecological strategy for sustainable food production and pollinator conservation in food-producing WBHs. In this we recognize native pollinators as internal inputs that cannot be replaced by IA technological packages and support the development of agroecological and biodiversity restorative practices to protect biodiversity. We suggest four fundamental pillars for food production change based on: (1) sharing the land, restoring and protecting; (2) ecological intensification; (3) localized knowledge, research, and technological development; and (4) territorial planning and implementation of socio-agroecological policies. This approach does not need modification of native pollination services that sustain the world with food and basic subsistence goods, but a paradigm change where the interdependency of nature and human wellbeing must be recognized for ensuring the world’s food security and sovereignty.

Keywords: agroecology; sacrifice zones; Apoidea; water deficit; pesticides



Citation: Henríquez-Piskulich, P.A.; Schapheer, C.; Vereecken, N.J.; Villagra, C. Agroecological Strategies to Safeguard Insect Pollinators in Biodiversity Hotspots: Chile as a Case Study. *Sustainability* **2021**, *13*, 6728. <https://doi.org/10.3390/su13126728>

Academic Editor: Sean Clark

Received: 5 May 2021

Accepted: 8 June 2021

Published: 14 June 2021

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Industrial agriculture (hereafter “IA”) promoted by the Green Revolution has arguably brought about significant increases in food production globally over the past 70 years [1]. These models involve the use of a «technical package» with strong dependency on fossil fuels, which include large-scale monocrop landscapes of improved/selected seeds, increased mechanization, and the incorporation of “external inputs” to enhance plant growth and yield such as the introduction of managed pollinators, synthetic fertilizers and pesticides [2]. Yet these welcomed apparent enhancements in production are also partly responsible for the ongoing massive release of greenhouse gases, the unsustainable use of water and land resources [3], and the contamination of soil as well as both surface and underground water reservoirs by fertilizers and pesticides. Under the current market model this intensive agriculture production is widely requested by “countries with higher developmental level” [4], driving unprecedented amounts of food waste [5]. IA functions at the expense of ever-increasing socio-ecological crises and has been recognized among the main drivers of irreparable biodiversity losses, especially in areas chiefly focused on producing and exporting food crops (i.e., “developing countries”) [6–9]. Biodiversity decline

has been associated with the above-mentioned negative externalities of IA such as habitat loss and fragmentation, pollution, and climate change [10–15]. This reduction in biological diversity is currently jeopardizing ecosystem functions and associated processes (including pollination, water, and nutrient cycling) and putting human wellbeing at risk [10,16–19]. Considering these problems derived from IA and its associated market model, several authors have stressed the need for a paradigm shift in agriculture if we are to meet future food demands while preserving the ecosystems that sustain this food production [20–22].

Agroecology (hereafter “AE”) is considered the most relevant alternative to IA by a wide range of actors involved in food production, such as stakeholders, farmers, scientists, NGOs, and policymakers [23–25]. AE is a scientific discipline as well as a practice that involves the development of diversified farming systems and short supply chains, the promotion of low external input schemes and conservation and regenerative agriculture [2,23,26–30]. As a political movement, AE promotes food security and sovereignty (see Glossary) as an essential human dimension of agricultural transitions in the world’s political agenda [28]. Agroecology goals include the application of ecologically based knowledge to agriculture, with the aim of a sustainable food production while at the same time reducing the environmental impact by spending less energy and resources in the process [31,32], for example by lessening agriculture dependency on the application of external inputs (such as exotic managed pollinators, pesticides, and fertilizers) to maintain food production [23]. AE is considered as the scientific rediscovery of some of the ancient agricultural practices developed and preserved by peasant and native cultures as alternatives to IA around the world [33–35]. AE takes advantage of local biotic components and abiotic conditions found in the agricultural landscape, seeking to match crops with local abiotic conditions and promote beneficial associated organisms [36]; highlighting the value of local knowledge and biodiversity that benefits agricultural production [37]. For instance, AE considers available organisms that improve crop productivity such as pollination, biological control, and decomposition as “resource biota” [38,39]. Through this lens, local diversity is regarded as a natural “internal input” (Figure 1; Figure 2), as opposed to “external inputs” required for IA production, enhancing sustainable food production in agroecologically-managed fields. Internal input provides different ecosystem services and ecological interactions [30,40]. The latter includes pollinators, predators, parasites, and herbivores as well as non-crop vegetation, soil invertebrates, and microorganisms, among other components of local biodiversity helping crop yield [41].




Figure 1. Schematic representation of *industrial agriculture* intensive management. Arrows and positive signs represent favorable influences between elements depicted by icons and titles. “T” ending lines and negative signs symbolize unfavorable impacts. *Landscape homogenization*, the simplification of rural ecosystems that takes place under industrial agriculture, is illustrated with a bulldozer. The application of *external inputs* such as pesticides, GMOs, and managed exotic biological control agents and pollinators, is shown as an operator spraying agrochemicals. Landscape homogenization and external inputs are used to sustain *crop yield* production (represented by various fruits) under industrialized schemes. Nonetheless industrial agriculture’s *landscape homogenization* and *external inputs* are at the same time causing a decline of local biodiversity (e.g., beneficial microorganisms, plants, and animals), which despite not being recognized by industrial agriculture, are contributing to crop yield as *internal inputs* (in calypso lines). This component is illustrated by a slide of soil showing different wild lifeforms and their positive influences by calypso color lines. Among beneficial organisms present in agricultural landscapes are wild pollinators, represented by *native bees*. These are being exemplified in this figure by three specimens (with large to small species) by genera: *Bombus*, *Anthidium*, and *Lasioglossum* native species. Native bees’ positive interactions with crop yield and the remaining internal inputs the other components of this diagram are shown with red lines and arrows. Images in grey highlight detrimental effects on illustrated components (e.g., internal inputs and native bees).




Figure 2. An agroecological strategy (in calypso) to counteract the effects of industrial agriculture managements currently in use for food production at world biodiversity hotspots. This plan is based on four main pillars: (i) *Land sharing, restoration and preservation* (illustrated by a wild plant community along a fruit orchard), (ii) the conservation and promotion of *internal inputs*, (iii) the recovery and development of *localized research and technology* (depicted by a local farmer and a scientist exchanging knowledges), as well as the implementation of (iv) *territorial planning and agroecological policies* (illustrated by a farm area map and pointing hand). We propose that these agroecological pillars may contribute to the survival and performance of native pollinators such as *native bees*. These wild pollinators contribute directly to crop yield as well as indirectly by its influences on the agroecological strategy. We suggest agroecological strategy may be able to buffer current LH and EE from IA, as a start point towards a gradual change towards the implementation of an agroecological food production system; not focused on international market needs only, but on food sovereignty and safety as the base for a true global sustainable food production. Illustrations by Cristian Villagra.

Insects provide several ecosystem services to food production and are considered irreplaceable resource biota under an AE approach [42–44]. For example, it has been established worldwide that native bee species can improve yield and production quality [22,45–49]. Insect pollinators can contribute to food production even in cases where crops are capable of autonomous self-reproduction [50]; as selfing can have detrimental effects on yield and quality due to inbreeding [51,52]. Regarding the economic relevance of insect pollination, it has been found that the productivity of five of the seven main crops of the USA are limited by unavailability of pollinators. The USA annual production value of native pollinator services to these crops has been estimated to be over \$1.5 billion [53,54]. In the case of pollinator species not visiting crops but associated with farmland hedgerow flora and/or wild plant patches, this additional diversity has also been found to contribute to agroecosystem functioning, thus the preservation of these native species must also be considered [54,55]. Pollinating insects are currently contributing to world food production, even though intensive IA practices are paying them back with detrimental effects on their

health and survival (Figure 1; Appendix A [44,56–58]). All this is currently negatively impacting food security [59,60], especially when managed pollinators may not be the solution to present wild pollination losses [61,62].

AE, by contrast, acknowledges the contribution of native pollinators and highlights them as priceless players for a lasting food production strategy [63]. Using the AE framework, this study proposes an agroecological strategy (AES) to face the current decline of native pollinators due to IA applying four fundamental AES pillars (Figure 2). AES can be put in practice in order to counteract known biodiversity threats produced by IA and overcome these negative impacts. AES aims to enhance crop production while maintaining healthy ecosystems and native pollinator diversity [64]. To support AES, we emphasize the relevance and urgency of AE research, practices, and policymaking, especially for areas of the planet currently considered reservoirs of pollinator diversity [65] such as WBHs [66]. The reason for this emphasis is the fact that WBHs often overlap with prime agriculture zones [67]. Therefore, the continuity of food supply may be at stake, considering that biodiversity is a key contributing factor to world agricultural production [4]. Ironically, WBHs' unique assembly of species and ecosystem services are being jeopardized by IA practices and its associated globalized market schemes [6,68,69]. As irreplaceable resource biota, native pollinator biodiversity could contribute to a sustainable long-term food production model [39,70–73], therefore it merits a place in the design of a modern food production system. In this study we briefly outline the main effects of IA on native pollinators and its implication on insect decline. We use Chile as a case study, a fruit-exporting OCDE developing country with considerable endemism that hosts an unprotected biodiversity hotspot [66,74,75]. IA practices in Chile are one of the main causes of environmental and social problems [76,77]. This pattern can also be found in other countries around the world hosting WBHs, where raw material export-oriented economies have been often maintained with disregard for social unrest and damage to the environment produced [78–80]. We discuss to what extent a change in food production procedures which includes AE pillars may contribute to ameliorate native pollinator biodiversity decline in WBHs and highlight the relevance to consider this not as local issues concerning agriculture-oriented economies but as a key matter for world environmental health and food security.

2. Effects of IA on Pollinators

In the last five decades there has been a significant global increase in land use changes for agricultural production purposes [81,82]. As a consequence, landscapes on Earth have been simplified and homogenized [83–86]. This is concerning as both human-managed and natural ecosystems rely on their biodiversity for the provision of diverse services that allow their functioning [87–89]. Industrialized agriculture manages agroecosystems through the constant application of external inputs with the goal to maximize the production of commodities based on a small variety of crop species, mostly to supply to international food market demands [6,90]. As was previously mentioned, this highly industrialized agribusiness is conducted largely in underdeveloped areas of the planet at the expense of reducing biodiversity, soil, and water sources quality as well as the wellbeing of workers and local communities [91–98]. For instance, pesticides and fertilizers are among several indispensable external inputs needed for the maintenance of IA goals. These are often applied in massive amounts to fields in order to attain high productivity [13,39], with disregard for the negative effects on biotic and abiotic components of these managed habitats [99,100], including the resources needed by different insect species to complete their life cycle (e.g., nesting materials, resting refuge, egg laying, and suitable larval development microhabitats) [101]. Industrialized agriculture has been recognized among the main threats to insect pollinators [58,102,103]. Great reductions in pollinator populations reported have been attributed to the IA practices for food production [9,44,104–108], causing a general decline in native pollinator richness and visitation rates not only in surrounding patches of native vegetation but also in croplands [109,110]. In this section we summarize how landscape changes as well as the incorporation of external inputs by IA affect native

insect pollinators, including native bees, drivers that act together and additively under an intensive agricultural scheme (Figure 1) [111].

2.1. Landscape

Landscape changes due to intensive agriculture may negatively impact pollinators [9,108,112], by changing their composition (percentage of natural/semi-natural habitats in the landscape) and/or configuration (i.e., patch density and interpatch connectivity) [113]. The effects on native pollinators will also depend on species-specific traits of these insects and the landscape context [114–122]. For example, nesting resource availability seems to explain 61% of the variation found in different nesting guilds such as ground nesters, pre-existing cavity nesters, carpenters, hollow stem nesters, and cleptoparasite bee species [123]. Thus, the effects of IA may cause both overall decline and community structure alternations.

IA farms are characterized by large-scale crops isolated from natural and/or semi-natural habitats, lacking enough floral and nesting resources as well as decreasing production and survival of insect pollinator offspring [124]. These industrialized croplands typically harbor low insect pollinator richness and abundance [125], reducing pollination services and functional diversity [115]. Vulnerable species have been found to be the most affected under these circumstances [126], losing millions of years of plant–pollinator evolution in the process [127]. Landscape homogenization (hereafter “LH”; Glossary; Figure 1; Figure 2), appears to be an important driver of IA effects on biodiversity [83]. This might be linked to agroecosystems’ reduction of natural habitat patches and/or natural habitat elements [115], decrease of available resources needed for the different components of biodiversity [128] and the loss in connectivity of farmland to natural remnant patches [120,129]. LH also affects mutually beneficial interactions between flowering plants and insect pollinator communities [130]. Pollinator diversity in agricultural habitats under LH might end up being replaced by the few species able to survive these depauperated conditions, leading to further biotic homogenization [131]. This is represented in Figure 1 by native bee specimens pictured in grayscales. The reduced surviving pollinators that remain may not guarantee the delivery of sufficient pollination services, both for human-managed and natural ecosystems [132]. For instance, coevolved associations between native insect pollinators and functionally specialized plants may become at risk of pollen limitation due to LH [133]. This evidence suggests that not only native pollinators currently visiting crop plants must be the focus of concern due to current agricultural practices, but also the whole wild bee guild that may also contribute to the maintenance of local plant diversity near agricultural landscapes [134].

2.2. External Inputs

Because IA simplifies landscapes and their biological diversity, hampering their contribution to agricultural production [13,135], it needs to incorporate “external inputs” (“EI” in Figure 1; Figure 2) to replace the lost regulatory and supporting ecological services in modified landscapes otherwise provided by internal inputs (II) [39]. External inputs include abiotic and biotic factors. For example, chemical formulations such as fertilizers and pesticides are often used along with modified seeds (e.g., herbicide-tolerant crops) capable of enduring these applications while most local biodiversity cannot [100]. IA also introduces exotic managed organisms to provide pollination and biological control. Both pesticides and the use of managed pollinators have been regarded as main external inputs responsible for the decline in native pollinators [136]. Below we detail this evidence.

2.2.1. Pesticides

Pollinators under IA food production plans are exposed to multiple pesticides [137], which have demonstrated deleterious effects in their nervous system, behavior, and cognition as well as their development, reproduction, and overall survival [138–147] (Appendix A). It has also been suggested that sublethal exposure to pesticides may pro-

duce immune suppression in pollinators [148], increasing their susceptibility to pathogens. Recent evidence regarding epigenetic inheritance has demonstrated that pesticides drive pathological alterations in insect pollinators [149], while in target organisms it has been reported the development of IA promotes epigenetic transgenerational resistance against pesticides [150]. Thus, while pesticide detrimental impact may last several generations on non-target organisms, their efficacy on pest species may be reduced as they became immune to their effects [151].

Pesticides reduce the richness and abundance of pollinators and other beneficial native insects [152–154], resulting in mid- and long-term declines and higher extinction rates, whether they forage in treated crops or not (Figure 1) [25,155,156]. Pesticide exposure routes are correlated with the different materials these insects need to complete their life cycles (e.g., nesting and food resources) [157–159]. Pesticide residues have been found in food items and substrates used by target and non-target insects [160–165]. This impairs the delivery of pollination services, reducing pollen collection efficiency and affecting crop yield [166–168]. It has also been demonstrated that native pollinators respond differently to pesticide exposure compared to managed pollinators such as honeybees [169], and in some cases they are more susceptible to their toxic effects [170,171]. Pollinator species may have different responses to pesticides (Appendix A), making it difficult to predict the adverse consequences of these chemicals on pollination services [172,173]. The availability of this kind of data for every species seems unfeasible in the short-term, and thus species-specific traits (such as nesting behavior and sociality type) could be used as proxies to predict pesticide response [116,158,174]. While there is a sustained use of large amounts of pesticides in IA schemes [67], claims of a reduction of their environmental damage have been questioned by researchers. For example, recent reports considering the toxic effects of several pesticides for eight non-target species groups revealed a noticeable increase in the toxicity of applied insecticide over the last 25 years for both aquatic invertebrates and pollinators [100]. This was mainly attributed to the contributions of pyrethroids and neonicotinoids, respectively. The increase of pesticide toxicity included studies in GM corn crops (towards aquatic invertebrates and pollinators) as well as in GM herbicide-tolerant soybeans, where coexisting plant species were also heavily affected [100]. These updated findings stress the urgency to change how food is being produced, leaving current dependency on these external inputs for the sake of the survival of pollinating insects and human health.

2.2.2. Managed Pollinators

Regarding this external input largely use in IA schemes to secure crop pollination, most studies have reported negative effects on native pollinators due to the introduction/spread of exotic competing managed bee species (e.g., *Apis mellifera*, *Bombus terrestris*) in agroecosystems [9,175–177]. Managed bees affect the development and reproduction of native bee species that are close to their colonies [178]. For instance, sometimes managed bees mate with local species, resulting in inviable hybrids [179,180]. EI pollinators introduced under IA management might also compromise food and nesting resources available to other native insect pollinators through competition [181]. When managed bees become naturalized outside of their native range they can adapt easily to varied nesting substrates, potentially being less susceptible to nesting site shortages [182], and overcoming this shortage by usurping closely related species' nests [183]. In the presence of greater floral abundance, the number of managed bees visiting floral species is higher than those of pollinators [184], potentially outcompeting them [185]. They are also able to amass a great amount of provisions rapidly [186], possibly depleting resources for other native insects [187].

Pathogen spillover might also be of concern in this context; managed bees are usually social insects and given their behavior could be more likely to host and spread pathogens [188]. These exotic pollinators are able to transport and spread pathogens to flower species while visiting [189]. These then are transmitted to other wildflower

visitors, including native pollinators [190]. Although pathogens have been indicated as one of the drivers of lower pollinator abundance [191], their impact on native and introduced bee species seems to be so widely distributed that is difficult to pinpoint the direction of these spillovers [192]. Nonetheless, this is a recognized source of deterioration of native pollinator wellbeing.

3. IA and AE in Biodiversity Hotspots: Chile, a Case Study

Biodiversity hotspots are highly endemic biogeographic regions threatened by human activity [66]. The Neotropical region includes several of these areas, hosting an outstanding diversity and richness of native pollinators [9]. This area of the planet produces a considerable portion of food crops by IA and it has been regarded as the zone which has suffered one of the greatest declines in biodiversity and ecosystem services [4,80]. Chile includes in its territory almost an entire hotspot, named the “Chilean Winter Rainfall-Valdivian Forests”. This consists of several biomes hosted within the Chilean Matorral and the Valdivian temperate rainforest [66,193]. The former could be considered a WBH and largely overlaps with IA food production areas [194,195]. Unfortunately, only 1.8% of Matorral land is under the Chilean national protection program [196]. The Chilean Matorral is also a region that hosts an important bee species diversity with elevated endemism [65,197].

Chile has subscribed to environmental treaties to know, conserve, and restore its biodiversity as well as reforest endangered areas [198–201], but so far there are no territorial management plans that aim to make agricultural production compatible with biodiversity conservation (Appendix A). This has resulted in a significant loss of natural habitats in a few decades [202,203]. National records report that approximately 70% (12,900,682 ha) of the land used for agriculture, livestock, and plantation forestry is within the “Chilean Winter Rainfall-Valdivian Forests” hotspot [204]. This hotspot holds an area of 30,000,000 ha, which means that nearly 43% of it has already been replaced by these production schemes [193]. Even more concerning is that habitat loss rate in this hotspot keeps growing [205].

Agricultural practices in Chile are deeply rooted in export-oriented IA models [67,206], directly attributed to the economic liberalization policies mandated by the military dictatorship after 1973 [207]. Measures established by force during this period included the privatization of the public sector, resulting in the concentration of agricultural land in the hands of a few and a significant exploitation of natural resources to supply international markets [208], creating a globalized and capitalized commercial IA scheme at the expense of neglecting and marginalizing small farmers and indigenous people [209,210].

The rediscovery of AE alternatives in Chile began as a reaction towards the economic crisis that unfolded immediately after the application of Milton Friedman’s neoliberal policies in the early 1980s, due to an exponential increase in rural poverty and abandonment of the urban and rural working classes [69,211]. Chilean AE advances were made by a small number of NGOs, small farmers, and academics [35]. Although some of these developments were recognized by the Food and Agriculture Organization of the United Nations [212], only in rare cases have AE practices been adopted by corporations and promoted by policy makers, and at present these practices are not used on a productive scale in Chile [69,213].

3.1. Pesticides

Around 9.6% of the pesticides approved by the Chilean government [214] have been already banned by the European Union (Appendix A), one of the main consumers of Chile’s fruit exports [67]. Most of these pesticides are highly toxic, with demonstrated negative effects on bees at sublethal doses (Appendix A). For example, while neonicotinoids (e.g., clothianidin, imidacloprid, thiamethoxam) are being questioned by experts around the world and have restricted use in Europe due to harmful effects on native and managed bees [98,156,215–218], they are widely used in Chile due to an alleged “absence of proof in the country of their negative effects” [219,220]. This is concerning given the chemical behavior of these pesticides, as these widely used formulations are adsorbed by mineral

clays and organic matter that form agricultural volcanic ash-derived soils [221], damaging biodiversity as a consequence [222], and most likely affecting native bee species directly, as nearly 70% of Chile's wild Apoidea nest in soil substrates [223]. Despite the aforementioned issues, current regulatory protocols for the approval of new formulations and maintenance of pesticide use in Chile have not been updated based on current scientific knowledge and do not require the development of local science-based risk-assessments over biodiversity for their approval for IA use [224].

3.2. Managed Pollinators

External biotic inputs in Chile are already impacting the environment; the main exotic bumblebee species commercially used for providing pollination services, the buff-tailed bumblebee *Bombus terrestris* Linnaeus, 1758 and *B. ruderatus* Fabricius, 1775 have rapidly replaced the Patagonian giant “moscardón” bumblebee *Bombus dahlbomii* Guérin-Ménéville, 1835 [225–229] (Figure 3). *B. dahlbomii* was a source of medicinal honey and considered a sacred being by Mapuche, one of the First Nations people of Chile [230]. Scientists have demonstrated that introduced bumblebee species are displacing native *B. dahlbomii* in Chile and Argentina, colonizing natural areas in most of the southern cone of South America, and have urged authorities to ban the imports of these IA-managed pollinators [231,232]. Even though this is a concerning situation, government policy still allows the importation of buff-tailed bumblebees for IA crop pollination in Chile [231].




Figure 3. Giant bumblebee: *Bombus dahlbomii* (Hymenoptera), native from Chile and Argentina, legitimately visiting blueberry flowers in November 2015, Villarica, X Region, Chile (scale: 1 cm). This species has been categorized as “endangered” by the IUCN Red List. Photography by Marianela Castillo Arias.

In the Mediterranean region of central Chile, a bee biodiversity hotspot [65], avocado orchards have been recorded to be profusely visited by managed *A. mellifera*, while five native bees species have also been reported visiting this crop [233]. Although this finding was proposed as a demonstration of the compatibility of IA avocado production with native bee biodiversity, these observations were conducted through one-season focal observations and with no additional collection methods or control of native vegetation or wild bee abundance comparisons. Scarce and often preliminary local research in combination with fast-paced habitat loss in Chile paints a concerning picture for pollinators and their ecosystem services in the agricultural production canvas. Chile probably hosts around

800 bee species, with more than 450 species described and 70% endemism [197]. Very little research has been published regarding native insect performance as pollinators for native and crop plant species. For example, in the case of the endangered *B. dahlbomii* [234], this native Apidae has been described as a possible pollinator of greenhouse tomatoes [235] and has been seen visiting blueberry and avocado orchards [233]. Considering this evidence and the research from neighboring countries [236–238], it seems likely that most native insect pollinators may already be pollinating crops of economic importance. The knowledge of wild bee species association with native plants is largely incomplete [197,239]. Chile may hold an irreplaceable pollinator workforce in its native bee pollinators, contributing both to crop yield and the preservation of unique biomes, nonetheless they are threatened by intensive IA production and neglected by government policy makers. This highlights the unsuitability of Chile's current agricultural production and market and jeopardizes the mid- and long-term contribution of this country to the production of fruit commodities and to its own resilience against future environmental and food crises. In the following section we develop our proposal to face these issues and be able to protect pollinators in agriculturally oriented WBHs like Chile.

4. Protecting Pollination: Strategies for the Future

Human practices, including agriculture, need to return within the limits that keep our planet habitable [89,240], for the sake of our own species and all living organisms [241,242]. Countries with invaluable biodiversity need to rethink critically the way they are doing agriculture and reevaluate local and native sustainable practices [243,244]. Understanding that native pollinator species are unique “resource biota” (see Glossary already contributing to current crop yield is to be aware of a strategic advantage compared to agriculture food production in non-WBH regions. Native pollinators are part of AE internal inputs that cannot be replaced by IA technological packages or external inputs [245]. Coexisting with our threatened local biodiversity (i.e., internal inputs) and valuing its cultural and biological wealth within productive ecosystems will protect the future of pollination services as well as contribute to food security and sovereignty. Here we focus on the development of an agriculture schemes in WBHs considering native biodiversity, and compile a strategy summarized in four pillars based on agroecological thinking as well as First Nations' knowledge: (1) sharing, restoring and protecting the land; (2) local biodiversity as fundamental AE internal inputs contributing to sustainable agriculture food production and pollinator protection; (3) the need for recovering local knowledges and developing localized research and technology; and (4) territorial planning and the implementation of AE policies (Figure 2).

4.1. Sharing, Restoring, and Protecting the Land

Natural ecosystems are far from simple, and to achieve sustainable agriculture there is a need to maintain their complexity [40]. Polycultures and florally diverse environments have been found to support native pollinator diversity due to a continuous supply of food resources [246]. Agricultural practices need to consider that pollinator functional diversity relies on these native habitats and that biodiversity hotspots by definition are already threatened, thus need to be considered with special care when conducting productive and extractive activities. A sustainable complex landscape matrix is needed to protect hotspots and ensure the delivery of pollination services to crops. This pillar should integrate restoration and protection of large areas of natural habitat and restoration of native land patches within agroecosystems to increase habitat quality (i.e., land sharing) [247]. Pollination services delivered by native insects have been shown to rely strongly on their proximity to natural habitats [109,248,249]. Protected natural areas host higher biodiversity [250] but are not enough to sustain ecological stability [251]. To achieve stability, habitats that have been altered by human activities, including urban zones and areas utilized for productive activities, need to be restored as much as possible [252], leading to effective conservation outcomes by assessing their coverage (i.e., the number and types of species included within

their limits) and management [253]. Restoring native patches of anthropized land improves habitat quality within agroecosystems, maintaining and securing native insects [254]. Native patches buffer the negative effects of pesticide application on pollinators [153,255], offer greater flower diversity and nesting sites [256] and are correlated with higher pollinator density [257]. In farmlands these patches also serve as wildlife corridors [114,181,258–260], promoting heterogeneous landscapes [261] and stabilizing crop pollination [262]. These patches could be implemented at field edges and should have mixed native plants with partial overlap in floral phenology to provide resources for bees during the whole flowering season [256]. Pollinators benefit from florally diverse environments due to a continuous supply of food resources [246], which are critical for ensuring their reproduction [124]. The size of these patches could be dependent on the crop type that they surround, and research should be carried out to define the appropriate cost-effective sizes within specific agroecosystems [101,263].

4.2. *AE Internal Inputs for Sustainability and Pollinator Protection*

Among the core principles of AE science and practice is the preservation and use of local diversity as natural inputs contributing to crop yield [264]. This approach also advocates for food sovereignty while reducing the negative effects of agriculture on the environment and society [265]. Monocultures, organic or not, reduce the functional diversity of pollinators [115]. Under an agroecological strategy (AES), biodiversity is incorporated into agroecosystems to mimic natural ecological processes [28] (Figure 2). With higher biodiversity, agroecosystem inner complexity grows and reduces the dependence of crops on destructive external inputs, allowing the system to maintain its own soil fertility, productivity, and protect itself from pests [266], benefiting insects and attracting pollinators [101]. All this allows native pollinators to visit crops safely and thrive in an agroecosystem with food and nesting resources free of pesticides. This higher pollinator biodiversity could even reduce the need to incorporate large numbers of managed pollinators within crops as additional external input. Nonetheless, this falls short of defining AE, as not only are academic, political, and cultural perspectives tightly knitted to this model, AE places small farmers and local knowledge as the key for food sovereignty [267] and does not agree with the new Green Revolution approach, which seeks to perpetuate an IA system for food production [268]. Instead, AE focuses on the dissemination of knowledge from farmer to farmer based on their historical backgrounds and on reviving their ancestral farming roots [269], strengthening communities and allowing them to become autonomous, securing local food production [268]. Mexican and Bolivian farmers are examples of how traditional low-intensity agriculture allows native bee species to provide successful pollination service [270,271]. There is no need for a new Green Revolution, as social vulnerability and income inequities are the main cause of hunger [5]. AES, summarized in this review, aim to protect pollinators not only by its effects in agroecosystems, but also by reducing poverty and improving people's livelihoods, by both recovering local knowledges and developing local research technologies as well as implementing territorial planning and AE policies considering the needs of local communities (explained further in following sections, Figure 2) [32]. People can only protect or be concerned about biodiversity and its conservation once their basic needs have been met. Thus, the world does not need more food commodities to be traded globally; it needs equal access to nutritive food and production not focused only on market and profits [272,273].

4.3. *Localized Research and Technology*

IA is leading a steady biodiversity decline and exceeding the planetary boundaries that allow humans to survive on Earth [89]. The IA production and market scheme keeps low-income WBH countries of the world relying on the import of technological packages and depending on globalized markets to achieve their productivity goals. Technological packages should not be imported without knowing their consequences to ecosystems, local communities, and economies [274,275]. Critical knowledge gaps still exist regarding

taxonomy, ecosystem services, and socio-ecological vulnerability in order to implement production alternatives considering native pollinators [276]. This is especially urgent in WBH countries risking their biodiversity, food sovereignty, and human wellbeing [277].

Localized studies need to be conducted in regions where biodiversity knowledge is scarce, and nature is heavily under threat due to industrialized food production activities [80]. As a starting point it is necessary to fill the current knowledge gap on species and their ecological associations [278]. There are still a great number of organisms and ecological interactions left to describe [279], largely in WBH zones. Taxonomy is one of the foundations of the applied sciences. If species have not been described, it becomes challenging to understand how they respond to ecological changes and be able to monitor them [280]. This is especially urgent for insects, a group underrepresented in conservation research [281] and under global decline [8]. Research in WBH countries is also key to assess native pollinator contribution to crop and native plant species reproduction, their nesting needs, and different behaviors. Pollinator species have diverse life histories and traits, responding differently to the same threats [83,114]. Assessing how pollinators respond to potential dangers will allow for the modeling of proper AE production programs. To illustrate the urgent need of information we use the Apoidea, highly charismatic native pollinators. It has been estimated the number of native bee species in the Neotropical Region would be above 15,100, stressing that current knowledge on actual species richness would represent roughly one third of this total [9]. This is worrisome considering the high rates of biodiversity and ecosystem services losses reported for this region of the world [80], largely composed of WBHs focused on agriculture exports [4]. This may imply the potential extinction of many pollinators before even their description, “Centinelan Extinction” [282], and its neglect represents a threat to both local and global food security [283]. Therefore, the local study of bee biodiversity and conservation in these regions must be a global concern. Local farmers and first nations have been recognized as “local knowledge holders” for already possessing the understanding regarding their pollinators and their pollination services in their local food production [284–286]. This wisdom needs to be recovered and applied, as they are key to implement and ensure AES allowing a gradual transition towards a sustainable global food production scheme (Figure 2) [287].

Research will improve our understanding on how insect pollinators respond to agricultural practices in WBH countries and provide alternatives with the goal to advance towards the sustainability of socio-ecological systems, allowing for the development of AE tools and technology as part of the production chain as well as conservation in several food and plant-derived goods needed by our species. Developing local AE knowledge will not only protect biodiversity (e.g., native pollinating insects) and agricultural productivity but also reduce the dependency on IA external drivers and inputs and contribute to the coupling of ecosystems and human wellbeing [277]. With this design, pollinator conservation will not be considered a trade-off against agroecosystems or society but as a partnership for our coexistence.

As was already explained, AES for WBHs must consider the political and cultural perspectives along with the research program. Therefore, the rediscovery of AE must link human wellbeing and ecosystem integrity, thus the collecting of information about the ecological vulnerability of pollination services needs to be coupled with gathering information on social inequalities in this food-producing WBH [277], as a link between economic vulnerability and biodiversity loss has been demonstrated [288]. Integrating all this will allow the development of local AE research and technology in consideration of the societal and ecological conditions of different WBH regions of the planet [276,289,290], providing AE schemes for each socio-agroecosystem [37].

4.4. Territorial Planning and AE Policies

World biodiversity hotspots are strongly threatened by the loss of their species and resource depletion (e.g., water scarcity) due to IA business, currently representing sacrifice zones that provide food and goods to global markets, so the developed side of the world can

“go green” [6]. This needs to change. AE’s local biodiversity “internal inputs” such as native pollination services [73] cannot be labeled as commodities (e.g., “natural capital” [73]), as its “exchange” threatens the sustainability of food production and commerce⁴. This is likely currently happening in a “Centinelan” pollination consumption (not a “trade”), as native bees cannot be replaced or recovered once species go extinct. Moreover, there is not a fair planetary-level exchange and interdependency between WBH exporters and international food commerce, as the resulting benefits have been demonstrated to be distributed globally in a both socially and economically unequal way [4]. For instance, in Chile IA is coupled with sustained social inequalities and unrest, local communities driven to unsanitary water deficit and unique biomes shrinking as IA expands, leading pollinating species to decline before having a chance to be studied [207,211,291]. These are the challenges policy makers need to face; if we want to keep the remaining biodiversity of native pollinators in food-producing countries, intensive industrialized agriculture schemes must be first buffered by AES and gradually replaced by true sustainable food production [9,72].

In order to translate knowledge into policies, first the gathering of information needs to be supported. Science and local knowledge holders can provide a roadmap to make well-informed decisions (Figure 2), but their work needs to be properly funded and listened to [292]. These policies can provide the data science and technology need to assess, propose, and apply the best cost-effective strategy for pollinator conservation, food security, and sovereignty [293]. This is already happening in main food consumer countries of the European Union and the United States [8]. Unfortunately, this is not true for most WBH countries [80], where not taking the steps in this direction will have global consequences.

Strong environmental governments will be required in order to change IA schemes and prioritize the conservation of native pollinators and wildlife. Ecosystems, especially those belonging to biodiversity hotspots, need to be within an international legal framework of protection that starts by recognizing the context-specific complexity of agricultural systems and the irreplaceable relevance of local diversity, both biological and cultural [286]. Local deterioration of biodiversity due to extractivism has global consequences on the health of the Earth’s system and food security [89]. Small-scale farming applying AE schemes, such as that proposed in this work, must be prioritized in WBH [69]. Agricultural businesses should be required to follow AES and sustainability standards [273], including coherence with crop and climatic conditions of local biomes, diversified farming, and to prioritize the use of AE’s internal inputs. As a complement, rural and urban public awareness policies and AE education must be considered to provide tools towards conservation and food sovereignty [294]. Traditional ecological knowledge of local agricultural practice and native pollinators must be outreached to the public and applied, preferring small, diversified AE farms instead of large monocrop IA. Moreover, urban AE initiatives and native plant gardening must be promoted as additional patches for native reforestation [33,273,295]. All these urgently need to be assessed and overseen, to ensure sustainable management practices and the conservation of biodiversity [213].

Agroecological management reduces the need for pesticide use and their undesirable consequences (Appendix A), which is an opportunity for WBH countries to ban harmful pesticides, already done in main food consumer countries [8]. Given that insect decline is a global threat, taking sustainable measures in richer countries will not make this crisis disappear without a global commitment [60]. WBH governments also urgently need to implement AE-inspired territorial management plans, including the protection of people’s livelihoods over large corporately owned agricultural areas (e.g., in Chile watering avocado orchards owned by a few cannot hamper entire communities’ access to water).

Developed main food consumer countries need to consider that being climate neutral at the expense of importing food crops from underdeveloped countries does little to solve the negative effects of IA and completely ignores that the loss of ecosystem services will not make distinctions between geopolitical borders [296]. When trading with other nations, developed countries need to have policies that hold the same standards of sustainable production (including bans on GMOs and pesticides) as those applied to their own countries,

and not insist on requiring “yield increases in many low-income countries” [297]. These low-income areas are often also world reservoirs of biodiversity (including pollinators). To consider WBS as sacrifice zones, for the sake of meeting current market needs, are putting in peril not only biodiversity itself, but also global food security and Earth system health [6].

5. Conclusions

A new deal considering AE approaches must be implemented globally, considering WBH as key areas both for the preservation of native pollinator biodiversity and rights and wellbeing of local communities. The implementation of agroecological strategies in WBHs as starting point and buffer for IA may facilitate the transition towards a true sustainable food production. AES will improve our understanding of ecological dynamics in agroecosystems, allowing sustainable development over time, ensuring local development and food sovereignty of WBH, for the sake of keeping native pollinator biodiversity and the wellbeing of the whole planet [88,89].

Author Contributions: Conceptualization, C.V., C.S., P.A.H.-P. and N.J.V.; investigation, C.V., C.S. and P.A.H.-P.; resources, C.V. and C.S.; data curation, C.V., C.S. and P.A.H.-P. writing—original draft preparation, C.V., C.S. and P.A.H.-P.; writing—review and editing, C.V., C.S., P.A.H.-P. and N.J.V.; visualization, C.V. project administration and funding acquisition, C.V. and C.S. All authors have read and agreed to the published version of the manuscript.

Funding: Cristian Villagra was supported by DIUMCE 02-2019-PGI and NGS-64895T-19. Constanza Schapheer was funded by Rufford Booster Grant 29177-B. MDPI’s APC was partially supported by DIUMCE UMCE.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: We would like to thank the three anonymous reviewers who provided insightful comments on our manuscript. We are also grateful with Marianela Castillo Arias for providing an original photograph of *B. dahlbomii* visiting blueberries and with Lafayette Eaton for his thoughtful comments and detailed native English speaker revision of our work.

Conflicts of Interest: The authors declare no conflict of interest.

Glossary

Agroecology	Agronomic discipline focused on an environmental and socially responsible agricultural management. This is achieved through the study of ecological processes inside agroecosystems and the application of this knowledge to agricultural practices.
Agricultural intensification	Agricultural scheme that seeks to maximize crop yield per unit of area using external inputs.
Ecological intensification	Replacement of external inputs used in intensive agriculture (e.g., insecticides, fertilizers, and growth regulators) by ecosystem services to maximize crop yield with minimum environmental impacts.
Ecosystem services	Ecological functions that benefit and are essential for human beings.
Habitat	Environment inhabited by a particular species.

Landscape homogenization	Simplification and reduction of biotic components inside an area of land, which leads to a community of similar functional and structural traits.
Natural habitat	Pristine environment inhabited by native species.
Organic agriculture	agricultural scheme that does not use fertilizers and pesticides.
Patch	Area of land with the same characteristics, regardless of its size.
Seminatural habitat	A native environment partially modified by human activities.
Sustainable agriculture	Agricultural scheme that efficiently maximizes production while protecting the habitat and natural resources from which it depends, safeguarding biodiversity in the long term.

Appendix A

Table A1. Active ingredients with effects in bees still used in Chile and not approved by the European Union.

Use Classification in Chile ¹	Active Ingredient ²	Pesticide Class	Effect ³	Reference
I, R, A	Acephate	Organophosphate	Highly toxic to bees and other beneficial insects.	[298]
H	Atrazine	Triazine	Oxidative stress responses and alteration acetylcholinesterase activity in honeybees; pesticide detected in native bee tissue; found in stored pollen of honeybees; decreases survival, reduces food consumption, and negatively affects behavior in stingless bees.	[137,299–302]
H	Atrazine/S-metolachlor	Triazine/ Chloroacetamide	Oxidative stress responses and alteration acetylcholinesterase activity in honeybees; pesticide detected in native bee tissue; found in stored pollen of honeybees; decreases survival, reduces food consumption, and negatively affects behavior in stingless bees.	[137,299–302]
F, B	Benomyl	Benzimidazole	Moderately toxic to honeybees	[303]
I, R, A	Cadusafos	Organophosphate	Highly toxic to bees	[304]
I, R, A	Carbaryl	Carbamate	Highly toxic to honeybees; found in stored pollen of honeybees	[299,305]
F, B	Carbendazim	Benzimidazole	May alter the immune response and P450-mediated detoxification of honeybees	[306]
F, B	Carbendazim/ Epoxiconazole	Benzimidazole/ Triazole	May alter the immune response and P450-mediated detoxification of honeybees; detected in corbicular pollen loads of honeybees	[306,307]

Table A1. Cont.

Use Classification in Chile ¹	Active Ingredient ²	Pesticide Class	Effect ³	Reference
F, B	Carbendazim/ Mancozeb	Benzimidazole/ Carbamate	May alter the immune response and P450-mediated detoxification of honeybees	[306]
F, B	Tebuconazole/ Carbendazim	Triazole/ Benzimidazole	May alter the immune response and P450-mediated detoxification of honeybees; pesticide detected in native bee tissue	[137,306]
I, R, A	Cartap hydrochloride	Carbamate	Toxic to bumblebees	[308]
I, R, A	Cartap monohydrochloride	Carbamate	Highly toxic to insects	[309]
I, R, A	Chlorfenapyr	Pyrrole	Highly toxic to honeybees	[310]
F, B	Chlorothalonil/ Carbendazim	Chloronitrile/ Benzimidazole	May alter the immune response and P450-mediated detoxification of honeybees; found in stored pollen of honeybees	[299,306]
F, B	Copper 8-quinolinolate/ Carbendazim	Organometallic compound/ Benzimidazole	May alter the immune response and P450-mediated detoxification of honeybees	[306]
F, B	Copper oxychloride/Dibasic copper sulfate/Iprodione/ Sulphur	Copper salt/Copper salt/Dicarboximide/ Chalcogen	Decrease in honeybees' forager survival; found in stored pollen of honeybees	[299,311]
I, R, A	Diazinon	Organophosphate	Precocious foraging in honeybees; Impaired olfactory learning in honeybees; found in stored pollen of honeybees	[299,312,313]
I, R, A	Fenprothrin	Pyrethroid	Highly toxic to honeybees	[314]
I, R, A	Fenvalerate	Pyrethroid	Highly toxic to honeybees; hazardous to leafcutter bees	[315]
I, R, A	Fipronil	Phenylpyrazole	Highly toxic to honeybees; Impaired olfactory learning in honeybees; toxic to leafcutter bees; pesticide detected in native bee tissue; found in stored pollen of honeybees; causes lethargy, motor difficulty, paralysis and hyperexcitation in stingless bees	[137,299,316–319]
H	Glufosinate-ammonium	Phosphinic acid	Low toxicity in honeybees	[320]
H	Imazamox/Imazapyr	Imidazolinone/ Imidazolinone	Low toxicity in honeybees	[321]
F, B	Iprodione	Dicarboximide	Decrease in honeybees' forager survival; found in stored pollen of honeybees	[299,311]
F, B	Iprodione/Propiconazole	Dicarboximide/ Triazole	Decrease in honeybees' forager survival; pesticide detected in native bee tissue; detected in corbicular pollen loads of honeybees; found in stored pollen of honeybees	[137,299,307,311]

Table A1. Cont.

Use Classification in Chile ¹	Active Ingredient ²	Pesticide Class	Effect ³	Reference
F, B	Iprodione/Sulphur	Dicarboximide/ Chalcogen	Decrease in honeybees' forager survival; found in stored pollen of honeybees	[299,311]
H	Isoproturon	Phenylurea	High mortality in honeybees; detected in corbicular pollen loads of honeybees	[307,322]
I, R, A	Methidathion	Organophosphate	Highly toxic to honeybees; found in beeswax of honeybees	[323,324]
I, R, A	Novaluron	Benzoylurea	Highly toxic to honeybees	[325]
H	Paraquat dichloride	Bipyridylum	Highly toxic to honeybees; changes the size of honeybee oenocytes	[326,327]
H	Paraquat dichloride/Diquat (dibromide)	Bipyridylum/ Bipyridylum	Highly toxic to honeybees; changes the size of honeybee oenocytes	[326,327]
I, R, A	Permethrin	Pyrethroid	Highly toxic to honeybees; disorientation and disruption of normal behavior in honeybees; pesticide detected in native bee tissue	[137,328–330]
F, B	Tebuconazole/ Propiconazole/Permethrin	Pyrethroid	Highly toxic to honeybees; disorientation and disruption of normal behavior in honeybees; pesticide detected in native bee tissue	[137,328–330]
F, B	Procymidone	Dicarboximide	Low toxicity to bees; found in stored pollen and beeswax of honeybees	[299,323,331]
I, R, A	Profenofos	Organophosphate	Highly toxic to honeybees; high mortality in honeybees	[332,333]
H	Saflufenacil	Pyrimidinedione	Low toxicity to honeybees	[334]
I, R, A	Thiocyclam hydrogen oxalate	Oxalate salt	Highly toxic to bees	[335]
I, R, A	Acetamiprid/Novaluron	Neonicotinoid/ Benzoylurea	Highly toxic to honeybees; detected in corbicular pollen loads of honeybees; impaired long-term retention of olfactory learning and increased locomotor activity in honeybees; ataxia in bees; slow to no movements and ataxia in bumble bees and leafcutter bees; occur in sufficient quantities in natural bee food to have adverse effects on bees.	[307,325,336,337]

Table A1. Cont.

Use Classification in Chile ¹	Active Ingredient ²	Pesticide Class	Effect ³	Reference
I, R, A	Dinotefuran	Neonicotinoid	Highly toxic to honeybees; higher number of bouts of behavior in honeybees	[338,339]
I, R, A	Fipronil/Imidacloprid	Phenylpyrazole/ Neonicotinoid	Highly toxic to honeybees; impaired olfactory learning in honeybees; honeybees line up in perfect rows or clusters; pesticide detected in native bee tissue; found in stored pollen of honeybees; honeybees loose postural control and spent more time laying on their backs; inhibited grooming, reduced walking and lower righting reflex in honeybees; increased foraging and homing flight times in honeybees; detected in corbicular pollen loads of honeybees; trembling, excessive grooming, uncontrolled proboscis extension, slow to no movements, ataxia and reduced survival in bumble bees and leafcutter bees; toxic to leafcutter bees; occur in sufficient quantities in natural bee food to have adverse effects on bees.	[137,299,307,317–319,336,339,340]
I, R, A	Fipronil/Thiamethoxam	Phenylpyrazole/ Neonicotinoid	Highly toxic to honeybees; Impaired olfactory learning in honeybees; toxic to leafcutter bees; pesticide detected in native bee tissue; found in stored pollen of honeybees; honeybees loss postural control and spent more time laying on their backs; honeybees spend more time grooming; impaired homing ability in honeybees; hyperactivity, ataxia, excessive grooming, permanent late-onset neuromuscular dysfunction and reduced survival in bumble bees and leafcutter bees; occur in sufficient quantities in natural bee food to have adverse effects on bees.	[137,299,317–319,336,339,341]
F, B	Orthoboric acid/Borax	Inorganic compound/ Inorganic compound	Toxic to honeybees	[342]

Table A1. Cont.

Use Classification in Chile ¹	Active Ingredient ²	Pesticide Class	Effect ³	Reference
F, B	Orthoboric acid/Fenpropimorph/Propiconazole	Inorganic compound/Morpholine/Triazole	Toxic to honeybees; detected in corbicular pollen loads of honeybees; found in stored pollen of honeybees	[299,307,342]
F, B	Picoxystrobin/Cyproconazole	Strobilurin/Triazole	Decreased survival, slight changes in pericardial cells and fat bodies in africanized honeybees; detected in corbicular pollen loads of honeybees	[307,343]
F, B	Tributyltin naphthenate/Permethrin	Organotin/Pyrethroid	Highly toxic to honeybees; found in honeybees and beeswax; associated with winter losses of honeybee colonies; disorientation and disruption of normal behavior in honeybees; pesticide detected in native bee tissue	[137,328–330,344]

¹ A = acaricide; B = bactericide; F = fungicide; H = herbicide; I = insecticide; R = rodenticide; ² Mixed active ingredients were considered not approved with one active ingredient not approved by the EU; ³ Effect can correspond to one or more of the mixed active ingredients. NA = Not Applicable.

References

- Khush, G.S. Green Revolution: Preparing for the 21st Century. *Genome* **1999**, *42*, 646–655. [[CrossRef](#)] [[PubMed](#)]
- Lin, B.B.; Chappell, M.J.; Vandermeer, J.; Smith, G.; Quintero, E.; Bezner-Kerr, R.; Griffith, D.M.; Ketcham, S.; Latta, S.C.; McMichael, P.; et al. Effects of Industrial Agriculture on Climate Change and the Mitigation Potential of Small-Scale Agro-Ecological Farms. *CAB Rev. Perspect. Agric. Vet. Sci. Nutr. Nat. Resour.* **2011**, *6*, 1–19. [[CrossRef](#)]
- Pfister, S.; Bayer, P.; Koehler, A.; Hellweg, S. Projected Water Consumption in Future Global Agriculture: Scenarios and Related Impacts. *Sci. Total Environ.* **2011**, *409*, 4206–4216. [[CrossRef](#)] [[PubMed](#)]
- Silva, F.; Carvalheiro, L.G.; Aguirre-Gutiérrez, J.; Lucotte, M.; Guidoni-Martins, K.; Mertens, F. Virtual Pollination Trade Uncovers Global Dependence on Biodiversity of Developing Countries. *AAAS Sci. Adv.* **2021**, *7*, eabe6636. [[CrossRef](#)] [[PubMed](#)]
- Tscharntke, T.; Clough, Y.; Wanger, T.C.; Jackson, L.; Motzke, I.; Perfecto, I.; Vandermeer, J.; Whitbread, A. Global Food Security, Biodiversity Conservation and the Future of Agricultural Intensification. *Biol. Conserv.* **2012**, *151*, 53–59. [[CrossRef](#)]
- Wilting, H.C.; Schipper, A.M.; Bakkenes, M.; Meijer, J.R.; Huijbregts, M.A.J. Quantifying Biodiversity Losses Due to Human Consumption: A Global-Scale Footprint Analysis. *Environ. Sci. Technol.* **2017**, *51*, 3298–3306. [[CrossRef](#)]
- Green, J.M.H.; Croft, S.A.; Durán, A.P.; Balmford, A.P.; Burgess, N.D.; Fick, S.; Gardner, T.A.; Godar, J.; Suavet, C.; Virah-Sawmy, M.; et al. Linking Global Drivers of Agricultural Trade to On-the-Ground Impacts on Biodiversity. *Proc. Natl. Acad. Sci. USA* **2019**, *116*, 26085–26086. [[CrossRef](#)]
- Wagner, D.L.; Grames, E.M.; Forister, M.L.; Berenbaum, M.R.; Stopak, D. Insect Decline in the Anthropocene: Death by a Thousand Cuts. *Proc. Natl. Acad. Sci. USA* **2021**, *118*, e2023989118. [[CrossRef](#)]
- Freitas, B.M.; Imperatriz-Fonseca, V.L.; Medina, L.M.; Kleinert, A.D.M.P.; Galetto, L.; Nates-Parra, G.; Javier, J. Diversity, Threats and Conservation of Native Bees in the Neotropics. *Apidologie* **2009**, 332–346. [[CrossRef](#)]
- Tscharntke, T.; Klein, A.M.; Kruess, A.; Steffan-Dewenter, I.; Thies, C. Landscape Perspectives on Agricultural Intensification and Biodiversity—Ecosystem Service Management. *Ecol. Lett.* **2005**, *8*, 857–874. [[CrossRef](#)]
- Matson, P.A.; Parton, W.J.; Power, A.G.; Swift, M.J. Agricultural Intensification and Ecosystem Properties. *Science* **1997**, *277*, 504–509. [[CrossRef](#)]
- Medan, D.; Torretta, J.P.; Hodara, K.; de la Fuente, E.B.; Montaldo, N.H. Effects of Agriculture Expansion and Intensification on the Vertebrate and Invertebrate Diversity in the Pampas of Argentina. *Biodivers. Conserv.* **2011**, *20*, 3077–3100. [[CrossRef](#)]
- Tsiafouli, M.A.; Thébault, E.; Sgardelis, S.P.; de Ruiter, P.C.; van der Putten, W.H.; Birkhofer, K.; Hemerik, L.; de Vries, F.T.; Bardgett, R.D.; Brady, M.V.; et al. Intensive Agriculture Reduces Soil Biodiversity across Europe. *Glob. Chang. Biol.* **2015**, *21*, 973–985. [[CrossRef](#)]
- Muñoz-Sáez, A.; Perez-Quezada, J.F.; Estades, C.F. Agricultural Landscapes as Habitat for Birds in Central Chile. *Rev. Chil. Hist. Nat.* **2017**, *90*, 3. [[CrossRef](#)]
- Stanton, R.; Clark, R.G.; Morrissey, C.A. Intensive Agriculture and Insect Prey Availability Influence Oxidative Status and Return Rates of an Aerial Insectivore. *Ecosphere* **2017**, *8*, e01746. [[CrossRef](#)]
- Berka, C.; Schreier, H.; Hall, K. Linking Water Quality with Agricultural Intensification in a Rural Watershed. *Water. Air. Soil Pollut.* **2001**, *127*, 389–401. [[CrossRef](#)]
- Kennish, M.J. Environmental Threats and Environmental Future of Estuaries. *Environ. Conserv.* **2002**, *29*, 78–107. [[CrossRef](#)]

18. Rohr, J.R.; Schotthoefer, A.M.; Raffel, T.R.; Carrick, H.J.; Halstead, N.; Hoverman, J.T.; Johnson, C.M.; Johnson, L.B.; Lieske, C.; Piwoni, M.D.; et al. Agrochemicals Increase Trematode Infections in a Declining Amphibian Species. *Nature* **2008**. [[CrossRef](#)] [[PubMed](#)]
19. Vermeulen, S.J.; Campbell, B.; Ingram, J.S. Climate Change and Food Systems. *Annu. Rev. Environ. Resour.* **2012**, *37*, 195–222. [[CrossRef](#)]
20. Godfray, H.C.J.; Beddington, J.R.; Crute, I.R.; Haddad, L.; Lawrence, D.; Muir, J.F.; Pretty, J.; Robinson, S.; Thomas, S.M.; Toulmin, C. Food Security: The Challenge of Feeding 9 Billion People. *Science* **2010**, *327*, 812–818. [[CrossRef](#)] [[PubMed](#)]
21. Hinrichs, C.C. Regionalizing Food Security? Imperatives, Intersections and Contestations in a Post-9/11 World. *J. Rural Stud.* **2013**, *29*, 7–18. [[CrossRef](#)]
22. Melathopoulos, A.P.; Cutler, G.C.; Tyedmers, P. Where Is the Value in Valuing Pollination Ecosystem Services to Agriculture? *Ecol. Econ.* **2015**, *109*, 59–70. [[CrossRef](#)]
23. Wezel, A.; Herren, B.G.; Kerr, R.B.; Barrios, E.; Gonçalves, A.L.R.; Sinclair, F. Agroecological Principles and Elements and Their Implications for Transitioning to Sustainable Food Systems. A Review. *Agron. Sustain. Dev.* **2020**, *40*. [[CrossRef](#)]
24. HLPE. *Agroecological and Other Innovative Approaches for Sustainable Agriculture and Food Systems That Enhance Food Security and Nutrition*; HLPE: Rome, Italy, 2019; p. 163.
25. Kluser, S.; Peduzzi, P. *Global Pollinator Decline: A Literature Review*; UNEP/GRID Europe: Geneva, Switzerland, 2007; p. 12.
26. Merrill, M.C. Eco-Agriculture: A Review of Its History and Philosophy. *Biol. Agric. Hortic.* **1983**, *1*, 181–210. [[CrossRef](#)]
27. Smith, L.; Williams, A.G.; Pearce, B.D. The Energy Efficiency of Organic Agriculture: A Review. *Renew. Agric. Food Syst.* **2014**, *30*, 1–22. [[CrossRef](#)]
28. Altieri, M.A. *Agroecology: The Science of Sustainable Agriculture*, 2nd ed.; CRC Press Taylor & Francis Group: New York, NY, USA, 2018. [[CrossRef](#)]
29. Kremen, C.; Iles, A.; Bacon, C. Diversified Farming Systems: An Agroecological, Systems-Based Alternative to Modern Industrial Agriculture. *Ecol. Soc.* **2012**, *17*. [[CrossRef](#)]
30. Maran, A.M.; Weintraub, M.N.; Pelini, S.L. Does Stimulating Ground Arthropods Enhance Nutrient Cycling in Conventionally Managed Corn Fields? *Agric. Ecosyst. Environ.* **2020**, *297*, 106934. [[CrossRef](#)]
31. Wezel, A.; Bellon, S.; Doré, T.; Francis, C.; Vallod, D.; David, C. Agroecology as a Science, a Movement and a Practice. *Sustain. Agric.* **2009**, *2*, 27–43. [[CrossRef](#)]
32. Altieri, M.A. Agroecology, Small Farms, and Food Sovereignty. *Mon. Rev.* **2009**, *61*. [[CrossRef](#)]
33. Berkes, F.; Colding, J.; Folke, C. Rediscovery of Traditional Ecological Knowledge as Adaptive Management. *Ecol. Appl.* **2000**, *10*, 1251–1262. [[CrossRef](#)]
34. Holdridge, G.A.; Sarmiento, F.O.; Birch, S.E.P.; Boley, B.; Reap, J.K.; Macdonald, E.A.; Navarro, M.; Hitchner, S.L.; Schelhas, J.W. Chapter 15: Feeding Futures Framed: Rediscovering Biocultural Diversity in Sustainable Foodscapes. In *The Elgar Companion to Geography, Transdisciplinarity and Sustainability*; Frolich, F.O.S., Ed.; Edward Elgar Publishing: Cheltenham, UK, 2020; pp. 235–251. [[CrossRef](#)]
35. Montalba, R.; Infante, A.; Contreras, A.; Vieli, L. Agroecology in Chile: Precursors, Pioneers, and Their Legacy. *Agroecol. Sustain. Food Syst.* **2017**, *41*, 416–428. [[CrossRef](#)]
36. Gurr, G.; Wratten, S.; Altieri, M. *Ecological Engineering for Pest Management. Advances in Habitat Manipulation for Arthropods*; CSIRO: Collingwood, VIC, Canada, 2004. [[CrossRef](#)]
37. González-Chang, M.; Dörner, J.; Zúñiga, F. Agroecología y Sistemas Agrícolas Sustentables. *Agro Sur* **2018**, *46*, 1–2. [[CrossRef](#)]
38. Southwood, R.E.; Way, M.J. Ecological Background to Pest Management. In *Concepts of Pest Management*; Rabb, R.C., Guthrie, F.E., Eds.; North Carolina State University: Raleigh, NC, USA, 1970; pp. 6–29.
39. Altieri, M.A. The Ecological Role of Biodiversity in Agroecosystems. *Agric. Ecosyst. Environ.* **1999**, *74*, 19–31. [[CrossRef](#)]
40. Bommarco, R.; Kleijn, D.; Potts, S.G. Ecological Intensification: Harnessing Ecosystem Services for Food Security. *Trends Ecol. Evol.* **2013**, *28*, 230–238. [[CrossRef](#)] [[PubMed](#)]
41. Altieri, M.A. *Biodiversity and Pest Management in Agroecosystems*; Haworth Press: New York, NY, USA, 1994.
42. Xin, C.; JianJun, T. Utilization of Biodiversity in Agriculture: Today and Tomorrow. *Chin. J. Eco-Agric.* **2013**, *21*, 54–60.
43. Kremen, C.; Chaplin-Kramer, R. Insects as Providers of Ecosystem Services: Crop Pollination and Pest Control. In *Insect Conservation Biology*; Stewart, A.J.A., New, T.R., Lewis, O.T., Eds.; CABI Publishing: Cambridge, MA, USA, 2007; pp. 349–382.
44. Rader, R.; Bartomeus, I.; Garibaldi, L.A.; Garratt, M.P.D.; Howlett, B.G.; Winfree, R.; Cunningham, S.A.; Mayfield, M.M.; Arthur, A.D.; Andersson, G.K.S.; et al. Non-Bee Insects Are Important Contributors to Global Crop Pollination. *Proc. Natl. Acad. Sci. USA* **2016**, *113*, 146–151. [[CrossRef](#)]
45. Costanza, R.; D'Arge, R.; De Groot, R.; Farber, S.; Grasso, M.; Hannon, B.; Limburg, K.; Naeem, S.; O'Neill, R.V.; Paruelo, J.; et al. The Value of the World's Ecosystem Services and Natural Capital. *Nature* **1997**, *387*, 253–260. [[CrossRef](#)]
46. Lautenbach, S.; Seppelt, R.; Liebscher, J.; Dormann, C.F. Spatial and Temporal Trends of Global Pollination Benefit. *PLoS ONE* **2012**. [[CrossRef](#)]
47. Chaplin-Kramer, R.; Dombeck, E.; Gerber, J.; Knuth, K.A.; Mueller, N.D.; Mueller, M.; Ziv, G.; Klein, A.M. Global Malnutrition Overlaps with Pollinator-Dependent Micronutrient Production. *Proc. R. Soc. B Biol. Sci.* **2014**, *281*. [[CrossRef](#)]
48. Gallai, N.; Salles, J.M.; Settele, J.; Vaissière, B.E. Economic Valuation of the Vulnerability of World Agriculture Confronted with Pollinator Decline. *Ecol. Econ.* **2009**, *68*, 810–821. [[CrossRef](#)]

49. Garibaldi, L.A.; Steffan-dewenter, I.; Winfree, R.; Aizen, M.A.; Bommarco, R.; Cunningham, S.A.; Kremen, C.; Carvalheiro, L.G. Wild Pollinators Enhance Fruit Set of Crops Regardless of Honey Bee Abundance. *Science* **2013**, *339*, 1608–1611. [[CrossRef](#)] [[PubMed](#)]
50. Kremen, C.; Bugg, R.L.; Nicola, N.; Smith, S.A.; Thorp, R.W.; Williams, N.M. Native Bees, Native Plants and Crop Pollination in California. *Fremontia* **2002**, *30*, 41–49.
51. Stein, K.; Coulibaly, D.; Stenchly, K.; Goetze, D.; Porembski, S.; Lindner, A.; Konaté, S.; Linsenmair, E.K. Bee Pollination Increases Yield Quantity and Quality of Cash Crops in Burkina Faso, West Africa. *Sci. Rep.* **2017**, *7*, 17691. [[CrossRef](#)]
52. Klein, A.M.; Steffan-Dewenter, I.; Tscharrnke, T. Bee Pollination and Fruit Set of *Coffea Arabica* and *C. Canephora* (Rubiaceae). *Am. J. Bot.* **2003**, *90*, 153–157. [[CrossRef](#)] [[PubMed](#)]
53. Reilly, J.R.; Artz, D.R.; Biddinger, D.; Bobiwash, K.; Boyle, N.K.; Brittain, C.; Brokaw, J.; Campbell, J.W.; Daniels, J.; Elle, E.; et al. Crop Production in the USA Is Frequently Limited by a Lack of Pollinators. *Proc. R. Soc. B* **2020**, *287*, 2–9. [[CrossRef](#)]
54. Venturini, E.M.; Drummond, F.A.; Hoshida, A.K.; Dibble, A.C.; Stack, L.B. Pollination Reservoirs for Wild Bee Habitat Enhancement in Cropping Systems: A Review. *Agroecol. Sustain. Food Syst.* **2017**, *41*, 101–142. [[CrossRef](#)]
55. Albrecht, M.; Kleijn, D.; Williams, N.M.; Tschumi, M.; Blaauw, B.R.; Bommarco, R.; Campbell, A.J.; Dainese, M.; Drummond, F.A.; Entling, M.H.; et al. The Effectiveness of Flower Strips and Hedgerows on Pest Control, Pollination Services and Crop Yield: A Quantitative Synthesis. *Ecol. Lett.* **2020**, *23*, 1488–1498. [[CrossRef](#)]
56. Belsky, J.; Joshi, N.K. Impact of Biotic and Abiotic Stressors on Managed and Feral Bees. *Insects* **2019**, *10*, 233. [[CrossRef](#)]
57. Sánchez-Bayo, F.; Wyckhuys, K.A.G. Worldwide Decline of the Entomofauna: A Review of Its Drivers. *Biol. Conserv.* **2019**, *8–27*. [[CrossRef](#)]
58. Steffan-Dewenter, I.; Potts, S.G.; Packer, L.; Ghazoul, J. Pollinator Diversity and Crop Pollination Services Are at Risk [3] (Multiple Letters). *Trends Ecol. Evol.* **2005**, *20*, 651–652. [[CrossRef](#)]
59. Ellis, A.M.; Myers, S.S.; Ricketts, T.H. Do Pollinators Contribute to Nutritional Health? *PLoS ONE* **2015**, *10*. [[CrossRef](#)] [[PubMed](#)]
60. Van Der Sluijs, J.P. Insect Decline, an Emerging Global Environmental Risk. *Curr. Opin. Environ. Sustain.* **2020**. [[CrossRef](#)]
61. Aizen, M.A.; Arbetman, M.P.; Chacoff, N.P.; Chalcoff, V.R.; Feinsinger, P.; Garibaldi, L.A.; Harder, L.D.; Morales, C.L.; Sáez, A.; Vanbergen, A.J. *Invasive Bees and Their Impact on Agriculture*, 1st ed.; Elsevier Ltd.: Amsterdam, The Netherlands, 2020. [[CrossRef](#)]
62. Cock, M.J.W.; Biesmeijer, J.C.; Cannon, R.J.C.; Gerard, P.J.; Gillespie, D.; Jimenez, J.J.; Lavelle, P.M.; Raina, S.K. The Positive Contribution of Invertebrates to Sustainable Agriculture and Food Security. *CAB Rev. Perspect. Agric. Vet. Sci. Nutr. Nat. Resour.* **2012**, *7*. [[CrossRef](#)]
63. Cepeda-Valencia, J.; Gómez, P.D.; Nicholls, C. The Structure Matters: Bees Visitors of Coffee Flowers and Agroecological Main Structure (MAS). *Rev. Colomb. Entomol.* **2014**, *40*, 241–250.
64. Garibaldi, L.A.; Pérez-Méndez, N.; Garratt, M.P.D.; Gemmill-Herren, B.; Miguez, F.E.; Dicks, L.V. Policies for Ecological Intensification of Crop Production. *Trends Ecology Evolution.* **2019**, 282–286. [[CrossRef](#)]
65. Orr, M.C.; Hughes, A.C.; Chesters, D.; Pickering, J.; Zhu, C.D.; Ascher, J.S. Global Patterns and Drivers of Bee Distribution. *Curr. Biol.* **2021**, *31*, 451–458.e4. [[CrossRef](#)] [[PubMed](#)]
66. Myers, N.; Mittermeier, R.A.; Mittermeier, C.G.; da Fonseca, G.A.B.; Kent, J. Biodiversity Hotspots for Conservation Priorities. *Nature* **2000**, *403*, 853–858. [[CrossRef](#)]
67. Food and Agriculture Organization of the United Nations. FAOSTAT. Available online: <http://www.fao.org/faostat/en/> (accessed on 1 June 2021).
68. Brooks, T.M.; Mittermeier, R.A.; Mittermeier, C.G.; Fonseca, G.A.B.D.A.; Rylands, A.B.; Konstant, W.R.; Flick, P.; Pilgrim, J.; Oldfield, S.; Magin, G.; et al. Habitat Loss and Extinction in the Hotspots of Biodiversity. *Conserv. Biol.* **2002**, *16*, 909–923. [[CrossRef](#)]
69. Altieri, M.A. Una Estrategia Agroecológica En Chile Como Base Para La Soberanía Alimentaria. *Ambient. Desarro. CIPMA* **2010**, *24*, 25–29.
70. Dai, P.; Zhang, X.; Liu, Y. Conserving Pollinator Diversity and Improving Pollination Services in Agricultural Landscapes. *Biodivers. Sci.* **2015**, *23*, 408–418. [[CrossRef](#)]
71. Seitz, N.; van Engelsdorp, D.; Leonhardt, S.D. Are Native and Non-Native Pollinator Friendly Plants Equally Valuable for Native Wild Bee Communities? *Ecol. Evol.* **2020**, *10*, 12838–12850. [[CrossRef](#)]
72. Pérez-Méndez, N.; Andersson, G.K.S.; Requier, F.; Hipólito, J.; Aizen, M.A.; Morales, C.L.; García, N.; Gennari, G.P.; Garibaldi, L.A. The Economic Cost of Losing Native Pollinator Species for Orchard Production. *J. Appl. Ecol.* **2020**, *57*, 599–608. [[CrossRef](#)]
73. Birch, K.; Levidow, L.; Papaioannou, T. Sustainable Capital? The Neoliberalization of Nature and Knowledge in the European “Knowledge-Based Bio-Economy”. *Sustainability* **2010**, *2*, 2898–2918. [[CrossRef](#)]
74. Tognelli, M.F.; De Arellano, P.I.R.; Marquet, P.A. How Well Do the Existing and Proposed Reserve Networks Represent Vertebrate Species in Chile? *Divers. Distrib.* **2008**, *14*, 148–158. [[CrossRef](#)]
75. Plischoff, P.; Fuentes-Castillo, T. Representativeness of Terrestrial Ecosystems in Chile’s Protected Area System. *Environ. Conserv.* **2011**, *38*, 303–311. [[CrossRef](#)]
76. Silva, E. Democracy, Market Economics, and Environmental Policy in Chile. *J. Inter. Am. Stud. World Aff.* **2007**, *38*, 1. [[CrossRef](#)]
77. Ribbe, L.; Delgado, P.; Salgado, E.; Flügel, W.A. Nitrate Pollution of Surface Water Induced by Agricultural Non-Point Pollution in the Pochay Watershed, Chile. *Desalination* **2008**, *226*, 13–20. [[CrossRef](#)]

78. Crook, M.; Short, D.; South, N. Ecocide, Genocide, Capitalism and Colonialism: Consequences for Indigenous Peoples and Global Ecosystems Environments. *Theor. Criminol.* **2018**, *22*, 298–317. [[CrossRef](#)]
79. Bambrick, H. Resource Extractivism, Health and Climate Change in Small Islands. *Int. J. Clim. Chang. Strateg. Manag.* **2018**, *10*. [[CrossRef](#)]
80. Larterra, P.; Nahuelhual, L.; Vallejos, M.; Berrouet, L.; Arroyo Pérez, E.; Enrico, L.; Jiménez-Sierra, C.; Mejía, K.; Meli, P.; Rincón-Ruiz, A.; et al. Linking Inequalities and Ecosystem Services in Latin America. *Ecosyst. Serv.* **2019**, *36*, 100875. [[CrossRef](#)]
81. DeFries, R.; Birkenholtz, T.; Uriarte, M.; Hecht, S.; Grau, R.; Lambin, E.F.; Baptista, S.; Schneider, L.; Lawrence, D.; Rudel, T.K.; et al. Agricultural Intensification and Changes in Cultivated Areas, 1970–2005. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 20675–20680. [[CrossRef](#)]
82. Alexander, P.; Rounsevell, M.D.A.; Dislich, C.; Dodson, J.R.; Engström, K.; Moran, D. Drivers for Global Agricultural Land Use Change: The Nexus of Diet, Population, Yield and Bioenergy. *Glob. Environ. Chang.* **2015**, *35*, 138–147. [[CrossRef](#)]
83. Gámez-Virués, S.; Perović, D.J.; Gossner, M.M.; Börschig, C.; Blüthgen, N.; De Jong, H.; Simons, N.K.; Klein, A.M.; Krauss, J.; Maier, G.; et al. Landscape Simplification Filters Species Traits and Drives Biotic Homogenization. *Nat. Commun.* **2015**, *6*, 8568. [[CrossRef](#)] [[PubMed](#)]
84. Flynn, D.F.B.; Gogol-Prokurat, M.; Nogeire, T.; Molinari, N.; Richers, B.T.; Lin, B.B.; Simpson, N.; Mayfield, M.M.; DeClerck, F. Loss of Functional Diversity under Land Use Intensification across Multiple Taxa. *Ecol. Lett.* **2009**, *12*, 22–33. [[CrossRef](#)] [[PubMed](#)]
85. Reidsma, P.; Tekelenburg, T.; Van Den Berg, M.; Alkemade, R. Impacts of Land-Use Change on Biodiversity: An Assessment of Agricultural Biodiversity in the European Union. *Agric. Ecosyst. Environ.* **2006**, *114*, 86–102. [[CrossRef](#)]
86. dos Santos, J.S.; Dodonov, P.; Oshima, J.E.F.; Martelloc, F.; Santos de Jesuse, A.; Ferreirae, M.E.; Silva-Netof, C.M.; Ribeiroc, M.C.; Garcia Collevatti, R. Landscape Ecology in the Anthropocene: An Overview for Integrating Agroecosystems and Biodiversity Conservation. *Perspect. Ecol. Conserv.* **2021**, *158*, 21–32. [[CrossRef](#)]
87. Mace, G.M.; Norris, K.; Fitter, A.H. Biodiversity and Ecosystem Services: A Multilayered Relationship. *Trends Ecol. Evol.* **2012**, *27*, 19–26. [[CrossRef](#)]
88. Hamilton, C.; Bonneuil, C.; Gemenne, F. *The Anthropocene and the Global Environmental Crisis*; Hamilton, C., Bonneuil, C., Gemenne, F., Eds.; Routledge, Taylor and Francis Group: London, UK; New York, NY, USA, 2015. [[CrossRef](#)]
89. Rockström, J.; Steffen, W.; Noone, K.; Persson, Å.; Chapin, F.S.; Lambin, E.; Lenton, T.M.; Scheffer, M.; Folke, C.; Schellnhuber, H.J.; et al. Planetary Boundaries: Exploring the Safe Operating Space for Humanity. *Ecol. Soc.* **2009**, *14*. [[CrossRef](#)]
90. Karp, D.S.; Rominger, A.J.; Zook, J.; Ranganathan, J.; Ehrlich, P.R.; Daily, G.C. Intensive Agriculture Erodes β -Diversity at Large Scales. *Ecol. Lett.* **2012**, *15*, 963–970. [[CrossRef](#)]
91. Horrigan, L.; Lawrence, R.S.; Walker, P. How Sustainable Agriculture Can Address the Environmental and Human Health Harms of Industrial Agriculture. *Environ. Health Perspect.* **2002**, 445–456. [[CrossRef](#)]
92. Shah, A.N.; Tanveer, M.; Shahzad, B.; Yang, G.; Fahad, S.; Ali, S.; Bukhari, M.A.; Tung, S.A.; Hafeez, A.; Souliyanonh, B. Soil Compaction Effects on Soil Health and Crop productivity: An Overview. *Environ. Sci. Pollut. Res.* **2017**, *24*, 10056–10067. [[CrossRef](#)]
93. Scotti, R.; Bonanomi, G.; Scelza, R.; Zoina, A.; Rao, M.A. Organic Amendments as Sustainable Tool to Recovery Fertility in Intensive Agricultural Systems. *J. Soil Sci. Plant Nutr.* **2015**, 333–352. [[CrossRef](#)]
94. De Roos, A.J.; Blair, A.; Rusiecki, J.A.; Hoppin, J.A.; Svec, M.; Dosemeci, M.; Sandler, D.P.; Alavanja, M.C. Cancer Incidence among Glyphosate-Exposed Pesticide Applicators in the Agricultural Health Study. *Environ. Health Perspect.* **2005**. [[CrossRef](#)]
95. Lerro, C.C.; Beane Freeman, L.E.; DellaValle, C.T.; Andreotti, G.; Hofmann, J.N.; Koutros, S.; Parks, C.G.; Shrestha, S.; Alavanja, M.C.R.; Blair, A.; et al. Pesticide Exposure and Incident Thyroid Cancer among Male Pesticide Applicators in Agricultural Health Study. *Environ. Int.* **2021**, *146*, 106187. [[CrossRef](#)] [[PubMed](#)]
96. Picó, Y.; Alvarez-Ruiz, R.; Alfarhan, A.H.; El-Sheikh, M.A.; Alshahrani, H.O.; Barceló, D. Pharmaceuticals, Pesticides, Personal Care Products and Microplastics Contamination Assessment of Al-Hassa Irrigation Network (Saudi Arabia) and Its Shallow Lakes. *Sci. Total Environ.* **2020**, *701*, 135021. [[CrossRef](#)] [[PubMed](#)]
97. Hussain, S.; Siddique, T.; Saleem, M.; Arshad, M.; Khalid, A. *Chapter 5 Impact of Pesticides on Soil Microbial Diversity, Enzymes, and Biochemical Reactions*, 1st ed.; Elsevier Inc.: Amsterdam, The Netherlands, 2009; Volume 102. [[CrossRef](#)]
98. Hoshi, N. Chapter 12. Adverse Effects of Pesticides on Regional Biodiversity and Their Mechanisms. In *Risks and Regulation of New Technologies*; Matsuda, T., Wolff, J., Yanagawa, T., Eds.; Springer & Kobe University: Kobe, Japan, 2021; pp. 235–247.
99. González-Varo, J.P.; Biesmeijer, J.C.; Bommarco, R.; Potts, S.G.; Schweiger, O.; Smith, H.G.; Steffan-Dewenter, I.; Szentgyörgyi, H.; Woyciechowski, M.; Vilà, M. Combined Effects of Global Change Pressures on Animal-Mediated Pollination. *Trends Ecol. Evol.* **2013**, *28*, 524–530. [[CrossRef](#)] [[PubMed](#)]
100. Schulz, R.; Bub, S.; Petschick, L.; Stehle, S.; Wolfram, J. Applied Pesticide Toxicity Shifts toward Plants and Invertebrates, Even in GM Crops. *Science* **2021**, *372*, 6537. [[CrossRef](#)]
101. Nicholls, C.I.; Altieri, M.A. Plant Biodiversity Enhances Bees and Other Insect Pollinators in Agroecosystems. A Review. *Agron. Sustain. Dev.* **2013**, *33*, 257–274. [[CrossRef](#)]
102. Stavert, J.R.; Pattenmore, D.E.; Gaskett, A.C.; Beggs, J.R.; Bartomeus, I. Exotic Species Enhance Response Diversity to Land-Use Change but Modify Functional Composition. *Proc. R. Soc. B Biol. Sci.* **2017**, *284*. [[CrossRef](#)] [[PubMed](#)]

103. Biesmeijer, J.C.; Roberts, S.P.M.; Reemer, M.; Ohlemüller, R.; Edwards, M.; Peeters, T.; Schaffers, A.P.; Potts, S.G.; Kleukers, R.; Thomas, C.D.; et al. Parallel Declines in Pollinators and Insect-Pollinated Plants in Britain and the Netherlands. *Science* **2006**, *313*, 351–354. [[CrossRef](#)] [[PubMed](#)]
104. Grixti, J.C.; Wong, L.T.; Cameron, S.A.; Favret, C. Decline of Bumble Bees (*Bombus*) in the North American Midwest. *Biol. Conserv.* **2009**, *142*, 75–84. [[CrossRef](#)]
105. Benedek, P. Possible Indirect Effect of Weed Control on Population Changes of Wild Bees Pollinating Lucerne. *Acta Phytopathol. Acad. Sci. Hungaricae* **1972**, *7*, 267–278.
106. Kevan, P.G.; Viana, B.F. The Global Decline of Pollination Services. *Biodiversity* **2003**, *4*, 3–8. [[CrossRef](#)]
107. Eeraerts, M.; Meeus, I.; Van Den Berge, S.; Smagghe, G. Landscapes with High Intensive Fruit Cultivation Reduce Wild Pollinator Services to Sweet Cherry. *Agric. Ecosyst. Environ.* **2017**, *239*, 342–348. [[CrossRef](#)]
108. Kremen, C.; Williams, N.M.; Thorp, R.W. Crop Pollination from Native Bees at Risk from Agricultural Intensification. *Proc. Natl. Acad. Sci. USA* **2002**, *99*, 16812–16816. [[CrossRef](#)]
109. Ricketts, T.H.; Regetz, J.; Steffan-Dewenter, I.; Cunningham, S.A.; Kremen, C.; Bogdanski, A.; Gemmill-Herren, B.; Greenleaf, S.S.; Klein, A.M.; Mayfield, M.M.; et al. Landscape Effects on Crop Pollination Services: Are There General Patterns? *Ecol. Lett.* **2008**, *11*, 499–515. [[CrossRef](#)]
110. Ockinger, E.; Smith, H.G. Seminatural Grasslands as Population Sources for Pollinating Insects in Agricultural Landscapes. *J. Appl. Ecol.* **2007**, *44*, 50–59. [[CrossRef](#)]
111. Stuligross, C.; Williams, N.M. Pesticide and Resource Stressors Additively Impair Wild Bee Reproduction. *Proc. R. Soc. B Biol. Sci.* **2020**, *287*, 20201390. [[CrossRef](#)]
112. Hendrickx, F.; Maelfait, J.P.; Van Wingerden, W.; Schweiger, O.; Speelmans, M.; Aviron, S.; Augenstein, I.; Billeter, R.; Bailey, D.; Bukacek, R.; et al. How Landscape Structure, Land-Use Intensity and Habitat Diversity Affect Components of Total Arthropod Diversity in Agricultural Landscapes. *J. Appl. Ecol.* **2007**, *44*, 340–351. [[CrossRef](#)]
113. Senapathi, D.; Goddard, M.A.; Kunin, W.E.; Baldock, K.C.R. Landscape Impacts on Pollinator Communities in Temperate Systems: Evidence and Knowledge Gaps. *Funct. Ecol.* **2017**, *31*, 26–37. [[CrossRef](#)]
114. Hopfenmüller, S.; Steffan-Dewenter, I.; Holzschuh, A. Trait-Specific Responses of Wild Bee Communities to Landscape Composition, Configuration and Local Factors. *PLoS ONE* **2014**, *9*. [[CrossRef](#)]
115. Forrest, J.R.K.; Thorp, R.W.; Kremen, C.; Williams, N.M. Contrasting Patterns in Species and Functional-Trait Diversity of Bees in an Agricultural Landscape. *J. Appl. Ecol.* **2015**, *52*, 706–715. [[CrossRef](#)]
116. Williams, N.M.; Crone, E.E.; Roulston, T.H.; Minckley, R.L.; Packer, L.; Potts, S.G. Ecological and Life-History Traits Predict Bee Species Responses to Environmental Disturbances. *Biol. Conserv.* **2010**, *143*, 2280–2291. [[CrossRef](#)]
117. De Palma, A.; Kuhlmann, M.; Roberts, S.P.M.; Potts, S.G.; Börger, L.; Hudson, L.N.; Lysenko, I.; Newbold, T.; Purvis, A. Ecological Traits Affect the Sensitivity of Bees to Land-Use Pressures in European Agricultural Landscapes. *J. Appl. Ecol.* **2015**, *52*, 1567–1577. [[CrossRef](#)] [[PubMed](#)]
118. Fitzpatrick, Ú.; Murray, T.E.; Paxton, R.J.; Breen, J.; Cotton, D.; Santorum, V.; Brown, M.J.F. Rarity and Decline in Bumblebees—A Test of Causes and Correlates in the Irish Fauna. *Biol. Conserv.* **2007**, *136*, 185–194. [[CrossRef](#)]
119. Basu, P.; Parui, A.K.; Chatterjee, S.; Dutta, A.; Chakraborty, P.; Roberts, S.; Smith, B. Scale Dependent Drivers of Wild Bee Diversity in Tropical Heterogeneous Agricultural Landscapes. *Ecol. Evol.* **2016**, *6*, 6983–6992. [[CrossRef](#)] [[PubMed](#)]
120. Hass, A.L.; Liese, B.; Heong, K.L.; Settele, J.; Tscharrntke, T.; Westphal, C. Plant-Pollinator Interactions and Bee Functional Diversity Are Driven by Agroforests in Rice-Dominated Landscapes. *Agric. Ecosyst. Environ.* **2018**, *253*, 140–147. [[CrossRef](#)]
121. Ballantyne, G.; Baldock, K.C.R.; Rendell, L.; Willmer, P.G. Pollinator Importance Networks Illustrate the Crucial Value of Bees in a Highly Speciose Plant Community. *Sci. Rep.* **2017**, *7*, 8389. [[CrossRef](#)]
122. Wood, T.; Holland, J.; Goulson, D. Diet Characterisation of Solitary Bees on Farmland: Dietary Specialisation Predicts Rarity. *Biodivers. Conserv.* **2016**, *25*, 2655–2671. [[CrossRef](#)]
123. Potts, S.G.; Vulliamy, B.; Roberts, S. Role of Nesting Resources in Organising Diverse Bee Communities in a Mediterranean Landscape. *Ecol. Entomol.* **2005**, *30*, 78–85. [[CrossRef](#)]
124. Williams, N.M.; Kremen, C. Resource Distributions among Habitats Determine Solitary Bee Offspring Production in a Mosaic Landscape. *Ecol. Appl.* **2007**, *17*, 910–921. [[CrossRef](#)]
125. Kennedy, C.M.; Lonsdorf, E.; Neel, M.C.; Williams, N.M.; Ricketts, T.H.; Winfree, R.; Bommarco, R.; Brittain, C.; Burley, A.L.; Cariveau, D.; et al. A Global Quantitative Synthesis of Local and Landscape Effects on Wild Bee Pollinators in Agroecosystems. *Ecol. Lett.* **2013**, *16*, 584–599. [[CrossRef](#)]
126. Carré, G.; Roche, P.; Chifflet, R.; Morison, N.; Bommarco, R.; Harrison-Cripps, J.; Krewenka, K.; Potts, S.G.; Roberts, S.P.M.; Rodet, G.; et al. Landscape Context and Habitat Type as Drivers of Bee Diversity in European Annual Crops. *Agric. Ecosyst. Environ.* **2009**, *133*, 40–47. [[CrossRef](#)]
127. Grab, H.; Branstetter, M.G.; Amon, N.; Urban-Mead, K.R.; Park, M.G.; Gibbs, J.; Blitzer, E.J.; Poveda, K.; Loeb, G.; Danforth, B.N. Agriculturally Dominated Landscapes Reduce Bee Phylogenetic Diversity and Pollination Services. *Science* **2019**, *363*, 282–284. [[CrossRef](#)]
128. Hines, H.M.; Hendrix, S.D. Bumble Bee (Hymenoptera: Apidae) Diversity and Abundance in Tallgrass Prairie Patches: Effects of Local and Landscape Floral Resources. *Environ. Entomol.* **2005**, *34*, 1477–1484. [[CrossRef](#)]

129. Giannini, T.C.; Tambosi, L.R.; Acosta, A.L.; Jaffé, R.; Saraiva, A.M.; Imperatriz-Fonseca, V.L.; Metzger, J.P. Safeguarding Ecosystem Services: A Methodological Framework to Buffer the Joint Effect of Habitat Configuration and Climate Change. *PLoS ONE* **2015**, *10*. [[CrossRef](#)]
130. Memmott, J.; Waser, N.M.; Price, M.V. Tolerance of Pollination Networks to Species Extinctions. *Proc. R. Soc. Lond. Ser. B Biol. Sci.* **2004**, *2605*–2611. [[CrossRef](#)]
131. McKinney, M.; Lockwood, J. Biotic Homogenization: A Few Winners Replacing Many Losers in the next Mass Extinction. *Trends Ecol. Evol.* **1999**, *14*, 450–453. [[CrossRef](#)]
132. Martins, K.T.; Gonzalez, A.; Lechowicz, M.J. Pollination Services Are Mediated by Bee Functional Diversity and Landscape Context. *Agric. Ecosyst. Environ.* **2015**, *200*, 12–20. [[CrossRef](#)]
133. Bennett, J.M.; Steets, J.A.; Burns, J.H.; Burkle, L.A.; Vamosi, J.C.; Wolowski, M.; Arceo-Gómez, G.; Burd, M.; Durka, W.; Ellis, A.G.; et al. Land Use and Pollinator Dependency Drives Global Patterns of Pollen Limitation in the Anthropocene. *Nat. Commun.* **2020**, *11*, 3999. [[CrossRef](#)]
134. Kleijn, D.; Winfree, R.; Bartomeus, I.; Carvalheiro, L.G.; Henry, M.; Isaacs, R.; Klein, A.M.; Kremen, C.; M'Gonigle, L.K.; Rader, R.; et al. Delivery of Crop Pollination Services Is an Insufficient Argument for Wild Pollinator Conservation. *Nat. Commun.* **2015**, *6*. [[CrossRef](#)]
135. Cunningham, S.; Lindenmayer, D.; Young, A. *Land Use Intensification: Effects on Agriculture, Biodiversity and Ecological Processes*; CSIRO Publishing: Collingwood, VIC, Australia, 2012.
136. Winfree, R.; Aguilar, R.; Vázquez, D.P.; Lebuhn, G.; Aizen, M.A. A Meta-Analysis of Bees' Responses to Anthropogenic Disturbance. *Ecology* **2009**, *90*, 2068–2076. [[CrossRef](#)]
137. Hladik, M.L.; Vandever, M.; Smalling, K.L. Exposure of Native Bees Foraging in an Agricultural Landscape to Current-Use Pesticides. *Sci. Total Environ.* **2016**, *542*, 469–477. [[CrossRef](#)] [[PubMed](#)]
138. Jeschke, P.; Nauen, R. Review Neonicotinoids—From Zero to Hero in Insecticide Chemistry. *Pest Manag. Sci.* **2008**, *64*, 1084–1098. [[CrossRef](#)] [[PubMed](#)]
139. James, R.R.; Xu, J. Mechanisms by Which Pesticides Affect Insect Immunity. *J. Invertebr. Pathol.* **2012**, *109*, 175–182. [[CrossRef](#)] [[PubMed](#)]
140. Tomé, H.V.V.; Martins, G.F.; Lima, M.A.P.; Campos, L.A.O.; Guedes, R.N.C. Imidacloprid-Induced Impairment of Mushroom Bodies and Behavior of the Native Stingless Bee *Melipona quadrifasciata* Anthidioides. *PLoS ONE* **2012**. [[CrossRef](#)] [[PubMed](#)]
141. Ken, T.; Chen, W.; Dong, S.; Liu, X.; Wang, Y.; Nieh, J.C. Imidacloprid Alters Foraging and Decreases Bee Avoidance of Predators. *PLoS ONE* **2014**. [[CrossRef](#)]
142. Morandin, L.A.; Winston, M.L.; Franklin, M.T.; Abbott, V.A. Lethal and Sub-Lethal Effects of Spinosad on Bumble Bees (*Bombus impatiens* Cresson). *Pest Manag. Sci.* **2005**, *61*, 619–626. [[CrossRef](#)] [[PubMed](#)]
143. Ken, T.; Chen, W.; Dong, S.; Liu, X.; Wang, Y.; Nieh, J.C. A Neonicotinoid Impairs Olfactory Learning in Asian Honey Bees (*Apis cerana*) Exposed as Larvae or as Adults. *Sci. Rep.* **2015**, *5*. [[CrossRef](#)]
144. Woodcock, B.A.; Bullock, J.M.; Shore, R.F.; Heard, M.S.; Pereira, M.G.; Redhead, J.; Ridding, L.; Dean, H.; Sleep, D.; Henrys, P.; et al. Country-Specific Effects of Neonicotinoid Pesticides on Honey Bees and Wild Bees. *Science* **2017**, *356*, 1393–1395. [[CrossRef](#)]
145. Crall, J.D.; Switzer, C.M.; Oppenheimer, R.L.; Ford Versypt, A.N.; Dey, B.; Brown, A.; Eyster, M.; Guérin, C.; Pierce, N.E.; Combes, S.A.; et al. Neonicotinoid Exposure Disrupts Bumblebee Nest Behavior, Social Networks, and Thermoregulation. *Science* **2018**, *362*, 683–686. [[CrossRef](#)]
146. Sandrock, C.; Tanadini, L.G.; Pettis, J.S.; Biesmeijer, J.C.; Potts, S.G.; Neumann, P. Sublethal Neonicotinoid Insecticide Exposure Reduces Solitary Bee Reproductive Success. *Agric. For. Entomol.* **2014**, *16*, 119–128. [[CrossRef](#)]
147. Gill, R.J.; Raine, N.E. Chronic Impairment of Bumblebee Natural Foraging Behaviour Induced by Sublethal Pesticide Exposure. *Funct. Ecol.* **2014**, *28*, 1459–1471. [[CrossRef](#)]
148. Wu, J.Y.; Anelli, C.M.; Sheppard, W.S. Sub-Lethal Effects of Pesticide Residues in Brood Comb on Worker Honey Bee (*Apis mellifera*) Development and Longevity. *PLoS ONE* **2011**, *6*. [[CrossRef](#)] [[PubMed](#)]
149. Bebane, P.S.A.; Hunt, B.J.; Pegoraro, M.; Jones, A.R.C.; Marshall, H.; Rosato, E.; Mallon, E.B. The Effects of the Neonicotinoid Imidacloprid on Gene Expression and DNA Methylation in the Buff-Tailed Bumblebee *Bombus terrestris*. *Proc. R. Soc. B Biol. Sci.* **2019**, *286*. [[CrossRef](#)] [[PubMed](#)]
150. Brevik, K.; Bueno, E.M.; McKay, S.; Schoville, S.D.; Chen, Y.H. Insecticide Exposure Affects Intergenerational Patterns of DNA Methylation in the Colorado Potato Beetle, *Leptinotarsa decemlineata*. *Evol. Appl.* **2020**. [[CrossRef](#)]
151. Brevik, K.; Lindström, L.; McKay, S.D.; Chen, Y.H. Transgenerational Effects of Insecticides—Implications for Rapid Pest Evolution in Agroecosystems. *Curr. Opin. Insect Sci.* **2018**, *26*, 34–40. [[CrossRef](#)]
152. Kevan, P.G. Forest Application of the Insecticide Fenitrothion and Its Effect on Wild Bee Pollinators (Hymenoptera: Apoidea) of Lowbush Blueberries (*Vaccinium* spp.) in Southern New Brunswick, Canada. *Biol. Conserv.* **1975**, *7*, 301–309. [[CrossRef](#)]
153. Park, M.G.; Blitzer, E.J.; Gibbs, J.; Losey, J.E.; Danforth, B.N. Negative Effects of Pesticides on Wild Bee Communities Can Be Buffered by Landscape Context. *Proc. R. Soc. B Biol. Sci.* **2015**, *282*. [[CrossRef](#)] [[PubMed](#)]
154. Lüscher, G.; Jeanneret, P.; Schneider, M.K.; Turnbull, L.A.; Arndorfer, M.; Balázs, K.; Baldi, A.; Bailey, D.; Bernhardt, K.G.; Choisis, J.P.; et al. Responses of Plants, Earthworms, Spiders and Bees to Geographic Location, Agricultural Management and Surrounding Landscape in European Arable Fields. *Agric. Ecosyst. Environ.* **2014**, *186*, 124–134. [[CrossRef](#)]

155. Woodcock, B.A.; Isaac, N.J.B.; Bullock, J.M.; Roy, D.B.; Garthwaite, D.G.; Crowe, A.; Pywell, R.F. Impacts of Neonicotinoid Use on Long-Term Population Changes in Wild Bees in England. *Nat. Commun.* **2016**. [[CrossRef](#)]
156. Tasei, J. Impact of Agrochemicals on Non-Apis Bees. In *Honey Bees*; Taylor & Francis Group: London, UK, 2010; pp. 101–131. [[CrossRef](#)]
157. Kopit, A.M.; Pitts-Singer, T.L. Routes of Pesticide Exposure in Solitary, Cavity-Nesting Bees. *Environ. Entomol.* **2018**, *47*, 499–510. [[CrossRef](#)]
158. Sgolastra, F.; Hinarejos, S.; Pitts-Singer, T.L.; Boyle, N.K.; Joseph, T.; Luckmann, J.; Raine, N.E.; Singh, R.; Williams, N.M.; Bosch, J. Pesticide Exposure Assessment Paradigm for Solitary Bees. *Environ. Entomol.* **2019**, *48*, 22–35. [[CrossRef](#)] [[PubMed](#)]
159. Krupke, C.H.; Hunt, G.J.; Eitzer, B.D.; Andino, G.; Given, K. Multiple Routes of Pesticide Exposure for Honey Bees Living near Agricultural Fields. *PLoS ONE* **2012**, *7*. [[CrossRef](#)] [[PubMed](#)]
160. Botías, C.; David, A.; Horwood, J.; Abdul-Sada, A.; Nicholls, E.; Hill, E.; Goulson, D. Neonicotinoid Residues in Wildflowers, a Potential Route of Chronic Exposure for Bees. *Environ. Sci. Technol.* **2015**, *49*, 12731–12740. [[CrossRef](#)] [[PubMed](#)]
161. Goulson, D. An Overview of the Environmental Risks Posed by Neonicotinoid Insecticides. *J. Appl. Ecol.* **2013**, *50*, 977–987. [[CrossRef](#)]
162. Main, A.R.; Webb, E.B.; Goynes, K.W.; Mengel, D. Agriculture, Ecosystems and Environment Reduced Species Richness of Native Bees in Field Margins Associated with Neonicotinoid Concentrations in Non-Target Soils. *Agric. Ecosyst. Environ.* **2020**, *287*, 106693. [[CrossRef](#)]
163. Bredeson, M.M.; Lundgren, J.G. Neonicotinoid Insecticidal Seed-Treatment on Corn Contaminates Interseeded Cover Crops Intended as Habitat for Beneficial Insects. *Ecotoxicology* **2019**, *28*, 222–228. [[CrossRef](#)]
164. Stewart, S.D.; Lorenz, G.M.; Catchot, A.L.; Gore, J.; Cook, D.; Skinner, J.; Mueller, T.C.; Johnson, D.R.; Zawislak, J.; Barber, J. Potential Exposure of Pollinators to Neonicotinoid Insecticides from the Use of Insecticide Seed Treatments in the Mid-Southern United States. *Environ. Sci. Technol.* **2014**, *48*, 9762–9769. [[CrossRef](#)] [[PubMed](#)]
165. Long, E.Y.; Krupke, C.H. Non-Cultivated Plants Present a Season-Long Route of Pesticide Exposure for Honey Bees. *Nat. Commun.* **2016**, *7*, 11629. [[CrossRef](#)]
166. Gill, R.J.; Ramos-Rodriguez, O.; Raine, N.E. Combined Pesticide Exposure Severely Affects Individual- and Colony-Level Traits in Bees. *Nature* **2012**, *491*, 105–108. [[CrossRef](#)] [[PubMed](#)]
167. Stanley, D.A.; Garratt, M.P.D.; Wickens, J.B.; Wickens, V.J.; Potts, S.G.; Raine, N.E. Neonicotinoid Pesticide Exposure Impairs Crop Pollination Services Provided by Bumblebees. *Nature* **2015**, *528*, 548–550. [[CrossRef](#)] [[PubMed](#)]
168. Feltham, H.; Park, K.; Goulson, D. Field Realistic Doses of Pesticide Imidacloprid Reduce Bumblebee Pollen Foraging Efficiency. *Ecotoxicology* **2014**, *23*, 317–323. [[CrossRef](#)] [[PubMed](#)]
169. Biddinger, D.J.; Robertson, J.L.; Mullin, C.; Frazier, J.; Ashcraft, S.A.; Rajotte, E.G.; Joshi, N.K.; Vaughn, M. Comparative Toxicities and Synergism of Apple Orchard Pesticides to *Apis mellifera* (L.) and *Osmia cornifrons* (Radoszkowski). *PLoS ONE* **2013**, *8*. [[CrossRef](#)] [[PubMed](#)]
170. Tomé, H.V.V.; Ramos, G.S.; Araújo, M.F.; Santana, W.C.; Santos, G.R.; Guedes, R.N.C.; Maciel, C.D.; Newland, P.L.; Oliveira, E.E. Agrochemical Synergism Imposes Higher Risk to Neotropical Bees than to Honeybees. *R. Soc. Open Sci.* **2017**, *4*. [[CrossRef](#)] [[PubMed](#)]
171. Ladurner, E.; Bosch, J.; Kemp, W.P.; Maini, S. Assessing Delayed and Acute Toxicity of Five Formulated Fungicides to *Osmia lignaria* Say and *Apis mellifera*. *Apidologie* **2005**, *36*, 449–460. [[CrossRef](#)]
172. Soares, H.M.; Jacob, C.R.O.; Carvalho, S.M.; Nocelli, R.C.F.; Malaspina, O. Toxicity of Imidacloprid to the Stingless Bee *Scaptotrigona postica* Latreille, 1807 (Hymenoptera: Apidae). *Bull. Environ. Contam. Toxicol.* **2015**, *94*, 675–680. [[CrossRef](#)]
173. Arena, M.; Sgolastra, F. A Meta-Analysis Comparing the Sensitivity of Bees to Pesticides. *Ecotoxicology* **2014**, *23*, 324–334. [[CrossRef](#)]
174. Brittain, C.; Potts, S.G. The Potential Impacts of Insecticides on the Life-History Traits of Bees and the Consequences for Pollination. *Basic Appl. Ecol.* **2011**, *12*, 321–331. [[CrossRef](#)]
175. Mallinger, R.E.; Gaines-Day, H.R.; Gratton, C. Do Managed Bees Have Negative Effects on Wild Bees?: A Systematic Review of the Literature. *PLoS ONE* **2017**, *12*, e0189268. [[CrossRef](#)]
176. Paini, D.R. Impact of the Introduced Honey Bee (*Apis mellifera*) (Hymenoptera: Apidae) on Native Bees: A Review. *Austral Ecology* **2004**, 399–407. [[CrossRef](#)]
177. Goulson, D. Effects of Introduced Bees on Native Ecosystems. *Annu. Rev. Ecol. Evol. Syst.* **2003**, *34*, 1–26. [[CrossRef](#)]
178. Thomson, D. Competitive Interactions between the Invasive European Honey Bee and Native Bumble Bees. *Ecology* **2004**, *85*, 458–470. [[CrossRef](#)]
179. Kanbe, Y.; Okada, I.; Yoneda, M.; Goka, K.; Tsuchida, K. Interspecific Mating of the Introduced Bumblebee *Bombus terrestris* and the Native Japanese Bumblebee *Bombus hypocrita sapporoensis* Results in Inviabile Hybrids. *Naturwissenschaften* **2008**, *95*, 1003–1008. [[CrossRef](#)] [[PubMed](#)]
180. Kondo, N.I.; Yamanaka, D.; Kanbe, Y.; Kunitake, Y.K.; Yoneda, M.; Tsuchida, K.; Goka, K. Reproductive Disturbance of Japanese Bumblebees by the Introduced European Bumblebee *Bombus terrestris*. *Naturwissenschaften* **2009**, *96*, 467–475. [[CrossRef](#)] [[PubMed](#)]
181. Alomar, D.; González-Estévez, M.A.; Traveset, A.; Lázaro, A. The Intertwined Effects of Natural Vegetation, Local Flower Community, and Pollinator Diversity on the Production of Almond Trees. *Agric. Ecosyst. Environ.* **2018**, *264*, 34–43. [[CrossRef](#)]

182. Matsumura, C.; Yokoyama, J.; Washitani, I. Invasion Status and Potential Ecological Impacts of an Invasive Alien Bumblebee, *Bombus Terrestris* L. (Hymenoptera: Apidae) Naturalized in Southern Hokkaido, Japan. *Glob. Environ. Res.* **2004**, *8*, 51–66.
183. Inoue, M.N.; Yokoyama, J.; Washitani, I. Displacement of Japanese Native Bumblebees by the Recently Introduced *Bombus Terrestris* (L.) (Hymenoptera: Apidae). *J. Insect Conserv.* **2008**, *12*, 135–146. [CrossRef]
184. Hung, K.-L.J.; Kingston, J.M.; Lee, A.; Holway, D.A.; Kohn, J.R. Non-Native Honey Bees Disproportionately Dominate the Most Abundant Floral Resources in a Biodiversity Hotspot. *Proc. R. Soc. B Biol. Sci.* **2019**, *286*, 20182901. [CrossRef]
185. Magrach, A.; González-Varo, J.P.; Boiffier, M.; Vilà, M.; Bartomeus, I. Honeybee Spillover Reshuffles Pollinator Diets and Affects Plant Reproductive Success. *Nat. Ecol. Evol.* **2017**, *1*, 1299–1307. [CrossRef] [PubMed]
186. Cane, J.H.; Tepedino, V.J. Gauging the Effect of Honey Bee Pollen Collection on Native Bee Communities. *Conserv. Lett.* **2017**, *10*, 205–210. [CrossRef]
187. Carneiro, L.T.; Martins, C.F. Africanized Honey Bees Pollinate and Preempt the Pollen of *Spondias mombin* (Anacardiaceae) Flowers. *Apidologie* **2012**, *43*, 474–486. [CrossRef]
188. Chen, Y.; Evans, J.; Feldlaufer, M. Horizontal and Vertical Transmission of Viruses in the Honey Bee, *Apis mellifera*. *J. Invertebr. Pathol.* **2006**, *92*, 152–159. [CrossRef] [PubMed]
189. Alger, S.A.; Burnham, P.A.; Brody, A.K. Flowers as Viral Hot Spots: Honey Bees (*Apis mellifera*) Unevenly Deposit Viruses across Plant Species. *PLoS ONE* **2019**, *14*. [CrossRef]
190. Purkiss, T.; Lach, L. Pathogen Spillover from *Apis mellifera* to a Stingless Bee. *Proc. R. Soc. B Biol. Sci.* **2019**, *286*. [CrossRef] [PubMed]
191. McMahon, D.P.; Fürst, M.A.; Caspar, J.; Theodorou, P.; Brown, M.J.F.; Paxton, R.J. A Sting in the Spit: Widespread Cross-Infection of Multiple RNA Viruses across Wild and Managed Bees. *J. Anim. Ecol.* **2015**, *84*, 615–624. [CrossRef] [PubMed]
192. Fürst, M.A.; McMahon, D.P.; Osborne, J.L.; Paxton, R.J.; Brown, M.J.F. Disease Associations between Honeybees and Bumblebees as a Threat to Wild Pollinators. *Nature* **2014**, *506*, 364–366. [CrossRef] [PubMed]
193. Arroyo, M.K.; Marquet, P.A.; Simonetti, J.A.; Cavieres, L.A. Chilean Winter Rainfall-Valdivian Forests. In *Hotspots: Earth's Biological Richest and most Endangered Terrestrial Ecoregions*; Mittermeier, R.A., Robles, P., Hoffmann, M., Pilgrim, J., Brooks, T., Mittermeier, C., et al., Eds.; CEMEX: Mexico City, Mexico, 2004; pp. 99–103.
194. Harvey, C.A.; Komar, O.; Chazdon, R.; Ferguson, B.G.; Finegan, B.; Griffith, D.M.; Martínez-Ramos, M.; Morales, H.; Nigh, R.; Soto-Pinto, L.; et al. Integrating Agricultural Landscapes with Biodiversity Conservation in the Mesoamerican Hotspot. *Conserv. Biol.* **2008**, *8*–15. [CrossRef]
195. Kehinde, T.; Samways, M.J. Endemic Pollinator Response to Organic vs. Conventional Farming and Landscape Context in the Cape Floristic Region Biodiversity Hotspot. *Agric. Ecosyst. Environ.* **2012**, *146*, 162–167. [CrossRef]
196. Schutz, J. Creating an Integrated Protected Area Network in Chile: A GIS Assessment of Ecoregion Representation and the Role of Private Protected Areas. *Environ. Conserv.* **2018**, *45*, 269–277. [CrossRef]
197. Montalva, J.; Ruz, L. Actualización de La Lista Sistemática de Las Abejas Chilenas (Hymenoptera: Apoidea). *Rev. Chil. Entomol.* **2010**, *35*, 15–52.
198. United Nations. Paris Agreement (COP21). 2015. Available online: <https://www.un.org/en/climatechange/paris-agreement> (accessed on 1 June 2021).
199. NYDF Global Platform. *New York Declaration on Forests*. 2014. Available online: <https://forestdeclaration.org/> (accessed on 1 June 2021).
200. UN Convention on Biological Diversity. Aichi Biodiversity Targets. Chile National Targets. 2010. Available online: <https://www.cbd.int/> (accessed on 1 June 2021).
201. WRI. 20x20 Initiative. Healthy Lands for Food, Water and Climate. 2014. Available online: <https://initiative20x20.org/es> (accessed on 1 June 2021).
202. Carmona, A.; Nahuelhual, L.; Echeverría, C.; Báez, A. Linking Farming Systems to Landscape Change: An Empirical and Spatially Explicit Study in Southern Chile. *Agric. Ecosyst. Environ.* **2010**, *139*, 40–50. [CrossRef]
203. Del Pozo, A.; Lavin, A.; Etienne, M.; Ovalle, C.; Avendaño, J.; Aronson, J. Land Use Changes and Conflicts in Central Chile. In *Landscape Disturbance and Biodiversity in Mediterranean-Type Ecosystems*; Rundel, P.W., Montenegro, G., Jaksic, F., Eds.; Springer: Berlin/Heidelberg, Germany, 2013; pp. 155–168. [CrossRef]
204. ODEPA. *Panorama de La Agricultura Chilena (Chilean Agriculture Overview)*; A Impresores: Santiago, Chile, 2019; Available online: <https://www.odepa.gob.cl/wp-content/uploads/2019/09/panorama2019Final.pdf> (accessed on 1 June 2021).
205. Ministerio del Medio Ambiente. Sexto Informe Nacional de Biodiversidad de Chile. *Ministerio del Medio Ambiente*. **2019**, 220. Available online: https://mma.gob.cl/wp-content/uploads/2020/01/6NR_FINAL_ALTA-web.pdf (accessed on 1 June 2021).
206. Gwynne, R.N. Globalisation, Commodity Chains and Fruit Exporting Regions in Chile. *Tijdschr. Econ. Soc. Geogr.* **1999**. [CrossRef]
207. Altieri, M.A.; Rojas, A. Ecological Impacts of Chile's Neoliberal Policies, with Special Emphasis on Agroecosystems. *Environ. Dev. Sustain.* **1999**, *1*, 55–72. [CrossRef]
208. Carruthers, D. Environmental Politics in Chile: Legacies of Dictatorship and Democracy. *Third World Q.* **2001**, *22*, 343–358. [CrossRef]
209. Muñoz, O.; Ortega, H. Chilean Agriculture and Economic Policy. In *Modernization and Stagnation: Latin American Agriculture into the 1990's*; Helwege, M.J.T., Ed.; Greenwood Press: New York, NY, USA, 1991; pp. 161–188.

210. Clark, T.D. Putting the Market in Its Place: Food Security in Three Mapuche Communities in Southern Chile. *Lat. Am. Res. Rev.* **2011**, *46*, 154–179. [CrossRef]
211. Kay, C. Chile's Neoliberal Agrarian Transformation and the Peasantry. *J. Agrar. Chang.* **2002**, *2*, 464–501. [CrossRef]
212. FAO. GIAHS: Globally Important Agricultural Heritage Systems; Chiloé Agriculture, Chile. Available online: <http://www.fao.org/3/bp773e/bp773e.pdf> (accessed on 1 June 2021).
213. Wratten, S.D.; Shields, M.W.; González-Chang, M. Prospects for Regenerative Agriculture in Chile. *Agro Sur* **2019**, *47*, 1–6. [CrossRef]
214. SAG. Lista de Plaguicidas Autorizados. Available online: <https://www.sag.gob.cl/ambitos-de-accion/plaguicidas-y-fertilizantes/78/registros> (accessed on 1 June 2021).
215. Samson-Robert, O.; Labrie, G.; Chagnon, M.; Fournier, V. Planting of Neonicotinoid-Coated Corn Raises Honey Bee Mortality and Sets Back Colony Development. *PeerJ* **2017**, *5*, e3670. [CrossRef]
216. European Commission. Neonicotinoids. Available online: https://ec.europa.eu/food/plants/pesticides/approval-active-substances/renewal-approval/neonicotinoids_en (accessed on 1 June 2021).
217. Paleolog, J.; Wilde, J.; Siuda, M.; Bąk, B.; Wójcik, Ł.; Strachecka, A. Imidacloprid Markedly Affects Hemolymph Proteolysis, Biomarkers, DNA Global Methylation, and the Cuticle Proteolytic Layer in Western Honeybees. *Apidologie* **2020**, *51*, 620–630. [CrossRef]
218. Colin, T.; Meikle, W.G.; Wu, X.; Barron, A.B. Traces of a Neonicotinoid Induce Precocious Foraging and Reduce Foraging Performance in Honey Bees. *Environ. Sci. Technol.* **2019**, *53*, 8252–8261. [CrossRef]
219. Acta Reunión de Panel de Discusión de Expertos Proyecto Ley Apícola. *Legislado Para La Protección de La Salud y Hábitat de Las Abejas*, 2015; 1–6.
220. SAG. *Informe de Venta de Plaguicidas de Uso Agrícola En Chile, Año 2012*; Servicio Agrícola y Ganadero División Protección Agrícola y Forestal Subdepartamento de Plaguicidas y Fertilizantes, Gobierno de Chile: Chile, 2012.; Available online: http://www.sag.cl/sites/default/files/declaracion_de_venta_de_plaguicidas_ano_2012.pdf (accessed on 1 June 2021).
221. Caceres-Jensen, L.; Rodriguez-Becerra, J.; Escudey, M.; Joo-Nagata, J.; Villagra, C.A.; Dominguez-Vera, V.; Neira-Albornoz, A.; Cornejo-Huentemilla, M. Nicosulfuron Sorption Kinetics and Sorption/Desorption on Volcanic Ash-Derived Soils: Proposal of Sorption and Transport Mechanisms. *J. Hazard. Mater.* **2020**, *385*, 121576. [CrossRef]
222. Aparicio, V.C.; De Geronimo, E.; Marino, D.; Primost, J.; Carriquiriborde, P.; Costa, J.L. Environmental Fate of Glyphosate and Aminomethylphosphonic Acid in Surface Waters and Soil of Agricultural Basins. *Chemosphere* **2013**, *93*, 1866–1873. [CrossRef]
223. Michener, C.D. *The Bees the World*, 2nd ed.; The Johns Hopkins University Press: Baltimore, MD, USA, 2007.
224. Ministerio de Salud; Subsecretaría de Salud Pública. *Decreto 157. Reglamento de Pesticidas de Uso Sanitario y Doméstico*; Biblioteca del Congreso Nacional de Chile: Santiago de Chile, Chile, 2007; p. 19.
225. Linnaeus, C. *Systema Naturae*, 10th ed.; Salvii: Stockholm, Sweden, 1758; Volume 1.
226. Fabricius, J.C. *Systema Entomologiae, Sistens Insectorum Classes, Ordines, Genera, Species, Adjectis Synonymis, Locis, Descriptionibus, Observationibus*, Flensbvrge et Lipsiae: In Officina Libraria Kortii. 1775.
227. Guérin-Méneville, F.-É. *Bombus Dahlbomii* Pp. Pl. 75, Fig. 3. In *Iconographie du Règne Animal de G. Cuvier, ou Représentation D'après Nature de l'une des Espèces les Plus Remarquables, et Souvent non Encore Figurées, de Chaque Genre D'animaux; Pouvant Servir D'atlas à Tous les Traités de Zoologie*; Guérin-Méneville, F.E., Ed.; Baillière: Paris, France, 1835; p. 576.
228. Madjidian, J.A.; Morales, C.L.; Smith, H.G. Displacement of a Native by an Alien Bumblebee: Lower Pollinator Efficiency Overcome by Overwhelmingly Higher Visitation Frequency. *Oecologia* **2008**, *156*, 835–845. [CrossRef] [PubMed]
229. Morales, C.L.; Arbetman, M.P.; Cameron, S.A.; Aizen, M.A. Rapid Ecological Replacement of a Native Bumble Bee by Invasive Species. *Front. Ecol. Environ.* **2013**, *11*, 529–534. [CrossRef]
230. Montalva, J.; Dudley, L.S.; Sepúlveda, J.E.; Smith-Ramírez, C. The Giant Bumble Bee (*Bombus dahlbomii*) in Mapuche Cosmovision. *Ethnoentomology* **2020**, *4*, 1–11.
231. Smith-Ramírez, C.; Vieli, L.; Barahona-Segovia, R.M.; Montalva, J.; Cianferoni, F.; Ruz, L.; Fontúrbel, F.E.; Valdivia, C.E.; Medel, R.; Pauchard, A.; et al. Las Razones de Por Qué Chile Debe Detener La Importación Del Abejorro Comercial *Bombus terrestris* (Linnaeus) y Comenzar a Controlarlo. *Gayana (Concepción)* **2018**, *82*, 118–127. [CrossRef]
232. Geslin, B.; Morales, C.L. New Records Reveal Rapid Geographic Expansion of *Bombus terrestris* Linnaeus, 1758 (Hymenoptera: Apidae), an Invasive Species in Argentina. *Check List* **2015**, *11*, 1620. [CrossRef]
233. Monzón, V.H.; Avendaño-Soto, P.; Araujo, R.O.; Garrido, R.; Mesquita-Neto, J.N. Avocado Crops as a Floral Resource for Native Bees of Chile. *Rev. Chil. Hist. Nat.* **2020**, *93*. [CrossRef]
234. Morales, C.; Montalva, J.; Arbetman, M.; Aizen, M.; Smith-Ramírez, C.; Vieli, L.; Hatfield, R. *Bombus dahlbomii*. *The IUCN Red List of Threatened Species* **2016**. [CrossRef]
235. Estay, P.; Wagner, A.; Escaff, M. Evaluación DE *Bombus dahlbomii* (GUÉR.) como agente polinizador de flores de tomate (*Lycopersicon esculentum* (mill)), bajo condiciones de invernadero. *Agric. Técnica Scienclocl.* **2001**, 113–119. [CrossRef]
236. Santos, E.; Daners, G.; Morelli, E.; Galván, G.A. Diversity of Bee Assemblage (Family Apidae) in Natural and Agriculturally Intensified Ecosystems in Uruguay. *Environ. Entomol.* **2020**, *49*, 1232–1241. [CrossRef]
237. Le Féon, V.; Poggio, S.L.; Torretta, J.P.; Bertrand, C.; Molina, G.A.R.; Burel, F.; Baudry, J.; Ghersa, C.M. Diversity and Life-History Traits of Wild Bees (Insecta: Hymenoptera) in Intensive Agricultural Landscapes in the Rolling Pampa, Argentina. *J. Nat. Hist.* **2016**, *50*, 1175–1196. [CrossRef]

238. Rosanigo, M.P.; Marrero, H.J.; Torretta, J.P. Limiting Resources on the Reproductive Success of a Cavity-Nesting Bee Species in a Grassland Agroecosystem. *J. Apic. Res.* **2020**, *59*, 583–591. [\[CrossRef\]](#)
239. Allendes, J.L.; Montalva, J.; Castro, B. Las Abejas (Hymenoptera: Apoidea) Del Jardín Botánico Chagual. Estudio de Caso de Abejas Nativas En Zonas Urbanas de Santiago de Chile. *Rev. Chagual* **2010**, *8*, 13–23.
240. Steffen, W.; Richardson, K.; Rockström, J.; Cornell, S.E.; Fetzer, I.; Bennett, E.M.; Biggs, R.; Carpenter, S.R.; De Vries, W.; De Wit, C.A.; et al. Planetary Boundaries: Guiding Human Development on a Changing Planet. *Science* **2015**, *347*. [\[CrossRef\]](#)
241. Desing, H.; Brunner, D.; Takacs, F.; Nahrath, S.; Frankenberger, K.; Hirschier, R. A Circular Economy within the Planetary Boundaries: Towards a Resource-Based, Systemic Approach. *Resour. Conserv. Recycl.* **2020**, *155*, 104673. [\[CrossRef\]](#)
242. Heck, V.; Hoff, H.; Wirsenius, S.; Meyer, C.; Kreft, H. Land Use Options for Staying within the Planetary Boundaries—Synergies and Trade-Offs between Global and Local Sustainability Goals. *Glob. Environ. Chang.* **2018**, *49*, 73–84. [\[CrossRef\]](#)
243. Parraguez-Vergara, E.; Contreras, B.; Clavijo, N.; Villegas, V.; Paucar, N.; Ther, F. Does Indigenous and Campesino Traditional Agriculture Have Anything to Contribute to Food Sovereignty in Latin America? Evidence from Chile, Peru, Ecuador, Colombia, Guatemala and Mexico. *Int. J. Agric. Sustain.* **2018**, *16*, 326–341. [\[CrossRef\]](#)
244. Mickey, S. Learning Native Wisdom: What Traditional Cultures Teach Us About Subsistence, Sustainability, and Spirituality. *Worldviews Glob. Relig. Cult. Ecol.* **2013**, *13*, 136–139. [\[CrossRef\]](#)
245. Grey, S.; Patel, R. Food Sovereignty as Decolonization: Some Contributions from Indigenous Movements to Food System and Development Politics. *Agric. Human Values* **2015**, *32*, 431–444. [\[CrossRef\]](#)
246. Kaluza, B.F.; Wallace, H.M.; Heard, T.A.; Minden, V.; Klein, A.; Leonhardt, S.D. Social Bees Are Fitter in More Biodiverse Environments. *Sci. Rep.* **2018**, *8*, 12353. [\[CrossRef\]](#)
247. Kremen, C. Reframing the Land-sparing/Land-sharing Debate for Biodiversity Conservation. *Ann. N. Y. Acad. Sci.* **2015**, *1355*, 52–76. [\[CrossRef\]](#) [\[PubMed\]](#)
248. Kremen, C.; Williams, N.M.; Bugg, R.L.; Fay, J.P.; Thorp, R.W. The Area Requirements of an Ecosystem Service: Crop Pollination by Native Bee Communities in California. *Ecol. Lett.* **2004**, *7*, 1109–1119. [\[CrossRef\]](#)
249. Bailey, S.; Requier, F.; Nusillard, B.; Roberts, S.P.M.; Potts, S.G.; Bouget, C. Distance from Forest Edge Affects Bee Pollinators in Oilseed Rape Fields. *Ecol. Evol.* **2014**, *4*, 370–380. [\[CrossRef\]](#)
250. Gray, C.L.; Hill, S.L.L.; Newbold, T.; Hudson, L.N.; Boirger, L.; Contu, S.; Hoskins, A.J.; Ferrier, S.; Purvis, A.; Scharlemann, J.P.W. Local Biodiversity Is Higher inside than Outside Terrestrial Protected Areas Worldwide. *Nat. Commun.* **2016**, *7*. [\[CrossRef\]](#) [\[PubMed\]](#)
251. Felipe Viana, B. How Well Do We Understand Landscape Effects on Pollinators and Pollination Services? *J. Pollinat. Ecol.* **2012**, *7*, 31–40. [\[CrossRef\]](#)
252. Sobreiro, A.I. Recover and They'll Come: Flower Visiting Bees Benefit from the Continuous of Micro- Environments Set by Regenerating Forest Fragments. *Sociobiology* **2021**, *68*, 1–17. [\[CrossRef\]](#)
253. Chape, S.; Harrison, J.; Spalding, M.; Lysenko, I. Measuring the Extent and Effectiveness of Protected Areas as an Indicator for Meeting Global Biodiversity Targets. *Philos. Trans. R. Soc. B Biol. Sci.* **2005**, *360*, 443–455. [\[CrossRef\]](#)
254. Steffan-Dewenter, I.; Tschardtke, T. Effects of Habitat Isolation on Pollinator Communities and Seed Set. *Oecologia* **1999**, *121*, 432–440. [\[CrossRef\]](#)
255. Carrié, R.; Andrieu, E.; Ouin, A.; Steffan-Dewenter, I. Interactive Effects of Landscape-Wide Intensity of Farming Practices and Landscape Complexity on Wild Bee Diversity. *Landsc. Ecol.* **2017**, *32*, 1631–1642. [\[CrossRef\]](#)
256. Morandin, L.A.; Kremen, C. Hedgerow Restoration Promotes Pollinator Populations and Exports Native Bees to Adjacent Fields. *Ecol. Appl.* **2013**, *23*, 829–839. [\[CrossRef\]](#)
257. Blaauw, B.R.; Isaacs, R. Larger Patches of Diverse Floral Resources Increase Insect Pollinator Density, Diversity, and Their Pollination of Native Wildflowers. *Basic Appl. Ecol.* **2014**, *15*, 701–711. [\[CrossRef\]](#)
258. Riojas-López, M.E.; Díaz-Herrera, I.A.; Fierros-López, H.E.; Mellink, E. The Effect of Adjacent Habitat on Native Bee Assemblages in a Perennial Low-Input Agroecosystem in a Semiarid Anthropized Landscape. *Agric. Ecosyst. Environ.* **2019**, *272*, 199–205. [\[CrossRef\]](#)
259. Burkle, L.A.; Delphia, C.M.; O'Neill, K.M. A Dual Role for Farmlands: Food Security and Pollinator Conservation. *J. Ecol.* **2017**, *105*, 890–899. [\[CrossRef\]](#)
260. Krewenka, K.M.; Holzschuh, A.; Tschardtke, T.; Dormann, C.F. Landscape Elements as Potential Barriers and Corridors for Bees, Wasps and Parasitoids. *Biol. Conserv.* **2011**, *144*, 1816–1825. [\[CrossRef\]](#)
261. Boscolo, D.; Tokumoto, P.M.; Ferreira, P.A.; Ribeiro, J.W.; Santos, J.S. dos. Positive Responses of Flower Visiting Bees to Landscape Heterogeneity Depend on Functional Connectivity Levels. *Perspect. Ecol. Conserv.* **2017**, *15*, 18–24. [\[CrossRef\]](#)
262. Klein, A.M.; Vaissière, B.E.; Cane, J.H.; Steffan-Dewenter, I.; Cunningham, S.A.; Kremen, C.; Tschardtke, T. Importance of Pollinators in Changing Landscapes for World Crops. *Proc. R. Soc. B Biol. Sci.* **2007**, *274*, 303–313. [\[CrossRef\]](#) [\[PubMed\]](#)
263. Wojtjowski, P.A. *Agroecological Economics. Sustainability and Biodiversity*; Elsevier: London, UK, 2008.
264. Nicholls, C.I.; Altieri, M.A.; Vazquez, L. Agroecology: Principles for the Conversion and Redesign of Farming Systems. *J. Ecosyst. Ecography* **2016**, *01*. [\[CrossRef\]](#)
265. Altieri, M.A.; Toledo, V.M. The Agroecological Revolution in Latin America: Rescuing Nature, Ensuring Food Sovereignty and Empowering Peasants. *J. Peasant Stud.* **2011**, *38*, 587–612. [\[CrossRef\]](#)

266. Altieri, M.A.; Rosset, P. Agroecology and the Conversion of Large-Scale Conventional Systems to Sustainable Management. *Int. J. Environ. Stud.* **1996**, *50*, 165–185. [[CrossRef](#)]
267. Jacobi, J.; Mathez-Stiefel, S.L.; Gambon, H.; Rist, S.; Altieri, M. Whose Knowledge, Whose Development? Use and Role of Local and External Knowledge in Agroforestry Projects in Bolivia. *Environ. Manage.* **2017**, *59*, 464–476. [[CrossRef](#)]
268. Holt-Giménez, E.; Altieri, M.A. Agroecology, Food Sovereignty, and the New Green Revolution. *Agroecol. Sustain. Food Syst.* **2013**, *37*, 90–102. [[CrossRef](#)]
269. Nicholls, C.I.; Altieri, M.A. Pathways for the Amplification of Agroecology. *Agroecol. Sustain. Food Syst.* **2018**, *42*, 1170–1193. [[CrossRef](#)]
270. Landaverde-González, P.; Quezada-Euán, J.J.G.; Theodorou, P.; Murray, T.E.; Husemann, M.; Ayala, R.; Moo-Valle, H.; Vandame, R.; Paxton, R.J. Sweat Bees on Hot Chillies: Provision of Pollination Services by Native Bees in Traditional Slash-and-Burn Agriculture in the Yucatán Peninsula of Tropical Mexico. *J. Appl. Ecol.* **2017**, *54*, 1814–1824. [[CrossRef](#)] [[PubMed](#)]
271. Catacora-Vargas, G.; Piepenstock, A.; Sotomayor, C.; Cuentas, D.; Cruz, A.; Delgado, F. Brief Historical Review of Agroecology in Bolivia. *Agroecol. Sustain. Food Syst.* **2017**, *41*, 429–447. [[CrossRef](#)]
272. Pinstrup-Andersen, P. Food Security: Definition and Measurement. *Food Secur.* **2009**, *1*, 5–7. [[CrossRef](#)]
273. Altieri, M.A.; Nicholls, C.I. Agroecology: Challenges and Opportunities for Farming in the Anthropocene. *Int. J. Agric. Nat. Resour.* **2020**, *47*, 204–215. [[CrossRef](#)]
274. Matsuda, T.; Wolff, J.; Yanagawa, T. *Risk and the Regulation of New Technologies*; Unive, K., Yanagawa, T., Eds.; Springer & Kobe University: Kobe, Japan, 2021. [[CrossRef](#)]
275. Shaw, A.; Wilson, K. The Bill and Melinda Gates Foundation and the Necro-Populationism of ‘Climate-Smart’ Agriculture. *Gend. Place Cult.* **2020**, *27*, 370–393. [[CrossRef](#)]
276. Shields, M.W.; Johnson, A.C.; Pandey, S.; Cullen, R.; González-Chang, M.; Wratten, S.D.; Gurr, G.M. History, Current Situation and Challenges for Conservation Biological Control. *Biol. Control.* **2019**, 25–35. [[CrossRef](#)]
277. Lattera, P.; Barral, P.; Carmona, A.; Nahuelhual, L. Focusing Conservation Efforts on Ecosystem Service Supply May Increase Vulnerability of Socio-Ecological Systems. *PLoS ONE* **2016**, *11*, 100875. [[CrossRef](#)]
278. Cock, M.J.W. Strategic Entry Points for Funding Taxonomic Support to Agriculture in Developing Countries. CABI Working Paper 3; CABI: Wallingford, UK, 2011; p. 32.
279. Mora, C.; Tittensor, D.P.; Adl, S.; Simpson, A.G.B.; Worm, B. How Many Species Are There on Earth and in the Ocean? *PLoS Biol.* **2011**, *9*, 1–8. [[CrossRef](#)]
280. Cardoso, P.; Erwin, T.L.; Borges, P.A.V.; New, T.R. The Seven Impediments in Invertebrate Conservation and How to Overcome Them. *Biol. Conserv.* **2011**, *144*, 2647–2655. [[CrossRef](#)]
281. Clark, J.A.; May, R.M. Taxonomic Bias in Conservation Research. *Science* **2002**, *297*, 191–192. [[CrossRef](#)]
282. Wilson, E.O. *The Diversity of Life*; Harvard University Press: Cambridge, MA, USA, 1992.
283. Cardoso, P.; Barton, P.S.; Birkhofer, K.; Chichorro, F.; Deacon, C.; Fartmann, T.; Fukushima, C.S.; Gaigher, R.; Habel, J.C.; Hallmann, C.A.; et al. Scientists’ Warning to Humanity on Insect Extinctions. *Biol. Conserv.* **2020**, 242. [[CrossRef](#)]
284. Dewalt, B. Using Indigenous Knowledge to Improve Agriculture and Natural Resource Management. *Hum. Organ.* **1994**, *53*, 123–131. [[CrossRef](#)]
285. Johnson, J.T.; Howitt, R.; Cajete, G.; Berkes, F.; Louis, R.P.; Kliskey, A. Weaving Indigenous and Sustainability Sciences to Diversify Our Methods. *Sustain. Sci.* **2016**, *11*, 1–11. [[CrossRef](#)]
286. Gemmill-Herren, B.; Garibaldi, L.A.; Kremen, C.; Ngo, H.T. Building Effective Policies to Conserve Pollinators: Translating Knowledge into Policy. *Curr. Opin. Insect Sci.* **2021**, *46*, 64–71. [[CrossRef](#)]
287. Lyver, P.; Perez, E.; Carneiro da Cunha, M.; Roué, M. Indigenous and Local Knowledge about Pollination and Pollinators Associated with Food Production: Outcomes from the Global Dialogue Workshop. In *Outcomes from the Global Dialogue Workshop USDA IPBES UNEP UNESCO FAO UNDP*; Lyver, R., Perez, E., Carneiro da Cunha, M., Roué, M., Eds.; United Nations Educational, Scientific and Cultural Organization: Panama City, Panama, 2014; p. 106.
288. Mikkelsen, G.M.; Gonzalez, A.; Peterson, G.D. Economic Inequality Predicts Biodiversity Loss. *PLoS ONE* **2007**, *2*. [[CrossRef](#)]
289. Altieri, M.A. Linking Ecologists and Traditional Farmers in the Search for Sustainable Agriculture. *Front. Ecol. Environ.* **2004**, *2*, 35–42. [[CrossRef](#)]
290. Richard, E. *Bioregionalism and Global Ethics: A Transactional Approach to Achieving Ecological Sustainability, Social Justice, and Human Well-Being*; Routledge, Taylor and Francis Group: New York, NY, USA, 2011. [[CrossRef](#)]
291. Aitken, D.; Rivera, D.; Godoy-Faúndez, A.; Holzapfel, E. Water Scarcity and the Impact of the Mining and Agricultural Sectors in Chile. *Sustainability* **2016**, *8*, 128. [[CrossRef](#)]
292. Hipólito, J.; Coutinho, J.; Mahlmann, T.; Santana, T.B.R.; Magnusson, W.E. Legislation and Pollination: Recommendations for Policymakers and Scientists. *Perspect. Ecol. Conserv.* **2021**, *19*, 1–9. [[CrossRef](#)]
293. Wittman, H. Food Sovereignty: A New Rights Framework for Food and Nature? *Environ. Soc.* **2012**, *2*, 87–105. [[CrossRef](#)]
294. Seminar, A.; Sarwoprasodjo, S.; Santosa, D.; Rilus, A. Agroecological Education Aimed at Achieving Food Sovereignty. *J. Dev. Sustain. Agric.* **2017**, *12*, 34–44. [[CrossRef](#)]
295. Baker, A.M.; Redmond, C.T.; Malcolm, S.B.; Potter, D.A. Suitability of Native Milkweed (*Asclepias*) Species versus Cultivars for Supporting Monarch Butterflies and Bees in Urban Gardens. *PeerJ* **2020**, *8*, e9823. [[CrossRef](#)] [[PubMed](#)]

296. Fuchs, R.; Brown, C.; Rounsevell, M. *Europe's Green Deal Offshores Environmental Damage to Other Nations*; Nature Publishing Group: London, UK, 2020; Volume 586, pp. 671–673. [CrossRef]
297. Garnett, T.; Appleby, M.C.; Balmford, A.; Bateman, I.J.; Benton, T.G.; Bloomer, P.; Burlingame, B.; Dawkins, M.; Dolan, L.; Fraser, D.; et al. Sustainable Intensification in Agriculture: Premises and Policies. *Science* **2013**, *33*, 33–34. [CrossRef] [PubMed]
298. Christiansen, A.; Gervais, J.; Buhl, K.; Stone, D. Acephate General Fact Sheet. 2011. Available online: <http://npic.orst.edu/factsheets/acephagen.html> (accessed on 16 September 2018).
299. Bernal, J.L.; Garrido-Bailon, E.; Del Nozal, J.B.; González-Porto, A.V.; Hernández, R.M.; Diego, J.C.; Jimenez, J.J.; Higes, M. Overview of Pesticide Residues in Stored Pollen and Their Potential Effect on Bee Colony (*Apis mellifera*) Losses in Spain. *J. Econ. Entomol.* **2010**, *103*, 1964–1971. [CrossRef]
300. Boily, M.; Sarrasin, B.; DeBlois, C.; Aras, P.; Chagnon, M. Acetylcholinesterase in honey bees (*Apis mellifera*) exposed to neonicotinoids, atrazine and glyphosate: Laboratory and field experiments. *Environ. Sci. Pollut. Res.* **2013**, *20*, 5603–5614. [CrossRef]
301. Araújo, R.D.S.; Bernardes, R.C.; Martins, G.F. A mixture containing the herbicides Mesotrione and Atrazine imposes toxicological risks on workers of *Partamona helleri*. *Sci. Total. Environ.* **2021**, *763*, 142980. [CrossRef]
302. Williams, J.R. Biomarkers of Oxidative Stress in Atrazine-Treated Honey Bees: A Laboratory and In-Hive Study. Ph.D. Thesis, Virginia Tech, Blacksburg, VA, USA, 2016; p. 78. Available online: <http://hdl.handle.net/10919/72949> (accessed on 1 June 2021).
303. North Carolina Department of Agriculture & Consumer Services. Pesticide Toxicity to Bees “Traffic Light”. 2016. Available online: <http://www.ncagr.gov/pollinators/documents/BeePesticideRiskTrafficLight3-2-17.pdf> (accessed on 1 June 2021).
304. European Food Safety Authority. Conclusion regarding the peer review of the pesticide risk assessment of the active substance cadusafos. *EFSA Sci. Rep.* **2006**, *1*–70.
305. Bond, C.; Hallman, A.; Buhl, K.; Stone, D. Carbaryl General Fact Sheet. 2016. Available online: <http://npic.orst.edu/factsheets/carbarylgen.html> (accessed on 16 September 2018).
306. Shi, T.; Burton, S.; Zhu, Y.; Wang, Y.; Xu, S.; Yu, L. Effects of Field-Realistic Concentrations of Carbendazim on Survival and Physiology in Forager Honey Bees (Hymenoptera: Apidae). *J. Insect Sci.* **2018**, *18*, 1–5. [CrossRef] [PubMed]
307. Böhme, F.; Bischoff, G.; Zebitz, C.P.W.; Rosenkranz, P.; Wallner, K. Pesticide residue survey of pollen loads collected by honeybees (*Apis mellifera*) in daily intervals at three agricultural sites in South Germany. *PLoS ONE* **2018**, *13*, e0199995. [CrossRef] [PubMed]
308. Marletto, F.; Patetta, A.; Manino, A. Laboratory assessment of pesticide toxicity to bumblebees. *Bull. Insectology* **2003**, *56*, 155–158.
309. Kegley, S.E.; Hill, B.R.; Orme, S.; Choi, A.H. Cartap Monohydrochloride. 2016. Available online: http://www.pesticideinfo.org/Detail_Chemical.jsp?Rec_Id=PC38464 (accessed on 16 September 2018).
310. Rhodes, J.; Scott, M. *Pesticides: A Guide to Their Effects on Honey Bees*; New South Wales Department of Primary Industries: Orange, Australia, 2006.
311. Fisher, A., II; Coleman, C.; Hoffmann, C.; Fritz, B.; Rangel, J. Apiculture & Social Insects the Synergistic Effects of Almond Protection Fungicides on Honey Bee (Hymenoptera: Apidae) Forager Survival. *J. Econ. Entomol.* **2017**, *110*, 802–808. [PubMed]
312. Mackenzie, K.A.E.; Winston, A.L. Effects of Sublethal Exposure to Diazinon on Longevity and Temporal Division of Labor in the Honey Bee (Hymenoptera: Apidae). *J. Econ. Entomol.* **1989**, *82*, 75–82. [CrossRef]
313. Weick, J.; Thorn, R.S. Effects of acute sublethal exposure to coumaphos or diazinon on acquisition and discrimination of odor stimuli in the honey bee (Hymenoptera: Apidae). *J. Econ. Entomol.* **2002**, *95*, 227–236. [CrossRef]
314. Gromisz, M. Gastric toxicity to bees of fenopatratin—The active ingredient of Danitol and Danirun formulas. *J. Apic. Sci.* **2001**, *45*, 51–60.
315. National Research Council of Canada, Associate Committee on Scientific Criteria for Environmental Quality. *Pesticide–pollinator interactions*. NCCC, Publication N° 18477 of the Environmental Secretariat; Subcommittee on Pesticides and Industrial Organic Chemicals: Ottawa, Canada, 1981; p. 190.
316. De Moraes, C.R.; Travençolo, B.A.N.; Carvalho, S.M.; Beletti, M.; Santos, V.S.V.; Campos, E.; Júnior, E.O.D.C.; Pereira, B.B.; Naves, M.P.C.; De Rezende, A.A.A.; et al. Ecotoxicological effects of the insecticide fipronil in Brazilian native stingless bees *Melipona scutellaris* (Apidae: Meliponini). *Chemosphere* **2018**, *206*, 632–642. [CrossRef]
317. El Hassani, A.K.; Dacher, M.; Gauthier, M.; Armengaud, C. Effects of sublethal doses of fipronil on the behavior of the honeybee (*Apis mellifera*). *Pharmacol. Biochem. Behav.* **2005**, *82*, 30–39. [CrossRef]
318. Mayer, D.F.; Lunden, J.D. Effects of imidacloprid insecticide on three bee pollinators. *Hortic Sci* **1997**, *29*, 93–97.
319. Pisa, L.W.; Amaral-Rogers, V.; Belzunces, L.P.; Bonmatin, J.M.; Downs, C.A.; Goulson, D.; Kreutzweiser, D.P.; Krupke, C.; Liess, M.; McField, M.; et al. Effects of neonicotinoids and fipronil on non-target invertebrates. *Environ. Sci. Pollut. Res.* **2015**, *22*, 68–102. [CrossRef]
320. European Food Safety Authority. Conclusion regarding the peer review of the pesticide risk assessment of the active substance glufosinate finalized: 14 March 2005. *Eur. Food Safety Auth. J.* **2005**. [CrossRef]
321. European Food Safety Authority. Peer Review of the Pesticide Risk Assessment of the Active Substance Imazamox. European Food Safety Authority Journal 14:n/a-n/a. 2016. Available online: <http://10.0.11.87/j.efsa.2016.4432%5Cnhttp://search.ebscohost.com/login.aspx?direct=true&AuthType=ip,url,cookie,uid&db=aph&AN=115160811&site=ehost-live&scope=site> (accessed on 1 June 2021).
322. Abrol, D.P.; Andotra, R.S. Field Toxicity of Pesticides to Honeybee, *Apis mellifera* L., Foragers. *Journal of Apiculture* **2001**, *1*, 19–26. Available online: <http://dev02.dbpia.co.kr/0/95/59/955925.pdf?article=NODE00955926> (accessed on 1 June 2021).

323. Chauzat, M.; Faucon, J. Pesticide residues in beeswax samples collected from honey bee colonies (*Apis mellifera* L.) in France. *Pest. Manag. Sci.* **2007**, *1106*, 1100–1106. [CrossRef]
324. EPA. Reregistration Eligibility Decision for Methidathion. United States Environmental Protection Agency: Washington, DC, USA, 2006. Available online: <http://www.epa.gov/opp00001/methods/atmpmethods/QC-23-01.pdf> (accessed on 1 June 2021).
325. Fine, J.D.; Mullin, C.A.; Frazier, M.T.; Reynolds, R.D. Field Residues and Effects of the Insect Growth Regulator Novaluron and Its Major Co-Formulant N-Methyl-2-Pyrrolidone on Honey Bee Reproduction and Development. *J. Econ. Entomol.* **2017**, *110*, 1993–2001. [CrossRef] [PubMed]
326. Cousin, M.; Silva-Zacarin, E.; Kretzschmar, A.; El Maataoui, M.; Brunet, J.L.; Belzunces, L.P. Size Changes in Honey Bee Larvae Oenocytes Induced by Exposure to Paraquat at Very Low Concentrations. *PLoS ONE* **2013**, *8*. [CrossRef] [PubMed]
327. Moffett, J.O.; Morton, H.L.; Macdonald, R.H. Toxicity of Herbicides to Newly Emerged Honey Bees. *Environ. Entomol.* **1972**, *1*, 102–104. [CrossRef]
328. Cox, R.L.; Wilson, W.T. Effects of Permethrin on the Behavior of Individually Tagged Honey Bees, *Apis mellifera* L. (Hymenoptera: Apidae). *Environ. Entomol.* **1984**, *13*, 375–378. [CrossRef]
329. Hagler, J.R.; Waller, G.D.; Lewis, B.E. Mortality of honeybees (Hymenoptera: Apidae) exposed to permethrin and combinations of permethrin with piperonyl butoxide. *J. Apic. Res.* **1989**, *28*, 208–211. [CrossRef]
330. Sanchez-Bayo, F.; Goka, K. Pesticide Residues and Bees—A Risk Assessment. *PLoS ONE* **2014**, *9*, e94482. [CrossRef]
331. FAO. PROCYMIDONE. FAO Specifications for Plant. Protection Products; 2001; p. 97. Available online: http://www.fao.org/fileadmin/templates/agphome/documents/Pests_Pesticides/Specs/procymid.pdf (accessed on 1 June 2021).
332. Melisie, D.; Damte, T.; Thakur, A.K. Effects of some insecticidal chemicals under laboratory condition on honeybees [*Apis mellifera* L. (Hymenoptera: Apidae)] that forage on onion flowers. *Environ. Sci. Pollut. Res.* **2015**, *10*, 1295–1300.
333. Stanley, J.; Sah, K.; Jain, S.; Bhatt, J.; Sushil, S. Evaluation of pesticide toxicity at their field recommended doses to honeybees, *Apis cerana* and *A. mellifera* through laboratory, semi-field and field studies. *Chemosphere* **2015**, *119*, 668–674. [CrossRef] [PubMed]
334. Australian Pesticides and Veterinary Medicines Authority. Australian Government 2012. *Public Release Summary on the Evaluation of the New Active Saflufenacil in the Product SHARPEN WG HERBICIDE (Previously Heat Herbicide)*; APVMA Product Number 62853; Australian Pesticides and Veterinary Medicines Authority: Armidale, NSW, Australia, 2012; p. 66.
335. Jimenez, D.R. Efecto letal agudo de los insecticidas en formulación comercial Imidacloprid, Spinosad y Thiocyclam hidrogenoxalato en obreras de *Bombus atratus* (Hymenoptera: Apidae). *Rev. Biol. Trop.* **2016**, *64*, 1737–1745. [CrossRef]
336. Baines, D.; Wilton, E.; Pawluk, A.; De Gorter, M.; Chomistek, N. Neonicotinoids act like endocrine disrupting chemicals in newly-emerged bees and winter bees. *Sci. Rep.* **2017**, *7*, 1–18. [CrossRef]
337. El Hassani, A.K.; Dacher, M.; Gary, V.; Lambin, M.; Gauthier, M.; Armengaud, C. Effects of Sublethal Doses of Acetamiprid and Thiamethoxam on the Behavior of the Honeybee (*Apis mellifera*). *Arch. Environ. Contam. Toxicol.* **2007**, *54*, 653–661. [CrossRef]
338. EPA. Pesticide Fact Sheet: Dinotefuran. United States Environmental Protection Agency: Washington, DC, USA, 2004. Available online: https://www3.epa.gov/pesticides/chem_search/reg_actions/registration/fs_PC-044312_01-Sep-04.pdf (accessed on 1 June 2021).
339. Williamson, S.M.; Willis, S.J.; Wright, G.A. Exposure to neonicotinoids influences the motor function of adult worker honeybees. *Ecotoxicology* **2014**, *23*, 1409–1418. [CrossRef]
340. Schneider, C.W.; Tautz, J.; Grünwald, B.; Fuchs, S. RFID Tracking of Sublethal Effects of Two Neonicotinoid Insecticides on the Foraging Behavior of *Apis mellifera*. *PLoS ONE* **2012**, *7*, e30023. [CrossRef]
341. Henry, M.; Béguin, M.; Requier, F.; Rollin, O.; Odoux, J.F.; Aupinel, P.; Aptel, J.; Tchamitchian, S.; Decourtye, A. A common pesticide decreases foraging success and survival in honey bees. *Science* **2012**, *336*, 348–350. [CrossRef]
342. Taylor, M.A.; Goodwin, M.; McBrydie, H.; Cox, H.M. Destroying managed and feral honey bee (*Apis mellifera*) colonies to eradicate honey bee pests. *N. Z. J. Crop. Hortic. Sci.* **2007**, *35*, 313–323. [CrossRef]
343. Domingues, C.E.; Abdalla, F.C.; Balsamo, P.J.; Pereira, B.V.; Hausen, M.D.A.; Costa, M.J.; Silva-Zacarin, E.C. Thiamethoxam and picoxystrobin reduce the survival and overload the hepato-nephrotoxic system of the Africanized honeybee. *Chemosphere* **2017**, *186*, 994–1005. [CrossRef] [PubMed]
344. Kalnins, M.A.; Detroy, B.F. The effect of wood preservative treatment of beehives on honey bees and hive products. *J. Agric. Food Chem.* **1984**, *32*, 1176–1180. [CrossRef]



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Henríquez-Piskulich, PA; Schapheer, C; Vereecken, NJ; Villagra, C

Title:

Agroecological Strategies to Safeguard Insect Pollinators in Biodiversity Hotspots: Chile as a Case Study

Date:

2021-06-14

Citation:

Henríquez-Piskulich, P. A., Schapheer, C., Vereecken, N. J. & Villagra, C. (2021). Agroecological Strategies to Safeguard Insect Pollinators in Biodiversity Hotspots: Chile as a Case Study. *Sustainability*, 13 (12), pp.6728-6728. <https://doi.org/10.3390/su13126728>.

Persistent Link:

<http://hdl.handle.net/11343/281144>

File Description:

Published version

License:

CC BY