



# Draft Genome Sequences of Eight *Campylobacter volucris* Isolates from Freshwater Sources in Victoria, Australia

Agnus M. Davis,<sup>a</sup> David. T. McCarthy,<sup>a</sup> Dieter M. Bulach,<sup>b</sup>  Rebekah Henry<sup>a</sup>

<sup>a</sup>Environmental and Public Health Microbiology Laboratory (EPHM Lab), Department of Civil Engineering, Monash University, Clayton, Victoria, Australia

<sup>b</sup>Microbiological Diagnostic Unit Public Health Laboratory, The Peter Doherty Institute, The University of Melbourne, Parkville, Victoria, Australia

**ABSTRACT** *Campylobacter* spp. can survive and be transmitted from a range of environments. Here, we examine eight draft genome sequences of *Campylobacter volucris*, identified as part of an examination of waterborne *Campylobacter* species. This is the first report of environmental survival of *C. volucris* outside gull species.

**C**ampylobacter-associated infection is often spread to humans via consumption of poultry, beef, lamb, or pork products (1). However, there are increasing reports of environmental pathways (water, soil) to disease transmission (2–5). *Campylobacter* species can be introduced into these environments through direct defecation by zoonotic disease vectors (e.g., birds, dogs, livestock). For example, sampling of the cloaca of black-headed gulls (*Larus ridibundus*) in Sweden revealed carriage of a *Campylobacter lari*-like strain (6). Further DNA analysis identified the bacterium to be a distinct *Campylobacter* species, *Campylobacter volucris*. To date, *C. volucris* has been minimally reported within the literature, with eight published genome sequences primarily associated with black-headed gulls (7).

In this study, waterborne strains were isolated from five freshwater locations across Greater Melbourne (Table 1). The samples underwent membrane filtration onto 0.45- $\mu\text{m}$  cellulose nitrate filters (Sartorius, Germany). Isolation was undertaken as described in Australian/New Zealand standard 4276.19:2001 (AS/NZS, 2001), with membranes placed into 25 ml Preston broth and resuscitated aerobically for 2 h at 37°C, prior to the addition of *Campylobacter* selective supplement (Oxoid, UK) and microaerobic incubation (5% O<sub>2</sub>, 10% CO<sub>2</sub>, and 85% N<sub>2</sub>) for 48 h at 42°C. Postenrichment (48 h at 42°C), 2  $\mu\text{l}$  of each sample was plated onto modified charcoal-cefoperazone deoxycholate agar (CCDA)-Preston medium and incubated for 48 h at 42°C (Oxoid). The colonies underwent biochemical confirmation using the Oxoid Biochemical Identification System (OBIS) Campy kit, followed by selection on horse blood agar (HBA) (Oxoid) and microaerophilic growth for 48 h at 42°C.

Genomic DNA was extracted using the RBC genomic DNA extraction kit (Real Biotech, Taiwan). The DNA quality and quantity were assessed using a Qubit fluorometer (Thermo Fisher Scientific, Waltham, MA, USA) following the manufacturer's instructions. Sequencing libraries were prepared using 1 ng DNA for the Nextera XT kit (Illumina, USA) and 100 ng DNA for the MGIeasy DNA FS kit (MGI, China). Sequencing of the Nextera XT libraries was conducted on the NextSeq 500 platform (Illumina) using v2 2  $\times$  150-bp paired-end (PE) chemistry, while the MGI libraries underwent sequencing on an MGITech DNBSEQ-G400 (MGI) instrument using the 150PE DNBSEQ high-throughput sequencing set FCL v2 chemistry (MGI). Sequence data were quality processed, analyzed, and *de novo* assembled using the tools within the Nullarbor v2.0 pipeline (8). Default parameters were applied to all software contained therein unless otherwise specified.

The draft genome sequence and assembly statistics are provided in Table 1. *De*

**Citation** Davis AM, McCarthy DT, Bulach DM, Henry R. 2021. Draft genome sequences of eight *Campylobacter volucris* isolates from freshwater sources in Victoria, Australia. *Microbiol Resour Announc* 10:e01086-20. <https://doi.org/10.1128/MRA.01086-20>.

**Editor** Julia A. Maresca, University of Delaware

**Copyright** © 2021 Davis et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](#).

Address correspondence to Rebekah Henry, rebekah.henry@monash.edu.

**Received** 7 October 2020

**Accepted** 9 December 2020

**Published** 7 January 2021

**TABLE 1** Australian *Campylobacter volucris* isolation, sequencing, and genome assembly data

Parameter	Data for strain:						
	MON0058	MON0078	MON0115	MON0166	MON0180	MON1184	MON1267
Sequencing technology	Illumina NextSeq 500 Nextera XT NextSeq 500 v2						
Library chemistry							
Sequencing chemistry							
Read length (bp)	PE150	PE75	PE75	PE75	PE150	PE150	MGITech DNaseQ-G400
No. of reads	2,404,222	3,743,990	913,212	3,969,814	1,203,542	1,195,848	MGIEasy DNA FS
No. of contigs	36	292	61	281	340	28	DNBSEQ high-throughput sequencing set FCL
Genome length (bp)	1,486,257	1,467,362	1,508,747	1,548,798	1,413,139	1,497,020	PE100
N <sub>50</sub> (bp)	83,244	9,015	66,645	11,468	6,494	109,393	5,627,484
Coverage (×)	220	174	83	184	147	112	27
GC content (%)	30.3	33.4	30.8	33.6	32.7	30.7	1,637,888
SRA accession no.	<a href="#">SRX9038520</a>	<a href="#">SRX9038521</a>	<a href="#">SRX9038522</a>	<a href="#">SRX9038523</a>	<a href="#">SRX9038524</a>	<a href="#">SRX9038525</a>	1,519,422
BioSample accession no.	<a href="#">SAMN15944590</a>	<a href="#">SAMN15944591</a>	<a href="#">SAMN15944592</a>	<a href="#">SAMN15944593</a>	<a href="#">SAMN15944594</a>	<a href="#">SAMN15944595</a>	PRJNA660282
BioProject accession no.	<a href="#">PRJNA660282</a>		<a href="#">PRJNA660282</a>		<a href="#">PRJNA660282</a>	<a href="#">PRJNA660282</a>	Urban river
Source	Rural river						
GPS location	37°42'25.1"S, 145°19'19.5"E	37°42'25.1"S, 145°19'19.5"E	37°40'25.3"S, 145°29'20.1"E	37°42'25.1"S, 145°19'19.5"E	37°45'17.7"S, 145°40'46.8"E	37°45'17.7"S, 145°40'46.8"E	37°47'38.2"S, 145°00'06.8"E

*novo* assembly achieved an overall coverage of 135× ( $\pm 43\times$ ), with the strains having an average GC content of 31% ( $\pm 2\%$ ) and size of 1.5 Mb ( $\pm 62$  kb). These characteristics are similar to those previously reported for *C. volucris* isolates.

This study reports the draft genome sequence for eight *C. volucris* isolates from waterways in Victoria, Australia; this is the first non-host-associated isolation of this species. Thus, this report extends our understanding of the genetic diversity within the species and is the first demonstration that this organism has the capacity to exist in environmental waters. We also note that black-headed gulls are not part of the indigenous fauna of the region, indicating a broader avian host range, possibility associated with other waterfowl species.

**Data availability.** The genome sequences and raw data have been deposited in GenBank within BioProject accession number [PRJNA660282](#), under the accession numbers outlined in Table 1.

## ACKNOWLEDGMENTS

We gratefully acknowledge the funding and data contribution from Melbourne Water and the Australian Research Council (LP160100408). We also acknowledge use of the services and facilities of Micromon Genomics at Monash University.

## REFERENCES

- Varrone L, Stafford RJ, Lilly K, Selvey L, Glass K, Ford L, Bulach D, Kirk MD, The CampySource Project Team. 2018. Investigating locally relevant risk factors for *Campylobacter* infection in Australia: protocol for a case-control study and genomic analysis. *BMJ Open* 8:e026630. <https://doi.org/10.1136/bmjopen-2018-026630>.
- Shrestha RD, Midwinter AC, Marshall JC, Collins-Emerson JM, Pleydell EJ, French NP. 2019. *Campylobacter jejuni* strains associated with wild birds and those causing human disease in six high-use recreational waterways in New Zealand. *Appl Environ Microbiol* 85:e01228-19. <https://doi.org/10.1128/AEM.01228-19>.
- Nilsson A, Johansson C, Skarp A, Kaden R, Bertilsson S, Rautelin H. 2018. Survival of *Campylobacter jejuni* and *Campylobacter coli* water isolates in lake and well water. *APMIS* 126:762–770. <https://doi.org/10.1111/apm.12879>.
- Gibney KB, O'Toole J, Sinclair M, Leder K. 2017. Burden of disease attributed to waterborne transmission of selected enteric pathogens, Australia, 2010. *Am J Trop Med Hyg* 96:1400–1403. <https://doi.org/10.4269/ajtmh.16-0907>.
- Mughini-Gras L, Penny C, Ragimbeau C, Schets FM, Blaak H, Duim B, Wagenaar JA, de Boer A, Cauchie H-M, Mossong J, van Pelt W. 2016. Quantifying potential sources of surface water contamination with *Campylobacter jejuni* and *Campylobacter coli*. *Water Res* 101:36–45. <https://doi.org/10.1016/j.watres.2016.05.069>.
- Debruyne L, Broman T, Bergstrom S, Olsen B, On SLW, Vandamme P. 2010. *Campylobacter volucris* sp. nov., isolated from black-headed gulls (*Larus ridibundus*). *Int J Syst Evol Microbiol* 60:1870–1875. <https://doi.org/10.1099/ijsm.0.013748-0>.
- Kweon OJ, Lim YK, Yoo B, Kim HR, Kim T-H, Lee M-K. 2015. First case report of *Campylobacter volucris* bacteremia in an immunocompromised patient. *J Clin Microbiol* 53:1976–1978. <https://doi.org/10.1128/JCM.00442-15>.
- Seemann T, Goncalves da Silva A, Bulach DM, Schultz MB, Kwong JC, Howden BP. 2018. Nullarbor. <https://github.com/tseemann/nullarbor>.

# University Library



MINERVA  
ACCESS

A gateway to Melbourne's research publications

Minerva Access is the Institutional Repository of The University of Melbourne

**Author/s:**

Davis, AM; McCarthy, DT; Bulach, DM; Henry, R

**Title:**

Draft Genome Sequences of Eight *Campylobacter volucris* Isolates from Freshwater Sources in Victoria, Australia

**Date:**

2021-01-01

**Citation:**

Davis, A. M., McCarthy, D. T., Bulach, D. M. & Henry, R. (2021). Draft Genome Sequences of Eight *Campylobacter volucris* Isolates from Freshwater Sources in Victoria, Australia. *MICROBIOLOGY RESOURCE ANNOUNCEMENTS*, 10 (1), <https://doi.org/10.1128/MRA.01086-20>.

**Persistent Link:**

<http://hdl.handle.net/11343/281659>

**File Description:**

Published version

**License:**

CC BY