

DR. MICHAEL R. RICKELS (Orcid ID : 0000-0002-9253-838X)

DR. LORENZO PIEMONTE (Orcid ID : 0000-0002-2172-2198)

PROF. THIERRY BERNEY (Orcid ID : 0000-0002-4230-9378)

Article type : Review Article

## **Defining Outcomes for $\beta$ -Cell Replacement Therapy in the Treatment of Diabetes: a Consensus Report on the Igls Criteria from the IPITA/EPITA Opinion Leaders Workshop**

Michael R. Rickels, MD,<sup>1</sup> Peter G. Stock, MD, PhD,<sup>2</sup> Eelco J. P. de Koning, MD, PhD,<sup>3</sup> Lorenzo Piemonti, MD,<sup>4</sup> Johann Pratschke, MD, PhD,<sup>5</sup> Rodolfo Alejandro, MD,<sup>6</sup> Melena D. Bellin, MD,<sup>7</sup> Thierry Berney, MD,<sup>8</sup> Pratik Choudhary, MD,<sup>9</sup> Paul R. Johnson, MD,<sup>10</sup> Raja Kandaswamy, MD,<sup>11</sup> Thomas W. H. Kay, MBBS, PhD,<sup>12</sup> Bart Keymeulen, MD, PhD,<sup>13</sup> Yogish C. Kudva, MBBS,<sup>14</sup> Esther Latres, PhD,<sup>15</sup> Robert M. Langer, MD, PhD,<sup>16</sup> Roger Lehmann, MD,<sup>17</sup> Barbara Ludwig, MD, PhD,<sup>18</sup> James F. Markmann, MD, PhD,<sup>19</sup> Marjana Marinac, PharmD,<sup>15</sup> Jon S. Odorico, MD,<sup>20</sup> François Pattou, MD,<sup>21</sup> Peter A. Senior, MBBS, PhD,<sup>22</sup> James A. M. Shaw, FRCP, PhD,<sup>23</sup> Marie-Christine Vantyghem, MD, PhD,<sup>24</sup> and Steven White, MD<sup>23</sup>

<sup>1</sup>Division of Endocrinology, Diabetes & Metabolism, Department of Medicine, and Institute for Diabetes, Obesity & Metabolism, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA.

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/tri.13138](https://doi.org/10.1111/tri.13138)

This article is protected by copyright. All rights reserved

<sup>2</sup>Division of Transplantation, Department of Surgery, University of California at San Francisco, San Francisco, CA.

<sup>3</sup>Department of Medicine, Leiden University Medical Center, Leiden, The Netherlands.

<sup>4</sup>Diabetes Research Institute, San Raffaele Scientific Institute, Milan, Italy.

<sup>5</sup>Department of Surgery, Charité Medical School Berlin, Berlin, Germany

<sup>6</sup>Division of Endocrinology, Diabetes & Metabolism, Department of Medicine, and Diabetes Research Institute, University of Miami Miller School of Medicine, Miami, FL.

<sup>7</sup>Division of Endocrinology, Department of Pediatrics, and the Schulze Diabetes Institute, University of Minnesota, Minneapolis, MN.

<sup>8</sup>Division of Transplantation and Visceral Surgery, Department of Surgery, Geneva University Hospital, Geneva, Switzerland.

<sup>9</sup>Diabetes Research Group, King's College London, London, United Kingdom

<sup>10</sup>Nuffield Department of Surgical Sciences, University of Oxford, Oxford, United Kingdom.

<sup>11</sup>Division of Transplantation, Department of Surgery, and the Schulze Diabetes Institute, University of Minnesota, Minneapolis, MN.

<sup>12</sup>Department of Medicine, St. Vincent's Hospital, and St. Vincent's Institute of Medical Research, University of Melbourne, Melbourne, Victoria, Australia.

<sup>13</sup>Diabetes Research Center, Vrije Universiteit Brussel, Brussels, Belgium.

<sup>14</sup>Division of Endocrinology, Diabetes, Metabolism & Nutrition, Department of Internal Medicine, Mayo Clinic, Rochester, MN

<sup>15</sup>JDRF International, New York, NY.

<sup>16</sup>Ordensklinikum Elisabethin Hospital, Linz, Austria.

<sup>17</sup>Department of Endocrinology and Diabetology, University Hospital Zurich, Zurich, Switzerland.

<sup>18</sup>Division of Endocrinology and Diabetes, Department of Medicine III, University Hospital Carl Gustav Carus at the Technische Universität Dresden, Dresden, Germany.

<sup>19</sup>Division of Transplantation, Department of Surgery, Massachusetts General Hospital, Boston, MA.

<sup>20</sup>Division of Transplantation, Department of Surgery, University of Wisconsin School of Medicine and Public Health, Madison, WI.

<sup>21</sup>Department of General and Endocrine Surgery, Centre Hospitalier Universitaire de Lille, and Inserm, Université de Lille, Lille, France.

<sup>22</sup>Division of Endocrinology & Metabolism, Department of Medicine, University of Alberta, Edmonton, Alberta, Canada.

<sup>23</sup>Institute of Transplantation, The Freeman Hospital and Newcastle University, Newcastle upon Tyne, United Kingdom.

<sup>24</sup>Department of Endocrinology, Diabetology and Metabolism, Centre Hospitalier Universitaire de Lille, and Inserm, Université de Lille, Lille, France.

Address correspondence and requests for reprints to: Michael R. Rickels, MD, MS, University of Pennsylvania Perelman School of Medicine, 12-134 Smilow Center for Translational Research, 3400 Civic Center Boulevard, Philadelphia, PA 19104-5160, USA; phone: (215)746-0025; fax (215)898-5408; e-mail: [rickels@penncmedicine.upenn.edu](mailto:rickels@penncmedicine.upenn.edu)

M.R.R. and P.G.S. co-chaired the organizing committee, presented at, chaired, and moderated sessions, and wrote the manuscript; E.J.P.de K., L.P., and J.P. served on the organizing committee, presented at, chaired, and/or moderated sessions, and reviewed/edited the manuscript; R.A., M.D.B., T.B., P.C., P.R.J., R.K. T.W.H.K., B.K., Y.C.K., E.L., R.M.L., R.L., B.L., J.F.M., M.M., J.S.O., F.P., P.A.S., J.A.M.S., M.-C.V., and S.W. presented at, chaired, and/or moderated sessions, and reviewed/edited the manuscript. All authors and the additional workshop participants listed in the Appendix approved the manuscript.

The workshop was supported in part by educational grants received from The Transplantation Society and JDRF International. M.R.R. is supported in part by U.S. Public Health Services research grants R01 DK091331, R01 DK97830, and U01 DK070430. Y.C.K. is supported in part by U.S. Public Health Services research grant UC4 DK108483.

The authors thank Sondra Livingston of The Transplantation Society and Chiara Parisotto and Giovanna Rossi of the European Society for Organ Transplantation for their assistance with organization of the workshop, and Aristeia Slikas of the University of Pennsylvania Institute for Diabetes, Obesity & Metabolism for assistance with preparation of the manuscript.

The authors declare no conflicts of interest relevant to this work.

Word count: 3,284

Tables/Figures: 3/0 (out of 3 total allowed)

Abbreviations:

AP	Artificial pancreas (closed loop automated insulin delivery)
CFRD	Cystic Fibrosis Related Diabetes
CGM	Continuous glucose monitoring
CV	Coefficient of variation (= SD/mean)
EPITA	European Pancreas & Islet Transplant Association
HbA <sub>1c</sub>	Glycated hemoglobin
IPITA	International Pancreas & Islet Transplant Association
JDRF	Juvenile Diabetes Research Foundation
MMTT	Mixed-meal tolerance test
OGTT	Oral glucose tolerance test
PRO	Patient-reported outcome
SD	Standard deviation
SMBG	Self-monitoring blood glucose
T1D	Type 1 Diabetes
T2D	Type 2 Diabetes

# Author Manuscript

**Abstract:**  $\beta$ -cell replacement therapy, available currently as pancreas or islet transplantation, has developed without a clear definition of graft functional and clinical outcomes. The International Pancreas & Islet Transplant Association (IPITA) and European Pancreas & Islet Transplantation

Association (EPITA) held a workshop to develop consensus for an IPITA/EPITA Statement on the definition of function and failure of current and future forms of  $\beta$ -cell replacement therapy. There was consensus that  $\beta$ -cell replacement therapy could be considered as a treatment for  $\beta$ -cell failure, regardless of etiology and without requiring undetectable C-peptide, accompanied by glycemic instability with either problematic hypoglycemia or hyperglycemia. Glycemic control should be assessed at a minimum by glycated hemoglobin (HbA<sub>1c</sub>) and the occurrence of severe hypoglycemia. Optimal  $\beta$ -cell graft function is defined by near-normal glycemic control (HbA<sub>1c</sub>  $\leq$ 6.5% [48 mmol/mol]) without severe hypoglycemia or requirement for insulin or other antihyperglycemic therapy, and with an increase over pre-transplant measurement of C-peptide. Good  $\beta$ -cell graft function requires HbA<sub>1c</sub>  $<$ 7.0% (53 mmol/mol) without severe hypoglycemia and with a significant ( $>$ 50%) reduction in insulin requirements and restoration of clinically significant C-peptide production. Marginal  $\beta$ -cell graft function is defined by failure to achieve HbA<sub>1c</sub>  $<$ 7.0% (53 mmol/mol), the occurrence of any severe hypoglycemia, or less than 50% reduction in insulin requirements when there is restoration of clinically significant C-peptide production documented by improvement in hypoglycemia awareness/severity, or glycemic variability/lability. A failed  $\beta$ -cell graft is defined by the absence of any evidence for clinically significant C-peptide production. Optimal and good function are considered successful clinical outcomes.

[250/250 words]

Pancreas and islet transplantation are established approaches for providing  $\beta$ -cell replacement therapy in the treatment of diabetes, and stem-cell derived and xenogeneic sources of islet cell tissue for transplantation have entered early-phase clinical trials. Understanding the therapeutic effectiveness of existing and future forms of  $\beta$ -cell replacement therapy is currently limited by the lack of a clear definition of graft functional and clinical outcomes. Moreover, glycemic control metrics have been poorly aligned with the field of artificial pancreas (AP) development. This limitation was identified as a significant barrier to progress in the field of pancreas and islet transplantation at the IPITA-TTS Opinion Leaders Meeting on the Future of  $\beta$ -Cell Replacement.<sup>1,2</sup> As AP systems become available that promise to provide improved glycemic control, similar metrics for assessing glycemic control are needed to compare effectiveness across  $\beta$ -cell replacement and AP approaches. The current lack of clear definitions for clinical

success or failure of available  $\beta$ -cell replacement therapies and glycemic metrics has impacted acceptance from the endocrinology community that has turned attention away from cellular treatment with potential to cure diabetes in hopes that a technologic solution may provide acceptable glycemic control for most patients. Only with comparable methods of assessment for the various approaches to achieving glycemic control available now and in the future can we identify those patients most likely to derive benefit from each type of therapy.

In order to address the lack of standardized outcome definitions for  $\beta$ -cell replacement therapy, the International Pancreas & Islet Transplant Association (IPITA) joined with the European Pancreas & Islet Transplant Association (EPITA) for a two-day workshop on “Defining Outcomes for  $\beta$ -Cell Replacement Therapy in the Treatment of Diabetes” in January 2017 in Igls, Austria. The workshop objectives were to: develop consensus for an IPITA/EPITA Statement on the definition of function and failure of current and future forms of  $\beta$ -cell replacement therapies; review the metabolic and immunologic outcome measures used to select patients and assess the efficacy of  $\beta$ -cell replacement therapies and guide therapeutic decisions; ensure consistency of definitions for glycemic control metrics with the field of artificial pancreas device development; and build a network of collaborators to foster scientific synergy in the clinical investigation of various  $\beta$ -cell replacement and artificial insulin delivery approaches to diabetes.

In order to review relevant information required to formulate a consensus definition for functional and clinical outcomes for  $\beta$ -cell replacement therapy, individual sessions were designed with specified objectives (Table 1). Historically, success in pancreas transplantation has been defined by independence from exogenous insulin, without consideration of the resultant degree of glycemic control, while in islet transplantation success has been defined by near-normal glycemic control determined by glycated hemoglobin ( $HbA_{1c}$ ) in the absence of severe hypoglycemia. Recently, JDRF International (formerly known as the Juvenile Diabetes Research Foundation) led an initiative to identify and define clinically meaningful outcomes for patients with type 1 diabetes (T1D) beyond  $HbA_{1c}$ , prioritizing standardization of outcomes such as hypoglycemia, hyperglycemia, time in range (based on continuous glucose monitoring [CGM]), and diabetic ketoacidosis. This T1D Outcomes Program also evaluated patient reported outcomes (PROs) but existing evidence were not able to support the selection of any specific PRO for the assessment of T1D-related care or research.<sup>3</sup> The T1D research community is also

emphasizing the need to assess benefit beyond reduction in HbA<sub>1c</sub>, arguing that even an increase in HbA<sub>1c</sub> may be acceptable with an AP system if previously frequent hypoglycemia was improved.<sup>4</sup> With the International Hypoglycemia Study Group providing further consensus on definitions of hypoglycemia for clinical trials,<sup>5</sup> the evaluation of hypoglycemia in addition to some average metric of glycemic control such as HbA<sub>1c</sub> will be necessary for the selection of patients for and assessment of all forms of  $\beta$ -cell replacement and AP therapies, as has already been established for islet transplantation.<sup>6</sup> Moreover, consistent outcomes definitions are needed for quality assurance in the performance assessment of programs offering various forms of  $\beta$ -cell replacement and AP therapies.

### **Indications for and Approaches to $\beta$ -cell Replacement Therapy**

The principal indications for  $\beta$ -cell replacement therapy have been to treat insulin-dependent patients (T1D and insulin-requiring type 2 diabetes [T2D]) with end-stage renal disease or experiencing problematic hypoglycemia.<sup>7</sup> Success following a pancreas or islet transplant has been judged in part by the elimination of insulin requirements; however, discontinuation of insulin should not be at the expense of sub-optimal glycemic control. A reasonable expectation for insulin-independence is the maintenance of non-diabetic levels of glycemic control (HbA<sub>1c</sub>  $\leq$  6.5% [48 mmol/mol]) off exogenous insulin or other antihyperglycemic therapy.<sup>8</sup> Importantly, use of insulin or other antihyperglycemic therapy following pancreas or islet transplantation is not synonymous with graft loss or failure, as patients may require low doses of exogenous insulin or other glucose lowering agents to maintain glycemic control in the non-diabetic range, which is only possible to achieve when a portion of the insulin requirement is provided endogenously from a functioning graft.

Such “partial” function of a  $\beta$ -cell replacement therapy has been viewed as successful when particular challenges in glycemic control, such as the occurrence of severe hypoglycemia, are eliminated following restoration of endogenous insulin secretion. Indeed, patients with problematic hypoglycemia, defined by two or more episodes per year of severe hypoglycemia or as one episode in the context of impaired awareness of hypoglycemia, extreme glycemic lability, or major fear and maladaptive behavior, should be considered for either pancreas or islet transplantation.<sup>7</sup> Other patients to consider are those with problematic hyperglycemia, defined by the presence of recurrent episodes of diabetic ketoacidosis or severe, rapidly progressing



secondary complications of diabetes. All patients should have completed a structured education program on basal-bolus insulin delivery with flexible dosing of modern insulin analogs using pump or multi-dose injection delivery based on frequent self-monitoring of blood glucose (SMBG) with or without continuous glucose monitoring (CGM).

A unifying concept is the consideration of  $\beta$ -cell replacement therapy as treatment for  $\beta$ -cell failure, regardless of etiology, when  $\beta$ -cell failure is associated with glycemic instability and either problematic hypoglycemia or hyperglycemia despite availability of and adherence to optimized medical care. This allows consideration of candidates beyond T1D to include some with advanced insulinopenic T2D, or any cause of insulin-deficient diabetes such as cystic fibrosis-related diabetes and other pancreatogenic forms of diabetes (e.g. chronic pancreatitis or following pancreatectomy). While it is expected that C-peptide levels in such individuals would be low, the importance of assessing C-peptide levels (as well as insulin requirements) is to identify elevated levels consistent with insulin resistance that might impart stress on a  $\beta$ -cell graft and compromise the potential for benefit from replacement therapy. Undetectable levels of C-peptide, while making it easier to attribute post-transplant C-peptide to graft function, should not be required. Thus, levels of C-peptide should be measured prior to transplantation in order to determine post-transplant graft function.

### **Outcome Measures of Glucose Homeostasis**

Average glycemic control, particularly over the long-term, remains best assessed by measurement of HbA<sub>1c</sub>. However, for shorter term assessment of average glycemia, mean glucose can be assessed from frequent SMBG (most valid with 5 times daily monitoring)<sup>9</sup> or CGM. Average blood or CGM glucose can be used to estimate the HbA<sub>1c</sub> under situations such as marked anemia or use of dapson<sup>10</sup> when the HbA<sub>1c</sub> is not accurate. While there is inter-individual variability in the mean glucose-HbA<sub>1c</sub> relationship, the relationship within an individual is very reproducible and most influenced by the prior month of glycemia.<sup>11</sup> Consistency of average glucose measures depends, however, on the duration of observation and becomes most reliable with 14 or more days. The frequency and duration of SMBG and CGM are also important for measures of glycemic variability, which are readily assessed from the SD

of glucose measurements or glucose CV ( $=SD/mean$ ). The glucose SD has been validated against clinical assessment of glycemic lability using only 48 hours of CGM data.<sup>12</sup> Glycemic lability incorporates the temporal aspect to glycemic variability and may also be assessed by the glycemic lability index (LI) using at least 4 times daily SMBG over a 4 week period.<sup>13,14</sup> LI has been validated against clinical assessment of glycemic lability,<sup>13</sup> and is highly reproducible over time.<sup>14</sup> Glucose time in range, available only from CGM and being promoted for shorter-term assessment of AP systems,<sup>4</sup> requires further study to understand and validate its use.

The most important measure of hypoglycemia is the occurrence of severe hypoglycemia, defined as an event associated with loss of consciousness or requiring third party assistance for recovery.<sup>15</sup> A recent history of experiencing severe hypoglycemia, impaired awareness of hypoglycemia, and marked glycemic lability are established risk factors for experiencing future severe hypoglycemia. Thus, problematic hypoglycemia has been defined as two or more episodes per year of severe hypoglycemia or as one episode associated with impaired awareness of hypoglycemia, extreme glycemic lability, or major fear and maladaptive behavior.<sup>7</sup> Impaired awareness of hypoglycemia is assessed by validated questionnaires concerning the glucose threshold at which symptom recognition occurs, with the Clarke survey assessing thresholds at both 50 and 60 mg/dl (2.8 and 3.3 mmol/l)<sup>16</sup> and the Gold survey assessing thresholds at an intermediate 54 mg/dl (3.0 mmol/l).<sup>17</sup> Both questionnaires provide a score up to 7 with a score of  $\geq 4$  indicating impaired awareness of hypoglycemia that is highly correlated, supporting either survey as an appropriate assessment tool even if not directly comparable. While more laborious to collect and requiring 4 weeks of prospective diary keeping together with SMBG records, the HYPO score also captures hypoglycemia severity by tabulating the frequency and associated symptoms of, and assistance required for treating a glucose level of  $<54$  mg/dl ( $<3.0$  mmol/l). The HYPO score can be used to identify those with problematic hypoglycemia,<sup>13</sup> and is reproducible.<sup>14</sup> More practically, the frequency of episodes or percent time with glucose  $<54$  mg/dl ( $<3$  mmol/l) can be assessed using either SMBG or CGM. The International Hypoglycemia Study Group defined a glucose level  $<54$  mg/dl ( $<3$  mmol/l) as sufficiently low to indicate serious, clinically important hypoglycemia that should be reported in clinical trials.<sup>5</sup>

The goal, then, for glycemic control outcomes of  $\beta$ -cell replacement therapies should be attainment of target levels of HbA<sub>1c</sub>  $<7.0\%$  (53 mmol/mol), and ideally near-normal HbA<sub>1c</sub>  $\leq 6.5\%$  (48 mmol/mol), in the absence of severe hypoglycemia (Table 2). Additional goals may

be driven by the indication for treatment: impaired awareness of hypoglycemia (Clarke or Gold score  $\geq 4$ ) should be resolved (score  $< 4$ ), serious, clinically important hypoglycemia (glucose  $< 54$  mg/dl [ $< 3$  mmol/l]) should be lessened or eliminated; marked glycemic variability or lability should be improved. Where CGM data are available, time with serious hypoglycemia (glucose  $< 54$  mg/dl [ $< 3$  mmol/l]), time with any hypoglycemia ( $< 70$  mg/dl [ $< 3.9$  mmol/l]), time on-target (70 – 140 mg/dl [3.9 – 7.8 mmol/l]), time-in-range (70 – 180 mg/dl [3.9 – 10 mmol/l] or 54 – 180 mg/dl [3 – 10 mmol/l]), and time with any hyperglycemia ( $> 180$  mg/dl [ $> 10$  mmol/l]) should be considered and maybe useful for making comparisons to AP systems.<sup>4</sup> While safety considerations differ between  $\beta$ -cell replacement and AP system approaches, their detailed assessment is critical and qualitative assessment of patient satisfaction will need to be part of future treatment comparisons. In particular, the complicated patient groups so far treated with pancreas and islet transplantation, those with end-stage renal disease or experiencing problematic hypoglycemia, have been excluded from clinical trials of AP systems, and may not derive similar benefit with AP as trial participants with relatively uncomplicated diabetes. Future assessment of AP systems in patients with end-stage renal disease and those with problematic hypoglycemia is needed. Furthermore, use of PROs including health-related quality of life, diabetes distress, and fear of hypoglycemia requires further attention.<sup>18-20</sup>

### **Outcome Measures of $\beta$ -cell Graft Function and Demand**

Both insulin requirements and levels of stimulated C-peptide reflect the contribution of  $\beta$ -cell replacement therapy to the resultant state of glycemic control, but at the same time are dependent on the degree of glycemic control and underlying insulin sensitivity. With improvement in glycemic control, and consequently insulin sensitivity,<sup>21</sup> following  $\beta$ -cell replacement therapy, a reduction in insulin requirements can be attributed to restoration of endogenous insulin secretion from the  $\beta$ -cell graft. However, in the absence of meeting glycemic control targets, a measured reduction in insulin requirements cannot be attributed to the effectiveness of  $\beta$ -cell replacement therapy. Moreover, a patient withdrawn from insulin who is not meeting glycemic control targets as defined above should not be considered insulin-independent, since insulin therapy would be indicated to achieve appropriate glycemic control.

C-peptide levels, when undetectable prior to treatment, can be used to assess function of a  $\beta$ -cell graft, but depend on the metabolic demand for secretion (fasted or stimulated, underlying

insulin sensitivity, and glucose level) and renal clearance. With increasing sensitivity of assays for detection of C-peptide, low levels of questionable clinical significance are often detected (e.g.  $<0.3$  ng/ml [ $<0.1$  nmol/l] fasting or  $<0.6$  ng/ml [ $<0.2$  nmol/l] post-prandial) despite clinical  $\beta$ -cell failure,<sup>22</sup> and may be even higher in the presence of uremia or sub-total  $\beta$ -cell loss (e.g. with cystic fibrosis-related diabetes and advanced insulinopenic T2D). Nevertheless, pre-transplant testing of C-peptide is critical to inform post-transplant monitoring, and should be performed fasting together with a concomitant glucose level with or without stimulated measures. Testing of C-peptide should be done the same way before as after transplant.

$\beta$ -cell replacement therapy aims to restore non-diabetic fasting and post-prandial glucose without hypoglycemia. Oral glucose tolerance can be assessed by a standardized liquid nutrient meal containing a reasonable amount of carbohydrate (~50 grams) in place of the standard 75 gram oral glucose tolerance test (OGTT) used for diagnosis of diabetes. The 90-minute glucose during the standard mixed-meal tolerance test (MMTT) is highly correlated with the 120-minute glucose during the OGTT.<sup>23</sup> The post-transplant ratio of fasting C-peptide-to-glucose is predictive of the 90-minute glucose,<sup>24</sup> and so may allow for more frequent assessment of  $\beta$ -cell graft function while the MMTT may be most useful to resolve uncertainty regarding the interpretation of more routine clinical assessment.

The  $\beta$ -score is a composite measure of  $\beta$ -cell graft function that incorporates the HbA<sub>1c</sub>, insulin requirement, fasting glucose, and C-peptide, and so may be calculated during routine clinical assessment, although C-peptide assessment may require a stimulation test.<sup>25</sup> More recently, the  $\beta$ 2-score models the same variables but requires only the fasting C-peptide and provides a continuous rather than categorical metric.<sup>26</sup> The  $\beta$ -score was initially validated against the 90-minute glucose derived from the MMTT and has also been shown to relate to CGM metrics of mean glucose, glucose variability, time spent with serious, clinically important hypoglycemia ( $<54$  mg/dl [ $3.0$  mmol/l]), and time spent with hyperglycemia ( $>180$  mg/dl [ $10$  mmol/l]).<sup>27</sup> While helpful for longitudinal monitoring, the  $\beta$ -score remains limited by its summative derivation and absence of including a direct measure of hypoglycemia. The  $\beta$ 2-score may have potential utility as a continuous variable rather than the categorical quantification provided by the  $\beta$ -score.

The goal, then, for functional outcomes of  $\beta$ -cell replacement therapies should be at a minimum to achieve a 50% reduction in insulin requirements (and which should be  $<0.5$  units

per kg body weight per day), assuming adequate glycemic control ( $HbA_{1c} < 7.0\%$  [ $53 \text{ mmol/mol}$ ]), that is associated with an increase from pre-transplant measures of C-peptide (and which should be at least  $>0.5 \text{ ng/ml}$  [ $>0.17 \text{ nmol/l}$ ]) interpreted with a concomitant glucose level. More accurate assessment of functional  $\beta$ -cell mass requires determination of glucose-potentiation of insulin or C-peptide release in response to a non-glucose insulin secretagogue such as arginine or glucagon;<sup>22,28,29</sup> however, this gold-standard testing of  $\beta$ -cell secretory capacity is not widely available.

### **Outcome Measures of Immunologic Mechanisms**

While the success of  $\beta$ -cell replacement therapy ultimately depends on the prevention of alloimmune rejection and autoimmune recurrence, and which themselves depend on the source of tissue for transplantation and whether the initial cause of  $\beta$ -cell failure was type 1 (autoimmune) diabetes, the assessment of immune markers was not felt directly relevant to the definition of outcomes, but rather to the understanding of unsuccessful outcomes or declining functional status.

## **CONCLUSION**

### **Defining Successful Outcomes**

There was consensus that categorizing  $\beta$ -cell graft function would not be synonymous with defining the clinical success of a  $\beta$ -cell replacement therapy. A  $\beta$ -cell graft that provides some function but without clinical benefit relative to the indication for treatment should be considered a failure. On the other hand, a marginal  $\beta$ -cell graft associated with clear evidence of improvement in hypoglycemia or glycemic variability/lability even in the absence of achieving target glycemic control may be clinically important, but such an outcome would not be considered a success in terms of function since the overall treatment goals were not accomplished. This is an important distinction from a functionally failed  $\beta$ -cell graft where in the absence of any evidence for clinically significant C-peptide production consideration should be made to abandoning further monitoring and support of the failed graft. In particular, continuation of immunosuppression may no longer be indicated unless to support another allograft (e.g. a transplanted kidney) or to prevent possible sensitization to HLA antigens expressed by the  $\beta$ -cell graft in the case that another transplant is being considered. Thus, we

sought to define the functional status and clinical success of a  $\beta$ -cell graft separately, but using the same components of assessment: the HbA<sub>1c</sub>, severe hypoglycemic events, insulin requirements, and C-peptide. We did not define a duration required for correction of HbA<sub>1c</sub>, protection from hypoglycemia, restoration of C-peptide, or insulin-independence as these measures are fluid, and should each be evaluated together at any time of post-transplant graft functional assessment. Any reported change in glycemic control noted by SMBG and/or CGM should prompt such an assessment in order to identify a functionally stressed or declining graft.

We propose that functional and clinical outcomes can be assigned using a 4 tiered system as outlined in Table 3. Optimal  $\beta$ -cell graft function is defined by the presence of near-normal glycemic control assessed by non-diabetic HbA<sub>1c</sub>  $\leq 6.5\%$  (48 mmol/mol), the absence of any severe hypoglycemia, the absence of any requirement for exogenous insulin or other antihyperglycemic therapy, and documentation of an increase over pre-transplant measurement of C-peptide. Good  $\beta$ -cell graft function is defined by the presence of on-target glycemic control assessed by an HbA<sub>1c</sub>  $< 7.0\%$  (53 mmol/mol), the absence of any severe hypoglycemia, a reduction by more than 50% from baseline in insulin requirements or the use of non-insulin antihyperglycemic agents, and documentation of an increase over pre-transplant measurement of C-peptide. Both optimal and good functional outcomes are considered successful clinical outcomes. Marginal  $\beta$ -cell graft function is defined by the failure to achieve an HbA<sub>1c</sub>  $< 7.0\%$  (53 mmol/mol), the occurrence of any severe hypoglycemia, or less than 50% reduction in insulin requirements when there is documentation of an increase over pre-transplant measurement of C-peptide. If documented impairment in hypoglycemia awareness, frequent occurrence or exposure to SH, or marked glycemic variability/lability is convincingly improved, then it may be appropriate to consider that the  $\beta$ -cell graft is clinically impactful. In the absence of any evidence for a clinical impact, reassessment of the C-peptide status is warranted as clinically insignificant levels, even if quantifiably higher than before transplant, should be considered  $\beta$ -cell graft failure. Neither a marginal  $\beta$ -cell graft nor a failed  $\beta$ -cell graft is considered a clinically successful. Finally, the ultimate success for a  $\beta$ -cell replacement therapy in any individual patient requires the functional clinical benefits to outweigh any potential harm from the transplantation procedure or adverse effects of any required immunosuppression.

In conclusion, to be deemed successful  $\beta$ -cell replacement therapies should require the HbA<sub>1c</sub>  $< 7.0\%$  (53 mmol/mol) in the absence of severe hypoglycemia associated with a

Manuscript

significant >50% reduction in insulin requirements and restoration of clinically significant C-peptide production (>0.5 ng/ml or >0.17 nmol/l). Baseline assessment of hypoglycemia awareness, hypoglycemia severity, and glycemic variability/lability is helpful for monitoring whether a marginally functioning graft is continuing to provide any clinical impact. This proposed classification of function and clinical outcomes for  $\beta$ -cell replacement therapies is a work in progress and should be validated and further refined based on results from implementation in future prospective investigation.

## APPENDIX

Workshop participants:

Rodolfo Alejandro, University of Miami, Miami, FL, USA

Helmut Arbogast, University of Munich, Munich, Germany

Marcel Bassil, Benta Pharma Industries, Dbayeh, Lebanon

Melena Bellin, University of Minnesota, Minneapolis, MN, USA

Kanza Benomar, Centre Hospitalier Universitaire de Lille, Lille, France

Andrzej Berman, Warsaw Medical University, Warsaw, Poland

Thierry Berney, Geneva University Hospital, Geneva, Switzerland

Pratik Choudary, King's College London, London, United Kingdom

Jan de Boer, Eurotransplant International Foundation, Leiden, The Netherlands

Eelco De Koning, Leiden University, Leiden, The Netherlands

Jason Doppenberg, Leiden University, Leiden, The Netherlands  
Eva Dovolilova, Institute for Clinical & Experimental Medicine, Prague, Czech Republic  
Anneliese Flatt, Newcastle University, Newcastle upon Tyne, United Kingdom  
Justyna Gołębiewska, Medical University of Gdańsk, Gdańsk, Poland  
Aiste Gulla, Vilnius University, Vilnius, Lithuania  
Anna Högvall, Uppsala University, Uppsala, Sweden  
Paul Johnson, University of Oxford, Oxford, United Kingdom  
Andreas Kahl, Charité – Universitätsmedizin Berlin, Berlin, Germany  
Raja Kandaswamy, University of Minnesota, Minneapolis, MN, USA  
Thomas Kay, University of Melbourne, Melbourne, Victoria, Australia  
Bart Keymeulen, Vrije Universiteit Brussel, Brussels, Belgium  
Tomas Koblas, Institute for Clinical & Experimental Medicine, Prague, Czech Republic  
Lucie Kosinová, Institute for Clinical & Experimental Medicine, Prague, Czech Republic  
Jan Kriz, Institute for Clinical & Experimental Medicine, Prague, Czech Republic  
Yogish Kudva, Mayo Clinic, Rochester, MN, USA  
Robert Langer, Ordensklinikum Elisabethin Hospital, Linz, Austria  
Esther Latres, JDRF International New York, NY, USA  
Roger Lehmann, University Hospital Zurich, Zurich, Switzerland  
Ivan Leontovyc, Institute for Clinical & Experimental Medicine, Prague, Czech Republic  
Barbara Ludwig, University Hospital Carl Gustav Carus, Dresden, Germany  
Marjana Marinac, JDRF International New York, NY, USA  
James Markmann, Massachusetts General Hospital, Boston, MA, USA  
Marius Miglinas, Vilnius University, Vilnius, Lithuania  
Lenka Némětová, Institute for Clinical & Experimental Medicine, Prague, Czech Republic  
Michiel Nijhoff, Leiden University, Leiden, The Netherlands  
Jon Odorico, University of Wisconsin, Madison, WI, USA  
Ingalill Ort, Uppsala University, Uppsala, Sweden  
Alzbeta Patikova, Institute for Clinical & Experimental Medicine, Prague, Czech Republic  
Francois Pattou, Université de Lille, Lille, France  
Lorenzo Piemonti, San Raffaele Scientific Institute, Milan, Italy  
Johann Pratschke, Charité Medical School Berlin, Berlin, Germany



Michael Rickels, University of Pennsylvania, Philadelphia, PA, USA  
Charles Saab, Benta Pharma Industries, Dbayeh, Lebanon  
Frantisek Saudek, Institute for Clinical & Experimental Medicine, Prague, Czech Republic  
Tim Scholz, Oslo University Hospital, Oslo, Norway  
Hanne Scholz, Oslo University Hospital, Oslo, Norway  
Peter Senior, University of Alberta, Edmonton, Alberta, Canada  
James Shaw, Newcastle University, Newcastle upon Tyne, United Kingdom  
Ioannis Spiliotis, University of Oxford, Oxford, United Kingdom  
Peter Stock, University of California at San Francisco, San Francisco, CA, USA  
Gunnar Tufveson, Uppsala University, Uppsala, Sweden  
Aart Van Apeldoorn, Maastricht University, Maastricht, The Netherlands  
Marie-Christine Vantuyghem, Université de Lille, Lille, France  
Barbara Voglova, Institute for Clinical & Experimental Medicine, Prague, Czech Republic  
Steve White, Newcastle University, Newcastle upon Tyne, United Kingdom  
Michal Wszola, Warsaw Medical University, Warsaw, Poland  
Ming Han Yao, Karolinska Hospital, Stockholm, Sweden

## REFERENCES

1. Bartlett ST, Markmann JF, Johnson P, et al. Report from IPITA-TTS Opinion Leaders Meeting on the Future of Beta-Cell Replacement. *Transplantation*. 2016;100 Suppl 2:S1-44.
2. Markmann JF, Bartlett ST, Johnson P, et al. Executive Summary of the IPITA-TTS Opinion Leaders Meeting on the Future of Beta-Cell Replacement. *Transplantation*. 2016;100(7):e25-e31.
3. Agiostratidou G AH, Ball D, Blonde L, Gourgari E, Griffen S, Harriman KN, Kowalski A, Madden P, McAuliffe-Fogarty AH, McElwee-Malloy M, Peters A, Raman S, Reifschneider K, Rubin K, Weinzimer SA, for the T1D Outcomes Program Steering Committee. Standardizing clinically meaningful outcome measures beyond HbA1c for type 1 diabetes. *Diabetes Care*. 2017;40(12):1622-1630.
4. Maahs DM, Buckingham BA, Castle JR, et al. Outcome measures for artificial pancreas clinical trials: a consensus report. *Diabetes Care*. 2016;39(7):1175-1179.
5. International Hypoglycemia Study Group. Glucose concentrations of less than 3.0 mmol/L (54 mg/dL) should be reported in clinical trials: a joint position statement of the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetes Care*. 2017;40(1):155-157.
6. Hering BJ, Clarke WR, Bridges ND, et al. Phase 3 trial of transplantation of human islets in type 1 diabetes complicated by severe hypoglycemia. *Diabetes Care*. 2016;39(7):1230-1240.
7. Choudhary P, Rickels MR, Senior PA, et al. Evidence-informed clinical practice recommendations for treatment of type 1 diabetes complicated by problematic hypoglycemia. *Diabetes Care*. 2015;38(6):1016-1029.
8. Vantyghem MC, Kerr-Conte J, Arnalsteen L, et al. Primary graft function, metabolic control, and graft survival after islet transplantation. *Diabetes Care*. 2009;32(8):1473-1478.
9. Sivasubramaniyam S, Amiel SA, Choudhary P. Proportion of daily capillary blood glucose readings required in the target range for target glycaemic control: shift of focus from target range to proportion in range. *Diabet Med*. 2017;34(10):1456-1460.

10. Froud T, Faradji RN, Gorn L, et al. Dapsone-induced artifactual A1c reduction in islet transplant recipients. *Transplantation*. 2007;83(6):824-825.
11. Beck RW, Connor CG, Mullen DM, Wesley DM, Bergenstal RM. The fallacy of average: how using HbA1c alone to assess glycemic control can be misleading. *Diabetes Care*. 2017;40(8):994-999.
12. Whitelaw BC, Choudhary P, Hopkins D. Evaluating rate of change as an index of glycemic variability, using continuous glucose monitoring data. *Diabetes Technol Ther*. 2011;13(6):631-636.
13. Ryan EA, Shandro T, Green K, et al. Assessment of the severity of hypoglycemia and glycemic lability in type 1 diabetic subjects undergoing islet transplantation. *Diabetes*. 2004;53(4):955-962.
14. Senior PA, Bellin MD, Alejandro R, et al. Consistency of quantitative scores of hypoglycemia severity and glycemic lability and comparison with continuous glucose monitoring system measures in long-standing type 1 diabetes. *Diabetes Technol Ther*. 2015;17(4):235-242.
15. Seaquist ER, Anderson J, Childs B, et al. Hypoglycemia and diabetes: a report of a workgroup of the American Diabetes Association and The Endocrine Society. *J Clin Endocrinol Metab*. 2013;98(5):1845-1859.
16. Clarke WL, Cox DJ, Gonder-Frederick LA, Julian D, Schlundt D, Polonsky W. Reduced awareness of hypoglycemia in adults with IDDM. A prospective study of hypoglycemic frequency and associated symptoms. *Diabetes Care*. 1995;18(4):517-522.
17. Gold AE, Macleod KM, Frier BM. Frequency of severe hypoglycemia in patients with type-1 diabetes with impaired awareness of hypoglycemia. *Diabetes Care*. 1994;17(7):697-703.
18. Radosevich DM, Jevne R, Bellin M, Kandaswamy R, Sutherland DE, Hering BJ. Comprehensive health assessment and five-yr follow-up of allogeneic islet transplant recipients. *Clin Transplant*. 2013;27(6):E715-724.
19. Barnard KD, Hood KK, Weissberg-Benchell J, Aldred C, Oliver N, Laffel L. Psychosocial assessment of artificial pancreas (AP): commentary and review of existing measures and their applicability in AP research. *Diabetes Technol Ther*. 2015;17(4):295-300.
20. Speight J, Woodcock AJ, Reaney MD, et al. Well, I wouldn't be any worse off, would I, than I am now? A qualitative study of decision-making, hopes, and realities of adults with type 1 diabetes undergoing islet cell transplantation. *Transplant Direct*. 2016;2(5):e72.
21. Rickels MR, Kong SM, Fuller C, et al. Improvement in insulin sensitivity after human islet transplantation for type 1 diabetes. *J Clin Endocrinol Metab* 2013;98(11):E1780-E1785.

22. Gillard P, Hilbrands R, Van de Velde U, et al. Minimal functional beta-cell mass in intraportal implants that reduces glycemic variability in type 1 diabetic recipients. *Diabetes Care*. 2013;36(11):3483-3488.
23. Ryan EA, Lakey JR, Paty BW, et al. Successful islet transplantation: continued insulin reserve provides long-term glycemic control. *Diabetes*. 2002;51(7):2148-2157.
24. Faradji RN, Monroy K, Messinger S, et al. Simple measures to monitor beta-cell mass and assess islet graft dysfunction. *Am J Transplant*. 2007;7(2):303-308.
25. Ryan EA, Lakey JRT, Paty BW, Bigam D, Senior PA, Shapiro AMJ. beta-Score - An assessment of beta-cell function after islet transplantation. *Diabetes Care*. 2005;28(2):343-347.
26. Forbes S, Oram RA, Smith A, et al. Validation of the BETA-2 score: an improved tool to estimate beta cell function after clinical islet transplantation using a single fasting blood sample. *Am J Transplant*. 2016;16(9):2704-2713.
27. Vantghem MC, Raverdy V, Balavoine AS, et al. Continuous glucose monitoring after islet transplantation in type 1 diabetes: an excellent graft function (beta-score greater than 7) is required to abrogate hyperglycemia, whereas a minimal function is necessary to suppress severe hypoglycemia (beta-score greater than 3). *J Clin Endocrinol Metab*. 2012;97(11):E2078-E2083.
28. Rickels MR, Liu C, Shlansky-Goldberg RD, et al. Improvement in beta-cell secretory capacity after human islet transplantation according to the CIT07 protocol. *Diabetes*. 2013;62(8):2890-2897.
29. Robertson RP, Bogachus LD, Oseid E, et al. Assessment of beta-cell mass and alpha- and beta-cell survival and function by arginine stimulation in human autologous islet recipients. *Diabetes*. 2015;64(2):565-572.

**TABLE 1.**  
**Workshop session objectives**

<b>Indications for and Approaches to <math>\beta</math>-cell Replacement Therapy</b>
Patient candidates and available forms of $\beta$ -cell replacement vary. The goal of this session was to define those patient characteristics that directly influence the type of $\beta$ -cell graft and measures of glycemic control and graft function, and so lay the framework for how definitions of successful outcomes may be tailored by indication.
<b>Outcome Measures of Glucose Homeostasis</b>
Regulation of glucose homeostasis involves the maintenance and return of glucose excursions to a non-diabetic range of glycemia. Various measures of glycemic control capture average glycemia, glycemic variability, and exposure to hyper- and hypoglycemia, as well as hypoglycemia awareness and severity. The goal of this session was to define successful outcomes for glycemic control, and align definitions with those used in the field of artificial insulin delivery/artificial pancreas development.
<b>Outcome Measures of <math>\beta</math>-cell Graft Function and Demand</b>
Measures of $\beta$ -cell graft function may vary by the stimulus for secretion, differences in metabolic clearance, demands for secretion imposed by differences in insulin sensitivity or the use of insulin, as well as any possible residual native $\beta$ -cell function. The goal of this session was to define a meaningful reduction in insulin requirements attributable to $\beta$ -cell graft function, necessary confirmatory testing, relationship to standardized measures of glucose tolerance, and differences between type 1 and type 2 diabetic recipients.
<b>Outcome Measures of Immunologic Mechanisms</b>
Distinguishing immunologic from metabolic mechanisms for $\beta$ -cell graft dysfunction and/or failure is paramount to understanding the mechanisms underlying current graft status and implications of functional $\beta$ -cell graft monitoring. The goal of this session was to define useful assays of allo- and autoimmune reactivity and when they should be employed to complement the metabolic evaluation of $\beta$ -cell replacement therapies.
<b>Defining Successful Outcomes</b>
Clear definitions for success or failure of available $\beta$ -cell replacement therapies require incorporation

of both metrics for glycemic control and  $\beta$ -cell graft function. The goal of this session was to establish a practical consensus definition for  $\beta$ -cell graft functional and efficacy outcomes for  $\beta$ -cell replacement therapies.

**TABLE 2.**  
**Indications and goals for beta cell replacement therapies expressed in relation to various glycemic control measures**

Metric	Indication <sup>a</sup>	Goal	Ideal
HbA1c, % (mmol/mol) <sup>b</sup>	>7.5-8.0 (58-64)	<7.0 (53)	≤6.5 (48)
SH, events per yr	One or more	None	None
Clarke or Gold score <sup>c</sup>	≥4	<4	0
Time <54 mg/dl (3.0 mmol/l), % <sup>d</sup>	≥5	<1	0

Glucose SD, mg/dl (mmol/l) <sup>e</sup>	≥40 (2.2)	<40 (2.2)	NE
Glucose CV, % <sup>f</sup>	≥30	<30	NE
Time <70 mg/dl (3.9 mmol/l), % <sup>g</sup>	NE	<5	<5
Time 70 – 180 mg/dl (3.9 – 10 mmol/l), % <sup>h</sup>	NE	>70	>90
Time >180 mg/dl (10 mmol/l), % <sup>i</sup>	NE	<20-30	<5

HbA<sub>1c</sub>, glycated hemoglobin; SH, severe hypoglycemia; SD, standard deviation; CV, coefficient of variation = mean/SD; NE, not established.

<sup>a</sup>Typically more than one measure is used to define indications for  $\beta$ -cell replacement therapy and establish a baseline prior to treatment.

<sup>b</sup>Mean glucose should be used to provide an estimate of the HbA<sub>1c</sub> in the setting of marked anemia or administration of dapsone.<sup>10</sup>

<sup>c</sup>Used to assess impaired awareness of hypoglycemia.<sup>14,15</sup>

<sup>d</sup>Used to assess exposure to serious, clinically important hypoglycemia,<sup>4</sup> which can also be defined by frequency of episodes or using the HYPO score.<sup>11</sup>

<sup>e</sup>Used to assess glycemic variability,<sup>10</sup> which can also be assessed as glycemic lability using the lability index or LI.<sup>11</sup>

<sup>f</sup>Used for comparison to artificial pancreas systems.<sup>3</sup>

**TABLE 3.****Igls definition of functional and clinical outcomes for  $\beta$ -cell replacement therapy**

<b><math>\beta</math>-cell graft functional status</b>	<b>HbA<sub>1c</sub>, % (mmol/mol)<sup>a</sup></b>	<b>Severe hypoglycemia, events per yr</b>	<b>Insulin requirements, U·kg<sup>-1</sup>·d<sup>-1</sup></b>	<b>C-peptide</b>	<b>Treatment success</b>
Optimal	≤6.5 (48)	None	None	>Baseline <sup>b</sup>	Yes
Good	<7.0 (53)	None	<50% baseline <sup>c</sup>	>Baseline <sup>b</sup>	Yes
Marginal	Baseline	<Baseline <sup>d</sup>	≥50% baseline	>Baseline <sup>b</sup>	No <sup>e</sup>
Failure	Baseline	Baseline <sup>f</sup>	Baseline	Baseline <sup>g</sup>	No

HbA<sub>1c</sub>, glycated hemoglobin. Baseline, pre-transplant assessment.

<sup>a</sup>Mean glucose should be used to provide an estimate of the HbA<sub>1c</sub> in the setting of marked anemia or administration of dapsone.<sup>10</sup>

<sup>b</sup>Should also be >0.5 ng/ml (>0.17 nmol/l) fasting or stimulated.

<sup>c</sup>Should also be <0.5 U·kg<sup>-1</sup>·d<sup>-1</sup>; might include the use of non-insulin antihyperglycemic agents.

<sup>d</sup>Should severe hypoglycemia occur following treatment, then continued benefit may require assessment of hypoglycemia awareness, exposure to serious hypoglycemia (<54 mg/dl [3.0 mmol/l]), and/or glycemic variability/lability with demonstration of improvement from baseline.

<sup>e</sup>Clinically, benefits of maintaining and monitoring  $\beta$ -cell graft function may outweigh risks of maintaining immunosuppression.

<sup>f</sup>If severe hypoglycemia was not present before  $\beta$ -cell replacement therapy, then a return to baseline measures of glycemic control used as the indication for treatment (Table 2) may be consistent with  $\beta$ -cell graft failure.

<sup>g</sup>May not be reliable in uremic patients and/or in those patients with evidence of C-peptide production prior to  $\beta$ -cell replacement therapy.





Minerva Access is the Institutional Repository of The University of Melbourne

**Author/s:**

Rickels, MR; Stock, PG; de Koning, EJP; Piemonti, L; Pratschke, J; Alejandro, R; Bellin, MD; Berney, T; Choudhary, P; Johnson, PR; Kandaswamy, R; Kay, TWH; Keymeulen, B; Kudva, YC; Latres, E; Langer, RM; Lehmann, R; Ludwig, B; Markmann, JF; Marinac, M; Odorico, JS; Pattou, F; Senior, PA; Shaw, JAM; Vantghem, M-C; White, S

**Title:**

Defining outcomes for  $\beta$ -cell replacement therapy in the treatment of diabetes: a consensus report on the Igls criteria from the IPITA/EPITA opinion leaders workshop

**Date:**

2018-04-01

**Citation:**

Rickels, M. R., Stock, P. G., de Koning, E. J. P., Piemonti, L., Pratschke, J., Alejandro, R., Bellin, M. D., Berney, T., Choudhary, P., Johnson, P. R., Kandaswamy, R., Kay, T. W. H., Keymeulen, B., Kudva, Y. C., Latres, E., Langer, R. M., Lehmann, R., Ludwig, B., Markmann, J. F. ,... White, S. (2018). Defining outcomes for  $\beta$ -cell replacement therapy in the treatment of diabetes: a consensus report on the Igls criteria from the IPITA/EPITA opinion leaders workshop. *TRANSPLANT INTERNATIONAL*, 31 (4), pp.343-352.  
<https://doi.org/10.1111/tri.13138>.

**Persistent Link:**

<http://hdl.handle.net/11343/283647>

**File Description:**

Accepted version