

Research Article

No Additive Effects of Polyphenol Supplementation and Exercise Training on White Adiposity Determinants of High-Fat Diet-Induced Obese Insulin-Resistant Rats

Karen Lambert ¹, Marie Hokayem,¹ Claire Thomas,^{1,2} Odile Fabre,¹ Cécile Cassan,¹ Annick Bourret,¹ Florence Bernex,^{3,4} Jessica Lees,¹ Marie Demion,¹ Pascal Seyer,¹ Gérald Hugon,¹ Jacques Mercier,^{1,5} Antoine Avignon,^{1,5} and Catherine Bisbal¹

¹*PhyMedExp, CNRS UMR 9214, INSERM U1046, University of Montpellier, Montpellier, France*

²*Département STAPS, University of Evry, University of Paris Saclay, Evry, France*

³*IRCM, Inserm, University of Montpellier, ICM, Montpellier, France*

⁴*RHEM, BioCampus Montpellier, CNRS, INSERM, University of Montpellier, Montpellier, France*

⁵*Centre Hospitalier Universitaire (CHU), Montpellier, France*

Correspondence should be addressed to Karen Lambert; karen.lambert@univ-montp1.fr

Received 24 November 2017; Revised 26 January 2018; Accepted 15 February 2018; Published 15 April 2018

Academic Editor: Paola Venditti

Copyright © 2018 Karen Lambert et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

One of the major insulin resistance instigators is excessive adiposity and visceral fat depots. Individually, exercise training and polyphenol intake are known to exert health benefits as improving insulin sensitivity. However, their combined curative effects on established obesity and insulin resistance need further investigation particularly on white adipose tissue alterations. Therefore, we compared the effects on different white adipose tissue depot alterations of a combination of exercise and grape polyphenol supplementation in obese insulin-resistant rats fed a high-fat diet to the effects of a high-fat diet alone or a nutritional supplementation of grape polyphenols (50 mg/kg/day) or exercise training (1 hr/day to 5 days/wk consisting of treadmill running at 32 m/min for a 10% slope), for a total duration of 8 weeks. Separately, polyphenol supplementation and exercise decreased the quantity of all adipose tissue depots and mesenteric inflammation. Exercise reduced adipocytes' size in all fat stores. Interestingly, combining exercise to polyphenol intake presents no more cumulative benefit on adipose tissue alterations than exercise alone. Insulin sensitivity was improved at systemic, epididymal, and inguinal adipose tissues levels in trained rats thus indicating that despite their effects on adipocyte morphological/metabolic changes, polyphenols at nutritional doses remain less effective than exercise in fighting insulin resistance.

1. Introduction

An obesogenic lifestyle of an excessive caloric intake versus a low energy expenditure via physical inactivity favors obesity by increasing fat mass. This detrimental state of obesity/extreme fatness is at the center of the pathophysiology of type 2 diabetes (T2D) by causing insulin resistance (IR) to appear several decades before T2D. T2D prevalence is increased with aging (prevalence of 12–25% among older adults (>65 years)) and is characterized by IR, metabolic alterations, and chronic inflammation [1]. The initiating mechanisms of IR are partially known with major inducers being oxidative stress and

chronic inflammation [2–5]. Increased levels of reactive oxygen species and inflammatory cytokines have been found to negatively impact insulin signaling [1]. Interestingly, adipose tissue contributes to IR via adipocyte morphology dysfunction, adipocytokine secretion, or metabolic alterations. It has been shown that IR is linked to specific adipose tissue locations in the body [6]. In fact, the increase of metabolically active visceral adiposity is associated with IR development [7] while subcutaneous adiposity is associated with a lower risk of T2D development [8]. Consistent with the notion that subcutaneous fat is metabolically beneficial, its transplantation into the visceral cavity has been shown to improve

insulin sensibility [9]. Exercise training, a potent insulin sensitizer, is associated with a greater loss of visceral versus subcutaneous fat [10, 11]. Only recently, investigations aiming at understanding the effects of exercise training on nonmuscle tissues and more specifically adipose tissue depots have been designed [10–13]. Exercise training has been shown to reduce adiposity, adipocyte size [10], oxidative stress [14], and inflammation levels [15] while inducing an increase in mitochondrial enzyme activities in white adipose tissue [16]. Thus, the positive health effects of exercise training in preventing metabolic chronic diseases could be due to its metabolic regulation of fat stores and not only to its undeniable impact on skeletal muscle metabolism.

As mentioned above, the health-promoting effects of physical activity are well documented, leading the scientific community to discover and even create “exercise-mimetic” molecules at the center of which are the phytochemical mega family of polyphenols. Polyphenols due to their antioxidant/anti-inflammatory properties [17] have been used to fight against IR with contradictory effects. Some studies report beneficial effects [18–22] while others do not [23–26]. In sedentary rodents, resveratrol, a polyphenol found in colored berries, induces metabolic adaptations normally seen in trained animals [19]. However, due to its very low concentration in fruits and vegetables, this polyphenol is not a good representative of dietary polyphenol intake [27]. Moreover, the fact that polyphenol-rich diets positively improve glucose homeostasis [28] and obesity parameters in an elderly population [29] highlights how polyphenols’ health benefits cannot be attributed to a single compound but rather to a complex synergistic molecular effect. Accordingly to this data, we previously demonstrated the metabolic advantages of a mixture of grape polyphenol supplementation in nutritional amounts against fructose-induced IR and oxidative stress in healthy volunteers at high metabolic risk [30]. Interestingly, the combination of training and polyphenol supplementation on adipose tissue characteristics has been poorly investigated in obese IR rats [31]. The aim of our study was therefore to better understand the potential curative effects of a nutritional supplementation of grape polyphenols alone and in combination with exercise on the different adipose tissue depot characteristics in high-fat diet obese and IR rats in comparison with exercise training and grape polyphenol supplementation alone.

2. Materials and Methods

2.1. Ethical Approval. All animal experimentation procedures were in accordance with the Directive 2010/63/EU that was adopted on September 22, 2010, for the protection of animals used for scientific purposes (agreement number: A34–172–38) and were approved by the local research ethics committee (protocol number: CEEA-LR-1062).

2.2. Animals. After acclimation to the facility, a total of forty 6-week- (wk-) old male Sprague-Dawley rats (Janvier Laboratories) were fed a high-fat ad libitum diet (HFD, D12330 Research Diets, containing 25.5% carbohydrate, 58% fat, and 16.4% protein) for the 12-week duration of

the experimental protocol. The rats were single or double housed in a temperature-controlled room and maintained with food and drink ad libitum in a 12:12 h light-dark cycle, lights on at 8:00 postmeridian. The training and *in vivo* tests were performed during the rats’ dark cycle exposure. Rats’ body mass was monitored every two days throughout the experimental period.

2.3. Experimental Design. After 4 wks of HFD, previously shown to induce skeletal muscle IR [32], the animals were divided into 4 groups ($n = 10/\text{group}$) while simultaneously continuing the HFD for an additional 8-week period (Figure 1). Group specificities were the following: HFD alone (HF), HFD, and supplementation with grape polyphenol extract (PP) at 50 mg/kg/d per 50 ml of drinking water (PP), HFD, and exercise training 1 hr/day to 5 days/wk consisting of running on a treadmill set at 32 m/min for a 10% slope (EXO), HFD, and PP supplementation combined with exercise training (EXOPP). The grape polyphenol extract supplemented in this study (64.4% total polyphenol with 17.9% procyanidins, 4.3% anthocyanins, and 533 ppm resveratrol) had been previously successfully used in humans and in mice [30, 33]. The polyphenols’ mixture intake was 33.0 ± 2.1 ml/d for PP and 31.9 ± 1.7 ml/d for EXOPP groups compared to a water intake of 26.5 ± 1.3 ml/d for HF and EXO groups, indicating that the polyphenol mixture was well accepted by the animals.

Glucose tolerance tests were performed on animals at the start of the protocol and then after 4 and 7 wks of HFD (Figure 1). During the first 4 wks, all animals were handled daily and familiarized 5 days a week/5–10 min/day with the treadmill set at 10 m/min at 10% slope. This exercise duration is insufficient to significantly increase aerobic performance [34, 35]. Graded exercise tests were performed after 4wks and then at 11 wks (Figure 1). The rats selected for the exercise groups (EXO and EXOPP) reached the required training intensity in 1 wk. Animals were euthanized 48 hrs after the last training session to avoid the acute effect of the last exercise training session. Euthanasia was done by exsanguination after isoflurane anesthesia on 12 hr-fasted animals. Immediately after death, epididymal, perirenal, inguinal, anterior subcutaneous, and mesenteric adipose tissues were rapidly excised as previously detailed [36], weighed, and directly frozen in liquid nitrogen. Total fat mass (g) was estimated by summing the weights of the different fat stores excised. Adiposity (%) was calculated as the sum of the different adipose tissue depots (g) divided by rat’s body weight and multiplied by 100. Soleus, tibialis anterior, and extensor digitorum longus muscles were also rapidly surgically removed and weighted to determine lean mass.

2.4. Exercise Tests

2.4.1. Graded Exercise Test (GET). This test consisted of a progressive exercise test in which each rat initially ran at a speed of 10 m/min, up a 5% slope, for 3 min. Thereafter, the speed was increased by 3 m/min every 2 min until the rat was unable/unwilling to keep pace with the treadmill belt




FIGURE 1: Experimental design. After 4 weeks of HFD, 40 male Sprague-Dawley rats were divided into 4 groups ($n = 10/\text{group}$) while simultaneously continuing for an 8-week period the HFD alone (HF), HFD, and supplementation with grape polyphenol extract (PP) at 50 mg/kg/d per 50 ml of drinking water (PP), HFD, and exercise training 1 hr per day/5 days/wk consisting of running on a treadmill set at 32 m/min for a 10% slope (EXO), HFD, and PP supplementation combined with exercise training (EXOPP). Glucose tolerance tests (GTT) were performed at T0, 4, and 11 weeks. A graded exercise test was completed between the 4th and the 5th week and between the 11th and the 12th week where a time-to-fatigue test was also done.

despite encouragement to do so by application of manual solicitation at the hind legs.

2.5. Blood Analysis

2.5.1. Glucose Tolerance Test (GTT). After a 12 hr fast, rats were injected intraperitoneally with a 30% glucose solution (2 g/kg body weight), as previously described [37]. Glucose levels were determined using QuantiChrom™ Glucose Assay Kit (BioAssay Systems) on blood collected from the tail, at baseline and following glucose injection. Insulin levels were determined at baseline using Ultrasensitive Insulin ELISA (Mercodia). Homeostasis model assessment of IR (HOMA-IR) index was calculated using the formula: $\text{HOMA-IR} = [\text{fasting insulin } (\mu\text{IU/ml}) \times \text{fasting glucose (mmol/ml)}] / 22.5$. Kg represented the exponential decrease in glycaemia during glucose tolerance test and thus glucose disposal [38].

2.5.2. Determination of Free Fatty Acid, Total Cholesterol, Low- and Very Low-Density Lipoproteins, High-Density Lipoproteins, and Leptin Plasma Levels. Plasma samples from 12 hr-fasted rats were collected under anesthesia from the carotid artery and centrifuged for 10 min at 2000 \times g. Free fatty acid (FFA), total cholesterol, low- and very low-density lipoproteins (LDL/VLDL), and high-density lipoproteins (HDL) levels were determined using EnzyChrom™ Free Fatty Acid Assay Kit (EFFA, BioAssay System) and EnzyChrom HDL and LDL/VLDL Assay Kit (EHDL, BioAssay System). Leptin concentration was quantified by ELISA (BioVendor).

2.6. Adipose Tissue Analysis. Adipose tissue insulin response was assessed by measuring P-Akt/Akt protein ratio in epididymal, inguinal, peritoneal, and mesenteric adipose portions incubated or not with insulin, as previously described [39, 40]. Two 50–100 mg pieces of adipose tissue slices were washed in phosphate buffer saline (PBS). Then, one entire explant was incubated at 37°C for 15 min with PBS only or supplemented with 1 μM human insulin (Umuline RAPIDE)

[30] before being snap frozen in liquid nitrogen. The adipose tissue portions were disrupted and homogenized in a hypotonic lysis buffer [41] with protease and phosphatase inhibitors (Sigma-Aldrich). Protein concentration was measured in the extracts using the Pierce BCA Protein Assay Kit (ThermoFisher Scientific). Protein expression was assessed on protein extracts by Western blot, using antibodies diluted in Odyssey Blocking Buffer (LI-COR Biosciences, Bad Homburg, Germany). Insulin response was determined using anti-Phospho-Akt (P-Akt) (Ser473) and anti-Akt antibodies at a 1:1000 dilution (both from Cell Signaling Technology Inc). Similarly, phospho-AMPK (P-AMPK, Thr172), AMPK, I κ B α , and SOCS3 proteins expression were analyzed in adipose tissues using anti P-AMPK, AMPK at a 1:500 and 1:1000 dilution, anti-I κ B α primary antibody at a 1:500 dilution, anti-SOCS3 primary antibody at a 1:1000 dilution, and α/β -tubulin antibody at a 1:1000 dilution (all five antibodies used were from Cell Signaling Technology Inc.). Following primary antibody incubation, nitrocellulose membranes were incubated with secondary antibodies that were directed against the species of the primary antibody while also being conjugated to IRDye800 or IrDye700, at a 1:30,000 dilution (LI-COR Biosciences). Specific bands were visualized with a LI-COR Odyssey CLx Imaging System (LI-COR Biosciences). Protein expression levels were quantified using ImageJ software. Finally, levels of protein expression in each sample were corrected using the respective quantified levels of the corresponding loading control (Akt for P-Akt, AMPK for P-AMPK, or α/β -tubulin for I κ B α and SOCS3) analyzed on the same membrane.

2.7. Adipocyte Size. Relative adipocyte size was estimated at 20-fold magnified hematoxylin and eosin-stained sections by counting the number of adipocytes in 5 random fields of each sample with two sections by sample. Relative cell size was expressed as adipocyte count per section. An increase in adipocyte count per section is related to a decrease in adipocyte size.




FIGURE 2: Glucose tolerance test scores. (a) Glycaemia expressed in percentage of basal glycaemia and (b) area under the curve (AUC) during glucose tolerance test of 16 male Sprague-Dawley rats before (baseline, white squares) and after 4 weeks of an ad libitum high-fat diet (black squares). Data are expressed as mean \pm SEM. Significant difference: * $p < 0.05$ versus baseline.

TABLE 1: Metabolic changes after polyphenol supplementation and exercise training.

	HF	PP	EXO	EXOPP	Two-way ANOVA
Weight (g)	603 \pm 19	570 \pm 15	539 \pm 15 *	542 \pm 15 *	Exercise $p < 0.01$
Treadmill speed (m/min)	28 \pm 1.0	28 \pm 1.3	41.5 \pm 1.5	42.25 \pm 1.1	Exercise $p < 0.01$
Leptin/adiposity	0.26 \pm 0.02	0.19 \pm 0.03	0.23 \pm 0.03	0.21 \pm 0.03	NS
HOMA-IR	2.3 \pm 0.1	2.2 \pm 0.2 ^s	2.1 \pm 0.2 ^s	1.7 \pm 0.1*	Exercise $p < 0.01$
FFA (μ M)	346.6 \pm 77.1	218.9 \pm 27.0	147.0 \pm 31.6	174.7 \pm 36.2	Exercise $p < 0.05$
HDL (mg/dl)	28.8 \pm 1.8	31.1 \pm 2.3	30.5 \pm 2.3	36.7 \pm 1.6*	Exercise $p = 0.074$ Supplementation $p < 0.05$
Cholesterol (mg/dl)	62.8 \pm 3.6	63.1 \pm 3.3	57.7 \pm 7.2	69.7 \pm 4.3	$p > 0.05$
LDL/VLDL (mg/dl)	28.7 \pm 2.7	24.5 \pm 2.8	29.2 \pm 3.0	26.1 \pm 2.7	$p > 0.05$
Citrate synthase activity (μ mol/min/g tissue)	13.7 \pm 0.7	16.9 \pm 1.2	19.9 \pm 1.9	16.5 \pm 0.4	Exercise $p < 0.05$

* $p < 0.05$ versus HF; ^s $p < 0.05$ versus EXOPP.

2.8. *Citrate Synthase (CS) Activity.* Enzymatic activity of citrate synthase was determined by spectrometry, as previously described on soleus homogenates [42].

2.9. *Statistical Analysis.* Data are expressed as means \pm standard error of the mean (SEM). Comparisons between treatment groups were performed using two-way analysis of variance (ANOVA) with a Bonferroni post hoc test of pairwise comparisons between groups or using three-way analysis of variance (ANOVA) with a Bonferroni post hoc test of pairwise comparisons between groups or Student's unpaired, two-tailed t -test, when required. A probability value of <0.05 was considered significant.

3. Results

3.1. *Animal Characteristics.* After one month of the high-fat diet (HF), rats presented an increase in both body weight 250.6 g \pm 1.7 to 422.2 g \pm 4.8 ($p < 0.001$) and glycaemia during glucose tolerance test measurements compared to baseline ($p < 0.05$ for all points) (Figures 2(a) and 2(b)) thus

demonstrating an impairment in glucose disposal associated with obesity. During the eight following weeks, the animals were subjected to high-fat diet alone (HF) or HF with a supplementation of grape polyphenol extract at nutritional doses (PP) or to exercise training (EXO) or to the combination of exercise and polyphenol supplementation (EXOPP). As expected, we observed an increase in treadmill speed during the graded exercise test (GET) and an increase in muscle citrate synthase activity in trained rats compared to sedentary rats (Table 1). Free fatty acid (FFA) were decreased only in trained rats (EXO and EXOPP versus HF and PP, $p < 0.05$), and high-density lipoproteins (HDL) were increased after polyphenol supplementation (PP and EXOPP versus HF and EXO, $p < 0.05$) (Table 1). Total cholesterol, low-density lipoproteins (LDL), and very low-density lipoproteins (VLDL) levels remained unaltered in all experimental groups tested (Table 1).

3.2. *Metabolic Characterization.* Whole-body glucose tolerance was determined with a glucose tolerance test (GTT) (Figure 3(a)). Glucose disposal during GTT was increased




FIGURE 3: Measure of whole-body glucose homeostasis and adipose tissue insulin sensitivity. (a) Glycaemia expressed in percentage of fasted blood glucose and (b) glucose disposal (Kg) calculated from GTT. Data are expressed as mean \pm SEM with $n = 8 - 10$ rats per group. P-Akt and Akt expression levels were measured by immunoblotting after insulin stimulation in epididymal (c), inguinal (d), mesenteric (e), and perirenal (f) fat stores. A representative blot is shown for each adipose tissue depot. Quantified values are presented as P-Akt/Akt fold induction relative to HF group's values, which were set at 1. Data are expressed as mean \pm SEM with $n = 5$ rats per group. Bars not sharing a common letter are significantly different at $p < 0.05$.

in trained rats (EXO and EXOPP versus HF and PP) as attested by increased of Kg glycaemia constant (Figure 3(b), $p < 0.05$), indicating that whole-body glucose tolerance was improved by exercise training (EXO and EXOPP versus HF and PP). Moreover, EXOPP rats presented the lowest HOMA-IR compared to HF, PP, and EXO rats, demonstrating a higher insulin sensitivity (Table 1). To further our understanding, we investigated the insulin response of each adipose tissue depots obtained (Figures 3(c), 3(d), 3(e), and 3(f)) by measuring P-Akt/Akt ratio before and after incubation with insulin. P-Akt/Akt ratio was increased only in epididymal and inguinal fat stores of trained rats (EXO and EXOPP versus HF and PP). When analyzed together, all adipose tissues collected presented significant increases

in P-Akt/Akt ratios following exercise training. Alone, polyphenol supplementation did not increase P-Akt/Akt ratio. We also investigated P-AMPK/AMPK activation in the epididymal, inguinal, mesenteric, and perirenal adipose tissue depots (Figures 4(a), 4(b), 4(c), and 4(d)) in order to evaluate GLUT4 translocation [43]. We found no notable change in P-AMPK/AMPK ratio in any of the experimental conditions (exercise and/or polyphenol supplementation) or any of the adipose tissue depots.

3.3. Adipose Tissue Depot Characteristic Modification. After eight weeks of high-fat diet associated with exercise training (EXO) alone or combined with polyphenol supplementation (EXOPP), trained rats had a decrease in body weight




FIGURE 4: Effect of exercise training and polyphenol supplementation on P-AMPK/AMPK levels in epididymal, inguinal, mesenteric, and perirenal fat stores. P-AMPK and AMPK expression levels were measured by immunoblotting in epididymal (a), inguinal (b), mesenteric (c), and perirenal (d) fat stores. A representative blot is shown for each adipose tissue depot. Values are measured as P-AMPK/AMPK ratios relative to HF group's values, which were set at 1. Data are expressed as mean \pm SEM with $n = 4$ rats per group.

compared to sedentary rats (HF and PP), without any beneficial effect of polyphenol supplementation on this parameter (Table 1). This decrease is mainly due to a reduced fat mass (Table 2) that is significantly correlated to body weight ($r = 0.726$, $p < 0.0001$). Total fat mass decrease was observed in exercise-trained groups (EXO and EXOPP, $p < 0.001$) (Table 2) with a tendency to decrease also observed in the PP group ($p = 0.059$, Table 2). Although not significantly different from EXO rats, EXOPP rats exhibited the smallest fat mass and adiposity values compared to all other animal groups (Table 2). Exercise training decreased the quantity of all fat pads: inguinal, epididymal, perirenal, mesenteric, and anterior subcutaneous (Table 2). In addition, as shown in Figures 5(a), 5(b), 5(c), and 5(d), exercise training significantly increased adipocyte number per section ($p < 0.05$) in all adipose tissue depots which indicates an overall decrease in adipocyte size. When combined together, the three-way ANOVA indicated that polyphenol supplementation

lowered the mass of all types of white adipose tissue studied (Table 2, $p = 0.006$). An interaction was also found between all adipose tissue depots and exercise training ($p = 0.003$). Nevertheless, the amplitude of the decrease was not the same for all fat stores with inguinal and mesenteric adipose tissues presenting the lowest decrease due to exercise training, as opposed to anterior subcutaneous, perirenal, and epididymal adipose tissues. We investigated visceral fat pad (total sum of epididymal, perirenal, and mesenteric adipose tissues) weight as it has previously been shown that during high-fat feeding, exercise training preferentially affects this type of fat store [10]. Here, due to exercise training, we observed a decrease in visceral adipose tissue ($p < 0.001$) and subcutaneous adipose tissue stores ($p < 0.033$) (Table 2). Combination of polyphenol supplementation with exercise training did not further amplify this reduction compared to exercise alone. Interestingly, we found an important decrease in the ratio of fat mass on lean (muscle) mass with training. The effect

TABLE 2: Impact of polyphenol supplementation and exercise on fat stores.

	HF	PP	EXO	EXOPP	Two-way ANOVA	Three-way ANOVA
Fat mass (g)	63.5 ± 5.5	49.8 ± 4.9	38.0 ± 4.3	35.1 ± 3.2	Exercise $p < 0.001$ Suppl. effect $P = 0.059$	NA
Adiposity (%)	10.1 ± 0.7	8.7 ± 0.7 [§]	6.4 ± 0.5 *	6.2 ± 0.5 *	Exercise $p < 0.001$	NA
Leptin (ng/ml)	2.5 ± 0.4	1.6 ± 0.2*	1.4 ± 0.1*	1.2 ± 0.1*	Exercise and suppl. effect $p < 0.05$	NA
Inguinal fat (g)	9.2 ± 1.5	4.9 ± 1.17.507	3.9 ± 1.8	3.6 ± 1.7	Exercise $p = 0.056$	
Epididymal fat (g)	15.3 ± 1.0	13.3 ± 1.2 [§]	9.0 ± 1.2*	8.4 ± 1.2*	Exercise $p < 0.001$	Suppl. effect $p = 0.006$ interaction fat stores and exercise $p = 0.003$
Perirenal fat (g)	20.7 ± 1.6	16.6 ± 1.8 [§]	11.4 ± 1.9*	10.3 ± 1.8*	Exercise $p < 0.001$	
Mesenteric fat (g)	17.2 ± 1.1	14.2 ± 1.2	13.1 ± 1.3*	12.4 ± 1.2*	Exercise $p < 0.05$	
Anterior subcutaneous fat (g)	1.2 ± 0.1	0.9 ± 0.1 [§]	0.6 ± 0.1*	0.5 ± 0.1*	Exercise $p = 0.001$	
Visceral fat (g)	53.2 ± 4.0	44.0 ± 4.4 [§]	33.5 ± 3.7 *	31.1 ± 2.9 *	Exercise $p < 0.001$	NA
Subcutaneous fat (g)	10.3 ± 1.6	5.8 ± 1.7	4.5 ± 1.8 * $p = 0.091$	4.0 ± 1.7*	Exercise $p = 0.033$	NA
Fat mass/lean mass*100	4.5 ± 0.4	3.4 ± 0.3 * [§]	2.3 ± 0.2*	2.2 ± 0.2*	Exercise $p < 0.001$ Suppl. effect $p = 0.056$	NA

* $p < 0.05$ versus HF, [§] $p < 0.05$ versus EXOPP. Suppl. effect: polyphenol supplementation effect. Exercise: exercise training effect. NA: not applicable

of polyphenol supplementation was close to significance ($p = 0.056$), which implies that polyphenol supplementation, as exercise, may induce a decrease in fat mass associated with an increase in muscle mass.

3.4. Leptin Levels, $\text{I}\kappa\text{B}$, and SOCS3 Expression According to White Adipose Tissue Fat Depot Type. Leptin is a major adipokine related to the amount of energy stored as fat in white adipose tissue and involved in the pathogenesis of chronic inflammation [44]. Interestingly, we observed a decrease in leptin levels after training (Table 2); however, leptin/adiposity ratio was constant between groups (Table 2), indicating that the decrease in leptin levels must be related to adiposity decline. Confirming this result, we found strong correlations between leptin levels and both subcutaneous ($R = 0.668$, $p < 0.00001$) and visceral ($R = 0.628$, $p < 0.0001$) fat pad mass. Seeing that leptin is at the interface between metabolism and inflammation [44], we investigated $\text{I}\kappa\text{B}$ protein levels in the different adipose tissue depots as $\text{I}\kappa\text{B}$ is able to sequester the transcription factor $\text{NF}\kappa\text{B}$ in the cytosol and thus counteracting its proinflammatory action [45] (Figures 6(a)–6(d)). Individually, only mesenteric fat stores presented an increase in $\text{I}\kappa\text{B}$ protein levels due to both exercise ($p = 0.010$) and polyphenol supplementation ($p = 0.033$) (Figure 6(c)). However, when analyzed together, the three-way ANOVA showed a significant effect of both exercise ($p < 0.001$) and polyphenol supplementation ($p = 0.005$). Thus, both interventions were able to improve adipose tissue inflammation due to high-fat feeding. Again, there was no additive effect of exercise and polyphenol on this parameter. Since suppressor of cytokine signaling 3 (SOCS3) is a protein involved in a negative feedback loop of cytokine activity [46] but also in insulin signaling regulation [47], we explored its expression in the different adipose

tissue depots (Figures 7(a)–7(d)) and found that SOCS3 protein content only significantly decreased in inguinal fat depots ($p = 0.040$) (Figure 7(b)) of trained rats. Polyphenol supplementation did not modify SOCS3 protein content in any adipose tissue depot studied.

4. Discussion

In this study, we investigated, in high-fat IR obese rats, the potential curative effects of exercise training combined or not with a daily nutritional supplementation of a grape polyphenol mixture on different adipose tissue depots. The main result of our study is that compared to exercise alone, combining exercise training to a supplementation of polyphenols has no cumulative beneficial effect on adipose tissue alterations of IR obese rats whatever their location.

In our model of obese IR rats, exercise training decreases the total amount of white adipose tissue with a more significant decrease in visceral ($p < 0.001$) than subcutaneous ($p < 0.05$) depots. Interestingly, polyphenol supplementation also induces a decrease in the total amount of adipose tissue depots. However, even though there is an improvement in insulin sensitivity at both systemic and epididymal/inguinal adipose tissue levels with exercise training, such improvement is not observed with polyphenol supplementation alone. Thus, the reduction in adipose tissue mass observed with our nutritional polyphenol supplementation is not associated with an improvement of insulin sensitivity at both systemic and tissular levels as observed in previous study reporting a decrease in body weight without improvement of insulin sensitivity [48]. However, in our study, such surprising result could be due to an insufficient decrease in fat pads with polyphenol supplementation (-22% with PP versus -40% with EXO). Another hypothesis




FIGURE 5: Effect of exercise training and polyphenol supplementation on adipose tissue. Representative sections of epididymal (a), inguinal (b), mesenteric (c), and perirenal (d) adipose tissues stained with hematoxylin and eosin are shown. Relative cell size was expressed as adipocyte count per section. An increase in adipocyte count per section is related to a decrease in adipocyte size. Data are expressed as mean \pm SEM with $n = 8 - 10$ rats per group. Bars not sharing a common letter are significantly different at $p < 0.05$.

is that the beneficial effect of exercise training on adipose tissue is not only related to its lowering effect on adipose tissue mass and adipocyte size but may involve other mechanisms. Interestingly, we extend in obese insulin-resistant rat the previous data from Stanford et al. [13] showing that exercise training was able to remodel inguinal adipose tissue in normal weight rats. Moreover, we did not find significant variation in AMPK activation in any adipose tissue location; even

if there is a tendency of P-AMPK/AMPK ratio to increase after training.

Adipose tissue was previously thought to be a mere storage depot for lipids and energy; several studies have now shown that adipocytes secrete biologically active molecules, that is, adipokines as leptin with complex autocrine, paracrine, and endocrine functions [49]. A dysregulation in such adipokine levels and leptin resistance has been implicated in




FIGURE 6: IκB levels in fat stores. IκB protein expression levels were measured by immunoblotting in epididymal (a), inguinal (b), mesenteric (c), and perirenal (d) fat stores. Quantification values are presented as IκB protein levels normalized to tubulin protein level, used as protein loading control, and are expressed relative to HF group's values, which were set at 1. Data are expressed as mean \pm SEM with $n = 5 - 6$ rats per group. Bars not sharing a common letter are significantly different at $p < 0.05$.

the pathophysiology of IR and T2D [50]. However, we have recently shown that such dysregulation was not found in women with grade I obesity since IR and insulin sensitive women presented the same adipokine levels [39]. As previously shown in several human studies [51–53], our results point out that leptin is not involved in the beneficial effect of exercise as there is no difference in leptin levels between groups when adjusted to adipose tissue mass. However, the decreased adipocytes' size observed with exercise training could participate in the improvement of insulin sensitivity. In accordance with this data, we did not observe an improvement in insulin sensitivity with polyphenol supplementation or a decrease in adipocyte size. In fact, large adipocytes have been associated with ROS production leading to IR and inflammatory reactions [54, 55]. Exercise training and polyphenol supplementation improve only mesenteric adipose tissue inflammation whereas there is no improvement in insulin sensitivity in this fat pad depot. The three-way

ANOVA demonstrates that the other adipose tissue depots also present metabolic improvement with exercise only when analyzed together. SOCS3 diminishes insulin-stimulated IRS1 phosphorylation leading to a decrease in insulin response [56–58]. Interestingly, we observed only in inguinal fat depots a decrease in SOCS3 protein content in trained rat which is associated with an increase in insulin sensitivity in this fat depot after training. Thus, mesenteric fat stores are more sensitive to our polyphenol supplementation while inguinal and epididymal adipose tissue depots are more sensitive to training, demonstrating a depot-specific response as previously found by others [15, 59].

Our polyphenol supplementation presents some similar beneficial effects of exercise training as a reduction of adipose tissue depots and of mesenteric inflammation, which has been both described as positive regulators of insulin sensitivity. More interestingly, exercise training has been shown in our study to improve systemic and adipose tissue insulin




FIGURE 7: SOCS3 levels in fat stores. SOCS3 protein expression levels were measured by immunoblotting in epididymal (a), inguinal (b), mesenteric (c), and perirenal (d) fat stores. Quantification values are presented as SOCS3 protein levels normalized to tubulin protein levels, used as protein loading control, and are expressed relative to HF group's values, which were set at 1. Data are expressed as mean \pm SEM with $n = 4 - 5$ rats per group. Bars not sharing a common letter are significantly different at $p < 0.05$.

sensitivity while the combination of exercise and polyphenol supplementation (EXOPP) results in an improvement in systemic insulin sensitivity attested by a significant decrease in HOMA-IR without further altering any adipose tissue markers tested in this study.

Taken together, our data seem to indicate that the beneficial effects of exercise training are not only due to alteration in adipose tissue. Recent data in mice [60] and obese patients [61] also confirm that adipose tissue alterations in the development of insulin resistance and T2D play a less important role than once thought. We previously found in obese women that skeletal muscle seems to be the primary tissular target of insulin signaling defects in obesity development [39]. Exercise training and polyphenol supplementation individually or in combination have been found to increase insulin sensitivity at whole-body but also at skeletal muscle levels [30, 62, 63]. Thus, the improvement of systemic insulin sensitivity that we observed after training could also be due to its effect on skeletal muscle.

In conclusion, our study indicates that in high-fat diet-induced obese and IR rats, the combination of polyphenol

supplementation with exercise training presents no more cumulative benefit on white adipose tissue alterations than exercise training or polyphenol supplementation alone.

Conflicts of Interest

The authors declare that there is no conflict of interest regarding the publication of this article.

Acknowledgments

The authors thank M. Blaqui re (PhyMedExp, INSERM U1046) for the technical assistance. Financial support for this study was obtained from INSERM.

References

- [1] G. Verdile, K. N. Keane, V. F. Cruzat et al., "Inflammation and oxidative stress: the molecular connectivity between insulin resistance, obesity, and Alzheimer's disease," *Mediators of Inflammation*, vol. 2015, Article ID 105828, 17 pages, 2015.

- [2] A. Avignon, M. Hokayem, C. Bisbal, and K. Lambert, "Dietary antioxidants: do they have a role to play in the ongoing fight against abnormal glucose metabolism?," *Nutrition*, vol. 28, no. 7-8, pp. 715-721, 2012.
- [3] C. Bisbal, K. Lambert, and A. Avignon, "Antioxidants and glucose metabolism disorders," *Current Opinion in Clinical Nutrition and Metabolic Care*, vol. 13, no. 4, pp. 439-446, 2010.
- [4] O. Fabre, C. Breuker, C. Amouzou et al., "Defects in TLR3 expression and RNase L activation lead to decreased MnSOD expression and insulin resistance in muscle cells of obese people," *Cell Death & Disease*, vol. 5, no. 3, article e1136, 2017.
- [5] G. Hotamisligil, N. Shargill, and B. Spiegelman, "Adipose expression of tumor necrosis factor- α : direct role in obesity-linked insulin resistance," *Science*, vol. 259, no. 5091, pp. 87-91, 1993.
- [6] J. Vague, "Sexual differentiations and distribution of fat," *La Semaine des Hôpitaux*, vol. 26, no. 49, pp. 2387-2390, 1950.
- [7] C. Zhang, K. M. Rexrode, R. M. van Dam, T. Y. Li, and F. B. Hu, "Abdominal obesity and the risk of all-cause, cardiovascular, and cancer mortality: sixteen years of follow-up in US women," *Circulation*, vol. 117, no. 13, pp. 1658-1667, 2008.
- [8] V. J. Carey, E. E. Walters, G. A. Colditz et al., "Body fat distribution and risk of non-insulin-dependent diabetes mellitus in women. The nurses' health study," *American Journal of Epidemiology*, vol. 145, no. 7, pp. 614-619, 1997.
- [9] T. T. Tran, Y. Yamamoto, S. Gesta, and C. R. Kahn, "Beneficial effects of subcutaneous fat transplantation on metabolism," *Cell Metabolism*, vol. 7, no. 5, pp. 410-420, 2008.
- [10] K. S. C. Gollisch, J. Brandauer, N. Jessen et al., "Effects of exercise training on subcutaneous and visceral adipose tissue in normal- and high-fat diet-fed rats," *American Journal of Physiology. Endocrinology and Metabolism*, vol. 297, no. 2, pp. E495-E504, 2009.
- [11] D. Vissers, W. Hens, J. Taeymans, J. P. Baeyens, J. Poortmans, and L. Van Gaal, "The effect of exercise on visceral adipose tissue in overweight adults: a systematic review and meta-analysis," *PLoS One*, vol. 8, no. 2, article e56415, 2013.
- [12] K. I. Stanford and L. J. Goodyear, "Exercise regulation of adipose tissue," *Adipocytes*, vol. 5, no. 2, pp. 153-162, 2016.
- [13] K. I. Stanford, R. J. W. Middelbeek, K. L. Townsend et al., "A novel role for subcutaneous adipose tissue in exercise-induced improvements in glucose homeostasis," *Diabetes*, vol. 64, no. 6, pp. 2002-2014, 2015.
- [14] T. Sakurai, T. Izawa, T. Kizaki et al., "Exercise training decreases expression of inflammation-related adipokines through reduction of oxidative stress in rat white adipose tissue," *Biochemical and Biophysical Research Communications*, vol. 379, no. 2, pp. 605-609, 2009.
- [15] F. S. Lira, J. C. Rosa, A. S. Yamashita, C. H. Koyama, M. L. Batista Jr, and M. Seelaender, "Endurance training induces depot-specific changes in IL-10/TNF- α ratio in rat adipose tissue," *Cytokine*, vol. 45, no. 2, pp. 80-85, 2009.
- [16] B. Stallknecht, J. Vinten, T. Ploug, and H. Galbo, "Increased activities of mitochondrial enzymes in white adipose tissue in trained rats," *The American Journal of Physiology*, vol. 261, no. 3, pp. E410-E414, 1991.
- [17] Y. Zhao, B. Chen, J. Shen et al., "The beneficial effects of quercetin, curcumin, and resveratrol in obesity," *Oxidative Medicine and Cellular Longevity*, vol. 2017, Article ID 1459497, 8 pages, 2017.
- [18] P. Brasnyo, G. A. Molnar, M. Mohas et al., "Resveratrol improves insulin sensitivity, reduces oxidative stress and activates the Akt pathway in type 2 diabetic patients," *The British Journal of Nutrition*, vol. 106, no. 03, pp. 383-389, 2011.
- [19] M. Lagouge, C. Argmann, Z. Gerhart-Hines et al., "Resveratrol improves mitochondrial function and protects against metabolic disease by activating SIRT1 and PGC-1 α ," *Cell*, vol. 127, no. 6, pp. 1109-1122, 2006.
- [20] I. Momken, L. Stevens, A. Bergouignan et al., "Resveratrol prevents the wasting disorders of mechanical unloading by acting as a physical exercise mimetic in the rat," *The FASEB Journal*, vol. 25, no. 10, pp. 3646-3660, 2011.
- [21] J. Most, G. H. Goossens, J. W. E. Jocken, and E. E. Blaak, "Short-term supplementation with a specific combination of dietary polyphenols increases energy expenditure and alters substrate metabolism in overweight subjects," *International Journal of Obesity*, vol. 38, no. 5, pp. 698-706, 2014.
- [22] S. Timmers, E. Konings, L. Bilet et al., "Calorie restriction-like effects of 30 days of resveratrol supplementation on energy metabolism and metabolic profile in obese humans," *Cell Metabolism*, vol. 14, no. 5, pp. 612-622, 2011.
- [23] L. Gliemann, J. F. Schmidt, J. Olesen et al., "Resveratrol blunts the positive effects of exercise training on cardiovascular health in aged men," *The Journal of Physiology*, vol. 591, no. 20, pp. 5047-5059, 2013.
- [24] N. Hart, L. Sarga, Z. Csenge et al., "Resveratrol attenuates exercise-induced adaptive responses in rats selectively bred for low running performance," *Dose Response*, vol. 12, no. 1, pp. 57-71, 2014.
- [25] J. Olesen, L. Gliemann, R. Bienso, J. Schmidt, Y. Hellsten, and H. Pilegaard, "Exercise training, but not resveratrol, improves metabolic and inflammatory status in skeletal muscle of aged men," *The Journal of Physiology*, vol. 592, no. 8, pp. 1873-1886, 2014.
- [26] T. D. Scribbans, J. K. Ma, B. A. Edgett et al., "Resveratrol supplementation does not augment performance adaptations or fibre-type-specific responses to high-intensity interval training in humans," *Applied Physiology, Nutrition, and Metabolism*, vol. 39, no. 11, pp. 1305-1313, 2014.
- [27] C. Manach, A. Scalbert, C. Morand, C. Remesy, and L. Jimenez, "Polyphenols: food sources and bioavailability," *The American Journal of Clinical Nutrition*, vol. 79, no. 5, pp. 727-747, 2004.
- [28] K. Hanhineva, R. Torronen, I. Bondia-Pons et al., "Impact of dietary polyphenols on carbohydrate metabolism," *International Journal of Molecular Sciences*, vol. 11, no. 12, pp. 1365-1402, 2010.
- [29] X. Guo, A. Tresserra-Rimbau, R. Estruch et al., "Polyphenol levels are inversely correlated with body weight and obesity in an elderly population after 5 years of follow up (the randomised PREDIMED study)," *Nutrients*, vol. 9, no. 12, 2017.
- [30] M. Hokayem, E. Blond, H. Vidal et al., "Grape polyphenols prevent fructose-induced oxidative stress and insulin resistance in first-degree relatives of type 2 diabetic patients," *Diabetes Care*, vol. 36, no. 6, pp. 1454-1461, 2013.
- [31] T. Sakurai, J. Ogasawara, T. Kizaki et al., "The effects of exercise training on obesity-induced dysregulated expression of adipokines in white adipose tissue," *International Journal of Endocrinology*, vol. 2013, Article ID 801743, 28 pages, 2013.

- [32] B. B. Yaspelkis III, S. J. Lessard, D. W. Reeder et al., "Exercise reverses high-fat diet-induced impairments on compartmentalization and activation of components of the insulin-signaling cascade in skeletal muscle," *American Journal of Physiology. Endocrinology and Metabolism*, vol. 293, no. 4, pp. E941–E949, 2007.
- [33] K. Lambert, M. Coisy-Quivy, C. Bisbal et al., "Grape polyphenols supplementation reduces muscle atrophy in a mouse model of chronic inflammation," *Nutrition*, vol. 31, no. 10, pp. 1275–1283, 2015.
- [34] K. M. Baldwin, D. A. Cooke, and W. G. Cheadle, "Time course adaptations in cardiac and skeletal muscle to different running programs," *Journal of Applied Physiology*, vol. 42, no. 2, pp. 267–272, 1977.
- [35] G. A. Dudley, W. M. Abraham, and R. L. Terjung, "Influence of exercise intensity and duration on biochemical adaptations in skeletal muscle," *Journal of Applied Physiology*, vol. 53, no. 4, pp. 844–850, 1982.
- [36] J. M. A. de Jong, O. Larsson, B. Cannon, and J. Nedergaard, "A stringent validation of mouse adipose tissue identity markers," *American Journal of Physiology. Endocrinology and Metabolism*, vol. 308, no. 12, pp. E1085–E1105, 2015.
- [37] M. Elferchichi, J. Mercier, A. Bourret et al., "Is static magnetic field exposure a new model of metabolic alteration? Comparison with Zucker rats," *International Journal of Radiation Biology*, vol. 87, no. 5, pp. 483–490, 2011.
- [38] R. N. Bergman, "Toward physiological understanding of glucose tolerance. Minimal-model approach," *Diabetes*, vol. 38, no. 12, pp. 1512–1527, 1989.
- [39] C. Amouzou, C. Breuker, O. Fabre et al., "Skeletal muscle insulin resistance and absence of inflammation characterize insulin-resistant grade I obese women," *PLoS One*, vol. 11, no. 4, article e0154119, 2016.
- [40] J. R. Zierath, D. Galuska, L. A. Nolte, A. Thörne, J. S. Kristensen, and H. Wallberg-Henriksson, "Effects of glycaemia on glucose transport in isolated skeletal muscle from patients with NIDDM: in vitro reversal of muscular insulin resistance," *Diabetologia*, vol. 37, no. 3, pp. 270–277, 1994.
- [41] T. Salehzada, L. Cambier, N. V. Thi, L. Manchon, L. Regnier, and C. Bisbal, "Endoribonuclease L (RNase L) regulates the myogenic and adipogenic potential of myogenic cells," *PLoS One*, vol. 4, no. 10, article e7563, 2009.
- [42] P. Srere, *Citrate Synthase*, 1969.
- [43] M. Daval, F. Foufelle, and P. Ferre, "Functions of AMP-activated protein kinase in adipose tissue," *The Journal of Physiology*, vol. 574, no. 1, pp. 55–62, 2006.
- [44] N. Iikuni, Q. K. Lam, L. Lu, G. Matarese, and A. Cava, "Leptin and inflammation," *Current Immunology Reviews*, vol. 4, no. 2, pp. 70–79, 2008.
- [45] S. T. Whiteside and A. Israel, "I κ B proteins: structure, function and regulation," *Seminars in Cancer Biology*, vol. 8, no. 2, pp. 75–82, 1997.
- [46] R. Starr, T. A. Willson, E. M. Viney et al., "A family of cytokine-inducible inhibitors of signalling," *Nature*, vol. 387, no. 6636, pp. 917–921, 1997.
- [47] B. Emanuelli, P. Peraldi, C. Filloux et al., "SOCS-3 inhibits insulin signaling and is up-regulated in response to tumor necrosis factor- α in the adipose tissue of obese mice," *The Journal of Biological Chemistry*, vol. 276, no. 51, pp. 47944–47949, 2001.
- [48] S. Talukdar, Y. Zhou, D. Li et al., "A long-acting FGF21 molecule, PF-05231023, decreases body weight and improves lipid profile in non-human primates and type 2 diabetic subjects," *Cell Metabolism*, vol. 23, no. 3, pp. 427–440, 2016.
- [49] T. Sakurai, J. Ogasawara, K. Shirato et al., "Exercise training attenuates the dysregulated expression of adipokines and oxidative stress in white adipose tissue," *Oxidative Medicine and Cellular Longevity*, vol. 2017, Article ID 9410954, 12 pages, 2017.
- [50] H. Tilg and A. R. Moschen, "Adipocytokines: mediators linking adipose tissue, inflammation and immunity," *Nature Reviews. Immunology*, vol. 6, no. 10, pp. 772–783, 2006.
- [51] W. M. Kohrt, M. Landt, and S. J. Birge Jr., "Serum leptin levels are reduced in response to exercise training, but not hormone replacement therapy, in older women," *The Journal of Clinical Endocrinology & Metabolism*, vol. 81, no. 11, pp. 3980–3985, 1996.
- [52] L. Perusse, G. Collier, J. Gagnon et al., "Acute and chronic effects of exercise on leptin levels in humans," *Journal of Applied Physiology*, vol. 83, no. 1, pp. 5–10, 1997.
- [53] F. S. L. Thong, R. Hudson, R. Ross, I. Janssen, and T. E. Graham, "Plasma leptin in moderately obese men: independent effects of weight loss and aerobic exercise," *American Journal of Physiology. Endocrinology and Metabolism*, vol. 279, no. 2, pp. E307–E313, 2000.
- [54] G. S. Hotamisligil and E. Erbay, "Nutrient sensing and inflammation in metabolic diseases," *Nature Reviews Immunology*, vol. 8, no. 12, pp. 923–934, 2008.
- [55] N. Houstis, E. D. Rosen, and E. S. Lander, "Reactive oxygen species have a causal role in multiple forms of insulin resistance," *Nature*, vol. 440, no. 7086, pp. 944–948, 2006.
- [56] H. Shi, I. Tzamelis, C. Bjorbaek, and J. S. Flier, "Suppressor of cytokine signaling 3 is a physiological regulator of adipocyte insulin signaling," *The Journal of Biological Chemistry*, vol. 279, no. 33, pp. 34733–34740, 2004.
- [57] H. Shi, B. Cave, K. Inouye, C. Bjorbaek, and J. S. Flier, "Overexpression of suppressor of cytokine signaling 3 in adipose tissue causes local but not systemic insulin resistance," *Diabetes*, vol. 55, no. 3, pp. 699–707, 2006.
- [58] R. Palanivel, M. D. Fullerton, S. Galic et al., "Reduced *Socs3* expression in adipose tissue protects female mice against obesity-induced insulin resistance," *Diabetologia*, vol. 55, no. 11, pp. 3083–3093, 2012.
- [59] T. Tchkonina, T. Thomou, Y. Zhu et al., "Mechanisms and metabolic implications of regional differences among fat depots," *Cell Metabolism*, vol. 17, no. 5, pp. 644–656, 2013.
- [60] W. T. Peppler, L. K. Townsend, C. M. Knuth, M. T. Foster, and D. C. Wright, "Subcutaneous inguinal white adipose tissue is responsive to, but dispensable for, the metabolic health benefits of exercise," *American Journal of Physiology. Endocrinology and Metabolism*, vol. 314, no. 1, pp. E66–E77, 2018.
- [61] N. Barbarroja, C. Lopez-Pedreria, L. Garrido-Sanchez et al., "Progression from high insulin resistance to type 2 diabetes does not entail additional visceral adipose tissue inflammation," *PLoS One*, vol. 7, no. 10, article e48155, 2012.
- [62] A. Shimotoyodome, S. Haramizu, M. Inaba, T. Murase, and I. Tokimitsu, "Exercise and green tea extract stimulate fat oxidation and prevent obesity in mice," *Medicine and Science in Sports and Exercise*, vol. 37, no. 11, pp. 1884–1892, 2005.
- [63] M. C. Venable, C. J. Hulston, H. R. Cox, and A. E. Jeukendrup, "Green tea extract ingestion, fat oxidation, and glucose tolerance in healthy humans," *The American Journal of Clinical Nutrition*, vol. 87, no. 3, pp. 778–784, 2008.



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Lambert, K; Hokayem, M; Thomas, C; Fabre, O; Cassan, C; Bourret, A; Bernex, F; Lees, J; Demion, M; Seyer, P; Hugon, G; Mercier, J; Avignon, A; Bisbal, C

Title:

No Additive Effects of Polyphenol Supplementation and Exercise Training on White Adiposity Determinants of High-Fat Diet-Induced Obese Insulin-Resistant Rats.

Date:

2018

Citation:

Lambert, K., Hokayem, M., Thomas, C., Fabre, O., Cassan, C., Bourret, A., Bernex, F., Lees, J., Demion, M., Seyer, P., Hugon, G., Mercier, J., Avignon, A. & Bisbal, C. (2018). No Additive Effects of Polyphenol Supplementation and Exercise Training on White Adiposity Determinants of High-Fat Diet-Induced Obese Insulin-Resistant Rats.. *Oxid Med Cell Longev*, 2018, pp.7406946-. <https://doi.org/10.1155/2018/7406946>.

Persistent Link:

<http://hdl.handle.net/11343/287568>

File Description:

Published version

License:

CC BY