



Journal of Contemporary Medicine
and Dentistry

www.jcmad.com

ISSN [P-2347-4513]
ISSN [O-2349-0799]
Year: 2020, Volume: 8
Issue: 2, p: 05-09


Attribution-NonCommercial 4.0
International (CC BY-NC 4.0)

A Community Based Study on the Role of Maternal Education on Antenatal Care Services and Child Care at Various Tribal Villages, Adilabad, Telangana State

A Ravinder¹, G Sreekanth²

1. Tutor cum Lecturer, Statistician, Department of Community Medicine, Rajiv Gandhi Institute of Medical Sciences, Adilabad Telangana State
2. Associate Professor, Department of Community Medicine, Rajiv Gandhi Institute of Medical Sciences, Adilabad, Telangana State

Abstract

Background: Educated women tend to have a greater awareness of the existence of ANC services, more aware of health problems, know more about the availability of health care services, and utilize the information more effectively than non-educated women. Moreover, higher levels of education tend to positively affect health-seeking behaviors, and education may increase a woman's control over her pregnancy and expose women to more health education messages and campaigns, enabling them to recognize danger signs and complications and take appropriate action. In this study, we tried to analyze to what level the maternal education can influence the antenatal care services and childcare. **Methods:** A community-based cross-sectional study was conducted on 93 randomly selected mothers who have children less than 7 years by using a pre-tested structured questionnaire for data collection at various tribal villages, at an average 25 kms away from Adilabad town from April-October 2019. Analysis was done using SPSS for windows version 16, Microsoft excel and Open epi website. **Results:** Out of 93 study subjects, 19.3% (18) of the study subjects were not registered for the antenatal services. Higher the maternal education higher the age at pregnancy ($p < 0.05$), was associated with more preference for deliveries by doctors ($p < 0.05$). More the frequency of exclusive breast feeding, long lasting breast feeding was associated more than a year ($p < 0.05$), a smaller number of children with wasting ($p < 0.05$). **Conclusion:** This study revealed that utilization of ANC services was relatively better for the mother's educated higher than secondary school, but they are still low. Educational status is important in having more health seeking behavior. In this study, it proves that health education is more important than the mere school education which can help to improve knowledge on ANC.

Keywords: Antenatal care, maternal education, health service utilization, tribal

Address for correspondence: Dr. G Srikanth, Associate Professor, Department of Community Medicine, Rajiv Gandhi Institute of Medical Sciences, Adilabad, Telangana state. Email id:

Date of Acceptance: 06/06/2020

Introduction

Antenatal care services were considered to be key element in the primary health care delivery system of a country which aims for a healthy society. Utilization of Antenatal Care (ANC) services and maternal and child health programs were critically important in a country like India which is experiencing high infant and child

mortality rate and maternal mortality rate. As per NFHS-4, Under-5 mortality rate declined in India from 74 in 2005-06 to 54 in 2015-16. Under-5 mortality rate in India was 54 in 2015-16 and in urban, rural areas, Infant mortality rate was 34, 56 in 2015-16 simultaneously^[1]. Over the past sixty years maternal health situation in the country has been staggering despite several changes in a rapidly evolving socio-economic

environment. In the last decade, as per the national data, health indicators including utilization of antenatal care services were as poor as 30% in tribal India [2]. Understanding the effect of infant and young child feeding (IYCF) practices on improving the nutritional status of children under two years of age, the World Health Organization (WHO) developed a set of core indicators to assess IYCF practices [3]. These indicators incorporated both breastfeeding and complementary feeding linked practices. Appropriate feeding practices, therefore, include timely initiation of feeding of solid and semi-solid foods from age 6 months and to improve the quantity and quality of foods children consume, while maintaining breastfeeding [4].

Previous studies conducted elsewhere on factors associated with appropriate complementary feeding practices of children aged 6–23 months show higher maternal and paternal education, better household wealth, exposure to media, adequate antenatal and post-natal contacts, child's sex and age, institutional delivery, low parity, maternal occupation, urban residence, knowledge & frequency of complementary feeding and receiving feeding advice in immunization as determinant factors for appropriate complementary feeding [5]. To improve complementary feeding practice through this essential time of growth and development of the child, assessment of complementary feeding practices and its factors are vital [2].

Among the various determinants of nutritional status, parent's education is probably the next most important factor after the socio-economic status. A literate mother uses scarce resources in 2 better manners for the child's welfare than an illiterate mother with higher resources does [6]. D'souza et al; believe that the effect of women's education on the nutritional status of their children is exerted through their roles as providers of household health and nutrition [7]. While the relationship between the mother's education and the child's nutritional status is well documented, that between the education of father or the education of both the parents put together and the child's nutritional status is not well established. Benerji et al; says that nutrition education of mothers of infants has a positive

effect on the nutritional status of their children [8].

Glewwe (1999) highlights three links through which education may affect child health [9]. Other studies have found a strong link between maternal education, social economic status and child nutritional status. This is because educated women are more likely to get steadier, higher paying jobs; to get married to men with higher education and higher income; and to live in better neighborhoods, which have influence on child health and survival [10].

Materials and Methods

A community-based cross-sectional study was conducted on mothers who have children less than 7 years by using a pre-tested structured questionnaire for data collection at Arjuni, Kolamguda, Lohara, Jamini, Mamidiguda, Nandigama, Kandala, Pothaguda, Dahiguda, Gunjala, Linguguda, Tekdiguda, Chinna maleboregav, Pendurluddi tribal villages, at an average 25kms away from Adilabad town from April-October 2019. The study was conducted on 93 randomly selected women on convenience, house to house visit and taken consent. Analysis was done using SPSS for windows version 16, Microsoft excel and Open Epi website.

Inclusion criteria:

Mothers who have children less than 7 years and Mothers who have undergone the recent pregnancy.

Exclusion criteria:

Mothers who did not give consent were excluded.

Results

The study subjects selected for the study were mothers educated more than secondary school were 12(12.9%), primary educated were 28 (30%) and illiterate were 53(56.3%).

The subjects selected for the study were statistically different in their age groups and education ie Higher the maternal education higher was the age at pregnancy ($p < 0.05$). 19.3 % (12) of the study subjects were not registered for the antenatal services. The maternal education and the duration of birth spacing were not statistically associated.

Figure- 1: Mothers according to education status




Table- 1: Role of maternal education

	Education			Total	p value
	Secondary [n=12]	Primary [n=28]	Illiterate [n=53]		
Age					
< 21yrs	5	5	5	15	0.05*
22-30 yrs	1	6	6	13	
31-35	6	17	42	65	
Registration of Pregnancy					
Yes	11	24	40	75	0.316
No	1	4	13	18	
Birth spacing					
Primi	5	9	18	32	0.447
12-24 m	3	15	21	39	
>24 m	4	4	14	22	

Table- 2: Role of maternal education on number of antenatal visits, person conducting delivery and place of delivery

	Education			Total	p value
	Secondary [n=12]	Primary [n=28]	Illiterate [n=53]		
Number of Antenatal Visits					
< 4	9	21	45	75	0.49
Above 4	3	7	8	18	
Person Conducted Delivery					
Dai & Other	3	18	18	39	0.03
Doctor	7	9	22	38	
ANM	2	1	13	16	
Place of Delivery					
Hospital	11	21	36	68	0.24
Home	1	7	17	25	

86% (45 of 53) of illiterate study subjects have less than 4 antenatal visits. Higher the education more the preference of doctor and the association is statistically significant ($p < 0.05$). 32% (17) of illiterate women were delivered at home.

The maternal education and the birth weight of the child were not associated significantly. 7.5% (7) of the study subjects feed their children with pre-lacteal feeds. Higher the maternal education more children were feed more than 8 times in a day, stopped exclusive breast feeding upto 6 months, feed their children with breast milk for

more than 1 year and the association is statistically significant.

Table 3: Role of maternal education on birth weight, number of breast feedings per day, exclusive breast feeding, time of stoppage of breast feeding.

	Education			Total	p value
	Secondary [n=12]	Primary [n=28]	Illiterate [n=53]		
Birth weight	2.75±0.62	2.77±0.42	2.65±0.44	93	0.499
Pre-lacteal feeds					
Yes	1	0	6	7	0.474
No	11	28	47	86	
Number of breast feeding per day					
<8	2	6	29	37	0.004*
8 to 12	6	16	21	43	
>12	4	6	3	13	
Exclusive breastfeeding					
less than 6 months	2	6	29	37	0.025*
above 6 months	6	16	21	43	
Total stoppage of breast feeding					
<1yr	2	6	45	53	<0.001*
>1yr	10	22	8	40	

Table- 4: Role of maternal education on baby hospitalization for sickness, proportion of wasting and stunted children

	Education			Total	p value
	Secondary [n=12]	Primary [n=28]	Illiterate [n=53]		
Baby hospitalization					
Yes	1	1	8	10	0.27
No	11	27	45	83	
Weight/age of child					
Wasting	8	27	39	74	0.03*
Normal	4	1	14	19	
Height/age of the child					
Stunting	7	19	40	66	0.45
Normal	5	9	13	27	
Total	12	28	53	93	

15% (8 of 45) of illiterate women had hospitalized their children during sickness episodes. The children of illiterate mothers were most wasted, and the association is statistically significant ($p = 0.026$). 75% [39 of 53] children of illiterate mothers were wasted and 74[79.5%] of study subjects' children were wasted. 75.5% (45 of 53) children of illiterate mother child were stunted and 66[71%] of study subjects' children were stunted.

Discussion

The subjects selected for the study were statistically different in their age groups and education. Therefore, higher the maternal education higher was the age at pregnancy ($p < 0.05$). 19.3% (12) of the study subjects were not registered for the antenatal services compared to Bhanwar Singh et al 28% of

females not received antenatal care ^[11]. The maternal education and the duration of birth spacing were not statistically associated. 86% (45 of 53) of illiterate study subjects have less than 4 antenatal visits compared to 40 % of illiterate subjects have less than 4 antenatal visits in the study by Hiyeswar Borah et al; ^[12]. Higher the education more the preference of doctor and the association is statistically significant ($p < 0.05$) is similar ($p < 0.001$) to the study of Rodgers O Moindi et al; ^[13] 32% (17) of illiterate women were delivered at home which is almost similar to the study of Rodgers O Moindi et al; ^[13] which is 25.5% (26 of 103). The maternal education and the birth weight of the child were not associated significantly. 23.54% of high educated mothers feed their children with pre-lacteal feeds in the study of Vishnu khalal et al; ^[14] compared it is 7.5 % (7) of the study subjects feed their children with pre-lacteal feeds. Higher the maternal education more children were feed more than 8 times in a day, stopped exclusive breast feeding up to 6 months, feed their children with breast milk for more than 1 year and the association is statistically significant which is similar to the study by Tigist Kassa et al; ^[15]. Sive et al; ^[6], while comparing 53 children hospitalized for kwashiorkor with 106 children hospitalized for non-nutritional diseases observed that the major difference between the two groups was the educational status of mother. Only 57% of the mothers of the children with kwashiorkor were literate as compared to 93% of the controls. In our study, 15% (8 of 45) of illiterate women had hospitalized their children during sickness episodes. 75% (39 of 53) children of illiterate mothers were wasted and the association is statistically significant ($p = 0.03$) and is similar to the study carried out at Parbhani, Marathwada (Maharashtra), Arya et al; ^[16] have shown that the children of literate mothers have better anthropometric measurements than children of illiterate mothers. 74 (79.5%) of study subjects' children were wasted compared to 28 % of children for total Telangana state according to NFHS-IV (17) and 22.1% for total Adilabad district according to NFHS-IV ^[18] which was comparatively very high. 75.5% (45 of 53) children of illiterate mother child were stunted compared to 33% of children in total Telangana

state according to NFHS –IV ^[17] were stunted and 38.3% for total Adilabad district according to NFHS-IV ^[18] which was comparatively very high.

Conclusion

This study revealed that utilization of ANC services was relatively better for the mother's educated higher than secondary school, but they are still low. This study proves that literate mothers perform much better if they are educated, motivated. Educational status is important in having more health seeking behavior. In this study, it proves that health education is more important than the mere school education which can help to improve knowledge on ANC. Providing IEC and household level discussion is important for ANC service utilization in the district.

Conflict of Interest: None declared

Source of Support: Nil

Ethical Permission: Obtained

References

1. <http://rchiips.org/NFHS/pdf/NFHS4/India.pdf> (Accessed dt: 16/03/2020)
2. Roy MP, Mohan U, Singh SK, Singh VK, Srivastava AK. Determinants of utilization of antenatal care services in rural Lucknow, India. *Journal of family medicine and primary care*. 2013 Jan;2(1):55
3. WHO, UNICEF, USAID, FANTA, AED, UC DAVIS, and IFPRI. Indicators for assessing infant and young child feeding practices part 2: measurement. Geneva: The World Health Organization; 2010. page no 2-12
4. Lack RE, Victora CG, Walker SP, Bhutta ZA, et al, for the Maternal and Child Under nutrition Group. Maternal and child under nutrition: maternal and child under nutrition and overweight in low-income and middle-income countries. *Lancet*. 2013; 382(9890): 427-451.
5. Senarath U, et al. Comparisons of complementary feeding indicators and associated factors in children aged 6–23 months across five South Asian countries. *Matern Child Nutr*. 2012; 8(Suppl1):89–106.

6. Sive AA, Subotzky EF, Malan H. The social, family and medical backgrounds of children with kwashiorkor presenting at a teaching hospital. *South Africa Medical Journal*. 1993; 83(3):180–183.
7. D'souza S, Bhuiya AL. Socio-economic mortality differentials in a rural area of Bangladesh, population and development review. 1982;8: 753–09.
8. B. Banerjee, O.N. Mandal An Intervention Study in Malnutrition Among Infants in a rural Community of West Bengal *Indian Journal of Community Medicine* 2005;30(1):1-3.
9. Glewwe P. Why Does Mother's Schooling Raise Child Health in Developing Countries? Evidence from Morocco. *J. Human Res.* 1999;34(1):124–159.
10. Lassi ZS, Das JK, Zahid G, Imdad A, Bhutta ZA. Impact of education and provision of complementary feeding on growth and morbidity in children less than 2 years of age in developing countries: a systematic review. *BMC Public Health*. 2013;13(Suppl 3):S13.
11. Bhanwar Singh, Meely Panda et al. Practices related to delivery and antenatal care among females in rural block of Haryana, *International Journal of Therapeutic Applications* 2013;10: 23-27.
12. Hiyeswar Borah, Parash Jyoti Bora Sch. *J. App. Med. Sci.*, Aug 2017; 5(8C):3179-3184.
13. Rodgers O Moindi, Moses M Ngari, et al. Why mothers still deliver at home: understanding factors associated with home deliveries and cultural practices in rural coastal Kenya, a cross-section study. *BMC Public Health* 2016;16:114.
14. Vishnu Khanal, Mandira Adhikari et al. Factors associated with introduction of prelacteal feeds in Nepal: Findings from the Nepal demographic and health survey 2011. *Int Breast feed J* 2013; 8: 9.
15. Tigist Kassa, Berhan Meshesha, Yusuf Haji, Jemal Ebrahim. Appropriate complementary feeding practices and associated factors among mother of children aged 6-23 months in Southern Ethiopia 2015. *BMC Pediatr* 2016;16:131.
16. Arya A, Devi R. Influence of maternal literacy on the nutritional status of pre-school children. *Indian Journal of Pediatrics*. 1991;58:265–268
17. http://rchiips.org/nfhs/pdf/NFHS4/TG_FactSheet.pdf [accessed on 16/03/2020]
18. http://rchiips.org/nfhs/FCTS/TG/TG_FactSheet_532_Adilabad.pdf [Accessed on 16/03/2020]