



^b
**UNIVERSITÄT
BERN**

Faculty of Business, Economics and
Social Sciences

Department of Social Sciences

University of Bern Social Sciences Working Paper No. 39

Entropy balancing as an estimation command

Ben Jann

Current version: August 16, 2021

First version: August 3, 2021

<http://ideas.repec.org/p/bss/wpaper/39.html>

<http://econpapers.repec.org/paper/bsswpaper/39.htm>

Entropy balancing as an estimation command

Ben Jann
Institute of Sociology
University of Bern
ben.jann@unibe.ch

Abstract. Entropy balancing is a popular reweighting technique that provides an alternative to approaches such as, for example, inverse probability weighting (IPW) based on a logit or probit model. Even if the balancing weights resulting from the procedure will be of primary interest in most applications, it is noteworthy that entropy balancing can be represented as a simple regression-like model. An advantage of treating entropy balancing as a parametric model is that it clarifies how the reweighting affects statistical inference. In this article I present a new Stata command called `ebalfit` that estimates such a model including the variance-covariance matrix of the estimated coefficients. The balancing weights are then obtained as model predictions. Variance estimation is based on influence functions, which can be stored for further use, for example, to obtain consistent standard errors for statistics computed from the reweighted data.

Keywords: `st0001`, Stata, `ebalfit`, entropy balancing, reweighting, inverse probability weighting, IPW, influence function

1 Introduction

The goal of entropy balancing, a procedure made popular by [Hainmueller \(2012\)](#), is to find a vector of weights that balances the data between two subsamples with respect to specific moments (e.g. the means and variances of a given set of covariates). For example, in order to estimate an “average treatment effect on the treated” (ATET) from observational data we might want to reweight a “control group” such that the means of observed pre-treatment variables match the means of these variables in the “treatment group”. Entropy balancing thus provides an alternative to other reweighting techniques commonly used in the treatment effects literature, such as inverse probability weighting (IPW) or matching (see, e.g., [Imbens and Wooldridge 2009](#) for an overview), some of which are implemented in Stata’s `teffects` command ([TE] `teffects`). An advantage of entropy balancing over classic IPW or matching is that it leads to perfect balance (if perfect balance is possible given the degree to which the common support assumption is violated); classic IPW and matching typically balance the data only approximately (unless the balancing problem is very simple). Perfect balance means that modeling the outcome (e.g. using regression adjustment) after the data have been balanced will lead to no refinements in the treatment effect estimate, implying that entropy balancing has the “doubly robust” property (also see [Zhao and Percival 2017](#)).

Entropy balancing can also be useful for other types of applications. For example, we may employ entropy balancing to construct weights for population surveys, say, by

adjusting the sample to a known population distribution or by fitting weights based on sampling frame data (see, e.g., Chapters 13 and 14 in [Valliant et al. 2013](#) for an overview of survey weighting). Related applications would be to use entropy balancing to compensate selective attrition in a randomized controlled trial (RCT) (assuming that selection is on observables) or to generalize experimental results from a selective sample of participants to the population (assuming that treatment effect heterogeneity is conditional on observables). Furthermore, entropy balancing may be used for more peculiar purposes, such as constructing weights that make the data orthogonal.

From a statistical point of view, entropy balancing boils down to a relatively simple regression-like parametric model. A first key contribution of this paper is to show how entropy balancing can be expressed as a system of moment equations and how, based on this representation, influence functions can be derived for the parameters of the model. These influence functions can then be used to obtain a consistent estimate of the variance matrix of the entropy balancing coefficients, but also to adjust the variance estimates of statistics computed from the reweighted data to take account of the uncertainty implied by the estimation of the weights. A second key contribution is to provide a new command called `ebalfit` that implements the described methods. Advantages of `ebalfit` over existing implementations¹ are that `ebalfit` behaves like official Stata’s estimation commands (similar syntax, output, and returns), that it provides standard errors and confidence intervals for the estimated coefficients, and that influence functions can be stored for further use in analyses employing the balancing weights.

This article is structured as follows. Drawing on preliminary work by [Jann \(2020b\)](#) and [Jann \(2020c\)](#), I first describe the entropy balancing model and its estimation, including the derivation of influence functions and an approach to adjust the standard errors of reweighted statistics. I then describe the syntax and options of the new command. Finally, I provide a set of examples illustrating the practical application of the new command.

2 The entropy balancing model

2.1 Two-sample balancing

In two-sample entropy balancing the goal is to reweight a sample of interest such that it has the same characteristics as some reference sample. Let $i = 1, \dots, N$ be the index of observations across both samples. Indicator variable S_i is equal to 1 for observations that belong to the primary sample, that is, to the sample that is to be reweighted, and 0 else. Furthermore, \mathcal{S} is the set of indices of observations that belong to this sample (i.e. observations for which $S_i = 1$). Likewise, indicator variable R_i is equal to 1 for observations that belong to the reference sample, and \mathcal{R} contains its indices. Note

1. See command `ebalance` by [Hainmueller and Xu \(2011, 2013\)](#). Note that entropy balancing can also be performed by command `psweight` by [Kranker \(2019\)](#), a command that implements “covariate-balancing propensity score” (CBPS) estimation as proposed by [Imai and Ratkovic \(2014\)](#). Entropy balancing is formally equivalent to just-identified CBPS, leading to the same coefficients and the same balancing weights.

that \mathcal{S} and \mathcal{R} do not need to be disjoint nor exhaustive (for example, the two samples may overlap). Each observation has a base weight w_i (e.g. a sampling weight based on the survey design) and a $k \times 1$ vector \mathbf{x}_i of data. Furthermore, $W = \sum_{i=1}^N w_i$ is the sum of weights across the joint sample; $W_S = \sum_{i=1}^N S_i w_i = \sum_{i \in \mathcal{S}} w_i$ and $W_R = \sum_{i=1}^N R_i w_i = \sum_{i \in \mathcal{R}} w_i$ are the sums of weights in the primary sample and the reference sample, respectively.

Given the target sum of weights $\hat{\tau} = W_R = \sum_{i \in \mathcal{R}} w_i$ (i.e. the size of the reference sample) and the $k \times 1$ vector of target moments $\hat{\mu} = \frac{1}{W_R} \sum_{i \in \mathcal{R}} w_i \mathbf{x}_i$ (i.e. the means of the data in the reference sample), entropy balancing looks for an estimate of $(\beta', \alpha)'$ such that

$$\frac{1}{\hat{\tau}} \sum_{i \in \mathcal{S}} \hat{w}_i \mathbf{x}_i = \hat{\mu} \quad \text{and} \quad \sum_{i \in \mathcal{S}} \hat{w}_i = \hat{\tau} \quad \text{with} \quad \hat{w}_i = w_i \exp(\mathbf{x}'_i \hat{\beta} + \hat{\alpha}) \quad (1)$$

Note that α is just a normalizing constant ensuring that the sum of balancing weights is equal to $\hat{\tau}$. We could also set the target sum to some other (strictly positive) value, say, 1 or W_S . This would only affect α , but not β .

Let $\gamma = (\mu', \tau, \beta', \alpha)'$ be the complete vector of estimates involved in the entropy balancing problem. Rearranging the above formulas for the different elements in γ we can express the model as a system of moment equations given as

$$\frac{1}{W} \sum_{i=1}^N w_i \mathbf{h}_i^\gamma(\gamma) = \mathbf{0} \quad \text{with} \quad \mathbf{h}_i^\gamma(\gamma) = \begin{bmatrix} \mathbf{h}_i^\mu(\gamma) \\ h_i^\tau(\gamma) \\ \mathbf{h}_i^\beta(\gamma) \\ h_i^\alpha(\gamma) \end{bmatrix} = \begin{bmatrix} R_i(\mathbf{x}_i - \mu) \\ WR_i - \tau \\ S_i \exp(\mathbf{x}'_i \beta + \alpha)(\mathbf{x}_i - \mu) \\ S_i \left(\exp(\mathbf{x}'_i \beta + \alpha) - \frac{\tau}{W_S} \right) \end{bmatrix} \quad (2)$$

Following the approach outlined in [Jann \(2020b\)](#), the influence function for $\hat{\gamma}$ can thus be obtained as

$$\text{IF}_i^{\hat{\gamma}} = \mathbf{G}^{-1} \mathbf{h}_i^\gamma(\hat{\gamma}) \quad \text{where} \quad \mathbf{G} = -\frac{1}{W} \sum_{i=1}^N w_i \left. \frac{\partial \mathbf{h}_i^\gamma(\gamma)}{\partial \gamma'} \right|_{\gamma=\hat{\gamma}} \quad (3)$$

Solving the derivatives we get

$$\mathbf{G} = \begin{bmatrix} \mathbf{G}^\mu & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & G^\tau & \mathbf{0} & \mathbf{0} \\ \mathbf{G}^\beta_\mu & \mathbf{0} & \mathbf{G}^\beta & \mathbf{G}^\beta_\alpha \\ \mathbf{0} & G^\alpha_\tau & \mathbf{G}^\alpha_\beta & G^\alpha \end{bmatrix} = -\frac{1}{W} \sum_{i=1}^N w_i \begin{bmatrix} -R_i \mathbf{I}_k & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & -1 & \mathbf{0} & \mathbf{0} \\ -S_i \hat{v}_i \mathbf{I}_k & \mathbf{0} & \mathbf{h}_i^\beta(\hat{\gamma}) \mathbf{x}'_i & \mathbf{h}_i^\beta(\hat{\gamma}) \\ \mathbf{0} & -S_i \frac{1}{W_S} & S_i \hat{v}_i \mathbf{x}'_i & S_i \hat{v}_i \end{bmatrix} \quad (4)$$

where \mathbf{I}_k is the identity matrix of size k and $\hat{v}_i = e^{\mathbf{x}'_i \hat{\beta} + \hat{\alpha}}$. We are only interested in the influence functions for $\hat{\beta}$ and $\hat{\alpha}$. Hence, noting that $\mathbf{G}^\mu = \frac{W_R}{W} \mathbf{I}_k$, $G^\tau = 1$, and $G^\alpha_\tau = \frac{1}{W}$, we collapse the system to

$$\begin{bmatrix} \text{IF}_i^{\hat{\beta}} \\ \text{IF}_i^{\hat{\alpha}} \end{bmatrix} = \begin{bmatrix} \mathbf{G}^\beta & \mathbf{G}^\beta_\alpha \\ \mathbf{G}^\alpha_\beta & G^\alpha \end{bmatrix}^{-1} \begin{bmatrix} \mathbf{h}_i^\beta(\hat{\gamma}) - \frac{W}{W_R} \mathbf{G}^\beta_\mu \mathbf{h}_i^\mu(\hat{\gamma}) \\ h_i^\alpha(\hat{\gamma}) - \frac{1}{W} h_i^\tau(\hat{\gamma}) \end{bmatrix} \quad (5)$$

Furthermore, applying rules for the inversion of a block matrix we can write

$$\begin{bmatrix} \mathbf{G}^\beta & \mathbf{G}_\alpha^{\beta} \\ \mathbf{G}_\beta^\alpha & G^\alpha \end{bmatrix}^{-1} = \begin{bmatrix} \mathbf{A} & -d(\mathbf{G}^\beta)^{-1}\mathbf{G}_\alpha^\beta \\ -\mathbf{G}_\beta^\alpha\mathbf{A}/G^\alpha & d \end{bmatrix} \quad \text{with} \quad \mathbf{A} = \left(\mathbf{G}^\beta - \mathbf{G}_\alpha^\beta\mathbf{G}_\beta^\alpha/G^\alpha\right)^{-1} \\ d = \frac{1}{G^\alpha - \mathbf{G}_\beta^\alpha(\mathbf{G}^\beta)^{-1}\mathbf{G}_\alpha^\beta}$$

such that the influence functions can be expressed as

$$\text{IF}_i^{\hat{\beta}} = \mathbf{A} \left(\mathbf{h}_i^\beta(\hat{\gamma}) - \frac{W}{W_R} \mathbf{G}_\mu^\beta \mathbf{h}_i^\mu(\hat{\gamma}) \right) - d (\mathbf{G}^\beta)^{-1} \mathbf{G}_\alpha^\beta \left(h_i^\alpha(\hat{\gamma}) - \frac{1}{W} h_i^\tau(\hat{\gamma}) \right) \quad (6)$$

$$\text{IF}_i^{\hat{\alpha}} = d \left(h_i^\alpha(\hat{\gamma}) - \frac{1}{W} h_i^\tau(\hat{\gamma}) \right) - \frac{\mathbf{G}_\beta^\alpha \mathbf{A}}{G^\alpha} \left(\mathbf{h}_i^\beta(\hat{\gamma}) - \frac{W}{W_R} \mathbf{G}_\mu^\beta \mathbf{h}_i^\mu(\hat{\gamma}) \right) \quad (7)$$

If balance is achieved, then $\mathbf{G}_\mu^\beta = \frac{\hat{\tau}}{W} \mathbf{I}_k$, $\mathbf{G}_\alpha^\beta = \mathbf{0}$, and $G^\alpha = -\frac{\hat{\tau}}{W}$ such that the influence functions simplify to

$$\text{IF}_i^{\hat{\beta}} = (\mathbf{G}^\beta)^{-1} \left(\mathbf{h}_i^\beta(\hat{\gamma}) - \frac{\hat{\tau}}{W_R} \mathbf{h}_i^\mu(\hat{\gamma}) \right) \quad (8)$$

$$\text{IF}_i^{\hat{\alpha}} = \frac{W}{-\hat{\tau}} \left(h_i^\alpha(\hat{\gamma}) - \frac{1}{W} h_i^\tau(\hat{\gamma}) - \mathbf{G}_\beta^\alpha \text{IF}_i^{\hat{\beta}} \right) \quad (9)$$

In the current setup, note that $\hat{\tau}/W_R = 1$, but we may wish to normalize the weights using some other value for $\hat{\tau}$, in which case $\hat{\tau}/W_R$ would no longer be equal to 1. For example, we may set $\hat{\tau}$ to the sum of base weights in the primary sample, that is, $\hat{\tau} = W_S = \sum_{i \in \mathcal{S}} w_i$. In this case, use $h_i^\tau(\hat{\gamma}) = WS_i - \tau$ in (7) or (9) instead of $h_i^\tau(\hat{\gamma}) = WR_i - \tau$. Alternatively, we may want to set τ to some fixed value, such as $\tau = 1$. In this case, $h_i^\tau(\hat{\gamma}) = 0$. Yet, an advantage of using $\hat{\tau} = W_R$ is that, in this case, $\hat{p}_i = \exp(\mathbf{x}_i' \hat{\beta} + \hat{\alpha}) / (1 + \exp(\mathbf{x}_i' \hat{\beta} + \hat{\alpha}))$ can be interpreted as a propensity score, that is, as an estimate of the conditional probability of belonging to \mathcal{R} rather than \mathcal{S} given \mathbf{x}_i .

In general, it seems justifiable to assume τ as fixed even when it is set to sample quantities such as W_R or W_S . First, the moment condition for τ will only affect the influence function of $\hat{\alpha}$, which is typically only of minor interest (for example, the influence function of $\hat{\alpha}$ is typically not needed when correcting the standard errors of statistics computed from the reweighted data). Second, also for the influence function of $\hat{\alpha}$ the bias introduced by assuming τ as fixed will typically be small. This is why command `ebalfit` discussed below will treat τ as fixed when computing influence functions and standard errors.

2.2 One-sample balancing

In one-sample entropy balancing, the data is adjusted to given values from an external source (e.g. known population averages). To obtain the influence functions for this situation, replace $\hat{\mu}$ by fixed vector μ and replace $\hat{\tau}$ by fixed value τ (e.g. the population size) and let $\gamma = (\beta', \alpha)'$. Compared to the two-sample case, all components related to

the estimation of μ and τ drop out of the system. Hence, we get

$$\text{IF}_i^{\hat{\beta}} = \mathbf{A}\mathbf{h}_i^\beta(\hat{\gamma}) - d(\mathbf{G}^\beta)^{-1}\mathbf{G}_\alpha^\beta h_i^\alpha(\hat{\gamma}) \quad (10)$$

$$\text{IF}_i^{\hat{\alpha}} = d h_i^\alpha(\hat{\gamma}) - \frac{\mathbf{G}_\beta^\alpha \mathbf{A}}{G^\alpha} \mathbf{h}_i^\beta(\hat{\gamma}) \quad (11)$$

and, if balance is achieved,

$$\text{IF}_i^{\hat{\beta}} = (\mathbf{G}^\beta)^{-1} \mathbf{h}_i^\beta(\hat{\gamma}) \quad (12)$$

$$\text{IF}_i^{\hat{\alpha}} = \frac{W}{-\tau} \left(h_i^\alpha(\hat{\gamma}) - \mathbf{G}_\beta^\alpha \text{IF}_i^{\hat{\beta}} \right) \quad (13)$$

2.3 Alternative formulation of the problem

Acknowledging that α is just a normalizing constant, we can also rephrase the problem such that β is estimated independently of α in a first step and α is then determined taken the estimate of β as given. In this case, we first look for β that solves

$$\frac{1}{\hat{\Omega}} \sum_{i \in \mathcal{S}} w_i \exp(\mathbf{x}'_i \hat{\beta}) \mathbf{x}_i = \hat{\mu} \quad (14)$$

where $\hat{\Omega} = \sum_{i \in \mathcal{S}} w_i \exp(\mathbf{x}'_i \hat{\beta})$, and then set α to $\hat{\alpha} = \ln(\hat{\tau}) - \ln(\hat{\Omega})$. The moment conditions for $\gamma = (\mu', \beta)'$, the first part of the problem in the two sample setting, can be written as

$$\mathbf{h}_i^\gamma(\gamma) = \begin{bmatrix} \mathbf{h}_i^\mu(\gamma) \\ \mathbf{h}_i^\beta(\gamma) \end{bmatrix} = \begin{bmatrix} R_i(\mathbf{x}_i - \mu) \\ S_i \frac{1}{\hat{\Omega}} \exp(\mathbf{x}'_i \hat{\beta})(\mathbf{x}_i - \mu) \end{bmatrix} \quad (15)$$

with

$$\begin{aligned} \mathbf{G} &= \begin{bmatrix} \mathbf{G}^\mu & \mathbf{0} \\ \mathbf{G}_\mu^\beta & \mathbf{G}^\beta \end{bmatrix} \\ &= -\frac{1}{W} \sum_{i=1}^N w_i \begin{bmatrix} -R_i \mathbf{I}_k & \mathbf{0} \\ -S_i \frac{1}{\hat{\Omega}} \exp(\mathbf{x}'_i \hat{\beta}) \mathbf{I}_k & \mathbf{h}_i^\beta(\hat{\gamma}) \left(\mathbf{x}'_i - \sum_{i \in \mathcal{S}} w_i \frac{1}{\hat{\Omega}} \exp(\mathbf{x}'_i \hat{\beta}) \mathbf{x}'_i \right) \end{bmatrix} \end{aligned} \quad (16)$$

The influence function of $\hat{\beta}$ can thus be obtained as

$$\text{IF}_i^{\hat{\beta}} = (\mathbf{G}^\beta)^{-1} \left(\mathbf{h}_i^\beta(\hat{\gamma}) - \mathbf{G}_\mu^\beta (\mathbf{G}^\mu)^{-1} \mathbf{h}_i^\mu(\hat{\gamma}) \right) = (\mathbf{G}^\beta)^{-1} \left(\mathbf{h}_i^\beta(\hat{\gamma}) - \frac{1}{W_R} \mathbf{h}_i^\mu(\hat{\gamma}) \right) \quad (17)$$

which reduces to

$$\text{IF}_i^{\hat{\beta}} = (\mathbf{G}^\beta)^{-1} \mathbf{h}_i^\beta(\hat{\gamma}) \quad (18)$$

in the one-sample setting. If balance is achieved, these expressions are formally equivalent to (8) and (12), respectively. The moment condition for α is still the same as before, such that its influence function is given by (9) with $\text{IF}_i^{\hat{\beta}}$ as defined in (17) or, in the one-sample setting, by (13) with $\text{IF}_i^{\hat{\beta}}$ as defined in (18).

2.4 Estimation

We could use `gmm` ([R] `gmm`) to estimate the entropy balancing coefficients based on the moment equations provided in section 2.1. However, given that α is simply a normalization constant, it may be more convenient to first run an optimization algorithm to fit $\hat{\beta}$ and then determine $\hat{\alpha}$ as

$$\hat{\alpha} = \ln(\tau) - \ln\left(\sum_{i \in \mathcal{S}} w_i \exp(\mathbf{x}_i' \hat{\beta})\right) \quad (19)$$

as discussed in section 2.3. This ensures that the sum of balancing weights will always match the target sum of weights. Furthermore, in the two-sample case, the complexity of the estimation can be reduced by computing the target means μ and the target sum of weights τ upfront instead of including them in a joint optimization problem.

To obtain an estimate for β , we can run a standard Newton-Raphson algorithm that minimizes

$$L^\omega = \ln\left(\sum_{i \in \mathcal{S}} w_i \exp((\mathbf{x}_i - \mu)' \beta)\right) = \ln\left(\sum_{i \in \mathcal{S}} \tilde{w}_i\right) \quad \text{where} \quad \tilde{w}_i = w_i \exp((\mathbf{x}_i - \mu)' \beta) \quad (20)$$

with respect to β (also see Hainmueller 2012). The vector of first derivatives of L^ω (the gradient vector) and the matrix of second derivatives (the Hessian), which are required by the Newton-Raphson procedure, are given as

$$\mathbf{g} = \frac{1}{\sum_{i \in \mathcal{S}} \tilde{w}_i} \sum_{i \in \mathcal{S}} \tilde{w}_i (\mathbf{x}_i - \mu) \quad \text{and} \quad \mathbf{H} = \frac{1}{\sum_{i \in \mathcal{S}} \tilde{w}_i} \sum_{i \in \mathcal{S}} \tilde{w}_i (\mathbf{x}_i - \mu) (\mathbf{x}_i - \mu)' \quad (21)$$

In practice, to avoid numerical overflow, we may want to change the definition of \tilde{w} to

$$\tilde{w}_i = w_i \exp((\mathbf{x}_i - \mu)' \beta - c) \quad \text{where} \quad c = \max((\mathbf{x}_i - \mu)' \beta) \quad (22)$$

and redefine L^ω as

$$L^\omega = \ln\left(\sum_{i \in \mathcal{S}} \tilde{w}_i\right) + c \quad (23)$$

Furthermore, instead of using L^ω , one may also determine convergence based on a loss criterion that is directly defined in terms of achieved balance, while still employing the gradient vector and Hessian given in (21) for updating β . For example, we could use the maximum absolute difference

$$L^{\text{absdif}} = \max(\text{abs}(\mathbf{g})) \quad (24)$$

the maximum relative difference

$$L^{\text{reldif}} = \max\left(\frac{\text{abs}(\mathbf{g})}{\text{abs}(\mu) + 1}\right) \quad (25)$$

or the Euclidean norm

$$L^{\text{norm}} = \|\mathbf{g}\| = \sqrt{\mathbf{g}'\mathbf{g}} \quad (26)$$

where \mathbf{g} is the gradient vector as defined above. As can easily be seen from its definition, the gradient vector

$$\mathbf{g} = \frac{1}{\sum_{i \in \mathcal{S}} \tilde{\omega}_i} \sum_{i \in \mathcal{S}} \tilde{\omega}_i (\mathbf{x}_i - \mu) = \left(\frac{1}{\sum_{i \in \mathcal{S}} \tilde{\omega}_i} \sum_{i \in \mathcal{S}} \tilde{\omega}_i \mathbf{x}_i \right) - \mu \quad (27)$$

is equal to the difference between the means of the reweighted data and the target values μ , given the current current values of β . That is, \mathbf{g} quantifies for each variable how well the balancing has been achieved up to that point in the algorithm.

Practical experience indicates that using one of these balancing loss criteria instead of L^ω makes the algorithm more robust in situations where perfect balance is not possible. However, as the optimization criterion is no longer fully consistent with the used gradient and Hessian, the algorithm profits from some standardization of the data (so that the different variables have similar scales). For example, we may obtain the standard deviations

$$\sigma_S = \sqrt{\frac{1}{W_S} \sum_{i \in \mathcal{S}} w_i (\mathbf{x}_i - \bar{\mathbf{x}}_S)^2} \quad \text{with} \quad \bar{\mathbf{x}}_S = \frac{1}{W_S} \sum_{i \in \mathcal{S}} w_i \mathbf{x}_i \quad (28)$$

from the primary sample and then use \mathbf{x}_i/σ_S and μ/σ_S instead of \mathbf{x}_i and μ in equations (20) to (27). Before computing $\hat{\alpha}$ in (19), back-transform the resulting estimate for β by dividing it by σ_S .

Furthermore, as usual, collinear terms have to be excluded from estimation. These terms, however, are relevant for the evaluation of final quality of the achieved balancing (collinear terms may remain unbalanced). My suggestion thus is to use \mathbf{x}_i^{nc} , a variant of \mathbf{x}_i without elements that are collinear in \mathcal{S} , for estimation of β (with elements corresponding to collinear terms set to 0) and then evaluate the final fit based on the complete data by applying one of the above loss functions to

$$\hat{\mathbf{g}} = \frac{1}{\sum_{i \in \mathcal{S}} \hat{\omega}_i} \sum_{i \in \mathcal{S}} \hat{\omega}_i \mathbf{x}_i - \mu \quad \text{with} \quad \hat{\omega}_i = w_i \exp(\mathbf{x}_i' \hat{\beta} + \hat{\alpha}) \quad (29)$$

The variance-covariance matrix of $\hat{\gamma} = (\hat{\beta}', \hat{\alpha})'$ can be estimated by taking the total of the squared influence functions divided by the sum of weights. To be precise, if the base weights are frequency weights (or if there are no base weights, i.e. $w_i = 1$ for all observations and $W = N$), then

$$\hat{V}(\hat{\gamma}) = \frac{W}{W - k - 1} \sum_{i=1}^N w_i \lambda_i \lambda_i' \quad \text{with} \quad \lambda_i = \frac{1}{W} \begin{bmatrix} \text{IF}_i^{\hat{\beta}} \\ \text{IF}_i^{\hat{\alpha}} \end{bmatrix} \quad (30)$$

If the base weights are probability (sampling) weights, then

$$\hat{V}(\hat{\gamma}) = \frac{N}{N - k - 1} \sum_{i=1}^N w_i^2 \lambda_i \lambda_i' \quad \text{with} \quad \lambda_i = \frac{1}{W} \begin{bmatrix} \text{IF}_i^{\hat{\beta}} \\ \text{IF}_i^{\hat{\alpha}} \end{bmatrix} \quad (31)$$

Complex survey design such as clustering or stratification can be taken into account by appropriately modifying the aggregation. In practice, variance estimates can be obtained by applying command [R] **total** to λ_i , possibly including the [SVY] **svy** prefix.

□ **Technical note**

Instead of running a Newton-Raphson algorithm as described above we may also mimic the behavior of [R] **gmm** and apply the Gauss-Newton method to criterion

$$Q^\beta = (\mathbf{q}^\beta)' \mathbf{q}^\beta \quad \text{with} \quad \mathbf{q}^\beta = \sum_{i \in \mathcal{S}} \mathbf{h}_i^\beta \quad (32)$$

where \mathbf{h}_i^β is as given in (15). This is equivalent to a Newton-Raphson algorithm with gradient and Hessian set to

$$\mathbf{g} = \mathbf{q}^\beta \mathbf{G}^\beta \quad \text{and} \quad \mathbf{H} = (\mathbf{G}^\beta)' \mathbf{G}^\beta \quad (33)$$

with \mathbf{G}^β given in (16). □

2.5 Balancing of higher-order moments and covariances

In the exposition above I only considered balancing of first moments (i.e. the means), but entropy balancing can easily be extended to higher moments such as the variance or the skewness or even to covariances. The balancing constraints for higher moments and covariances are equivalent to first-moment balancing constraints for specific transformations of the variables. Rather than extending the above exposition to cover higher moments, we may thus simply change the definition of the data. To balance the variance of a variable X in addition to its mean, add x_i^2 to the data. To also balance the skewness, add x_i^3 . To balance the covariance between two variables X_1 and X_2 , include the product $x_{i1}x_{i2}$ in the data.

2.6 Correcting standard errors of reweighted estimators

Define $\hat{\omega}_i = w_i \hat{v}_i$ with

$$\hat{v}_i = \begin{cases} \exp(\mathbf{x}'_i \hat{\beta} + \hat{\alpha}) & \text{if } S_i = 1 \\ 1 & \text{else} \end{cases} \quad (34)$$

That is, for observations within the reweighted sample, $\hat{\omega}_i$ is equal to the balancing weight, for all other observations, $\hat{\omega}_i$ is equal to the base weight. Most estimators can be expressed as a system of moment equations

$$\frac{1}{\sum_{i=1}^N \hat{\omega}_i} \sum_{i=1}^N \hat{\omega}_i \mathbf{h}_i^\theta(\theta) = \mathbf{0} \quad (35)$$

such that $\hat{\omega}_i$ does not appear in $\mathbf{h}_i^\theta(\theta)$. For such estimators, the necessary correction to take account of the uncertainty imposed by the estimation of the balancing weights has a very simple form. Re-expressing the system as

$$\frac{1}{W} \sum_{i=1}^N w_i \left(\frac{\hat{v}_i}{c} \mathbf{h}_i^\theta(\theta) \right) = \mathbf{0} \quad \text{with} \quad c = \frac{1}{W} \sum_{i=1}^N \hat{\omega}_i \quad (36)$$

we see that we can obtain the adjusted influence function as

$$\text{IF}_i^{\hat{\theta}} = \frac{\hat{v}_i}{c} \tilde{\text{IF}}_i^{\hat{\theta}} - \mathbf{G}_{\beta}^{\tilde{\text{IF}}} \text{IF}^{\hat{\beta}} - \mathbf{G}_{\alpha}^{\tilde{\text{IF}}} \text{IF}^{\hat{\alpha}} \quad (37)$$

with

$$\mathbf{G}_{\beta}^{\tilde{\text{IF}}} = -\frac{1}{W} \sum_{i=1}^N w_i S_i \frac{\hat{v}_i}{c} \tilde{\text{IF}}_i^{\hat{\theta}} \mathbf{x}_i' \quad \mathbf{G}_{\alpha}^{\tilde{\text{IF}}} = -\frac{1}{W} \sum_{i=1}^N w_i S_i \frac{\hat{v}_i}{c} \tilde{\text{IF}}_i^{\hat{\theta}} \quad (38)$$

where $\tilde{\text{IF}}_i^{\hat{\theta}}$ is the influence function of $\hat{\theta}$ assuming the weights $\hat{\omega}_i$ as fixed. Since $\mathbf{G}_{\alpha}^{\tilde{\text{IF}}} = \mathbf{0}$ by definition, the corrected influence function simplifies to

$$\text{IF}_i^{\hat{\theta}} = \frac{\hat{v}_i}{c} \tilde{\text{IF}}_i^{\hat{\theta}} - \mathbf{G}_{\beta}^{\tilde{\text{IF}}} \text{IF}^{\hat{\beta}} \quad (39)$$

To summarize, we can first compute the influence function for $\hat{\theta}$ in the usual way, as if balancing weights were fixed, and then adjust the influence function using equation (39). Naturally, we need a way to obtain the (unadjusted) influence function of our estimator in the first place, but in many cases this is not very difficult (for example, see [Jann 2020b](#) for practical instruction on how to obtain influence functions for maximum-likelihood models given the results returned by Stata).²

3 Stata implementation

Command `ebalfit`, available from the SSC Archive, implements the methods described above. To install the command on your system, type

```
. ssc install ebalfit
```

The heavy lifting is done by Mata function `mm_ebalance()` that is provided as part of the `moremata` library ([Jann 2005](#)), also available from the SSC Archive. To be able to run `ebalfit`, the latest update of `moremata` is required. To install `moremata`, type

```
. ssc install moremata, replace
```

2. In the above derivation I assumed c , which depends on the relative size of the reweighted group (i.e. the sum of balancing weights) with respect to the size (sum of base weights) of the rest of the data, to be fixed. This is valid as long as the statistic conditions on S_i such that the sum of balancing weights does not matter or if $\tau = W_S$ such that c is always equal to 1. In other cases the true correction would be more complicated, but the bias introduced by assuming c as fixed should be negligible in most situations.

The exposition below focuses on Stata command `ebalfit` and does not provide details on Mata function `mm_ebalance()`. Users interested in applying `mm_ebalance()` directly can type `help mata mm_ebalfit()` after installation to view its documentation.

3.1 Syntax

Syntax 1: adjust a subsample to values from another subsample (two-sample balancing)

```
ebalfit varlist [if] [in] [weight], by(varname) [options]
```

Syntax 2: adjust a sample to population values (one-sample balancing)

```
ebalfit varlist [if] [in] [weight], population([size:]numlist) [options]
```

Replay results

```
ebalfit [, reporting_options]
```

where *reporting_options* are as described under “Reporting” in Section [3.2](#)

Generate predictions after estimation

```
predict [type] newvar [if] [in] [, predict_options]
```

where *predict_options* are

<code>w</code>	generate balancing weights (the default)
<code>u</code>	generate raw balancing weights (i.e. without base weights)
<code>pr</code>	generate propensity scores
<code>pscore</code>	synonym for <code>pr</code>
<code>xb</code>	generate linear predictions

Generate influence functions after estimation

```
predict [type] { stub* | newvarlist } [if] [in], ifs [IF_options]
```

where *IF_options* are

<code>nocons</code>	skip influence function for α ; only relevant with <i>stub*</i>
<code>noalpha</code>	synonym for <code>nocons</code>

In both syntax 1 and syntax 2, *varlist* may contain factor variables (see [U] [11.4.3 Factor variables](#)). `fweights`, `pweights`, and `iweights` are allowed (see [U] [11.1.6 weight](#)).

3.2 Options

Main

`by(groupvar)` is required in syntax 1 and identifies the subsamples. *groupvar* must be integer and nonnegative and must identify exactly two groups. By default, the lower value identifies the subsample to be reweighted and the higher value identifies the reference subsample. Also see option `swap`.

`swap` swaps the subsamples (only allowed in syntax 1). By default, the lower value of *groupvar* identifies the subsample to be reweighted. Specify `swap` to use the higher value of *groupvar* as the subsample to be reweighted.

`pooled` uses the pooled sample across both groups as the reference sample (only allowed in syntax 1). If `pooled` is specified, the selected subsample will be reweighted with respect to the overall sample (rather than with respect to the other subsample).

`population(spec)` is required in syntax 2. Use this option to specify the size of the population as well the population averages to which the sample should be reweighted. The syntax of *spec* is

$$[\textit{popsiz}e :] \textit{numlist}$$

where *popsiz*e is the size of the population and *numlist* provides the population averages of the variables. *numlist* must contain one value for each variable. If *popsiz*e is omitted, it will be set to the sum of weights in the sample.

`tau(spec)` specifies a custom target sum of weights for the balancing weights within the reweighted sample. *spec* may either be real number ($\# > 0$) or one of `Wref` (sum of base weights in the reference sample), `W` (sum of base weights in the reweighted sample), `Nref` (number of rows the reference sample), or `N` (number of rows the reweighted sample). The default is `Wref`.

`scales(spec)` determines the scales to be used for standardization during estimation (unless `nostd` is specified) and for computation of standardized differences in the balancing table. *spec* may either be a *numlist* containing custom values (one for each term in the model; the values must be positive) or, alternatively, `main` (use standard deviations from the main sample), `reference` (use standard deviations from the reference sample), `average` (use standard deviations averaged between the two samples), `waverage` (use standard deviations averaged between the two samples, weighted by sample size), `pooled` (use standard deviations from the pooled sample). `reference`, `average`, `waverage`, and `pooled` are only allowed in syntax 1. Standard deviations are computed using population formulas (division by N rather than $N-1$). Scales equal to 0 will be reset to 1. The default is `main`.

`targets(options)` specifies the types of moments to be balanced. *options* are:

<code>mean</code>	balance means (the default)
<code>variance</code>	balance variances (implies <code>mean</code>)
<code>skewness</code>	balance skewnesses (implies <code>mean</code> and <code>variance</code>)
<code>covariance</code>	balance covariances (implies <code>mean</code>)

By default, only the means of the specified variables will be balanced. If you type, for example, `targets(variance)`, then the variances of the variables will be balanced in addition to the means. Balancing of higher moments and covariances is implemented by adding extra terms to `varlist` before running the balancing algorithm. For example, `variance` will add `c.varname#c.varname` for each continuous variable in `varlist` (skipping omitted terms). Likewise, `covariance` will add `c.varname1#c.varname2` for each combination of continuous variables. Factor variables will be ignored by `variance` and `skewness`, but `covariance` will consider them and add appropriate interaction terms such as `1.fvvar#c.varname` (skipping base levels).

If option `targets()` is specified, interaction terms such as `i.fvvar#c.varname` are not allowed in `varlist`. However, interactions are allowed if option `targets()` is omitted. For example, you could type

```
c.hours##c.tenure i.south i.south#c.tenure
```

to balance the means of `hours` and `tenure`, the covariance between `hours` and `tenure`, the proportions of the levels of `south`, as well as the averages of `tenure` within levels of `south` (see [U] 11.4.3 **Factor variables** for details on notation). That is, you can use custom interactions as an alternative to option `targets()` if you want to have more control over the exact configuration of moments to be balanced.

[`no`] `adjust(numlist)` selects the terms to be balanced. Use this option if you want to construct weights such that only a subset of terms is adjusted, while keeping the others fixed. `numlist` provides the indices of the relevant terms. For example, in a model with three variables, to adjust the means of the first two variables and keep the mean of the third variable fixed, type `adjust(1 2)` or, equivalently, `noadjust(3)`. Keeping terms fixed leads to different results than excluding the terms from the model.

Reporting

`level(#)` specifies the confidence level, as a percentage, for confidence intervals. The default is `level(95)` or as set by `set level` (see [R] `level`).

`noheader` suppresses the display of the header.

`nowtable` suppress the display of the summary table of balancing weights.

`notable` suppresses the display of the coefficient table.

`display_options` are standard reporting options to be applied to the coefficient table, such as `eformat()`, or `coeflegend`; see [R] *eform_option* and the Reporting options in [R] **Estimation options**.

`bal` displays a balancing table in addition to the table of coefficients. The balancing table contains for each term the target value, the unbalanced value, the standardized difference between the target value and the unbalanced value, the balanced value, and the standardized difference between the target value and the balanced value.

VCE/SE

`vce(vcetype)` determines how standard errors are computed. `vcetype` may be:

```

robust
cluster clustvar
none

```

`vce(robust)`, the default, computes standard errors based on influence functions. Likewise, `vce(cluster clustvar)` computes standard errors based on influence function allowing for intragroup correlation, where `clustvar` specifies to which group each observation belongs. `vce(none)` omits the computation of standard errors.

`cluster(clustvar)` can be used as a synonym for `vce(cluster clustvar)`.

`nose` omits the computation of standard errors. Use this option to save computer time. `nose` is a synonym for `vce(none)`.

Generate

`generate(newvar)` stores the balancing weights in `newvar`. Alternatively, use command `predict` to generate the balancing weights after estimation.

In syntax 1, weights will be filled in for both the reweighted subsample and the reference subsample, using a copy of the base weights for the latter (or 1 if there are no base weights).

`ifgenerate(names)` stores the influence functions of the coefficients. `names` is either a list of (new) variable names or `stub*` to create names `stub1`, `stub2`, etc. Alternatively, use command `predict` with option `ifs` to generate the influence functions after estimation. In any case, the influence functions will be scaled in a way such that command [R] `total` can be used to estimate the variance-covariance matrix (that is, compared to the expressions provided above, the stored influence functions will be divided by the sum of weights in the overall sample).

`nodescribe` suppresses the list of generated variables that is displayed in the output by default when `generate()` or `ifgenerate()` is specified.

`replace` allows replacing existing variables.

Optimization

`btolerance(#)` sets the balancing tolerance. Balance is achieved if the balancing loss is smaller than the balancing tolerance. The default is `btolerance(1e-6)`.

`ltype(ltype)` sets the type of loss function to be used to evaluate balancing. `ltype` can be `reldif` (maximum relative difference), `absdif` (maximum absolute difference), or `norm` (norm of differences). The default is `reldif`.

`etype(etype)` selects the evaluator to be used to fit the coefficients. `etype` can be `bl` (evaluator based on the balancing loss), `wl` (evaluator based on distribution of weights, i.e. criterion L^ω from equation 20), `mm` (method of moments evaluator), or `mma` (method of moments evaluator including the intercept). The default is `bl`. Irrespective of the choice of evaluator, balancing loss will be used to evaluate the final fit.

`iterate(#)` specifies the maximum number of iterations. Error will be returned if convergence is not reached within the specified maximum number of iterations. The default is as set by `set maxiter` ([R] `set.iter`).

`ptolerance(#)` specifies the convergence tolerance for the coefficient vector. Convergence is reached if `ptolerance()` or `vtolerance()` is satisfied. See [M-5] `optimize()` for details. The default is `ptolerance(1e-6)`.

`vtolerance(#)` specifies the convergence tolerance for the balancing loss. Convergence is reached if `ptolerance()` or `vtolerance()` is satisfied. See [M-5] `optimize()` for details. The default is `vtolerance(1e-7)` in case of `etype(bl)` and `vtolerance(1e-10)` else.

`difficult` uses a different stepping algorithm in nonconcave regions. See the singular H methods in [M-5] `optimize()` and the description of the `difficult` option in [R] `Maximize`.

`nostd` omits standardization of the data during estimation. Specifying `nostd` is not recommended.

`nolog` suppresses the display of progress information.

`relax` causes `ebalfit` to proceed even if convergence or balance is not achieved. `ebalfit` uses formulas assuming balance when computing influence functions and standard errors. The stored influence functions and reported standard errors will be invalid if balance has not been achieved.

`nowarn` suppresses any “convergence not achieved” or “balance not achieved” messages. This is only relevant if option `relax` has been specified.

3.3 Stored results

`ebalfit` stores its results in `e()` similar to any other estimation command (see [R] `Stored results`). See `help ebalfit` for a complete list of saved results.

4 Examples

4.1 Balancing two samples

Consider the data from [LaLonde \(1986\)](#), provided by [Dehejia and Wahba \(1999\)](#) at <http://users.nber.org/~rdehejia/nswdata.html>. The following code combines a subset of the treatment group from the NSW training program with one of the PSID comparison groups.

```
. use http://users.nber.org/~rdehejia/data/nsw_dw.dta, clear
. keep if treat==1
(260 observations deleted)
. append using http://users.nber.org/~rdehejia/data/psid_controls2.dta
```

For purpose of exposition, I additionally generate some sampling weights (normalized such that the group sizes are preserved). I also set the default storage type for new variables to `double` so that some of the results below will have less roundoff error.

```
. set type double
. set seed 32387939
. generate w0 = runiform()
. summarize w0 if treat==0, meanonly
. quietly replace w0 = w0 * r(N) / r(sum) if treat==0
. summarize w0 if treat==1, meanonly
. quietly replace w0 = w0 * r(N) / r(sum) if treat==1
```

The focus of the LaLonde data lies on the comparison of `re78` (real earnings in 1978 after the program intervention) between the (experimental) treatment group and the (non-experimental) control group. The comparison is not straight forward as there are substantial differences between the two groups in terms of pre-treatment characteristics. Members of the treatment group are younger, more often black, less often married, more often without college degree, and have lower pre-treatment earnings than members of the control group:

```
. table () (treat) [pw=w0], stat(mean age-re75) nototal
```

	treat	
	0	1
age	37.39198	26.02386
education	10.50566	10.52379
black	.4359494	.8572367
hispanic	.0722888	.0567015
married	.7481573	.1944541
nodegree	.5295137	.6608205
re74	10401.03	2230.392
re75	7230.567	1682.258

Various techniques such as matching or inverse probability weighting (IPW) have been

proposed in the literature to address the problem of making the groups comparable such that the average effect of program participation (the ATET) can be estimated consistently. Inverse probability weights, for example, could be obtained as follows:

```
. logit treat age-re75 [pw=w0], nolog
Logistic regression                               Number of obs =   438
                                                Wald chi2(8)    =  93.08
                                                Prob > chi2    =  0.0000
Log pseudolikelihood = -159.20379                Pseudo R2      =  0.4663
```

treat	Robust		z	P> z	[95% conf. interval]	
	Coefficient	std. err.				
age	-.092245	.0158144	-5.83	0.000	-.1232408	-.0612493
education	.0642543	.0870386	0.74	0.460	-.1063382	.2348467
black	1.932721	.4336358	4.46	0.000	1.08281	2.782631
hispanic	1.671454	.543556	3.08	0.002	.6061038	2.736804
married	-1.290829	.3192307	-4.04	0.000	-1.91651	-.665148
nodegree	.2890979	.4642715	0.62	0.533	-.6208574	1.199053
re74	-.0000947	.0000395	-2.40	0.017	-.0001721	-.0000172
re75	-.0000944	.000078	-1.21	0.226	-.0002472	.0000584
_cons	1.649338	1.529141	1.08	0.281	-1.347723	4.646399

```
. predict pscore if treat==0, pr
(185 missing values generated)
. generate ipw = w0 * cond(treat==0, pscore/(1-pscore), 1)
. drop pscore
. table () (treat) [pw=ipw], stat(mean age-re75 re78) nottotal
```

	treat	
	0	1
age	25.21257	26.02386
education	10.70952	10.52379
black	.8905226	.8572367
hispanic	.0234441	.0567015
married	.1699626	.1944541
nodegree	.5972221	.6608205
re74	2814.793	2230.392
re75	2433.187	1682.258
re78	5088.788	6004.657

This worked quite well and much of the group differences disappeared, but there are still some non-negligible discrepancies, especially with respect to pre-treatment earnings. We can now try to improve the reweighting using entropy balancing:

```
. ebalfit age-re75 [pw=w0], by(treat)
Iteration 0: balancing loss = .88095577
Iteration 1: balancing loss = .20574871
Iteration 2: balancing loss = .11227971
Iteration 3: balancing loss = .01088361
Iteration 4: balancing loss = .00056568
Iteration 5: balancing loss = 1.833e-06
Iteration 6: balancing loss = 1.884e-11
Iteration 7: balancing loss = 9.108e-17
```

```

Final fit:      balancing loss = 2.038e-16
Entropy balancing      Number of obs   =      438
                        Wald chi2(8)    =      60.71
                        Prob > chi2     =      0.0000
                        Evaluator       =      bl
Main      = 0.treat (253 obs)
Reference = 1.treat (185 obs)
                        Loss type       =      reldif
                        Balancing loss   =      2.038e-16

```

balancing weights					
minimum	average	maximum	total	CV	DEFF
6.722e-09	.7312253	28.603379	185	3.3617903	12.301634

	Coefficient	Robust std. err.	z	P> z	[95% conf. interval]	
age	-.0962814	.0299907	-3.21	0.001	-.155062	-.0375008
education	.0894269	.1486163	0.60	0.547	-.2018557	.3807096
black	1.640353	.5943787	2.76	0.006	.4753918	2.805314
hispanic	2.386918	.7496026	3.18	0.001	.917724	3.856112
married	-1.079562	.4866648	-2.22	0.027	-2.033407	-.1257165
nodegree	.8138536	.7268795	1.12	0.263	-.6108041	2.238511
re74	-.000121	.0000509	-2.38	0.017	-.0002208	-.0000212
re75	-.000174	.0000818	-2.13	0.033	-.0003343	-.0000136
_cons	1.585729	2.51717	0.63	0.529	-3.347834	6.519292

Option `by()` identifies the groups to be compared; the specified variable must be dichotomous (e.g. 0 and 1). By default, `ebalfit` takes the group with the lower value as the group to be reweighted and takes the other group as the reference group. Specify option `swap` to switch the groups.

The coefficients displayed by `ebalfit` are similar to the coefficients of the `logit` model above. In fact, the coefficients do have a similar interpretation: a positive effect means that people with high values on the respective variable tend to be overrepresented in the reference group (and vice versa).

The output contains some more information that is relevant. For example, the “balancing loss” is a measure of how well `ebalfit` managed to balance the data. In the current situation, perfect balancing could be achieved as the balancing loss is essentially zero.³ Furthermore, some information on the distribution of the weights is provided. `CV` is the coefficient of variation of the weights, defined as

$$CV = \frac{\sqrt{\frac{1}{N_S} \sum_{i \in S} (\hat{\omega}_i - \bar{\omega}_S)^2}}{\bar{\omega}_S} \quad \text{with} \quad \bar{\omega}_S = \frac{1}{N_S} \sum_{i \in S} \hat{\omega}_i$$

where summation is across the reweighted group (N_S is the number of observations in the reweighted group); `DEFF` is the “design effect” of the weights based on Kish’s

3. `ebalfit` returns error if perfect balance cannot be achieved, unless option `relax` is specified. The critical value for “perfect balance” can be set using option `btolerance()`. By default, the critical value is set to 10^{-6} , that is, a solution is considered as balanced if balancing loss, the maximum relative difference between the reweighted means and the target values, is smaller than 0.000001.

formula for the effective sample size (Kish 1965), that is

$$\text{DEFF} = \frac{N_S \sum_{i \in S} \hat{\omega}_i^2}{\left(\sum_{i \in S} \hat{\omega}_i\right)^2}$$

Both statistics indicate that there is large variation in the weights. Apparently, the two groups are very different and balancing them is an ambitious exercise.

As mentioned, however, despite the difficulty of the problem, the output by `ebalfit` tells us that perfect balance has been achieved. We can confirm that this is true by replaying results with option `baltable` to displaying the balancing table that is provided by `ebalfit` (but is suppressed in the output by default):

```
. ebalfit, baltable noheader nowtable notable
```

```
Balancing table
```

	Target value	Unbalanced		Balanced	
		value	std. dif.	value	std. dif.
age	26.02386	37.39198	.9464147	26.02386	-2.96e-16
education	10.52379	10.50566	-.0055215	10.52379	0
black	.8572367	.4359494	-.8495743	.8572367	-2.24e-16
hispanic	.0567015	.0722888	.0601908	.0567015	1.61e-16
married	.1944541	.7481573	1.275604	.1944541	0
nodegree	.6608205	.5295137	-.2630724	.6608205	0
re74	2230.392	10401.03	.775686	2230.392	-4.32e-17
re75	1682.258	7230.567	.6519779	1682.258	-2.67e-17

Options `noheader`, `nowtable`, and `notable` have been specified so that the default output is not displayed again. As is evident, the reweighted means (column “Balanced value”) perfectly match the target values (column “Target value”). The standardized difference between the target value and and the balanced value is essentially zero for all variables.

If we still do not trust this result, we can use `predict` to generate the balancing weights and then construct a balancing table manually:

```
. predict wbal
```

```
. table () (treat) [pw=wbal], stat(mean age-re75) nottotal
```

	treat	
	0	1
age	26.02386	26.02386
education	10.52379	10.52379
black	.8572367	.8572367
hispanic	.0567015	.0567015
married	.1944541	.1944541
nodegree	.6608205	.6608205
re74	2230.392	2230.392
re75	1682.258	1682.258

A comparison of the weights from IPW and the weights from entropy balancing reveals that the latter contain more variation⁴

```
. dstat (cv0) ipw wbal if treat==0
cv0                                Number of obs   =           253
```

	Coefficient	Std. err.	[95% conf. interval]	
ipw	3.069078	.2628062	2.551501	3.586654
wbal	3.36179	.4408397	2.493591	4.22999

```
. program DEFF
1.  syntax varname [if]
2.  tempvar x2
3.  quietly generate `x2' = `varlist'^2
4.  summarize `x2' `if', meanonly
5.  local NX2 = r(sum) * r(N)
6.  summarize `varlist' `if', meanonly
7.  display as res `NX2'/r(sum)^2
8.  end
. DEFF ipw if treat==0
10.419239
. DEFF wbal if treat==0
12.301634
```

Apparently, the better balance came at the cost of more variation in the weights. Large variation in weights generally reduces statistical efficiency so that weights with lower variation may be preferable. As illustrated below, however, this is not necessarily true for treatment effect analyses because the degree to which the weights balance the data also plays a role for the efficiency of the estimate. Yet, for some applications, for example when using entropy balancing to construct sampling weights, we might want to apply some trimming to the resulting weights to reduce the design effect without sacrificing too much precision in balance⁵

4.2 Computing a treatment effect with corrected standard errors

We now continue estimating the treatment effect on post-treatment earnings. The naive estimate of the ATET (average treatment effect on the treated) is negative:

```
. mean re78 [pw=w0], over(treat)
Mean estimation                                Number of obs = 438
```

	Mean	Std. err.	[95% conf. interval]	
c.re78@treat				
0	9104.129	758.2113	7613.935	10594.32

4. I use command `dstat` (Jann 2020a), available from the SSC Archive, because it allows computing the CV in the same way as `ebalfit` does. The CV could also be computed using [R] `tabstat`, which applies a slightly different definition (division by $N - 1$ rather than N in the variance).

5. Also see Kranker et al. (2020) who propose penalized CBPS to address this issue (on CBPS see footnote 1).

	1	6004.657	567.8919	4888.518	7120.796
<code>. lincom _b[c.re78@1.treat] - _b[c.re78@0bn.treat]</code>					
<code>(1) - c.re78@0bn.treat + c.re78@1.treat = 0</code>					
Mean	Coefficient	Std. err.	t	P> t	[95% conf. interval]
(1)	-3099.472	947.3043	-3.27	0.001	-4961.311 -1237.633

However, as seen above, the two groups are very different in terms of pre-treatment characteristics. Using IPW or entropy balancing to remove these discrepancies, the treatment effect estimate becomes positive:

```
. mean re78 [pw=ipw], over(treat)
Mean estimation                               Number of obs = 438
```

	Mean	Std. err.	[95% conf. interval]	
c.re78@treat				
0	5088.788	943.9743	3233.493	6944.082
1	6004.657	567.8919	4888.518	7120.796

```
. lincom _b[c.re78@1.treat] - _b[c.re78@0bn.treat]
( 1) - c.re78@0bn.treat + c.re78@1.treat = 0
```

Mean	Coefficient	Std. err.	t	P> t	[95% conf. interval]	
(1)	915.8695	1101.63	0.83	0.406	-1249.282	3081.021

```
. drop ipw
. mean re78 [pw=wbal], over(treat)
Mean estimation                               Number of obs = 438
```

	Mean	Std. err.	[95% conf. interval]	
c.re78@treat				
0	4174.016	999.5839	2209.426	6138.605
1	6004.657	567.8919	4888.518	7120.796

```
. lincom _b[c.re78@1.treat] - _b[c.re78@0bn.treat]
( 1) - c.re78@0bn.treat + c.re78@1.treat = 0
```

Mean	Coefficient	Std. err.	t	P> t	[95% conf. interval]	
(1)	1830.641	1149.639	1.59	0.112	-428.8669	4090.15

```
. drop wbal
```

The two effect estimates are not statistically significant, but note that we did not yet correct the standard errors for the fact that the balancing weights are estimated. To do so for the estimate based on entropy balancing, we can use the formulas provided in sec-

tion 2.6. As inputs we need the influence functions of the entropy balancing coefficients as well as the influence functions of the mean estimates assuming the balancing weights as fixed. The former we can obtain by applying command `predict` after `ebalfit`; the latter we can compute as $IF^{\hat{\mu}} = \frac{W}{W_S} S_i(x_i - \hat{\mu})$ where x_i is the variable of interest, S_i is an indicator for the analyzed subsample, W_S is the sum of weights in the subsample, and W is the overall sum of weights. In the computations below I omit the leading W because this is how `ebalfit` defines influence functions and because it implies that factor c in the correction formulas will be equal to 1 and can be omitted. To obtain standard errors from influence functions that are scaled in this way command `[R] total` can be used (rather than command `[R] mean`).

```
. ebalfit age-re75 [pw=w0], by(treat)
      (output omitted)
. predict wbal
. predict IFeb*, ifs noalpha // the IF for the constant is not needed
. summarize re78 if treat==0 [aw=wbal], meanonly
. generate IFy0 = (treat==0) * (re78 - r(mean)) / r(sum_w)
. summarize re78 if treat==1 [aw=wbal], meanonly
. generate IFy1 = (treat==1) * (re78 - r(mean)) / r(sum_w)
. total IFy0 IFy1 [pw=wbal]
```

Total estimation		Number of obs = 438		
	Total	Std. err.	[95% conf. interval]	
IFy0	-5.68e-13	999.5839	-1964.59	1964.59
IFy1	5.26e-13	567.8919	-1116.139	1116.139

Note how `total` applied to the influence functions of the two mean estimates reproduces the standard errors reported by `mean` above. We can now correct the influence functions using the formulas from section 2.6. We only need to correct `IFy0`, the influence function of the mean estimate in the control group because in the treatment group we did not apply any reweighting.

```
. mata:
----- mata (type end to exit) -----
: // data
: grp = st_data(., "treat")
: X = st_data(., "age-re75")
: IFy0 = st_data(., "IFy0")
: IFeb = st_data(., "IFeb*")
: wbal = st_data(., "wbal")
: w0 = st_data(., "w0")
: // compute (negative of) G
: G = colsum(select(wbal :* IFy0 :* X, grp==0))'
: // adjust IF
: st_store(., st_addvar("double", "IFy0c"), wbal ./ w0 :* IFy0 + IFeb * G)
: end
```

To compute the corrected standard error of the reweighted mean difference take the total of the difference between the (corrected) influence functions of the two means:

```
. generate IFte = IFy1 - IFy0c
. total IFy0c IFy1 IFte [pw=w0]
Total estimation                               Number of obs = 438
```

	Total	Std. err.	[95% conf. interval]	
IFy0c	-8.53e-14	750.6493	-1475.332	1475.332
IFy1	5.26e-13	567.8919	-1116.139	1116.139
IFte	5.40e-13	906.1741	-1781.001	1781.001

```
. drop IF*
```

We see how taking account of the estimated nature of the balancing weights reduces the standard error of the mean estimate in the control group and also brings down the standard error of the treatment effect estimate, such that the treatment effect estimate is now statistically significant ($t = 1830.6/906.2 = 2.02$, $p = 0.043$).⁶

As mentioned above, entropy balancing is doubly-robust so that applying a regression adjustment model to the reweighted data does not change the estimate of the treatment effect (as long as the same covariates are used in the regression adjustment). I illustrate this here by running [TE] **teffects ra** including the entropy balancing weights:

```
. teffects ra (re78 age-re75) (treat) [pw=wbal], atet
Iteration 0:  EE criterion = 3.671e-22
Iteration 1:  EE criterion = 1.249e-24
Treatment-effects estimation           Number of obs   =       438
Estimator      : regression adjustment
Outcome model  : linear
Treatment model: none
```

re78	Coefficient	Robust std. err.	z	P> z	[95% conf. interval]	
ATET treat (1 vs 0)	1830.641	905.139	2.02	0.043	56.60155	3604.681
POmean treat 0	4174.016	749.7919	5.57	0.000	2704.45	5643.581

```
. display _se[ATET:r1vs0.treat] * sqrt(e(N) / (e(N)-1))
906.17408
```

The estimate is still the same and also the standard error is identical even though regression adjustment assumed the balancing weights as fixed (the small difference is

6. The appendix illustrates how a similar correction can be implemented for IPW.

because `teffects` divides by N while `total` divides by $N - 1$ when computing the variance). That is, if regression adjustment is used to analyze the balanced data no further correction is needed for the standard errors (as long as the same covariates are used in both models). This may seem remarkable, but it is a direct consequence of the doubly-robust property.

4.3 Balancing variances and covariance

In the example above we just balanced the means of the different covariates. To also balance higher moments or covariances we can add powers interactions to the model. Here is an example using a reduced set of covariates:

```
. ebalfit c.age##c.age c.education##c.education 1.black ///
> c.age#c.education 1.black#c.(age education), by(treat) nolog vsquish
Entropy balancing          Number of obs   =      438
                          Wald chi2(8)    =      80.49
                          Prob > chi2     =      0.0000
                          Evaluator       =      b1
Main      = 0.treat (253 obs)           Loss type   =      reldif
Reference = 1.treat (185 obs)          Balancing loss = 2.532e-15
```

balancing weights					
minimum	average	maximum	total	CV	DEFF
.00023148	.7312253	13.072914	185	1.7253741	3.9769159

	Coefficient	Robust std. err.	z	P> z	[95% conf. interval]	
age	-.3454953	.1819311	-1.90	0.058	-.7020737	.0110831
c.age#c.age	-.0007966	.0017921	-0.44	0.657	-.004309	.0027159
education	-.6033715	.8552482	-0.71	0.481	-2.279627	1.072884
c.education#						
c.education	-.0240512	.0360925	-0.67	0.505	-.0947912	.0466887
1.black	-3.866887	2.519995	-1.53	0.125	-8.805986	1.072212
c.age#						
c.education	.0232092	.0105351	2.20	0.028	.0025608	.0438577
black#c.age						
1	.0905916	.0420067	2.16	0.031	.00826	.1729231
black#						
c.education						
1	.2766195	.1750993	1.58	0.114	-.0665689	.6198078
_cons	11.74866	6.392829	1.84	0.066	-.7810508	24.27838

```
. predict wbal2
```

```
. summarize age education black if treat==0 [iw=wbal2]
```

Variable	Obs	Weight	Mean	Std. dev.	Min	Max
age	253	185	25.81622	7.155019	18	55
education	253	185	10.34595	2.01065	0	17
black	253	185	.8432432	.3645579	0	1

```
. summarize age education black if treat==1 [iw=wbal2]
```

Variable	Obs	Weight	Mean	Std. dev.	Min	Max
age	185	185	25.81622	7.155019	17	48

Entropy balancing as an estimation command

```

education |      185      185   10.34595   2.01065      4      16
black     |      185      185    .8432432   .3645579      0      1
. corr age education black if treat==0 [aw=wbal2]
(sum of wgt is 185)
(obs=253)

```

	age	educat-n	black
age	1.0000		
education	-0.0080	1.0000	
black	0.0535	-0.0368	1.0000

```

. corr age education black if treat==1 [aw=wbal2]
(sum of wgt is 185)
(obs=185)

```

	age	educat-n	black
age	1.0000		
education	-0.0080	1.0000	
black	0.0535	-0.0368	1.0000

We see that means, standard deviations (and variances), as well as correlations (and covariances) have been perfectly balanced. Alternatively, option `targets()` can be used to generate the necessary terms automatically. `ebalfit` will then expand the variable list accordingly, taking account of the types of the variables (e.g., no terms for variances of categorical variables will be included as balancing the mean of a 0/1 variable also balances its variance):

```

. ebalfit age education 1.black, by(treat) targets(variance covariance) ///
> nolog vsquish
Entropy balancing                               Number of obs   =      438
                                                Wald chi2(8)       =      80.49
                                                Prob > chi2        =      0.0000
                                                Evaluator          =      bl
Main      = 0.treat (253 obs)                  Loss type          =      reldif
Reference = 1.treat (185 obs)                  Balancing loss     = 2.532e-15

```

	minimum	average	maximum	total	CV	DEFF
	.00023148	.7312253	13.072914	185	1.7253741	3.9769159

```


```

	Coefficient	Robust std. err.	z	P> z	[95% conf. interval]	
age	-.3454953	.1819311	-1.90	0.058	-.7020737	.0110831
c.age#c.age	-.0007966	.0017921	-0.44	0.657	-.004309	.0027159
education	-.6033715	.8552482	-0.71	0.481	-2.279627	1.072884
c.education#						
c.education	-.0240512	.0360925	-0.67	0.505	-.0947912	.0466887
1.black	-3.866887	2.519995	-1.53	0.125	-8.805986	1.072212
c.age#						
c.education	.0232092	.0105351	2.20	0.028	.0025608	.0438577
black#c.age						
1	.0905916	.0420067	2.16	0.031	.00826	.1729231
black#						
c.education						
1	.2766195	.1750993	1.58	0.114	-.0665689	.6198078
_cons	11.74866	6.392829	1.84	0.066	-.7810508	24.27838

				Evaluator	=	b1
				Loss type	=	reldif
Population size = 1,360,000				Balancing loss	=	1.547e-14
				balancing weights		
minimum	average	maximum	total	CV	DEFF	
587.87243	3105.0228	14976.219	1360000	.68869277	1.4742977	

	Coefficient	Robust std. err.	z	P> z	[95% conf. interval]	
age	-.0392513	.0058811	-6.67	0.000	-.0507781	-.0277246
education	-.1730964	.0264454	-6.55	0.000	-.2249285	-.1212643
black	-1.206872	.1398756	-8.63	0.000	-1.481023	-.9327208
hispanic	-.3027049	.2401386	-1.26	0.207	-.7733679	.167958
_cons	11.65529	.4453534	26.17	0.000	10.78241	12.52816

The target values do not necessarily need to be true values from a population. We can also use entropy balancing to construct as-if scenarios by setting the targets to theoretically interesting values, as long as the targets are not too far away from the center of the data such that no balancing solution exists.

4.5 Partial balancing

The `adjust()` option can be used to select the terms to be balanced, while keeping the means of the remaining variables fixed at their original values. Here is an example where the control group is reweighted such that the racial distribution is adjusted to the distribution in the treatment group, but the means of age and education are preserved:⁷

```
. ebalfit age education 1.black, by(treat) adjust(3) ///
> nolog noheader nowtable notable baltab
```

Balancing table

	Target value	Unbalanced value	std. dif.	Balanced value	std. dif.
age	36.09486	36.09486	0	36.09486	1.12e-14
education	10.7668	10.7668	0	10.7668	2.80e-14
1.black	.8432432	.3913043	-.9260241	.8432432	-6.85e-14

Such partial reweighting can be useful, for example, to study the “contributions” of individual covariates to an overall group difference in an outcome variable.⁸ In the following example we see that the group difference in 1978 earnings is reduced substantially if the racial distribution is adjusted while keeping age and education fixed. Additionally adjusting age and education only leads to a minor further decrease in the difference.

7. The same result could also be obtained by typing `noadjust(1 2)` instead of `adjust(3)`.

8. See Fortin et al. (2011) for an overview of counterfactual decomposition methods; see Jann (2008) for an implementation of the popular Oaxaca-Blinder decomposition in Stata.

```
. table () (treat), stat(mean re78) nottotal
```

	treat	
	0	1
Mean	9995.95	6349.144

```
. ebalfit age education 1.black, by(treat) generate(wbal) replace adjust(3)
(output omitted)
```

```
. table () (treat) [pw=wbal], stat(mean re78) nottotal
```

	treat	
	0	1
Mean	8160.198	6349.144

```
. ebalfit age education 1.black, by(treat) generate(wbal) replace
(output omitted)
```

```
. table () (treat) [pw=wbal], stat(mean re78) nottotal
```

	treat	
	0	1
Mean	8104.765	6349.144

5 Conclusions

Entropy balancing is a powerful alternative to other reweighting techniques such as inverse probability weighting based on logistic regression. In this paper I defined the model, derived influence functions for the parameters of the model, and illustrated how consistent standard errors can be obtained for statistics based on the reweighted data. I further presented software that implements the discussed methods.

The software provides a convenient tool for estimating balancing weights and generating influence functions, but adjusting statistical inference for reweighted statistics still requires a good understanding of the problem at hand and some programming skills on the side of the user. Based on the results presented in this paper, entropy balancing could be integrated into other estimation commands as a preprocessing device, such that reweighted estimates with consistent statistical inference would be readily available for a variety of applications without the need to write lengthy code. Some steps in this direction have already been taken. Command `dstat`, available from the SSC Archive, offers a `balance()` option that applies reweighting to a large collection of summary statistics (Jann 2020a). In fact, the treatment effect estimate in section 4.2 can be replicated by `dstat` in a single line of code:

```
. dstat re78 [pw=w0], over(treat, contrast) balance(eb:age-re75, reference(1))
Difference in mean          Number of obs    =      438
                          Contrast                    =      0.treat
```

```
Balancing:
      method =      eb
      reference =    1.treat
      controls = e(balance)
```

re78	Coefficient	Std. err.	t	P> t	[95% conf. interval]
1.treat	1830.641	906.1741	2.02	0.044	49.64029 3611.643

Similar support for `reweightig` is provided in `reldist`, a command for relative distribution analysis (Jann 2020c). Furthermore, the matching and reweighting command `kmatch` (Jann 2017) supports entropy balancing, albeit without specific adjustment of statistical inference. Since regression adjustment after entropy balancing yields consistent standard errors, however, command `kmatch` can still be used to obtain valid results. Here is a replication of the treatment effect from section 4.2 using `kmatch`. The trick is to include the full vector of covariates also in the outcome equation.

```
. kmatch eb treat age-re75 (re78=age-re75) [pw=w0], nomtable att
(fitting balancing weights ... done)

Entropy balancing
      Number of obs = 438
      Balance tolerance = .00001

Treatment   : treat = 1
Targets     : 1
Covariates  : age education black hispanic married nodegree re74 re75
RA equations: re78 = age education black hispanic married nodegree re74 re75 ...

Treatment-effects estimation
```

re78	Coefficient	Std. err.	t	P> t	[95% conf. interval]
ATT	1830.641	906.1741	2.02	0.044	49.64029 3611.643

Implementing entropy balancing as a reweighting and standardization option may also be useful for various other types of estimators, but such work still has to be done. Another topic that deserves further investigation is whether the efficiency of reweighted statistics based on entropy balancing can further be improved by optimized trimming techniques, and how this would affect statistical inference. See Kranker et al. (2020) for work that goes in this direction.

6 Appendix: Correction of IPW standard errors

For IPW we can adjust the standard errors of statistics based on the reweighted data in a similar way as for entropy balancing. In fact, working through the details (see Jann 2020c) shows that for IPW based on a logit model the correction has the exact same form as for entropy balancing. We simply need to replace the entropy balancing influence functions by the influence functions of the parameters of the logit model. A convenient way to compute influence functions for maximum likelihood models makes use of the scores and the model-based variance matrix (see Jann 2020b). The correction goes as follows:

```

. logit treat age-re75 [pw=w0]
  (output omitted)
. predict pscore if treat==0, pr
(185 missing values generated)
. generate ipw = w0 * cond(treat==0, pscore/(1-pscore), 1)
. predict score, score
. forvalues i=1/9 {
  2.   quietly generate IFipw`i' = .
  3. }
. mata:
----- mata (type end to exit) -----
:   X      = st_data(., "age-re75")
:   X      = X, J(rows(X), 1, 1) // add constant
:   score  = st_data(., "score")
:   Ginv   = st_matrix("e(V_modelbased)")
:   st_view(IF=., ., "IFipw*")
:   IF[.,.] = (X :* score) * Ginv`
: end
-----
. drop score
. drop IFipw9 // drop the IF for the constant; it is not needed
. summarize re78 if treat==0 [aw=ipw], meanonly
. generate IFy0 = (treat==0) * (re78 - r(mean)) / r(sum_w)
. summarize re78 if treat==1 [aw=ipw], meanonly
. generate IFy1 = (treat==1) * (re78 - r(mean)) / r(sum_w)
. mata:
----- mata (type end to exit) -----
:   // data
:   grp    = st_data(., "treat")
:   X      = st_data(., "age-re75")
:   IFy0   = st_data(., "IFy0")
:   IFipw  = st_data(., "IFipw*")
:   ipw    = st_data(., "ipw")
:   w0     = st_data(., "w0")
:   // compute (negative of) G
:   G = colsum(select(ipw :* IFy0 :* X, grp:==0))`
:   // adjust IF
:   st_store(., st_addvar("double", "IFy0c"), ipw ./ w0 :* IFy0 + IFipw * G)
: end
-----
. generate IFte = IFy1 - IFy0c
. total IFy0c IFy1 IFte [pw=w0]
Total estimation                               Number of obs = 438
-----

```

	Total	Std. err.	[95% conf. interval]	
IFy0c	-6.88e-08	943.3969	-1854.159	1854.159
IFy1	5.26e-13	567.8919	-1116.139	1116.139
IFte	6.88e-08	1071.73	-2106.385	2106.385

. drop IF*

The effect of the correction is less pronounced than for entropy balancing. This is related to the finding that conditioning on the estimated propensity score is more efficient than conditioning on the true propensity score, because random imbalance is partially removed. For entropy balancing this efficiency gain is stronger than for IPW because entropy balancing completely removes random imbalance.

7 References

- Dehejia, R. H., and S. Wahba. 1999. Causal Effects in Non-Experimental Studies: Reevaluating the Evaluation of Training Programs. *Journal of the American Statistical Association* 94(448): 1053–1062.
- Fortin, N., T. Lemieux, and S. Firpo. 2011. Decomposition Methods in Economics. In *Handbook of Labor Economics*, ed. O. Ashenfelter and D. Card, vol. 4, chap. 1, 1–102. Amsterdam: Elsevier. <http://ideas.repec.org/h/eee/labchp/4-01.html>.
- Hainmueller, J. 2012. Entropy Balancing for Causal Effects: A Multivariate Reweighting Method to Produce Balanced Samples in Observational Studies. *Political Analysis* 20(1): 25–46.
- Hainmueller, J., and Y. Xu. 2011. ebalance: Stata module to perform Entropy reweighting to create balanced samples. Statistical Software Components S457326. Available from <https://ideas.repec.org/c/boc/bocode/s457326.html>.
- . 2013. ebalance: A Stata Package for Entropy Balancing. *Journal of Statistical Software* 54(7): 1–18.
- Imai, K., and M. Ratkovic. 2014. Covariate balancing propensity score. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 76(1): 243–263.
- Imbens, G. W., and J. M. Wooldridge. 2009. Recent Developments in the Econometrics of Program Evaluation. *Journal of Economic Literature* 47(1): 5–86.
- Jann, B. 2005. moremata: Stata module (Mata) to provide various functions. Statistical Software Components S455001. Available from <http://ideas.repec.org/c/boc/bocode/s455001.html>.
- . 2008. The Blinder-Oaxaca decomposition for linear regression models. *The Stata Journal* 8(4): 453–479.
- . 2017. kmatch: Stata module for multivariate-distance and propensity-score matching, including entropy balancing, inverse probability weighting, (coarsened) exact matching, and regression adjustment. Statistical Software Components S458346. Available from <https://ideas.repec.org/c/boc/bocode/s458346.html>.

- . 2020a. DSTAT: Stata module to compute summary statistics and distribution functions including standard errors and optional covariate balancing. Statistical Software Components S458874,. Available from <https://ideas.repec.org/c/boc/bocode/s458874,.html>.
- . 2020b. Influence functions continued. A framework for estimating standard errors in reweighting, matching, and regression adjustment. University of Bern Social Sciences Working Papers 35. Available from <https://ideas.repec.org/p/bss/wpaper/35.html>.
- . 2020c. Relative distribution analysis in Stata. University of Bern Social Sciences Working Papers 37. Available from <https://ideas.repec.org/p/bss/wpaper/37.html>.
- Kish, L. 1965. *Survey Sampling*. New York: Wiley.
- Kranker, K. 2019. psweight: IPW- and CBPS-type propensity score reweighting, with various extensions. Statistical Software Components S458657. Available from <https://ideas.repec.org/c/boc/bocode/s458657.html>.
- Kranker, K., L. Blue, and L. Vollmer Forrow. 2020. Improving Effect Estimates by Limiting the Variability in Inverse Propensity Score Weights. *The American Statistician* .
- LaLonde, R. J. 1986. Evaluating the Econometric Evaluations of Training Programs with Experimental Data. *The American Economic Review* 76(4): 604–620.
- Valliant, R., J. A. Dever, and F. Kreuter. 2013. *Practical Tools for Designing and Weighting Survey Samples*. New York: Springer.
- Zhao, Q., and D. Percival. 2017. Entropy Balancing is Doubly Robust. *Journal of Causal Inference* 5: 2016–0010.