



## Late-glacial and early Holocene lake sediments, ground-water formation and climate in the Atacama Altiplano 22–24°S\*

Martin Grosjean<sup>1</sup>, Mebus A. Geyh<sup>2</sup>, Bruno Messerli<sup>3</sup> & Ueli Schotterer<sup>4</sup>

*Resource Management Science and Environmental Studies, University of British Columbia, 2206 East Mall, V6T 1Z3 Vancouver, Canada;* <sup>2</sup> *State Geological Survey Lower Saxony, 30655 Hannover, Germany;* <sup>3</sup> *Department of Geography, University of Bern, 3012 Bern, Switzerland;* <sup>4</sup> *Department of Environmental Physics, University of Bern, 3012 Bern, Switzerland*

Received 26 September 1994; accepted 20 February 1995

**Key words:** fossil groundwater, reservoir effect, seasonally-laminated sediments, subtropical Andes, paleomonsoon

### Abstract

Precipitation rates in the Atacama Altiplano 22–24°S were 400–500 mm yr<sup>-1</sup> during late glacial and early Holocene times as opposed to 200 mm yr<sup>-1</sup> today. This humid phase (Tauca phase) was likely due to strengthened tropical (monsoonal) circulation, which brought continental moisture to the Atacama Altiplano. The lake level of Laguna Lejía (23°30'S, 4350 m) at that time was up to 25 m higher than it is today. Mg/Ca and Sr/Ca data from lake sediments show that, what today is a highly saline lake was a freshwater lake at that time. Seasonally-laminated calcareous sediments were deposited between 13 500 and < 10 400 yr B.P. indicating the maximum of the humid phase. Climatic changes in the past are important for current groundwater resources. <sup>14</sup>C and <sup>3</sup>H data from lake-, ground- and well water suggest that modern groundwater formation (i.e. water < 40 years) in the Altiplano is very limited under current arid conditions. We conclude that significant amounts of the water resources in this area originated during the time of the late-glacial and early Holocene humid climate. Tritium data from snow samples show that the moisture in the Altiplano at 22–24°S is mainly of continental origin, whereas precipitation from the westerlies hardly contributes to the water supply in this area. This precipitation pattern matches the paleodata, and we suggest that current precipitation formation may provide an analogue framework for late-glacial circulation in this area.

### Introduction

Paleomonsoon on subtropical high mountain plateaus has become increasingly important in recent research on climate change. Most studies concentrate on the plateau of Tibet and on northwestern America (Rudiman & Kutzbach, 1991). Little is known about the central Andean plateau, the Altiplano. Although relatively small in size, the Altiplano may be important for the environmental history of the entire south American

continent and could, therefore, provide insight into the similarities and dissimilarities of past climates in both hemispheres.

The Atacama Altiplano is located between the modern tropical and extratropical rainfall belt. The current climate in this area is extremely dry with less than 200 mm yr<sup>-1</sup> precipitation. Therefore, this environment is highly sensitive to changes in humidity. Evidence of a more humid climate in this area during late-glacial and early Holocene times is found in lacustrine and glacial features, in the properties of paleosols, in the pollen spectra of peat bogs, and in the history of human habitation (Ochsenius, 1986; Nuñez, 1992; Baied & Wheeler, 1993; Messerli *et al.*, 1993; Grosjean, 1994; Grosjean & Nuñez, 1994).

\* This is the 3rd in a series of papers published in this special AMQUA issue. These papers were presented at the 1994 meeting of the American Quaternary Association held 19–22 June, 1994, at the University of Minnesota, Minneapolis, Minnesota, USA. Dr Linda C. K. Shane served as guest editor for these papers.




Fig. 1. Research area and map of Laguna Lejía showing the late-glacial maximum extent of the paleolake and the site of the sediment profile.

The lake level history of Laguna Lejía (23°30'S, 4350 m, Figs. 1 and 2) provides insight into long-term climatic conditions during late-glacial and early Holocene times (Grosjean, 1994). Based on the lake level history of this lake on a water and energy budget model, precipitation is estimated to have varied over the late Pleistocene/early Holocene as follows: less than 350 mm yr<sup>-1</sup> between c. 15 400 to c. 13 500 yr B.P., 450–500 mm yr<sup>-1</sup> between 13 500 to about 10 400 yr B.P., and about 400 mm yr<sup>-1</sup> during the early Holocene (Grosjean, 1994). The current precipitation rate is less than 200 mm yr<sup>-1</sup>. The lake surface increased during the maximum humid phase by a factor of six, and the lake level was 25 m higher than it is today (Fig. 1).

Markgraf (1989) and Kessler (1991) have concluded that strengthened monsoonal circulation accounted for the late-glacial humid phase in the Altiplano. *Alnus* and *Podocarpus* pollen in the late-glacial sediments of Laguna Lejía, and the lack of late-glacial high lake levels and glacial deposits further south between 25°S and 28°S provide evidence of strengthened monsoonal precipitation during late-glacial times as far south as 24°S (Messerli *et al.*, 1993; Grosjean, 1994).

However, little is known about short-term climatic fluctuations during this humid phase. Another question concerns the impact of this past humid climate on modern water resources: how renewable are the water resources in this dry area? Groundwater use is presently a subject of political controversy due to




Fig. 2. Stratigraphy of Laguna Lejía and the three major sediment units (southern shore, see also Fig. 3). The lake level history and the respective precipitation rates are modified after Grosjean (1994).

urbanisation, tourism, and rapidly increasing demands for copper.

The purpose of this article is to characterize the late-glacial and early Holocene climate based on lake sediments, and to discuss the significance of past humid conditions for present water resources in the Atacama Altiplano. The research area is shown in Fig. 1. The chemical and mineralogical composition of the laminated late-glacial lake deposits of Laguna Lejía is described in order to evaluate their potential as high-resolution records of former climatic conditions. The difficulty of <sup>14</sup>C dating of lake sediments due to the reservoir effect in a closed basin is illustrated by an example from sediments of the Laguna Lejía. <sup>3</sup>H tritium data on precipitation (snow), <sup>14</sup>C and <sup>3</sup>H data on groundwater, and surface water allow the modern pre-

cipitation component and its origin to be identified. The present-day recharge rate in the water cycle and/or old water components can be evaluated; this in turn provides insight into the linkage between the late-glacial lakes and current water resources in this area.

## Methods

Cores of lake sediment outcrops at the southern shore of Laguna Lejía were collected using PVC tubes. Powdered bulk samples were analyzed for mineralogy with X-ray diffraction (XRD). Bulk sediment samples were digested in 1% HCl, and Ca, Mg and Sr were analyzed using inductively coupled argon plasma atomic emission spectroscopy (ICAP-AES). SO<sub>4</sub><sup>2-</sup> was measured using a Lachat Autoanalyzer (turbidimetric method), and inorganic carbon with a Leco C Analyzer. Vertical thin-section were prepared using O'Sullivan's (1983) method. Mineralogy and chemical composition of individual laminae were determined by XRD and energy dispersive X-ray spectroscopy (SEM-EDX). The SEM-EDX analysis comprised the elements Na, Mg, Al, Si, S, Cl, K, Ca and Fe, giving 100 wt% (see sediment section 115–120 cm in Fig. 3). The results for Na, Cl, K, and Fe are not discussed in this study.

Chivas *et al.* (1986) used molar Mg/Ca, and Sr/Ca ratios in ostracode shells as indicators for paleotemperatures and paleosalinity. In order to obtain a high-resolution annual paleohydrological record, the Mg distribution in calcite within individual laminae was determined by measuring the d-spacings of the calcite lattice by XRD (Goldsmith & Graf, 1958). Among other trace elements, Mg and Sr may replace Ca in calcium carbonate lattice positions as well as in non-lattice sites (Tucker & Wright, 1990). The partitioning of Mg and Sr in the calcite lattice depends on the concentrations of Ca, Mg, and Sr in the host water, on kinetic effects such as precipitation rate, and probably on other factors (organic matter). Unlike Sr, partitioning of Mg is strongly influenced by temperature (Tucker & Wright, 1990: 299). Mg and Sr are useful as indicators of paleosalinity and paleotemperature because partitioning is not affected by redox state. However, in a natural system like Laguna Lejía it is most difficult to identify all controlling factors, and precipitation of gypsum is also likely to affect the Mg/Ca ratio in the host water. Therefore, changes in ratios of Ca, Mg, and Sr in magnesian calcite may be interpreted only qualitatively with respect to salinity and temperature changes.




Fig. 3. Sediment chemistry of the paleolake Laguna Leija. Mg/Ca and Sr/Ca ratios, CaCO<sub>3</sub> and SO<sub>4</sub><sup>2-</sup> content of the sediments in the core. The core shows the three major units: beginning (unit 1), maximum (unit 2) and end (unit 3) of the humid phase. The arrow shows an example of a dry period during the most humid phase with more saline conditions in the lake. The close-up section at a depth of 115–120 cm with the seasonally laminated sediments (e.g. 118.5–119.5 cm) shows a period of relatively stable environmental conditions during the maximum lake phase compared to the earlier deposits (e.g. 256–261 cm). Note: different methods (ICAP-AES, XRD and SEM-EDX) were used to determine the chemical compositions of the different sections (see 'Methods').

Macroscopic terrestrial organic matter (OM) is lacking in the sediments. Therefore, both the total OM and the inorganic carbonates were used separately for radiocarbon dating. Thermoluminescence dating of the polymineral fractions 4–12  $\mu\text{m}$  in volcanic ash layers within the lake sediments (Bürigi, 1992) was used to provide carbon-independent dates.

In order to estimate the reservoir effect, the dissolved inorganic carbon compounds (DIC) in groundwater, lake water and spring water were precipitated as  $\text{BaCO}_3$ , and analyzed for  $^{14}\text{C}$  (percent modern carbon pMC) and  $\delta^{13}\text{C}$  (‰ PDB). To minimize outgassing and fractionation of carbon as  $\text{CO}_2$ , the pH milieu was kept high ( $> 8$ ) during precipitation by addition of NaOH.

The modern component in the water cycle was identified by tritium measurements. Direct gas counting in a low-background underground laboratory allowed us to detect concentrations as low as 0.4 TU (tritium units).  $^3\text{H}$  originated in the atmosphere from nuclear weapon tests in the 1950s and is, therefore, a tracer for modern meteoric water, i.e. water younger than 40 years. Measurements of modern rainfall (IAEA/WMO, 1994) in Salta (NW Argentina) yielded more than 4 TU for tropical, continental moisture, whereas rainfall from the Pacific shows significantly lower tritium concentrations ( $< 2$  TU). Therefore, moisture advection from the continent can be clearly identified in precipitation samples with more than 2 TU.

## Results and discussion

### *Sedimentology, chemistry and mineralogy of Laguna Lejía*

Figures 2 and 3 show the sedimentology and selected chemical compounds of the paleolake Lejía.

The sediment section is divided into three major units: The sediments of the first unit (225–270 cm deep, *c.* 15 400 to *c.* 13 500  $^{14}\text{C}$  yr B.P. with reservoir correction) consist of a) a sublittoral sediment facies with laminated (1 mm) to banded ( $< 3$  cm) micrite with magnesian calcite, gypsum, traces of quartz and plagioclase, and diatoms, and b) a littoral facies with 1–3 cm thick sand layers with magnesian carbonate coatings of charophytae, and c) thin layers ( $< 0.5$  cm) of gypsum crystals. The relatively high Mg/Ca ratio in the HCl-soluble fraction of the sediments (Table 1), the strongly fluctuating calcite content, and frequent changes between littoral and sublittoral facies

suggest varying environmental conditions. The erosion rates on the adjacent slopes were considerable, and the environmental conditions in the lake and in the surrounding areas were unstable, although the lake was at a much higher level than today, and the climate was generally more humid. The late-glacial paleolake Lejía was lower in salinity than Laguna Miscanti (Table 1, Figs. 1 and 3), the freshest lake in this area today. Table 1 shows that modern sediments in Laguna Miscanti exhibit higher Mg/Ca ratios and higher Sr/Ca ratios than all of the sediments of paleolake Lejía. The aqueous electrical conductivity in the current water of Laguna Miscanti is 6.4  $\text{mS cm}^{-1}$  compared to 35  $\text{mS cm}^{-1}$  for Laguna Lejía today.

The sediments of the second unit (111–225 cm deep, *c.* 13 500 to  $< 10 400$   $^{14}\text{C}$  yr B.P. with reservoir correction) consist mainly of finely laminated calcareous sediments. They were deposited in a very calm environment corresponding to the maximum lake level, 25 m higher than today's level. Detritus is very sparse, consisting mainly of pyroclastic magnetite, feldspars, glasses, and a few lenses of fine sand. The 1.2 m thick, undisturbed and finely (0.5–2 mm) laminated carbonates have a lower Mg/Ca ratio than in unit one, and the Sr/Ca ratio is slowly decreasing (Table 1). Both suggest a trend towards fresher conditions, interpreted as being due to an increase in the lake level. Ripple marks and lateral slumping of sediments are very weak or absent, suggesting a calm, relatively deep-water environment (sublittoral to profundal facies). The lack of bioturbation suggests meromixis in the lake. As shown in the section between 118.5–119.5 cm (Fig. 3), the magnesian calcite layers, with occasional gypsum coprecipitation, alternate with deposits of diatoms and filamentous biogenic silica. These two phases are interpreted as seasonally-laminated calcareous sediments (O'Sullivan, 1983), with magnesian calcite precipitated during summer as the result of higher water temperatures and biologic activity, and bioclasts (mainly diatoms) accumulated during fall and winter due to the decay of plankton populations. Gypsum precipitation, as reflected by S content (SEM-EDX), is believed to be the result of the saturation in the water, and tends to follow calcite precipitation in the summer layer. However, the exact timing of the gypsum precipitation in the annual cycle is not yet known. Since gypsum precipitates at the end of the dry season when evaporation in the lake is greatest, this information could provide evidence for the timing of the rainy season, which is an important element of the paleoclimate. Clastic input, as represented by Al, correlates positively with Si from

*Table 1.* Mg/Ca and Sr/Ca ratios in calcites of the late-glacial/early Holocene sediments from lake Lejía (profile see Figs. 2 and 3) and the modern lake Miscanti. The low ratios in the sediments of Laguna Lejía suggest that the late-glacial paleolake Lejía was fresher than the modern lake Miscanti, the freshest lake in the Altiplano today (electrical conductivity  $6.4 \text{ mS cm}^{-1}$ )

	Material	Depth [cm]	1000 Mg/Ca average	1000 Sr/Ca average	
L. Lejía	calcite	0–110	$46 \pm 19$	$2.6 \pm 0.46$	$n = 35$
	calcite	111–223	$26 \pm 5$	$2.9 \pm 0.66$	$n = 23$
	calcite	225–270	$46 \pm 10$	$3.3 \pm 0.25$	$n = 11$
L. Miscanti	calcite	0–5	90	3.7	$n = 2$

diatom opal, and is characteristic for the fall/winter layer. The section that is between 115–120 cm deep (Fig. 3) gives an example of the sediments of unit 2 (111–225 cm). These 5 cm of laminated sediments are composed of 30–35 ‘annual’ cycles representing about three decades of lake history. The Mg and S contents suggest that relatively stable environmental conditions prevailed for a few decades. This is in contrast to the section that is 256–261 cm deep and represents the first sediment unit. However, fluctuations in the lake are observed for the period of laminated sediments: the high Mg/Ca and Sr/Ca ratio, and the high gypsum content at a depth of 190 cm (Fig. 3, arrows) suggest short-term saline conditions in the lake during a dry period.

The annual deposits are 0.5–2 mm thick. It follows that the deposition of the sediments at depths between 225–111 cm took more than 1000 years under high lake level conditions. This in turn suggests that the high lake levels in the Altiplano were the result of a shift in the precipitation belts, with a general increase in precipitation and reduction of evaporation due to greater cloudiness. Kessler (1991) and Grosjean (1994) concluded earlier that the high lake levels could not have been produced exclusively by single storms or melting glaciers. They argue that, under current climatic conditions, the paleolakes would have dried out within less than 50 years, which is in contrast to the 1000 years of high lake levels as determined by the laminated sediments.

The uppermost sediment unit (110–0 cm deep, < 10400 to *c.* 8500 (?)  $^{14}\text{C}$  yr B.P. with reservoir correction) consists of reworked, downwashed ‘older’ lake sediments. Oscillation ripples, alternating beds of carbonates and sand are abundant. The sandy interlayers sometimes show normal grading. All these fea-

tures provide evidence of fluctuating water levels. The mineralogy generally shows a mix of the underlying sediments, and the higher Mg/Ca ratios (Table 1) suggest more saline conditions in the lake due to higher evaporation rates and decreasing lake volume.

A mineralogically interesting feature was found at a sediment depth of 50 cm (Fig. 2): hydrated calcite needles ( $\text{CaCO}_3 \cdot 6\text{H}_2\text{O}$ , ikaite?) form a characteristic 10 cm thick layer. The monocline ikaite (?) needles, up to 6 cm long, are preserved as calcite pseudomorphs. This is typical for hydrated calcites because they are metastable mineral phases under current climatic conditions. The presence of ikaite is important since it indicates calcite precipitation at temperatures around  $0^\circ\text{C}$  (Tucker & Wright, 1990).

#### *The chronology of the Laguna Lejía sediments*

Discussion of the sediment chronology is purposely conservative, because determining the age of lake sediments in the extremely arid environment of the Altiplano is difficult. Therefore, the purpose of this section is to illustrate problems rather than to give a detailed chronology. The trouble is that no macroscopic terrestrial organic matter ( $\text{OM}_t$ ) could be found in the sediments. Therefore, we had to use both the aquatic  $\text{OM}_{aq}$  and the carbonate fraction of sediments separately for radiocarbon dating (Fig. 4). Today, all dissolved inorganic carbon DIC compounds in the lake water of Laguna Lejía are derived from modern, atmospheric  $\text{CO}_2$  (100 pMC), from  $\text{HCO}_3^-$  in groundwater inflow (e.g. 45.9 pMC, Hv-18084) of one sample, and dissolved carbonates from old lake sediments. The values of 35.3–36.1 pMC (Hv-17802, Hv-18083) seem to be representative for present lake water, which corresponds to 8200 conventional  $^{14}\text{C}$  years B.P. Since

the live aquatic plants found in the lake today contain 79.3 pMC (corresponding to 1900 conventional  $^{14}\text{C}$  years B.P., Hv-17880), they obviously assimilate a portion of the already aged carbon from the lake water. In conclusion, the radiocarbon ages of aquatic OM and precipitated carbonates in this lake are too large and have to be interpreted as maximum ages due to the discussed reservoir effect. This may be estimated by analysing modern lake water and live aquatic plants, where the correction values in Laguna Lejía are  $-2000$  years for  $\text{OM}_{\text{aq}}$  and  $-8000$  years for carbonates. Assuming a constant reservoir effect, the laminated sediments were deposited between 13 500 to  $\sim 10\,000$  yr B.P. However, since we are not yet able to give a carbon mass balance for the lake and the different sources, we do not know how good the assumption of a constant reservoir effect really is. Therefore, we conclude conservatively – at this stage of the work – a late-glacial, early Holocene age for the laminated sediments.

The similar  $^{14}\text{C}$  ages of the carbonate fractions in the lake sediments (Fig. 4, Grosjean, 1994) suggest post-sedimentary dissolution and reprecipitation of carbonates. This finding is not surprising, since magnesian calcites are metastable precipitates with high dislocation densities and compositional heterogeneities at the atomic level (Tucker & Wright, 1990: 301). SEM analysis showed that most of the calcite crystals are corroded. This observation suggests post-sedimentary dissolution, and this in turn puts limits on the interpretation of the Mg and Sr partitioning in the calcite lattice.

In light of these difficulties, it is necessary to compare the dating results from several basins using carbon-independent dating techniques, e.g. thermoluminescence dating of volcanic tuff layers in sediments (Bürgi, 1992). Reservoir effects seem to be less important in the hydrologically open system of the Salar de Aguas Calientes I (Fig. 5) and in the Salar de Ollagüe 21°S (Grosjean, 1994). Stromatolites corresponding to the maximum lake level, about 40 m above the present lake levels, yielded ages of  $13\,160 \pm 80$  yr B.P. (Beta-63356) and  $11\,300 \pm 70$  yr B.P. (Beta-63353), and laminated sediments were still deposited  $10\,400 \pm 75$  yr B.P. Since all of these ages are maximum ages due to an unknown reservoir effect, we conclude conservatively a late-glacial to early Holocene age for the maximum humid phase in the Altiplano.

### *Recharge of groundwater today*

Several ground- and surface waters were analyzed for their  $^3\text{H}$  and  $^{14}\text{C}$  content in order to estimate the proportions of old and young components in bodies of water in the Altiplano (Fig. 5). In order to interpret the radiocarbon data, different sources of carbon had to be evaluated (see also Fig. 4), including atmospheric  $\text{CO}_2$  of infiltrating rain water, biogenic  $\text{CO}_2$ , carbonate from weathered bedrock or dissolved lake sediments, soil  $\text{CO}_2$ , and volcanic-hydrothermal carbon.

The volcanic rocks in the study area are free of major carbonate sequences. Although chemical weathering is slow in this area, the alteration of abundant plagioclase may produce carbonates. In addition, volcanic  $\text{CO}_2$  may have serious implications for determination of groundwater age. The significance of this carbon source and of soil  $\text{CO}_2$  in the Atacama basin and the Pampa de Tamarugal is discussed in Fritz *et al.* (1979). In our study, chemical compositions help to identify the hydrothermal component in the water samples. Some typical results are listed in Table 2, and Fig. 5 shows the sample sites.

The  $^{14}\text{C}$  values of the water samples vary between 70.7 and 8.6 pMC. For instance, the high sulfate and chloride values of Quebrada Toro suggest a hydrothermal component in the water. This source adds a proportion of 'dead' C to the DIC reservoir, making the water appear older than it really is (i.e. age calculated starting with atmospheric values for  $^{14}\text{C}$ ). The sample from Salar Aguas Calientes I shows this effect to some extent, while the high mineralisation in the lake water of Laguna Lejía is due to concentration of ions in the brine. Although the surface water is in continuous carbon exchange with the 'modern' atmosphere, the level of  $^{14}\text{C}$  in open water bodies is low. We conclude that there is a significant contribution from an old water reservoir, which is recharged only on time scales of centuries to millenia. This situation seems to be typical for the Atacama area.

Tritium  $^3\text{H}$  was measured in order to evaluate the modern water component. Modern continental rainfall in south America at approximately 24°S can be expected to have concentrations  $> 4$  TU (tritium units), whereas rainfall from the Pacific without admixture of re-evaporated continental moisture is on the order of 2 TU (IAEA/WMO, 1994). All the analyzed snow and ice samples from different seasons (Table 3) contain tritium concentrations between 2.9 and 9.6 TU, suggesting that the moisture source is mainly continental. This finding is in agreement with Fuenzalida &








Fig. 5.  $^{14}\text{C}$  concentrations and  $\delta^{13}\text{C}$  data of water samples in the Atacama Altiplano. The data on water chemistry are listed in Table 2. \* data from Fritz *et al.*, 1979.

Rutllant (1986) and Aravena *et al.* (1989). Therefore, we conclude that water samples with concentrations below the detection limit  $< 0.4$  TU contain less than 10–20% of modern water, i.e. water which recharged within the last 40 years.

The tritium concentrations in the water samples follow the general precipitation pattern: values less than 1 TU or even below the detection limit are measured in groundwater and surface waters at lower elevations, generally more in the dry western part of the Andes down to the coast. At higher elevations and in smaller catchments situated in the Altiplano and to the more humid east, the modern component becomes increasingly important (Messerli *et al.*, 1993). The values for lakes, creeks and ponds in our research area in the Altiplano range between  $< 0.4$ – $< 2$  TU (Table 3). Worthy of note are the lakes which collect precipitation, and show tritium values up to 1.9 TU, which corresponds to  $< 50\%$  modern water. Springs and groundwater contain the lowest  $^3\text{H}$  concentrations of  $< 0.4$ – $< 1$  TU and, therefore, less than 10–20% modern water. The groundwater near Laguna Lejía is somewhat higher in tritium because it has a proportion of tritium from the lake.

Identification of the continent as the dominant moisture source is an important result in light of the atmospheric circulation during late-glacial times. Markgraf (1989), Kessler (1991), Messerli *et al.* (1993) and Grosjean (1994) postulate strengthened monsoonal circulation with moisture advection from the tropical continent to account for the humid climate in the Altiplano. Therefore, it is suggested that the moisture source likely was the same during late-glacial times as it is today, i.e. continental. Modern circulation patterns and precipitation processes (Fuenzalida & Rutllant, 1986; Aravena *et al.*, 1989) may, therefore, constitute a framework that is analogous to circulation during the late-glacial and early Holocene time in the Altiplano. This implies that the atmospheric flow pattern was the same as today, but heavily intensified, i.e. the modern precipitation-bearing circulation modes were statistically more abundant.

## Conclusions

Lake sediments from Laguna Legía in the Atacama Altiplano indicate climatic conditions that were significantly wetter during late-glacial and early Holocene times than they are today. Initially (*c.* 15 400–13 500 yr B.P.), the salinity of the lake was a little higher than dur-

ing the maximum lake phase, when freshwater conditions prevailed. During the main lake phase (*c.* 13 500– $< 10\,400$  yr B.P.), seasonally-laminated calcareous sediments were deposited in a 25 m deep lake. The environment was characterized by relative stability on a decadal to centennial time scale, as determined from the laminated sediments. However, some fluctuations in the environment can be observed. During the early Holocene, conditions in the lake were similar to those during the initial lake phase, still at a generally higher level than today.

The seasonally-laminated calcareous sediments suggest that the high lake levels were the result of a general shift of the climate to long-term-stable, more humid conditions. These high-lake-level conditions lasted for more than 1000 years, and were not the result of climatic peculiarities or melting glaciers.

The laminae represent seasonal deposits. Therefore, they provide a potential archive for high-resolution climatic information in an area where paleoenvironmental information is still scarce.

Terrestrial OM for radiocarbon dating was not found in these lake sediments. Therefore, aquatic OM or carbonates have to be used for  $^{14}\text{C}$  dating. Both materials are subject to the reservoir effect. For today's correction values in the Laguna Lejía we used –2000 yrs for aquatic OM and –8000 yrs for carbonates. A precise chronology of the major climatic events has yet to be fully established, and comparisons with other, C-independent dating techniques are needed. However, based on the data available today, we conclude a late-glacial to early Holocene age for the maximum humid phase.

We conclude from radiocarbon and tritium results that modern recharge of the main water bodies in the Atacama Altiplano is slow and limited. We suggest that the late-glacial/early Holocene humid climates, with more than twice as much precipitation as today accounted for much of the groundwater resources in this area today.

Tritium values of modern precipitation provide evidence of the continental origin of the water vapour. Since strengthened tropical rainfall is believed to have caused the humid phase in the Altiplano, it follows that studies of present-day circulation patterns and precipitation formation can be useful in reconstructing late-glacial atmospheric circulation in the Altiplano.

Table 3. Tritium values of snow, ice and firn samples, open water bodies, and spring and groundwater samples in the Atacama Altiplano. For sample sites, see Fig. 5 (\* data IAEA/WMO, 1994)

Date	Locality	Elevation material m	Tritium TU	$\pm 2\sigma$
snow, ice and firn samples				
	continental rainfall*		> 4	
	Pacific rainfall*		< 2	
20-Apr-90	Llullaillaco -2 cm	6200 snow -2 cm	5.27	$\pm 0.37$
20-Apr-90	Llullaillaco -50 cm	6200 ice -50 cm	9.60	$\pm 0.40$
20-Apr-90	Llullaillaco -300 cm	6200 firn -300 cm	6.37	$\pm 0.36$
20-Apr-90	Llullaillaco -320 cm	6200 firn -320 cm	6.49	$\pm 0.48$
2-Feb-93	Sairecabur	5800 fresh snow	5.33	$\pm 0.32$
20-May-90	Ojos del Salado	4500 fresh snow	6.22	$\pm 0.38$
16-Dec-88	Cerro Pili	5700 melt water	4.19	$\pm 0.51$
16-Dec-88	Cerro Pili	5700 snow -50 cm	4.85	$\pm 0.57$
16-Dec-88	Cerro Pili	5700 snow -25 cm	3.68	$\pm 0.44$
3-Apr-91	Cerro Toco N	4700 fresh snow	2.91	$\pm 0.29$
7-Apr-91	Cerro Tocopuri	5400 fresh snow	3.60	$\pm 0.33$
lakes, creeks and ponds				
10-Nov-91	Río Linzor	3980 creek	< 0.4	
19-Nov-91	Cerro Toco N	4580 pond	< 0.4	
3-Apr-91	Cerro Toco N	4640 creek	< 0.4	
10-Apr-91	Laguna Lejía	4300 lake	1.90	0.34
9-Dec-88	Laguna Lejía	4300 lake	0.94	0.38
8-Dec-88	Salar Aguas Calientes I	4100 salar water	0.73	0.46
3-Apr-91	Pena Blanca	4580 creek	1.37	0.32
11-Dec-88	Pena Blanca	4580 creek	1.69	0.49
springs and groundwater				
28-Mar-91	Talabre	3800 spring	< 0.4	
29-Mar-91	Laguna Lejía N, -170 cm	4300 groundwater	< 0.4	
20-Nov-91	Laguna Lejía N, -170 cm	4300 groundwater	< 0.4	
11-Apr-91	Laguna Lejía NW, -60 cm	4300 groundwater	< 0.4	
9-Dec-88	Laguna Lejía NW	4300 mix gw/lake	0.80	0.40
9-Dec-88	Laguna Lejía SW	4300 groundwater	1.78	0.35
9-Dec-88	Laguna Lejía SE	4300 groundwater	1.07	0.47
9-Dec-88	Laguna Lejía NE	4300 groundwater	0.76	0.49
3-Apr-91	Pena Blanca	4650 spring	< 0.4	
27-Nov-91	Toro Blanco (Hans)	4360 spring	< 0.4	
8-Dec-88	Salar Aguas Calientes I	4100 thermal spring	< 0.4	
8-Dec-88	Salar Aguas Calientes I	4100 spring 1, NE	0.70	0.46
8-Dec-88	Salar Aguas Calientes I	4100 spring 2, NE	0.89	0.38

### Acknowledgements

This study was financed with grants from the Swiss National Science Foundation (NF 20-36382.92) project 'Climate Change in the Arid Andes'. Les Lavkulich and Hans Schreier (Dept. of Soil Science,

UBC, Vancouver) generously provided the laboratory facilities for all the chemical and mineralogical analysis of the sediments. We thank Willi Egli and Marcela Espinoza N. (DIFROL) for logistical support and for granting permission to conduct research, and the two anonymous reviewers for constructive comments.

## References

- Aravena, R., H. Peña, A. Grilli, O. Suzuki & M. Mordeckai, 1989. Evolucion isotopica de las lluvias y origen de las masas de aire en el Altiplano chileno. IAEA-TECDOC-502: 129–142.
- Baied, C. A. & J. C. Wheeler, 1993. Evolution of high Andean ecosystems: Environment, climate, and cultural change over the last 12 000 years in the Central Andes. *Mountain Research and Development* 13: 145–156.
- Bürgi, A., 1992. Aufbau und Betrieb eines Thermolumineszenz-Labors zur Datierung quaternäreschichtlicher Proben. Ph.D. dissertation, University of Bern: 121 pp.
- Chivas, A. R., P. De Deckker & J. M. G. Shelly, 1986. Magnesium and strontium in non-marine ostracod shells as indicators of palaeosalinity and palaeotemperature. *Hydrobiologia* 143: 135–142.
- Fritz, P., C. H. Silva, O. Suzuki & E. Salati, 1979. Isotope Hydrology in Northern Chile. IAEA-SM-228/26: 525–543.
- Fuenzalida, H. & J. Rutllant, 1986. Estudio sobre el origen del vapor de agua que precipita en el invierno altiplanico. Informe final. Universidad de Chile: 51 pp.
- Goldsmith, J. R. & D. L. Graf, 1958. Relation between lattice constants and composition of the Ca-Mg carbonates. *Am. Mineralogist* 43: 84–101.
- Grosjean, M., 1994. Paleohydrology of the Laguna Lejía (Northchilean Altiplano) and climatic implications for late-glacial times. *Palaeogeography, Palaeoclimatology, Palaeoecology* 109: 89–100.
- Grosjean, M. & L. Nuñez, 1994. Late-glacial, Early and Middle Holocene Environments, Human Occupation and Resource Use in the Atacama (Northern Chile). *Geoarchaeology* 9: 271–286.
- IAEA/WMO, 1994. Network of Isotopes in Precipitation. NGDC/WDC-A Paleoenvironmental Data Sets. Boulder.
- Kessler, A., 1991. Zur Klimaentwicklung auf dem Altiplano seit dem letzten Pluvial. *Freiburger Geographische Hefte* 32: 141–148.
- Markgraf, V., 1989. Paleoclimates in Central and South America since 18 000 BP based on pollen and lake-level records. *Quat. Sci. Rev.* 8: 1–24.
- Messerli, B., M. Grosjean, G. Bonani, A. Bürgi, M. A. Geyh, K. Graf, K. Ramseyer, H. Romero, U. Schotterer, H. Schreier & M. Vuille, 1993. Climate Change and Natural Resource Dynamics of the Atacama Altiplano During the Last 18 000 Years: A Preliminary Synthesis. *Mountain Research and Development* 13: 117–127.
- Nuñez, L. A., 1992. Ocupacion arcaica en la Puna de Atacama: secuencia, movilidad y cambio. In B. J. Meggers (ed.), *Prehistoria Sudamericana. Nuevas Perspectivas*. Taraxacum, Washington: 283–307.
- Ruddiman, W. F. & J. E. Kutzbach, 1991. Plateaubildung und Klimaänderung. *Spektrum der Wissenschaften* 5/91: 114–125.
- Ochsenius, C., 1986. La glaciacion Puna durante el Wisconsin, deglaciacion y maximo lacustre en la tradicion Wisconsin-Holoceno y refugios de la megafauna postglaciales en la Puna y desierto de Atacama. *Revista de Geografia Norte Grande* 13: 29–58.
- O'Sullivan, P.E., 1983. Annually-laminated lake sediments and the study of quaternary environmental changes – a review. *Quat. Sci. Rev.* 2: 245–311.
- Tucker, M. E. & V. P. Wright, 1990. *Carbonate Sedimentology*. Blackwell Scientific Publications, Oxford: 481 pp.