

OPEN

## Demographic Changes in Pelvic Fracture Patterns at a Swiss Academic Trauma Center from 2007 to 2017

Karin Villiger<sup>1</sup>, Malin Kristin Meier<sup>2</sup>, Rebecca Maria Hasler<sup>3</sup>, Johannes Dominik Bastian<sup>2</sup>,  
Moritz Tannast<sup>4</sup>, Aristomenis Konstantinos Exadaktylos<sup>5</sup>, Simon Damian Steppacher<sup>6</sup>

<sup>1</sup>Department of Anesthesiology and Pain Therapy, Inselspital, Bern University Hospital,  
University of Bern, Bern, Switzerland

<sup>2</sup>Department of Orthopaedic Surgery and Traumatology, Inselspital, Bern University Hospital,  
Bern, Switzerland

<sup>3</sup>Department of Traumatology, Zürich University Hospital, Zürich, Switzerland

<sup>4</sup>Department of Orthopaedic Surgery and Traumatology, Fribourg Cantonal Hospital, University  
of Fribourg, Fribourg, Switzerland

<sup>5</sup>Department of Emergency Medicine, Inselspital, Bern University Hospital, Bern, Switzerland

<sup>6</sup>Department of Orthopaedic Surgery and Traumatology, Inselspital, Bern University Hospital,  
Freiburgstrasse 8, 3010 Bern, Switzerland

## **Corresponding author**

Simon Damian Steppacher, MD, Priv. Doz.

Department of Orthopaedic Surgery and Traumatology, Inselspital, Bern University Hospital,  
Freiburgstrasse 8, 3010 Bern, Switzerland

Tel. +41-31-632-2222

Fax +41-31-632-3600

Email: Simon.Steppacher@insel.ch

## **Conflict of Interest and Source of funding:**

The authors declare that they have no competing interests. No funding was received.

No funding was received from the National Institutes of Health, Welcome Trust or the Howard Hughes Medical Institute.

The results of the underlying manuscript have not been presented at any meetings or conferences.

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

*Background:* Increasing life expectancy has led to higher incidence of fragility fractures of the pelvis. These demographic changes may have a direct impact on fracture patterns. The goal was (1) to evaluate demographical trends in patients with pelvic ring injuries at a tertiary Swiss trauma center and (2) to analyze the influence on fracture patterns.

*Methods:* We performed a retrospective cross-sectional study including 958 patients (mean age  $57 \pm 21$  years, 48% women) with a pelvic ring injury between 2007 and 2017. Fractures were classified according to Tile, Young and Burgess or Rommens and Hofmann (fragility fractures) using conventional and CT imaging. Low-energy fractures were defined as fractures resulting from fall from standing height or less. Fracture classifications, age, gender, injury severity score (ISS) and trauma mechanism were compared using analysis of variance (ANOVA) or chi-square test. Cluster analysis was performed to identify groups with similarities in fracture patterns and demographic parameters.

*Results:* From 2007 to 2017, the frequency of pelvic ring injuries increased by 115% (increase per decade), mean age increased by 15% ( $p=0.031$ ). A trimodal age distribution was found; highest increase for fractures occurred in the 'older' (265%) patient group. Low-energy fracture was the most common trauma mechanism (43% of all fractures, an increase of 249%). Changes in fracture pattern showed a disproportioned increase of 'lateral compression (LC)' fractures (LC type 1 in 64%) or partially stable fracture (B2: with 39%). In patient aged over 65 years, the strongest increase was found for 'non-displaced posterior' fractures with an overall prevalence of 62%. Five clusters were found with the most frequent cluster representing 'older female patients with low-energy fracture (LC, Tile type B)' in 30%.

*Conclusion:* The current results corroborate the trend of increasing frequency of fragility fractures in an aging society. The demographic shift has a direct impact on fracture pattern with a disproportionate increase in partially stable compression fracture of the pelvis.

*Level of Evidence:* Level III

*Study type:* Retrospective cross sectional study

**Keywords:**

Pelvic ring injury, fragility fracture, orthogeriatrics, trauma, epidemiological trends

ACCEPTED

## **Background**

Increasing life expectancy has led to a higher incidence of fragility fractures of the pelvis in the elderly population. (1, 2) These demographic changes may have a direct impact on fracture patterns observed in pelvic ring injuries. Previous studies reported on demographic trends in patients with pelvic ring injuries, however, results were based on selected patient groups (geriatric patients(2–4) or high-energy trauma (5)) or did not include additional evaluation of fracture patterns (1, 3, 5). There is a lack of comprehensive reports of demographic trends and associated pelvic fracture patterns with a detailed and year-by-year evaluation. In addition, results of demographic trends and fracture patterns for the recently introduced classification system for pelvic fragility fractures are sparse. (1–4, 6–8)

In the past, we experienced an increase in the frequency of pelvic fragility fractures assigned to the Swiss tertiary trauma center at the author's institution. We hypothesized that changes in demographics such as age, gender distribution or trauma mechanism have a direct impact on the pelvic fracture patterns. Therefore, the objectives of the current study were (1) to evaluate demographical trends of patients with pelvic ring injuries (frequency, age, gender and trauma mechanism) over a study period from 2007 to 2017 at a tertiary Swiss trauma center and (2) to analyze the association between demographic changes and pelvic fracture patterns (classification of Tile (9), Young and Burgess (10) and fragility fractures of the pelvis classification according to Rommens and Hofmann (8)).

## Methods

### *Patient selection*

After institutional review board approval, we performed a retrospective cross-sectional study evaluating demographic trends and fracture patterns in patients with pelvic ring injuries assigned to the trauma center of the Inselspital, Bern University Hospital, between 2007 and 2017. It is a tertiary trauma center with 45,000 emergency consultations per year in an area of about 2 million inhabitants.<sup>(11)</sup> Patients aged less than 16 years were not included in the current study since they are referred to the pediatric emergency department in the hospital. Two different clinical reporting systems of the emergency department were used during the study period with cases recorded in Qualicare (Qualicare AG, Trimbach, Switzerland) up to May 2012 and subsequently in E-Care (E-Care BVBA, Turnhout, Belgium). Inclusion criterion was a documented pelvic fracture in the clinical reporting systems with a total of 1051 patients from February 2007 to December 2017. Exclusion criteria were missing or incomplete imaging of the pelvis (n=52 patients; 4.9%), isolated acetabulum fracture rather than the documented pelvic ring fracture (n=34 patients; 3.2%) and a pathological fracture of the pelvis due to a tumor (n= 7 patients; 0.7%). This left a total of 958 patients with pelvic ring injuries for evaluation (Figure 1). In 887 patients (93%) imaging included conventional radiographs and computer tomography (CT) of the pelvis and in 71 patients (7%) conventional imaging only existed. Additional data from the Trauma Audit and Research Network (TARN) were available for 556 out of the 892 patients (62%) from April 2009 to December 2017. Of those 556 patients the mean injury severity score (ISS) was  $27 \pm 15$  (4 – 75), the 30-day mortality rate was 6% (34 out 556 patients) and 309 patients (56%) were assigned to the resuscitation room at admission.

### *Fracture classification*

The fracture pattern of pelvic ring injuries were classified using the conventional images and CT scans according to the classification systems of Tile (9) or Young and Burgess (10). In patients aged over 65 years, fractures were additionally classified according to Rommens and Hofmann (8) for fragility fractures of the pelvis. The classifications were performed by two of the authors; one (KV) being a resident specifically trained in classifying pelvic ring injuries and one (SDS) being an experienced orthopaedic and trauma surgeon reviewing the cases. The classification systems have previously been evaluated for inter- and intraobserver reliability and showed a moderate to substantial agreement. (12, 13) However, differences in agreement were reported among observers with a different degree of experience regarding the treatment of pelvic trauma. (12, 13) For specialized pelvis surgeons, a level of agreement with a Kappa of 0.52 / 0.50 (interobserver / intraobserver agreement) for the Tile (9) classification, 0.60 / 0.55 for the Young and Burgess (10) classification and 0.47 / 0.49 for the Rommens and Hofmann (8) for fragility fractures of the pelvis has been reported.(12, 13)

### *Fracture classification according to Tile*

Classification according to Tile (9) was performed depending on the stability of the posterior arch of the pelvis including 'stable' (type A), 'partially stable' (type B) 'unstable' (type C). Each fracture type included three subtypes resulting in a total of nine fracture types according to Tile (9) (Supplemental Table 1, <http://links.lww.com/TA/C137>). The Tile classification (9) could be applied to 944 out of 958 pelvic injuries (99%; Supplemental Table 1, <http://links.lww.com/TA/C137>) with exclusions of 14 patients (1%) with isolated spinopelvic dissociation with an intact anterior arch of the pelvis.

### *Fracture classification according to Young and Burgess*

Classifications according to Young and Burgess (10) were performed according to the fractures mechanism including antero-posterior compression (APC; three subtypes), lateral compression (LC; three subtypes) and vertical shear (VS) injuries with a total seven subtypes. The Young and Burgess classification (10) could be applied in 772 fractures (81%) with exclusion of 172 fractures with an intact posterior arch of the pelvis (Tile type A; 18%) and 14 isolated spinopelvic dissociation (1%). Ambiguities with the classification systems were handled as follows: In cases with bilateral fractures (e.g. B3 or C3 according to Tile(9)), the more severe pelvic fracture was considered for classification according to Young and Burgess (10). Since symphyseal widening is reduced on images with a pelvic binder, the distinction between Young and Burgess (10) AP1 (<2.5cm widening) and AP2 (>2.5cm widening) was performed according to lesion of the posterior arch: a visible opening of the anterior part of the sacroiliac joint the fractures were classified as AP2, without widening as AP1. APC fractures with symphyseal rupture and a fracture of the posterior ileum were classified as AP3 according to Young and Burgess (10) (defined as symphyses widening with a fracture running through the sacroiliac joint). Isolated sacral compression fractures without an anterior ring fracture (as in fragility fractures type II according Rommens and Hofmann (8)) were classified as B2 according to Tile (9) or as LC 1 according to Young and Burgess (10).

### *Fracture classification according to Rommens and Hofmann*

Of the 365 patients aged over 65 years, 320 fractures (88%) were classified according to Rommens and Hofmann (8) for fragility fractures of the pelvis. Fractures were classified as anterior injury only (Type I), non-displaced posterior injury (Type II), displaced unilateral



posterior injury (Type III) or displaced bilateral posterior injury (Type IV). Exclusions included 27 patients with avulsion fractures or isolated iliac wing fractures (Tile (9) A 1/2) and 18 pelvic injuries with a symphyseal rupture (Young and Burgess (10) APC 1/2, Tile (9) B 1/3).

### *Outcome measures*

(1) For the first study objective, outcome measures to describe demographic trends included frequency, patient age at the time of pelvic fracture, gender and mechanism of trauma. Mechanism of trauma included 'low-energy fracture' with fall from standing height or less, 'motor vehicle accident (MVA)' including accidents of pedestrians, 'fall from height', 'sport accident' and 'crush injury'. The patients were separated into age groups using cluster analysis (see 'statistical analysis'). An increase / decrease was quantified with the correlation coefficient of the linear correlation model. In addition, the percentage of increase / decrease over the study period from 2007 to 2017 was calculated based on the linear regression model. (2) For the second study objective, outcome measures to describe fracture patterns included the previously described classification systems of Tile (9), Young and Burgess (10) or Rommens and Hofmann (8). The frequency of each fracture type was compared for each year over the study period of 2007 to 2017. Increase / decreased was also quantified using the correlation coefficient and percentage increase / decrease over 10 years.

### *Statistical analysis*

We tested normal distribution of continuous data (age, ISS) with the Kolmogorov-Smirnov test and since all continuous data were normally distributed, parametric tests were used. To compare continuous data over the study period, among age groups and fracture patterns the analysis of

variance (ANOVA) was performed. If significant differences existed, pairwise comparison was performed with the independent and two-tailed t-test with Bonferroni-correction. Binominal data were compared using the chi-square test. To quantify an increase / decrease in frequency over the study period a linear regression analysis was performed and the corresponding regression coefficient was calculated. The increase in frequency between 2007 and 2017 was calculated using the linear regression model. Increase / decrease in frequency of pelvic fractures were compared by comparing the slope of the linear regression among subgroups of fracture classifications, mechanisms of trauma or gender using the analysis of covariance (ANCOVA). Cluster analysis was performed for continuous parameters (age groups) using the Ward's method for hierarchical cluster analysis and for nominal data (fracture patterns) with the two-step cluster analysis. Statistical analysis was performed using WinStat® (Robert K. Fitch Software, Bad Krozingen, Germany) and IBM® SPSS® Statistics (Version 25, Armonk, New York, USA).

## **Results**

### *Patient characteristics*

Of the 958 included patients, a majority of 677 patients (71%) were primary referrals and 281 patients (29%) were referred from other hospitals. Overall, gender was equally distributed with 497 male patients (52%) and 461 female patients (48%; Table 1). The mean age was  $57 \pm 21$  years (range, 16 – 98 years) and 365 patients (38%) were aged over 65 years. A pelvic binder was present on admission in 269 patients (28%).

### *Demographic trends for frequency, age and gender of patients with pelvic injuries*

Over the study period from 2007 to 2017, the frequency for pelvic ring injuries increased by 157% (increase per decade; Figure 2A). No difference in frequency existed between male (regression coefficient of 2.99) and female patients (regression coefficient of 4.63;  $p = 0.347$ ; Figure 2B). The average age increased by 15% ( $p = 0.031$ ). Using cluster analysis, a trimodal age distribution was found for pelvic ring injuries (Figure 3A): the ‘young patients group’ (187 patients) had a mean age of  $25 \pm 6$  (16 – 38) years, the ‘middle-aged patient group’ (305 patients) had a mean age of  $49 \pm 6$  (36 – 62) years and the ‘older patient group’ (466 patients) had a mean age of  $74 \pm 10$  (58 – 98) years. In all age groups the frequency of pelvic fractures increased from 2007 to 2017 (Figure 3B), however, the increase in frequency among age groups differed ( $p = 0.008$ ) with the largest increase in the ‘older age group’ (regression coefficient of 4.76, increase of 265%). The percentage of female patients was higher in the ‘older age group’ (61% female patients) compared to the ‘middle-aged group’ (38% female patients;  $p = 0.001$ ) or the ‘younger age group’ (34% female patients;  $p = 0.001$ ; Table 1).

### *Changes in trauma mechanism for pelvic ring injuries*

Trauma mechanism for pelvic ring injuries changed over time (Table 1; Figure 4;  $p < 0.001$ ): the largest increase was found for low-energy fractures (regression coefficient of 3.55; increase of 249%), followed by fractures due to MVA, fall from height, sports accidents and crush injuries (Figure 4). Overall, ‘low-energy fractures’ accounted for 43%, MVA for 25%, fall from height for 13%, sport injuries for 10%, crush injuries for 5% and others for 4% of all pelvic ring fractures (Figure 4).

### *Changes in fracture patterns according to Tile (9)*

The frequency of all subtypes according to Tile (9) increased with a regression coefficient of 3.80, 2.07 and 1.47 for type B, A and C fractures, respectively (Figure 5A;  $p = 0.239$ ). This corresponds to an increase of 140%, 393% and 88% for type B, A and C fractures, respectively (Figure 5A). Overall, the most common injury type was the 'B2: lateral compression injury' with 39% (Supplemental Table 1, <http://links.lww.com/TA/C137>; Figure 5B). In the 'older patient group' the prevalence of type C fractures was decreased (22%;  $p < 0.001$ ) and the prevalence of type B fractures was increased (58%;  $p = 0.014$ ) compared to the other two age groups (Supplement Figure 1). The percentage of female patients was higher in fractures type A (51%) or B (53%) compared to type C (37%,  $p = 0.004$  and  $p < 0.001$ , respectively). Low-energy fractures were the most common mechanism for fractures for all three subgroups according to Tile (9) with a frequency of 55% for type A, 44% for type B and 31% for type C ( $p < 0.001$ ; Table 1). Fall from height was more common in fractures type C (20%) than type A (13%;  $p < 0.001$ ) or B (9%,  $p < 0.001$ ; Table 1).

### *Changes in fracture patterns according to Young and Burgess (10)*

LC fractures showed highest increase (regression coefficient of 4.48; increase of 158%; Figure 6A) compared to the other subtypes ( $p = 0.003$ ). No or minimal increase was found for APC (regression coefficient of 0.03; increase of 3%) or VS injuries (regression coefficient of 0.76; increase of 187%);  $p = 0.142$ ; Figure 6A). Overall, the most common pelvic ring injury was the LC injury type 1 with 64% (Figure 6B) (Supplemental Table 1, <http://links.lww.com/TA/C137>; Figure 6B). The mean age of LC fractures was  $58 \pm 21$  (16 – 98) and higher than in APC ( $53 \pm 15$  [17 – 84];  $p = 0.006$ ) or VS injuries ( $49 \pm 18$  [19 – 88];  $p < 0.001$ ; Table 1). In the 'older

patient group' the prevalence of LC fractures was increased (81%;  $p < 0.001$ ) and the prevalence of APC (12%;  $p < 0.001$ ) and VS (7%;  $p = 0.002$ ) injuries were decreased compared to the other two age groups (Supplemental Figure 1, <http://links.lww.com/TA/C139>). The percentage of female patients was higher in LC fractures (58%) than in APC (13%;  $p < 0.001$ ) or VS injuries (32%;  $p < 0.001$ ). Low-energy fractures were the most common reason for LC fractures (48% vs 15% in APC [ $p < 0.001$ ] or 23% in VS [ $p < 0.001$ ]). Fall from height was more common in VS fractures (28%) compared to APC (14%;  $p = 0.016$ ) or LC fractures (10%;  $p < 0.001$ ; Table 1).

#### *Changes in fragility fracture patterns according to Rommens and Hofmann (8)*

In patients aged over 65 years, a distinct increase in the frequency for 'non-displaced posterior injuries' could be found (regression coefficient of 2.48; increase of 453%;  $p < 0.001$ ; Figure 7). The other three subgroups showed a marginal increase in frequency over the study period (regression coefficient ranging from 0.15 to 0.45; Figure 7). Overall, the prevalence of 'non-displaced posterior injuries' was 62% (Figure 7).

#### *Cluster analysis of demographic characteristics and fracture patterns (evaluated for the entire patient sample)*

Using cluster analysis, we found five clusters for each the Tile (9) or Young and Burgess (10) classification (Table 2). The most prevalent cluster included predominantly female and older patients with type B (9) or LC (10) pelvic ring fractures (prevalence of 30% and 27%, respectively) due to low-energy fractures (Table 2). In younger and predominantly male patients, we found a prevalence of type C (9) or VS and APC injuries (10) pelvic ring fractures due to high-energy trauma (prevalence of 25% and 18%; Table 2). The third most prevalent cluster

included the middle-aged group with type B/C (9) or LC and APC injuries (10) due to high-energy trauma (prevalence of 12% to 23%; Table 2)

## **Discussion**

Due to the increasing life expectancy, osteoporotic fractures are on the rise. (14) For pelvic ring injuries an increased frequency has been reported for the older patient population. (3, 4) This is in accordance with the results from the current study from a Swiss tertiary trauma center. The strongest increase was found for the ‘older patients’ group. We found distinct patterns of pelvic ring injuries (Table 2): the most prevalent cluster comprised older and mostly female patients with a compression fracture due to low-energy fracture. The younger patient group included a higher percentage of instable, VS or APC injuries due to high-energy trauma (Supplemental Figure 1, <http://links.lww.com/TA/C139>). In contrast to classic publications reporting a bimodal age distribution for pelvic ring injuries (6, 7), we found a trimodal age distribution (Figure 3A). The middle-aged patients group comprised (Figure 3) both LC and APC injuries due to sports or motor vehicle accidents (Table 2).

We found an increasing frequency of pelvic fractures over the study period from 2007 to 2017 (Figure 2). This is in accordance with other reports (1–4) (Table 3): an increase of 26% in 27 years (1990 - 2007) has been reported for patients of any age (Table 3). (1) For patients aged over 65 years, an increase of 24% (2) or 37% (4) has been shown after an observation period of 17 and 25 years, respectively (study periods from 1980/90s to 2010s; Table 3). In patients from Finland aged over 80 years, an increase of 499% for an observation period of 43 years (1970 to 2013) was reported. (3) In the current study, an increase of 157% in 10 years (Figure 2) was

found which is comparable to the study from Finland (3) but higher than most reported rates (1, 2, 4) (Table 3). This could be due to the fact that osteopenic pelvis fractures are steeply rising in aging societies (3) and that the current results are from a more actual patient series starting in 2007 than in the other studies (starting between 1970s and 90s; Table 3). In the study period, the population in the closer area of the hospital (canton of Bern) rose by 7% and in the entire country by 12% (Data from 'Federal Statistical Office of Switzerland', <https://www.bfs.admin.ch/bfs/en/home/statistics/population.html>, accessed 16 November 2020; Supplemental Table 2, <http://links.lww.com/TA/C138>). Therefore, an increase of 155% (Figure 2) for pelvic injuries cannot be explained by the population increase alone. During the study period, the population aged over 65 years increased by 21% and 25% for the closer area and Switzerland, respectively (Data from 'Federal Statistical Office of Switzerland', <https://www.bfs.admin.ch/bfs/en/home/statistics/population.html>, accessed 16 November 2020; Supplemental Table 2, <http://links.lww.com/TA/C138>). This disproportional growth of the elderly population could be an explanation for the steep increase of pelvic injuries, especially low-energy and osteopenic fractures (Figure 6). Accordingly, the mean age of the patient series in the current study increased over the study period by 15% to  $55 \pm 20$  (16 – 89) years in 2017 (Figure 2A). In contrast to previous publications reporting a bimodal age distribution for pelvic ring injuries (6, 7), we found a trimodal age distribution (Figure 3). In addition, a female predominance has been reported (3, 4, 17), which we could not find with an overall prevalence of 48% female patients (Table 1). However, we found distinctive groups of pelvic ring injuries using the cluster analysis (Table 2): in older and predominantly female patients, we found LC (10) or partially stable (9) pelvic ring fractures due to low-energy fractures (Table 2; Supplemental Figure 1, <http://links.lww.com/TA/C139>). In younger and predominantly male

patients, we found a high prevalence of VS and APC injuries (10) or unstable (9) pelvic ring fractures due to high-energy trauma (Table 2; Supplemental Figure 1, <http://links.lww.com/TA/C139>). This bimodal distribution has been previously reported. (6, 7, 17) In addition, we found a relatively large middle-aged group (Figure 3) with a prevalence of 32% (Table 1) with LC and APC injuries (10) and (partially) unstable fractures (9) due to high energy trauma (Table 2; Supplemental Figure 1, <http://links.lww.com/TA/C139>).

The demographic changes have a direct influence on the fracture pattern of pelvic ring injuries (Figures 5, 6, 7) and are associated with different trauma mechanism (Figure 4). Both low-energy fractures and MVA (including pedestrian accidents) showed a strong increase over the study period (Figure 4). We found a lower percentage of high-energy trauma of 53% compared to literature ranging from 68% to 87% (Table 3). (15, 16, 19) Consequently, we had a higher percentage of low-energy trauma of 43% compared to literature (Table 3). According to the increasing number of older patients with osteopenic fractures (Figure 3B), we found an increasing number of LC fractures (9, 10) (Figure 6). Among the patients aged over 65 years, we found a substantial increase in the fractures with non-displaced posterior pelvic fractures (8) accounting for an overall 62% of all fragility fractures (Figure 7). According to the classification of Tile (9), the most prevalent fracture was the type B (partially stable posterior fracture) with 54% followed by type C (unstable posterior fracture) with 28% and type A (stable posterior fracture) with 18% (Table 1 and Figure 5). This in contrast to most results in literature (Table 3) reporting the highest prevalence for type A fractures followed by B and C. (6, 7, 15–17) The more prevalent type B and C fractures in the current study could also be due to a selection bias since more severe fractures potentially needing surgical treatment are usually referred to a



tertiary trauma center (see also limitations). Comparable results with a higher prevalence of type B and C fractures has been reported from another referral center for pelvic injuries in the Netherlands. (19)

This study has several limitations. First, there is a selection bias with 29% of the patients referred to the tertiary trauma center from smaller hospitals. These patients potentially had a more severe pelvic injury. This might be a reason why unstable fractures (type C according to Tile (9)) were more frequent than stable fractures (type A; Figure 5), which is in contrast to most publications (6, 7, 15–17) (Table 3) reporting a higher frequency for type A than C fractures. Despite this selection bias, the group with partially stable fractures (type B according to Tile (9)) and mainly conservative treatment (8) showed the largest increase over the study period. A second limitation is the reliability of the classification systems for pelvic ring injuries, which sometimes only showed a moderate level of reliability, especially for less experienced observers. (12, 13) In addition, CT imaging was missing in 7% of the patients which could have further decrease reliability of classification. (18) Also, some pelvic fractures cannot be classified without any doubts according to Tile (9) or Young and Burgess (10) (see methods section). A strength of the study is that all fractures were classified by the same observers (KV, SDS) and no classifications from medical records were used. Therefore, comparison among the years over the study period or subgroups is not susceptible for interobserver disagreement.

The current study shows in a year-by-year fashion the demographic shift in patients with pelvic ring injuries from 2007 to 2017. Low-energy fractures in elderly patients are on the rise, more than in any other patient subgroup (Figure 3B). In contrast to classic reports on pelvic ring

injuries from the 90s, we found a trimodal age distribution (Figure 3A): most prevalent are low-energy fractures in the elderly patients followed by high-energy trauma in younger patients (Table 2). In addition, we found a third subgroup with middle-aged patients with fractures due to mainly high-energy trauma. A strength of the study is the detailed and year-by-year evaluation of the fracture pattern spanning a decade. The demographic shift could be shown to have a direct impact on fracture pattern with a disproportionate increase in partially stable compression fracture of the pelvis. The current results corroborate the trend of increasing frequency of fragility fractures in an aging society, representing a major public health burden. The descriptive radiographic results serve as the basis for future interventional studies.

### **Supplemental digital content**

Supplemental digital content includes detailed information regarding frequency of subtypes of pelvic fractures classified according to Tile (9), Young and Burgess (10) and the fragility fracture of the pelvis (FFP) classification according to Rommens and Hofmann (8) in ‘Supplemental Table 1, <http://links.lww.com/TA/C137>’. In the ‘Supplemental Table 2, <http://links.lww.com/TA/C138>’ detailed information about changes in gender distribution or age in the reference population in the hospital area over the same period from 2007 to 2017 is summarized. The ‘Supplemental Figure 1, <http://links.lww.com/TA/C139>’ compares the frequencies of the classification of Tile (9) or Young and Burgess (10) in relation to the three age groups in a graphical way.

**Ethics approval**

The study protocol was approved by the local ethics committee (Approval no. KEK-BE 2016-00625).

**Conflict of interest and source of funding:**

The authors declare that they have no competing interests. No funding was received.

**Authors' contributions**

All authors have significantly contributed to either concept and study design, ethical approval, data acquisition, imaging analysis, interpretation of data and statistics, creation of figures, writing of initial draft, manuscript editing or critical review.

**Acknowledgements**

None.

## References

1. Buller LT, Best MJ, Quinnan SM. A Nationwide Analysis of Pelvic Ring Fractures: Incidence and Trends in Treatment, Length of Stay, and Mortality. *Geriatr Orthop Surg Rehabil.* 2016;7:9-17.
2. Sullivan MP, Baldwin KD, Donegan DJ, Mehta S, Ahn J. Geriatric fractures about the hip: divergent patterns in the proximal femur, acetabulum, and pelvis. *Orthopedics.* 2014;37:151-157.
3. Kannus P, Parkkari J, Niemi S, Sievänen H. Low-Trauma Pelvic Fractures in Elderly Finns in 1970-2013. *Calcif Tissue Int.* 2015;97:577-580.
4. Nanninga GL, de Leur K, Panneman MJ, van der Elst M, Hartholt KA. Increasing rates of pelvic fractures among older adults: The Netherlands, 1986-2011. *Age Ageing.* 2014;43:648-653.
5. Mann SM, Banaszek D, Lajkosz K, Brogly SB, Stanojev SM, Evans C, Bardana DD, Yach J, Hall S. High-energy trauma patients with pelvic fractures: Management trends in Ontario, Canada. *Injury.* 2018
6. Balogh Z, King KL, Mackay P, McDougall D, Mackenzie S, Evans JA, Lyons T, Deane SA. The epidemiology of pelvic ring fractures: a population-based study. *J Trauma.* 2007;63:1066-73; discussion 1072.
7. Pohlemann T, Tscherne H, Baumgärtel F, Egbers HJ, Euler E, Maurer F, Fell M, Mayr E, Quirini WW, Schlickewei W et al. [Pelvic fractures: epidemiology, therapy and long-term outcome. Overview of the multicenter study of the Pelvis Study Group]. *Unfallchirurg.* 1996;99:160-167.

8. Rommens PM, Hofmann A. Comprehensive classification of fragility fractures of the pelvic ring: Recommendations for surgical treatment. *Injury*. 2013;44:1733-1744.
9. Tile M. Pelvic ring fractures: should they be fixed. *J Bone Joint Surg Br*. 1988;70:1-12.
10. Young JW, Burgess AR, Brumback RJ, Poka A. Pelvic fractures: value of plain radiography in early assessment and management. *Radiology*. 1986;160:445-451.
11. Exadaktylos AK, Hautz WE. Emergency Medicine in Switzerland. *ICU Manag Pr*. 2015;15:1 - 3.
12. Berger-Groch J, Thiesen DM, Grossterlinden LG, Schaewel J, Fensky F, Hartel MJ. The intra- and interobserver reliability of the Tile AO, the Young and Burgess, and FFP classifications in pelvic trauma. *Arch Orthop Trauma Surg*. 2019;139:645-650.
13. Pieroh P, Höch A, Hohmann T, Gras F, Märdian S, Pflug A, Wittenberg S, Ihle C, Blankenburg N, Dallacker-Losensky K et al. Fragility Fractures of the Pelvis Classification: A Multicenter Assessment of the Intra-Rater and Inter-Rater Reliabilities and Percentage of Agreement. *J Bone Joint Surg Am*. 2019;101:987-994.
14. Compston JE, McClung MR, Leslie WD. Osteoporosis. *Lancet*. 2019;393:364-376.
15. Gänsslen A, Pohlemann T, Paul C, Lobenhoffer P, Tscherné H. Epidemiology of pelvic ring injuries. *Injury*. 1996;27 Suppl 1:S-A13.
16. Pereira GJC, Damasceno ER, Dinhane DI, Bueno FM, Leite JBR, Ancheschi BDC. Epidemiology of pelvic ring fractures and injuries. *Revista Brasileira de Ortopedia*. 2017;52:260-269.
17. Rollmann MF, Herath SC, Kirchhoff F, Braun BJ, Holstein JH, Pohlemann T, Menger MD, Histing T. Pelvic ring fractures in the elderly now and then - a pelvic registry study. *Arch Gerontol Geriatr*. 2017;71:83-88.

18. Koo H, Leveridge M, Thompson C, Zdero R, Bhandari M, Kreder HJ, Stephen D, McKee MD, Schemitsch EH. Interobserver reliability of the young-burgess and tile classification systems for fractures of the pelvic ring. *J Orthop Trauma*. 2008;22:379-384.
19. Hermans E, Biert J, Edwards MJR. Epidemiology of Pelvic Ring Fractures in a Level 1 Trauma Center in the Netherlands. *Hip Pelvis*. 2017;29:253-261.
20. Ojodu I, Pohlemann T, Hopp S, Rollmann MF, Holstein JH, Herath SC. Predictors of mortality for complex fractures of the pelvic ring in the elderly: a twelve-year review from a German level I trauma center. *Injury*. 2015;46:1996-1998.
21. Ward JH. Hierarchical Grouping to Optimize an Objective Function. *J Am Statist Assoc*. 1963;58:236-244.

## Figure legends

**Figure 1.** Flow diagram showing inclusion and exclusion of patients. After excluding 52 patients due to incomplete imaging, 34 patients due to isolated acetabular fracture and 7 patients due to fracture caused by tumor, the final sample size was 958.

**Figure 2.** (A) The Frequency of pelvic ring injuries at a Swiss tertiary trauma center increased from 2007 to 2017 by 157%. In addition, the average patient age per year increased over the study period by 15% ( $p=0.031$ ). (B) No difference for increase existed between male (regression coefficient of 2.99; increase 99%) and female patients (regression coefficient of 4.63; increase 246%;  $p$  for comparing slope of linear regression = 0.347).

**Figure 3.** (A) With the Ward's method for hierarchical cluster analysis (21) a trimodal age distribution was found for the pelvic ring injuries: the 'young patients group' (total of 187 patients) had a mean age of  $25 \pm 6$  (16 – 38) years, the 'middle-aged patient' group (total of 305 patients) had a mean age of  $49 \pm 6$  (36 – 62) years and the 'older patient group' (total of 466 patients) had a mean age of  $74 \pm 10$  (58 – 98) years. (B) In all age groups the frequency increased from 2007 to 2017, however, the increase differed among the three age groups ( $p$  slope linear regression = 0.008): the largest increase in frequency was found in the 'older age group' (regression coefficient of 4.76; increase of 265%) compared to the 'middle-aged group' ( $p$  0.038) or 'young age group' ( $p$  = 0.006). The 'middle-aged group' (regression coefficient of 1.76; increase of 96%) and the 'young age group' (regression coefficient of 1.01; increase of 86%) showed a comparable increase ( $p$  = 0.380).

**Figure 4.** (A) The increase in frequency differed among the mechanisms for pelvic ring injuries (see also Table 1;  $p$  slope linear regression  $< 0.001$ ): the largest increase in frequency of 249% was found for fractures due to low-energy fractures (regression coefficient of 3.55); the second largest increase of 110% was found for fractures due to motor vehicle accidents (MVA) including accidents of pedestrians (regression coefficient of 1.55); followed by fractures due to fall from height (regression coefficient of 1.02; increase of 216%). Minimal increase was found for sports accidents (regression coefficient of 0.85; increase of 191%) or crush accidents (regression coefficient of 0.15; increase of 39%). (B) Overall, low-energy fractures accounted for 43% of trauma.

**Figure 5.** (A) Pelvic ring injuries classified according to Tile (9) (total of 944 patients [99%]) with increasing frequency in all subgroups from 2007 to 2017 ( $p$  slope linear regression = 0.239): injuries classified as ‘B: partially stable’ showed the largest increase over the study period (regression coefficient of 3.80; increase of 140%), followed by injuries classified as ‘A: stable’ (regression coefficient of 2.07; increase of 393%) and ‘C: unstable’ (regression coefficient of 1.47; increase of 88%). (B) Overall, the most common injury type was the ‘B2: lateral compression injury’ with 39% followed by ‘C1: unilateral complete disruption of posterior arch’ with 20% and ‘A2: iliac wing or anterior arch fracture’ with 14%.

**Figure 6.** (A) Pelvic ring injuries classified according to Young and Burgess (10) (total of 772 patients [81%]) from 2007 to 2017 showed significant differences for increase of frequency among the subgroups ( $p$  slope linear regression = 0.003). The ‘lateral compression’ injuries showed a marked increase in frequency of pelvic fractures (regression coefficient of 4.48;



increase of 158%) compared to ‘anteroposterior compression’ ( $p = 0.009$ ) or ‘vertical shear’ ( $p = 0.025$ ) injuries. No or only marginal increase was found for ‘anteroposterior compression’ injuries (regression coefficient of 0.03; increase of 3%) or ‘vertical shear’ injuries (regression coefficient of 0.76; increase of 187%;  $p$  slope linear regression = 0.142). (B) Overall, the most common pelvic ring injury classified according to Young and Burgess was the ‘lateral compression’ injury type 1 with 64%.

**Figure 7.** (A) Fragility fractures of the pelvis in patients aged 65 years or more classified according to Rommens and Hofmann (8) are shown (total of 320 patients [33%]). A distinct increase in ‘non-displaced posterior injuries’ could be found between 2007 and 2017 (regression coefficient of 2.48; increase of 453%,  $p$  slope linear regression among subgroups  $< 0.001$ ). (B) The ‘non-displaced posterior injuries’ accounted for 62% of all fragility fractures.

**Supplemental Figure 1.** (A) The frequency of pelvic ring injuries according to Tile (9) or Young and Burgess (10) differed among the three age groups ( $p < 0.001$  for both): In the ‘older patient group’ the prevalence of ‘unstable’ pelvic injuries was decreased ( $p < 0.001$ ) and the prevalence of ‘partially stable’ injuries was increased ( $p = 0.014$ ) compared to the other two age groups. In the ‘older patient group’ the prevalence of ‘lateral compression’ injuries were increased ( $p < 0.001$ ) and the prevalence of ‘anteroposterior compression’ or ‘vertical shear’ injuries were decreased ( $p < 0.001$  and  $p = 0.002$ , respectively) compared to the other two age groups.

**Figure 1**




Figure 2A






Figure 2B



**B**

ACCEPTED

Figure 3A



A

Figure 3B




Figure 4

## Mechanism of Trauma




Figure 5






Figure 6

## Young and Burgess Classification





Figure 7

## Fragility Fractures of the Pelvis / Rommens and Hofmann Classification



**Table 1.** Demographic data including mechanisms of trauma and injury severity score (ISS) of the patients' series with pelvic injuries according the grading systems of Tile (9), Young and Burgess (10) or age groups

	Patients (Percent)	Age (years)	Gender (percent female)	Mechanism of trauma (patients [percent of all patients])						Injury severity score (ISS) <sup>†</sup>
				Low- energy	MVA	Fall from height	Sport accident	Crush accident	Others	
<b>Overall</b>	958	57 ± 21 (16 – 98)	48	410 (43)	242 (25)	123 (13)	97 (10)	50 (5)	36 (4)	27 ± 15 (4 – 75)
<b>Tile A</b>	172 (18)	57 ± 22 (16 – 96)	88 (51)	94 (55)	33 (19)	23 (13)	11 (6)	9 (5)	2 (1)	21 ± 14 (4 – 57)
<b>Tile B</b>	507 (54)	58 ± 20 (16 – 94)	268 (53)	225 (44)	136 (27)	44 (9)	63 (12)	22 (4)	19 (4)	23 ± 13 (4 – 66)
<b>Tile C</b>	265 (28)	53 ± 20 (19 – 98)	99 (37)	81 (31)	73 (28)	53 (20)	23 (9)	18 (7)	17 (6)	34 ± 14 (4 – 75)
p-value*		0.002 B vs C	<0.001 A/B vs C	<0.001 A vs B vs C	0.098	<0.001 A/B vs C	0.048 A vs B	0.325	0.022 A vs C	<0.001 A/B vs C
<b>Y&amp;B APC</b>	126 (16)	53 ± 15 (17 – 84)	16 (13)	19 (15)	38 (30)	18 (14)	33 (26)	15 (12)	3 (2)	25 ± 14 (4 – 75)
<b>Y&amp;B LC</b>	559 (72)	58 ± 21 (16 – 98)	323 (58)	267 (48)	146 (26)	55 (10)	46 (8)	21 (4)	26 (5)	29 ± 14 (4 – 59)
<b>Y&amp;B VS</b>	87 (11)	49 ± 18 (19 – 88)	28 (32)	20 (23)	25 (29)	24 (28)	7 (8)	4 (5)	7 (8)	36 ± 15 (4 – 66)
p-value*		<0.001 APC/VS vs LC	<0.001 APC vs LC vs VS	<0.001 APC/VS vs LC	0.610	<0.001 APC/LC vs VS	<0.001 LC/VS vs APC	<0.001 LC vs APC	0.156	<0.001 APC/LC vs VS
<b>Young Patient Group</b>	187 (20)	26 ± 5 (16 – 38)	63 (34)	42 (22)	73 (39)	37 (20)	21 (11)	11 (6)	4 (2)	29 ± 16 (4 – 66)
<b>Middle- aged Group</b>	305 (32)	48 ± 6 (36 – 62)	116 (38)	90 (30)	80 (26)	60 (20)	41 (13)	21 (7)	13 (4)	27 ± 14 (4 – 75)

<b>Older Patient Group</b>	466 (49)	74 ± 10 (58 – 98)	282 (61)	278 (60)	89 (19)	26 (6)	35 (8)	18 (4)	21 (5)	25 ± 15 (4 – 66)
p-value*		<0.001 1 vs 2 vs 3	<0.001 1/2 vs 3	<0.001 1/2 vs 3	<0.001 1 vs 2 vs 3	<0.001 1/2 vs 3	0.024 1/2 vs 3	0.164	0.356	0.028 1 vs 3

Continuous data are presented as mean ± standard deviation with range in parenthesis; Y&B = Young and Burgess; APC = anteroposterior compression; LC = lateral compression; VS = vertical shear; MVA = motor vehicle accident (including pedestrian accidents); From the total of 958 pelvic injuries 944 (99%) were classified according to Tile (9) and 772 (81%) were classified according to Young and Burgess (10). \*p-value for comparisons of all three groups with significant pairwise comparisons listed; †scores were available from 556 out of 892 patients (62%) from April 2009 till December 2017.

**Table 2.** Cluster groups for the Tile (9) and Young and Burgess (10) classifications of pelvic ring injuries

<b>Tile classification</b>					
<b>Age Group</b>	<b>Gender</b>	<b>Fracture type</b>	<b>Trauma mechanism</b>	<b>Prevalence (%)</b>	
Older	Female	B2, B3, A	Low-energy fracture	30	
Young, middle-aged	Male	C, A	Fall from height, MVA	25	
Young, middle-aged	Female	B2, C1	Sport, fall from height, MVA	18	
Older	=	C, B2	MVA, fall from height	15	
Middle-aged	Male	B1, C1	Sports, crush	12	
<b>Young and Burgess classification</b>					
<b>Age Group</b>	<b>Gender</b>	<b>Fracture type</b>	<b>Trauma mechanism</b>	<b>Prevalence (%)</b>	
Older	Female	LC1	Low-energy fracture	27	
Middle-aged, young	=	LC1	MVA, fall from height	23	
Young, middle-aged	=	VS, APC2, APC3	MVA, fall from height	18	
Middle-aged, young	Male	LC, APC	Sport	18	
Older	Male	LC, APC	Low-energy, MVA	14	

The three age groups were ‘young patients’ (187 patients with mean age of  $25 \pm 6$  [16 – 38] years), ‘middle-aged patients’ (305 patients with mean age of  $49 \pm 6$  [36 – 62] years) and ‘older patients’ (466 patients with mean age of  $74 \pm 10$  [58 – 98] years). See also Figure 3. MVA = motor vehicle accident. Tile(9) classification: A = stable, B = partially stable, C = unstable fractures. Young and Burgess classification(10): LC = lateral compression, APC = anteroposterior compression, VS = vertical shear.


**Table 3.** Selected literature on demographic and epidemiological trends in patients with pelvic ring injuries

<b>Author (year)</b>	<b>Number of patients (observation period)</b>	<b>Results</b>
Gänsslen et al (15) (1996)	1842 (1972 – 1993)	MVA account for 60%, fragility fracture 26%, falls from height for 8%. 55% Tile (9) type A, 25% type B and 21% type C fractures.
Pohlemann et al (7) (1996)	1140 (1991 - 1993)	Bimodal age distribution: First peak around age of 20 to 35 years, second peak for male patients around age of 50 years and for women around 80 years. 64% type A, 21% type B, 16% type C according to Tile (10).
Balogh et al (6) (2007)	157 (2005 – 2006)	61% Tile (9) type A, 26% type B and 13% type C. High energy fractures in predominantly male, younger and more severely injured patients. Type B and C fractures more common in high-energy fractures or fractures with prehospital death.
Nanninga et al (4) (2014)	34'307 (1986 – 2011)	Pelvic fractures in patients aged >65 years: increasing incidence of 0.52/1000 patients in 1986 to 0.71/1000 patients in 2011 (increase of 37% in 25 years); constant ratio of male:female of 1:4.
Sullivan et al (2) (2014)	522'831 (1993 - 2010)	Pelvic fractures in patients aged >65 years: 24% increase in pelvic fractures for the period of 17 years; in the same time increase in elderly population (>65 years) of 30%
Kannus et al (3) (2015)	n.a. (1970 - 2013)	Low-energy trauma in patients aged over 80 years: Increasing incidence over study period from 0.73 to 3.64/1000 (five-fold increase). Higher incidence for women and increasing age.
Ojodu et al (20) (2015)	84 (2001 – 2012)	Complex pelvic fractures in patients aged >70 years: 86% of Tile (9) type B and 14% type C. 10% mortality rate.
Buller et al (1) (2016)	1'464'458 (1990 – 2007)	Increasing incidence from 0.27 to 0.34/1000 (26% increase), declining mortality from 4.2% to 2.8%, increasing surgical fixation from 7.2% to 10.4%, and decreasing hospital stay from 11.2 to 6.5 days over the 17-year observation period.
Hermans et al (19) (2017)	537 (2004 – 2014)	39% Tile (9) type B, 35% type C and 26% type A fractures. Mean ISS of 26 (ISS 19 in type A, 26 in type B, 33 in type C). Fractures were due to high energy trauma in 87% (MVA, fall from height, crush accident), 13% low-energy trauma.
Pereira et al (16) (2017)	66 (2012 – 2014)	Trauma mechanism include MVA in 45%, fragility fracture in 25%, fall from height 6%, and 24% others (total 74% high-energy trauma). A majority were Tile (9) type A fractures with 55%, type B in 29%, and C in 17%.

Rollmann et al (17) (2017)	5665 (1991 – 2013)	Patients aged >60 years: Incidence of Tile (9) type A fractures decreased (from 85% to 44%) and increased for type B (14% to 42%) and C (7% to 8%) over the study period. Stable proportion of 75% of females. Type A more frequently in female, type B and C more in male.
Mann et al (5) (2018)	3915 (2005 – 2015)	High-energy pelvic ring fractures with an ISS >16: Constant incidence of 0.046/1000 over study period. Increased proportion of patients with ISS >50 over the study period. MVA and pedestrian struck by vehicle accounting for more than half of fractures.
Current study	958 (2007 – 2017)	Increasing frequency (female>male patients) and mean age over study period. Trimodal age distribution with strongest increase in frequency in older patients. Leading trauma mechanism include low-energy fractures and MVA with increasing frequency for both. Most common fracture with increasing incidence were B2 (partially stable) according to Tile (9) and LC1 (lateral compression) according to Young and Burgess (10)

n.a. = not applicable; MVA = motor vehicle accident; ISS = injury severity score. Assessed 16 November 2020.

**Supplemental-Figure-1**





**Supplemental Table 1.** Results for the pelvic ring injuries classified according to Tile (9), Young and Burgess (10) and the fragility fractures of the pelvis (FFP) classification according to Rommens and Hofmann (8)

<b>Parameter</b>	<b>Value</b>
Tile classification (patients [percent of all patients classified according to Tile])	
A	172 (18)
A1	8 (1)
A2	128 (14)
A3	36 (4)
B	507 (54)
B1	83 (9)
B2	366 (39)
B3	58 (6)
C	265 (28)
C1	190 (20)
C2	37 (4)
C3	38 (4)
Young and Burgess classification (patients [percent of all patients classified according to Young and Burgess])	
Anterior posterior compression (APC)	126 (16)
APC Type 1	16 (2)
APC Type 2	79 (10)
APC Type 3	31 (4)
Lateral compression (LC)	559 (72)
LC1 Type 1	492 (64)
LC2 Type 2	61 (8)
LC3 Type 3	6 (1)
Vertical shear (VS)	87 (11)
Fragility fractures of pelvis (FFP) classification (patients [percent of all patients classified according to FFP])	
Type 1 (anterior injury only)	41 (13)
Type 2 (non-displaced posterior injury)	197 (62)
Type 3 (displaced unilateral posterior injury)	52 (16)
Type 4 (displaced bilateral posterior injury)	30 (9)

From the total of 958 pelvic injuries 944 (99%) were classified according to Tile (9), 772 (81%) were classified according to Young and Burgess (10) and 320 patients (33%) were aged over 65 years and classified according to the fragility fractures of pelvis (FFP) classification according to Rommens and Hofmann (8).

**Supplemental Table 2.** Demographic changes in the resident population during the study period from 2007 to 2017

Area	Gender	Age	2007 [Population]	2017 [Population]	Changes [%]
<b>Canton of Bern</b>	All	All	958'897	1'026'513	+7.1
	Male	All	466'394	503'789	+8.0
	Female	All	492'503	522'724	+6.1
	All	> 65 years	171'901	208'441	+21.3
	Male	> 65 years	70'857	91'689	+29.4
	Female	> 65 years	101'044	116'752	+15.5
	<b>Switzerland</b>	All	All	7'508'739	8'419'550
Male		All	3'679'359	4'173'437	+13.4
Female		All	3'829'380	4'246'113	+10.9
All		> 65 years	1'216'662	1'523'039	+25.2
Male		> 65 years	508'644	674'610	+32.6
Female		> 65 years	708'018	848'449	+19.8

Data from 'Federal Statistical Office of Switzerland' (<https://www.bfs.admin.ch/bfs/en/home/statistics/population.html>)