

2021

Seagrass blue carbon stocks and sequestration rates in the Colombian Caribbean

Oscar Serrano
Edith Cowan University

Diana I. Gómez-López

Laura Sánchez-Valencia

Andres Acosta-Chaparro

Raul Navas-Camacho

See next page for additional authors

Follow this and additional works at: <https://ro.ecu.edu.au/ecuworkspost2013>



Part of the [Terrestrial and Aquatic Ecology Commons](#)

10.1038/s41598-021-90544-5

Serrano, O., Gómez-López, D. I., Sánchez-Valencia, L., Acosta-Chaparro, A., Navas-Camacho, R., González-Corredor, J., ... Marbà, N. (2021). Seagrass blue carbon stocks and sequestration rates in the Colombian Caribbean. *Scientific Reports*, 11, article 11067. <https://doi.org/10.1038/s41598-021-90544-5>

This Journal Article is posted at Research Online.

<https://ro.ecu.edu.au/ecuworkspost2013/10425>

Authors

Oscar Serrano, Diana I. Gómez-López, Laura Sánchez-Valencia, Andres Acosta-Chaparro, Raul Navas-Camacho, Juan González-Corredor, Cristian Salinas, Pere Masque, Cesar A. Bernal, and Núria Marbà



OPEN

Seagrass blue carbon stocks and sequestration rates in the Colombian Caribbean

Oscar Serrano^{1,2✉}, Diana Isabel Gómez-López³, Laura Sánchez-Valencia³, Andres Acosta-Chaparro³, Raul Navas-Camacho³, Juan González-Corredor³, Cristian Salinas¹, Pere Masque^{1,4}, Cesar A. Bernal³ & Núria Marbà⁵

Seagrass ecosystems rank amongst the most efficient natural carbon sinks on earth, sequestering CO₂ through photosynthesis and storing organic carbon (C_{org}) underneath their soils for millennia and thereby, mitigating climate change. However, estimates of C_{org} stocks and accumulation rates in seagrass meadows (blue carbon) are restricted to few regions, and further information on spatial variability is required to derive robust global estimates. Here we studied soil C_{org} stocks and accumulation rates in seagrass meadows across the Colombian Caribbean. We estimated that *Thalassia testudinum* meadows store 241 ± 118 Mg C_{org} ha⁻¹ (mean ± SD) in the top 1 m-thick soils, accumulated at rates of 122 ± 62 and 15 ± 7 g C_{org} m⁻² year⁻¹ over the last ~ 70 years and up to 2000 years, respectively. The tropical climate of the Caribbean Sea and associated sediment run-off, together with the relatively high primary production of *T. testudinum*, influencing biotic and abiotic drivers of C_{org} storage linked to seagrass and soil respiration rates, explains their relatively high C_{org} stocks and accumulation rates when compared to other meadows globally. Differences in soil C_{org} storage among Colombian Caribbean regions are largely linked to differences in the relative contribution of C_{org} sources to the soil C_{org} pool (seagrass, algae *Halimeda tuna*, mangrove and seston) and the content of soil particles < 0.016 mm binding C_{org} and enhancing its preservation. Despite the moderate areal extent of *T. testudinum* in the Colombian Caribbean (661 km²), it sequesters around 0.3 Tg CO₂ year⁻¹, which is equivalent to ~ 0.4% of CO₂ emissions from fossil fuels in Colombia. This study adds data from a new region to a growing dataset on seagrass blue carbon and further explores differences in meadow C_{org} storage based on biotic and abiotic environmental factors, while providing the basis for the implementation of seagrass blue carbon strategies in Colombia.

Seagrass meadows rank amongst the most prized ecosystems on Earth due to the provision of vital ecosystem services such as nutrient cycling, biodiversity and contribution to climate change mitigation and adaption through carbon sequestration and coastal protection despite they occupy less than 0.1% of the global marine surface^{1,2}. Facing the urgency to reduce atmospheric greenhouse gases to mitigate climate change, the strategy known as blue carbon builds up on the conservation and restoration of coastal vegetated ecosystems (mangrove forests, tidal marshes, seagrass meadows and macroalgae beds) to enhance and preserve their capacity to act as natural carbon sinks³. It is estimated that seagrass globally store 140 Mg organic carbon (C_{org}) per hectare in the top meter of soils, accumulated over centennial-millennial time scales, being up to 40 times more efficient at capturing organic carbon (C_{org}) than land forests soils^{4,5}.

Despite those benefits and services provided, seagrasses are among the most threatened ecosystems on the planet, and widespread die-off of seagrasses has been estimated at 0.9% year⁻¹, due to a diversity of pressures including coastal development and coupled nutrient and sediment discharges causing eutrophication and siltation, respectively⁶. More recently, local conservation and management actions resulted in the deceleration and reversal of seagrass declining trends in Europe⁷ and USA⁸. However, it has been estimated that 0.15 Pg of CO₂ may be released annually from disturbed seagrass ecosystems, which is equivalent to 3% of those from

¹School of Science and Centre for Marine Ecosystems Research, Edith Cowan University, Joondalup, WA, Australia. ²Centro de Estudios Avanzados de Blanes, Consejo Superior de Investigaciones Científicas, Blanes, Spain. ³Marine and Coastal Research Institute "José Benito Vives De Andrés" INVEMAR, Calle 25 No. 2-55, Santa Marta, Colombia. ⁴International Atomic Energy Agency, 98000 Principality of Monaco, Monaco. ⁵Global Change Research Group, IMEDEA (CSIC-UIB), Institut Mediterrani D'Estudis Avançats, Miquel Marquès 21, 07190 Esporles, Spain. ✉email: o.serranogras@ecu.edu.au

deforestation globally⁹. The inclusion of seagrass conservation and restoration projects into carbon crediting markets could provide a financial incentive to preserve the ecosystem services they provide, including their role in C_{org} storage and climate change mitigation and adaptation¹⁰.

Seagrass meadows are found along the shores of all continents except Antarctica occupying a global area estimated to range between 0.27 to 1.65 million^{2,11,12} km², and encompass about 70 seagrass species¹³. Seagrass species have a broad dissimilarity in traits including differences in biomass and primary production rates¹⁴, that also vary across environmental conditions such as water depth and geomorphology^{15,16}. Such differences in biotic and abiotic habitat characteristics result in up to 18-fold variability in C_{org} storage capacity across seagrass ecosystems¹⁷. An increasing number of studies are reporting seagrass blue carbon stocks and accumulation rates at local, regional and global scales^{5,18–23}. However, there is a scarcity of seagrass blue carbon estimates in American countries, with the exception of Mexico, Canada, the United States in North America, and a few studies in Central and South America (Brazil, Panama and Dutch Caribbean)^{5,20,23–29}. The scarcity of seagrass blue carbon studies in these key regions is limiting our capacity to derive robust global estimates while precluding their incorporation into national carbon accounting and the implementation of blue carbon strategies within Nationally Declared Contributions to mitigate climate change.

The Caribbean Sea, a tropical region with climates ranging from semi-arid to rain forests, provides a case for a region which is currently not represented in global estimates of seagrass blue carbon. Seagrasses grow in reef lagoons between the beaches and coral reefs in the Caribbean, and can form extensive meadows in protected embayment's and estuaries. Out of the seven seagrass species found within the Caribbean Sea, *Thalassia testudinum* (turtle grass) is the most abundant species and forms persistent and climax seagrass ecosystems in the region¹³. The current area of *T. testudinum* seagrass beds in the Colombian Caribbean has been estimated in 661 km², but extensive losses have also been documented³⁰. For instance, Cartagena Bay experienced more than 90% reduction in seagrass extent over the past decades (from 2.5 to 0.18 km²) attributed to local anthropogenic disturbances³¹. Past and current anthropogenic threats to seagrass in the Caribbean region include coastal development, mining, sediment run-off and pollution, while natural threats are mainly related to hurricanes³².

This study aims to provide estimates of seagrass soil C_{org} stocks and accumulation rates for the underrepresented Colombian Caribbean (Fig. 1), based on biotic and abiotic environmental factors interacting at regional scales. We combine estimates of C_{org} density in 1 m-thick soils from *T. testudinum* meadows with soil accumulation rates derived from sediment chronologies determined with ²¹⁰Pb and ¹⁴C to estimate soil C_{org} stocks within the top meter of soil and short- (last 70 years) and long-term (last 2000 years) soil C_{org} accumulation rates. We also assess differences in soil C_{org} storage among habitats based on the contribution of seagrass- C_{org} and allochthonous- C_{org} sources to the soil C_{org} pool, and sediment grain-size. The information gathered provides baseline values for future development of blue carbon strategies in Colombia as well as key information for regional and national marine planning and governance based on seagrass blue carbon ecosystem service.

Results

Compression of seagrass soils during coring operations averaged $37 \pm 16\%$ (mean \pm SD; Supplementary Information Table A). The top meter of seagrass soils of the Colombian Caribbean had a mean \pm SD dry bulk density (DBD) of 0.87 ± 0.32 g cm⁻³, containing $6.6 \pm 4.0\%$ organic matter (OM), $1.2 \pm 0.9\%$ organic carbon (C_{org}) and 7.9 ± 3.5 mg C_{org} cm⁻³ (Table 1). Seagrass soils were mainly constituted of particles $>0.016 < 0.125$ mm ($33 \pm 24\%$), with a relatively high abundance of particles <0.016 mm ($11 \pm 14\%$) and coarse sands >0.5 mm ($21 \pm 21\%$). Soil C_{org} stocks in the top first meter ranged from 39 to 673 Mg C_{org} ha⁻¹ (241 ± 118 Mg C_{org} ha⁻¹; Table 1).

A total of 11 cores were processed to determine the sedimentation rates during the last decades using the ²¹⁰Pb dating method. Due to the high degree of mixing presented in five of the cores analyzed, it was only possible to determine sediment accumulation rates in six cores. It was possible to estimate sedimentation rates during the last 500–2000 years using the ¹⁴C dating method in five cores, based on one to three radiocarbon results per core (Supplementary Information Table B). Short- and long-term soil accretion rates (SAR) ranged from 0.9 to 7.0 mm year⁻¹ (3.9 ± 0.4 mm year⁻¹) and 0.37 to 3.4 mm year⁻¹ (1.8 ± 0.9 mm year⁻¹), respectively, while short- and long-term C_{org} accumulation rates (CAR) ranged from 34 to 195 g C_{org} m⁻² year⁻¹ (122 ± 62 g C_{org} m⁻² year⁻¹) and from 2.2 to 27.5 g C_{org} m⁻² year⁻¹ (14.9 ± 7.2 g C_{org} m⁻² year⁻¹), respectively (Table 2).

The $\delta^{13}\text{C}$ values of sedimentary organic C_{org} in seagrass soils averaged $-19.4 \pm 2.8\text{‰}$. The $\delta^{13}\text{C}$ values of seagrass, mangrove, seston and *Halimeda tuna* (Supplementary Information Table C) were used to run mixing models to estimate the contribution of potential sources into the sedimentary C_{org} pool. Seagrass detritus and seston were the most important source of C_{org} in seagrass soils from the Colombian Caribbean ($39 \pm 13\%$ and $36 \pm 24\%$, respectively), with the exception of meadows at San Andrés where macroalgae *H. tuna* contributed $47 \pm 29\%$. Mangrove matter contributed $27 \pm 15\%$ across Alta Guajira, Tayrona and Cartagena Bay regions where mangroves are present (Fig. 2). Overall, C_{org} (% and mg C_{org} cm⁻³) contents increased with increasing fine particle contents ($\% < 0.016$ mm) in all regions except Alta Guajira (Fig. 3). There was a lack of relationship between $\delta^{13}\text{C}$ and C_{org} (% and mg C_{org} cm⁻³) contents.

The DBD increased with soil depth in the seagrass meadows at Alta Guajira, Media Guajira and San Andrés, but remained relatively stable at Tayrona and Cartagena Bay (Supplementary Information Figure A). OM and C_{org} contents decreased with soil depth at all sites, while $\delta^{13}\text{C}$ remained stable with soil depth. The content of particles <0.016 mm remained relatively stable along soil depth, with the exception of higher content below 100 cm depth at Cartagena Bay. Coarse sand (>0.5 mm) content was relatively higher within the upper 20 to 90 cm at Alta Guajira, Media Guajira and Cartagena Bay, while oscillated between 5 to 40% along the soil cores at San Andrés.

Soil biogeochemical characteristics (DBD, % C_{org} , %OM, $\delta^{13}\text{C}$, and particles <0.016 mm, $>0.016 < 0.125$ mm and >0.5 mm) and soil C_{org} stocks were significantly different ($P < 0.05$) among regions (Alta Guajira, Media Guajira, Tayrona, Cartagena Bay and San Andrés; Table 3). Soil DBD in meadows at Alta Guajira (0.7 ± 0.3 g cm⁻³)




Figure 1. Seagrass meadow distribution and location of the sampling sites in the Colombian Caribbean. Left panel shows the location of the five regions studied (Alta Guajira, Media Guajira, Cartagena Bay, Tayrona and San Andrés). (A–E) show the sampling sites within each region. The size of the circles indicates the number of cores studied at each sampling site. The maps were created in ArcGIS Online (<https://www.esri.com/en-us/arcgis/products/arcgis-online/overview>).

and San Andrés ($0.7 \pm 0.2 \text{ g cm}^{-3}$) were significantly lower than at the other regions (average ranging from 1.0 to 1.1 g cm^{-3} ; Fig. 4). The soil OM content was significantly higher in Alta Guajira ($9.7 \pm 4.6\%$ OM) compared to meadows at San Andrés ($6.9 \pm 2.5\%$ OM), while OM content at both Alta Guajira and San Andrés was higher than at Media Guajira, Tayrona and Cartagena Bay (ranging from 3.9 to 5.3%; Table 3). The soil C_{org} content followed a similar pattern, with significantly higher values at Alta Guajira ($1.54 \pm 1.02\%$ C_{org}) and San Andrés ($1.47 \pm 0.80\%$ C_{org}) compared to Media Guajira ($0.52 \pm 0.16\%$ C_{org}). The C_{org} stocks within the top meter of soil were up to twofold higher in Alta Guajira ($353 \pm 125 \text{ Mg } C_{\text{org}} \text{ ha}^{-1}$) compared to Cartagena Bay ($142 \pm 61 \text{ Mg } C_{\text{org}} \text{ ha}^{-1}$), while the other regions had intermediate soil C_{org} stocks with average values ranging from 210 to $241 \text{ Mg } C_{\text{org}} \text{ ha}^{-1}$. The lower number of SAR and CAR data across regions precluded comparisons. The $\delta^{13}\text{C}$ values of soil C_{org} were significantly lower in Alta and Media Guajira (average ranging from -21 to -22‰) than in San Andrés ($-16 \pm 1.4\text{‰}$). Based on isotope mixing models, seagrass was the major contributor to the soil C_{org} pool in the meadows located at Alta Guajira, Tayrona and Cartagena Bay (46% in all cases), with $\sim 25\%$ contributions of each seston and mangrove matter (Fig. 2). At Media Guajira, seston contributed 69%, while seagrass contributed the remaining 31%. The contribution of *H. tuna* to soil C_{org} pool at San Andrés was estimated at 47%, with seagrass and seston contributing 27% each. Seagrass soils at Alta Guajira had significantly higher amount of particles $< 0.016 \text{ mm}$ ($30 \pm 15\%$) than the other regions (ranging from 2 to 13%), while particles $> 0.016 < 0.125 \text{ mm}$ were more abundant in Media Guajira ($67 \pm 17\%$) and Alta Guajira ($46 \pm 11\%$) (ranging from 12 to 29%; Table 1). The concentration of coarse sands $> 0.5 \text{ mm}$ was significantly higher in Cartagena Bay ($40 \pm 15\%$) compared to the other regions (ranging from 4 to 19%).

Region	Location	Core ID	DBD (g cm ⁻³)		C _{org} (%)		OM (%)		δ ¹³ C (‰)		<0.016 mm (%)		>0.016<0.125 mm (%)		>0.5 mm		Soil C _{org} stock in 1 m-thick (Mg C _{org} ha ⁻¹)		
			Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD			
Alta Guajira	Cabo de la Vela	AGCV1	1.0	0.2			5.0	0.8										224	
		AGCV2	1.0	0.0			4.9	0.7										246	
		AGCV3	0.9	0.0			4.2	0.6										203	
		AGCV4	1.1	0.1			4.5	1.8										279	
		AGCV5	0.9	0.0			6.2	0.9										329	
	Portete Bay	AGBP1	0.6	0.2			16.8	5.2										673	
		AGBP2	0.4	0.2			14.8	4.7										344	
		AGBP3	0.7	0.1			10.7	5.0										470	
		AGBP4	0.6	0.1			10.3	2.8										369	
		AGBP5	0.7	0.1			5.7	0.1										235	
		Hondita Bay	AGBH1	0.8	0.0			10.3	3.6										415
			AGBH2	0.8	0.1			8.4	3.0										370
			AGBH3	0.8	0.3	1.3	1.1	8.9	3.2	-21.4	1.3	16.7	5.7	41.5	12.5	11.8	14.9		266
	AGBH4		0.9	0.2			10.8	4.1										479	
	AGBH5		0.8	0.2			8.2	1.7										381	
	Media Guajira	Carrizal	AGBH6	0.5	0.1	1.8	0.9	11.6	3.4	-20.3	2.0	45.6	2.0	52.0	1.9	0.0	0.0		206
AGBH7			0.5	0.0			12.1	2.6										350	
AGBH8			0.6	0.1			15.9	3.5										522	
MGCZ1			1.2	0.1			3.3	2.0										151	
MGCZ2			1.1	0.1			6.5	1.8										357	
MGCZ3			1.1	0.3			5.2	1.2										258	
MGCZ4			1.1	0.2	0.5	0.1	2.4	0.7	-21.5	0.6	4.8	0.1	64.9	1.9	1.2	2.3		108	
MGCZ5			1.1	0.1			2.9	0.6										153	
MGCZ6		1.0	0.1			3.5	1.2										159		
MGCZ7		0.9	0.2			3.6	4.1										130		
MGCZ8		1.0	0.2			5.7	3.9										203		
MGCZ9		1.1	0.1			5.3	0.9										252		
MGCZ10	1.0	0.3	0.5	0.2	3.6	0.6	-20.0	0.9	3.4	1.7	71.0	18.3	2.3	6.4		145			
MGCZ11	1.0	0.0			5.6	2.5											309		
MGCZ12	1.1	0.1			4.7	3.6											267		
Musichi	MGMC1	1.2	0.3	0.6	0.1	3.2	1.9	-22.8	1.0	3.5	0.8	64.4	21.1	6.8	10.0		139		
	MGMC2	1.3	0.1			2.9	0.9										176		
	MGMC3	1.1	0.3			5.2	2.6										338		
San Andrés	Mar Azul	SAMA1	0.6	0.1	1.5	0.5	7.6	2.8	-16.5	1.0	23.5	3.0	35.9	4.1	0.0	0.0		193	
		SAMA2	0.7	0.1			7.5	0.9										251	
		SAMA3	0.7	0.1			5.9	0.8										209	
	Hooker Bay	SABH1	0.6	0.2	1.7	0.9	8.2	1.7	-15.3	1.4	10.8	3.6	33.4	6.7	17.6	11.3		206	
		SABH2	0.8	0.2			7.4	3.2										258	
		SABH3	0.8	0.2			5.6	4.0										179	
	Cotton Cay	SACC1	0.9	0.2	1.1	0.8	4.9	1.6	-17.2	1.1	5.7	5.9	15.6	11.9	41.9	19.4		197	
		SACC2	1.0	0.1			6.5	1.2										331	
		SACC3	1.1	0.1			5.9	1.1										343	
Tayrona	Chengue Bay	SMCH1	0.7	0.1			6.4	3.1										217	
		SMCH2	1.0	0.1			5.0	1.4										229	
		SMCH3	1.2	0.1			5.2	1.5										240	
		SMCH4	1.2	0.1			5.0	0.9										238	
Cartagena	La Virgen	CLV1	0.8	0.0			5.2	0.0			2.4	0.9	11.5	1.1	44.4	8.4		148	
		CLV2	0.8	0.1			7.1	3.2			3.1	1.1	12.1	1.1	38.1	10.6		219	
		CLV3	0.7	0.0			5.2	0.3			1.6	1.0	10.0	2.1	43.6	10.0		138	
	Tierra Bomba	CTTB1	0.9	0.1			5.3	0.9			8.5	5.0	28.4	14.8	25.6	12.9		168	
		CTTB2	0.9	0.0			5.8	0.8			9.5	4.8	25.8	13.3	29.8	10.7		184	
		CTTB3	1.0	0.0			4.8	1.2			6.6	5.2	21.1	16.6	28.6	16.7		162	
	Punta Arena	CTPA1	0.8	0.0			4.1	0.3			0.2	0.5	6.8	3.8	47.6	7.7		119	
		CTPA2	1.1	0.0			4.4	0.3			0.9	1.2	8.6	4.4	42.8	8.4		159	
		CTPA3	0.9	0.0			4.2	0.3			0.9	1.2	8.2	3.4	44.8	2.8		142	

Continued

Region	Location	Core ID	DBD (g cm ⁻³)		C _{org} (%)		OM (%)		δ ¹³ C (‰)		<0.016 mm (%)		>0.016 <0.125 mm (%)		>0.5 mm		Soil C _{org} stock in 1 m-thick (Mg C _{org} ha ⁻¹)
			Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	
	Cocon Bay	CTBC1	0.9	0.1			4.5	0.8			0.0	0.0	2.7	3.2	62.9	6.7	141
		CTBC2	1.2	0.1			4.6	0.2			0.4	0.6	6.7	1.9	45.3	14.5	200
		CTBC3	1.4	0.0			4.4	0.8			0.5	0.5	8.2	4.2	59.0	12.5	226
	Bocagrande	CTBG1	1.3	0.0			1.4	0.2			0.0	0.0	6.4	1.4	33.0	8.3	39
		CTBG2	1.3	0.1			1.3	0.1			0.0	0.0	8.1	2.3	23.3	10.8	40
		CTBG3	1.9	0.5			1.0	0.2			0.0	0.0	5.1	2.9	31.0	6.5	40
OVERALL			0.9	0.3	1.2	0.9	6.6	4.0	-19.4	2.8	11.1	13.7	33.5	23.9	21.2	21.5	241 ± 117

Table 1. Mean (±SD) dry bulk density (g cm⁻³), organic carbon content (C_{org}, %), organic matter content (OM, %), δ¹³C (‰), sediment grain-size <0.016 mm, >0.016 <0.125 mm and >0.5 mm (%) and soil organic carbon stocks (in 1 m-thick, kg C_{org} m⁻²) in seagrass soils from all cores studied in the different locations and regions in the Colombian Caribbean.

Region	Location	Core ID	Short-term SAR	Long-term SAR	Short-term CAR	Long-term CAR
			(mm year ⁻¹)	(mm year ⁻¹)	(g C _{org} m ⁻² year ⁻¹)	(g C _{org} m ⁻² year ⁻¹)
Alta Guajira	Bahia Portete	AGBP2	1.8 ± 0.3		93.3 ± 17.1	
	Bahia Hondita	AGBH3	0.89 ± 0.16	0.44 ± 0.09	33.6 ± 5.9	4.5 ± 0.9
Media Guajira	Carrizal	MGCZ3	3.3 ± 0.8		79.5 ± 20.1	
	Musichi	MGMC1		0.37 ± 0.17		2.20 ± 1.01
San Andrés	Mar Azul	SAMA1	6.97 ± 0.61	3.36 ± 1.35	167 ± 15	27.5 ± 11.1
	Bahia Hooker	SABH1	5.97 ± 0.28	2.14 ± 1.67	164 ± 8	19.1 ± 14.9
	Cotton Cay	SACC1	4.55 ± 0.15	2.84 ± 1.11	195	21.2 ± 8.2
OVERALL			3.9 ± 0.4	1.8 ± 0.9	122 ± 62	14.9 ± 7.2

Table 2. Mean (±SD) short-term (last decades) and long-term (last 2000 years) sediment accumulation rates (SAR; mm year⁻¹) and organic carbon accumulation rates (CAR; g C_{org} m⁻² year⁻¹) in seagrass soils from all cores successfully dated in the different locations and regions in the Colombian Caribbean.




Figure 2. Box plots showing the results of the isotope mixing models to determine the contribution of sources (seagrass, seston, *Halimeda tuna* and mangrove) to the soil organic carbon pool in the five study sites.

Discussion

Seagrasses are sparsely distributed along the Colombian Caribbean, but their C_{org} stocks in 1 m-thick soils (241 Mg C_{org} ha⁻¹ on average) are well above the values from global estimates (140 Mg C_{org} ha⁻¹)⁵ and other tropical regions such as Mexico, US and Indonesia (ranging^{23,33–35} from 71 to 170 Mg C_{org} ha⁻¹). Estimates of soil short-term CAR (122 g C_{org} m⁻² year⁻¹) in Colombian seagrass meadows is higher than in the Dutch Caribbean (84 g C_{org} m⁻² year⁻¹)²⁵ and Australian (36 g C_{org} m⁻² year⁻¹)²¹ meadows. The relatively high C_{org} sink capacity of *Thalassia* seagrass meadows in Colombia could be explained by a combination of biogeochemical and environmental




Figure 3. Relationships between soil organic carbon content ($\% \text{C}_{\text{org}}$ and $\text{mg C}_{\text{org}} \text{cm}^{-3}$) and sediment particles $<0.016 \text{ mm}$ and $\delta^{13}\text{C}$ (‰) in the seagrass cores from San Andrés, Cartagena Bay and La Guajira in the Colombian Caribbean.

Variable	Factor	F	df	P value
Dry bulk density (g cm^{-3})	Region	74.23	4	<0.001
Organic carbon (%)	Region	21.25	2	<0.001
Organic matter (%)	Region	83.73	4	<0.001
C_{org} stocks ($\text{Mg C}_{\text{org}} \text{ha}^{-1}$)	Region	12.47	4	<0.001
$\delta^{13}\text{C}$ (‰)	Region	148.94	2	<0.001
$<0.016 \text{ mm}$ (%)	Region	93.36	3	<0.001
$>0.016 <0.125 \text{ mm}$ (%)	Region	180.86	3	<0.001
$>0.5 \text{ mm}$ (%)	Region	145.13	3	<0.001

Table 3. Results of the Generalized Linear Models (GLM) testing for significant effects of region (Alta Guajira, Media Guajira, San Andrés, Tayrona, Cartagena) on seagrass soil dry bulk density (g cm^{-3}), soil organic carbon content (%), soil organic matter content (%), soil organic carbon stocks ($\text{Mg C}_{\text{org}} \text{ha}^{-1}$ in 1-m thick), $\delta^{13}\text{C}$ (‰), and content of sediment grain-sizes $<0.016 \text{ mm}$, $>0.016 <0.125 \text{ mm}$ and $>0.5 \text{ mm}$ (%) in the Colombian Caribbean.

processes linked to habitat characteristics. The up to twofold higher above- and below-ground biomass and production of *T. testudinum* compared to the mean of seagrass species globally¹⁴, likely provides a relatively larger amount of seagrass detritus for burial. Indeed, the tropical wet climate in Colombia together with the numerous rivers discharging along the coastline results in the accumulation of fine-sized particles $<0.0125 \text{ mm}$ ($44 \pm 3\%$) within seagrass meadows (Fig. 4), which have been shown to be conducive of soil C_{org} preservation^{36,37}. In turn, terrestrial run-off and mangrove vegetation deliver allochthonous C_{org} into the coastal zone that contributes to the seagrass soil carbon pool ($54 \pm 0.1\%$)³⁸. SAR in Colombian seagrass meadows averaged $3.9 \pm 0.4 \text{ mm year}^{-1}$ over the last century, which are up to fourfold higher than previously reported for seagrass meadows globally¹, and for China, Japan, Mediterranean Sea, Arabian Gulf and Australia (ranging from $0.8^{19,21,39}$ to 2.1 mm year^{-1} , but similar to *Zostera marina* meadows²⁰). High SAR entails rapid burial of organic matter in anoxic conditions thereby reducing C_{org} remineralization rates and enhancing C_{org} storage¹⁵. The lack of excess ^{210}Pb in some of the cores analysed suggests that there is no net accumulation in some areas, or accumulation of reworked materials. Further studies (e.g. deployment of surface elevation tables to measure sediment accretion in situ) are required to overcome the limitations of radioisotopes⁴⁰. Overall, the high seagrass primary production together with the




Figure 4. Mean (\pm SE) dry bulk density (g cm^{-3}), organic matter (%), organic carbon (% C_{org} and $\text{kg C}_{\text{org}} \text{m}^{-2}$ in 1 m-thick soils), $\delta^{13}\text{C}$ (‰), and sediment grain-sizes <0.016 mm, $>0.016 <0.125$ mm and >0.5 mm (%) in seagrass soils from Alta Guajira, Cartagena Bay, Media Guajira and San Andrés in the Colombian Caribbean. Results of GLM tests to assess differences are denoted: different letters (a, b, c) indicate significant differences ($P < 0.05$).

rapid accumulation of suspended particles results in highly depositional environments that enhance carbon accumulation and preservation, which could explain the relatively high C_{org} storage capacity of seagrass meadows along the Colombian Caribbean compared to other meadows globally.

Differences in seagrass soil carbon storage across Colombian regions. The up to threefold variability in seagrass soil C_{org} stocks among regions in the Colombian Caribbean appears to be largely linked to the relative contribution of C_{org} sources to the soil C_{org} pool (seagrass, algae, mangrove and seston) and inputs of fine sediment particles binding C_{org} and enhancing its preservation. The limited number of cores that were successfully dated with ^{210}Pb and/or ^{14}C ($N = 6$ each) precluded comparison of CAR among regions.

La Guajira contains 88% of the country's *T. testudinum* seagrass extent (581 km^2) and seagrass soil C_{org} stocks ($12.4 \text{ Tg C}_{\text{org}}$). Although Alta Guajira only contains 18 km^2 of seagrass meadows, it holds the largest soil C_{org} stocks per unit area in Colombia ($352 \text{ Mg C}_{\text{org}} \text{ha}^{-1}$)⁴¹, which also rank among the highest values reported globally⁷. The high soil C_{org} stocks per unit area in Alta Guajira are likely related to the high depositional nature of their habitats, restricted to enclosed embayments protected from hydrodynamic energy and hurricanes, with the exception of the meadows located in Cabo de la Vela⁴². The seagrass soil in Alta Guajira are mainly composed of

soil inorganic particles < 0.125 mm (76%), which together with the contribution of mangrove and seston matter to the soil C_{org} pool (28% and 26%, respectively) likely led to enhanced soil C_{org} accumulation and preservation in seagrass meadows (Figs. 2 and 4). The lack of correlation between particles < 0.016 mm and C_{org} content in Alta Guajira can be explained by the saturation of particles with adsorbed organic C_{org} together with the presence of coarse seagrass detritus³⁷. The Alta Guajira region is exposed to extreme climate events driven by tropical storms and hurricanes, which can result in the sudden destruction of seagrass meadows^{41,42} and the erosion of seagrass soil C_{org} stocks⁴³. Since the onset of seagrass monitoring at Cabo de la Vela in 2015, one cyclone disruption event resulted in the erosion of > 0.5 ha of seagrass meadows and the top ~ 0.6 m of soils underneath the meadows⁴¹, likely resulting in CO_2 emissions from perturbed soil C_{org} stocks. However, seagrasses in the protected embayments of Hondita Bay and Portete Bay within Alta Guajira remained undisturbed⁴¹. Further studies are required to assess the risk of tropical storms for blue carbon projects in tropical regions, in particular in the face of increased frequency and intensity of extreme climate events under predicted climate change scenarios⁴⁴.

The relatively low seagrass soil C_{org} stocks in Media Guajira (210 Mg C_{org} ha⁻¹) compared to the Alta Guajira (352 Mg C_{org} ha⁻¹) is likely related to the severe exposure of meadows to hydrodynamic energy in the former. This region is constantly exposed to the currents and swell generated by strong winds from the East and Northeast, which results in the resuspension of particulate materials and turbid waters that limit the water depth limit for seagrass to up to 7 m. These processes are reflected in the relatively low content of sediment particles < 0.016 mm (4%) and dominance of particles > 0.016 < 0.125 mm (66%), which together with the low seagrass contribution to the soil C_{org} pool (31%) compared to seston (69%) suggests that meadows in this region are thriving in a highly dynamic environment compared to other Colombian Caribbean regions⁴² that could be limiting their productivity owing to reduced irradiance.

Meadows along the rocky coastline of Tayrona are restricted to enclosed environments protected from the swell (only 0.9 km² of meadows mapped)³¹. The soil C_{org} stocks at our study site were relatively high (231 Mg C_{org} ha⁻¹), likely related to the high productivity of the meadows thriving in clear waters, and the input of seston and mangrove detritus from adjacent creeks. However, the lack of stable C_{org} isotope and grain size analyses in the cores from this region preclude further interpretations.

Seagrass meadows have been severely decimated at Cartagena Bay over the past decades (from 100 km² in 1930s to 9 km² in 2001⁴⁵). In turn, only small patches of seagrass meadows (< 0.1 km² each) remain at Cartagena Bay, and their soil C_{org} stocks are the lowest found across Colombia (142 Mg C_{org} ha⁻¹). Previous studies showed that soil C_{org} stocks are lower in patchy seagrass landscapes compared to continuous and extensive meadows^{36,46}. Indeed, coarse sand and gravel (40%) originated from adjacent corals dominate the soil inorganic particles in Cartagena Bay meadows, which together with the high degree of anthropization in the region could explain the weakening of seagrass C_{org} sinks in this region⁴⁷.

Seagrass meadows in San Andrés and Providencia Archipelago (5 km²) are located in a coastal lagoon basin protected from currents and waves, and to a lesser degree from tropical storms and cyclones by a reef barrier, which facilitates sedimentation^{48,49} and likely explains the relatively high soil C_{org} stocks in this region (241 Mg C_{org} ha⁻¹). The short-term CAR reported for this region (175 g C_{org} m⁻² year⁻¹) are the highest reported to date for seagrass meadows, with the exception of two *P. oceanica* meadows from the Mediterranean Sea (ranging from 202 to 249 g C_{org} m⁻² year⁻¹)⁵⁰. The seagrass soils in San Andrés contain large amounts of *H. tuna* inorganic and organic (47%) matter. The calcareous macroalgae *Halimeda* spp. is common in coral reefs and has very high primary production rates^{51,52} that likely contribute to enhance SAR through high carbonate production, and CAR through rapid burial of C_{org} in anoxic conditions. Further studies are required to understand the specific role of *H. tuna* in blue carbon, including the implications of calcification and dissolution in the carbon cycle⁵³ as well as those enhancing CAR.

Seagrass blue carbon in the Colombian Caribbean. Seagrass *T. testudinum* in the Caribbean region of Colombia occupies 661 km², with 18 km² in Alta Guajira, 563 km² in Media and Baja Guajira, 0.9 km² in Tayrona, 9 km² in Cartagena Bay, 5 km² in San Andrés and surrounding islands, and 65 km² in the Caribbean coastline west of Cartagena Bay⁴⁵. Based on the spatially-explicit estimates of seagrass extent and soil C_{org} stocks and accumulation rates in the five regions studied (Fig. 4), we estimate that seagrass in the Colombian Caribbean contain 14 Tg of C_{org} in the top meter of soils. This is equivalent to 52 Tg CO_2 stored in the soils, which corresponds to 62% of the CO_2 emissions from fossil-fuel burning by Colombia in 2014⁵⁴. Based on average CAR estimates (122 g C_{org} m⁻² year⁻¹ over the last ~ 70 years), we estimated that seagrass meadows in the Colombian Caribbean accumulate 0.081 Tg C_{org} year⁻¹ (equivalent to 0.3 Tg CO_2 year⁻¹), which corresponds to ~ 0.4% of CO_2 emissions from fossil fuels in Colombia at 2014 rates⁵⁴. It is important to note that these estimates are likely underestimating seagrass soil C_{org} storage in the Colombian Caribbean, owing to poor seagrass mapping in some regions and the exclusion of meadows formed by small species meadows of the genera *Syringodium*, *Halophila*, *Halodule* and *Ruppia* that occur in Colombia. However, owing to the lack of excess ²¹⁰Pb in some meadows, further studies are required to determine whether some of the meadows studied continue to accumulate soil C_{org} ⁴⁰. Indeed, carbonate burial in seagrass meadows, which is particularly important in tropical regions, likely offsets a portion of the CO_2 sequestered through the burial of C_{org} ⁵³. Saderne et al⁵⁵ estimated that $CaCO_3$ burial within seagrass meadows may offset ~ 30% of the net CO_2 sequestration through photosynthesis, but acknowledged that the origin of $CaCO_3$ stored within seagrass meadows is mostly derived from adjacent ecosystems rather than by calcifying organisms inhabiting or supported by seagrass ecosystems. In addition, the dissolution of $CaCO_3$ linked to seagrass metabolism and associated biota, which is also common in tropical regions, results in the release of alkalinity and thereby CO_2 removal^{55,56}. Further studies are required to decipher the role of inorganic carbon processing in the overall CO_2 budget in seagrass meadows^{53,57}.

Our study demonstrates that Colombia host some of the largest seagrass soil carbon stocks per unit area globally (241 Mg C_{org} ha⁻¹ in the top meter of soils), sitting between global median values (140 Mg C_{org} ha⁻¹) and those of *Posidonia oceanica* meadows in the Mediterranean Sea (375 Mg C_{org} ha⁻¹)⁵⁰. This characteristic in combination with the large losses of seagrass extent over the last decades in Caribbean Coast^{41,45}, places Colombia among the countries that can largely benefit from blue carbon strategies to contribute to mitigate and adapt to climate change while preserving and restoring their natural heritage and enhancing the plethora of ecosystem services that vegetated coastal ecosystems provide². Colombia hosts one of the few ongoing blue carbon projects in the world, involving carbon trading by protecting 270 km² of mangrove forests from deforestation⁵⁸.

In summary, this study demonstrates that seagrass meadows along the Colombian Caribbean are hotspots for carbon sequestration and storage, provides data from a new region to a growing dataset on seagrass blue carbon and further explores differences in soil C_{org} storage based on biotic and abiotic drivers. The spatially explicit estimates of seagrass soil C_{org} stocks contribute to identify key regions for conservation and management actions (including restoration) aiming to enhance CO₂ sequestration and/or avoid greenhouse gas emissions resulting from seagrass loss, while providing the basis for the implementation of seagrass blue carbon strategies in Colombia. The reversal of historical seagrass losses can be traded in carbon markets and in this study, we provide baseline estimates of soil C_{org} stocks and accumulation rates in Colombian seagrasses to contribute to the conservation of seagrass ecosystems through seagrass restoration strategies at the national scale.

Materials and methods

Study sites and sampling. This study was conducted in five regions (Alta Guajira, Media Guajira, Tayrona, San Andrés, and Cartagena Bay) along 2250 km of the Colombian Caribbean (Fig. 1). Seagrass species *T. testudinum* forms the most extensive and apparent meadows along the Colombian coast. More than 85% of seagrass extent in Colombia is located within the Guajira peninsula (562 km² out of the estimated 661 km²)⁴⁵.

Seagrass meadows in Alta Guajira and Media Guajira grow not deeper than 7 m water depth on sandy habitats along the shoreline, and are exposed to regular wind-driven waves, with the exception of those meadows inhabiting the few existing embayments. The meadows in La Guajira are not exposed to severe anthropogenic disturbances because the remoteness of the region, despite small indigenous and non-indigenous settlements may cause localized impacts^{31,45}. Seagrass meadows in Tayrona National Natural Park grow in shallow sandy areas among corals and within embayments, and the study sites are not impacted by direct anthropogenic activities. Cartagena Bay encompasses a wide variety of seagrass habitats with different gradients of anthropogenic disturbances (mainly from industry, boating activities and dredging) and geomorphological settings (open coastlines and embayments)⁴³. Seagrasses in San Andrés grow on shallow soil of weathered coral and the study sites are located near the main touristic village of the island⁵⁹. Out of the studied regions, mangrove forests are only present in Alta Guajira, Tayrona and Cartagena Bay. La Guajira is characterized by a drier tropical climate (1100 annual precipitation) compared to the other studied regions located further west of the Colombian Caribbean (ranging from 1600 mm per year in Tayrona and Cartagena Bay to 1900 mm per year in San Andrés).

A total of 14 sites across the five regions were sampled in this study (Fig. 1). Sixty-one soil cores were sampled in 0.5 to 3 m-deep meadows dominated by *T. testudinum* by manual percussion and rotation of PVC pipes (150 cm long, 65 mm inner diameter; Supplementary Information Table A). Three to twelve replicate cores were sampled within 200 m² of each meadow at each site (61 cores in total). Soil compaction during coring was measured as the difference in surface soil elevation inside and outside the core⁶⁰. The length of all cores was corrected for compaction (Supplementary Information Table A), and all results presented hereafter refer to the amended 'uncompressed' depths. The cores were sealed at both ends, and stored vertically at 4 °C until further processing. Seagrass plants (*T. testudinum*), mangrove (*Rhizophora mangle*) leaves, algae (*H. tuna*) and seston (>0.7 µm) samples were collected across the study sites to determine origin of soil C_{org} in seagrass meadows using stable carbon isotopes (Supplementary Information Table C).

Laboratory procedures. The cores were opened longitudinally using an electric saw and cut at 1 to 15 cm-thick. The samples were dried at 60 °C until constant weight to estimate the dry weight density (g cm⁻³), and then divided in two by quartering for subsequent analyses. One set of subsamples were milled using an electric mortar and used for soil organic matter and C_{org} content analyses to estimate soil C_{org} storage, and stable carbon isotope (δ¹³C) analysis to determine the sources of soil C_{org}. The other set of subsamples was used for radioisotope (²¹⁰Pb and ¹⁴C) dating of the cores and soil grain-size analysis to determine distribution of particle sizes. Organic matter analyses were run in all samples from all cores sampled (N = 61), while C_{org} and grain-size analyses were only conducted in alternate samples along eight and 23 cores, respectively (Table 1).

For organic matter analyses, 4 g of milled sample was combusted at 550 °C for 4 h to estimate the proportionate loss of organic matter (loss on ignition, LOI)⁶¹. For the analyses of soil C_{org} and δ¹³C, 2 g of ground sample was acidified with 4% HCl to remove inorganic carbon, centrifuged (3400 revolutions per minute, for 5 min), and the supernatant with acid residues was carefully removed by pipette, avoiding resuspension. The sample was then washed with Milli-Q water, centrifuged and the supernatant removed. The residual samples were re-dried and then encapsulated for C_{org} and δ¹³C analyses using a Thermo Delta V ConFlo III coupled to a Costech 4010 at the UH Hilo Analytical Laboratory (USA). The C_{org} content was calculated for the bulk (pre-acidified) samples. Macrophyte samples containing inorganic carbon and seston were acidified prior to C_{org} analysis, following the procedures above for soils. Stable carbon isotope ratios (δ¹³C) are expressed as δ values in parts per thousand (‰) relative to the Vienna PeeDee Belemnite standard. Replicate assays and standards indicated measurement errors of 0.02% for C_{org} content and 0.07‰ for δ¹³C. Sediment grain size was measured with a Coulter LS230 laser-diffraction particle analyzer at the University of Barcelona (Spain), after digestion of organic matter with 30% H₂O₂. The grain size data was classified in the following fractions: <0.016 mm, >0.016 <0.125 mm and >0.5 mm.

Concentrations of ^{210}Pb were determined by alpha spectrometry through the measurement of its granddaughter ^{210}Po , assuming radioactive equilibrium between the two radionuclides⁶². Between 150 and 300 mg of each sample were spiked with known amounts of ^{209}Po and acid digested using a microwave oven. The Po isotopes were plated onto silver discs (99.99% purity) and their emissions were measured by alpha spectrometry using Passivated Implanted Planar Silicon (PIPS) detectors (CANBERRA, Mod. PD-450.18 A.M.) and silicon surface barrier detectors (ORTEC, mod. Ensemble). A selection of samples from some cores was measured for ^{226}Ra by gamma spectrometry using a high-purity Ge well-type detector (CANBERRA, Mod. GCW3523) to determine the concentrations of supported ^{210}Pb , used to calculate the excess ^{210}Pb concentrations were calculated. The ^{226}Ra concentrations were found to be comparable to those of ^{210}Pb at depth in those cores, and thus the later were taken as supported ^{210}Pb for the rest of the cores. The average sediment accumulation rates over the past 100 years could be estimated for six of the eleven dated sediment cores using the Constant Flux:Constant Sedimentation (CF:CS) model below the mixed surface layer when present⁶³. The other cores showed either evidence of intense mixing or negligible presence of excess ^{210}Pb that precluded estimating a sedimentation rate.

A total of 17 radiocarbon analyses were conducted in the bulk soil organic matter in seven cores sampled across the five regions (1–3 ^{14}C analyses per core; Supplementary Information Table B) at the AMS Direct Laboratory (USA) following standard procedures⁶⁴. The ^{14}C age-depth models were produced using the R routine “Bacon” for Bayesian chronology building⁶⁵, after calibration of ^{14}C ages using the marine13 radiocarbon age calibration curve, considering a local delta R of 25 ± 8 years⁶⁶, and assuming that the top core corresponded to the year of sampling. In two cores, the ^{14}C results showed that either the core was mixed or that the samples were younger than ~400 years and therefore, it was not possible to estimate age-depth chronologies based on ^{14}C in these cores. All ^{14}C ages reported in this study are expressed as radiocarbon calibrated years.

Numerical procedures. A linear regression between % C_{org} and % LOI ($R^2 = 0.78$) was used to estimate the % C_{org} content for subsamples that were not analyzed for C_{org} ⁵. The density of organic carbon ($\text{g } C_{\text{org}} \text{ cm}^{-3}$) was calculated for each core slice by multiplying the sediment dry bulk density (g cm^{-3}) by the C_{org} concentration (%). To allow direct comparisons among locations, the soil C_{org} stocks per unit area (cumulative stocks; $\text{Mg } C_{\text{org}} \text{ ha}^{-1}$) were standardized to 1 m-thick deposits. When sediment cores (32 out of 61) were shorter than 100 cm (65 to 90 cm), soil C_{org} stocks in the top 1 m were estimated by extrapolating the integrated values of C_{org} content (cumulative C_{org} stock; $\text{Mg } C_{\text{org}} \text{ ha}^{-1}$) linearly with depth. To validate this extrapolation approach, we assessed the correlation between extrapolated C_{org} stocks from 65 cm to 1 m and measured C_{org} stocks in 1 m soil cores across the 29 sampled cores that were equal or longer than 100 cm ($r = 0.97$; $P < 0.001$; Supporting Information Figure B). Soil C_{org} accumulation rates (expressed in $\text{g } C_{\text{org}} \text{ m}^{-2} \text{ year}^{-1}$) for the last century and the last 2500 years were estimated using ^{210}Pb and ^{14}C age-depth models, respectively. Short-term (based on ^{210}Pb) and long-term (based on ^{14}C) soil C_{org} accumulation were calculated in 6 and 5 out of the 61 cores sampled, respectively, by multiplying the C_{org} inventories in 1 m-thick soil by the respective average soil accretion rate (cm year^{-1}).

Pearson correlation analyses were run to assess significant relationships between soil C_{org} storage (stocks in $\text{Mg } C_{\text{org}} \text{ ha}^{-1}$ and short- and long-term accumulation rates in $\text{g } C_{\text{org}} \text{ m}^{-2} \text{ year}^{-1}$) and soil $\delta^{13}\text{C}$ and particles < 0.016 mm. Statistical analyses were performed using univariate General Linear Models (GLM) procedures in Statgraphics v16 software. The GLMs were run to test for significant effects of region (Alta Guajira, Media Guajira, San Andrés, Tayrona and Cartagena) on the soil organic carbon content (% C_{org}), soil C_{org} stocks ($\text{Mg } C_{\text{org}} \text{ ha}^{-1}$), short- and long-term soil C_{org} accumulation rates ($\text{g } C_{\text{org}} \text{ m}^{-2} \text{ year}^{-1}$), stable carbon isotope values of soil C_{org} ($\delta^{13}\text{C}$), % < 0.016 mm, % 0.016 to 0.125 mm and % > 0.5 mm (Table 3). Region was treated as fixed factor and all response variables were square-root transformed prior to analyses to meet homogeneity of variance. Normality and homoscedasticity of model residuals were assessed visually.

One-isotope, two- to three-source mixing models were run in R using simmr and rjags packages^{67,68}. The isotope used in this model was $\delta^{13}\text{C}$, and the types and values of source end-members used varied across regions based on the presence/absence of potential sources and the specific values obtained for each region. Seagrass, mangrove and seston were included for Alta Guajira, Tayrona and Cartagena Bay, while only seagrass and seston were included for the Media Guajira region, and seagrass, seston and *H. tuna* for San Andrés (Supplementary Information Table C). The reference isotopic signatures (mean \pm SD) used in this model comprised $\delta^{13}\text{C}$ values from this study.

Received: 27 November 2020; Accepted: 6 May 2021

Published online: 26 May 2021

References

- Duarte, C. M., Losada, I. J., Hendriks, I. E., Mazarrasa, I. & Marbà, N. The role of coastal plant communities for climate change mitigation and adaptation. *Nat. Clim. Change* **3**, 961–968 (2013).
- Cullen-Unsworth, L. C. *et al.* Seagrass meadows globally as a coupled social–ecological system: Implications for human wellbeing. *Mar. Pollut. Bull.* **83**, 387–397 (2014).
- Nellemann, C. *et al.* Blue Carbon: a rapid response assessment. United Nations Environment Programme, GRID-Arendal (2009).
- McLeod, E. *et al.* A blueprint for blue carbon: toward an improved understanding of the role of vegetated coastal habitats in sequestering CO_2 . *Front. Ecol. Environ.* **9**, 552–560 (2011).
- Fourqurean, J. W. *et al.* Seagrass ecosystems as a globally significant carbon stock. *Nat. Geosci.* **5**, 505–509 (2012).
- Waycott, M. *et al.* Accelerating loss of seagrasses across the globe threatens coastal ecosystems. *PNAS* **106**, 12377–12381 (2009).
- de los Santos, C. B. *et al.* Recent trend reversal for declining European seagrass meadows. *Nature Commun.* **10**, 1–8 (2019).
- Lefcheck, J. S. *et al.* Long-term nutrient reductions lead to the unprecedented recovery of a temperate coastal region. *PNAS* **115**, 3658–3662 (2018).

9. Pendleton, L. *et al.* Estimating global “blue carbon” emissions from conversion and degradation of vegetated coastal ecosystems. *PLoS ONE* **7**, e43542 (2012).
10. Macreadie, P. I. *et al.* The future of Blue Carbon science. *Nat. Commun.* **10**, 1–13 (2019).
11. Jayathilake, D. R. M. & Costello, M. J. A modelled global distribution of the seagrass biome. *Biol. Conserv.* **226**, 120–126 (2018).
12. McKenzie, L. *et al.* The global distribution of seagrass meadows. *Environ. Res. Lett.* **15**, 074041 (2020).
13. Den Hartog, C., & Kuo, J. Taxonomy and biogeography of seagrasses. In *Seagrasses: biology, ecology and conservation* (pp. 1–23). Springer, Dordrecht (2007).
14. Duarte, C. M. & Chiscano, C. L. Seagrass biomass and production: a reassessment. *Aquat. Bot.* **65**, 159–174 (1999).
15. Serrano, O., Lavery, P. S., Rozaimi, M. & Mateo, M. A. Influence of water depth on the carbon sequestration capacity of seagrasses. *Global Biogeochem. Cy.* **28**, 950–961 (2014).
16. Gullström, M. *et al.* Blue carbon storage in tropical seagrass meadows relates to carbonate stock dynamics, plant–sediment processes, and landscape context: insights from the western Indian Ocean. *Ecosystems* **21**, 551–566 (2018).
17. Lavery, P. S., Mateo, M. A., Serrano, O. & Rozaimi, M. Variability in the carbon storage of seagrass habitats and its implications for global estimates of blue carbon ecosystem service. *PLoS ONE* **8**, e73748 (2013).
18. Campbell, J. E., Lacey, E. A., Decker, R. A., Crooks, S. & Fourqurean, J. W. Carbon storage in seagrass beds of Abu Dhabi United Arab Emirates. *Estuaries Coast* **38**, 242–251 (2015).
19. Miyajima, T. *et al.* Geographic variability in organic carbon stock and accumulation rate in sediments of East and Southeast Asian seagrass meadows. *Global Biogeochem. Cy.* **29**, 397–415 (2015).
20. Röhr, M. E. *et al.* Blue carbon storage capacity of temperate eelgrass (*Zostera marina*) meadows. *Global Biogeochem. Cy.* **32**, 1457–1475 (2018).
21. Serrano, O. *et al.* Australian vegetated coastal ecosystems as global hotspots for climate change mitigation. *Nat. Commun.* **10**, 1–10 (2019).
22. Prentice, C. *et al.* A synthesis of blue carbon stocks, sources and accumulation rates in eelgrass (*Zostera marina*) meadows in the Northeast Pacific. *Global Biogeochem. Cy.* **34**, e2019GB006345 (2020).
23. Herrera-Silveira, J. A. *et al.* Blue carbon of Mexico, carbon stocks and fluxes: a systematic review. *PeerJ* **8**, e8790 (2020).
24. Carruthers, T. J. B., Barnes, P. A., Jacome, G. & Fourqurean, J. W. Lagoon scale processes in a coastally influenced Caribbean system: implications for the seagrass *Thalassia testudinum*. *Caribb. J. Sci.* **41**, 441–455 (2005).
25. Tamis, J. E., & Foekema, E. M. A review of blue carbon in the Netherlands. IMARES Report C151/15. 13 p. (2016).
26. Thorhaug, A. L. *et al.* Gulf of Mexico estuarine blue carbon stock, extent and flux: Mangroves, marshes, and seagrasses: A North American hotspot. *Sci. Total Environ.* **653**, 1253–1261 (2019).
27. Novak, A. B. *et al.* Factors influencing carbon stocks and accumulation rates in eelgrass meadows across New England, USA. *Estuaries Coast* **43**, 2076–2091 (2020).
28. Howard, J. L., Creed, J. C., Aguia, M. V. & Fourqurean, J. W. CO₂ released by carbonate sediment production in some coastal areas may offset the benefits of seagrass “Blue Carbon”. *Limnol. Oceanogr.* **63**, 160–172 (2018).
29. Nóbrega, G. N., Romero, D. J., Otero, X. L., & Ferreira, T. O. Pedological studies of subaqueous soils as a contribution to the protection of seagrass meadows in Brazil. *Rev. Bras. Cienc. Solo* **42** (2018).
30. Gómez-López, D., C. *et al.* Informe técnico Final Proyecto de Actualización cartográfica del atlas de pastos marinos de Colombia: Sectores Guajira, Punta San Bernardo y Chocó: Extensión y estado actual. PRY-BEM-005–13 FONADE-INVEMAR. Santa Marta. 136 pp. (2014).
31. Díaz, M., Barrios Suárez, L. M., & Gómez López, D. I. Las praderas de pastos marinos en Colombia: Estructura y distribución de un ecosistema estratégico, Instituto de Investigaciones Marinas y Costeras-INVEMAR (2003).
32. Green, E. P., Short, F. T., & Frederick, T. World atlas of seagrasses. Univ of California Press (2003).
33. Phang, V. X., Chou, L. M. & Friess, D. A. Ecosystem carbon stocks across a tropical intertidal habitat mosaic of mangrove forest, seagrass meadow, mudflat and sandbar. *Earth Surf Process Landf* **40**, 1387–1400 (2015).
34. Alongi, D. M. *et al.* Indonesia’s blue carbon: a globally significant and vulnerable sink for seagrass and mangrove carbon. *Wetl. Ecol. Manag.* **24**, 3–13 (2016).
35. Thorhaug, A., Poulos, H. M., López-Portillo, J., Ku, T. C. & Berlyn, G. P. Seagrass blue carbon dynamics in the Gulf of Mexico: Stocks, losses from anthropogenic disturbance, and gains through seagrass restoration. *Sci. Total Environ.* **605**, 626–636 (2017).
36. Oreska, M. P., McGlathery, K. J. & Porter, J. H. Seagrass blue carbon spatial patterns at the meadow-scale. *PLoS ONE* **12**, e0176630 (2017).
37. Serrano, O. *et al.* Can mud (silt and clay) concentration be used to predict soil organic carbon content within seagrass ecosystems?. *Biogeosciences* **17**, 4915–4926 (2016).
38. Chen, G. *et al.* Mangroves as a major source of soil carbon storage in adjacent seagrass meadows. *Sci. Rep.* **7**, 42406 (2017).
39. Cusack, M. *et al.* Organic carbon sequestration and storage in vegetated coastal habitats along the western coast of the Arabian Gulf. *Environ. Res. Lett.* **13**, 074007 (2018).
40. Lafratta, A. *et al.* Challenges to select suitable habitats and demonstrate ‘additionality’ in Blue Carbon projects: A seagrass case study. *Ocean Coast. Manag.* **197**, 105295 (2020).
41. Gómez-López, D. I. *et al.* Reporte del estado de los arrecifes coralinos y pastos marinos en Colombia (2016–2017). Serie de publicaciones Generales del Invemar # 101, Santa Marta. 100 pp. (2018).
42. Alvarez-León, R., Aguilera-Quiñonez, J., Andrade-Maya, C. C. & Novak, P. Caracterización general de la zona de surgencia en la Guajira Colombiana. *Rev. Acad. Colomb. Cienc.* **19**, 679–694 (1995).
43. Wilson, S. S., Furman, B. T., Hall, M. O. & Fourqurean, J. W. Assessment of Hurricane Irma impacts on South Florida seagrass communities using long-term monitoring programs. *Estuaries Coast* **43**, 1119–1132 (2020).
44. Hiraishi, T. *et al.* 2013 supplement to the 2006 IPCC guidelines for national greenhouse gas inventories: Wetlands. IPCC, Switzerland (2014).
45. Diaz-M, J. M. & Gómez-López, D. I. Historic changes in the abundance and distribution of seagrass beds in the Cartagena bay and neighbouring areas (Colombia). *Bol. Invest. Mar. Cost.* **32**, 57–74 (2003).
46. Ricart, A. M. *et al.* High variability of Blue Carbon storage in seagrass meadows at the estuary scale. *Sci. Rep.* **10**, 1–12 (2020).
47. Macreadie, P. I., Allen, K., Kelaher, B. P., Ralph, P. J. & Skilbeck, C. G. Paleoreconstruction of estuarine sediments reveal human-induced weakening of coastal carbon sinks. *Glob. Change Biol.* **18**, 891–901 (2012).
48. Geister, J., & Diaz-Merlano J. M. Reef environments and geology of an oceanic archipelago: San Andrés, Old Providence and Santa Catalina (Caribbean Sea, Colombia) with Field Guide. Ingeominas. 104 pp (2007).
49. Rodríguez-Ramírez, A. *et al.* Recent dynamics and condition of coral reefs in the Colombian Caribbean. *Rev. Biol. Trop.* **58**, 107–131 (2010).
50. Serrano, O., Lavery, P. S., Lopez-Merino, L., Ballesteros, E. & Mateo, M. A. Location and associated carbon storage of erosional escarpments of seagrass *Posidonia* mats. *Front. Mar. Sci.* **3**, 42 (2016).
51. Freile, D., & Hillis, L. Carbonate productivity by *Halimeda incrassata* land in a proximal lagoon, Pico Feo, San Blas, Panama. In *Proceedings 8th International Coral Reef Symposium* **1**, 767–772 (1997).
52. van Tussenbroek, B. I. & van Dijk, J. K. Spatial and temporal variability in biomass and production of Psammophytic *Halimeda incrassata* (Bryopsidales, Chlorophyta) in a Caribbean reef lagoon. *J. Phycol.* **43**, 69–77 (2007).

53. Macreadie, P. I., Serrano, O., Maher, D. T., Duarte, C. M. & Beardall, J. Addressing calcium carbonate cycling in blue carbon accounting. *Limnol. Oceanogr. Lett.* **2**, 195–201 (2017).
54. CDIAC, 2020. Carbon Dioxide Information Analysis Center
55. Saderne, V. *et al.* Role of carbonate burial in Blue Carbon budgets. *Nat. Commun.* **10**, 1–9 (2019).
56. Challener, R. C., Robbins, L. L. & McClintock, J. B. Variability of the carbonate chemistry in a shallow, seagrass-dominated ecosystem: implications for ocean acidification experiments. *Mar. Freshw. Res.* **67**, 163–172 (2016).
57. Gattuso, J. P. *et al.* Ocean solutions to address climate change and its effects on marine ecosystems. *Front. Mar. Sci.* **5**, 337 (2018).
58. Conservation International (2020). Accessed in November 2020. <https://www.conservation.org/stories/critical-investment-in-blue-carbon>
59. Coralina-Invemar. Gómez-López, D. I., C. Segura-Quintero, P. C. Sierra-Correa y J. Garay-Tinoco (Eds). Atlas de la Reserva de Biósfera Seaflower. Archipiélago de San Andrés, Providencia y Santa Catalina. Instituto de Investigaciones Marinas y Costeras “José Benito Vives De Andrés” -INVEMAR- y Corporación para el Desarrollo Sostenible del Archipiélago de San Andrés, Providencia y Santa Catalina -CORALINA-. Serie de Publicaciones Especiales de INVEMAR # 28. Santa Marta, Colombia 180 p. (2012).
60. Glew, J. R., Smol, J. P., & Last, W. M. Sediment Core Collection and Extrusion. In: Tracking Environmental Change Using Lake Sediments. Kluwer Academic Publishers, 73–105 (2005).
61. Heiri, O., Lotter, A. F. & Lemcke, G. Loss on ignition as a method for estimating organic and carbonate content in sediments: reproducibility and comparability of results. *J. Paleolimnol.* **25**, 101–110 (2001).
62. Sanchez-Cabeza, J. A., Masqué, P. & Ani-Ragolta, I. 210Pb and 210Po analysis in sediments and soils by microwave acid digestion. *J. Radioanal. Nucl. Chem.* **227**, 19–22 (1998).
63. Krishnaswamy, S., Lal, D., Martin, J. M. & Meybeck, M. Geochronology of lake sediments. *Earth Planet. Sci. Lett.* **11**, 407–414 (1971).
64. Stuiver, M. & Polach, H. A. Discussion reporting of 14 C data. *Radiocarbon* **19**, 355–363 (1977).
65. Blaauw, M. & Christen, J. A. Flexible paleoclimate age-depth models using an autoregressive gamma process. *Bayesian Anal.* **6**, 457–474 (2011).
66. Reimer, P. J. *et al.* IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years cal BP. *Radiocarbon* **55**, 1869–1887 (2013).
67. Parnell, A. C. *et al.* Bayesian stable isotope mixing models. *Environmetrics* **24**, 387–399 (2013).
68. Parnell, A. C. Package “simmr”: A Stable Isotope Mixing Model. <https://doi.org/10.1371/journal.pone.0009672> (2019).

Acknowledgements

The research reported in this paper was supported by MAPCO Action Grant contract for European Union external actions ENV/2016/380-526 cofounded by INVEMAR and Fundación Natura (Colombia,) with the support of ECU (Australia) and CSIC (project COOPB20366, I-COOP+ program, Spain). Its contents are the sole responsibility of the authors and do not necessarily reflect the views of the European Union. O.S. was supported by an ARC DECRA (DE170101524). C.S. was supported by ECU Higher Degree by Research Scholarship. P.M. was supported by the Australian Research Council LIEF Project (LE170100219). The IAEA is grateful for the support provided to its Environment Laboratories by the Government of the Principality of Monaco. This is the Contribution #1306 from INVEMAR. This study complies with national and international guidelines, legislation and permits for the collection of plant and soil specimens. The macrophytes used in this study were identified by D.I.G.-L. and O.S. and have been deposited in the public museum collection at INVEMAR (Santa Marta, Colombia).

Author contributions

O.S. drafted the first version of this manuscript. All authors collated and analyzed data and contributed to the writing of this manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-021-90544-5>.

Correspondence and requests for materials should be addressed to O.S.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher’s note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021