

Introduction

Background

- Livestock and dairy production are economically important in Rwanda, yet present challenges for conservation agriculture.
- The farmer-preferred forage Napier grass (*Cenchrus purpureus*) is an inefficient feed source.
- Shifting to low-nitrifying cropping systems can reduce nitrous oxide (N₂O) emissions and prevent the loss of leachable mineral nitrogen (N) from low-fertility soils.
- Little is known about N dynamics resulting from the use of 'climate smart' perennial forages, even while they are promoted as a conservation agriculture practice



Brachiaria prevents potential N loss through biological nitrification inhibition (BNI). Intensive harvesting can result in yield declines in unfertilized pastures.

Perennial forage legumes (*Desmodium distortum*) can maintain yields and promote soil fertility as an intercrop. However, legumes stimulate nitrification, potentially offsetting benefits from BNI.


Methods

Study Locations

RWANDA




Soil Sampling



Bulk soil was collected to a depth of 12cm after removing the top layer of leaf litter. Soil was collected six times at each location between September 2020 and March 2021. These sampling events spanned both dry and rainy seasons, occurring immediately before and two weeks following forage harvest to account for contrasting periods of crop N demand.


Forage Treatments



Forage trials were planted in October 2019 and harvested every ~8 weeks since establishment. Maize was not harvested during the course of data collection.

Soil Analyses

Nitrification Potential



Denitrification Enzyme Activity

Results & Discussion

Nitrification Potential

Karama: Nitrification Potential

Treatment	Predicted
Brachiaria + Desmodium	~1.5
Brachiaria cv. Mulato II	~1.5
Desmodium distortum	~1.5
Maize + Desmodium	~1.5
Annual Maize Monocrop	~1.5
Pennisetum + Desmodium	~1.5
Pennisetum purpureum	~1.5

Rubona: Nitrification Potential

Treatment	Predicted
Brachiaria + Desmodium	~1.5
Brachiaria cv. Mulato II	~1.5
Desmodium distortum	~1.5
Maize + Desmodium	~1.5
Annual Maize Monocrop	~1.5
Pennisetum + Desmodium	~1.5
Pennisetum purpureum	~1.5

Estimated marginal means of treatment effects from a mixed effects linear model. Treatment and collection timepoint were both treated as fixed effects, with block and timepoint as random effects. NP is expressed as mg NO₃⁻ kg⁻¹ soil hr⁻¹. Dashed vertical line: marginal mean effect of the *C. purpureus* monocrop treatment.

Denitrification Enzyme Activity


Karama: Denitrification Enzyme Activity

Treatment	Predicted
Brachiaria + Desmodium	~1.5
Brachiaria cv. Mulato II	~1.5
Desmodium distortum	~1.5
Maize + Desmodium	~1.5
Annual Maize Monocrop	~1.5
Pennisetum + Desmodium	~1.5
Pennisetum purpureum	~1.5

Rubona: Denitrification Enzyme Activity

Treatment	Predicted
Brachiaria + Desmodium	~1.5
Brachiaria cv. Mulato II	~1.5
Desmodium distortum	~1.5
Maize + Desmodium	~1.5
Annual Maize Monocrop	~1.5
Pennisetum + Desmodium	~1.5
Pennisetum purpureum	~1.5

Estimated marginal means of treatment effects from a mixed effects linear model. Treatment and collection timepoint were both treated as fixed effects, with block and timepoint as random effects. DEA was log-transformed and expressed as μmol N₂O kg⁻¹ soil hr⁻¹. Dashed vertical line: marginal mean effect of the *C. purpureus* monocrop treatment.



Discussion

- Brachiaria* plots tended to exhibit lower nitrification and denitrification potential than Napier (*P. purpureum*) plots in the dry season. Increasing soil moisture resulted in non-significant treatment differences in the rainy season.
- The legume *D. distortum* did not stimulate nitrification or denitrification compared to non-intercropped treatments, except for *P. purpureum*.
- Maize plots had low nitrification and denitrification activity; this may be due to 'priming' effects on microbial activity from intensive forage harvesting.
- Intercropping perennial legumes with a climate-smart forage such as *Brachiaria* is a promising smallholder strategy to improve soil fertility while preserving soil N cycling benefits from BNI.

Research Questions

- Does the climate-smart forage *Brachiaria* reduce potential N loss compared to preferentially grown non-BNI forage crops such as Napier grass (*Cenchrus purpureus*) or annual maize (*Zea mays*)?
- Is nitrification and denitrification stimulated by the presence of a legume intercrop?

Acknowledgements

This research was conducted as part of the CGIAR Research Program on Livestock and is supported by CGIAR Fund Donors. Additional support was provided by the National Science Foundation Graduate Research Fellowship Program (no. 00074041). We are grateful to our colleagues in the Grossman Lab, in addition to Dr. Rodney Venterea and Dr. Scott Mitchell for their technical assistance. Dr. Jessica Gutknecht also provided feedback on research methodologies. This work was made possible by a dedicated field support team in Rwanda, led by Paulin Mutanguha. Finally, we thank CIAT office staff in Kenya and Rwanda for managing research-associated logistics.