



12-2015

## Accurate spectral solutions of first and second-order initial value problems by the ultraspherical wavelets-Gauss collocation method

Y. H. Youssri  
*Cairo University*

W. M. Abd-Elhameed  
*University of Jeddah*

E. H. Doha  
*Cairo University*

Follow this and additional works at: <https://digitalcommons.pvamu.edu/aam>

### Recommended Citation

Youssri, Y. H.; Abd-Elhameed, W. M.; and Doha, E. H. (2015). Accurate spectral solutions of first and second-order initial value problems by the ultraspherical wavelets-Gauss collocation method, *Applications and Applied Mathematics: An International Journal (AAM)*, Vol. 10, Iss. 2, Article 13.  
Available at: <https://digitalcommons.pvamu.edu/aam/vol10/iss2/13>

This Article is brought to you for free and open access by Digital Commons @PVAMU. It has been accepted for inclusion in *Applications and Applied Mathematics: An International Journal (AAM)* by an authorized editor of Digital Commons @PVAMU. For more information, please contact [hvkoshy@pvamu.edu](mailto:hvkoshy@pvamu.edu).



## Accurate spectral solutions of first- and second-order initial value problems by the ultraspherical wavelets-Gauss collocation method

Y. H. Youssri<sup>1\*</sup>, W. M. Abd-Elhameed<sup>1,2</sup> and E. H. Doha<sup>1</sup>

<sup>1</sup>Department of Mathematics, Faculty of Science  
Cairo University, Giza-Egypt

<sup>2</sup>Department of Mathematics, Faculty of Science  
University of Jeddah, Jeddah, Saudi Arabia

\*Corresponding Author, E-mail: [youssri@sci.cu.edu.eg](mailto:youssri@sci.cu.edu.eg)

Received: June 17, 2014; Accepted: February 2, 2015

### Abstract

In this paper, we present an ultraspherical wavelets-Gauss collocation method for obtaining direct solutions of first- and second-order nonlinear differential equations subject to homogenous and nonhomogeneous initial conditions. The properties of ultraspherical wavelets are used to reduce the differential equations with their initial conditions to systems of algebraic equations, which then must be solved by using suitable numerical solvers. The function approximations are spectral and have been chosen in such a way that make them easy to calculate the expansion coefficients of the thought-for solutions. Uniqueness and convergence of the proposed function approximation is discussed. Four illustrative numerical examples are considered and these results are comparing favorably with the analytic solutions and proving more accurate than those discussed by some other existing techniques in the literature.

**Keywords:** Ultraspherical polynomials, wavelets, first- and second-order initial value problems, Gaussian quadrature, Bratu Equation

**MSC 2010 No.:** 65M70, 65N35, 35C10, 42C10

## 1. Introduction

In the fields of applied mathematics and physics, first- and second-order ordinary initial value problems (IVPs) are often encountered. An effective method is required to analyze the mathematical model which provides solutions conforming to physical reality, i.e., the real world of physics. Therefore, we should have the ability for solving nonlinear first- and second-order ordinary differential equations (ODEs).

Many practical problems arising in numerous branches of science and engineering require solving high even-order differential equations and, in particular, second-order differential equations.

For example, after spacial discretization, a large class of nonlinear wave equations, including Klein-Gordon and Sine-Gordon equations, are reduced to certain systems of second-order ODEs. Moreover, second-order ODEs appear in celestial mechanics, circuit theory, control theory, chemical kinetics, and biology. In the sequence of papers (Abd-Elhameed (2009); Abd-Elhameed et al. (2014); Abd-Elhameed et al. (2013); Doha and Abd-Elhameed (2002); Doha et al. (2013a)), the authors dealt with the high odd- and high even-order differential equations by the Galerkin and Pertov-Galerkin methods. They constructed suitable basis functions which satisfy the underlying boundary conditions of the given differential equation. The suggested algorithms in these articles are suitable for handling the high-order linear differential equations, while the authors in Doha et al. (2013b) used another technique based on employing operational matrix of derivatives for handling the so-called Lane-Emden equation.

Among the most important second-order differential equations is the one-dimensional Bratu problem, which has a long history. Due to its great importance in various applications, a substantial amount of research work has been done for studying this problem (see, for example, Boyd (1986); Boyd (2003); Buckmire (2004)). In Boyd (1968) and Boyd (2003) Boyd employed Chebyshev polynomial expansions and the Gegenbauer polynomials as base functions. Syam and Hamdan (2006) presented the Laplace Adomian decomposition method for solving such problem. Also, Aksoy and Pakdemirli (2010) suggested new perturbation iteration solutions for Bratu-type equations, while Wazwaz (2005) presented the Adomian Decomposition method for solving this problem. Also, Venkatesh et al. (2012) used Legendre wavelets for solving IVPs of Bratu-type and discussed the analysis concerned with the convergence and error analysis.

The subject of wavelets has recently drawn a great deal of attention from mathematical scientists in various disciplines. It is creating a common link between mathematicians, physicists, and electrical engineers. Wavelet theory is a relatively new and an emerging area in mathematical research. It has been applied to a wide range of engineering disciplines; in particular, wavelets are very successfully used in signal analysis for wave form representation and segmentations, time frequency analysis and fast algorithms for easy implementation. Wavelets permit the accurate representation of a variety of functions and operators. Moreover, wavelets establish a connection with fast numerical algorithms (see Constantinides (1987); Newland (1993)).

Recently, wavelets represent a significant approach in different fields of science and engineering. In literature, many authors have used various wavelets (Beylkin et al. (1991); Chen and Hsiao

(1997); Razzaghi and Yousefi (2000)) for analyzing problems of greater computational complexity, and they have pointed out that wavelets represent powerful tools to explore a new direction in solving differential equations. For example, Legendre wavelets are used to obtain an approximate solution of Lane-Emden type equations in Yousefi (2006).

The problem of approximating definite integrals is of central importance in many applications of mathematics. In practice, a mere approximation of an integral very often will not be satisfactory unless it is accompanied by an estimate of the error. For most quadrature formulas of practical interest, error bounds are available in the literature which use, for example, norms of higher-order derivatives or bounds for the integrand in the complex plane. However, in many practical situations such information about the integrand is not available. In particular, automatic quadrature routines are designed such that the user only has to insert the limits of integration, a routine for computing the integrand, a tolerance for the error, and an upper bound for the number of function evaluations (see Krommer and Ueberhuber (1994); Krommer and Ueberhuber (1998); Davis and Rabinowitz (1984)). Most quadrature methods used in modern numerical software packages like those of NAG (1998) and IMSL (1998) are based on Gaussian quadrature formulas. Furthermore, both numerical experience and theoretical results show the superiority of Gaussian formulas over many other quadrature formulas in many function classes (see in particular Brass et al. (1997) and the literature cited therein).

Collocation methods (see, for instance, Guo and Yan (2009); Doha et al. (2013b)) have become increasingly popular for solving ordinary and partial differential equations. In particular, they are very useful in providing highly accurate solutions to nonlinear differential equations.

In this paper, we aim to introduce a collocation wavelets algorithm for handling first- and second-order IVPs based on using ultraspherical wavelets collocation algorithm. This algorithm is based on making use of the ultraspherical wavelets together with the Gaussian integration formula. In this article, we discuss the convergence analysis and uniqueness of the proposed function approximations. Numerical examples and some applications are also considered.

The paper is organized as follows. In Section 2, we enumerate some properties of ultraspherical polynomials and their shifted ones. Section 3 describes the properties of ultraspherical wavelets. Sections 4 and 5 are devoted for solving first- and second-order IVPs based on using ultraspherical wavelets collocation method. Uniqueness and convergence theorems for second-order IVPs are discussed in detail in Section 6. In Section 7, we give some numerical examples to demonstrate the validity and applicability of our proposed algorithm. Some concluding remarks are given in Section 8.

## **2. Some properties of ultraspherical polynomials and their shifted ones**

In this section we give some useful properties of ultraspherical polynomials and their shifted forms.

## 2.1. Some properties of ultraspherical polynomials

The ultraspherical polynomials (a special type of Jacobi polynomials) associated with the real parameter ( $\lambda > -\frac{1}{2}$ ), are a sequence of orthogonal polynomials defined on  $(-1, 1)$ , with respect to the weight function  $w(x) = (1 - x^2)^{\lambda - \frac{1}{2}}$ . The orthogonality relation is given by

$$\int_{-1}^1 (1 - x^2)^{\lambda - \frac{1}{2}} C_m^{(\lambda)}(x) C_n^{(\lambda)}(x) dx = \begin{cases} \frac{\sqrt{\pi} n! \Gamma(\lambda + \frac{1}{2})}{(2\lambda)_n (n + \lambda) \Gamma(\lambda)}, & m = n, \\ 0, & m \neq n, \end{cases} \quad (1)$$

where  $(2\lambda)_n$  is the Pochhammer notation, i.e.,  $(2\lambda)_n = \frac{\Gamma(n + 2\lambda)}{\Gamma(2\lambda)}$ .

It should be noted here that the ultraspherical polynomials  $C_n^{(\lambda)}(x)$  are normalized such that  $C_n^{(\lambda)}(1) = 1$ . This normalization is characterized by an advantage that the polynomials  $C_n^{(0)}(x)$  are identical with the Chebyshev polynomials of the first kind  $T_n(x)$ ,  $C_n^{(\frac{1}{2})}(x)$  are the Legendre polynomials  $L_n(x)$ , and  $C_n^{(1)}(x)$  is equal to  $(1/(n + 1))U_n(x)$ , where  $U_n(x)$  are the Chebyshev polynomials of the second kind.

The polynomials  $C_n^{(\lambda)}(x)$  may be generated by using the recurrence relation

$$(n + 2\lambda) C_{n+1}^{(\lambda)}(x) = 2(n + \lambda) x C_n^{(\lambda)}(x) - n C_{n-1}^{(\lambda)}(x), \quad n = 1, 2, 3, \dots,$$

with the initial values:  $C_0^{(\lambda)}(x) = 1$  and  $C_1^{(\lambda)}(x) = x$ .

For more properties and relations of ultraspherical polynomials, see, for instance, Andrews et al. (1999).

The shifted ultraspherical polynomials  $\tilde{C}_n^{(\lambda)}(x) = C_n^{(\lambda)}(2x - 1)$  are a sequence of orthogonal polynomials defined on  $(0, 1)$ , with respect to the weight function  $\tilde{w}(x) = (x - x^2)^{\lambda - \frac{1}{2}}$ , i.e.,

$$\int_0^1 (x - x^2)^{\lambda - \frac{1}{2}} \tilde{C}_m^{(\lambda)}(x) \tilde{C}_n^{(\lambda)}(x) dx = \begin{cases} \frac{\pi 2^{1-4\lambda} \Gamma(n + 2\lambda)}{n! (n + \lambda) (\Gamma(\lambda))^2}, & m = n, \\ 0, & m \neq n. \end{cases} \quad (2)$$

They also may generated by using the recurrence relation

$$(n + 2\lambda) \tilde{C}_{n+1}^{(\lambda)}(x) = 2(n + \lambda) (2x - 1) \tilde{C}_n^{(\lambda)}(x) - n \tilde{C}_{n-1}^{(\lambda)}(x), \quad n = 1, 2, 3, \dots,$$

with the initial values:  $\tilde{C}_0^{(\lambda)}(x) = 1$  and  $\tilde{C}_1^{(\lambda)}(x) = 2x - 1$ .

All relations and properties of ultraspherical polynomials can be easily transformed to give the corresponding relations and properties of the shifted ultraspherical polynomials.

## 2.2 Some properties of shifted ultraspherical polynomials

The shifted ultraspherical polynomials (a special type of shifted Jacobi polynomials) associated with the real parameter ( $\lambda > -\frac{1}{2}$ ) are a sequence of orthogonal polynomials on the interval  $(0, 1)$ ,

with respect to the weight function  $w(x) = (x - x^2)^{\lambda - \frac{1}{2}}$ , i.e.,

$$\int_0^1 (x - x^2)^{\lambda - \frac{1}{2}} C_m^{(\lambda)}(x) C_n^{(\lambda)}(x) dx = \begin{cases} \frac{\pi 2^{1-4\lambda} \Gamma(n + 2\lambda)}{n! (n + \lambda) (\Gamma(\lambda))^2}, & m = n, \\ 0, & m \neq n. \end{cases} \quad (3)$$

Here, we present some properties of ultraspherical polynomials (see, for instance, Andrews et al. (1999)). These polynomials are eigenfunctions of the following singular Sturm-Liouville equation

$$(x - x^2) D^2 \phi_k(x) - \left( \lambda + \frac{1}{2} \right) (2x - 1) D \phi_k(x) + k(k + 2\lambda) \phi_k(x) = 0,$$

and they may be generated with the aid of the recurrence relation

$$(k + 2\lambda) C_{k+1}^{(\lambda)}(x) = 2(k + \lambda) (2x - 1) C_k^{(\lambda)}(x) - k C_{k-1}^{(\lambda)}(x), \quad k = 1, 2, 3, \dots,$$

starting from  $C_0^{(\lambda)}(x) = 1$  and  $C_1^{(\lambda)}(x) = 2\lambda(2x - 1)$ , or obtained from the Rodrigues' formula

$$C_n^{(\lambda)}(x) = \frac{(-4)^n \Gamma(n + \lambda) \Gamma(n + 2\lambda)}{n! \Gamma(\lambda) \Gamma(2n + 2\lambda)} (x - x^2)^{\frac{1}{2} - \lambda} D^n [(x - x^2)^{n + \lambda - \frac{1}{2}}],$$

where  $D \equiv \frac{d}{dx}$ .

### 3. Ultraspherical wavelets

Wavelets constitute a family of functions constructed from dilation and translation of single function called the mother wavelet. If the dilation parameter  $a$  and the translation parameter  $b$  vary continuously, then the following family of continuous wavelets are obtained:

$$\psi_{a,b}(t) = |a|^{-1/2} \psi \left( \frac{t - b}{a} \right) \quad a, b, \in \mathbb{R}, \quad a \neq 0. \quad (4)$$

We define the ultraspherical wavelets  $\psi_{nm}^{(\lambda)}(t) = \psi(k, n, m, \lambda, t)$  as they have five arguments:  $k, n$  can be assumed to be any positive integer,  $m$  is the order for the ultraspherical polynomial,  $\lambda$  is the known ultraspherical parameter, and  $t$  is the normalized time. Explicitly, they are defined on the interval  $[0, 1]$  as:

$$\psi_{nm}^{(\lambda)}(t) = \begin{cases} 2^{\frac{k}{2}} \xi_{m,\lambda} C_m^{(\lambda)}(2^k t - 2n + 1), & t \in \left[ \frac{n-1}{2^{k-1}}, \frac{n}{2^{k-1}} \right], \\ 0, & \text{otherwise,} \end{cases} \quad (5)$$

where  $0 \leq m \leq M - 1$ ,  $1 \leq n \leq 2^{k-1}$ , and

$$\xi_{m,\lambda} = 2^\lambda \Gamma(\lambda) \sqrt{\frac{m! (m + \lambda)}{2 \pi \Gamma(m + 2\lambda)}}. \quad (6)$$

*Remark 1:* It is worthy noting here that  $\psi_{nm}^{(\frac{1}{2})}(t)$  is identical to Legendre wavelets (Razzaghi and Yousefi (2000); Yousefi (2006)),  $\psi_{nm}^{(0)}(t)$  is identical to first kind Chebyshev wavelets (Babolian and Fattahzadeh (2007); Li (2010)) and  $\psi_{nm}^{(1)}(t)$  is identical to second kind Chebyshev wavelets (Maleknejad et al. (2007)).

Now, assume a function  $f(t)$  defined on  $[0, 1]$  and suppose that it may be expanded in terms of ultraspherical wavelets as

$$f(t) = \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} c_{nm} \psi_{nm}^{(\lambda)}(t), \quad (7)$$

where

$$c_{nm} = (f(t), \psi_{nm}^{(\lambda)}(t))_w = \int_0^1 (t - t^2)^{\lambda - \frac{1}{2}} f(t) \psi_{nm}^{(\lambda)}(t) dt.$$

Assume that  $f(t)$  is approximated in terms of ultraspherical wavelets as

$$f(t) \approx \sum_{n=1}^{2^{k-1}} \sum_{m=0}^{M-1} c_{nm} \psi_{nm}^{(\lambda)}(t) = \mathbf{C}^T \Psi^{(\lambda)}(t), \quad (8)$$

where  $\mathbf{C}$  and  $\Psi^{(\lambda)}(t)$  are  $2^{k-1}M \times 1$  matrices given by

$$\mathbf{C} = [c_{1,0}, c_{1,1}, \dots, c_{1,M-1}, \dots, c_{2^{k-1},M-1}, \dots, c_{2^{k-1},1}, \dots, c_{2^{k-1},M-1}]^T, \quad (9)$$

$$\Psi^{(\lambda)}(t) = [\psi_{1,0}^{(\lambda)}, \psi_{1,1}^{(\lambda)}, \dots, \psi_{1,M-1}^{(\lambda)}, \psi_{2,0}^{(\lambda)}, \dots, \psi_{2,M-1}^{(\lambda)}, \dots, \psi_{2^{k-1},0}^{(\lambda)}, \dots, \psi_{2^{k-1},M-1}^{(\lambda)}]^T. \quad (10)$$

#### 4. Solution of first-order initial value problem

In this section, we present an ultraspherical Gaussian quadrature wavelets method (UGQWM) for solving the following first-order IVP:

$$y'(x) = f(x, y(x)), \quad y(0) = a, \quad x \in [0, 1]. \quad (11)$$

If we define the integral operator  $\mathcal{L}(\cdot) = \int_0^x (\cdot) dt$ , then applying  $\mathcal{L}$  to both sides of Equation (11) yields,

$$y(x) = a + \int_0^x f(t, y(t)) dt.$$

Let  $y_{k,M}(x) = \sum_{n=1}^{2^{k-1}} \sum_{m=0}^{M-1} c_{nm} \psi_{nm}^{(\lambda)}(x) = \mathbf{C}^T \Psi^{(\lambda)}(x)$  be an approximation of  $y(x)$ . Then we have

$$\mathbf{C}^T \Psi^{(\lambda)}(x) \approx a + \int_0^x f(t, \mathbf{C}^T \Psi^{(\lambda)}(t)) dt. \quad (12)$$

Now, we collocate Equation (12) at the points  $x_i$ , to obtain

$$\mathbf{C}^T \Psi^{(\lambda)}(x_i) = a + \int_0^{x_i} f(t, \mathbf{C}^T \Psi^{(\lambda)}(t)) dt, \quad i = 1, 2, 3, \dots, 2^{k-1}M, \quad (13)$$

where  $x_i$  are the distinct  $(2^{k-1}M)$  roots of  $C_{2^{k-1}M}^{(\lambda)}(2x - 1)$ . For the sake of applying Gaussian integration formula on Equation (13), we make use of the transformation:  $t = \frac{x_i}{2}(z + 1)$ , which

transforms the intervals  $[0, x_i]$  into the interval  $[-1, 1]$ , and therefore Equation (13) may be written in the form

$$\mathbf{C}^T \Psi^{(\lambda)}(x_i) = a + \frac{x_i}{2} \int_{-1}^1 f\left(\frac{x_i}{2}(z+1), \mathbf{C}^T \Psi^{(\lambda)}\left(\frac{x_i}{2}(z+1)\right)\right) dz, \quad i = 1, 2, 3, \dots, 2^{k-1}M. \tag{14}$$

If we apply the Gaussian integration formula on the right hand side of (14), then we get

$$\mathbf{C}^T \Psi^{(\lambda)}(x_i) \approx a + \frac{x_i}{2} \sum_{j=0}^s \omega_j f\left(\frac{x_i}{2}(z_j+1), \mathbf{C}^T \Psi^{(\lambda)}\left(\frac{x_i}{2}(z_j+1)\right)\right), \quad i = 1, 2, 3, \dots, 2^{k-1}M, \tag{15}$$

where  $z_j$ 's are the  $(s+1)$  zeros of the polynomial  $C_{s+1}^{(\lambda)}(x)$ , and the weights  $\omega_j$  (see Ralston and Rabinowitz (1978)) can be computed by the formula

$$\omega_j = \int_{-1}^1 \prod_{\substack{j=0 \\ j \neq i}}^s \left(\frac{z - z_j}{z_i - z_j}\right) dz. \tag{16}$$

The idea behind the approximation in Equation (15) is the exactness of the Gaussian integration formula for polynomials of degree not exceeding  $(2s+1)$ .

### 5. Solution of second-order initial value problem

In this section, we are interested in solving the following second-order IVP subject to homogenous initial conditions

$$y''(x) + \alpha f(x) y'(x) + \beta g(x, y(x)) = 0, \quad y(0) = y'(0) = 0, \quad x \in [0, 1], \tag{17}$$

where  $\alpha, \beta$  are real constants.

Applying  $\mathcal{L}$  on both sides of Equation (17) yields

$$y'(x) + \alpha \int_0^x f(t) y'(t) dt + \beta \int_0^x g(t, y(t)) dt = 0. \tag{18}$$

Equation (18) gives after integration by parts

$$y'(x) = -\alpha f(x) y(x) + \int_0^x [\alpha f'(t) y(t) - \beta g(t, y(t))] dt. \tag{19}$$

Now, the application of the operator  $\mathcal{L}$  again on both sides of Equation (19), and performing some manipulations, yield

$$y(x) = - \int_0^x \alpha f(t) y(t) dt + \int_0^x (x-t) [\alpha f'(t) y(t) - \beta g(t, y(t))] dt. \tag{20}$$

Therefore we have

$$\mathbf{C}^T \Psi^{(\lambda)}(x) \approx \int_0^x (x-t) [\alpha f'(t) \mathbf{C}^T \Psi^{(\lambda)}(t) - \beta g(t, \mathbf{C}^T \Psi^{(\lambda)}(t))] dt - \int_0^x \alpha f(t) \mathbf{C}^T \Psi^{(\lambda)}(t) dt. \tag{21}$$

We now collocate Equation (21) (see Canuto et al. (1988)) as



$$\begin{aligned} \mathbf{C}^T \Psi^{(\lambda)}(x_i) &= \int_0^{x_i} (x_i - t) [\alpha f'(t) \mathbf{C}^T \Psi^{(\lambda)}(t) - \beta g(t, \mathbf{C}^T \Psi^{(\lambda)}(t))] dt \\ &\quad - \int_0^{x_i} \alpha f(t) \mathbf{C}^T \Psi^{(\lambda)}(t) dt, \quad 1 \leq i \leq 2^{k-1}M, \end{aligned} \quad (22)$$

where  $x_i$  are the distinct  $(2^{k-1}M)$  roots of  $C_{2^{k-1}M}^{(\lambda)}(2x-1)$ . As in Section 4, and if we make use of the transformation  $t = \frac{x_i}{2}(z+1)$ , then after applying Gaussian integration formula, Equation (22) may be written in the form

$$\begin{aligned} \mathbf{C}^T \Psi^{(\lambda)}(x_i) &\approx \frac{x_i^2}{4} \sum_{j=0}^s \omega_j (1 - z_j) \left[ \alpha f' \left( \frac{x_i}{2}(z_j + 1) \right) \mathbf{C}^T \Psi^{(\lambda)} \left( \frac{x_i}{2}(z_j + 1) \right) \right. \\ &\quad \left. - \beta g \left( \frac{x_i}{2}(z_j + 1), \mathbf{C}^T \Psi^{(\lambda)} \left( \frac{x_i}{2}(z_j + 1) \right) \right) \right] \\ &\quad - \frac{x_i}{2} \sum_{j=0}^s \omega_j \alpha f \left( \frac{x_i}{2}(z_j + 1) \right) \mathbf{C}^T \Psi^{(\lambda)} \left( \frac{x_i}{2}(z_j + 1) \right), \quad 1 \leq i \leq 2^{k-1}M, \end{aligned} \quad (23)$$

where  $z_j$ 's are the  $(s+1)$  zeros of  $C_{s+1}^{(\lambda)}(x)$ , and the weights  $\omega_j$  can be computed from the formula in Equation (16).

*Remark 2:* For the second-order IVP with nonhomogeneous initial conditions

$$y''(x) + \alpha f(x) y'(x) + \beta g(x, y(x)) = 0, \quad y(0) = a_0, \quad y'(0) = a_1, \quad x \in [0, 1], \quad (24)$$

we use the transformation

$$u(x) = y(x) - a_0 - a_1 x,$$

which turns Equation (24) into the homogeneous second order IVP

$$u''(x) + \alpha f(x) u'(x) + \beta \bar{g}(x, u(x)) = 0, \quad u(0) = u'(0) = 0, \quad x \in [0, 1], \quad (25)$$

where

$$\bar{g}(x, u(x)) = \alpha a_1 f(x) + \beta g(x, u(x) + a_0 + a_1 x).$$

Now Equation (25) can be treated as Equation (17).

## 6. Uniqueness and convergence theorems

In this section, the theoretical analysis of uniqueness and convergence of the proposed algorithms is discussed in detail. The uniqueness of the solution of Equation (11) was discussed in Burden and Faires (2011).

Now, we state and prove two theorems. In the first we prove the uniqueness of the solution of the second-order initial value problem (17), while in the second we show that the truncated series given in (8) converges to the exact solutions of the two initial value problems (11) and (17).

**Theorem 1.** (Uniqueness Theorem)

Assume that  $|\alpha f^{(i)}(x)| \leq L_i$ , where  $L_0, L_1$  are two positive constants,  $g(x, y)$  is continuous and satisfies Lipschitz condition in the variable  $y$  with Lipschitz constant  $L_2$ . Then Equation (17) has

a unique solution whenever  $0 < L < 1$ , where  $L = x \sum_{i=0}^2 L_i$ .

**Proof:**

Let  $y_1$  and  $y_2$  be two solutions of Equation (17). Then,

$$y_1(x) = - \int_0^x \alpha f(t) y_1(t) dt + \int_0^x (x-t) \alpha f'(t) y_1(t) dt - \int_0^x (x-t) \alpha f'(t) \beta g(t, y_1(t)) dt,$$

and

$$y_2(x) = - \int_0^x \alpha f(t) y_2(t) dt + \int_0^x (x-t) \alpha f'(t) y_2(t) dt - \int_0^x (x-t) \alpha f'(t) \beta g(t, y_2(t)) dt.$$

Now

$$\begin{aligned} |y_1 - y_2| &\leq \int_0^x |\alpha f(t)| |y_1 - y_2| dt + \int_0^x (x-t) |\alpha f'(t)| |y_1 - y_2| dt \\ &+ \int_0^x (x-t) |\beta g(t, y_1) - \beta g(t, y_2)| dt \\ &\leq L|y_1 - y_2|, \end{aligned}$$

and hence

$$|y_1 - y_2|(1 - L) \leq 0,$$

but since  $0 < L < 1$ , one should has  $|y_1 - y_2| = 0$ . This proves the uniqueness and hence completes the proof.  $\square$

**Theorem 2.** (Convergence Theorem)

The truncated series expansion of Equations (11) and (17) converge to the exact ones.

**Proof:** Let

$$\begin{aligned} y(t) &= \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} c_{nm} \psi_{nm}^{(\lambda)}(t), \\ y_{k,M}(t) &= \sum_{n=1}^{2^{k-1}} \sum_{m=0}^{M-1} c_{nm} \psi_{nm}^{(\lambda)}(t), \\ y_{k,N}(t) &= \sum_{n=1}^{2^{k-1}} \sum_{m=0}^{N-1} c_{nm} \psi_{nm}^{(\lambda)}(t), \end{aligned}$$

be the exact and approximate solutions (partial sums) to Equations (11) and (17) with  $N \geq M$ . Then we have

$$\begin{aligned} \left( y(t), y_{k,N}(t) \right)_w &= \left( y(t), \sum_{n=1}^{2^{k-1}} \sum_{m=0}^{N-1} c_{nm} \psi_{nm}^{(\lambda)}(t) \right)_w \\ &= \sum_{n=1}^{2^{k-1}} \sum_{m=0}^{N-1} \bar{c}_{nm} \left( y(t), \psi_{nm}^{(\lambda)}(t) \right)_w \\ &= \sum_{n=1}^{2^{k-1}} \sum_{m=0}^{N-1} \bar{c}_{nm} c_{nm} \\ &= \sum_{n=1}^{2^{k-1}} \sum_{m=0}^{N-1} |c_{nm}|^2. \end{aligned}$$

We show that  $y_{k,N}(t)$  is a Cauchy sequence in the complete Hilbert space  $L^2(\mathbb{R})$  and hence converges.

Now

$$\begin{aligned} \|y_{k,N}(t) - y_{k,M}(t)\|_w^2 &= \left\| \sum_{n=1}^{2^{k-1}} \sum_{m=M+1}^{N-1} c_{nm} \psi_{nm}^{(\lambda)}(t) \right\|_w^2 \\ &= \sum_{n=1}^{2^{k-1}} \sum_{m=M+1}^{N-1} |c_{nm}|^2. \end{aligned}$$

Bessel's inequality implies that  $\sum_{n=1}^{\infty} \sum_{m=0}^{\infty} |c_{nm}|^2$  is convergent and this leads to

$$\|y_{k,N}(t) - y_{k,M}(t)\|_w^2 \rightarrow 0 \quad \text{as } M, N \rightarrow \infty,$$

and hence  $y_{k,N}(t)$  converges to say  $s(t)$ . We prove that  $s(t) = y(t)$ ,

$$\begin{aligned} \left( s(t) - y(t), \psi_{nm}^{(\lambda)}(t) \right)_w &= \left( s(t), \psi_{nm}^{(\lambda)}(t) \right)_w - \left( y(t), \psi_{nm}^{(\lambda)}(t) \right)_w \\ &= \left( \lim_{N \rightarrow \infty} y_{k,N}(t), \psi_{nm}^{(\lambda)}(t) \right)_w - c_{nm} \\ &= \lim_{N \rightarrow \infty} \left( y_{k,N}(t), \psi_{nm}^{(\lambda)}(t) \right)_w - c_{nm} \\ &= 0, \end{aligned}$$

and this proves that  $\sum_{n=1}^{2^{k-1}} \sum_{m=0}^{M-1} c_{nm} \psi_{nm}^{(\lambda)}(t)$  converges to  $y(t)$ .  $\square$

## 7. Illustrative examples

In this section, we apply UGQWM to solve first- and second-IVPs as well as Bratu's equation.

**Example 1.**

Consider the first-order nonlinear IVP (see, Rajabi et al. (2007); Domairry and Nadim (2008); Caruntu and Bota (2012)):

$$y' + y + y^4 = 0; \quad x \in [0, 1], \quad y(0) = 1. \tag{26}$$

We solve Equation (26) using UGQWM for the case corresponds to  $\lambda = k = 1, M = 8$  and  $s = 10$ . Table I illustrates the maximum absolute error of Example 1. The comparison between the error resulting from the numerical solution obtained by UGQWM with the error resulting from the application of the following methods:

- Homotopy perturbation method (HPM) in Rajabi et al. (2007),
- Homotopy analysis method (HAM) in Domairry and Nadim (2008),
- Squared remainder minimization method (SRMM) in Caruntu and Bota (2012),

is presented in Table I. Since no exact solution for Equation (26) is known, the relative error is computed as the difference (in absolute value) between the approximate solution and the numerical solution given by the Mathematica Wolfram software. In addition, in Figure 1, we give a comparison between different approximate solutions of (26). This figure shows that the bigger values of  $M$  attains accurate solution.

Table I: Comparison of HPM, HAM, SRMM, and UGQWM for Example 1

$x$	HPM	HAM	SRMM	UGQWM
0.1	$2.24219 \times 10^{-3}$	$1.18868 \times 10^{-4}$	$6.08673 \times 10^{-5}$	$9.20708 \times 10^{-12}$
0.2	$9.47661 \times 10^{-3}$	$8.71208 \times 10^{-4}$	$1.25683 \times 10^{-3}$	$2.82149 \times 10^{-11}$
0.3	$1.79917 \times 10^{-2}$	$1.25548 \times 10^{-3}$	$3.12286 \times 10^{-4}$	$4.99824 \times 10^{-11}$
0.4	$2.51387 \times 10^{-2}$	$1.54052 \times 10^{-3}$	$9.67612 \times 10^{-4}$	$7.33763 \times 10^{-11}$
0.5	$3.00464 \times 10^{-2}$	$2.09832 \times 10^{-3}$	$1.12410 \times 10^{-3}$	$9.40933 \times 10^{-11}$
0.6	$3.27857 \times 10^{-2}$	$3.04594 \times 10^{-3}$	$1.78443 \times 10^{-4}$	$1.13213 \times 10^{-10}$
0.7	$3.37783 \times 10^{-2}$	$4.35783 \times 10^{-3}$	$8.14607 \times 10^{-4}$	$1.33089 \times 10^{-10}$
0.8	$3.35023 \times 10^{-2}$	$5.95087 \times 10^{-3}$	$7.43428 \times 10^{-4}$	$1.48971 \times 10^{-10}$
0.9	$3.23752 \times 10^{-2}$	$7.72966 \times 10^{-3}$	$3.59748 \times 10^{-4}$	$1.84797 \times 10^{-10}$

**Example 2.**

Consider the first-order linear IVP (see Caruntu and Bota (2012)):

$$y' = 100 (\sin x - y); \quad x \in [0, 1], \quad y(0) = 0, \tag{27}$$

with the exact solution  $y(x) = 0.9999 (\sin x - 0.01 \cos x + 0.01 e^{-100x})$ . We solve Equation (27) using UGQWM for the case corresponds to  $k = 2, M = 4$  and  $s = 7$ .

Integrating Equation (27), we get

$$y(x) = 100 \left( 1 - \cos x - \int_0^x y(t) dt \right). \tag{28}$$




Figure 1: Different solutions of Example 1

The approximate solution for (27) is given by

$$y_{2,4}(x) = c_{1,0} \psi_{1,0}^{(\lambda)}(x) + c_{1,1} \psi_{1,1}^{(\lambda)}(x) + \cdots + c_{2,3} \psi_{2,3}^{(\lambda)}(x),$$

then the application of formula (15) yields

$$\mathbf{C}^T \Psi^{(\lambda)}(x_i) \approx 100(1 - \cos x) + 50 x_i \sum_{j=0}^s \omega_j \mathbf{C}^T \Psi^{(\lambda)}\left(\frac{x_i}{2}(z_j + 1)\right), \quad i = 1, 2, \dots, 8, \quad (29)$$

and hence a system of equations in the unknown expansion coefficients  $c_{n,m}$  is obtained and then solved by a suitable numerical solver. Table II illustrates the maximum absolute error of Example 2 for various values of  $\lambda$ , while in Table III, we give a comparison between the best errors obtained by using the methods in Caruntu and Bota (2012) with UGQWM. This table shows that our method is more accurate if compared with all methods developed in Caruntu and Bota (2012). In addition, a comparison between the exact solution and some approximate solutions is displayed in Figure 2.

Table II: Maximum absolute error of Example 2

$\lambda$	1	$\frac{1}{2}$	0	-0.49
$E$	$7.658 \cdot 10^{-8}$	$6.298 \cdot 10^{-8}$	$4.927 \cdot 10^{-8}$	$3.851 \cdot 10^{-8}$

### Example 3.

Consider the second-order nonlinear IVP, (see Al-Khaled and Anwar (2007)):

$$y'' + e^{-2y} = 0; \quad x \in [0, 1], \quad y(0) = 1, \quad y'(0) = \frac{1}{e}, \quad (30)$$

Table III: Different solutions of Example 2

Method	Best error
UGQWM	$3.85 \cdot 10^{-8}$
Cheb I	$2.59 \cdot 10^{-5}$
Cheb II	$2.00 \cdot 10^{-5}$
Eq. pts	$4.45 \cdot 10^{-5}$
Lobatto III A	$7.52 \cdot 10^{-5}$
Runge-Kutta	$8.82 \cdot 10^{-6}$




Figure 2: Different solutions of Example 3

with the exact solution  $y(x) = \ln(e + x)$ . Equation (30) is solved using UGQWM for the case corresponds to  $k = 1, M = 9$  and  $s = 8$ . Table IV illustrates the maximum absolute error of Example 3 for various values of  $\lambda$ , and in Table V we give a comparison between the best errors obtained by using Adomian decomposition method (ADM), quintic spline method ( $C^2$ -spline) in Al-Khaled and Anwar (2007) with the best error obtained by UGQWM.

Table IV: Maximum absolute error of Example 3

$\lambda$	1	$\frac{1}{2}$	0	-0.49
$E$	$8.322 \times 10^{-13}$	$5.284 \times 10^{-13}$	$5.293 \times 10^{-13}$	$6.004 \times 10^{-14}$

**Example 4.**

Consider the second-order nonlinear Bratu problem (see Venkatesh et al. (2012)):

$$y'' - 2e^y = 0; \quad x \in [0, 1], \quad y(0) = y'(0) = 0, \tag{31}$$

Table V: Different solutions of Example 3

Method	Best error
UGQWM	$6.004 \times 10^{-14}$
ADM	$2.20 \times 10^{-6}$
$C^2$ -spline	$1.68 \times 10^{-13}$

with the exact solution  $y(x) = -2 \log(\cos x)$ . Equation (31) is solved using UGQWM for the case corresponds to  $k = 1$ ,  $M = 9$  and  $s = 8$ . Table VI lists some approximate solutions of (31) for various values of  $\lambda$  if UGQWM is applied, and the corresponding maximum absolute error for each case is displayed also in Table VII. In Table VIII, we give a comparison between the best absolute error obtained in Venkatesh et al. (2012) by using Legendre wavelets method (LWM) with that obtained with UGQWM.

Table VI: The approximate solution and the maximum absolute error of Example 4

$\lambda$	Numerical solution by UGQWM	$E$
$\frac{1}{2}$	$0.216 x^8 - 0.607 x^7 + 0.82 x^6 - 0.533 x^5 + 0.37 x^4 - 0.045 x^3 + 1.005 x^2 - 0.00023 x$	$6.470 \times 10^{-6}$
1	$0.212 x^8 - 0.598 x^7 + 0.81 x^6 - 0.533 x^5 + 0.38 x^4 - 0.048 x^3 + 1.005 x^2 - 0.00031 x$	$9.553 \times 10^{-6}$
0	$0.222 x^8 - 0.623 x^7 + 0.83 x^6 - 0.533 x^5 + 0.39 x^4 - 0.042 x^3 + 1.004 x^2 - 0.00016 x$	$3.162 \times 10^{-6}$

Table VII: Maximum pointwise error of Example 4

$x$	0	0.2	0.4	0.6	0.8	1
Exact	0.	0.0402695	0.164458	0.38393	0.722781	1.23125
$\lambda = \frac{1}{2}$	$2.607 \times 10^{-6}$	0.0402694	0.164459	0.383929	0.722782	1.23125
$\lambda = 1$	$4.731 \times 10^{-6}$	0.0402697	0.164459	0.38393	0.722781	1.23124
$\lambda = 0$	$1.021 \times 10^{-6}$	0.040269	0.164459	0.383929	0.722782	1.23125

Table VIII: Different solutions of Example 4

Method	Best error
UGQWM	$1.02 \times 10^{-6}$
LWM	$1.38 \times 10^{-4}$

*Remark 3:* It is worthy noting here that the obtained numerical results in the previous solved four examples are very accurate, although the number of retained modes in the spectral expansion is few, and the numerical results are comparing favorably with the known analytical solutions. Moreover, the numerical results show that our method is more accurate than those discussed by some other existing techniques in literature.

## 8. Conclusion

In this paper, a new ultraspherical wavelets algorithm for obtaining numerical spectral solutions for first- and second-order IVPs is presented and analyzed. The derivation of this algorithm is essentially based on ultraspherical wavelets basis functions and the Gaussian quadrature formula. The spectral collocation method is used for handling these equations. One of the main advantages of the presented algorithms is their availability for application on both linear and non linear first- and second-order IVPs including some important equations and also a Bratu-type equation. Another advantage of the developed algorithms is highly accurate approximate solutions are achieved by using a few number of terms of the suggested expansion.

## REFERENCES

- Abd-Elhameed, W. M. (2009). Efficient spectral Legendre dual-Petrov-Galerkin algorithms for the direct solution of  $(2n+1)$ th-order linear differential equations, *J. Egypt Math. Soc.*, Vol. 17, pp. 189-211.
- Abd-Elhameed, W. M., Doha, E. H. and Bassuony, M. A. (2014). Two Legendre-dual-Petrov-Galerkin algorithms for solving the integrated forms of high odd-order boundary value problems, *The Scientific World Journal*, Vol. 2014, Article ID 309264, 11 pages.
- Abd-Elhameed, W.M., Doha, E.H. and Youssri, Y.H. (2013a). Efficient spectral-Petrov-Galerkin methods for third- and fifth-order differential equations using general parameters generalized Jacobi polynomials, *Quaest. Math.*, Vol. 36, pp. 15-38 .
- Abd-Elhameed, W. M., Doha, E. H. and Youssri, Y. H. (2013b). New wavelets collocation method for solving second-order multipoint boundary value problems using Chebyshev polynomials of third and fourth kinds, *Abstract and Applied Analysis*, Vol. 2013, Article ID 542839, 9 pages.
- Abd-Elhameed, W. M., Doha, E. H. and Youssri, Y. H. (2013c). New spectral second kind Chebyshev wavelets algorithm for solving linear and nonlinear second-order differential equations involving singular and Bratu type equations, *Abstract and Applied Analysis*, Vol. 2013, Article ID 715756, 9 pages.
- Aksoy, Y. and Pakdemirli, M. (2010). New perturbation iteration solutions for Bratu-type equations, *Comput. Math. Appl.*, Vol. 59, pp. 2802-2808.
- Al-Khaled, K. and Anwar, M. N. (2007). Numerical comparison of methods for solving second-order ordinary initial value problems, *Appl. Math. Model.*, Vol. 31, pp. 292-301.
- Andrews, G.E., Askey, R., and Roy, R. (1999). *Special Functions*, Cambridge University Press, Cambridge.
- Babolian, E. and Fattahzadeh, F. (2007). Numerical solution of differential equations by using Chebyshev wavelet operational matrix of integration, *Appl. Math. Comp.*, Vol. 188, No. 1, pp. 417-426.
- Beylkin, G., Coifman, R. and Rokhlin, V. (1991). Fast wavelet transforms and numerical algorithms, *Comm. Pure Appl.*, Vol. 44, pp. 141-183.
- Boyd, J.P. (2003). Chebyshev polynomial expansions for simultaneous approximation of two branches of a function with application to the one dimensional Bratu equation, *Appl. Math.*



- Comput., Vol. 142, pp. 189-200.
- Boyd, J.P. (1986). An analytical and numerical study of the two-dimensional Bratu equation, *J. Sci. Comput.*, Vol. 1, No. 2, pp. 183-206.
- Brass, H., Fischer, J.-W. and Petras, K. (1997). The Gaussian quadrature method, *Abh. Braunschweig. Wiss. Ges.*, Vol. 47, pp. 115-150.
- Buckmire, R. (2004), Application of a Mickens finite-difference scheme to the cylindrical Bratu-Gelfand problem, *Numer. Meth. Part. D. E.*, Vol. 20, No. 3, pp. 327-337.
- Burden, R. L. and Faires, J. D. (2011). *Numerical Analysis*, Ninth Edition, Youngstown State University.
- Canuto, C., Hussaini, M.Y., Quarteroni, A. and Zang, T.A. (1988), *Spectral Methods on Fluid Dynamics*, Springer-Verlag.
- Caruntu, B. and Bota, C. (2012). Approximate polynomial solutions for nonlinear heat transfer problems using the squared remainder minimization method, *Int. Commun. Heat Mass*, Vol. 39, pp. 1336-1341.
- Constantinides, A. (1987). *Applied Numerical Methods with Personal Computers*, McGraw-Hill, New York.
- Costabile, F. and Napoli, A. (2011). A class of collocation methods for numerical integration of initial value problems, *Comput. Math. Appl.*, Vol. 62, pp. 3221-3235.
- Chen, C. F. and Hsiao, C. H. (1997). Haar wavelet method for solving lumped and distributed-parameter systems, *IEEE Proceeding Control Theory and Applications*, Vol. 144, No. 1, pp. 87-94.
- Davis, P. and Rabinowitz, P. (1984). *Methods of Numerical Integration*, Second Edition, Academic Press, New York.
- Doha, E. H. and Abd-Elhameed, W. M. (2002). Efficient spectral-Galerkin algorithms for direct solution of second-order equations using ultraspherical polynomials, *SIAM J. Sci. Comput.*, Vol. 24, pp. 548-571.
- Doha, E.H., Abd-Elhameed, W.M. and Bassuony, M.A. (2013a). New algorithms for solving high even-order differential equations using third and fourth Chebyshev-Galerkin methods, *J. Comput. Phys.*, Vol. 236, pp. 563-579.
- Doha, E.H., Abd-Elhameed, W. M. and Youssri, Y. H. (2013b). Second kind Chebyshev operational matrix algorithm for solving differential equations of Lane-Emden type, *New Astron.*, Vol. 23-24, pp. 113-117.
- Domairry, G. and Nadim, N. (2008). Assessment of homotopy analysis method and homotopy perturbation method in non-linear heat transfer equation, *Int. Commun. Heat Mass*, Vol. 35, pp. 93-102.
- Guo, Ben-yu and Yan, Jian-ping. (2009). Legendre-Gauss collocation method for initial value problems of second order ordinary differential equations. *Appl. Numer. Math.*, Vol. 59, pp. 1386-408.
- IMSL. (1998). *Fortran Numerical Libraries*, Version 3.0 (Visual Numerics).
- Krommer, A.R. and Ueberhuber, C.W. (1994). *Numerical Integration: on Advanced Computer Systems (Lecture Notes in Computer Science)*, Vol. 848, Springer, Berlin.
- Krommer, A.R. and Ueberhuber, C.W. (1998). *Computational Integration*, SIAM, Philadelphia,

PA.

- Li, Y. (2010). Solving a nonlinear fractional differential equation using Chebyshev wavelets, *Commun. Nonlinear Sci. Numer. Simulat.*, Vol. 15, No. 9, pp. 2284-2292.
- Maleknejad, K., Sohrabi, S. and Rostami, Y. (2007). Numerical solution of nonlinear Volterra integral equations of the second kind by using Chebyshev polynomials, *Appl. Math. Comp.*, Vol. 188, No. 1, pp. 123-128.
- NAG (1998). *Fortran Library, Mark 18*, NAG, Inc.
- Newland, D. E. (1993). *An Introduction to Random Vibrations, Spectral and Wavelet Analysis*, Longman Scientific and Technical, New York.
- Rajabi, A., Ganji, D.D. and Taherian, H. (2007) Application of homotopy perturbation method in nonlinear heat conduction and convection equations, *Phys. Lett. A*, Vol. 360, pp. 570-573.
- Ralston, A. and Rabinowitz, P. (1978). *A First Course in Numerical Analysis*, Second Edition, McGraw-Hill, New York.
- Razzaghi, M. and Yousefi, S. (2000). Legendre wavelets direct method for variational problems, *Math. Comput. Simulat.*, Vol. 53, pp. 185-192.
- Syam, M.I. and Hamdan, A. (2006). An efficient method for solving Bratu equations, *Appl. Math. Comput.*, Vol. 176, pp. 704-713.
- Venkatesh, S. G., Ayyaswamy, S. K. and Balachandar, S. R. (2012). The Legendre wavelet method for solving initial value problems of Bratu-type, *Comput. Math. Appl.*, Vol. 63, pp. 1287-1295.
- Wazwaz, A.M. (2005). Adomian decomposition method for a reliable treatment of the Bratu-type equations, *Appl. Math. Comput.*, Vol. 166, pp. 652-663.
- Yousefi, S. (2006). Legendre wavelets method for solving differential equations of Lane-Emden type, *Appl. Math. Comput.*, Vol. 181, pp. 1417-1422.