



12-2015

Solution to Some Open Problems on E-super Vertex Magic Total Labeling of Graphs

G. Marimuthu
The Madura College

M. S. Raja Durga
The Madura College

G. D. Devi
The Madura College

Follow this and additional works at: <https://digitalcommons.pvamu.edu/aam>

 Part of the [Discrete Mathematics and Combinatorics Commons](#)

Recommended Citation

Marimuthu, G.; Raja Durga, M. S.; and Devi, G. D. (2015). Solution to Some Open Problems on E-super Vertex Magic Total Labeling of Graphs, *Applications and Applied Mathematics: An International Journal (AAM)*, Vol. 10, Iss. 2, Article 33.

Available at: <https://digitalcommons.pvamu.edu/aam/vol10/iss2/33>

This Article is brought to you for free and open access by Digital Commons @PVAMU. It has been accepted for inclusion in *Applications and Applied Mathematics: An International Journal (AAM)* by an authorized editor of Digital Commons @PVAMU. For more information, please contact hvkoshy@pvamu.edu.



Solution to Some Open Problems on *E*-super Vertex Magic Total Labeling of Graphs

G. Marimuthu¹, M.S. Raja Durga², G. Durga Devi³

Department of Mathematics
 The Madura College
 Madurai- 625011
 Tamilnadu, India

¹yellowmuthu@yahoo.com; ²msrajadurga17@gmail.com; ³punithadurga1991@gmail.com

Received: January 15, 2015; Accepted: December 7, 2015

Abstract

Let G be a finite graph with p vertices and q edges. A *vertex magic total labeling* is a bijection f from $V(G) \cup E(G)$ to the consecutive integers $1, 2, \dots, p+q$ with the property that for every $u \in V(G)$, $f(u) + \sum_{v \in N(u)} f(uv) = k$ for some constant k . Such a labeling is *E-super* if $f : E(G) \rightarrow \{1, 2, \dots, q\}$. A graph G is called *E-super vertex magic* if it admits an *E-super vertex magic labeling*. In this paper, we solve two open problems given by Marimuthu, Suganya, Kalaivani and Balakrishnan (Marimuthu et al., 2015).

Keywords: super vertex magic labeling; E-super vertex magic labeling; E-super vertex magic graph

2010 AMS Subject classification: 05C78

1. Introduction

In this paper, we consider only finite simple undirected graphs with order p and size q . For graph theoretic notations we follow (Harary, 1969; Marr and Wallis, 2013). A labeling of a graph G is a mapping that carries a set of graph elements, usually the vertices and edges into a set of numbers, usually integers. Many kinds of labelings have been studied and an excellent survey of graph labeling can be found in (Gallian, 2014).

Sedláček (Sedláček, 1963) introduced the concept of magic labeling. Sedláček defined a graph to be magic if it had an edge-labeling, with range the real numbers, such that the sum of the labels around any vertex equalled constant, independent of the choice of vertex. MacDougall et al. (MacDougall et al., 2002) introduced the notion of *vertex magic total labeling*. If G is a

finite simple undirected graph with p vertices and q edges, then a *vertex magic total labeling* of G is a bijection f from $V(G) \cup E(G)$ to the integers $1, 2, \dots, p+q$ with the property that for every u in $V(G)$,

$$f(u) + \sum_{v \in N(u)} f(uv) = k,$$

for some constant k , where

$$N(u) = \{v \in V(G) : uv \in E(G)\}.$$

They studied the basic properties of vertex magic total graphs and showed some families of graphs having vertex magic total labeling.

MacDougall et al. (MacDougall et al., 2004) further introduced the *super vertex magic total labeling*. They called a vertex magic total labeling f is *super* if $f : V(G) \rightarrow \{1, 2, \dots, p\}$. Swaminathan and Jeyanthi (Swaminathan and Jeyanthi, 2003) introduced a concept with the name *super vertex magic total labeling*, but with different notion. They called a *vertex magic total labelling* f to be *super* if $f : E(G) \rightarrow \{1, 2, \dots, q\}$. To avoid confusion, Marimuthu and Balakrishnan (Marimuthu and Balakrishnan, 2012) called a total labeling f as an *E-super vertex magic total labeling* if $f : E(G) \rightarrow \{1, 2, \dots, q\}$. They studied the *E-super vertex magicness* of *even regular graphs*. Most recently Wang and Zhang (Wang and Zhang, 2014) extended the results found in the article (Marimuthu and Balakrishnan, 2012).

Theorem 1.1. (Swaminathan and Jeyanthi, 2003)

A path P_n is *E-super vertex magic* if and only if n is odd and $n \geq 3$.

Theorem 1.2. (Swaminathan and Jeyanthi, 2003)

If a non-trivial graph G of order p and size q is *super vertex magic*, then the magic constant k is given by

$$k = q + \frac{p+1}{2} + \frac{q(q+1)}{p}.$$

Theorem 1.3. (Marimuthu and Balakrishnan, 2012)

Every tree T of even order is not *E-super vertex magic*.

In this article, we partially solve the following open problems given by Marimuthu, Suganya, Kalaivani and Balakrishnan (Marimuthu et al., 2015).

Open problem 1.4. Discuss the *E-super vertex magicness* of $B_{n,n-t}$, when $t \neq 2$.

Open problem 1.5. Find all *E-super vertex magic wounded suns*, $C_n^+ - me$, $m \neq 3$.

2. Solution to the Problems

Definition 2.1.

A broom $B_{n,d}$ is defined by attaching $n-d$ pendant edges with any one of the pendant vertices of the path P_d .

Theorem 2.2.

The broom $B_{n,n-1}$ is E -super vertex magic if and only if n is odd and $n \geq 3$.

Proof:

As the broom $B_{n,n-1}$ is isomorphic to a path P_n the results follow immediately from Theorem 1.1. \square

Theorem 2.3.

The broom $B_{n,n-t}$ is not E -super vertex magic for $n-t \geq 2, t \geq 3$.

Proof:

Let

$$V(B_{n,n-t}) = \{v_1, v_2, \dots, v_{n-t}, u_1, u_2, \dots, u_t\}$$

and let

$$E(B_{n,n-t}) = \{v_i v_{i+1}, 1 \leq i \leq n-t-1\} \cup \{v_{n-t} u_j, 1 \leq j \leq t\}.$$

Then, $B_{n,n-t}$ has $p = n$ vertices and $q = n-1$ edges.

As $B_{n,n-t}$ is a tree, according to Theorem 1.3, n can't be even.

Assume that n is odd.

Suppose $B_{n,n-t}$ is E -super vertex magic. Then, there exists a E -super vertex magic total labeling of $B_{n,n-t}$, say, f . According to Theorem 1.2 the magic constant k is given by

$$\begin{aligned} k &= q + \frac{p+1}{2} + \frac{q(q+1)}{p} \\ &= n-1 + \frac{n+1}{2} + \frac{(n-1)n}{n} \\ &= n-1 + \frac{n+1}{2} + n-1 \\ &= 2n-2 + \frac{n+1}{2} \\ &= \frac{5n-3}{2}. \end{aligned}$$

Now, we determine the smallest possible value of the sum

$$f(v_{n-t}) + \sum_{u \in N(v_{n-t})} f(v_{n-t}u) = f(v_{n-t}) + \sum_{j=1}^t f(v_{n-t}u_j) + f(v_{n-t-1}v_{n-t}).$$

The smallest possible value of the set $\{f(v_{n-t}u_j): 1 \leq j \leq t\}$ is

$$\{k - (p + q) + j - 1 = k - 2n + j : 1 \leq j \leq t\}.$$

Also, the smallest possible values of $f(v_{n-t-1}v_{n-t})$ and $f(v_{n-t})$ are 1 and n , respectively. Therefore, the smallest possible value of the above sum is:

$$\begin{aligned} n + \sum_{j=1}^t (k - 2n + j) + 1 &= n + 1 + tk - 2nt + (1 + 2 + \dots + t) \\ &= n + 1 + tk - 2nt + \frac{t(t+1)}{2} \\ &= n + 1 + t \left(\frac{5n-3}{2} \right) - 2nt + \frac{t^2+t}{2} \\ &= \frac{(t+2)n + (t-1)^2 + 1}{2}, t \geq 3 \\ &> \frac{5n-3}{2} \\ &= k. \end{aligned}$$

Therefore, the value of the sum at v_{n-t} is at least $n + \sum_{j=1}^t (k - 2n + j) + 1$ exceeds k , a contradiction. Hence, the result follows.

Note that

$$n + \sum_{j=1}^t (k - 2n + j) + 1$$

exceeds k by

$$\frac{(t-3)n + (t-1)^2 + 4}{2}.$$

Definition 2.4.

The sun graph C_n^+ is defined as follows:

$$V(C_n^+) = \{v_1, v_2, \dots, v_{2n}\}.$$

$$E(C_n^+) = \{v_i v_{i+1}, 1 \leq i \leq n-1\} \cup \{v_n v_1\} \cup \{v_i v_{n+i}, 1 \leq i \leq n\}.$$

The graph $C_n^+ - me, m \leq n$ is obtained by removing $\{v_{2n}, v_{2n-1}, \dots, v_{2n-m+1}\}$ and the edges adjacent to them from C_n^+ . This is referred as a *wounded sun*.

Theorem 2.5.

A wounded sun $C_n^+ - me$, $3 \neq m \leq n$, is E -super vertex magic if and only if m is odd and $m \neq 1$.

Proof:

Let $C_n^+ - me$ be a wounded sun, where m is a positive integer such that $3 \neq m \leq n$. Then,

$$V(C_n^+ - me) = \{v_1, v_2, \dots, v_n\} \cup \{v_{n+1}, v_{n+2}, \dots, v_{2n-m}\}$$

and

$$E(C_n^+ - me) = \{v_i v_{i+1}, 1 \leq i \leq n-1\} \cup \{v_n v_1\} \cup \{v_i v_{n+i}, 1 \leq i \leq n-m\}.$$

Thus, $C_n^+ - me$ has $p = 2n - m$ vertices and $q = 2n - m$ edges.

Assume that $C_n^+ - me$, $m \neq 3$ is E -super vertex magic. Then, by Theorem 1.3, the magic constant k is given by

$$\begin{aligned} k &= q + \frac{p+1}{2} + \frac{q(q+1)}{p} \\ &= p + \frac{p+1}{2} + p+1 \\ &= 2p+1 + \frac{p+1}{2} \\ &= 2(2n-m)+1 + \frac{2n-m+1}{2} \\ &= 4n-2m+1+n - \left(\frac{m-1}{2}\right) \\ &= 5n+1 - \left(\frac{5m-1}{2}\right), \end{aligned}$$

which is an integer only when m is odd.

Suppose $m = 1$. Then.

$$V(C_n^+ - e) = \{v_1, v_2, \dots, v_n\} \cup \{v_{n+1}, v_{n+2}, \dots, v_{2n-1}\}$$

and

$$E(C_n^+ - e) = \{v_i v_{i+1}, 1 \leq i \leq n-1\} \cup \{v_n v_1\} \cup \{v_i v_{n+i}, 1 \leq i \leq n-1\}.$$

Thus, $C_n^+ - e$ has $p = 2n - 1$ vertices and $q = 2n - 1$ edges. Since $C_n^+ - e$ is E -super vertex magic, it has a E -super vertex magic labeling. Therefore, for each vertex $v \in V(C_n^+ - e)$, the total weight $wt(v)$, (the sum of the labels of the vertex v and that of the edges incident to v) is a constant $k = 5n - 1$. Since there are $n-1$ pendent vertices and $n-1$ pendent edges, the only

possible set of the labels of the pendent vertices is $\{3n, 3n+1, \dots, 4n-2 = p+q\}$ and the only possible set of the labels of the pendent edges is $\{n+1, n+2, \dots, 2n-1 = q\}$. Therefore, at the vertex v_n , the largest possible label of the vertex v_n is $3n-1$ and the largest possible labels of the two edges incident to this vertex are $n-1$ and n . Thus, the largest possible value $wt(v_n)$ is

$$(3n-1) + (n-1) + (n) = 5n-2 < 5n-1 = k.$$

That is, the largest possible value of $wt(v_n)$ is less than k . Therefore, $wt(v_n) < k$, a contradiction. Hence, m is odd and $m \neq 1$. Conversely, assume that $m \leq n$ is odd, $m \neq 1$ and $m \neq 3$. Then, $m \geq 5$.

If $m = n$, then the wounded sun $C_n^+ - me$ is the cycle C_n . By Theorem 1.2, $C_n^+ - me$ is E -super vertex magic.

Now let $5 \leq m \leq n-1$. Define a total labeling

$$f : V \cup E \rightarrow \{1, 2, \dots, 4n-2m\}$$

as follows:

$$\begin{aligned} f(v_1) &= 2n-1, \\ f(v_n) &= 3n-m, \\ f(v_{n-i}) &= 2n-m+i, 1 \leq i \leq m-2, \\ f(v_{i+1}) &= 3n-m-i, 1 \leq i \leq n-m, \\ f(v_{n+i}) &= 3n-m+i, 1 \leq i \leq n-m, \\ f(v_i v_{i+1}) &= i, 1 \leq i \leq n-m, \\ f(v_1 v_n) &= n-m+1, \\ f(v_i v_{n+i}) &= 2n-i-3\left(\frac{m-1}{2}\right), 1 \leq i \leq n-m, \\ f(v_{n-2i} v_{n-2i+1}) &= 2n-m+1-i, 1 \leq i \leq \frac{m-1}{2}, \\ f(v_{n-2i+1} v_{n-2i+2}) &= n-\left(\frac{m-3}{2}\right)-i, 1 \leq i \leq \frac{m-1}{2}. \end{aligned}$$

From the above labeling, we have,

$$V(C_n^+ - me) = \{v_1\} \cup \{v_n\} \cup \{v_{n-i}, 1 \leq i \leq m-2\} \cup \{v_{i+1}, 1 \leq i \leq n-m\} \cup \{v_{n+i}, 1 \leq i \leq n-m\}.$$

Now,

$$\begin{aligned} f(v_1) + \sum_{u \in N(v_1)} f(v_1 u) &= f(v_1) + f(v_1 v_2) + f(v_1 v_n) + f(v_1 v_{n+1}) \\ &= (2n-1) + (1) + (n-m+1) + \left(2n-1-3\left(\frac{m-1}{2}\right)\right) \\ &= 5n - \frac{5m}{2} + \frac{3}{2} = 5n + \frac{3-5m}{2}. \end{aligned}$$

$$\begin{aligned}
f(v_n) + \sum_{u \in N(v_n)} f(v_n u) &= f(v_n) + f(v_n v_1) + f(v_n v_{n-1}) \\
&= (3n - m) + (n - m + 1) + \left(n - \left(\frac{m-3}{2} \right) - 1 \right) \\
&= 5n + \frac{3-5m}{2}.
\end{aligned}$$

$$\begin{aligned}
f(v_{n-i}) + \sum_{u \in N(v_{n-i})} f(v_{n-i} u) &= f(v_{n-i}) + f(v_{n-i} v_{n-i-1}) + f(v_{n-i} v_{n-i+1}) \\
&= (2n - m + i) + \left(2n - m + 1 - \left(\frac{i+1}{2} \right) \right) + \left(n - \left(\frac{m-3}{2} \right) - \left(\frac{i+1}{2} \right) \right) \\
&= 5n + \frac{3-5m}{2}, 1 \leq i \leq m-2.
\end{aligned}$$

$$\begin{aligned}
f(v_{i+1}) + \sum_{u \in N(v_{i+1})} f(v_{i+1} u) &= f(v_{i+1}) + f(v_{i+1} v_i) + f(v_{i+1} v_{i+2}) + f(v_{i+1} v_{n+i+1}) \\
&= (3n - m - i) + (i) + (i+1) + \left(2n - m - (i+1) - \left(\frac{m-3}{2} \right) \right) \\
&= 5n + \frac{3-5m}{2}, 1 \leq i \leq n-m.
\end{aligned}$$

$$\begin{aligned}
f(v_{n+i}) + \sum_{u \in N(v_{n+i})} f(v_{n+i} u) &= f(v_{n+i}) + f(v_i v_{n+i}) \\
&= (3n - m + i) + \left(2n - i - 3 \left(\frac{m-1}{2} \right) \right) \\
&= 5n + \frac{3-5m}{2}, 1 \leq i \leq n-m.
\end{aligned}$$

Therefore,

$$\text{for every } v \in V(C_n^+ - me), f(v) + \sum_{u \in N(v)} f(uv) = 5n + \frac{3-5m}{2}.$$

Hence, f is an E -super vertex magic labeling of $V(C_n^+ - me)$ with magic constant

$$k = 5n + \frac{3-5m}{2},$$

where m is odd, $m \neq 1$ and $m \neq 3$.

An illustration for Theorem 2.5 is given in Figure 1.




Figure 1. An E -super vertex magic labeling of $C_{15}^+ - 9e$

3. Conclusion and Scope

In this article, we have solved two open problems given by Marimuthu, Suganya, Kalaivani and Balakrishnan (Marimuthu et al., 2015). There is another open problem for further investigation in the same paper which is given as follows:

Open problem 1. Characterize all E -super vertex magic trees of odd order.

Acknowledgment

The authors would like to express their gratitude to the anonymous referees for their kind suggestions and useful comments on the original manuscript, which resulted in this final version.

REFERENCES

Gallian, J. A. (2014). A dynamic survey of graph labeling, *Electron. J. Combin.* 16: #DS6.
 Harary, F. (1969). *Graph Theory*, Addison- Wesley.
 MacDougall, J. A, Miller, M, Slamin, Wallis, W. D. (2002). Vertex magic total labelings of graphs, *Util. Math.*, 61: 3-21.
 MacDougall, J. A, Miller, M, Sugeng, K. A. (2004). Super vertex magic total labelings of graphs, Proc. of the 16th Australian Workshop on Combinatorial Algorithms, pp. 222-229.
 Marimuthu, G and Balakrishnan, M. (2012). E -super vertex magic labelings of graphs, *Discrete Appl. Math.*, 160: 1766-1774.

- Marimuthu, G, Suganya, B, Kalaivani, S, Balakrishnan, M. (2015). E-super vertex magic labelling of graphs and some open problems, *Appl. Appl. Math.*, 10 (1): 536-543.
- Marr, A. M and Wallis, W. D. (2013). *Magic Graphs*, Birkhauser - Springer, Second edition, Boston.
- Sedláček, J., Problem 27. (1963). *Theory of Graphs and its Applications*, Proc. Symposium, pp. 163-167.
- Swaminathan, V and Jeyanthi, P. (2003). Super vertex magic labelling, *Indian J. Pure Appl. Math.*, 34 (6): 935-939.
- Wang and Zhang, (2014). Note on E-super vertex magic graphs, *Discrete Appl. Math.*, 178: 160-162.