



12-2018

Some results on Lupaş (p; q)-Bernstein operators and its limit form

Asif Khan
Aligarh Muslim University

Vinita Sharma
Aligarh Muslim University

Follow this and additional works at: <https://digitalcommons.pvamu.edu/aam>



Part of the [Numerical Analysis and Computation Commons](#)

Recommended Citation

Khan, Asif and Sharma, Vinita (2018). Some results on Lupaş (p; q)-Bernstein operators and its limit form, *Applications and Applied Mathematics: An International Journal (AAM)*, Vol. 13, Iss. 2, Article 33.
Available at: <https://digitalcommons.pvamu.edu/aam/vol13/iss2/33>

This Article is brought to you for free and open access by Digital Commons @PVAMU. It has been accepted for inclusion in *Applications and Applied Mathematics: An International Journal (AAM)* by an authorized editor of Digital Commons @PVAMU. For more information, please contact hvkoshy@pvamu.edu.



Some results on Lupaş (p, q) -Bernstein operators and its limit form

¹ Asif Khan and ² Vinita Sharma

Department of Mathematics
 Aligarh Muslim University
 Aligarh-202002, India

asifjnu07@gmail.com; vinita.sha23@gmail.com

Received: September 22, 2017; Accepted: April 19, 2018

Abstract

In this paper, statistical approximation properties of Lupaş (p, q) -analogue of Bernstein operators are studied. Rate of statistical convergence by means of modulus of continuity and Lipschitz type maximal functions has been investigated. Further Limit Lupaş (p, q) -Bernstein operators are defined and some results are investigated.

Keywords: (p, q) -integers; Lupaş (p, q) -Bernstein operators; Limit Lupaş (p, q) -Bernstein operators; Positive linear operators; Korovkin type approximation; Statistical convergence.

2010 MSC No.: 41A10, 41A25, 41A36, 40A30

1. Introduction

Mursaleen et al. (2015) introduced the concept of (p, q) -calculus (post quantum calculus) in approximation theory and constructed (p, q) -analogue of Bernstein operators (a generalisation of Phillips q -Bernstein polynomials) based on (p, q) -integers. They also introduced and studied approximation properties for (p, q) -analogue of Bernstein-Stancu operators (2015), Bleimann-Butzer-Hahn operators defined by (p, q) -integers (2016), (p, q) -analogue of Bernstein-Kantorovich operator (2016), (p, q) -analogue of Lorentz polynomials on a compact disk in (2016) etc.

The concept of post quantum calculus in Approximation theory and Computer aided geometric design got attention when Khalid et al. (2015) further used basis of (p, q) -analogue of Bernstein operators to study (p, q) -Bézier curves and surfaces which generalises q -Bézier curves and surfaces [Phillips (2000); Oruk and Phillips (2003); Phillips (2003)].

Khalid and Lobiyal (2017) also introduced Lupaş (p, q) -Bernstein operators (a generalisation of well known rational Lupaş q -Bernstein operators). They initially put all these results on arXiv (2015), later on it get published in 2017 in *Journal of Computational and Applied Mathematics*. They constructed Lupaş type (p, q) -Bézier curves and surfaces based on (p, q) -integers which is further generalization of q -Bézier curves and surfaces. In this paper, they have given a nice application of the extra parameter p in constructing Bézier curves and surfaces which provides more flexibility in controlling the shapes of curves and surfaces. These (p, q) -Bézier curves and surfaces mimics the control polygon much better than q -Bézier curves [Hana et al. (2014)] and classical Bézier curves. After this initiation, many researchers started working in this area.

For similar works based on (p, q) -integers, one can refer [Acar (2016); Acar et al. (2016); Ansari and Karaisa (2017); Cai and Zhou (2016); Kadak (2016); Kadak (2017); Mishra and Pandey (2017); Wafi and Rao (2016); Mursaleen et al. (2016); Mishra and Pandey (2016); Wafi and Rao (2017); Wafi and Rao (2019)].

For works related to constructive Approximation theory and Bézier curves, one can see [Agarwal et al. (2013); Aral et al. (2013); Dalmanoglu et al. (2007); Khalid et al. (2015); Mursaleen and Khan (2013); Mursaleen et al. (2017); Devdhara and Mishra (2017); Gairola et al. (2016); Pandey (2016); Mishra and Pandey (2017); Kac and Cheung (2002); Farouki and Rajan (1988); Hana et al. (2014); Oruk and Phillips (2003); Rababah and Manna (2011); Korovkin (1960); Ostrovska(2006); Wafi and Rao (2016); Wafi and Rao (2019)].

In 1912, Bernstein [Bernstein (1912, 1913)] introduced the operators $B_n : C[0, 1] \rightarrow C[0, 1]$ defined for any $n \in \mathbb{N}$ and for any function $f \in C[0, 1]$.

$$B_n(f; x) = \sum_{k=0}^n \binom{n}{k} x^k (1-x)^{n-k} f\left(\frac{k}{n}\right), \quad x \in [0, 1]. \quad (1)$$

and named it Bernstein polynomials to prove the constructive Weierstrass theorem.

In 1987, Lupaş [Lupaş (1987)] introduced the first q -analogue of Bernstein operators (rational) as follows

$$L_{n,q}(f; x) = \sum_{k=0}^n \frac{f\left(\frac{[k]_q}{[n]_q}\right) \begin{bmatrix} n \\ k \end{bmatrix}_q q^{\frac{k(k-1)}{2}} x^k (1-x)^{n-k}}{\prod_{j=1}^n \{(1-x) + q^{j-1}x\}} \quad (2)$$

and investigated its approximating and shape-preserving properties.

Let us recall certain notations of (p, q) -calculus. For any $p > 0$ and $q > 0$, the (p, q) integers $[n]_{p,q}$ are defined by

$$[n]_{p,q} = p^{n-1} + p^{n-2}q + p^{n-3}q^2 + \dots + pq^{n-2} + q^{n-1} = \begin{cases} \frac{p^n - q^n}{p - q}, & \text{when } p \neq q \neq 1, \\ n p^{n-1}, & \text{when } p = q \neq 1, \\ [n]_q, & \text{when } p = 1, \\ n, & \text{when } p = q = 1, \end{cases}$$

where $[n]_q$ denotes the q -integers and $n = 0, 1, 2, \dots$.

The formula for (p, q) -binomial expansion is as follows:

$$(ax + by)_{p,q}^n := \sum_{k=0}^n p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_{p,q} a^{n-k} b^k x^{n-k} y^k.$$

For $a = b = 1$,

$$(x + y)_{p,q}^n := \sum_{k=0}^n p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_{p,q} x^{n-k} y^k.$$

On using,

$$(x + y)_{p,q}^n = (x + y)(px + qy)(p^2x + q^2y) \dots (p^{n-1}x + q^{n-1}y),$$

one can write,

$$(x)_{p,q}^n = (x)(px)(p^2x) \dots (p^{n-1}x) = p^{\frac{n(n-1)}{2}} x^n,$$

which implies

$$(1)_{p,q}^n = (1)(p)(p^2) \dots (p^{n-1}) = p^{\frac{n(n-1)}{2}}.$$

From (p, q) -binomial expansion, or by using induction on n , one can easily verify that

$$\sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_{p,q} p^{\frac{k(k-1)}{2}} x^k \prod_{s=0}^{n-k-1} (p^s - q^s x) = p^{\frac{n(n-1)}{2}}, \quad x \in [0, 1]. \tag{3}$$

Details on (p, q) -calculus can be found in [Hounkonnou, Désiré and Kyemba (2013); Mursaleen et al. (2015); Jagannathan and Srinivasa (2005)].

The (p, q) -Bernstein operators introduced by Mursaleen et al. for $0 < q < p \leq 1$ in [Mursaleen et al. (2015, 2016)] are as follows:

$$B_{n,p,q}(f; x) = \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_{p,q} p^{\frac{k(k-1)}{2}} x^k \prod_{s=0}^{n-k-1} (p^s - q^s x) f\left(\frac{[k]_{p,q}}{p^{k-n}[n]_{p,q}}\right), \tag{4}$$

for $x \in [0, 1]$.

Note when $p = 1$, (p, q) -Bernstein operators given by (4) turns out to be Phillips q -Bernstein operators.

Also, we have (p, q) -analogue of Euler’s identity as:

$$\begin{aligned} (1-x)_{p,q}^n &= \prod_{s=0}^{n-1} (p^s - q^s x) = (1-x)(p-qx)(p^2-q^2x)\dots(p^{n-1}-q^{n-1}x) \\ &= \sum_{k=0}^n (-1)^k p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_{p,q} x^k. \end{aligned}$$

Another needed formulae, which can be easily derived from Euler’s identity for $|\frac{q}{p}| < 1$ is :

$$\sum_{k=0}^{\infty} \frac{q^{\frac{k(k-1)}{2}} x^k}{(p-q)^k [k]_{p,q}!} = \prod_{j=0}^{\infty} \left\{ 1 + \left(\frac{q}{p}\right)^{j-1} x \right\}.$$

Khalid and Lobiyal (2017) introduced Lupaş type (p, q) -analogue of the Bernstein functions (rational) as follows:

For any $p > 0$ and $q > 0$,

$$b_{p,q}^{k,n}(t) = \frac{\begin{bmatrix} n \\ k \end{bmatrix}_{p,q} p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} t^k (1-t)^{n-k}}{\prod_{j=1}^n \{p^{j-1}(1-t) + q^{j-1}t\}}, \tag{5}$$

where $b_{p,q}^{0,n}(t), b_{p,q}^{1,n}(t), \dots, b_{p,q}^{n,n}(t)$ are the (p, q) -analogue of the Lupaş q -Bernstein functions of degree n on the interval $[0, 1]$.

$$\sum_{k=0}^n b_{p,q}^{k,n}(t) = 1. \tag{6}$$

When $p = 1$, Lupaş (p, q) -Bernstein functions turns out to be Lupaş q -Bernstein functions as given in [Hana et al. (2014)], whereas when $p = q = 1$, Lupaş (p, q) -Bernstein functions turns out to be classical Lupaş Bernstein functions.

Similarly [Khalid and Lobiyal (2017)], Lupaş (p, q) -Bernstein operators are defined as follows:

For any $p > 0$ and $q > 0$, the linear operators $L_{p,q}^n : C[0, 1] \rightarrow C[0, 1]$,

$$L_{p,q}^n(f; x) = \sum_{k=0}^n \frac{f\left(\frac{p^{n-k} [k]_{p,q}}{[n]_{p,q}}\right) \begin{bmatrix} n \\ k \end{bmatrix}_{p,q} p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} x^k (1-x)^{n-k}}{\prod_{j=1}^n \{p^{j-1}(1-x) + q^{j-1}x\}}, \tag{7}$$

is (p, q) -analogue of Lupaş Bernstein operators.

Again when $p = 1$, Lupaş (p, q) -Bernstein operators turns out to be Lupaş q -Bernstein operators as given in [Mahmudov and Sabancigil (2010); Ostrovska (2006)].

When $p = q = 1$, Lupaş (p, q) -Bernstein operators turns out to be classical Bernstein operators.

Now we recall that (p, q) -analogue of Lupaş operator reproduces linear and constant functions [Khalid and Lobiyal (2017)] are as follows:

Some auxillary results:

$$(1) L_{p,q}^n(1, \frac{u}{u+1}) = 1,$$

$$(2) L_{p,q}^n(t, \frac{u}{u+1}) = \frac{u}{u+1},$$

$$(3) L_{p,q}^n(t^2, \frac{u}{u+1}) = \frac{u}{u+1} \frac{p^{n-1}}{[n]_{p,q}} + \frac{qu}{u+1} (\frac{qu}{p+qu}) \frac{[n-1]_{p,q}}{[n]_{p,q}},$$

or equivalently for $x = \frac{u}{u+1}$,

(1)

$$L_{p,q}^n(1, x) = 1, \quad (8)$$

(2)

$$L_{p,q}^n(t, x) = x, \quad (9)$$

(3)

$$L_{p,q}^n(t^2, x) = \frac{p^{n-1}x}{[n]_{p,q}} + \frac{q^2 x^2}{p(1-x) + qx} \frac{[n-1]_{p,q}}{[n]_{p,q}}. \quad (10)$$

Remark.

For $q \in (0, 1)$ and $p \in (q, 1]$, it is obvious that $\lim_{n \rightarrow \infty} [n]_{p,q} = 0$ or $\frac{1}{1-q}$. In order to reach to convergence results of the operator $L_{p,q}^n(f, x)$, we take a sequence $q_n \in (0, 1)$ and $p_n \in (q, 1]$ and such that $\lim_{n \rightarrow \infty} p_n = 1$, $\lim_{n \rightarrow \infty} q_n = 1$ and $\lim_{n \rightarrow \infty} q_n^n = 1$. so we get $\lim_{n \rightarrow \infty} [n]_{p_n, q_n} = \infty$.

2. Statistical approximation

The statistical version of Korovkin theorem for sequence of positive linear operators has been given by Gadjiev and Orhan [Gadjiev and Orhan (2002)].

Also this type approximation has been studied by many authors, the reader may refer to [Belen and Mohiuddine (2013); Fast (1951); Mursaleen and Khan (2013); Mahmudov and Sabancigil (2010); Edely et al. (2010); Dogru and Kanat (2012); Ostrovska(2006); Braha et al. (2014); Kadak (2016); Kadak et al. (2017); Kadak (2017)].

Let K be a subset of the set \mathbb{N} of natural numbers. Then, the asymptotic density $\delta(K)$ of K is defined as $\delta(K) = \lim_n \frac{1}{n} |\{k \leq n : k \in K\}|$ and $|\cdot|$ represents the cardinality of the enclosed set.

A sequence $x = (x_k)$ said to be statistically convergent to the number L if for each $\varepsilon > 0$, the set $K(\varepsilon) = \{k \leq n : |x_k - L| > \varepsilon\}$ has asymptotic density zero i.e.,

$$\lim_n \frac{1}{n} |\{k \leq n : |x_k - L| \geq \varepsilon\}| = 0.$$

In this case, we write $st - \lim x = L$.

Let us recall the following theorem:

Theorem 2.1.

Let S_n be the sequence of linear positive operators from $C[0, 1]$ to $C[0, 1]$ satisfies the conditions

$st - \lim_n \|S_{n,p,q}(t^\nu; x) - (x)^\nu\|_{C[0,1]} = 0$ for $\nu = 0, 1, 2$, then for any function $f \in C[0, 1]$,

$$st - \lim_n \|S_{n,p,q}(f) - f\|_{C[0,1]} = 0.$$

2.1. Korovkin Type statistical approximation properties

The main aim of this paper is to obtain the Korovkin type statistical approximation properties of operators defined in (7) with the help of Theorem (2.1).

Theorem 2.2.

let $L_{p,q}^n$ be the sequence of operators and the sequence $p = p_n$ and $q = q_n$ satisfying Remark 1.1, then for any function $f \in C[0, 1]$

$$st - \lim_n \|L_{p,q}^n(f, \cdot) - f\| = 0.$$

Proof:

Clearly for $\nu = 0$,

$$L_{p,q}^n(1, x) = 1,$$

which implies

$$st - \lim_n \|L_{p,q}^n(1; x) - 1\| = 0.$$

For $\nu = 1$ we get

$$st - \lim_n \|L_{p,q}^n(t; x) - x\| = 0,$$

Lastly for $\nu = 2$, we have

$$\|L_{p,q}^n(t^2; x) - x^2\| \leq \left| \frac{q^2 [n-1]_{p,q}}{p [n]_{p,q}} - 1 \right| + \left| \frac{p^{n-1}}{[n]_{p,q}} \right|.$$

If we choose

$$\alpha_n = \frac{q^2 [n-1]_{p,q}}{p [n]_{p,q}} - 1,$$

$$\beta_n = \frac{p^{n-1}}{[n]_{p,q}},$$

then

$$st - \lim_n \alpha_n = st - \lim_n \beta_n = 0.$$

Now given $\epsilon > 0$, we define the following three sets:

$$U = \{n : \|L_{p,q}^n(t^2; x) - x^2\| \geq \epsilon\}$$

$$U_1 = \{n : \alpha_n \geq \frac{\epsilon}{2}\}$$

$$U_2 = \{n : \beta_n \geq \frac{\epsilon}{2}\}$$

It is obvious that $U \subseteq U_1 \cup U_2$. Thus, we obtain

$$\begin{aligned} & \delta\{K \leq n : \|S_{n,p,q}(t^2; x) - x^2\| \geq \epsilon\} \\ & \leq \delta\{K \leq n : \alpha_n \geq \frac{\epsilon}{2}\} + \delta\{K \leq n : \beta_n \geq \frac{\epsilon}{2}\}. \end{aligned}$$

So the right hand side of the inequalities is zero.

Then,

$$st - \lim_n \|L_{p,q}^n(t^2; x) - x^2\| = 0.$$

This gives the proof.

3. The Rates of Convergence

In this section, we compute the rates of convergence of the operators $L_{p,q}^n$ to the function f by means of modulus continuity.

The modulus of continuity for the space of function $f \in C[0, 1]$ denotes by $w(f; \delta)$ is defined to be

$$w(f; \delta) = \sup_{x,t \in C[0,1], |t-x| < \delta} |f(t) - f(x)|,$$

where $w(f; \delta)$ satisfies the following conditions: For all $f \in C[0, 1]$,

$$\lim_{\delta \rightarrow 0} w(f; \delta) = 0 \tag{11}$$

and

$$|f(t) - f(x)| \leq w(f; \delta) \left(\frac{|t-x|}{\delta} + 1 \right). \quad (12)$$

For every $f \in C[0, 1]$ and $\delta > 0$, Lupas obtained the following rate of convergence for the operators

$$|L_{p,q}^n f(x) - f(x)| \leq w(f; \delta) \left\{ 1 + \frac{1}{\delta} \sqrt{\frac{x(1-x)}{[n]_{p,q}}} \right\}. \quad (13)$$

Theorem 3.1.

Let $0 < q_n < p_n \leq 1$ such that $\lim_{n \rightarrow \infty} p_n = 1$, $\lim_{n \rightarrow \infty} q_n = 1$ and $\lim_{n \rightarrow \infty} p_n^n = 1$, $\lim_{n \rightarrow \infty} q_n^n = 1$. Then for each $f \in C[0, 1]$, $L_{p,q}^n(f; x)$ and $\delta_n \geq 0$, we have

$\|L_{p,q}^n f(x) - f(x)\|_{C[0,1]} \leq 2\omega(f, \delta_n)$, where

$$\delta_n = \left(\frac{p^{n-1}}{[n]_{p,q}} + \frac{q^2}{p} \frac{[n-1]_{p,q}}{[n]_{p,q}} - 1 \right)^{\frac{1}{2}}. \quad (14)$$

Proof:

From (8), (9) and (10), we have

$$L_{p,q}^n((t-x)^2; x) = \frac{p^{n-1}x}{[n]_{p,q}} + \frac{q^2 x^2}{p(1-x) + qx} \frac{[n-1]_{p,q}}{[n]_{p,q}} - x^2, \quad (15)$$

$$\|L_{p,q}^n(f; x) - f(x)\|_{C[0,1]} \leq 2\omega(f, \delta) \left\{ 1 + \frac{1}{\delta} (L_{p,q}^n((t-x)^2; x)^{\frac{1}{2}}) \right\},$$

$$|L_{p,q}^n(t; x) - f(x)| \leq w(f; \delta) \left\{ 1 + \frac{1}{\delta_n} [(L_{p,q}^n(t-x)^2; x)^{\frac{1}{2}}] \right\}.$$

Then, we get

$$\|L_{p,q}^n(f; x) - f(x)\|_{C[0,1]} \leq \omega(f; \delta) \left\{ 1 + \frac{1}{\delta} \left(\frac{p^{n-1}}{[n]_{p,q}} + \frac{q^2}{p} \frac{[n-1]_{p,q}}{[n]_{p,q}} - 1 \right)^{\frac{1}{2}} \right\}. \quad (16)$$

If we choose

$$\delta_n = \left(\frac{p^{n-1}}{[n]_{p,q}} + \frac{q^2}{p} \frac{[n-1]_{p,q}}{[n]_{p,q}} - 1 \right)^{\frac{1}{2}}, \quad (17)$$

then we have

$$\|L_{p,q}^n(f; x) - f(x)\|_{C[0,1]} \leq 2\omega(f; \delta_n).$$

So we have the desired result.

4. Limit Lupaş (p, q)-Bernstein functions and operators

For $p > q > 0$, Limit Lupaş (p, q)-Bernstein functions are defined as follows:

$$b_{p,q}^{k,\infty}(u) = \frac{q^{\frac{k(k-1)}{2}} u^k}{(p-q)^k [k]_{p,q}! \prod_{j=0}^{\infty} \left\{ 1 + \left(\frac{q}{p}\right)^{j-1} u \right\}}. \quad (18)$$

From Eulers identity, we have

$$\sum_{k=0}^{\infty} b_{p,q}^{k,\infty}(u) = 1. \quad (19)$$

Further, we define Limit Lupaş (p, q)-Bernstein operators $L_{p,q}^{\infty}$ on $C[0, 1]$ as

$$L_{p,q}^n = \begin{cases} \sum_{k=0}^{\infty} f\left(1 - \left(\frac{q}{p}\right)^k\right) b_{p,q}^{k,\infty}, & \text{if } x \in [0, 1), \\ f(1), & \text{if } x = 1. \end{cases} \quad (20)$$

which is (p, q)-analogue of the limit Lupaş Bernstein operators.

Note that the function $L_{p,q}^{\infty}(f; x)$ is well-defined on $C[0, 1]$ whenever $f(x)$ is bounded on $[0, 1]$.

It follows directly from the definition that operators $L_{p,q}^{\infty}(f, t)$ posses the end point interpolation property, that is

$$L_{p,q}^{\infty}(f, 0) = f(0), \quad L_{p,q}^{\infty}(f, 1) = f(1). \quad (21)$$

Corollary 1.

Let $f \in C[0, 1]$, and $g(x) = f(1 - x)$. Then, for any $p, q > 0$, $L_{p,q}^n(f; t)$

$$L_{p,q}^n(f; t) = L_{\frac{1}{q}, \frac{1}{p}}^n(g; 1 - t), \quad \text{for } t \in [0, 1]. \quad (22)$$

Corollary 2.

Let $p, q \neq 1$ be fixed, $f \in C[0, 1]$, and $g(x) = f(1 - x)$. Then, for $x \in [0, 1]$, $L_{p,q}^n(f; t)$ converges uniformly to $L_{p,q}^{\infty}(f; t)$ for any $p > 0$ being fixed,

where

$$L_{p,q}^{\infty}(f; t) = \begin{cases} L_{p,q}^{\infty}(f; t), & \text{if } 0 < q < p < 1, \\ L_{\frac{1}{q}, \frac{1}{p}}^{\infty}(f; t), & \text{if } p > q > 1. \end{cases} \quad (23)$$

Theorem 4.1.

Let $p, q \in (0, 1)$. Then, for any $f \in C[0, 1]$, $L_{p,q}^n(f; x)$ uniformly converges to $L_{p,q}^\infty(f; x)$ for $x \in [0, 1]$.

Remark:

It is worth mentioning that the results above admit a probabilistic interpretation. Indeed, since $b_{p,q}^{k,n} \geq 0$ for $x \in [0, 1]$ and by (6) $\sum_{k=0}^n b_{p,q}^{k,n} = 1$, we may consider a sequence of discrete random variables $\{X_n\}$ with the distribution ρ_n defined by

$$P\left\{X_n = \frac{p^{n-k}[k]_{p,q}}{[n]_{p,q}}\right\} = b_{p,q}^{k,n}(x), \quad k = 0, 1, 2, \dots, n.$$

Then, $L_{p,q}^n(f; x) = E[f(X_n)]$. For $p, q \in (0, 1)$ consider a discrete random variable X_∞ with the distribution ρ defined by

$$\begin{cases} P\{X_\infty = 1 - (\frac{q}{p})^k\} = b_{p,q}^{k,\infty}(x), & \text{if } x \in [0, 1), \\ P\{X_\infty = 1\} = 1, & \text{if } x = 1. \end{cases}$$

The distribution is well-defined due to (19) and the fact that all $b_{p,q}^{k,\infty}(x) \geq 0$ on $[0, 1)$.

Then, $L_{p,q}^\infty(f; x) = E[f(X_\infty)]$ and Theorem 5.1 means that ρ is a limit distribution for the sequence ρ_n .

Proof:

From (21) it suffices to prove that $L_{p,q}^n(f; x)$ uniformly converges to $L_{p,q}^\infty(f; x)$ for $x \in [0, 1]$.

Consider

$$\Delta = |L_{p,q}^n(f; x) - L_{p,q}^\infty(f; x)|,$$

for $x \in [0, 1]$

$$\left| \sum_{k=0}^n f\left(\frac{p^{n-k}[k]_{p,q}}{[n]_{p,q}}\right) b_{p,q}^{k,n}(t) - \sum_{k=0}^{\infty} f\left(1 - \left(\frac{q}{p}\right)^k\right) b_{p,q}^{k,\infty}(t) \right|.$$

Let $\epsilon > 0$ be given, we choose $a \in (0, 1)$ in such a way that $\omega_f(1-a) < \frac{\epsilon}{3}$, where ω_f denotes the modulus of continuity of f .

Let R be a positive integer satisfying the condition $1 - \frac{q^{R+1}}{p} \geq a$. Then, $\frac{p^{n-k}[k]_{p,q}}{[n]_{p,q}} \geq a$, for all $K \geq R + 1$.

Using (6) and(19), we get

$$\begin{aligned} \Delta &= \left| \sum_{k=0}^n \left(f\left(\frac{p^{n-k}[k]_{p,q}}{[n]_{p,q}}\right) - f(1) \right) b_{p,q}^{k,n}(t) - \sum_{k=0}^n \left(f\left(1 - \left(\frac{q}{p}\right)^k\right) - f(1) \right) b_{p,q}^{k,\infty}(t) \right| \\ &\leq \left| \sum_{k=0}^R \left(f\left(\frac{p^{n-k}[k]_{p,q}}{[n]_{p,q}}\right) - f(1) \right) b_{p,q}^{k,n}(t) - \sum_{k=0}^R \left(f\left(1 - \left(\frac{q}{p}\right)^k\right) - f(1) \right) b_{p,q}^{k,\infty}(t) \right| \\ &+ \sum_{k=R+1}^n \left| f\left(\frac{p^{n-k}[k]_{p,q}}{[n]_{p,q}}\right) - f(1) \right| b_{p,q}^{k,n}(t) + \sum_{k=R+1}^{\infty} \left| f\left(1 - \left(\frac{q}{p}\right)^k\right) - f(1) \right| b_{p,q}^{k,\infty}(t) \\ &= \delta_1 + \delta_2 + \delta_3. \end{aligned} \tag{24}$$

Since $f\left(\frac{p^{n-k}[k]_{p,q}}{[n]_{p,q}}\right) \rightarrow f\left(1 - \left(\frac{q}{p}\right)^k\right)$ as $n \rightarrow \infty$, due to the fact that $b_{p,q}^{k,n}(t) \geq 0$ for $x \in [0, 1]$, we get the following estimate for δ_2 :

$$\begin{aligned} \delta_2 &\leq \omega_f(1 - a) \sum_{k=R+1}^n b_{p,q}^{k,n} \leq \omega_f(1 - a) \sum_{k=0}^n b_{p,q}^{k,n} \\ &= \omega_f(1 - a) < \frac{\epsilon}{3}, \end{aligned}$$

because of (6). Similarly, using (19) we get $\delta_3 \leq \omega_f(1 - a) < \frac{\epsilon}{3}$. Thus, $\Delta < \epsilon$ for n large enough.

Theorem 4.2.

Let $p > q > 0$ and $q \neq 1$ be fixed and $f \in C[0, 1]$. Then, $L_{p,q}^\infty(f; x) = f(x)$ for all $x \in [0, 1]$ if and only if $f(x) = ax + b$ for some $a, b \in R$.

Proof:

If $f(x) = ax + b$, then $L_{p,q}^\infty(f; x) = ax + b$ for all $n = 1, 2, \dots$ and, therefore,

$$L_{p,q}^\infty(f; x) = \lim_{n \rightarrow \infty} L_{p,q}^n(f; x) = ax + b = f(x).$$

Now, suppose that $f \in C[0, 1]$ and $L_{p,q}^\infty(f; x) = f(x)$, for all $x \in [0, 1]$.

Due to Theorem 5.1 it suffices to prove the statement in the case $q \in (0, 1)$.

Consider the function

$$\varphi(x) = f(x) - (f(1) - f(0))x.$$

Obviously, $\varphi(0) = \varphi(1)$ and $L_{p,q}^\infty(f; x) = \varphi(x)$. We will prove that $\varphi(x) = \varphi(0) = \varphi(1)$, for all $x \in [0, 1]$. Let

$$M = \max_{x \in [0,1]} \varphi(x).$$

Assume that $M > \varphi(1)$. Then, $M = \varphi(z)$ for some $z \in (0, 1)$ and $\varphi(1 - (\frac{q}{p})^k) < M$, for $k > N_s 0$. Using Eulers identity and positivity of $L_{p,q}^\infty(f; x)$, $k = 0, 1, 2 \dots$, on $(0, 1)$, we get

$$M = \varphi(z) = \sum_{k=0}^{\infty} \varphi(1 - (\frac{q}{p})^k) L_{p,q}^\infty(z) < M.$$

The contradiction shows that $\varphi(x) \leq \varphi(1)$, for all $x \in [0, 1]$. Likewise, we prove that $\varphi(x) \geq \varphi(1)$, for all $x \in [0, 1]$. Thus, $\varphi(x) \equiv \varphi(1) \equiv b$, for some $b \in R$, and finally $f(x) = ax + b$ with $a = f(1) - f(0)$.

5. Conclusion:

In real life applications, not every sequences are convergent, so in order to evaluate that to what number most of the terms of sequence are approaching, for this the role of statistical limit is very useful. Thus statistical approximation properties of Lupaş (p, q) -analogue of Bernstein operators has been studied. Rate of statistical convergence by means of modulus of continuity and Lipschitz type maximal functions has been investigated. Limit Lupaş (p, q) -Bernstein operators are defined and some results are investigated. The extra parameter p has applications in terms of flexibility in approximation.

REFERENCES

- Acar, T. (2016). (p, q) -Generalization of Szász-Mirakyan operators. *Math. Methods Appl. Sci.* 39(10), 2685-2695.
- Acar, T. Aral, A. Mohiuddine, S.A. (2016). On Kantorovich modification of (p, q) -Baskakov operators, *Journal of Inequalities and Applications*, 98: DOI:10.1186/s13660-016-1045-9.
- Agarwal, P. N. Gupta, V., Kumar, S. (2013). On q -analogue of Bernstein-Schurer-Stancu operators, *Appl. Math. Comput.*, 219, 7754-7764.
- Ansari, K. J. Karaisa, A. (2017). On the approximation by Chlodowsky type generalization of (p, q) -Bernstein operators, *Int. J. Nonlinear Anal. Appl.* 8, No. 2, 181-200.
- Aral, A. Gupta, V. Agarwal, R.P. (2013). Applications of q -Calculus in Operator Theory, *Springer*, New York.
- Belen, C. Mohiuddine, S. (2013). Generalized weighted statistical convergence and application. *Applied Mathematics and Computation*, 219(18): 9821-9826 (2013).
- Bernstein, S. N. (1912-1913). Constructive proof of Weierstrass approximation theorem, *Comm. Kharkov Math. Soc.* 13(1-2).
- Braha, N. L. Srivastava, H. M. Mohiuddine, S. (2014). A Korovkin's type approximation theorem for periodic functions via the statistical summability of the generalized de la Vallée Poussin mean. *Applied Mathematics and Computation*, 228, 162-169.

- Cai, Q. Zhou, G. (2016). On (p, q) -analogue of Kantorovich type Bernstein-Schurer operators, *Applied Mathematics and Computation*, Volume 276, Pages 12-20.
- Dalmanoglu, O. (2007). Approximation by Kantorovich type q -Bernstein operators, in Proceedings of the 12th WSEAS International Conference on Applied Mathematics, Cairo, Egypt, pp. 113-117.
- Devdharma, A. R. Mishra, V. N. (2017). Stancu variant of (p, q) -Szász-Mirakyan operators, *Journal of inequalities and special functions*, Vol. 8, Issue 5, pp. 1-7.
- Dogru, O. Kanat, K. (2012). On statistical approximation properties of the Kantorovich type Lupas operators, *Mathematical and Computer Modelling* 55, 1610-1620.
- Edely, O. H. Mohiuddine, S. A. Noman, A. (2010). Korovkin type approximation theorems obtained through generalized statistical convergence, *Appl. Math. Lett.* 23(11), 1382-1387.
- Fast, H. la Sur. (1951). convergence statistique, *Colloq. Math.* 2, 241-244.
- Farouki, R. T. Rajan, V. T. (1988). Algorithms for polynomials in Bernstein form, *Computer Aided Geometric Design*, Volume 5, Issue 1.
- Gadjiev, A. D. Orhan, C. (2002) Some approximation theorems by statistical convergence, *Rocky Mountain J.math.* 32 , 129-138.
- Gairola, A. R. Deepmala. Mishra, L. N. (2016). Rate of Approximation by Finite Iterates of q -Durrmeyer Operators, *Proc. Natl. Acad. Sci., India, Sect. A Phys. Sci.* 86(2):229-234 (2016). doi: 10.1007/s40010-016-0267-z.
- Hana, L.W. Chu, Y. Qiu, Z. Y. (2014). Generalized Bézier curves and surfaces based on Lupaş q -analogue of Bernstein operator, *Journal of Computational and Applied Mathematics* 261, 352-363.
- Hounkonnou, M. N. Désiré J. Kyemba, B. (2013). $\mathcal{R}(p, q)$ -calculus: differentiation and integration, *SUT Journal of Mathematics*, Vol. 49, No. 2, 145-167.
- Jagannathan, R. Srinivasa, R. K. (2005). Two-parameter quantum algebras, twin-basic numbers, and associated generalized hypergeometric series, *Proceedings of the International Conference on Number Theory and Mathematical Physics*, 20-21.
- Kac, V. Cheung, P. (2002). *Quantum Calculus*, in: Universitext Series, vol. IX, Springer-Verlag.
- Kadak, U. (2016). On weighted statistical convergence based on (p, q) -integers and related approximation theorems for functions of two variables, *Journal of Mathematical Analysis and Applications*, DOI: 10.1016/j.jmaa.2016.05.062.
- Kadak, U. Braha, N. L. Srivastava, H. M. (2017). Statistical Weighted B-Summability and Its Applications to Approximation Theorems, 302, 80-96 , *Applied Mathematics and Computation*.
- Kadak, U. (2017). Weighted statistical convergence based on generalized difference operator involving (p, q) -Gamma function and its applications to approximation theorems, *Journal of Mathematical Analysis and Applications*, Volume 448, Issue 2, Pages 1633-1650.
- Khan, K. Lobiyal, D. K. Kilicman, A. A de Casteljou Algorithm for Bernstein type Polynomials based on (p, q) -integers, *arXiv* 1507.04110v4.
- Khan, K. Lobiyal, D. K. (2017). Bézier curves based on Lupas (p, q) -analogue of Bernstein functions in CAGD, *Journal of Computational and Applied Mathematics* Volume 317, Pages 458-477.
- Khan, K. Lobiyal, D.K. (2015). Bézier curves and surfaces based on modified Bernstein polyno-

- mials, *arXiv:1511.06594v1*. .
- Korovkin, P. P. (1960). Linear operators and approximation theory, *Hindustan Publishing Corporation, Delhi*.
- Lupaş, A. (1987). A q -analogue of the Bernstein operator, Seminar on Numerical and Statistical Calculus, *University of Cluj-Napoca*, 9(85-92).
- Mahmudov N. I. and Sabancigil , P. (2010). Some approximation properties of Lupaş q -analogue of Bernstein operators, *arXiv:1012.4245v1 [math.FA]*.
- Mishra, V.N. Pandey, S. (2017). On (p, q) Baskakov-Durrmeyer-Stancu Operators, *Advances in Applied Clifford Algebras*, Vol. 27, Issue 2, pp. 1633-1646 DOI:10.1007/s00006-016-0738-y.
- Mishra, V.N. Pandey, S. (2016). On Chlodowsky variant of (p, q) Kantorovich-Stancu-Schurer operators, *International Journal of Analysis and Applications*, Vol. 11, No. 1, pp. 28-39.
- Mishra, V.N. Khatri, K. Mishra, L. N. Deepmala. (2013). Inverse result in simultaneous approximation by Baskakov-Durrmeyer-Stancu operators, *Journal of Inequalities and Applications*, 2013:586. doi:10.1186/1029-242X-2013-586
- Mursaleen, M. Ansari, K. J. Khan, A. (2015). On (p, q) -analogue of Bernstein Operators, *Appl. Math. Comput.*, 266, 874-882, [Erratum: *Appl. Math. Comput.*(2016), 278, 70-71.]
- Mursaleen, M. and Nasiruzzaman M. and Nurgali, A. (2015). Some approximation results on Bernstein-Schurer operators defined by (p, q) -integers, *Journal of Inequalities and Applications* , 249, DOI 10.1186/s13660-015-0767-4.
- Mursaleen, M. Ansari, K. J. and Khan, A. (2015). Some Approximation Results by (p, q) -analogue of Bernstein-Stancu Operators, *Appl. Math. Comput.*, 264, 392-402. (Corrigendum: *Appl. Math. Comput.*269(2015) 744-746).
- Mursaleen, M. Ansari, K. J. and Khan, A. (2016). Some approximation results for Bernstein-Kantorovich operators based on (p, q) -calculus, *U.P.B. Sci. Bull., Series A Vol. 78, Iss. 4, ISSN 1223-7027*.
- Mursaleen, M. Nasiruzzaman, M. Khan, A. and Ansari, K. J. (2016). Some approximation results on Bleimann-Butzer-Hahn operators defined by (p, q) -integers, *Filomat* 30:3 , 639-648, DOI 10.2298/FIL1603639M.
- Mursaleen, M. Khan, A. (2013). Generalized q -Bernstein-Schurer Operators and Some Approximation Theorems, *Journal of Function Spaces and Applications*, Article ID 719834, 7 pages <http://dx.doi.org/10.1155/2013/719834>
- Mursaleen, M. Khan, F and Khan, A. (2016). Approximation by (p, q) -Lorentz polynomials on a compact disk, *Complex Anal. Oper. Theory*, 10(8), 1725-1740.
- Mursaleen, M. Nasiruzzaman, M. Khan, F. Khan, A. (2017). On (p, q) -analogue of divided differences and Bernstein operators, Vol. (2017), Article ID 25, pp. 1 – 13.
- Oruk, H. Phillips, G. M. (2003). q -Bernstein polynomials and Bézier curves, *Journal of Computational and Applied Mathematics* 151, 1-12.
- Ostrowska, S. (2006). On the Lupaş q -analogue of the Bernstein operator, *Rocky mountain journal of mathematics* Volume 36, Number 5.
- Pandey, S. (2016). On Chlodowsky variant of (p, q) –Kantorovich-Stancu-Schurer operators, *International Journal of Analysis and Applications*, Vol. 11, No. 1, pp. 28-39.
- Phillips, G. M. (2000). A generalization of the Bernstein polynomials based on the q -integers *ANZIAMJ* 42, 79-86.

- Phillips, G. M. (2003). Interpolation and Approximation by Polynomials, *Springer*.
- Rababah, A. Manna, S. (2011). Iterative process for G2-multi degree reduction of Bézier curves, *Appl. Math. Comput.* 217, 8126-8133.
- Wafi, A. Rao, N. and Deepmala. (2019). Approximation properties of (p, q) -variant of Stancu-Schurer, *Boletim da Sociedade Paranaense de Matemática*, Vol 37, No 4.
- Wafi, A. Rao, N. (2017). Bivariate-Schurer-Stancu operators based on (p, q) -integers, *general mathematics*.
- Wafi, A. Rao, N. (2016). (p, q) -Bivariate-Bernstein-Chlowdosky operators *Filomat* , (accepted)<http://journal.pmf.ni.ac.rs/filomat/index.php/filomat/author/>.
- Wafi, A. Rao, N. (2016). A generalization of Szasz-type operators which preserves constant and quadratic test functions, *cogent mathematics*, 3 (1), 1227023.
- Wafi, A. Rao, N. (2016) Modified Szasz operators involving Charlier polynomials based on two parameters, *Thai J. of Maths*.
- Wafi, A. Rao, N. (2016) Approximation properties by generalized Baskakov Kantorovich Stancu type operators, *Appl. Maths Inf. Sci. Lett.*, Vol. 4, No. 3, PP:111-118.