



Applications and Applied Mathematics: An International Journal (AAM)

Volume 14 | Issue 1

Article 31

6-2019

Integral Inequalities of Hermite-Hadamard Type Via Green Function and Applications

Tuba Tunç

Duzce University

Sümeyye Sönmezoglu

Duzce University

Mehmet Z. Sarıkaya

Duzce University

Follow this and additional works at: <https://digitalcommons.pvamu.edu/aam>

 Part of the Analysis Commons

Recommended Citation

Tunç, Tuba; Sönmezoglu, Sümeyye; and Sarıkaya, Mehmet Z. (2019). Integral Inequalities of Hermite-Hadamard Type Via Green Function and Applications, Applications and Applied Mathematics: An International Journal (AAM), Vol. 14, Iss. 1, Article 31.

Available at: <https://digitalcommons.pvamu.edu/aam/vol14/iss1/31>

This Article is brought to you for free and open access by Digital Commons @PVAMU. It has been accepted for inclusion in Applications and Applied Mathematics: An International Journal (AAM) by an authorized editor of Digital Commons @PVAMU. For more information, please contact hvkoshy@pvamu.edu.



On Integral Inequalities of Hermite-Hadamard Type Via Green Function and Applications

¹Tuba Tunç, ²Sümeyye Sönmezoglu and ³Mehmet Zeki Sarıkaya

Department of Mathematics
 Faculty of Science and Arts
 Duzce University
 Duzce-Turkey

[¹tubatunc03@gmail.com](mailto:tubatunc03@gmail.com); [²sarikayamz@gmail.com](mailto:sarikayamz@gmail.com); [³smyy.mat@gmail.com](mailto:smyy.mat@gmail.com)

Received: January 14, 2019; Accepted: April 29, 2019

Abstract

In this study, we establish some Hermite- Hadamard type inequalities for functions whose second derivatives absolute value are convex. In accordance with this purpose, we obtain an identity using Green's function. Then using this equality we get our main results.

Keywords: Hermite-Hadamard's inequality; Convex function; Green function; Holder inequality

MSC 2010 No.: 26A33, 26A51, 26D15

1. Introduction

The following inequality discovered by Hermite and Hadamard for convex functions is well known in the literature as the Hermite-Hadamard inequality (see, e.g., Dragomir and Pearce (2000)):

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}, \quad (1)$$

where $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ is a convex function on the interval I of real numbers and $a, b \in I$ with $a < b$.

Hermite-Hadamard inequality provides a lower and an upper estimations for the integral average of any convex functions defined on a compact interval. This inequality has a notable place in mathematical analysis, optimization and so on. However, many studies have been established to demonstrate its new proofs, refinements, extensions and generalizations. A few of these studies are Azpeitia (1994), Erden and Sarikaya (2016), Husain et al. (2009), Pečarić et al. (1991), Pearce and Pečarić (2000), Sarikaya et al. (2012)-Xi and Qi (2013) referenced works and also the references included there. Particularly, see in the references Barani et al. (2012), Husain et al. (2009), Sarikaya et al. (2012), Set and Korkut (2016) that Hermite-Hadamard type inequalities for functions whose second derivatives absolute value are convex, s -convex and so on.

In this study, we establish new inequalities that are connected with the right-hand side of Hermite-Hadamard inequality by using Green's function and functions whose second derivatives absolute value are convex,

2. Preliminaries

In this section, we gave some works about the right hand side of the Hermite-Hadamard inequality (1) for twice differentiable mappings.

Dragomir and Pearce (2000) gave the following lemma related to the right hand side of the inequality (1) using twice differentiable mapping:

Lemma 2.1.

Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be a twice differentiable mapping on I° , $a, b \in I$ with $a < b$ and f'' of integrable on $[a, b]$, the following equality holds:

$$\frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx = \frac{(b-a)^2}{2} \int_0^1 t(1-t) f''(ta + (1-t)b) dt. \quad (2)$$

Hussain et al. (2009) proved some inequalities related to a trapezoid inequality of Hermite-Hadamard for s -convex functions by using Lemma 2.1:

Theorem 2.2.

Let $f : I \subset [0, \infty) \rightarrow \mathbb{R}$ be twice differentiable mapping on I° such that $f'' \in L_1[a, b]$, where $a, b \in I$ with $a < b$. If $|f''|$ is s -convex on $[a, b]$ for some fixed $s \in [0, 1]$ and $q \geq 1$, then the following inequality holds:

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{(b-a)^2}{2 \times 6^{\frac{1}{p}}} \left[\frac{|f''(a)|^q + |f''(b)|^q}{(s+2)(s+3)} \right]^{\frac{1}{q}}, \quad (3)$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

Remark 2.3. (Hussain et al. (2009))

If we take $s = 1$ in (3), then we have

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{(b-a)^2}{12} \left[\frac{|f''(a)|^q + |f''(b)|^q}{2} \right]^{\frac{1}{q}}. \quad (4)$$

Sarikaya and Aktan (2011) gave the following trapezoid inequality of Hermite-Hadamard inequality (1):

Theorem 2.4.

Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be twice differentiable function on I° with $f'' \in L_1[a,b]$. If $|f''|$ is a convex on $[a,b]$, then

$$\left| \frac{1}{b-a} \int_a^b f(x) dx - \frac{f(a) + f(b)}{2} \right| \leq \frac{(b-a)^2}{12} \left[\frac{|f''(a)| + |f''(b)|}{2} \right]. \quad (5)$$

In the following we will give some necessary definitions and mathematical preliminaries of fractional calculus theory which are used further in this paper:

Definition 2.5. [Gorenflo and Mainardi (1997), Miller and Ross (1993), Kilbas et al. (2006) Podlubny (1999)]

Let $f \in L_1[a,b]$. The Riemann-Liouville fractional integrals $J_{a+}^\alpha f$ and $J_{b-}^\alpha f$ of order $\alpha > 0$ with $a \geq 0$ are defined by

$$J_{a+}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f(t) dt, \quad x > a \quad (6)$$

and

$$J_{b-}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} f(t) dt, \quad x < b, \quad (7)$$

respectively. Here, $\Gamma(\alpha)$ is the Gamma function and $J_{a+}^0 f(x) = J_{b-}^0 f(x) = f(x)$.

It is remarkable that Sarikaya et al.(2013) first give the following interesting integral inequalities of Hermite-Hadamard type involving Riemann-Liouville fractional integrals.

Theorem 2.6.

Let $f : [a,b] \rightarrow \mathbb{R}$ be a positive function with $0 \leq a < b$ and $f \in L_1[a,b]$. If f is a convex function on $[a,b]$, then the following inequalities for fractional integrals hold:

$$f\left(\frac{a+b}{2}\right) \leq \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{a+}^\alpha f(b) + J_{b-}^\alpha f(a)] \leq \frac{f(a) + f(b)}{2} \quad (8)$$

with $\alpha > 0$.

On the other hand, Wang et al. (2012) proved the following identity for twice differentiable function involving Riemann-Liouville fractional integrals:

Lemma 2.7.

Let $f : [a,b] \rightarrow \mathbb{R}$ be a twice differentiable mapping on (a,b) with $0 \leq a < b$. If $f'' \in L[a,b]$, then the following equality for fractional integrals holds:

$$\begin{aligned} & \frac{f(a) + f(b)}{2} - \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{a+}^\alpha f(b) + J_{b-}^\alpha f(a)] \\ &= \frac{(b-a)^2}{2(\alpha+1)} \int_0^1 [1 - (1-t)^{\alpha+1} - t^{\alpha+1}] f''[ta + (1-t)b] dt. \end{aligned} \quad (9)$$

For recent results connected with fractional integral inequalities see Kunt et al. (2016), Sarikaya and Budak (2016)-Set et al. (2017), Wang et al. (2012), Wang and Qi (2017).

3. Main Results

In order to prove our main results we need the following lemma:

Lemma 3.1.

Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be a twice differentiable function on I° (the interior of the interval I) such that $f'' \in L[a,b]$, where $a, b \in I^\circ$ with $a < b$. Then the following identity holds:

$$\frac{\alpha}{2(b-a)^\alpha} \iint_a^b G(x,y) f''(y) dy dx = \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{a+}^\alpha f(b) + J_{b-}^\alpha f(a)] - \frac{f(a) + f(b)}{2}, \quad (10)$$

where

$$G(x;y) = \begin{cases} (x - a)^{\alpha-1} (y - b); & a < x < y < b; \\ (b - x)^{\alpha-1} (a - y); & a < x < y < b \end{cases}$$

Proof:

By integration by parts, we have

$$\int_a^b G(x, y) f''(y) dy = G(x, y) f'(y) \Big|_a^b - \int_a^b \frac{\partial G(x, y)}{\partial y} f'(y) dy.$$

Since $G(x, b) = G(x, a) = 0$, it is easy to see, we get

$$\begin{aligned} & \int_a^b G(x, y) f''(y) dy \\ &= \int_a^x (b-x)^{\alpha-1} f(y) dy - \int_x^b (x-a)^{\alpha-1} f(y) dy \\ &= (b-x)^{\alpha-1} f(x) + (x-a)^{\alpha-1} f(x) - (b-x)^{\alpha-1} f(a) - (x-a)^{\alpha-1} f(b). \end{aligned} \tag{11}$$

Integrating both sides of (11) with respect to x over $[a, b]$, we obtain

$$\begin{aligned} & \int_a^b \int_a^b G(x, y) f''(y) dy dx \\ &= \int_a^b (b-x)^{\alpha-1} f(x) dx + \int_a^b (x-a)^{\alpha-1} f(x) dx - \frac{(b-a)^\alpha}{\alpha} [f(a) + f(b)] \end{aligned} \tag{12}$$

Multiplying both sides of (12) by $\frac{\alpha}{2(b-a)^\alpha}$ and rearranging the last identity, we get desired inequality.

Theorem 3.2.

Let $f : I^\circ \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a twice differentiable mapping on I° , $a, b \in I^\circ$ with $a < b$. If $|f''|$ is convex on $[a, b]$, then the following inequality holds:

$$\left| \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{a+}^\alpha f(b) + J_{b-}^\alpha f(a)] - \frac{f(a) + f(b)}{2} \right| \leq \frac{(b-a)^2}{(\alpha+1)(\alpha+2)} \left(\frac{|f''(a)| + |f''(b)|}{2} \right). \tag{13}$$

Proof:

From Lemma 3.1, we have

$$\begin{aligned}
& \left| \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} \left[J_{a+}^\alpha f(b) + J_{b-}^\alpha f(a) \right] - \frac{f(a) + f(b)}{2} \right| \\
& \leq \frac{\alpha}{2(b-a)^\alpha} \int_a^b \int_a^b |G(x,y)| |f''(y)| dy dx \\
& = \frac{\alpha}{2(b-a)^\alpha} \\
& \quad \times \left[\int_a^b \int_a^x (b-x)^{\alpha-1} (y-a) |f''(y)| dy dx + \int_a^b \int_x^b (b-y) (x-a)^{\alpha-1} |f''(y)| dy dx \right].
\end{aligned}$$

Because $|f''(y)|$ is convex on $[a,b]$, we can write

$$\left| f'' \left(\frac{b-y}{b-a} a + \frac{y-a}{b-a} b \right) \right| \leq \frac{b-y}{b-a} |f''(a)| + \frac{y-a}{b-a} |f''(b)|. \quad (14)$$

Using (14), it follows that

$$\begin{aligned}
& \left| \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} \left[J_{a+}^\alpha f(b) + J_{b-}^\alpha f(a) \right] - \frac{f(a) + f(b)}{2} \right| \\
& \leq \frac{\alpha}{2(b-a)^{\alpha+1}} \\
& \quad \times \left[|f''(a)| \int_a^b \int_a^x (b-x)^{\alpha+1} (y-a)(b-y) dy dx + |f''(b)| \int_a^b \int_a^x (b-x)^{\alpha+1} (y-a)^2 dy dx \right. \\
& \quad \left. + |f''(a)| \int_a^b \int_x^b (x-a)^{\alpha+1} (b-y)^2 dy dx + |f''(b)| \int_a^b \int_x^b (x-a)^{\alpha+1} (b-y)(y-a) dy dx \right].
\end{aligned}$$

If we calculate the above integrals and also use elementary analysis, we obtain desired expression (13).

Theorem 3.3.

Let $f : I^o \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a twice differentiable mapping on I^o , $a, b \in I^o$ with $a < b$. If $|f''|^q$ is convex on $[a,b]$, $q > 1$, then the following inequality holds:

$$\begin{aligned}
& \left| \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} \left[J_{a+}^\alpha f(b) + J_{b-}^\alpha f(a) \right] - \frac{f(a) + f(b)}{2} \right| \\
& \leq \frac{\alpha(b-a)^2}{2^{\frac{1}{q}} (p+1)^{\frac{1}{p}}} \left(\frac{|f''(a)|^q + |f''(b)|^q}{2} \right)^{\frac{1}{q}} [B(p+2, p(\alpha-1)+1)]^{\frac{1}{p}}, \quad (15)
\end{aligned}$$

where $\frac{1}{p} + \frac{1}{q} = 1$ and $B(p, q) = \int_0^1 u^{p-1}(1-u)^{q-1} du$, $(p, q > 0)$ is Euler's Beta function.

Proof:

From Lemma 3.1 and using Holder's inequality, we find that

$$\begin{aligned} & \left| \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{a+}^\alpha f(b) + J_{b-}^\alpha f(a)] - \frac{f(a) + f(b)}{2} \right| \\ & \leq \frac{\alpha}{2(b-a)^\alpha} \left(\int_a^b \int_a^b |G(x, y)|^p dy dx \right)^{\frac{1}{p}} \left(\int_a^b \int_a^b |f''(y)|^q dy dx \right)^{\frac{1}{q}}. \end{aligned} \quad (16)$$

In order to proceed further to complete the proof, we evaluate the above integrals.

$$\begin{aligned} & \int_a^b \int_a^b |G(x, y)|^p dy dx \\ &= \int_a^b \int_a^x (b-x)^{(\alpha-1)p} (y-a)^p dy dx + \int_a^b \int_x^b (b-y)^p (x-a)^{(\alpha-1)p} dy dx \\ &= \int_a^b \frac{(b-x)^{(\alpha-1)p} (x-a)^{p+1}}{(p+1)} dx + \int_a^b \frac{(x-a)^{(\alpha-1)p} (b-x)^{p+1}}{(p+1)} dx. \end{aligned}$$

Using the change of the variables $x-a = (b-a)u$ and $b-x = (b-a)u$ for the above integrals, we get

$$\int_a^b \int_a^b |G(x, y)|^p dy dx = \frac{2(b-a)^{\alpha p+2}}{p+1} B(p+2, p(\alpha-1)+1). \quad (17)$$

Since $|f''(y)|^q$ is convex on $[a, b]$, we can write

$$\left| f''\left(\frac{b-y}{b-a}a + \frac{y-a}{b-a}b\right) \right|^q \leq \frac{b-y}{b-a} |f''(a)|^q + \frac{y-a}{b-a} |f''(b)|^q. \quad (18)$$

Using the inequality (18), it follows that

$$\begin{aligned} & \int_a^b \int_a^b |f''(y)|^q dy dx \\ & \leq |f''(a)|^q \int_a^b \int_a^b \frac{b-y}{b-a} dy dx + |f''(b)|^q \int_a^b \int_a^b \frac{y-a}{b-a} dy dx \\ &= \frac{|f''(a)|^q + |f''(b)|^q}{2} (b-a)^2. \end{aligned} \quad (19)$$

Finally by substituting (17) and (19) in (16), then we easily obtain required inequality (15). This completes the proof.

Now, we establish trapezoid in a different way by using convexity of $|f''|^q$.

Theorem 3.4.

Let $f : I^o \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a twice differentiable mapping on I^o , $a, b \in I^o$ with $a < b$. If $|f''|^q$ is convex on $[a, b]$, $q > 1$, then the following inequality holds:

$$\left| \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{a+}^\alpha f(b) + J_{b-}^\alpha f(a)] - \frac{f(a) + f(b)}{2} \right| \quad (20)$$

$$\leq \frac{\alpha(b-a)^2}{2^{1+\frac{1}{q}}(p+1)^{\frac{1}{p}}} [B(p+2, p(\alpha-1)+1)]^{\frac{1}{p}} \\ \times \left[\frac{2|f''(a)|^q + |f''(b)|^q}{3} \right]^{\frac{1}{q}} + \left[\frac{|f''(a)|^q + 2|f''(b)|^q}{3} \right]^{\frac{1}{q}},$$

where $\frac{1}{p} + \frac{1}{q} = 1$ and $B(p, q) = \int_0^1 u^{p-1} (1-u)^{q-1} du$, $(p, q > 0)$ is Euler's Beta function.

Proof:

From Lemma 3.1 and using Holder's inequality, we have

$$\begin{aligned} & \left| \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{a+}^\alpha f(b) + J_{b-}^\alpha f(a)] - \frac{f(a) + f(b)}{2} \right| \\ & \leq \frac{\alpha}{2(b-a)^\alpha} \left\{ \left[\int_a^b \int_a^x (b-x)^{(\alpha-1)p} (y-a)^p dy dx \right]^{\frac{1}{p}} \left[\int_a^b \int_a^x |f''(y)|^q dy dx \right]^{\frac{1}{q}} \right. \\ & \quad \left. + \left[\int_a^b \int_x^b (b-y)^p (x-a)^{(\alpha-1)p} dy dx \right]^{\frac{1}{p}} \left[\int_a^b \int_x^b |f''(y)|^q dy dx \right]^{\frac{1}{q}} \right\}. \end{aligned} \quad (21)$$

By calculating each of integrals in (21) with utilizing convexity of $|f''(y)|^q$, we obtain

$$\begin{aligned}
& \left| \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{a+}^\alpha f(b) + J_{b-}^\alpha f(a)] - \frac{f(a) + f(b)}{2} \right| \\
& \leq \frac{\alpha}{2(b-a)^\alpha} \\
& \times \left[\left[\frac{(b-a)^{\alpha p+2}}{p+1} B(p+2, p(\alpha-1)+1) \right]^{\frac{1}{p}} \left[\frac{(b-a)^2}{2} \times \frac{2|f''(a)|^q + |f''(b)|^q}{3} \right]^{\frac{1}{q}} \right. \\
& \left. + \left[\frac{(b-a)^{\alpha p+2}}{p+1} B(p+2, p(\alpha-1)+1) \right]^{\frac{1}{p}} \left[\frac{(b-a)^2}{2} \times \frac{|f''(a)|^q + 2|f''(b)|^q}{3} \right]^{\frac{1}{q}} \right]. \tag{22}
\end{aligned}$$

The proof is thus completed.

Now, we proved the following theorem using Holder's inequality different way:

Theorem 3.5.

Let $f : I^o \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a twice differentiable mapping on I^o , $a, b \in I^o$ with $a < b$. If $|f''|^q$ is convex on $[a, b]$, $q \geq 1$, then the following inequality holds:

$$\left| \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{a+}^\alpha f(b) + J_{b-}^\alpha f(a)] - \frac{f(a) + f(b)}{2} \right| \leq \frac{(b-a)^2}{12 \times 2^{\frac{1}{p}}} \left(\frac{|f''(a)|^q + |f''(b)|^q}{2} \right)^{\frac{1}{q}}, \tag{23}$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

Proof:

From Lemma 3.1 and using Holder's inequality, we find that

$$\begin{aligned}
& \left| \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{a+}^\alpha f(b) + J_{b-}^\alpha f(a)] - \frac{f(a) + f(b)}{2} \right| \\
& \leq \frac{\alpha}{2(b-a)^\alpha} \left[\int_a^b \int_a^b |G(x, y)| dy dx \right]^{\frac{1}{p}} \left[\int_a^b \int_a^b |G(x, y)| |f''(y)|^q dy dx \right]^{\frac{1}{q}}. \tag{24}
\end{aligned}$$

We calculate the above integrals respectively:

$$\int_a^b \int_a^b |G(x, y)| dy dx = \frac{2}{\alpha(\alpha+1)(\alpha+2)} (b-a)^{\alpha+2}. \tag{25}$$

Because $|f''(y)|^q$ is convex on $[a, b]$, we obtain

$$\int_a^b \int_a^b |G(x,y)| |f''(y)|^q dy dx = \frac{(b-a)^2}{2^{\frac{1}{q}}(\alpha+1)(\alpha+2)} \left(\frac{|f''(a)|^q + |f''(b)|^q}{2} \right). \quad (26)$$

Substituting the equalities (25) and (26) in (24), we obtain the inequality (23) which completes the proof.

4. Conclusion

In this paper, using a special function and Riemann-Liouville fractional integrals, we obtain several Hermite-Hadamard type inequalities for functions whose second derivatives absolute value are convex. It is important because of giving Hermite-Hadamard type inequalities different way. Similarly, you can obtain such as inequalities using special functions.

REFERENCES

- Azpeitia, A.G. (1994). Convex functions and the Hadamard inequality, Rev. Colombiana Math., 28, pp. 7-12.
- Bakula, M. K. and Pečarić, J. (2004). Note on some Hadamard-type inequalities, Journal of Inequalities in Pure and Applied Mathematics, 5(3), article 74, 9 pages.
- Barani, A., Ghanzafari, A.G. and Dragomir S.S. (2012). Hermite-Hadamard inequality for functions whose derivatives absolute values are preinvex, Journal of Inequalities and Applications, 2012:247, 9 pages.
- Dragomir, S.S. and Agarwal, R.P. (1998). Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula, Appl. Math. lett., 11(5), pp. 91-95.
- Dragomir, S.S. and Pearce, C.E.M. (2000). *Selected Topics on Hermite-Hadamard Inequalities and Applications*, RGMIA Monographs, Victoria University.
- Erden, S. and Sarikaya, M.Z. (2016). New Hermite Hadamard type inequalities for twice differentiable convex mappings via Green function and applications, Moroccan J. Pure and Appl. Anal., 2(2), pp. 107-117.
- Gorenflo, R., Mainardi, F. (1997). *Fractional calculus: integral and differential equations of fractional order*, Springer Verlag, Wien, pp. 223-276.
- Miller, S. and Ross, B. (1993). *An introduction to the Fractional Calculus and Fractional Differential Equations*, John Wiley&Sons, USA, pp.2.
- Hussain, S., Bhatti, M.I. and Iqbal M. (2009). Hadamard-type inequalities for s-convex functions I, Punjab Univ. Jour. of Math., Vol.41, pp.51-60.
- Kilbas, A.A., Srivastava, H.M. and Trujillo, J.J. (2006). *Theory and Applications of Fractional Differential Equations*, North-Holland Mathematics Studies, 204, Elsevier Sci. B.V., Amsterdam.
- Kunt, M., İşcan, İ., Yazıcı, N. And Gözütok U. (2016). On new inequalities of Hermite-Hadamard Fejer type for harmonically convex functions via fractional integrals, SpringerPlus, 5:635, 19 pages.

- Pečarić, J., Proschan, F. and Tong, Y.L. (1991). *Convex functions, partial ordering and statistical applications*, Academic Press, New York.
- Pearce, C.E.M. and Pečarić, J. (2000). Inequalities for differentiable mappings with application to special means and quadrature formulae, *Appl. Math. Lett.*, 13(2), pp. 51-55.
- Podlubni, I.(1999). *Fractional Differential Equations*, Academic Press, San Diego.
- Sarikaya, M.Z, Saglam, A. and Yıldırım, H. (2012). New inequalities of Hermite-Hadamard type for functions whose second derivatives absolute values are convex and quasi-convex, *Int. J. Open Problems Comput. Math.*, 5(3).
- Sarikaya, M.Z. and Aktan, N. (2011). On the generalization of some integral inequalities and their applications, *Mathematical and Computer Modelling*, vol. 54, pp. 2175-2182.
- Sarikaya, M.Z. and Budak, H. (2016). Generalized Hermite-Hadamard type integral inequalities for fractional integrals, *Filomat*, 30:5, pp. 1315-1326.
- Sarikaya, M.Z., Set, E., Yaldiz, H. and Basak, N.(2013). Hermite -Hadamard's inequalities for fractional integrals and related fractional inequalities, *Mathematical and Computer Modelling*, DOI:10.1016/j.mcm.2011.12.048, vol. 57, pp. 2403-2407.
- Set, E., Özdemir, M.E., Sarikaya, M.Z. and Karakoç, F. (2017). Hermite-Hadamard type inequalities for (α, m) -convex functions via fractional integrals, *Moroccan J. Pure and Appl. Anal.*, 3(1), pp. 15-21.
- Set, E. And Korkut, N. (2016). On new inequalities of Hermite Hadamard type for functions whose second derivatives in absolute value are s-convex, *AIP Conference Proceedings* 1726, 020040, doi: <http://dx.doi.org/10.1063/1.4945866>.
- Wang, J., Li, X., Feckan, M. and Zhou, Y. (2012). Hermite-Hadamard-type inequalities for Riemann-Liouville fractional integrals via two kinds of convexity, *Appl. Anal.*, doi:10.1080/00036811.2012.727986.
- Wang, S-H and Qi, F. (2017). Hermite-Hadamard type inequalities for s -convex functions via Riemann--Liouville fractional integrals, *J. Computational Analysis and Applications*, 22(6), pp. 1124-1134.
- Xi, B-Y and Qi, F. (2013). Some Hermite-Hadamard type inequalities for differentiable convex functions and applications, *Hacet. J. Math. Stat.*, 42(3), pp. 243-257.