

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,500

Open access books available

136,000

International authors and editors

170M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Ecology of Plant Communities in Central Mexico

*Joaquín Sosa-Ramírez, Vicente Díaz-Núñez  
and Diego R. Pérez-Salicrup*

## Abstract

In Central Mexico converge three biogeographic provinces: Altiplano sur, Sierra Madre Occidental and Costa del Pacífico. Each one of them is composed by different plant communities: Thorn Forest, Temperate Mountain Forest and Dry Tropical Forest respectively. Our objective is to show, through phytoecological analysis, the species richness, diversity and the structure of the plant communities from the Temperate Mountain Forest and from the Tropical Dry Forest. In the Temperate Mountain Forest, 50 forest species were recorded, with a Shannon Wiener diversity index  $H' = 1.63$  on altitudes from 2400 to 2600 m. The Whittaker  $\beta$  index is  $B_w = 7.22$ . In the tropical dry forest, we identified 79 plants species with a mean diversity index  $H' = 3.49$  on altitudes from 1951 to 2100 m. In this ecosystem the  $B_w$  index is 8.12. This study offers important information for the establishment of management practices, considering the protection status from the areas in which this vegetation type is distributed.


**Keywords:** Aguascalientes, Sierra Fria, Temperate Mountain Forest, Tropical Dry Forest, biogeographic provinces

## 1. Introduction

Mexico is one of the five countries with the greatest biological diversity in the world, due, in part to the confluence of the Nearctic (North America) and Neotropical biogeographic zones (Mexico, Central and south America). As well as, the species evolutionary processes in its territory [1]. The Mexican territory represent only 1% of the earth's surface; nevertheless, Mexico belongs to the select group of the five countries considered megadiverse, along with Brazil, Colombia, China and Indonesia [2, 3]. Due to its geographic locations and its multiple landscapes, a large number and diversity of ecosystems converge in the national territory. For that reason, Mexico is ranked 12th in terms of global forest area [4]. Even though, multiple efforts have been made for the forest conservation during the last decade of the XXI century, on a global scale, forest have been transformed to other uses at a rate of  $1.3 \times 10^6$  million ha/yr. or they have been affected by natural disturbances that have partially or totally changed their structure. This amount represents a 19% decrease in comparison to the exchange rate registered in the last decade of the 20th century ( $1.6 \times 10^6$  million ha/yr) [5]. Temperate forests in Mexico are found mostly, although not exclusively, in the mountainous areas along the Sierra Madre Occidental (the area with the highest concentration of forest ecosystems in the country),

the mountains of Sierra Madre Oriental, the Sierra Norte de Oaxaca and the Altos de Chiapas, as well as in different mountain ranges and isolated mountains in the Altiplano and intermingled in the tropical plains [6]. The conifer and oak forest in Mexico represent the most extensive vegetation cover in terms of vegetation types dominated by woody species, this species covers 16.4% of the total surface of the country, being only surpassed by the xeric shrubland which is the vegetation type which has the largest extension [5]. These ecosystems are important both economically and ecologically, since they support productive activities, harbor great biological diversity and serves as a refuge for wildlife. Likewise, forest provide essential environmental goods and services for the human society subsistence [7, 8].

The State of Aguascalientes has a total extension of 555, 867.4 hectares, of which 291,792.4 hectares equivalent to 52.5% present some forest type [9]. According to the classifications issued by different sources [10, 11], the State of Aguascalientes is made up by three large ecoregions (biogeographic regions), the Temperate Mountain Forest, the Tropical Dry Forest (also known as lowland deciduous forest) and the thorn forest (including crasicuale shrubland and xeric shrubland). The first ecosystem type is mainly distributed in la Sierra Fria, Sierra del Laurel, Sierra de Tepezalá and Cerro de Juan el Grande in El Llano municipality (**Figure 1**). The largest area covered by Temperate Mountain Forest vegetation in Aguascalientes is located in an area locally known as Sierra Fria, this site is a Protected Natural Area by state and federal decree which covers close to 107,000 ha [12]. In the Temperate Mountain Forest, the plant communities the most common vegetation types are oak forests (*Quercus* spp.), pine trees (*Pinus* spp.), oak-pine, pine-oak, juniper (*Juniperus* spp.), manzanita shrubland (*Arcostaphylos pungens*) and different associations of these genera. The vegetation that has mainly colonized the sites that had been disturbed are *Juniperus deppeana* and *A. pungens*, although there has also been an increase in conifer populations [13]. The second largest formation where this ecosystem is found is located in la Sierra del Laurel in the Southwest corner of the State occupying close to 17,000 ha. This area presents similar plant communities



**Figure 1.** Distribution areas of the Temperate Mountain Forest in Aguascalientes state. (1) sierra Fria protected natural area (San José de Gracia, Pabellón de Arteaga, Rincón de Romos and Calvillo municipalities); (2) Mountain Hill of Tepezalá, and (3) Juan el Grande Mountain (El llano municipality).

but with a greater dominance of oak populations (*Quercus* spp.) and lower density of manzanita (*A. pungens*).

The largest area occupied by the tropical dry forest is mainly located in the Calvillo municipality, although, there are relics of vegetation indicative of this ecosystem in the Jesus Maria, San Jose de Gracia and Aguascalientes municipalities, which suggests a larger presence of this vegetation type in the past. In the tropical dry forest, forest structures made up of shrubs and trees between 2 and 8 m high and some relics of medium tropical forest. In Aguascalientes, this is one of the ecosystems with the highest species richness [14]. The most representative vegetation in this ecoregion corresponds mainly to the *Lysiloma*, *Bursera*, *Ipomoea*, *Acacia*, *Eysenhardtia*, *Opuntia*, *Mimosa* and *Agave* genera.

Our objective was to provide an overview of some ecological aspects (species richness, diversity and distribution) of woody species natural communities in the most representative ecosystems of the State of Aguascalientes, assuming that there would be a high similarity degree with the vegetation of neighboring sites, considering both the environmental and physiographic characteristics from this State.

## 2. Materials and methods

Three studies were conducted individually. During 2008–2015, the natural communities of the temperate mountain forest in the area commonly known as Sierra Fria, in the northwest of the State of Aguascalientes, as well as the main disturbances that have affected them in the past and present were analyzed [10, 13]. Likewise, during the period 2011–2015 a study was carried out to determine the diversity, dynamics and functioning of the tropical dry forest in the Calvillo municipality [14, 15].

### 2.1 Temperate Mountain Forests


#### 2.1.1 Study area and sampling design

This study was carried out in to the Sierra Fria Protected Natural Area (SF-PNA) which is 106,114.6 hectares in size and is located in the northwest of the Aguascalientes State. This area has an altitude ranging between 2,100 and 3,050 masl. The study area comprised 25 thousand hectares, in a polygon located between the coordinates 102°31'31" to 102°37'44" west longitude and 22°05'47" at 22°14'03" north latitude, assuming that the conditions both geographic, ecological and climatic are representative of the entire ANP (See **Figure 2**).

A stratified sampling strategy was developed [16]. The sampling strata were delimited based on the altitude, solar exposure, and geoform of the site (flat, concave and convex terrain). The first stratum was defined using a Digital Elevation Model (DEM) of the ANP SF, elaborating a spatial grid according to five altitudinal categories: i) 2,000-2,200, ii) 2,200-2,400, iii) 2,400-2,600, iv) 2,600-2,800, and v) >2 800 masl.

To establish the altitudinal strata, the level curves from study site were defined using the DEM. The solar exposure was approached using an exposure map made with a SPOT 2010® image on which the DEM of the site was superimposed. Subsequently, a mesh map was prepared using the ArcGis 10.2. The geoform was obtained based on the slope, where flat terrain = sites with a slope  $\leq 10\%$ , concave t. = slope  $\geq 10$  and  $\leq 25\%$  and convex t. = slope  $\geq 25\%$ .





**Figure 2.** Location of the protected natural area Sierra Fria, the study area of the Temperate Mountain Forest.

### 2.1.2 Identification, distribution and abundance of forest species

To identify the tree and shrub diversity in the study area, we conducted 60 phytoecological inventories in 60 different sites distributed randomly using the sampling scheme already described (Table 1).

The field samplings were performed in rectangular plots of 600 m<sup>2</sup>, with a central line 100 m in length and two lateral lines with three m of separation. In each inventory, the frequency of the tree and shrub species present were determined, as well as the site environmental variables. Individuals with DBH ≥ 5 cm and height ≥ 1.50 m were considered as trees. Individuals below these categories were

Altitude levels	Topographic position								Flat	Total
	Concave				Convexe					
	N	S	E	W	N	S	E	W		
2000–2200	0	0	0	0	1	1	0	0	1	3
2200–2400	0	1	0	0	3	2	0	0	0	6
2400–2600	8	4	1	1	8	3	0	0	3	28
2600–2800	2	0	0	1	4	3	0	0	11	21
> 2800	2	0	0	0	0	0	0	0	0	2
<b>Total</b>	<b>12</b>	<b>5</b>	<b>1</b>	<b>2</b>	<b>16</b>	<b>9</b>	<b>0</b>	<b>0</b>	<b>15</b>	<b>60</b>
<b>Total inventories</b>	<b>60</b>									

*The intersections between lines and columns whose value is zero, indicate areas with little representativeness in the landscape and consequently an absence of samplings.*

**Table 1.** Number of samplings performed at different altitudinal levels, topographic positions and solar exposures, derived from the sampling system.

considered as juveniles and shrubs. The variables recorded in the site were: altitude, slope (in %), solar exposure (N, S, E, W), physiography (flat land, hillock, plateau, middle slope, high slope, ravine bottom, creek), coverage ( $c1 = \leq 10\%$ ;  $c2 = 11-30\%$ ;  $c3 = 31-50\%$ ;  $c4 = 51-70\%$  y  $c5 = \geq 70\%$ ) and geofom. Management variables related to land use (no use, forest exploitation, wildlife management, grazing, agriculture and conservation) were considered as well as intensity of use (null, moderate, over-exploited and not determinable). Each one of the sampling points were geographically located by Transverse Mercator Units (UTM).

In order to identify the oak and conifer species in the field, keys generated by De la Cerda [17] and Siqueiros [18], respectively, were used. The unknown species were collected in botanical presses and identified at the Autonomous University of Aguascalientes herbarium (HUAA). To leave evidence of the new species records in the ANP SF, specimens were deposited in the HUAA.

### 2.1.3 Distribution and abundance of species

To estimate the distribution of tree and shrub forest species, the presence of each of the species found in each of the 60 sampling sites was quantified. In the case of species considered as restricted distribution (eg. *Quercus cocolobifolia*, *Pinus chihuahuana*, and *P. duranguensis* var. *quinquefoliata*), samples were taken at specific sites ( $n = 4$ ), according to the information provided by De la Cerda [17] and Siquéiros [18]. Species with a wide distribution were those that occurred in the greatest number of sites.

The frequency of the species found was determined on 100 m transect at ground level, observing 100 separate points every meter. The species found at each point were recorded (when there was more than one vegetation layer), counting the number of times that each species appeared (absolute frequency) [16] over the whole transect. Relative frequency was calculated using the Equation [19]:

$$\text{Relative frequency} = \left( \frac{\text{Species frequency}}{\sum \text{Frequency values of all species}} \right) \quad (1)$$

Where:

Frequency of the species x = absolute frequency obtained from each site sampling.

Subsequently, an abundance index was calculated using the equation:

$$\text{Spp.ai} = \frac{\sum \text{relative frequencies}}{\text{Number of sampled sites}} \quad (2)$$

Where:

Spp.ai = Identified Species abundance index.

With this data, distribution and abundance graphs of the main arboreal-shrub forest species were created. The phytoecological analysis was used to calculate the species richness and the Shannon index diversity ( $H'$ ) and the beta Whittaker's ( $\beta_w$ ) index respectively, the first were calculated as a function of the altitudinal level, the second also incorporating the geofom using the Species Diversity and Richness® (Pisces Conservation LTD) software. Pear calculate the indexes we used the equation:

$$H' = - \sum_{i=1}^S p_i \log_2 p_i \quad (3)$$

Where

$S$  = species richness;  $P_i$  = proportion of the individuals of species  $i$  with respect to the total number of individuals;  $n_i$  = number of individuals of species  $i$

$$\beta_w = \frac{S}{S} \quad (4)$$

Where:

$S$  = Species richness and  $S$  = mean richness of the site.


## 2.2 Dry Tropical Forest (DTF)

### 2.2.1 Study area

Although there are some studies that suggest the existence of relics of Dry Tropical Forest (DTF) vegetation in some municipalities of the Aguascalientes State [15, 20], this ecosystem has a greater representation both in surface area and in its conservation status in Calvillo municipality. The study was conducted in 26 sites with DTF vegetation cover in Terrero de la Labor ejido, located within the Sierra Fria Protected Natural Area, in the Municipality of Calvillo, State of Aguascalientes, in Central Mexico. The ejido polygon comprises an area of 5,861 ha. [21], of which, the DTF occupies 45% of its total area (**Figure 3**). It is located within the extreme coordinates: 102°43'58.88" West Longitude and 22°6'4.78" North Latitude and at the Southeast end 102°41'24.95" West Longitude and 21°44'27.61" North Latitude.

### 2.2.2 Selection of the study sites and sampling design

We used a stratified sampling design system [16]. Sampling strata were delimited based on geofoms, slope, exposure and altitude. To characterize geofoms, three criteria were used: concave, convex and flat terrain. A concave geofom was defined when the slope ranged between 10 and 25%, which usually corresponded to ravines and small depressions. When the sites had a slope between 25 and 60% they were characterized as convex sites. Flat terrains had slopes <10%.



**Figure 3.** Location of the study area. (A) Mexico, (B) state of Aguascalientes, (C) municipality of Calvillo and (D) Terrero de la labor Ejido.




Solar exposure was defined using an exposure map made with a Geographic Information System from a 2008 Spot® satellite image and a digital elevation model (MDE). Only the main cardinal points (North, South, East and West) were considered. To locate the altitudinal strata, the contours of the zone defined from the MDE were used. Subsequently, a grid map was developed for the identification of the sampling areas (See **Figure 4**).

### 2.2.3 Selection and characterization of sites to quantify of the composition and abundance of woody species

We established 26 sites to quantify phytoecological inventories, distributed in the landscape according to the above mentioned sampling system. At each point, the projected coordinates of the site were taken with GPS Garmin 48 XL line in UTM format, zone 13 North and with reference Datum WGS84 and with accuracies of 5 to 12 m with differential kinematic adjustment (WAAS). Subsequently, the points were placed on a SPOT 2010 satellite image (**Figure 5**). Site variables considered were the slope (%), solar exposure, physiography of the terrain, intensity and type of use and canopy coverage.


Slope at each sampling site was obtained by direct field measurement with a Brunton clinometer with a precision of  $\pm 2^\circ$  of variation for each 100 meters of length. This data in turn was contrasted with the data obtained from the digital elevation model with precision of 1 to 2 meters in the Z value. Five classes were used to define the slope: i)  $<10\%$ , ii) 11–30, iii) 31–50, iv) 51–70 and v)  $>70$ . Exposure to solar radiation was estimated considering the cardinal points North (N), South (S), East (E) and West (O), as well as their combinations.

The altitude of each site was obtained directly in the field with the support of a GPS with barometric adjustment to reduce the effect of mathematical variation of the Geoid model and with precision of 1 to 3 meters. This was compared with the data obtained from the prospecting of points against elevation level curves obtained



**Figure 4.**  
*Geographic representation of dry tropical Forest and the sampling points in Terrero de la labor Ejido, in the municipality of Calvillo, Aguascalientes.*





**Figure 5.** Ipsographic model of Ejido Terrero de la labor Ejido polygon, and distribution of the sampling points in the DTF.

from the digital elevation model to reduce the potential errors of direct measurements.

The physiography of the terrain was characterized considering flat terrain (slope < 10%), steep (without slope), medium slope (10–25%) and high slope (>60%). The exposure of the sites was quantified with a compass and the magnetic north was taken as reference for its definition in the previously defined ranges. Exposure for each stand of the sampling site was also analyzed along with the digital model of exposures generated from the digital elevation model. The **Table 2** shown the sample points distributed in the landscape of the Dry Tropical Forest.

Other characteristics considered in the description of the sites were the degree of modification (i.e. transformation of geographical space, introduction of species), its intensity (light, medium and overexploited), as well as the type of use by local inhabitants (hunting, grazing, gathering, etc.).

#### 2.2.4 Species richness

To describe species composition, we used a sampling design based on nested plots in an area of 1024 m<sup>2</sup> in each inventory, using the criteria of the minimum area [16]. We started with a plot of 1 x 1 m in a direction perpendicular to the slope in which all present species were recorded, and subsequently, the plot. Subsequently, the plot was increased in size to 2 X 1, 2 X 2, 2 X 4, 4 X 4 m etc. registering the new species for each increment in the area of the squares until reaching the maximum

Meters above sea level (masl)	Topographic position										Total
	Concave coverage (%)					Convex coverage (%)					
	<10	11-30	31-50	51-70	>71	<10	11-30	31-50	51-70	>71	
1800-1900	0	0	0	0	1	0	1	0	3	0	5
1901-2000	0	0	0	0	0	1	1	3	5	1	11
2001-2100	0	0	0	0	0	1	1	2	2	0	6
>2100	0	0	0	0	0	1	1	0	4	0	4
Total inventories	0	0	0	0	1	0	0	5	14	1	26

**Table 2.**  
 Distribution of samplings sites according to the proposed design.

extension (i.e.: 32 x 32 m = 1024 m<sup>2</sup>), to obtain an area/species curve. We then identified the area in which the present species stabilized. This sampling method increased the probability of finding rare species as the area increased, an effect known as Rarefaction [22].

Identification of species was estimated in the field by morphological characters described in previous studies. Specimens that could not be identified in the field were collected and later identified in the Herbarium of the Autonomous University of Aguascalientes (HUAA).

We used the linear intercept survey method (Canfield line). A 100 m long line was perpendicular to the slope, starting at the GPS coordinates of the sampling site, then intersection lines were defined where individuals of DTF species were counted at constant intervals of 1 meter. Shrub and tree individuals were categorized into five heights classes 0-1 m, 1.1-2 m, 2.1-4 m, 4-8 m, 8-15 m and > 16 m. For each class we measured canopy cover of each species by measuring the perpendicular projection of the crown and the frequency of species. To estimate crown cover the following formula was used:

$$\text{Cover (C)} = \frac{\sum \text{length of individuals of species } i}{\text{total length of intersections}} \times 100.$$

To estimate frequencies, we used the formula:

$$\text{Frequency (F)} = \frac{\sum \text{of number of times that individuals of the species intercepted by the line}}{\sum \text{total species intercepted}} \times 100.$$

### 2.2.5 Data analysis

Species composition was estimated through the identification of the species found in each of the sampling plots. To find a limit on the number of samples and to reduce the possibility of under- or over-sampling, we conducted a rarefaction analysis. The Shannon-Wiener alpha diversity ( $H'$ ) was calculated for each of the sites and for each altitudinal level using the Richness and diversity species® software, considering that there could be variation in diversity according to the change in environmental conditions in temperature and precipitation as mentioned in the Standard Atmospheric Index (decrease of 0.6°C/100 m altitude).

The formula of the Shannon index is:

$$H' = - \sum_{i=1}^S p_i \log_2 p_i \quad (5)$$

where:

- $S$ – Total number of species (species richness)
- $p_i$ – Proportion of individuals of species  $i$  in respect to total of individuals (i.e.: relative abundance of species  $i$ ):  $\frac{n_i}{N}$
- $n_i$ – number of individuals of species  $i$
- $N$ – Total number of individuals of all species

The index considers the number of species present in the study area (species richness), and the relative number of individuals of each of those species (abundance).

To estimate replacement rates of species Whitakker's  $\beta$  diversity was computed, using the diversity found for each altitudinal level analyzed as reference.

$$\beta = \frac{S}{\alpha - 1} \quad (6)$$

Where:  $\beta$  = Whitakker's  $\beta$  diversity.

$S$  = Total number of species in samples.

$\alpha$  = Mean number of species in samples.

### 3. Results

#### 3.1 Temperate Mountain Forest

##### 3.1.1 Richness and diversity species

In the 60 sites, 50 species were recorded, corresponding to 20 families and 27 genera (**Table 3**), of which, due to their structure, 47% ( $n = 24$ ) were considered trees (height  $\geq 3.5$  m) and 53% ( $n = 27$ ) shrubs and juveniles. The best represented families were *Fagaceae* (11 species), *Pinaceae* (8 species) and *Ericaceae* (5 species). The species *Q. obtusata* Bonpl., *J. duranguensis* Martínez and *Crataegus* sp. are new reports in the Sierra Fria.

On average, the highest  $H'$  diversity index is found in sites whose altitude ranges between 2400 and 2600 and 2600–2800 mamsl ( $H' = 1.48$  y  $1.63$ , respectively), the former associated with ravines and difficult access places; the second index corresponds to places with higher moisture content and without use. The lowest indexes ( $H = 1.22$  y  $1.36$ ) were found in altitudinal ranges of 2200–2400 and 2000–2200 m, respectively, located on flat lands with intensive management and high resource use rates. According to the geoform, the diversity Wittaker's  $\beta$  was greater on the convex sites ( $\beta_w = 5.80$ ), followed by the concave sites ( $\beta_w = 4.27$ ) and flat lands ( $\beta_w = 4.04$ ). According to the altitudinal level, the highest diversity was found in the sites whose altitude ranges between 2,400 and 2,600 m ( $\beta_w = 7.22$ ), mainly in ravines and places hard to access. In contrast, the lowest indexes were found on site with an altitude lower than 2, 400 m ( $\beta_w = 4.52$ ), located on flat lands, under intensive management and easy access.

In the **Figure 6**, we shown an example of dominant vegetation in Temperate Mountain Forest (in conifers, *Pinus leiophylla* and *P. teocote* in order) in the Sierra Fria Protected Natural Area.

Species	Key	Common name*	Family	Forest classification**	Use <sup>†</sup>	Report***
<i>Acacia farnesiana</i>	Acafar	Huizache	Leguminosae	Tr	Nu	Y
<i>Asclepias linearis</i>	Aline	Romerillo	Apocynaceae	Sh	Nu	Y
<i>Arbutus arizonica</i>	Aariz	Madroño	Ericaceae	Sh	Fe	Y
<i>Arbutus xalapensis</i>	Axala	Madroño rojo	Ericaceae	Sh	Fe	Y
<i>Arbutus glandulosa</i>	Aglan	Madroño blanco	Ericaceae	Sh	Fe	Y
<i>Arctostaphylos pungens</i>	Apun	Manzanita	Ericaceae	Sh	Fe	Y
<i>Buddleia scordioides</i>	Bsco	Vara blanca	Compositae	Sh	Fr	Y
<i>Buddleia cordata</i>	Bcor	Tepozan	Compositae	Tr	Fr	Y
<i>Bursera fagaroides</i>	Burfaga	Venadilla	Burseraceae	Sh	Nu	Y
<i>Comerostaphyllis spp.</i>	Comesp	Pacuato	Ericaceae	Sh	Med	Y
<i>Dalea bicolor</i>	Dabic	Engordacabra	Fabaceae	Sh	Fr	Y
<i>Dasyliirion acotriche</i>	Dasaco	Sotol	Agavaceae	Sh	Nu	Y
<i>Dodonaea viscosa</i>	Dovisc	Jarilla	Sapindaceae	Sh	Med	Y
<i>Eucaliptus camaldulensis</i>	Eucamal	Eucalipto	Myrtaceae	Tr	Nu	Y
<i>Fraxinus uhdei</i>	Frauhd	Fresno	Oleaceae	Tr	Nu	Y
<i>Garria ovata</i>	Garova	planta peluda	Garryaceae	Sh	Nu	Y
<i>Pinus chihuahuana</i>	Pinchi	Pino Prieto	Pinaceae	Tr	Nu	Y
<i>Pinus duranguensis Mart.</i>	PiduM	Pino verde	Pinaceae	Tr	Ew	Y
<i>Pinus duranguensis f. quinquefoliata</i>	PiduQ	Pino verde	Pinaceae	Tr	Nu	Y
<i>Pinus leiophylla</i>	Pile	Pino Prieto	Pinaceae	Tr	Nu	Y
<i>Pinus lumholtzii</i>	Pilum	Pino llorón	Pinaceae	Tr	Nu	Y
<i>Pinus michoacana</i>	Pimich	Pino barbón	Pinaceae	Tr	Nu	Y
<i>Pinus cembroides</i>	Picem	Pino chaparro	Pinaceae	Tr	Nu	Y
<i>Prosopis laevigata</i>	Prolae	Mesquite	Pinaceae	Sh	Nu	Y
<i>Pinus teocote</i>	Pinteo	Pino	Pinaceae	Tr	Ew	Y
<i>Jatropha dioica</i>	Jadio	Sangre de grado	Euphorbiaceae	Sh	Nu	Y
<i>Juniperus flacida</i>	Jufla	Olmo triste	Cupresaceae	Sh	Nu	Y
<i>Juniperus deppeana</i>	Judep	Táscate	Cupresaceae	Tr	Fe-Tu	Y
<i>Juniperus duranguensis</i>	Judur	Cedro chino	Cupresaceae	Sh	Nu	New
<i>Opuntia leucotricha</i>	Opuleu	Nopal duraznillo	Cactaceae	Sh	Nu	Y
<i>Opuntia streptacantha</i>	Opust	Nopal cardón	Cactaceae	Sh	Nu	Y
<i>Prunus serotina</i>	Pruser	Cerezo negro	Rosaceae	Sh	Ft	Y
<i>Quercus cocolobifolia</i>	Queco	Palo manzano	Fagaceae	Tr	Fe	Y
<i>Quercus chihuahuensis</i>	Quechih	Palo blanco	Fagaceae	Tr	Fe	Y



Species	Key	Common name*	Family	Forest classification**	Use <sup>†</sup>	Report***
<i>Quercus laeta</i>	Quela	Palo blanco	Fagaceae	Tr	Fe	Y
<i>Quercus grisea</i>	Quegri	Palo chino	Fagaceae	Tr	Fe	Y
<i>Quercus potosina</i>	Quepo	Palo chaparro	Fagaceae	Tr	Fe	Y
<i>Quercus microphylla</i>	Quemic	Chaparrito	Fagaceae	Sh	Nu	Y
<i>Quercus resinosa</i>	Queres	Encino hojudo	Fagaceae	Tr	Le	Y
<i>Quercus rugosa</i>	Querug	Palo blanco	Fagaceae	Tr	To	Y
<i>Quercus sideroxyla</i>	Quersid	Palo rojo	Fagaceae	Tr	Fe- le	Y
<i>Quercus eduardii</i>	Queredu	Palo rojo	Fagaceae	Tr	Fe- le	Y
<i>Quercus sp.</i>	Encino 1	Encino	Fagaceae	Sh	Fe	Y
<i>Quercus obtusata</i>	Querobt	Encino	Fagaceae	Sh	Fe	New
<i>Yucca filifera</i>	Yufi	Palma	Agavaceae	Sh	Nu	Y
<i>Odontotrichum amplum</i>	Adoamp	Vaquerilla	Asteraceae	Sh	Nu	Y
<i>Phytocellobium leptophyllum</i>	Phylep	Gatuño de la sierra	Leguminosae	Sh	Nu	Y
<i>Eisenhardtia polystachya</i>	Eipol	Varaduz	Fabaceae	Sh	Nu	Y
<i>Crataegus spp.</i>	Crasp	Tejocote	Rosaceae	Tr	Nu	New
<i>Quercus sp-2</i>	Encino 2	Encino	Fagaceae	Sh	Nu	Y
<i>Ipomoea stans</i>	Ipost	Galuzá	Convolvulaceae	Sh	Nu	Y

\*The common names were provided by the habitants of “La Congoja” community and do not necessarily correspond to the common name in other localities where these species could be found.

\*\*Within the forest classification, Tr = Tree and Sh = Shrub.

\*\*\*The reports correspond to the flora identified previously, the new reports correspond to the individuals identified in this study.

<sup>†</sup>The use of the forest species recorded, depends on the forest managers experience, in this way, Nu = no use; Fe = firewood extraction; Fr = use as forage plant; Med = Medicinal use; Pt = Timber use; for extraction as fence pole; Ew = extraction as wood stripe from P. teocote; le = leave extraction for ornamentals; To = tools.

**Table 3.**  
Forest species identified in an area of the SF-natural protected area, Aguascalientes.

### 3.1.2 Distribution and abundance of species


The most widely distributed species belong to the genus *Juniperus* (locally known as cedros or táscates), *Quercus* (oaks) and *Arbutus* (locally known as madrone). *J. deppeana* is the most widely distributed species followed by *Quercus potosina* and by *Arctostaphylos pungens*. The madrones (*Arbutus xalapensis* and *A. glandulosa*) appear in fourth and fifth place, respectively (**Figure 7**).

Out of 50 recorded species, 6 are the ones with the highest abundance indexes. *Q. potosina*, the species best represented in the landscape. This species presents the highest abundance index (ia = 0.1585), followed by *J. deppeana* (ia = 0.1102) which also presents the widest distribution. Inside the genus *Pinus*, *P. leiophylla* is the most abundant, even above manzanita (*Arctostaphylos pungens*) and red oaks (*Q. sideroxyla* y *Q. eduardii*; **Figure 8**).

There are species such as *Pinus chihuahuana*, *Pinus lumholtzii* and *Pinus duranguensis* that present restricted distribution, but are abundant in very specific




**Figure 6.**  
 Typical vegetation of the Temperate Mountain Forest. (a) Landscape dominated by conifers in the Sierra Fria, in this case, by *Pinus leiophylla*; (b) wild ash twig (*Fraxinus uhdei*); (c) Manzanita (*Arctostaphylos pungens*) specimen in a plateau of the Sierra Fria; (d) oak specimen locally known as Palo chino (*Quercus grisea*).



**Figure 7.**  
 Forest species with wide distribution in the study area inside the SF-natural protected area.

sites. The distribution analysis based on the altitudinal gradients and geofom suggests that the altitudinal stratum between 2 000 and 2 200 m is the one with the lowest tree and shrub species richness. The best represented species in this range belong to the xeric shrubland being three of them such as *Dodonaea viscosa*, *Phytocellobium leptophyllum*, and *Odontotrichum amplum*, considered as overgrazing indicator species [22]. From the second stratum (2 200 to 2 400 mamsl) *Pinus* and



**Figure 8.**

Abundance indexes from the species best represented in the SF-natural protected area, the most representative ecosystem of Temperate Mountain Forest in Ags. The X axis represents the abundance index which ranges between 0.0670 (*Quercus eduardii*) and 0.1585 (*Q. potosina*). The maximum value of the abundance index could be 1.

*Quercus* species begin to appear, although isolated *Quercus resinosa* individuals can be found at higher altitudes (Table 4).

Out of the dominant conifer species at the SF-Natural Protected Area, *Pinus leiophylla* and *P. teocote* are distributed at altitudes ranging from 2 400 to 2 600 masl. Between 2 600 and 2 800 masl these two species are more dispersed and located mainly in ravines. *P. leiophylla* is also located on plateaus at 2 700 m (e.g. Mesa del Águila and Mesa del Aserradero). Red oaks (*Q. eduardii* and *Q. sideroxyla*) are distributed at altitudes from 2 400 to 2 600 m, mainly along the ravines (Table 3).

In Figure 9 we shown some species of *Pinus* genera dominants in the intermediate altitudinal strata of SF-Protected Natural Area.

## 3.2 Dry Tropical Forest

### 3.2.1 Richness and diversity woody species

We identified 79 species of trees and shrubs, within 45 genera and 14 families (see Table 5). The best represented families were Fabaceae (13 genera), Asteraceae (11 genera) and Cactaceae (9 genera). The genera better represented were *Opuntia* (n = 4 spp.), *Acacia* (n = 4 spp.) and *Bursera* (n = 3 spp.). The genero *Salvia* is also important.

The  $H'$  diversity found in the DTF of the ejido Terrero de la Labor ejido is constant. The highest diversity index found was 3.49 in two of the 26 analyzed sites, which apparently are well conserved sites. On the contrary, three sites had the lowest  $H'$  diversity index with 2.77 (Table 6). Although there are apparently no differences, the highest diversity indexes are located mainly in ravines and north facing exposures, and in locations with difficult access (see Table 7).

### 3.2.2 Distribution and abundance of woody species in the DTF

Of the 79 species identified, eight are distributed in more than 70% of the plots of Terrero de la Labor ejido. The species with the greater distribution are the



Species	ALTITUDE (MASL) <sup>±</sup>														
	A1		A2			A3		A4		A5					
	2	2.1	2.19	2.2	2.3	2.39	2.4	2.5	2.59	2.6	2.7	2.79	2.8	2.9	3
<i>Arctostaphylos pungens</i>															
<i>Dodonaea viscosa</i>															
<i>Juniperus deppeana</i>															
<i>Quercus potosina</i>															
<i>Bursera fagaroides</i>															
<i>Eisenhardtia polystachya</i>															
<i>Juniperus flacida</i>															
<i>Acacia farnesiana</i>															
<i>Prosopis laevigata</i>															
<i>Arbutus glandulosa</i>															
<i>Quercus resinosa</i>															
<i>Yucca filifera</i>															
<i>Phytocellobium leptophyllum</i>															
<i>Asclepias linearis</i>															
<i>Quercus eduardii</i>															
<i>Odontotrichum amplum</i>															
<i>Pinus leiophylla</i>															
<i>Pinus teocote</i>															
<i>Quercus rugosa</i>															
<i>Quercus chihuahuensis</i>															
<i>Quercus sideroxyla</i>															
<i>Arbutus xalapensis</i>															
<i>Pinus lumholtzii</i>															
<i>Juniperus duranguensis</i>															
<i>Quercus coccolobifolia</i>															
<i>Quercus grisea</i>															
<i>Quercus laeta</i>															

\*The species distribution in different altitudinal gradients was as a function of the 10 dominant species (obtained from the frequency/site) at each altitudinal stratum.

<sup>±</sup>Altitudes (A1-A5) are calculated in m\* 1000.

<sup>‡</sup>The bars with gray shades indicates that this species is abundant at the altitudinal gradient where it was found. In contrast, the black shades indicate that although this species is not abundant, it was found in.

**Table 4.**  
 Dominant species distribution by altitudinal strata.

*Myrtillocactus geometrizans* (garambullo), *Ipomoea murucoides* (palo bobo), *Eysenhardtia polystachya* (varaduz), *Bursera fagaroides* (venadilla), and *Forestiera phillyreoides* (palo blanco) (Figure 10), which were located in 96, 92, 90, 88 and 86% of the plots respectively, assuming that the sampling sites are representative of the entire landscape.

On the other extreme, the rarest species were *Plumeria rubra*, *Ficus petiolaris* and *Fraxinus purpurea*. The first species was only located in one site, while the last two





**Figure 9.**

Populations of *Pinus* (*Pinus spp*) at the SF-natural protected area. The photograph on the left side shows a *Pinus leiophylla* population at the Barranca de Piletas. The photograph in the right side shows an image of *Pinus duranguensis*. Photographs as courtesy of Clemente Villalobos llamas and Vicente Díaz Núñez.

Species	Family	Common name
<i>Acacia berlandieri</i> Benth.	Fabaceae	Carbonera
<i>Acacia farnesiana</i> (L.) Willd.	Fabaceae	Tepame
<i>Acacia pennatula</i> (Schltdl. & Cham.) Benth.	Fabaceae	Huizache o Cascalote
<i>Agave angustifolia</i> Haw.	Asparagaceae	Lechuguilla
<i>Albizia plurijuga</i> (Standl.) Britton & Rose	Fabaceae	Tepeguaje blanco
<i>Alnus acuminata</i> Kunth	Betulaceae	Aile
<i>Amelanchier denticulata</i> (Kunth) K. Koch	Rosaceae	Duraznillo
<i>Amphipterygium molle</i> (Hemsl.) Hemsl. & Rose	Anacardiaceae	Cuachalalate
<i>Asclepias linaria</i> Cav.	Apocynaceae	Algodoncillo
<i>Ayenia mexicana</i> Turcz.	Sterculioideae	
<i>Baccharis heterophylla</i> Kunth	Asteraceae	Escobilla
<i>Bouvardia multiflora</i> (Cav.) Schult. & Schult. f.	Rubiaceae	Clavelito
<i>Brickellia veronicifolia</i> (Kunth) A. Gray	Asteraceae	Orégano de monte
<i>Buddleja cordata</i> Kunth	Buddlejaceae	Tepozán blanco
<i>Buddleja sessiliflora</i> Kunth	Buddlejaceae	Tepozán verde
<i>Bursera bipinnata</i> Donn. Sm.	Burseraceae	Lantrisco
<i>Bursera fagaroides</i> (Kunth) Engl.	Burseraceae	Venadilla
<i>Bursera penicillata</i> (DC.) Engl.	Burseraceae	Arbol de chicle
<i>Calliandra eriophylla</i> Benth.	Fabaceae	Calandria
<i>Castilleja tenuifolia</i> M. Martens & Galeotti	Scrophulariaceae	Hierba del cancer
<i>Cedrela dugesii</i> S. Watson	Meliaceae	Cedro

Species	Family	Common name
<i>Ceiba aesculifolia</i> (Kunth) Britten & Baker f.	Malvaceae	Pochote
<i>Celtis caudata</i> Planch.	Ulmaceae	Capulincillo
<i>Celtis pallida</i> Torr.	Ulmaceae	Vara en cruz
<i>Chusquea</i> sp	Poaceae	Camalote
<i>Colubrina triflora</i> Brongn. Ex G. Don	Rhamnaceae	Algodoncillo
<i>Cordia sonora</i> Rose	Boraginaceae	Amapa o Vara prieta
<i>Croton ciliatoglandulifer</i> Ortega	Euphorbiaceae	Algodoncillo
<i>Dasyilirion acrotrichum</i> (Schiede) Zucc.	Asparagaceae	Sotol
<i>Dodonaea viscosa</i> Jacq.	Sapindaceae	Jarilla
<i>Erythrina flabelliformis</i> Kearney	Fabaceae	Colorín
<i>Eupatorium</i> sp	Asteraceae	Copalillo
<i>Eysenhardtia polystachya</i> (Ortega) Sarg.	Fabaceae	Palo azul o Varaduz
<i>Eysenhardtia punctata</i> Pennell	Fabaceae	Palo cuate
<i>Ferocactus histrix</i> Lindsay	Cactaceae	Biznaga costillona
<i>Ficus petiolaris</i> Kunth	Moraceae	Ficus silvestre
<i>Forestiera phillyreoides</i> (Benth.) Torr.	Oleaceae	Palo blanco
<i>Fraxinus purpusii</i> Brandege	Oleaceae	Saucillo
<i>Gymnosperma glutinosum</i> (Spreng.) Less.	Asteraceae	Cola de zorra
<i>Heliocarpus terebinthinaceus</i> (DC.) Hochr.	Meliaceae	Cicuito o Cuero de indio
<i>Ipomoea murucoides</i> Roem. & Schult.	Convolvulaceae	Palo bobo
<i>Iresine</i> sp.	Amaranthaceae	Cola de zorra
<i>Jatropha dioica</i> Sessé	Euphorbiaceae	Sangregrado
<i>Koanophyllon solidaginifolium</i> (A. Gray) R. M. King & H. Rob.	Asteraceae	Caballito
<i>Karwinskia humboldtiana</i> (Schult.) Zucc.	Rhamnaceae	Coyotillo
<i>Leucaena esculenta</i> (Moc. & Sessé ex DC.) Benth.	Fabaceae	Guaje rojo
<i>Lippia inopinata</i> Moldenke	Verbenaceae	Palo oloroso
<i>Lysiloma acapulcense</i> (Kunth) Benth.	Fabaceae	Ébano o Palo fierro Tepeguaje
<i>Lysiloma microphyllum</i> Benth.	Fabaceae	Tepeguaje
<i>Mammillaria bombycina</i> Quehl	Cactaceae	Biznaga de seda
<i>Mammillaria</i> sp.	Cactaceae	Biznaga
<i>Manihot caudata</i> Greenm.	Euphorbiaceae	Pata de gallo
<i>Mimosa monancistra</i> Benth.	Fabaceae	Gatuño o Uña de gato
<i>Mimosa</i> sp.	Fabaceae	Huizache
<i>Myrtillocactus geometrizans</i> (Mart. ex Pfeiff.) Console	Cactaceae	Garambullo
<i>Montanoa leucantha</i> (Lag.) S.F. Blake	Asteraceae	Talacao o Vara blanca
<i>Opuntia leucotricha</i> DC.	Cactaceae	Nopal chaveño o duraznillo
<i>Opuntia robusta</i> J.C. Wendl.	Cactaceae	Tuna tapona
<i>Opuntia</i> sp.	Cactaceae	Nopal
<i>Opuntia streptacantha</i> Lem.	Cactaceae	Nopal cardón

Species	Family	Common name
<i>Perymenium mendezii</i> DC.	Asteraceae	
<i>Pistacia mexicana</i> Kunth	Anacardiaceae	Lantrisco
<i>Pittocaulon filare</i> (McVaugh) H. Rob. & Brettell	Asteraceae	Palo loco
<i>Plumbago pulchella</i> Boiss	Plumbaginaceae	Chilillo medicinal
<i>Plumeria rubra</i> L.	Apocynaceae	Flor de mayo
<i>Prosopis laevigata</i> (Humb. & Bonpl. ex Willd.) M.C. Johnst.	Fabaceae	Mezquite
<i>Ptelea trifoliata</i> L.	Rutaceae	Naranja agrio o Zorrillo
<i>Quercus laeta</i> Liebm.	Fagaceae	Roble blanco
<i>Salvia mexicana</i> L.	Labiatae	Tlacote
<i>Salvia</i> sp.	Labiatae	Salvias
<i>Stachys coccínea</i> Ortega	Labiatae	Mirto
<i>Stenocereus queretaroensis</i> (F. A. C. Weber) Buxb.	Cactaceae	Pitahaya
<i>Tecoma stans</i> (L.) Juss. ex Kunth	Bignoniaceae	Tronadora
<i>Trixis angustifolia</i> DC.	Asteraceae	Vara verde
<i>Verbesina serrata</i> Cav.	Asteraceae	Vara blanca
<i>Viguiera quinqueradiata</i> (Cav.) A. Gray ex S. Watson	Asteraceae	Vara amarilla
<i>Wimmeria confusa</i> Hemsl.	Celastraceae	Algodoncillo
<i>Yucca filifera</i> Chabaud	Asparagaceae	Palma
<i>Zanthoxylum fagara</i> (L.) Sarg.	Rutaceae	Rabo lagarto

**Table 5.**  
List of species identified in the dry tropical Forest of Terrero de la labor Ejido, Calvillo, Ags.


Altitud level (masl)	Sampled sites	H'
1851–1900	7	3.08
101–1950	5	2.57
1951–2000	7	3.60
2001–2100	5	3.14
>2100	2	3.25

**Table 6.**  
Average H' diversity indices associated to different altitudinal ranges in the DTF of Terrero de la labor Ejido.


were only found in two and three sampling sites, respectively. Their low frequency could be associated to their presence in mid statured forests. The most abundant species are those that, even though they are not those with a wide distribution in the landscape, in the places where they are located their frequency is higher than the rest of the identified species. In the DTF of the Terrero de la Labor, the most abundant species belonged to five different genera, of which the most important are *Lysiloma microphylla* (tepeguaje), *Ipomoea murucoides* (palo bobo), and *Bursera fagaroides* (locally known as venadilla) (**Figure 7**). In the case of *Ipomoea murucoides*, it occupies the second place in both distribution and abundance (see **Figure 11**).

Slope range (%)	Sampled sites	H'
0–9	6	2.94
9–25	3	2.88
26–37	5	2.92
37–49	4	3.26
49–64	5	3.34
>65	3	3.39

**Table 7.**  
 Diversity indexes associated to different slopes of the sites.



**Figure 10.**  
 Species best represented in the DTF Terrero de la labor Ejido.




**Figure 11.**  
 List of species with the highest abundance in Terrero de la labor Ejido.

The **Figure 12** shown some species of the dominant vegetation in tropical dry forest, in this case, of the Terrero de la labor and las Moras ejidos in the Municipality of Calvillo, Aguascalientes State.



#### 4. Discussion

The loss of biodiversity is one of the environmental problems that has managed to arouse broad global interest in the last two decades [4, 23]. Some of the main



**Figure 12.** Diversity of forest species in the tropical dry Forest. (a) Landscape of the tropical dry Forest in the Terrero de la labor and las Moras Ejido; (b) an example of *Manihot caudata*, locally known as jaboncillo; (c) specimen of *Bursera fagaroides* (locally known as venadilla or papelillo). Photographs courtesy of Vicente Díaz Núñez, Joaquín Sosa-Ramírez and Jesús Argumedo-Espinoza.

causes are related to human activities, either directly (overexploitation) or indirectly (habitat alteration), although there is generally an interaction between them. The communication systems have impacted in such a way that both the government and the private sector, as well as society in general, consider a priority to direct greater efforts towards conservation programs. The basis for an objective analysis of biodiversity and its change lies in its correct evaluation and monitoring.

In the Temperate Mountain Forest, the 50 woody species identified show a high species richness in comparison with other mountain regions. The best represented genera correspond to oak trees (*Quercus spp.*) and pines (*Pinus spp.*). The studied area harbors a small portion (6.8%) of the oak species that inhabit Mexico (161 species) [24]; although, this percentage is lower than those reported in areas with a greater territorial surface and higher rainfall, such as the case of San Luis Potosi and Jalisco States, which has identified 45 and 51 oak species respectively, that represent 27.95 and 36.9% of the total oak species registered in Mexico [25], the SF-NPA represent less 5% of the territorial surface in the mentioned states. In relation to pines, the studied area has about 17% of the species identified in Mexico [26]. This proportion is similar to that reported by Márquez-Linares et al., [27] in an area of pine-oak forest, in Durango, Mexico, where they recorded 8 pine species. In relation to “Las Joyas” scientific station, in the Sierra de Manantlán Biosphere Reserve, the *Quercus* diversity (16 spp.) is similar to the one in the Sierra Fria, although the area of las Joyas is smaller (Ca. 3600 ha.).

In the Sierra Fría, the most widely distributed and abundant species are the potosine oak (*Q. potosina*) and alligator juniper (*J. deppeana*). In the case of *Q. potosina*, its distribution and abundance may be related to the dominant physiography in this area, as well as to the mean annual precipitation (650 mm). The appearance of *J. deppeana* is possibly related to the disturbances that occurred in the Sierra Fria during the period between 1920 and 1940 [28]. This species has probably been a pioneer in the recovery of the vegetation cover, although the presence of manzanita (*A. pungens*) has also been documented colonizing sites where disturbances occurred, either natural, as in the case of fires or, anthropogenic, such as forest clearance and harvesting. Pines population is restricted to the Sierra Fría and the Sierra del Laurel. In the Sierra Fria, *Pinus teocote* (locally known as pino ocote) and *Pinus leiophylla* (locally known as pino prieto) are the two most abundant pine species. Its population is abundant in humid places and altitudes higher than 2,500 masl. *P. leiophylla* isolated specimens have been found on flat lands, which suggests that in the past this species had a greater distribution. In the Sierra del Laurel only two pine species have been identified, the pino triste (*Pinus lumholtzii*) and the pino piñonero (*Pinus cembroides var. cembroides*) in isolated populations, which suggests that in the past they were more abundant; however, the existing information is incipient.

The  $H'$  diversity indexes for each altitudinal stratum suggest that, between 2,400 and 2,600 masl, the plant richness of the SF-Natural Protected Area is similar to temperate forests, similar to what the  $\beta$  Whitakker index showed.

The distribution of species such as *J. deppeana* and *Q. potosina*, the most abundant and widely distributed, are influenced by flat sites and canopy covers that vary between 30 and 50%. One explanation is that *Q. potosina* tolerates high drought rates and *J. deppeana* is a pioneer species in disturbed sites, as suggested by Minnich et al. (1994) [28]. On the other hand, the presence or absence of the species may also be dictated due to their dispersal capacity or to the presence or absence of dispersers [19]. The results obtained contribute to describe the habitat of the species, which is an essential factor in programs aiming the restoration and management of temperate climate forests [8, 29], actions that, at least in the case of Mexico, have shown few results.



The species richness in BTS is generally lower than in humid tropical forests [30], although higher than in Temperate Mountain Forests [25]. The BTS is dominated by relatively short trees, most of which lose all their foliage during the dry season. In this community, herbaceous life form, thin woody species, and vines are common, but epiphytes and thick lianas are less abundant and diverse than in humid forests [31]. Diversity is generally higher without a clear dominance of any species, to the point that many of them are rare [32]. In this type of ecosystem, it is common to identify some genera such as *Bursera*, *Lonchocarpus*, *Lysiloma* and *Jatropha*, as well as emerging columnar cacti [33].

The species richness found at the Terrero de la labor Ejido BTS (N = 79) is similar to that reported by Trejo (2005) [33], where he points out that on average the tropical dry forest in Mexico harbors around 74 species with a DBH  $\geq 1$  cm in 0.1 ha. However, in the study site, some species considered “rare” which are indicators of medium forest (e.g *Amphipterygium molle*) were found in ravines and better preserved sites, suggesting that at some point this ecosystem had a greater presence in the landscape.

The analysis of the diversity, distribution and abundance associated with the Tropical Dry Forest in Aguascalientes has been little addressed, so the study conducted in the BTS of the Municipality of Calvillo represents one of the first efforts to understand this ecosystem natural heritage [14]. Previously, partial floristic studies had been carried out, studies which mainly referred to the dominant vegetation types and some important species, however on these studies there were gaps in relation to the ecology of the plant communities [20]. On the other hand, other studies mention some factors related to the mortality of these natural communities [15], but there is no information on vegetation diversity which reflects the real tropical dry forest importance.

This work contributes directly to the management of the ecosystems analyzed. Knowledge about species richness and their distribution provides an overview of the territory's conservation state, considering that both the Temperate Mountain Forest and the Tropical Dry Forest studied are part of the Sierra Fria Protected Natural Area, which is the protected area with the biggest extension in the State. On the other hand, the bases are established for the restoration of degraded ecosystems, either through active restoration or through mechanisms of ecological succession (passive restoration) [29].

## Acknowledgements

The authors acknowledge the participation of Jesus Argumedo-Espinoza for his cartographic support. Likewise, we thank the facilities provided of the owners of the Sierra Fria, as well as Jesus Velasco Serna of the Terrero de la Labor ejido for in the gathering of field information.

# IntechOpen

## Author details

Joaquín Sosa-Ramírez<sup>1\*</sup>, Vicente Díaz-Núñez<sup>2</sup> and Diego R. Pérez-Salicrup<sup>3</sup>

<sup>1</sup> Agricultural Sciences Center, Autonomous University of Aguascalientes, Av. Universidad, Ciudad Universitaria, Aguascalientes, Ags., México

<sup>2</sup> Scientific Research Invited by the Agricultural Sciences Center, Autonomous University of Aguascalientes, Avenida Universidad, Ciudad Universitaria, Aguascalientes, Ags., México

<sup>3</sup> Ecosystem Research Institute, National Autonomous University of Mexico, Morelia, Mich, Mexico

\*Address all correspondence to: [jsosar@correo.uaa.mx](mailto:jsosar@correo.uaa.mx)

## IntechOpen

---

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 



## References

- [1] Sarukhán, J., P. Koleff, J. Carabias, J. Soberón, R. Dirzo, J. Llorrente-Busquets, G. Halfter, R. González, I. March, A. Mohar, S. Anta, J. de la Maza. 2009. Natural Capital of Mexico: current knowledge, evaluation and perspectives of sustainability. Synthesis/Capital Natural de México: conocimiento actual, evaluación y perspectivas de sustentabilidad. Síntesis. México: CONABIO. 100 p.
- [2] Llorente-Bousquets, J., y S. Ocegueda. 2008. State of knowledge of biota. In: CONABIO (2008). Natural Capital of Mexico, vol. I: Current knowledge of biodiversity/Estado del conocimiento de la biota. En: CONABIO (2008). Capital natural de México, vol. I: Conocimiento actual de la biodiversidad. CONABIO, México, 2008.
- [3] Mora, F. 2019. The use of ecological integrity indicators within the natural capital index framework: The ecological and economic value of the remnant natural capital of Mexico. *Journal for Nature Conservation* 47:77–92
- [4] Organización de las Naciones Unidas para la Agricultura y la Alimentación (FAO). 2016. Global Forest Resources Assessment 2015; How are the world's changing? Second edition. FAO. 54 p. Available in: <http://www.fao.org/3/a-i4793e.pdf>
- [5] Organización de las Naciones Unidas para la Agricultura y la Alimentación (FAO). 2010. Evaluación de los Recursos Forestales Mundiales. Informe Principal/Global Forest Resources Assessment. Main Report. FAO. Roma. 381 p.
- [6] Challenger, A. y J. Soberón. 2008. Los ecosistemas terrestres de México. En: Capital natural de México, Vol. I: Conocimiento actual de la biodiversidad, J. Soberón, G. Halfter y J. Llorente (eds.)/Terrestrial ecosystems of Mexico. In: Natural Capital of Mexico, Vol. I: Current knowledge of biodiversity, J. Soberón, G. Halfter and J. Llorente (eds.). Comisión Nacional para el Conocimiento de la Biodiversidad –CONABIO–, México. ISBN 978–607-7607-03-8. p. 87–108.
- [7] Balvanera, P. 2012. Los servicios ecosistémicos que ofrecen los bosques tropicales/ The ecosystem services offered by tropical forests. *Ecosistemas* 21: (1). <http://www.revistaecosistemas.net/articulo.asp?Id=709>
- [8] Baskent, E.Z. 2020. A framework for characterizing and regulating ecosystem services in a management planning context. *Forests* 11: 102; doi:10.3390/f11010102
- [9] Comisión Nacional Forestal (CONAFOR). 2014. National Forest and Soil Inventory 2014–2018. Main report/ Inventario Nacional Forestal y de Suelos 2014–2018. Informe de resultados. SEMARNAT-CONAFOR. 200 p.
- [10] Díaz, V., J. Sosa-Ramírez y D.R. Pérez-Salicrup. 2012. Trees and shrubs Distribution and abundance in Sierra Fria, Aguascalientes, Mexico/ Distribución y abundancia de las especies arbóreas y arbustivas en la Sierra Fría, Aguascalientes, México/. *Polibotánica* 34: 99–126.
- [11] Inventario Estatal Forestal y de Suelos (IEFyS). 2012. State Forest and Soil Inventory of Aguascalientes/ Inventario Estatal Forestal y de Suelos de Aguascalientes. Gobierno del Estado de Aguascalientes-SEMARNAT-CONAFOR. 122 P.
- [12] Gobierno del Estado de Aguascalientes. 2015. Decree of the Protected Natural Area "Área Silvestre Estatal Sierra Fría"/Decreto del Área Natural Protegida "Área Silvestre Estatal

Sierra Fría”. Diario Oficial del Estado de Aguascalientes. Disponible en: <http://servicios2.aguascalientes.gob.mx/NormaTecaAdministrador/archivos/EDO-12-47.pdf>

[13] Díaz-Núñez, V., J. Sosa-Ramírez, and D.R. Pérez-Salicrup. 2016. Vegetation patch dynamics and tree diversity in a conifer and oak forests in Central Mexico. *Botanical Sciences* 94: 229–240.

[14] Argumedo-Espinoza, J., J. Sosa-Ramírez, V. Díaz-Núñez, D.R. Pérez-Salicrup, and M.E. Siqueiros-Delgado. 2017. Diversity, distribution and abundance of woody plants in a dry tropical forest: recommendation for its management. In: Ortega-Rubio (Ed). 2017. *Mexican Natural Resources Management and Biodiversity Conservation*. Springer. Pp. 479–500

[15] Díaz-Núñez, V., J. Sosa-Ramírez e I. P. Macías-Medina. 2014. Phytosanitary diagnosis of vegetation in priority ecosystems of Aguascalientes, Mexico/ Diagnóstico fitosanitario de la vegetación en ecosistemas prioritarios de Aguascalientes, México. Comisión Nacional Forestal-Secretaría de Medio Ambiente del Estado de Aguascalientes. 84 p.

[16] Daget Ph. y M. Godron. 1982. Analysis of the ecology of species in communities/ Analyse de l'écologie des espèces dans les communautés. Masson, Paris. 163p.

[17] De la Cerda, M. E. 1999. Oaks of Aguascalientes/Encinos de Aguascalientes. Universidad Autónoma de Aguascalientes. Segunda Edición. 77 p.

[18] Siqueiros, D.M.E. 1989. Conifers of Aguascalientes/Coníferas de Aguascalientes. Universidad Autónoma de Aguascalientes, 68 p.

[19] Krebs, C. K. 1993. Factors that limit distributions: Dispersal. In *Ecology*, C.

K. Krebs. John Wiley and Sons. p. 41–56.

[20] Siqueiros-Delgado, M.E., J.A. Rodríguez-Ávalos, J. Martínez-Ramírez y J.C. Sierra-Muñoz. 2016. Current status of the vegetation of Aguascalientes, Mexico/ Situación actual de la vegetación del Estado de Aguascalientes/. *Botanical Sciences* 94: 455–470

[21] INEGI. 2007. Agrarian nuclei. Municipality basic tables. Certification Program for Ejido Rights and Land Titling/Núcleos agrarios. Tabulados básicos por municipio. Programa de Certificación de Derechos Ejidales y Titulación de Solares. PROCEDE. México.

[22] Begon, M., C. T. Townsend y J. L. Harper. 2006. *Ecology: from individuals to ecosystems*. Blackwell Publishing, Oxford. 738 p.

[23] Brawman, K.A., and G.C. Daily. 2008. Ecosystem services. *Human ecology*. Elsevier. Pp. 1148–1154

[24] Valencia, A. S. 2004. Diversity of *Quercus* genera (Fagaceae) in Mexico/ Diversidad del género *Quercus* (Fagaceae) en México. *Boletín de la Sociedad Botánica de México* 75: 33–53.

[25] Sabás-Rosales, J.L., J. Sosa-Ramírez, and J.J. Luna-Ruiz. 2015. Diversity, distribution and basic habitat characterization of the Oaks (*Quercus*: Fagaceae) of San Luis Potosí, México/ Diversidad, distribución y caracterización básica del hábitat de los encinos (*Quercus*: Fagaceae) del Estado de San Luis Potosí, México. *Botanical Sciences* 93: 881–897. DOI 10.17129/botsoci.205

[26] Saénz-Romero, C., A. E. Snively y R. Lindig-Cisneros 2003. Conservation and restoration of pine forest genetic resources in Mexico. *Silvae Genetica* 52: 233–237.

- [27] Márquez-Linares, M. A., S. González-Elizondo y R. Álvarez-Zagoya. 1999. Components of tree diversity in pine-oak forests of Durango, Mex./ Componentes de la diversidad arbórea en bosques de pino encino de Durango, Mex. *Madera y Bosques* 5: 67–78.
- [28] Minnich, R. A., J. Sosa-Ramírez, V. E. Franco, W. J. Barry y M. E. Siqueiros. 1994. Preliminar recognizing of vegetation and human impacts in Sierra Fria, Aguascalientes/Reconocimiento preliminar de la vegetación y de los impactos de las actividades humanas en la Sierra Fría, Aguascalientes. *Investigación y Ciencia* 12: 23–29.
- [29] Rey-Benayas, J. M., J. M. Bullock y A. C. Newton. 2008. Creating woodland islets to reconcile ecological restoration, conservation, and agricultural land use. *Frontiers in Ecology and the Environment* 6: 329–336.
- [30] Gentry, A. H. 1995. Diversity and floristic composition of neotropical dry forests. In *Seasonally dry tropical forests*, S. H. Bullock, H. A. Mooney y E. Medina (eds.) Cambridge University Press, Cambridge. p. 146–190.
- [31] Pineda-García, F., L. Arredondo-Amezcuca y G. Ibarra-Manríquez. 2007. Richness and diversity of wood species in deciduous tropical forest of El Tarimo, basin of del Balsas, Guerrero/ Riqueza y diversidad de especies leñosas del bosque tropical caducifolio El Tarimo, Cuenca del Balsas, Guerrero/ Richness and diversity of Woody species in the tropical dry forests of El Tarimo, Cuenca del Balsas, Guerrero. *Revista Mexicana de Biodiversidad* 78: 129–139.
- [32] Durán, Z.V.H., M.J.R. Francia, P.C. R. Rodríguez, R.A. Martínez y R.B. Cárceles. 2006. Soil-erosion and runoff prevention by plant covers in a mountainous area (SE Spain): Implications for sustainable agriculture. *The Environmentalist* 26:309–31.
- [33] Trejo, I. 2005. Analysis of the diversity of the deciduous tropical forest in Mexico. In: Halfter et al., (2005). *On biological diversity: the meaning of alpha, beta and gamma diversities/ Análisis de la diversidad de la selva baja caducifolia en México*. En: Halfter et al., (2005). *Sobre diversidad biológica: el significado de las diversidades alfa, beta y gama*. CONABIO, SEA, CONACyT. ISBN 84–932807–7-1. Pp. 111–122.