

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,600

Open access books available

137,000

International authors and editors

170M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Chapter

The Regenerative Effect of Intra-Articular Injection of Autologous Fat Micro-Graft in Treatment of Chronic Knee Osteoarthritis

*Mohammed Mesfer Al Kahtani, Ali H. Al Yami,
Sarah Saleh Al Qahtani and Sihem Aouabdi*

Abstract

Osteoarthritis (OA) is one of the most prevalent conditions resulting to disability particularly in elderly population. About 13% of women and 10% of men aged 60 years and older have symptomatic knee OA. The proportions of people affected with symptomatic knee OA is likely to increase due to the aging of the population and the rate of obesity or overweight in the general population. There are multiple factors associated with this progressive disease such as obesity, female gender, and repetitive trauma. Pain is the most common symptom in knee OA, a leading cause of chronic disability, clinical diagnosis will be supported by certain radiological findings. There are numerous conservative therapies that help to relieve symptoms depend on severity of Osteoarthritis, and knee replacement remains standard of care in advance disease. Fat Micrografting is evolving technique with promising result in selected patients with regenerative and reparative effect of adipocyte-derived stem cell toward damaged cartilage and bone, which supported by clinical evidence.

Keywords: autologous fat micrograft, knee osteoarthritis, intra-articular injection, cartilage degeneration, adipocyte stem cells

1. Introduction

Rheumatic and musculoskeletal diseases (RMDs) constitute a group of more than 150 Conditions that are commonly characterized by progressive lesions and painful symptomatology. Altogether, they account for the leading cause of morbidity and disability worldwide, giving rise to tremendous health expenditures and professional incapacity. Osteoarthritis (OA), one of the RMDs, is a degenerative condition that principally involves the joint's cartilage, leading to its progressive destruction. It is related to aging and lifelong continual stress on the most functional articulations such as the knees, hips, and fingers, and the lower spine region. OA ranks among the ten most disabling conditions in developed countries. The global prevalence of symptomatic OA is estimated as 9.6% in males and 18.0% in

females over 60 years of age. Further, 80% of individuals with OA would experience significant movement limitations, and 25% would have serious handicap to perform routine activities of the daily life [1].

The prevalence of OA varies in different regions of the world, with rates ranging from 3.8–70%, depending on the methodology of studies, whether clinical, radiographic, patient self-reporting, or physician diagnosis [2]. As the incidence and prevalence of OA increase with age, the extending life expectancy results in a growing number of people afflicted with OA, with a proportional risk of disability. In the United Kingdom, 20–30% of the elderly population (aged 60 years and above) are diagnosed with symptomatic OA [3]. In the Middle East countries, including Iraq, Yemen, Saudi Arabia, and Syria, more than one million people are estimated to have OA [4]. Approximately 85% of individuals over the age of 75 experience some symptoms of OA [5].

Knee pain represents more than 80% of the total burden of OA [6]. High body mass Index (BMI) has become an epidemic in the US in recent decades and is a well-known risk factor for knee OA [7]. In Saudi Arabia, a clinically based epidemiological study, at a primary healthcare clinic, by Al-Shammari et al. showed a prevalence of OA as high as 57.2%. Other data by Al Arfaj estimated the prevalence of knee OA as 53.3% and 60.9% in males and females respectively [8, 9].

The treatment of knee OA may use conservative measures including medications, Physiotherapy, and local injections, or surgical approach including total knee arthroplasty (TKA). TKA is highly effective in reducing articular pain and is associated with acceptable functional outcomes. The procedure of TKA is safe and is considered one of the most common and successful procedures in orthopedics.

2. Risk factors

Blagojevic et al. did a systemic review and meta-analysis to study the risk factors of Knee OA in older patients. They included 85 out of 2233 studies screened. The main risk factors found were obesity, previous trauma, hand OA, female gender and older age [10].

High body mass index (BMI) is associated with development of knee OA, and it was proven that the physical disability of the patients affected by knee OA reduced after weight reduction [11].

Another meta analysis conducted by Muthuri et al. showed that knee injury's history is one of the major risk factors associated with Knee OA and it should be included in any prevention program since it is preventable factor [12]. Genetic factor as shown in different studies is also associated with OA [13]. Increased loading or mal alignment of the joints is considered to be other factors that may lead to OA. So, in summary the risk factors can be classified into patient related factors like BMI, genetic, gender and age and joint related like previous injury, abnormal loading or malalignment [14].

3. Staging

In research, multiple variations of the Kellgren and Lawrence staging system have been used. However, the original one is: [15].

- grade 0 (none): definite absence of x-ray changes of osteoarthritis.
- grade 1 (doubtful): doubtful joint space narrowing and possible osteophytic lipping.

- grade 2 (minimal): definite osteophytes and possible joint space narrowing.
- grade 3 (moderate): moderate multiple osteophytes, definite narrowing of joint space and some sclerosis and possible deformity of bone ends.
- grade 4 (severe): large osteophytes, marked narrowing of joint space, severe sclerosis and definite deformity of bone ends.

4. Management

General and clinical assessment of the patient will help in determining the appropriate. Treatment of osteoarthritis. The assessment of OA effect on patient's function, daily activity, social relationship and quality of life should precede any treatment. The. Management plan has to be discussed with the patient thoroughly including the education of osteoarthritis, benefits and risks of various treatment options.

4.1 Non-surgical management

There are different non-surgical methods to treat joint OA. According to the Osteoarthritis Research society International (OARSI) guidelines published in 2019, patient education and land-based exercise with or without management of dietary weight are core treatments of knee related osteoarthritis. Non-steroid anti-inflammatory drugs (NSAIDs) are recommended in OA.

Management and its topical derivative are strongly recommended for patients with knee OA. COX-2 inhibitors or NSAIDs with proton pump inhibitor should be utilized in patients with gastrointestinal pathology. Oral NSAIDs are not recommended for patients with cardiovascular diseases. Intraarticular injection with corticosteroids or hyaluronic acid is mainly used for knee OA. It is not recommended for poly articular or even hip OA. All these non-surgical methods are associated with different level of evidence in the literature [16].

4.2 Surgical management

Before surgical intervention is considered, the patient should have received the core conservative treatment. Arthroscopic debridement and lavage are not routinely recommended unless there is a clear justification such as mechanical block that can be resolved arthroscopically [17]. After failure of all non-surgical treatments and in advanced joint arthritis, the total joint arthroplasty using artificial joint is the best option. It should be done before the patient gets advanced functional limitations. It is very safe procedure and associated with sound outcome. Pre- and post-operative patient engagement in terms of having proper education and well-structured physiotherapy is crucial to end up with great results. The American Academy of Orthopedic Surgeons (AAOS) in their 2nd edition evidence-based guideline for the treatment of knee OA has published 15 recommendations. This includes self-management exercise programs, weight reduction for painful knee and BMI < 25 and NSAIDs (oral or topical) or tramadol for symptomatic patients. They could not recommend the use of glucosamine and chondroitin, Intra articular Hyaluronic acid injection and arthroscopic lavage and or debridement for patient diagnosed with knee OA. They were unable to recommend for or against the using of knee corticosteroid, growth factor, platelet rich plasma (PRP) intraarticular injections and arthroscopic partial meniscectomy for a torn meniscus in patients with

symptomatic OA. They also have stated that high tibial valgus osteotomy might be performed for patients with painful medial knee joint OA [18].

5. Fat graft evolution

Neuber presented history of fat transplantation in literature initially in 1893 and he stated that smaller fat parcels tend to undergo less absorption [19], followed by communication from Czerny [20] Lexer [21], and Rehn [22]. In 1911, Bruning was the first to transfer autologous fat into the subcutaneous tissue for the purpose of soft-tissue augmentation [23]. 1950 Peer published 1st book about fat grafting in 300 pages, mentioned that survival rate of fat graft could be 50% and determined the viability of fat tissue which injected [24], in 1980 s liposuction technique evolved which improved technically over time, 1985 Illouz [25] and Fournier [26] developed an comprehensive approach to fat transfer by syringe harvesting, called “microlipoinjection”. Fat harvest from liposuction became the simplest and easiest method to pursue fat grafting, which indicated to treat soft tissue depression and contour deformity. 1990 Coleman described steps of fat injection procedure and coins the term Lipostructure [27].

There is numerous indications for fat graft in esthetic and regenerative medicine, Fat grafting technique evolved over period of time and become standard of care use in esthetic and reconstructive cases, in era of regenerative medicine, Lipoaspirate is consider source of fat micrograft which contains adipocyte derived stem cell (ADCS), growth factors, preadipocytes and cytokines demonstrate promising clinical application in wide spectrum of pathology to regenerate and reform damaged biological structure and improve outcome as in Osteoarthritis which consider leading cause to disability particularly in elderly population [28].

6. The use of microfat graft for cartilage repair

Cartilage contain a small number of cells known as chondrocytes, which are responsible for maintaining a large extracellular matrix, 85% of cartilage constitutes water and two categories of molecules: collagenous and noncollagenous [29], The main function of cartilage is to protect underlying bone from friction and act as gliding surface to enable motion, Articular cartilage is characterized as avascular, aneural, and alymphatic and, at maturity, of low metabolic activity [30], which entitle cartilage to special tissue with difficult task to repair it self, It has been demonstrated that early in the process of cartilage damage there is a rapid loss of glycosaminoglycans from the tissue [31], Thus, most large defects fail to heal, leading to a long-term prognosis of osteoarthritis [32].

7. Surgical technique to harvest fat

Fat is readily available and simple to harvest, with the fat grafting surgery itself shows a low donor-site morbidity, and is inexpensive and repeatable. Liposuction is considered one of the most frequently performed surgical procedures all over the world, Since its introduction in 1982 using a blunt cannula attached to a suction generating device, the procedure has been improved [33], Current technology for liposuction includes suction-assisted lipectomy, ultrasound-assisted, power-assisted, laser-assisted, and radiofrequency-assisted liposuction [34].

7.1 Harvesting of the adipose tissue

The first step in fat transfer to harvest fat by Many different liposuction techniques, all with the aim of minimizing adipocyte damage and increasing its survival Chosen Donor site is pretreated with a tumescent solution which containing an anesthetic mixture of Lidocaine, Sodium bicarbonate to overcome the acidity of mixture and to reduce pain and discomfort at injection site,epinephrine to control bleeding by vasoconstrictive action and Normal saline, proportionally mixed according to surgeon preference, volume to be injected determine by the desire volume of fat harvest and accordingly 2–3 cc of tumescent mixture to each 1 cc of anticipated fat harvest.

There are a variety of suction methods from which one may choose. The standard techniques are the manual (Coleman technique) or suction-assisted liposuction (SAL). Negative pressure is applied With the Coleman technique and SAL in combination with gentle forward and backward movement of the cannula, causing physical disruption, thus allowing fat tissue harvest. These methods represent the current gold standard, and reports show a lack of stem cell damage and preservation of their regenerative potential [35].

There are multiple modifications of this method, including power-assisted liposuction (PAL), water jet- assisted liposuction (WAL), laser-assisted liposuction (LAL), ultrasound-assisted liposuction (UAL), and VASER (Vibration Amplification of Sound Energy at Resonance).

They all are developed to further facilitate the process of suctioning, with minimal trauma to the donor site and maximal outcome in the requested esthetic result.

Large-bore cannulas decrease the mechanical sheer stress on the harvested cells, and subsequently increase the total number of viable aspirated cells, and Studies have showed an inverse relationship between cellular damage and the diameter of the instrument used to extract fat [36].

8. Processing of lipoaspirate

The goal of processing is to eliminate cellular debris, a cellular oil and excess of infiltrated solution [37]. These elements cause inflammation at the recipient site, which can be unfavorable for the fat graft. Also, blood must be removed as it accelerates the degradation of the transplanted fat.

Sedimentation: little traumatic and gives a large number of vital and intact adipocytes. However, this method contains smaller concentrations of stem cells and a substantial amount of cellular debris and thus making it harmful to graft survival [38].

Filtration techniques: more efficient in producing viable graft material for large-volume fat transfers. One example is Puregraft filtration system; which is a closed-membrane filtration system that was originally designed to prepare fat for isolation of the stromal vascular fraction. Another example of filtration is lipoaspirate filtered with cotton gauze; this results in concentrating the fat and separating it from the infiltrated solution, oil and cellular debris. This method, as compared to centrifugation, showed no significant differences in the viability of transplanted fat cells.

Washing with normal saline: this preserves mesenchymal stem cells as well as a great number of adipocytes.

Centrifugation: considered the most frequently used technique. It separates fat from substances that increase the degradation such as blood, proteases, lipids, lipases, and it may concentrate the adipose stem cell fraction, potentially enhancing

graft survival, Coleman suggested a processing method where centrifugation speed is 3000 rpm for 3 min. This creates multiple layers; the upper level is composed primarily of oil, the middle portion is fatty tissue, and the lowest portion, which is the densest layer, is composed of fluids and blood. This method obtains the highest possible concentration of stem cells within aspirate. It has also the increased content of angiogenic growth factors such as fibroblast growth factor (FGF) and vascular endothelial growth factor (VEGF). To note that excessive centrifugal force may damage intact adipocytes and on the other hand better graft viability with low centrifugal forces.

9. Microfat graft preparation

Processing micro fat graft that require to process harvested fat graft through multiple micro pores filters that allow micro fat graft to pass through, these filters have different sizes that connected to Leur-leur 1–10 cc syringe.

10. Fat graft injection

Adipose tissue is injected in to the transplantation site through cannulas, where multiple tunnels are created on insertion, and fat is injected during withdrawal of the cannula. Graft through nutrition by tissue fluid absorption can survive up to 48 h. Meanwhile, neovascularization is being established. Therefore, the diameter of the graft should not be more than 2 mm to avoid central necrosis. The osteoarthritic knee joint was injected with autologous intra-articular fat micrograft 15–20 mL through the lateral approach according to the case in an amount that did not produce high pressure inside the joint and did not produce pain to the patients due to tension of the joint capsule [39].

11. Postoperative care

Postoperative care include antibiotics for one week, pain killers, and garment pressure dressing at injection and donor sites, encourage exercise and physiotherapy, massage to reduce swelling.

12. Complications

Infection is the most devastating encounter complication which presented as redness, hotness, increase pain, and purulent collection, treatment depend on severity of infection ranging from I.V antibiotics in addition to incision and drainage.

13. Fat is source of adipocyte derived stem cells (ADSC)

Adipose tissue composed of mature adipocytes (>90%) and a stromal vascular fraction (SVF), which includes preadipocytes, fibroblasts, vascular smooth muscle cells, endothelial cells, resident monocytes/macrophages, lymphocytes, ADSC, cytokines and growth factors [40–46], led to a growing interest for the use fat graft as regenerative therapy for common bone and joint diseases, with promising

therapeutic clinical application of ADSC into skeletal system with underlying structure such as Muscle, cartilage, ligament, tendon, and bone with regenerative and reparative potential and ADCs are considered as an ideal source of cell therapy for different types of diseases including bone and joint diseases [47].

14. Adipocyte derived stem cell role in regenerate knee osteoarthritis

Adult stem cells, represented mainly by the mesenchymal stem cells (MSCs) are present in most organs and tissues of the human body; they intend to replace damaged cells as a normal process [48]. These stem cells do not have ethical or legal concerns as compared to the embryonic and fetal stem cells [49]. Mesenchymal stem cells are a promising tool for tissue regeneration. They are multipotent adult stem cells obtained from different sources like bone marrow (BM), adipose tissue, umbilical cord, placenta, synovial membrane [50]. These cells because they are located in fat and synovial membrane, they are most suitable for the treatment of osteoarthritis (OA) [51].

The International Society of Cellular Therapy (ISCT) defines mSCs by three characteristics [52]. They need to be plastic adherent, express specific markers like the CD105, CD90, and negative for CD34, CD45, HLA-DR [47, 53, 54]. MSCs can differentiate to mesodermal lineage cells (osteocytes, adipocytes, chondrocytes) [54]. These cells differentiate also into many other cell types, like myocytes, neurons [55], cardiomyocytes and hepatocytes [56] *in vitro* and *in vivo* [57–59].

They are non-immunogenic cells as they lack the expression HLA-DR receptor, which makes them suitable for allogeneic transplantation [59]. These cells are capable of suppressing lymphocyte reactivity [60] and inhibit the production of inflammatory cytokines *in vitro* [61]. MSCs express cytokine and chemokine receptors on their cell surface, which allows them to migrate to the site of injury [62]. The ability to suppress the immune response enabled their use in graft-versus-host disease and transplant rejection [63].

Each of the MSCs depending on their origins presents some differences. The BM-MSCs have high differentiation capability, but are difficult to get from bone marrow. The adipose MSCs are easily obtained from adipose tissue with high yield and strong suppressive capabilities [64]. The umbilical cord MSCs are easy to get after birth, they have high self-renewable and differentiation capacities. The synovial-MSCs have high proliferative and differentiation capacities and very low immunogenicity [64].

MSCs from BM from mouse and from human were the first to be identified and are the most studied [54, 58]. Adipose stem cells (ASCs) were first identified as stem cells in 2001, capable to differentiate into cartilage, bone and adipose cells [65].

The age of the donor reflects on the differentiation potential of the cells [66]. Umbilical cord MSCs (UC-MSCs) showed to be highly proliferative and with differentiation potentials [67]. The drawbacks of the BM-MSCs are that the procedure to obtain the cells is painful, costly and does not yield high number of cells for cell therapy [49, 68, 69]. Besides, the procedure for BM isolation can result in potential infections [54].

Synovial-MSCs (S-MSCs) isolated and characterized first in 2001, are also promising tools for the treatment OA due to their natural homing in this site [70]. They were shown to have high chondrogenic differentiation capacities, high expression of type II collagen, compared to other sources of MSCs [71, 72]. S-MSCs are isolated in low numbers from different sites, like styloid fossa and paralabral synovium [73, 74]. They have low immunogenicity with high proliferation potentials [75, 76].

The adipose MSCs (A-MSCs) represent many advantages compared to other sources of the MSCs. They are abundant in the adipose tissue, adipose tissue is easier to get compared to BM, they have strong immunosuppressive capacities [47].

More research is needed to unveil the differences in the traits of A-MSCs isolated from different sites and their implications in therapy. Previously, it was shown that A-MSCs could differentiate from one site to another in term of their cell markers. The A-MSCs isolated from abdominal adipose tissue had high expression in CD31, CD45 and HLA-DR compared to the cells isolated from orbital adipose and had lower expression of CD73, CD90, CD105 and CD146 [77]. MSCs from adipose tissue represent a good and a promising alternative to bone marrow MSCs. They share the same phenotype as MSCs from bone marrow apart from few. They differentiate similarly to BM-MSCs into the three lineages, chondrocytes, adipocytes and osteocytes. Most importantly, they are easily obtained from liposuction aspirate of patients undertaking plastic surgery [65]. Adipose tissue contains greater number of stem cells than bone marrow (10 times higher) [59]. A large number of cells are required for transplantation. Stem cells isolated from adipose tissue or another source like bone marrow, need to be scaled up. At least 2×10^6 of MSCs is required per kilogram body weight [78].

MSCs from different sites (BM, adipose) tissue are heterogeneous cell population [79]. Although MSCs from bone marrow and from adipose tissue have similar differentiation potential, there are minor differences. ASCs showed higher chondrogenic potential than MSCs from bone marrow [80]. Other study however, showed no significant differences between these two populations for chondrogenic differentiation in 2D culture but the BM-MSCs showed higher chondrogenic differentiation in 3D culture [81]. Proliferative capacities and osteogenic differentiation of the MSCs from bone marrow are reduced with age [82]. According to previous report, MSCs can be affected differently depending on the disease. For example, it was reported that the BM-MSCs from osteoporotic patients had reduced osteogenic activity [82], whereas the BM-MSCs from OA patients did not show any differences with the normal patient [83]. In another, the chondrogenic and adipogenic differentiation of the MSCs were reduced in OA patients but not the osteogenic differentiation [84].

The expression of CD36 differed between donors from no expression to highly expressed [85]. However, all ASC extracted from different donors showed the expression of the CD90, CD73 and CD105. ASCs markers can change in expression depending on the age of cells in culture. CD106 is expressed in MSCs from bone marrow but not in ASCs where this latter expresses CD49b [52]. ASCs and BM-MSCs express CD29 (beta-1 integrin, important factor in angiogenesis) [74] and CD44 (hyaluronate receptor, important for neoextracellular matrix) and CD49e (alpha-5 integrin, important for cell adhesion to fibronectin) [59]. ASCs and BM-MSCs have been shown to secrete angiogenic growth factors, like VEGF, P1GF, bFGF, angiogenin, GM-CSF, MCP-1 and SDF-1alpha (Rehman, 2004; Kinnaird et al.; 2004), these could be involved in increased angiogenesis in ischaemic tissue [59]. CD117 (stem cell factor) a marker for totipotency and pluripotency, was expressed in ASCs and BM-MSCs [85]. Based on the International Society for Cellular Therapy, the minimum criteria to define mesenchymal stem cells are CD105 and CD90, their potential to differentiate to adipocytes, chondrocytes and osteocytes and they are plastic adherent [86]. The expression of the CD34 is only reported on the adipose derived mesenchymal stem cells but the expression decreases in culture [87, 88] and the expression of CD105 increases [89] (Braun et al.; 2013). CD34 is expressed on ASCs before they are isolated from the stromal vascular fraction (SVF) [89] (Braun et al., 2013). The International Society for Cellular Therapy (ISCT) and the International Federation for Adipose Therapeutics

and Science (IFATS) agreed on these changes in phenotype of ASCs in SVF (uncultured) or when cultured [90]. The SVF is referred to the cellular pellet containing ASCs and endothelial progenitor cells without adipocytes and immune cells, [89]. When the SVF is plated, the ASCs adhere to the surface and the rest of the cell population including non-adherent and non-proliferating cells, are removed [89]. ASCs could be isolated from SVF by culture or by magnetic-activated cell sorting (MACS) [89]. The advantage of the MACS isolation is cells can be prepared in few hours as required in clinical application. When cells need to be expanded, isolating the ASCs with culture is the best method. ASCs and SVF could be used for transplant but the number of cells obtained from patients could limit the use of SVF. ASCs could be expanded to a large number of cells, which could be used for autologous transplant or banked for allogeneic transplant [89]. These differences could be due to site differences or due to the lack of standardized isolation method [91].

14.1 Clinical trials with ASCs

MSCs showed to have a big potential in clinical applications. The first clinical trials run on the application of bone marrow MSCs in patients with Hurler syndrome and metachromatic leukodystrophy (MLD) after allogeneic hematopoietic stem cells transplant, showed no toxicity secondary to the bone marrow MSCs transplant and the recovery of some of the symptoms were suggested to be due to the transplanted MSCs [92]. Since then, hundreds of MSCs clinical trials were run on many conditions including neurological diseases, cardiovascular, autoimmune and bone and cartilage diseases [93]. So far, there have been few approved clinical application for the MSCs in different countries. Few examples, are the application of allogeneic MSCs for the treatment of graft versus host disease (GVHD) in Japan, autologous bone marrow MSCs for amyotrophic lateral sclerosis and autologous adipose MSCs for Chron's fistula, human umbilical cord blood-derived MSCs for osteoarthritis in South Korea, and in Europe the application of allogeneic adipose MSCs for the treatment of fistulas in Crohn's disease [70].

There are few procedures for the treatment of cartilage injuries; the arthroplasty, microfat grafting and the autologous transplant of chondrocytes but a curative therapy still need to be developed [94, 95].

The autologous chondrocytes implantation is a surgical procedure that involves the isolation of autologous chondrocytes, expanding them in vitro then transplanting them back to the patient [96]. The application of MSCs for knee repair showed to be more effective with fewer side effects from the surgical procedure of the isolation of chondrocytes [96]. There are many clinical trials on the applications of the MSCs for OA [70].

OA is a very common condition in adults, which affects articular cartilage, subchondral bone, synovial tissue and meniscus of the joint. This leads to cartilage degeneration, osteophytes formation, subchondral sclerosis and synovial hyperplasia [97]. OA may be caused by joint injuries, obesity, aging and could be inherited condition [97].

Previous applications of MSCs for cartilage repair showed good outcomes. The BM-MSCs transplant in patients for defects in their knee cartilage and in athletes with the defect in femoral cartilage showed a good recovery of their functions [98].

MSC isolated from bone marrow, adipose tissue, umbilical cord, synovial membrane were previously used for the treatment of OA [64].

Since the discovery of ASCs, they have been used in many clinical trials for different diseases [90]. Success has been reported from different clinical trials using ASCs, but the mechanism of action is still not clear on whether cells would differentiate into the tissue or modulate the immune system [89]. Direct application of the

human ASCs improved cardiac function when injected in animals with myocardial infarction [99]. This effect was believed to be due to trophic factors released by the stem cells and differentiation of the ASCs. Local administration of ASCs accelerated the wound healing in normal and diabetic animals through differentiation into epithelial and endothelial lineage and neovascularization [100]. ASCs were applied to an injured skin due to radiotherapy, showed good healing process compared to the control [101]. ASCs although they have been used successfully in clinical trials, but their effects are not always achieved. There are many reasons that could impact the success of the use of ACS from one trial to another like cell separation, delivery methods, cell homing, engraftment and their survival [89]. Other factors might have an impact on the success of the use of ASCs like the type of liposuction procedure, site of liposuction, age of the patient and the body mass index (BMI). These factors are being investigated. Extraction of fat with different techniques might have an impact on cell viability of stem cells, their proliferation and the phenotype due to the trauma generated during the procedure. It is important to examine the different liposuction techniques and their impact on the ASCs phenotype, proliferation, and stress level for downstream applications either in research or in clinical applications.

15. Conclusion

Knee osteoarthritis is most prevalence musculoskeletal disease which cause functional limitations and affect a person's quality of life, there are known factors related to underling OA pathology such as Obesity, female gender and repetitive trauma to knee, Radiological finding such as subchondral cyst and narrowing of joint space, will support clinical diagnosis and staging of the disease using Kellgren and Lawrence staging system. Which will dictate the treatment modalities, in early stages conservative treatment such as modification of life style, rehabilitation, NSAID. Where advance stage surgical interventions ranging from arthroscopic debridement and lavage in case of blockage to total knee arthroplasty, which is being standard of care in severe condition.

Evolution of fat graft application in wide spectrum of clinical applications and have shown promising outcome to alleviate OA symptoms, where its contain adipocyte stem cells which known with reparative and regenerative process with presence of signal cofactors such as platelets derived growth factors and other growth factors to repaired damaged cartilage, fat graft being harvested from donor site mostly from abdomen, then;lipoaspirate will undergo further processing to isolate pure fat out from oil and fluid layer, then fat graft filtered with connector to reach pure micro fat graft ready to inject into affected knee under sterile process with local anesthesia.

Numerous clinical trials conducted to examine efficacy of microfat graft and ASC injection into knee with OA which demonstrated improvement in the overall condition, and further research with larger samples being in the process of publication to support clinical application and examine safety of patients.

IntechOpen

Author details

Mohammed Mesfer Al Kahtani^{1*}, Ali H. Al Yami¹, Sarah Saleh Al Qahtani²
and Sihem Aouabdi^{1,3}

1 College of Medicine, King Saud Bin Abdulaziz University for Health Sciences,
King Abdulaziz Medical City, Jeddah, Saudi Arabia

2 Department of Plastic Surgery, Université de Picardie Jules Verne, Amiens, France

3 King Abdullah International Medical Research Center, Jeddah, Saudia Arabia

*Address all correspondence to: drhababi@gmail.com

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] <https://www.who.int/chp/topics/rheumatic/en/>
- [2] Sowers M, Lanchance L, Hochberg M et al. (2000): Radiographically define osteoarthritis of the hand and knee in a young and middle aged African American and Caucasian women. *Osteoarthritis Cartilage*, 8: 69-77.
- [3] Heikkinen E (2006): Disability and physical activity in late life—research models and approaches. *Eur. Rev. Aging. Phys. Act.*, 3: 3.
- [4] Halter J, Ouslander J, Tinetti M et al. (2009): *Hazzard Geriatric Medicine and Gerontology*. 6th ed. New York: The McGraw-Hill companies.
- [5] Stevermer C (2005): Functional movement assessment for individuals with knee osteoarthritis. Published thesis, DSN, Iowa: Iowa State University, Faculty of the Graduate,. Available at <http://www.proquest.com>.
- [6] Vos T, et al. (2012) Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990-2010: A systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380:2163-2196.
- [7] Felson DT, et al. (2000) Osteoarthritis: New insights. Part 1: The disease and its risk factors. *Ann Intern Med* 133:635-646.
- [8] Alrushud AS, Rushton AB, Kanavaki AM, Greig CA. Effect of physical activity and dietary restriction interventions on weight loss and the musculoskeletal function of overweight and obese older adults with knee osteoarthritis: a systematic review and mixed method data synthesis. *BMJ Open*. 2017; 7(6): e014537. <https://doi.org/10.1136/bmjopen-2016-014537> PMID: 28600365.
- [9] Homoud AH. Knowledge, attitude, and practice of primary health care physicians in the management of osteoarthritis in Al-Jouf province, Saudi Arabia. *Niger Med J*. 2012; 53(4):213±9. <https://doi.org/10.4103/0300-1652.107556> PMID: 23661881.
- [10] Blagojevic, M., Jinks, C., Jeffery, A., & Jordan, 1. (2010). Risk factors for onset of osteoarthritis of the knee in older adults: a systematic review and meta-analysis. *Osteoarthritis and cartilage*, 18(1), 24-33.
- [11] Christensen, R., Bartels, E. M., Astrup, A., & Bliddal, H. (2007). Effect of weight reduction in obese patients diagnosed with knee osteoarthritis: a systematic review and meta-analysis. *Annals of the rheumatic diseases*, 66(4), 433-439.
- [12] Muthuri, S. G., McWilliams, D. F., Doherty, M., & Zhang, W. (2011). History of knee injuries and knee osteoarthritis: a meta-analysis of observational studies. *Osteoarthritis and Cartilage*, 19(11), 1286-1293.
- [13] Spector, T. D., & MacGregor, A. J. (2004). Risk factors for osteoarthritis: genetics. *Osteoarthritis and cartilage*, 12, 39-44.
- [14] Johnson, V. L., & Hunter, D. J. (2014). The epidemiology of osteoarthritis. *Best practice & research Clinical rheumatology*, 28(1), 5-
- [15] Kellgren JH, Lawrence JS. Radiological assessment of osteoarthrosis. *Ann Rheum Dis* 1957;16:494-502.
- [16] Bannuru, R. R., Osani, M. C., Vaysbrot, E. E., Arden, N. K., Bennell, K., Bierma-Zeinstra, S. M. A., ... & McAlindon, T. E. (2019). OARSI guidelines for the non-surgical management of knee, hip, and

- polyarticular osteoarthritis. *Osteoarthritis and cartilage*, 27(11), 1578-1589.
- [17] Conaghan, P. G., Dickson, J., & Grant, R. L. (2008). Care and management of osteoarthritis in adults: summary of NICE guidance. *Bmj*, 336(7642), 502-503.
- [18] Jevsevar, D. S. (2013). Treatment of osteoarthritis of the knee: evidence-based guideline. *JAAOS-Journal of the American Academy of Orthopaedic Surgeons*, 21(9), 571-576.
- [19] Neuber GA (1893) Fettransplantation. *Verh Dtsch Ges Chir* 22:66
- [20] Czerny A (1895) Plastischer Ersatz der BrustdrosedurcheinLipoma. *Chir Kongr Verhandl* 216:2
- [21] Lexer E (1910) Freirefettgewebstranplantation. *Dtsch Med Wochenschr* 36:46
- [22] Rehn E (1912) Die fettransplantation. *Arch Klin Chir* 98:1
- [23] Bruning P, Broeckaert TJ (1919) Contribution al'tude des greffesadipeuses. *Bull Acad R M.d Belg* 28:440
- [24] Peer LA (1950) Loss of weight and volume in human fat grafts. *Plast Reconstr Surg* 5:217-228
- [25] Illouz YG (1985) De l'utilisation de la graisse aspire pour combler les defects cutan.s. *Rev Chir Esthet Lang Franc* 10:13
- [26] Fournier PF (1985) Microlipo-extraction et microlipo-injection. *Rev Chir Esthet Lang Franc* 10:36-40
- [27] Coleman S.R. Structural fat grafts: the ideal filler? *Clin. Plast. Surg.* 2001;28:111
- [28] Behzad Heidari, Knee osteoarthritis prevalence, risk factors, pathogenesis and features: Part I , *Caspian J Intern Med.* 2011 Spring; 2(2): 205-212
- [29] Stoddart MJ, Grad S, Eglin D, Alini M (2009) Cells and biomaterials in cartilage tissue engineering. *Regen Med* 4:81-98
- [30] Cohen NP, Foster RJ, Mow VC (1998) Composition and dynamics of articular cartilage: structure, function, and maintaining healthy state. *J Orthop Sports Phys Ther* 28:203-215
- [31] Tiderius CJ, Olsson LE, Nyquist F, Dahlberg L (2005) Cartilage glycosaminoglycan loss in the acute phase after an anterior cruciate ligament injury: delayed gadolinium enhanced magnetic resonance imaging of cartilage and synovial fluid analysis. *Arthritis Rheum* 52:120-127
- [32] Fife RS, Brandt KD, Braunstein EM et al (1991) Relationship between arthroscopic evidence of cartilage damage and radiographic evidence of joint space narrowing in early osteoarthritis of the knee. *Arthritis Rheum* 34:377-382
- [33] Michele Zocchi M.D., Ph.D. *Aesthetic Plastic Surgery* volume 16, pages 287-298 (1992)
- [34] Sachin M Shridharani , Liposuction devices: technology update , *Med Devices (Auckl)*. 2014; 7: 241-251. Published online 2014 Jul 21. doi: 10.2147/MDER.S47322
- [35] S.R. Coleman, Structural fat grafting: more than a permanent filler, *Plast. Reconstr. Surg.* 118 (2006) 108S-120S
- [36] GL Campbell, N. Laudenslager, J. Newman, The effect of mechanical stress on Adipocyte morphology and metabolism, *Am. J. Cosmet. Surg.* 4 (1987) 89-94

- [37] K.A. Gutowski, Current applications and safety of autologous fat grafts: a report of the ASPS fat graft task force, *Plast. Reconstr. Surg.* 124 (2009) 272-280.
- [38] A. Condé-Green, N.F. Gontijo De Amorim, I. Pitanguy, Influence of decantation, washing and centrifugation on adipocyte and mesenchymal stem cell content of aspirated adipose tissue: a comparative study, *J. Plast. Reconstr. Aesthet. Surg.* 63 (2010) 1375-1381.
- [39] S. Moshref. The Regenerative Effect of Intra-Articular Injection of Autologous Fat Micro-Graft in Treatment of Chronic Knee Osteoarthritis, Intechopen
- [40] P. de Coppi, G. Bartsch Jr., M. M. Siddiqui et al., "Isolation of amniotic stem cell lines with potential for therapy," *Nature Biotechnology*, vol. 25, no. 1, pp. 100-106, 2007.
- [41] M. Barba, F. Pirozzi, N. Saulnier et al., "Lim mineralization protein 3 induces the osteogenic differentiation of human amniotic fluid stromal cells through Kruppel-like factor-4 downregulation and further bone-specific gene expression," *Journal of Biomedicine and Biotechnology*, vol. 2012, Article ID 813894, 11 pages, 2012.
- [42] Y.H. Chao, H. P. Wu, C.K. Chan, C. Tsai, C. T. Peng, and K.H. Wu, "Umbilical cord-derived mesenchymal stem cells for hematopoietic stem cell transplantation," *Journal of Biomedicine and Biotechnology*, vol. 2012, Article ID 759503, 5 pages, 2012.
- [43] U. G. Longo, M. Loppini, A. Berton, V. L. La, W. S. Khan, and V. Denaro, "Stem cells from umbilical cord and placenta for musculoskeletal tissue engineering," *Current Stem Cell Research & Therapy*, vol. 7, pp. 272-281, 2012.
- [44] S. Yang, S. Huang, C. Feng, and X. Fu, "Umbilical cord-derived mesenchymal stem cells: strategies, challenges, and potential for cutaneous regeneration," *Frontiers of Medicine*, vol. 6, no. 1, pp. 41-47, 2012.
- [45] D. peroni, I. Scambi, A. Pasini et al., "Stem molecular signature of adipose-derived stromal cells," *Experimental Cell Research*, vol. 314, no. 3, pp. 603-615, 2008.
- [46] K. Yoshimura, H. Suga, and H. Eto, "Adipose-derived stem/progenitor cells: roles in adipose tissue remodeling and potential use for soft tissue augmentation," *Regenerative Medicine*, vol. 4, no. 2, pp. 265-273, 2009.
- [47] Zuk PA, Zhu M, Mizuno H, Huang J, Futrell JW, Katz AJ, et al. Multilineage cells from human adipose tissue: Implications for cell based therapies. *Tissue Engineering*. 2001;7:211-228
- [48] Luyten, Frank P. Mesenchymal stem cells in osteoarthritis. *Current Opinion in Rheumatology*: September 2004 - Volume 16 - Issue 5 - p 599-603
- [49] Padoin AV, Braga-Silva J, Martins P, Rezende K, Rezende AR, Grechi B, Gehlen D, Machado DC. Sources of processed lipoaspirate cells: influence of donor site on cell concentration. *Plast Reconstr Surg*. 2008;122:614-8
- [50] Markarian C F; Frey G Z; Silveira M D; Chem E M; Milani A R; Ely P B; Horn A P; Nardi N B; Camassola M. Isolation of adipose-derived stem cells: a comparison among different methods. *Biotechnol Lett*. 2014; 36:69302.
- [51] Luyten, Frank P. Mesenchymal stem cells in osteoarthritis. *Current Opinion in Rheumatology*: September 2004 - Volume 16 - Issue 5 - p 599-603
- [52] Dominici M, Le Blanc K, Mueller I, Slaper-Cortenbach I, Marini F, Krause D, et al. Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society

for Cellular Therapy position statement. Cytotherapy. (2006) 8:315-7.

[53] Gimble JM, Katz AJ, Bunnell BA. Adipose-derived stem cells for regenerative medicine. *Circ Res.* 2007;100:1249-60. Gnanasegaran N, Govindasamy V, Musa S, Kasim NH. Different isolation methods alter the gene expression profiling of adipose derived stem cells. *Int J Med Sci.* 2014; 6:391-03.

[54] Pittenger M F, Mackay A M, et al. Multilineage Potential of Adult Human Mesenchymal Stem Cells. *Science* 1999;284;143-47.

[55] Ashjian PH, Elbarbary AS, Edmonds B, DeUgarte D, Zhu M, Zuk PA, Lorenz HP, Benhaim P, Hedrick MH. In vitro differentiation of human processed lipoaspirate cells into early neural progenitors. *Plast Reconstr Surg.* 2003; 111:1922-31.

[56] Seo MJ¹, Suh SY, Bae YC, Jung JS. Differentiation of human adipose stromal cells into hepatic lineage in vitro and in vivo. *Biochem Biophys Res Commun.* 2005;328:258-64

[57] Rodríguez JP, Astudillo P, Ríos S, Pino AM. Involvement of adipogenic potential of human bone marrow mesenchymal stem cells (MSCs) in osteoporosis. *Curr Stem Cell Res Ther.* 2008;3:208-18.

[58] Gnanasegaran N, Govindasamy V, Musa S, Kasim NH. Different isolation methods alter the gene expression profiling of adipose derived stem cells. *Int J Med Sci.* 2014; 6:391-03.

[59] Strem BM, Hicok KC, Zhu M, Wulur I, Alfonso Z, Schreiber RE, Fraser JK, Hedrick MH. Multipotential differentiation of adipose tissue-derived stem cells. *Keio J Med.* 2005;54:132-41.

[60] Cui L, Yin S, Liu W, Li N, Zhang W, Cao Y. Expanded adipose-derived stem

cells suppress mixed lymphocyte reaction by secretion of prostaglandin E2. *Tissue Eng.* 2007;13:1185-95.

[61] Gonzalez-Rey E, Gonzalez MA, Varela N, O'Valle F, Hernandez-Cortes P, Rico L, Büscher D, Delgado M. Human adipose-derived mesenchymal stem cells reduce inflammatory and T cell responses and induce regulatory T cells in vitro in rheumatoid arthritis. *Ann Rheum Dis.* 2010;69:241-8.

[62] Feisst V, Brooks AE, Chen CJ, Dunbar PR. Characterization of mesenchymal progenitor cell populations directly derived from human dermis. *Stem Cells Dev.* 2014; 15;23:631-42.

[63] Fang B, Song Y, Liao L, Zhang Y, Zhao RC. Favorable response to human adipose tissue-derived mesenchymal stem cells in steroid-refractory acute graft-versus-host disease. *Transplant Proc.* 2007;39:3358-62.

[64] Han Y, Li X, Zhang Y, Han Y, Chang F, Ding J. Mesenchymal stem cells for regenerative medicine. *Cells.* (2019) 8:886.

[65] Zuk PA¹, Zhu M, Ashjian P, De Ugarte DA, Huang JI, Mizuno H, Alfonso ZC, Fraser JK, Benhaim P, Hedrick MH. Human adipose tissue is a source of multipotent stem cells. *Mol Biol Cell.* 2002;13:4279-95

[66] Fan XL, Zhang Y, Li X, Fu QL. Mechanisms underlying the protective effects of mesenchymal stem cell-based therapy. *Cell Mol Life Sci.* (2020) 77:2771-94

[67] Chen JY, Mou XZ, Du XC, Xiang C. Comparative analysis of biological characteristics of adult mesenchymal stem cells with different tissue origins. *Asian Pac J Trop Med.* (2015) 8:739-46.

[68] Lu LL, Liu YJ, Yang SG, Zhao QJ, Wang X, Gong W, Han ZB, Xu ZS,

Lu YX, Liu D, Chen ZZ, Han ZC. Isolation and characterization of human umbilical cord mesenchymal stem cells with hematopoiesis-supportive function and other potentials. *Haematologica*. 2006;91:1017-26.

[69] Harasymiak-Krzyanowska I, Niedojado A, Karwat J, Kotua L, Gil-Kulik P, Sawiuk M, Kocki J. Adipose tissue-derived stem cells show considerable promise for regenerative medicine applications. *Cell Mol Biol Lett*. 2013;18:479-93.

[70] Joel Jihwan Hwang, Yeri Alice Rim, Yoojun Nam and Ji Hyeon Ju. Recent Developments in Clinical Applications of Mesenchymal Stem Cells in the Treatment of Rheumatoid Arthritis and Osteoarthritis *Front. Immunol.*, 08 March 2021

[71] Kubosch EJ, Lang G, Furst D, Kubosch D, Izadpanah K, Rolauuffs B, et al. The potential for synovium-derived stem cells in cartilage repair. *Curr Stem Cell Res Ther.* (2018) 13:174-84.

[72] Ogata Y, Mabuchi Y, Yoshida M, Suto EG, Suzuki N, Muneta T, et al. Purified human synovium mesenchymal stem cells as a good resource for cartilage regeneration. *PLoS ONE*. (2015) 10:e0129096

[73] Murata Y, Uchida S, Utsunomiya H, Hatakeyama A, Nakashima H, Chang A, et al. Synovial mesenchymal stem cells derived from the cotyloid fossa synovium have higher self-renewal and differentiation potential than those from the paralabral synovium in the hip joint. *Am J Sports Med.* (2018) 46:2942-53.

[74] Li TS, Ito H, Hayashi M, Furutani A, Matsuzaki M, Hamano K. Cellular expression of integrin-beta 1 is of critical importance for inducing therapeutic angiogenesis by cell

implantation. *Cardiovasc Res*. 2005; 65:64-72

[75] De Bari C, Dell'Accio F, Tylzanowski P, Luyten FP. Multipotent mesenchymal stem cells from adult human synovial membrane. *Arthritis Rheum.* (2001) 44:1928-42. doi: 10.1002/1529-0131(200108)44:8

[76] Li N, Gao J, Mi L, Zhang G, Zhang L, Zhang N, et al. Synovial membrane mesenchymal stem cells: past life, current situation, and application in bone and joint diseases. *Stem Cell Res Ther.* (2020) 11:381.

[77] Nepali S, Park M, Lew H, Kim O. Comparative analysis of human adipose-derived mesenchymal stem cells from orbital and abdominal fat. *Stem Cells Int.* (2018) 2018:3932615

[78] Govindasamy V, Ronald VS, Abdullah AN, Ganesan Nathan KR, Aziz ZA, Abdullah M, Zain RB, Kasim NH, Musa S, Bhonde RR. Human platelet lysate permits scale-up of dental pulp stromal cells for clinical applications. *Cytotherapy*. 2011;13:1221-33.

[79] Wang WZ, Fang XH, Williams SJ, Stephenson LL, Baynosa RC, Wong N, Khiabani KT, Zamboni WA. The effect of lipoaspirates cryopreservation on adipose-derived stem cells. *Aesthet Surg J*. 2013; 33:1046-55.

[80] De Ugarte DA, Morizono K, Elbarbary A, Alfonso Z, Zuk PA, Zhu M, Drago J, Ashjian P, Thomas B, Benhaim P, Chen I, Fraser J, Hedrick MH. Comparison of multi-lineage cells from human adipose tissue and bone marrow. *Cells Tissues Organs*. 2003;174:101-9.

[81] Winter A1, Breit S, Parsch D, Benz K, Steck E, Hauner H, Weber RM, Ewerbeck V, Richter W. Cartilage-like gene expression in differentiated human

stem cell spheroids: a comparison of bone marrow-derived and adipose tissue-derived stromal cells. *Arthritis Rheum.* 2003;48:418-29.

[82] Rodríguez JP, Astudillo P, Ríos S, Pino AM. Involvement of adipogenic potential of human bone marrow mesenchymal stem cells (MSCs) in osteoporosis. *Curr Stem Cell Res Ther.* 2008;3:208-18.

[83] J. Mary Murphy, Kenneth Dixon, Stephen Beck, Dennis Fabian, Andrew Feldman, Frank Barry. Reduced chondrogenic and adipogenic activity of mesenchymal stem cells from patients with advanced osteoarthritis. *Arthritis and Rheumatology.* Volume 46, Issue 3. 07 March 2002

[84] Baer PC, Geiger H. Adipose-derived mesenchymal stromal/stem cells: tissue localization, characterization, and heterogeneity. *Stem Cells Int.* 2012;2012: 812693.

[85] Aye MT, Hashemi S, Leclair B, Zeibdawi A, Trudel E, Halpenny M, Fuller V, Cheng G. Expression of stem cell factor and c-kit mRNA in cultured endothelial cells, monocytes and cloned human bone marrow stromal cells (CFU-RF). *Exp Hematol.* 1992;20:523-7.

[86] Dominici M, Le Blanc K, Mueller I, Slaper-Cortenbach I, Marini F, Krause D, et al. Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement. *Cytotherapy.* (2006) 8:315-7.

[87] Planat-Benard, V., Silvestre, JS., Cousin, B., André M., Nibbelink M., Tamarat R., Clergue M., Manneville C, Saillan-Barreau C., Duriez M., Tedgui A., Levy B., Pénicaud L., & Casteilla L. Plasticity of human adipose lineage cells toward endothelial cells: physiological and therapeutic

perspectives. *Circulation.* 2004;109: 656-63.

[88] Bajek A, Gurtowska N, Gackowska L, Kubiszewska I, Bodnar M, Marszaek A, Januszewski R, Michalkiewicz J, Drewa T. Does the liposuction method influence the phenotypic characteristic of human adipose-derived stem cells? *Biosci Rep.* 2015; 14;35.

[89] Feisst V, Brooks AE, Chen CJ, Dunbar PR. Characterization of mesenchymal progenitor cell populations directly derived from human dermis. *Stem Cells Dev.* 2014; 15;23:631-42.

[90] Bourin P, Bunnell BA, Casteilla L, Dominici M, Katz AJ, March KL, Redl H, Rubin JP, Yoshimura K, Gimble JM. Stromal cells from the adipose tissue-derived stromal vascular fraction and culture expanded adipose tissue-derived stromal/stem cells: a joint statement of the International Federation for Adipose Therapeutics and Science (IFATS) and the International Society for Cellular Therapy (ISCT). *Cytotherapy.* 2013; 15:641-8.

[91] Baer P. Adipose-derived mesenchymal stromal/stem cells: An update on their phenotype in vivo and in vitro. *World J. Stem Cells.* 2014; 6: 256-65.

[92] ON Koç, J Day, M Nieder, SL Gerson, HM Lazarus, W Krivit. Allogeneic mesenchymal stem cell infusion for treatment of metachromatic leukodystrophy (MLD) and Hurler syndrome (MPS-IH). *Bone Marrow Transplantation* volume 30, pages 215-222 (2002)

[93] Saeed H, Ahsan M, Saleem Z, Iqtedar M, Islam M, Danish Z, et al. Mesenchymal stem cells (MSCs) as

skeletal therapeutics - an update. *J Biomed Sci.* (2016) 23:41. doi: 10.1186/s12929-016-0254-3

[94] Zylinska B, Silmanowicz P, Sobczynska-Rak A, Jarosz L, Szponder T. Treatment of articular cartilage defects: focus on tissue engineering. *In Vivo.* (2018) 32:1289-300.

[95] Du D, Hsu P, Zhu Z, Zhang C. Current surgical options and innovation for repairing articular cartilage defects in the femoral head. *J Orthop Translat.* (2020) 21:122-8.

[96] Nejadnik H, Hui JH, Feng Choong EP, Tai BC, Lee EH. Autologous bone marrow-derived mesenchymal stem cells versus autologous chondrocyte implantation: an observational cohort study. *Am J Sports Med.* (2010) 38:1110-6

[97] Chen D, Shen J, Zhao W, Wang T, Han L, Hamilton JL, et al. Osteoarthritis: toward a comprehensive understanding of pathological mechanism. *Bone Res.* (2017) 5:16044.

[98] Kuroda R, Ishida K, Matsumoto T, Akisue T, Fujioka H, Mizuno K, et al. Treatment of a full-thickness articular cartilage defect in the femoral condyle of an athlete with autologous bone-marrow stromal cells. *Osteoarthritis Cartilage.* (2007) 15:226-31

[99] Cai L, Johnstone B H, Cook T G, Tan J, Fishbein M C, Chen PS and March K L. IFATS collection: Human adipose tissue-derived stem cells induce angiogenesis and nerve sprouting following myocardial infarction, in conjunction with potent preservation of cardiac function. *Stem Cells.* 2009;27:230-7.

[100] Nie C1, Yang D, Xu J, Si Z, Jin X, Zhang J. Locally administered adipose-derived stem cells accelerate wound healing through differentiation and

vasculogenesis. *Cell Transplant.* 2011;20:205-16.

[101] Rigotti G, Marchi A, Galiè M, Baroni G, Benati D, Krampera M, Pasini A, Sbarbati A. Clinical treatment of radiotherapy tissue damage by lipoaspirate transplant: a healing process mediated by adipose-derived adult stem cells. *Plast Reconstr Surg.* 2007;119:1409-22.