



# Last tesserae of a fading mosaic: floristic census and forest vegetation survey at Parche di Bilello (south-western Sicily, Italy), a site needing urgent protection measures

Alfonso La Rosa<sup>1</sup>, Lorenzo Gianguzzi<sup>2</sup>, Giuseppe Salluzzo<sup>3</sup>, Leonardo Scuderi<sup>4</sup>, Salvatore Pasta<sup>5</sup>

<sup>1</sup> *Cooperativa Silene, Via V. D'Ondes Reggio 8/a, I-90127, Palermo, Italy*

<sup>2</sup> *Department of Agricultural, Food and Forest Sciences, University of Palermo, Viale delle Scienze Ed. 4, I- 90128, Palermo, Italy*

<sup>3</sup> *Legambiente Circolo Crimiso, Piazza Regina Margherita 34, I-91022 Castelvetro, Italy*

<sup>4</sup> *Via Andromaca 60, I-91100, Trapani, Italy*

<sup>5</sup> *Institute of Biosciences and BioResources (IBBR), Italian National Council of Research (CNR), Unit of Palermo, Corso Calatafimi 414, I-90129, Palermo, Italy*

Corresponding author: Salvatore Pasta ([salvatore.pasta@ibbr.cnr.it](mailto:salvatore.pasta@ibbr.cnr.it))

Academic editor: Simonetta Bagella ♦ Received 24 December 2020 ♦ Accepted 12 May 2021 ♦ Published XX May 2021

## Abstract

This paper illustrates the botanic heritage of Parche di Bilello, a site located in the municipality of Castelvetro. The study area hosts several woodland fragments dominated by *Olea europaea* var. *sylvestris*, *Quercus suber* and *Quercus ilex*, respectively. According to historical data, these nuclei represent the last remnants of an open forestland which covered a much wider coastal area between Mazara del Vallo and Sciacca until the end of Middle Age. Phytosociological relevés were focused on these forest nuclei, probably the most representative of south-western Sicily, which correspond to three habitats included in the 92/43 EEC Directive (9320, 9330 and 9340, respectively) and represent the final stage of three different edaphic series. Wild olive forests probably dominated on sandy calcareous soils, holm oaks prevailed on steep calcareous and N-exposed slopes. In contrast, cork oaks mostly occurred on sandy subacid soils issuing from pedogenetic processes on palaeodunes. Moreover, field surveys allowed to list 331 vascular plant taxa. Among them, *Linaria multicaulis* subsp. *humilis*, *Orobanche balsensis* and *Serapias orientalis* subsp. *siciliensis* are new to Trapani Province. Detailed information on the current distribution and the synecology of several plant taxa of high biogeographic and conservation interest is also provided. The study site also hosts one habitat of priority interest (6220, i.e. xerophilous Mediterranean perennial grasslands and annual swards) and two species protected by international laws, i.e. the orchid *Ophrys lunulata* and the lichen *Teloschistes chrysophthalmus*. Considering the high value of its natural heritage, this territory deserves the adoption of more effective protection measures. For this reason the authors recommend its inclusion as a new Site of Community Interest within the Sicilian Natura 2000 network.

## Keywords

Endangered plant species, floristic inventories, forest vegetation, landscape history, nature conservation

## Introduction

Italy and Sicily have a long history of nature protection. They host hundreds of protected areas, sites of interest according to 92/43 EEC Directive and important plant areas (Blasi et al. 2010). The implementation and the planning of management and conservation measures within this

network is often informed by two-step gap analysis (Scott et al. 1993): the assessment of the extent, health, frequency, connectivity and distribution of the extant protected areas represents the starting point for the identification of knowledge and conservation gaps. These gaps may be filled in a second step by identifying new high priority areas for conservation (Rodriguez et al. 2004), for instance

by focusing on rare or endangered species and/or on habitat types, whose occurrence and distribution is also taken into account for further development of global, national or regional conservation strategies (Vimal et al. 2011).

In this framework, field research remains the main tool to deepen basic knowledge about the biological heritage of fragile and poorly investigated areas where endangered plants and habitats may co-occur. Moreover, the correct identification and the standardised description of rare vegetation units may play a key role in protecting locally endangered plant communities and preserving all the steps of local vegetation series. These actions improve the survival and the natural dynamics between and within the patches of natural and semi-natural habitats rare at the national (Rosati et al. 2007, 2008) or the regional scale (Viciani et al. 2016).

This paper presents the results from a research focused on the vascular flora and the forest vegetation of the locality 'Parche di Bilello'. This site is located between the localities 'Bresciana' and 'Madonna di Trapani' (South), 'Guardiola' (West), 'Atria' (East) and 'Canalotto' (North) and belongs to the municipality of Castelvetrano (Province of Trapani, south-western Sicily) (Figs. 1A–B). During the field survey campaign carried out between 2004 and 2014, also the occurrence and distribution of noteworthy lichens was recorded. Historic information was added in order to provide a more complete picture of the great biological and cultural value of this site.

## Location, geology, soil and climate

The study area (Figs. 1C and 2) is located between 65 and 90 m a.s.l. and corresponds to a system of gently sloping hills aligned from west to east. These hills are calcareous sandstone outcrops which survived in the form of semi-natural islands within a 'matrix' of cultivated lands.

The origin of toponym 'Parche di Bilello' remains unclear. On an old map printed by the Istituto Topografico Militare Italiano (ITMI 1872; Fig. 1C), the study site is indicated as 'Le Parche di Belleddu'. Le 'Parche' is almost certainly a mispronunciation of the vernacular Sicilian term 'Li parchi' (= the parks, i.e. the fenced/protected private areas). The study area is currently surrounded by an almost continuous wall made of calcareous sandstone blocks; this wall has been built either to protect its surface against burning and browsing by domestic herbivores or to create a closed hunting reserve for rabbits.

The almost flat coastal landscapes between Sciacca and Marsala are usually called Sciare, a vernacular term which seems to derive from the Arab word 'sha'raa' (= sterile and uncultivated area). Agricultural activities are difficult here due to poor soil availability. The term 'Sciare' has been broadly used to indicate local karst flatlands, characterized by a mosaic of outcropping calcareous rocks and sandy-loamy shallow lithosols supporting a mosaic-like patchwork of thermo-xerophilous vegetation including therophytic swards, perennial dry grasslands, garrigues and low maquis. The Sciare also host several small temporary ponds and rock pools rich in hygrophilous and



**Figure 1.** Location of the study area with respect to Italian (A) and Sicilian (B) territory; C) the study area and its close surroundings: particular from ITMI (1872, original scale 1:100,000).

aquatic communities. Until mid 1900s the local complex network of humid areas was fed by a rich and shallow aquifer, whose level was progressively lowered by more than 20 m during the last decades due to indiscriminate (and often illegal) water uptake (Bonanno et al. 2000).

According to Lentini and Carbone (2014), most of the Sciare are characterised by lower Pleistocene (1.8–0.8 Ma) marine sediments (the so-called ‘Calcareni di Marsala’), covering the so-called ‘Formazione Marnoso-Arenacea della Valle del Belice’, made of sandstones with interbedded marl and clay layers which filled this sector of the Sicilian foredeep during mid-upper Pliocene (3.6–1.8 Ma). According to USDA soil taxonomy (Fierotti 1988), most of the coastal plain and low hills between Marsala and Menfi hosts different combinations of lithic xerorthents (lithosols with  $\text{pH} \geq 7$ ), typical and/or lithic rhodoxeralfs (‘terra rossa’,  $\text{pH} < 7$ ).

Basing on the available thermo-pluviometric data from the stations of Castelvetrano, Mazara del Vallo and Sciac-

ca, according to Rivas-Martínez bioclimatic classification (Rivas-Martínez et al. 2004) the study area belongs to lower thermo-mediterranean thermotype and upper dry ombrottype (Bazan et al. 2015; Gianguzzi et al. 2016). Mean annual temperatures range between 17.5 and 18.0 °C, and the highest mean monthly temperatures reach 25.5–26 °C (July–August) and never go below 11 °C (January–February). More details on local climatic patterns are provided in Table 1.

### Historical evolution of local landscape

The pollen analysis from cores carried out along the south-western coasts of Sicily not far from our study site at Gorgo Basso and Lago Preola (municipality of Mazara del Vallo) allowed to trace the main phases of the evolution of local landscape over last 10,500 years (Tinner et al. 2009; Calò et al. 2012). According to these studies, local

**Table 1.** Main climatic data obtained from the nearest climatic stations (from Duro et al. 1996). A, local rainfall (mm); B, temperature (°C); AV = average values.

A														
Station	J	F	M	A	M	J	J	A	S	O	N	D	Year	Nb. rainy days
Campobello di Mazara	71.0	76.4	49.6	53.9	27.7	3.0	1.3	8.8	29.1	81.4	80.9	73.5	556.6	67
B														
Station	AV of the warmest month	AV of the coldest month	Annual AV	Range between the AV of the warmest and coldest month	Absolute max. value	Absolute min. value								
Mazara del Vallo	22.3	13.8	18.0	8.5	48.8	-0.1								



**Figure 2.** The site of ‘Parche di Bilello’ (source: Google Earth). White line = perimeter of the area where the floristic inventory was carried out; its protection as a new Site of Community Interest is here recommended.




forest ecosystems never recovered from a major collapse occurred under Roman rule, i.e. between 200 BC and 200 AD, and another reduction of tree cover eventually occurred some 500–400 years ago.

The study site falls within the wider territory of Birribaida. This toponym, still reported in the already mentioned ITMI (1872; Fig. 1C), probably derives from a settlement built in the Middle Age. In fact, a place called ‘Bellumreparum’, clearly recalling the Provençal ‘Bel-ripayre’ (= beautiful shelter), was first mentioned on some documents dating back to the XIII–XIV centuries AD (Bresc 1975, 1986; Sciascia 2002; Bresc and Sciascia 2016), and refers to a fortified residence where the Emperor Frederic II Hohenstaufen used to rest when visiting local hunting estate (Huillard-Bréholles 1860; Bresc 1980; Maurici 1995). During the following centuries, when the toponym was gradually corrupted into ‘Perribaida’ (Fazello 1560) and ‘Berribaida’ (Magini 1620), this territory was renowned for the richness of its game (large mammals like wildboars, roe and fallow deers; birds like rock partridges, quails and hazels). Birribaida was probably just a portion of an open and discontinuous forestland, occupying almost continuously the flatlands stretching from the present-day cities of Mazara del Vallo and Sciacca along the coast, the so-called ‘Basso Belice’ nowadays including

the inland territories of Menfi, Campobello di Mazara, Castelvetro and Partanna (Fig. 3).

Indeed, between 13th and 15th centuries local forest cover underwent substantial reduction and fragmentation. Several notarial documents confirm this; for instance, Bresc-Gautier (1983) reports that already in 1239 the farmers of Sciacca complained with Emperor Frederic II Hohenstaufen for the shortage of wood needed to build ploughs, and Simonsohn (2007) mentions the sale of many dozens of bags of charcoal obtained from the cork oaks of ‘Castelvetro’ in the years 1428 and 1429. In the meantime, many marshes and temporary ponds were drained and converted into cultivated lands. The Cistercian monks of Delia were responsible for the drainage of numerous swamps in the fiefs ‘Margio’ (vernacular name for marsh, temporary pond), ‘Fontanelle’ (= small springs), and ‘Marcita’ (= permanent, stagnant water body), toponyms which confirm the locally shallow water table and the high freshwater availability due to a remarkable frequency of springs (Noto 1732). The occurrence of wetlands (Fig. 3) is also confirmed by the occurrence of scattered nuclei of elms (*Ulmus* spp.), poplars (*Populus* spp.) and willows (*Salix* spp.) in the surroundings (G. Salluzzo, *pers. obs.*) by the local frequency of hackberry trees (*Celtis australis*; see Cusumano 1983; Scuderi 2006), and by other toponyms like ‘Margio rotondo’ (= rounded



**Figure 3.** The coastal area between Mazara del Vallo and Menfi during the Middle Ages (from Calamia et al. 2004, modified). The shaded surface coincides with the previous extension of the open forestland of Birribaida; lines refer to 50 and 100 m a.s.l. or to the main local watercourses. As for the Swabian castles, Bellumrepar (1), Bellumvider (2), and Burgimilluso (3), are currently located in the centre of Castelvetro, Campobello di Mazara and Menfi, respectively.

‘margio’) and ‘Gurra’ (a vernacular name referred to willow species) reported in medieval documents (Calamia et al. 2004).

During the XVI century, the noble family Aragona Tagliavia, counts (and since 1564 princes) of Castelvetro implemented a policy aiming at promoting local agricultural activities by means of mid- to long-term ground leasing contracts (emphyteusis); consequently local farmers could act as temporary land owners and gradually expanded cereal crop fields, vineyards, olive groves to the detriment of local woodland nuclei, and intensified the exploitation of local water for mills and to irrigate sugar cane plantations, rice fields and vegetable gardens (Scalisi 2012). The wide area still hosted some fragments of forest-maquis until the XVIII–XIX centuries AD (Noto 1732; Amico Statella 1759; Bivona-Bernardi 1845; Ferrigno 1931, 1933; Napoli 1932; Titone 1961, Caruso and Nobili 2001), as testified by the numerous local toponyms like ‘Bosco della Guardiola’, ‘Macchia di Lupo’, ‘Bosco nuovo’, ‘Bosco di Belice’ and ‘Bosco Tre Fontane’ (Calamia et al. 2004).

## History of botanical investigation in the surroundings

The first records concerning the vascular plants growing in the so-called ‘Sciare’ of Marsala and Mazara date back to Gussone (1827–1832, 1842–1845), Fanales (1899) and Lojaco-Pojero (1889–1909). A general and updated overview on the main botanical features of this territory was provided by Scuderi (2006) and Guarino and Pasta (2017). Moreover, during the last 50 years several papers concerning the vascular flora and the vegetation of the territories of Campobello di Mazara, Mazara del Vallo and Castelvetro have been published. Most of them were focused on the coastal habitats and wetlands (e.g., Frei 1937; Brullo et al. 1974; Brullo and Ronsisvalle 1975; Brullo and Furnari 1976, 1978; Ronsisvalle 1979; Bartolo and Brullo 1993; La Mantia and Gianguzzi 2003; Ottonello et al. 2003; La Rosa et al. 2007), on local woody communities, namely garrigues and maquis (Brullo and Marcenò 1985; Raimondo et al. 1991; Brullo et al. 1997, 2009), and on grasslands (Catanzaro 1983; Minissale 1995; Brullo et al. 2010). Additionally, many articles focused on/mentioning single species growing in the surroundings have been published during last decades (e.g., Ottonello and Catanzaro 1986; Troia 2006; Troia and Spallino 2009; Geraci et al. 2010; Gianguzzi et al. 2010, 2012; Troia and Raimondo 2010; Badalamenti et al. 2012; Gennai et al. 2015; Médail et al. 2016; Troia and Napolitano 2017; Musarella et al. 2018).

## Methods

Lichens identification was carried out by using the analytical keys proposed by Nimis (1987), while their nomenclature follows Nimis and Martellos (2017).

The taxonomic identity, the chorotypes and the life forms of the observed vascular plants follow the second edition of ‘Flora d’Italia’ (Pignatti et al. 2017–2019), whilst the nomenclature of the listed taxa mainly follows Galasso et al. (2018) and Bartolucci et al. (2018). Alien plants have been classified as ‘casual’, ‘naturalized’ or ‘invasive’ according to their degree of naturalization (Pyšek et al. 2004).

The syntaxonomy of the high rank syntaxa (i.e. alliances, orders and classes) follows the schemes proposed by Biondi et al. (2014) and Mucina et al. (2016), while the associations mainly follow the proposals of Brullo et al. (2002) and Guarino and Pasta (2017).

Forest vegetation was investigated applying the Zurich-Montpellier method (Braun-Blanquet 1964) implemented by Géhu and Rivas-Martínez (1981) and other authors (e.g., Rivas-Martínez 2005; Biondi 2011). Phytosociological relevés were focused on forest nuclei; plot sizes ranged from 80 to 200 m<sup>2</sup> according to the surface of the surveyed forest patches.

Moreover, the evaluation of the biogeographic and conservation interest of local vascular flora was supported by the most updated red-list (Raimondo et al. 2011), floristic inventory (Giardina et al. 2007) and biogeographic overviews (Brullo et al. 1995; Guarino and Pasta 2018) available at regional level.

Some dry specimens of the vascular plants and lichens collected in the study area are stored in the personal herbaria of A. La Rosa, L. Gianguzzi, while many exsiccata of L. Scuderi are deposited in the herbarium of the University of Catania (CAT).

## Results and Discussion

### Lichens

Field investigations aiming to list all the lichens occurring in the study area are still going on, and the final results will hopefully be published soon in a separate paper. During these surveys, around 50 individuals of an extremely rare lichen, *Teloschistes chrysophthalmus* (L.) Th. Fr. (Fig. 4A), were observed growing on the bark of a wild olive tree. *T. chrysophthalmus* is a fruticose lichen which prefers well lit and moderately nutrient-rich places. It usually colonizes the twigs of small trees and shrubs, and it may also grow on the wooden poles of fences (UK BAP 1999). Even if small-sized, this species is very showy due to the grey-greenish to golden colour of its thallus and the crown of orange cilia surrounding its apothecia. *T. chrysophthalmus* is a subcosmopolitan lichen: its distribution range stretches from tropical and sub-tropical to temperate countries on both hemispheres. By contrast, this species has recently experienced a dramatic shrinkage, triggered by air pollution, the indiscriminate use of non-organic fertilisers and the abandonment of extensive fruit-tree orchards (Ravera et al. 2011). As a consequence, it results to be severely endangered in several parts of cen-

tral and northern Europe such as in North Italy (Capozzi et al. 2013) and in North Ireland, where it was considered extinct (UK PAB 1999). To contrast this negative trend, special conservation measures have been adopted at the European level, such as the inclusion in the European red list (Serasiaux 1989). In the UK an ad hoc action plan aiming at its safeguard has been implemented, including protecting its preferential habitat (i.e. old fruit tree orchards and hedgerows: UK PAB 1999). Although this species is reported to occur throughout the entire Italian territory, it is very rare in many regions and subject to recent shrinking (Nimis and Martellos 2017). As for Sicily, during recent times the species was only observed in the cork oakwood referred to *Stipo bromoidis-Quercetum suberis* at Santo Pietro near Caltagirone (Capozzi et al. 2013).

### General traits of local vascular flora

As much as 331 infrageneric taxa were observed during the field surveys (see Table S1, Supplementary Material 1), belonging to 57 plant families, the most common being *Fabaceae* (46 taxa), *Asteraceae* (40), *Poaceae* (35), *Caryophyllaceae* (12), *Apiaceae* (11), *Lamiaceae* and *Orchidaceae* (10). As for the genera, the most represented were *Trifolium* (12 taxa), *Medicago* (7), *Allium*, *Convolvulus* and *Ophrys* (6), *Euphorbia* (5), *Erodium* and *Lotus* (4).

As for life-forms, the study site is characterized by a strong prevalence of annual ( $T = 170$  taxa) and perennial ( $H = 72$ ;  $G = 41$ ) herbs and grasses, representing about 85% of total vascular flora. Woody plant list includes 27 phanerophytes, whilst chamaephytes and nano-phanerophytes are both represented by 11 taxa.

The exotic (cultivated, naturalised and invasive) taxa are 13, corresponding to only 3.9% of all the observed vascular plants.

### Plants new to the Province of Trapani

*LINARIA MULTICAULIS* (L.) Mill. subsp. *HUMILIS* (Guss.) De Leon., Giardina & Zizza

This taxon (Fig. 4B) was never recorded before in the province of Trapani. It was previously reported as a narrow-ranged endemic to south-eastern Sicily (Madonna del Piano near Grammichele, Contrada Molara near Caltagirone, Contrada Dirillo near Acate: De Leonardi et al. 1999, 2003). Up to now, only the subsp. *multicaulis* was reported for the 'Sciare di Marsala' (Fanales 1899; confirmed by Scuderi 2006), Trapani and Bonagia (Ponzo 1900) and Alcamo (Ponzo 1903). More recently, subsp. *multicaulis* has been observed on the western foothills of Monte Cofano (S. Pasta, April 1998, unpubl.), while Scuderi (2006) reports it for Alcamo Marina, the 'sciare' of Campobello di Mazara and Mazara del Vallo, Castelve-trano (Cave di Cusa) and Lago Trinità as a common colonizing the sandy soils of the fossil dunes within the gaps of local shrublands. In south-eastern Sicily this species

grows in very similar conditions, dominating local therophytic swards recently framed into the alliance *Filagini asterisciflorae-Linarion humilis* Minissale & Sciandrello 2015 (order *Malcolmietalia* Rivas Goday 1958).

Local population counts few individuals observed growing in local fallow fields (79 m a.s.l., exposition East, 31.03.2013. A. La Rosa).

*OROBANCHE BALSENSIS* (J.A.Guim.) Carlón, M.Laínz, Moreno Mor. & O.Sánchez

This taxon (Fig. 4C) was never recorded before in the province of Trapani. Besides Sicily, where it has been reported to grow at Monte Catalfano (Bagheria, Palermo Province) and at Castiglione di Sicilia (Catania Province), the distribution range of this previously neglected taxon includes Spain, Portugal, southern France and Corsica (Carlón et al. 2015). Local population consists of few individuals, parasites on *Carlina sicula* subsp. *sicula* (79 m a.s.l., exposition East, 05.05.2013, A. La Rosa).

*SERAPIAS ORIENTALIS* (Greuter) H. Baumann & Künkele subsp. *SICILIENSIS* Bartolo & Pulv.

This taxon is new to Trapani province and to western Sicily (Fig. 4D). Originally described from the territory of Niscemi (Caltanissetta Province) by Bartolo and Pulvirenti (1993), so far it was considered to occur only there within open garrigue and grassland communities (Galesi 1996).

Local population counts about 20 individuals, colonizing a grassland patch occupying a gap of local shrubland, dominated by *Cytisus infestus*, referred to the association *Pyro amygdaliformis-Calicotometum infestae* (79 m a.s.l., exposition North, 28.04.2013, A. La Rosa).

### Other vascular plants of high biogeographic/conservation interest or poorly known for the Province of Trapani

*ACIS AUTUMNALIS* (L.) Sweet


The earliest records of this species (Fig. 4E) for the Province of Trapani date back to Gussone (1827), who reported it as growing at Alcamo, Trapani, Marsala and Mazara del Vallo; later on, it was observed near the Gorgi Tondi west of Mazara del Vallo by Lopriore (1900) and near the coast at Trapani (a specimen is still conserved in the herbarium of Palermo, PAL) and in locality Xiggiari near Paceco by Ponzo (1900). More recently, its occurrence has been recorded in the nature reserve of Zingaro (Raimondo and Schicchi 2000; Scuderi 2006), while Scuderi (2006) noticed it in the swards colonising the gaps of grassland and maquis communities of Montagna Grande di Salemi, as well as near Bosco di Scorace between Brucà and Buseto Palizzolo, in Contrada Fastuchera (south-west of Alcamo), and in the forest clearings near the top of Monte Inici (Castellamare del Golfo, L. Scuderi and L. Gianguzzi, unpubl.); it also occurs in the same habitats within the cork oakwood of Angimbè near Calatafimi (C.



Marcenò, October 2015, published in Pignatti et al. 2017–2019). Additional data and photos testifying the occurrence of this species in several localities of the Province of Trapani (Contrada Runza, municipality of Trapani, M. Aleo, October 2017; Gallitello, municipality of Calatafimi, G. Di Gregorio, November 2017; Marsala, G. Donadelli, September 2019; Rilievo, municipality of Trapani, M.

Aleo, October 2020) have been published by the members of the forum of 'Acta Plantarum' ([https://www.actaplantarum.org/galleria\\_flora/galleria1.php?view=1&id=2977](https://www.actaplantarum.org/galleria_flora/galleria1.php?view=1&id=2977)).

Local population counts few individuals concentrated in semi-shaded surface along a path with one of the forest nuclei referred to *Pistacio lentisci-Quercetum ilicis* (80 m a.s.l., exposition North-East, 14.10.2012, A. La Rosa).



**Figure 4.** A, *Teloschistes chrysophthalmus* (L.) Th.Fr. (photo: A. La Rosa, February 24, 2013); B, *Linaria multicaulis* (L.) Mill. subsp. *humilis* (Guss.) De Leon., Giardina & Zizza (photo: A. La Rosa, March 31, 2013); C, *Orobanche balsensis* (J.A.Guim.) Carlón, M.Laínz, Moreno Mor. & O.Sánchez (photo: A. La Rosa, May 5, 2013); D, *Serapias orientalis* (Greuter) H.Baumann & Künkele subsp. *siciliensis* Bartolo & Pulv. (photo: A. La Rosa, April 28, 2013); E, *Acis autumnalis* (L.) Sweet (photo: L. Gianguzzi, October 13, 2019); F, *Desmazeria sicula* (Jacq.) Dumort. (photo: A. La Rosa, April 20, 2013); G, *Ophrys lunulata* Parl. (photo: A. La Rosa, March 31, 2013).

*DESMAZERIA SICULA* (Jacq.) Dumort.

Within Trapani Province, this plant (Fig. 4F) grows in the annual therophytic coastal swards between M. Cofano and Bonagia (Ponzo 1900; confirmed here by Barbagallo et al. 1980), Alcamo (Ponzo 1903), Mazara del Vallo (Borzi and Franke in Nicotra and Campagna 1908), Trapani (Todaro in Nicotra and Campagna 1908). This species is featured among the characteristic taxa of the order *Stipo-Bupleuretalia semicompositi* and the alliance *Plantagini-Catapodion balearici* (Brullo 1985; Guarino and Pasta 2017). Lojacono-Pojero (1909) reports the var. *monostachya* Tineo for the locality 'Macchie di Alcamo'. Its occurrence has been reported also for Isola Lunga (Di Martino and Perrone 1970), San Vito Lo Capo (Bartolo and Brullo 1993). More recently, the species was recorded at San Teodoro in the Lagoon of Marsala and in the nature reserve of Zingaro (Scuderi 2006), at Castello di Pietra between Partanna and Castelvetrano (Scuderi, 2006; Pasta et al. 2008), on Levanzo Island (Romano et al. 2006) and near Torre Grimesi (Mazara del Vallo, L. Scuderi, unpubl.).

Local population counts few individuals growing along the borders of the road that takes to the local 'baglio' (= rural house) (80 m a.s.l., exposition North, 20.04.2013, A. La Rosa).

*OPHIGLOSSUM LUSITANICUM* L.

This small fern grows in the annual swards on nutrient-poor sandy soils (*Helianthemion guttati* Br.-Bl in Br.-Bl. & Wagner 1940) or in temporary ponds or seasonally wet rock pools referred to *Cicendion* (Rivas Goday in Rivas Goday & Borja 1961) Br.-Bl. 1967. In the province of Trapani it was first noticed by Gussone (1842–1845) who reported it at Calatafimi, Macchie di Alcamo and the coastal area between Trapani and Mazara del Vallo. It also occurs on Pantelleria Island (Sommier 1907; Médail et al. 2016), along the coast of the Stagnone and on Isola Lunga near Marsala (Brullo et al. 1994; Scuderi 2006), at Monte Cofano (Aleo and Ottonello 2000; Gianguzzi et al. 2005), Contrada San Nicola near Mazara del Vallo (Raimondo et al. 2004). Finally, Médail et al. (2016) reported five more populations, located in Contrada Ballottella (municipality of Trapani), in the temporary ponds of Contrada Anguillara near Calatafimi, in the rock pools of Pietra Colle (an isolated outcropping quartz sandstone rock between Valderice and Buseto Palizzolo) and 'Birribaida'.

More in detail, the population found at Parche di Bilello counts few individuals growing within a bryophytic community colonising a clearing within one of the main forest nuclei (80 m a.s.l., exposition East, 03.02.2013, A. La Rosa).

*OPHRYS LUNULATA* Parl.

Included in the Appendix I of Washington Convention and in the Appendix II of the 92/43 EU 'Habitats' Directive, this Sicilian endemic occurs almost everywhere in the pasturelands (garrigues, perennial grasslands, open shrubland, thermophilous forest edges) throughout the

Sicilian territory, albeit it usually forms small or very small populations. As for its distribution in the Province of Trapani (Fig. 4G), the current knowledge is probably underestimated. Based on published data, local population appears to be the first record for the southern part of Trapani Province. In fact, it was first reported for Monte Inici by Lojacono-Pojero (1909), then for several sites of the Zingaro nature reserve (Raimondo et al. 1986; Raimondo and Schicchi 2000; Scuderi 2006), near Buseto Palizzolo (Raimondo 1987) and for M. Cofano (Campo and Romano 2000; Gianguzzi et al. 2005).

Only three individuals were observed growing in a xeric grasslands along the main road (80 m a.s.l., exposition East, 31.03.2013, A. La Rosa).

## Remarks on the non-native plants observed in the study site

*Oxalis pes-caprae* dominates the winter facies of the weed communities referred to the alliance *Diploaxion eruroidis* Br.-Bl. in Br.-Bl., Gajewski, Wraber & Walas 1936 (Brullo and Marcenò 1980; Bernhardt 1986).

The local occurrence of *Galinsoga parviflora* confirms its establishment and spread during last decades. This ruderal thermophilous plant native to North America is listed among the characteristic species of the order *Eragrostietalia* J. Tx. ex Poli 1996 (Mucina et al. 2016), It was observed for the first time in the province of Trapani near Mazara del Vallo by Ottonello and Catanzaro (1986). As for *Chasmanthe floribunda*, probably most (if not all) the previous records for Sicily referred to *Chasmanthe aethiopica* (L.) N.E.Br. should be referred to this alien plant, as recently clarified by Grandis (2016), who reported its occurrence in the province on the base of photos testifying its presence in Pantelleria Island and near Rilievo (municipality of Trapani). This species grows under thermo-mediterranean climate, mostly occurring in sub-ruderal perennial grasslands typical to disturbed road verges and suburban areas and referred to *Bromo-Oryzopsis miliaceae* O. de Bolòs 1970.

## Vegetation, land-use and disturbance

This paragraph provides a brief description of the main features of the local landscape (Table 2) and of the disturbances in the territory surrounding the study area. A complete syntaxonomic scheme of the vegetation units observed in the study site and mentioned in the text is provided at the end of the paper.

The main disturbance factors recorded in the study area are wildfires, which frequently affect local vegetation, and the use of non-organic fertilisers and pesticides in the olive groves, citrus orchards and vineyards in the surroundings.

As for the semi-natural plant communities, particular attention has been paid to forest vegetation. Vegetation



**Table 2.** Soil cover units, phytosociological units and habitats (\* = priority) according to 92/43 EU Directive of the study area and the nearby territory.

Soil cover units	Association(s)	Habitats
<i>Climatophilous vegetation of semi-natural areas</i>		
Holm oakwood nuclei	<i>Pistacio lentisci-Quercetum ilicis</i>	9340
Cork oakwood nuclei	<i>Stipo bromoidis-Quercetum suberis</i> (impoverished)	9330
Dwarf palm-wild olive maquis	<i>Chamaeropo humilis-Oleetum sylvestris acanthetosum mollis</i>	9320
Rock-rose garrigues	<i>Rosmarino-Thymetum capitati</i>	-
Perennial grasslands	<i>Hyparrhenietum hirtum-pubescentis</i>	6220*
Annual swards	<i>Ononido breviflorae-Stipetum capensis, Thero-Sedetum caerulei</i>	6220*
<i>Hygrophilous vegetation occurring in the watercourses of the nearby territory</i>		
Riparian forests	<i>Salicetum albo-pedicellatae</i>	92A0
Scrub of the seasonally dry Mediterranean riverbeds	<i>Tamarix africana</i> -dominated community	92D0
Reed meadows	<i>Phragmitetum communis</i>	-

relevés (see Tables 3–5) allowed to identify three different communities, i.e. many patches of evergreen maquis dominated by *Olea europaea* var. *sylvestris* and *Chamaerops humilis* and few small remnant nuclei of evergreen forest dominated by *Quercus ilex* or *Quercus suber*.

Several patches of local evergreen sclerophyllous vegetation are dominated by wild olive trees, localised on calcareous sandstone outcrops. These nuclei shall be framed into the *Chamaeropo humilis-Oleetum sylvestris acanthetosum mollis* (Gianguzzi and Bazan 2019; Fig. 5A and Table 3), a tall and usually dense maquis dominated by *Olea europaea* var. *sylvestris* and characterized by many sclerophyllous evergreen small trees, shrubs and subshrubs characteristic of the alliance *Oleo-Ceratonion siliquae*, like *Chamaerops humilis*, *Pistacia lentiscus*, *Rhamnus alaternus*, *Teucrium fruticans*, *Asparagus albus*, *Osyris alba*, several lianas (e.g., *Asparagus acutifolius*, *Clematis cirrhosa*, *Rubia peregrina*, *Smilax aspera*) and geophytes typical to the maquis undergrowth (e.g., *Allium subhirsutum* subsp. *subhirsutum*, *Arisarum vulgare* subsp. *vulgare*). This community may represent the final stage of progressive succession on very nutrient-poor and thin lithosols under dry climatic conditions. The remnant fragments occur on South-exposed slopes on calcareous sandstones and fossil dunes along the coasts of southwestern Sicily (e.g., Sciacca, Menfi, Campobello di Mazara, Mazara del Vallo, Castelvetro) and near Palermo (northwestern Sicily) between 50 and 300 m a.s.l. (Gianguzzi and Bazan 2019), in close contact with the psammophilous dune communities (*Ammophiletea* Br.-Bl. et Tx. ex Westhoff et al. 1946) and/or with the communities colonizing the rocky shores subject to salt-spray (*Crithmo-Staticetea* Br.-Bl. in Br.-Bl. et al. 1952). Century-long agro-silvo-pastoral activities (forest clearing, overgrazing, wildfires, etc.) were responsible for the deep degradation of local soils, which in turn led to a remarkable regression of the head of series, dominated by *Olea europaea* var. *oleaster*. Consequently, pasturelands prevail in the local mosaic-like landscape, with some nuclei of low maquis dominated by *Chamaerops humilis* (*Pistacio lentisci-Chamaeropetum humilis*), intermingled with perennial xerophilous grasslands (*Hyparrhenietum hirtum-pubescentis*) and annual therophytic swards referred

to *Ononido breviflorae-Stipetum capensis* and *Thero-Sedetum caerulei*. No phytosociological relevés were carried out in these plant communities, locally represented by small-sized, discontinuous and degraded patches.

Local wild olive forest fragments, referred as habitat 9320 of the 92/43 ‘Habitats’ Directive, are among the best preserved in the entire Sicilian territory (Gianguzzi and Bazan 2019a, 2019b). The high frequency and cover rate of *Oxalis pes-caprae* within these stands are due to the high invasivity of this species, spreading from the surrounding areas characterised by large surfaces devoted to olive cultivation.

Few patches of *Quercus ilex* forest are localized on north-facing slopes; these woodland fragments (Fig. 5B and Table 4) belong to *Pistacio lentisci-Quercetum ilicis* (Brullo and Marcenò 1985; Brullo et al. 2009), an association characterised by the high cover of *Pistacia lentiscus* and evergreen and summer-deciduous small trees and shrubs characteristic of thermophilous maquis communities (order *Pistacio lentisci-Rhamnetalia alaterni*, alliance *Oleo-Ceratonion siliquae*). Another remarkable fragment of this mixed evergreen oakwood is Bosco del Cantarro near the Gorgi Tondi nature reserve, west of Mazara del Vallo. This community may represent the final stage of local progressive succession processes on the thin and base-rich lithosols found on compact limestone bedrocks. First observed in the locality Gorgi Tondi near Mazara del Vallo (Brullo and Ronsisvalle 1975), not far from our study site, this association was described by Brullo and Marcenò (1985) and his presence recorded in several localities of western Sicily, like the coastal area near Palermo (Gianguzzi et al. 1996), the Island of Marettimo (Gianguzzi et al. 2006), the southern sector of the Sicani Mts. (Gianguzzi et al. 2016), as well as the south-eastern Sicily at Bosco di Santo Pietro near Caltagirone and elsewhere in the Hyblaean Plateau and in Sardinia (Bartolo et al. 1992; Brullo et al. 2009). Mostly enjoying cooler microclimatic conditions and sites protected from the warm and dry winds coming from North Africa, under more xeric conditions this community is usually replaced by the thermo-xerophilous maquis communities referred to *Pistacio lentisci-Chamaeropetum humilis*.

**Table 3.** Dwarf Palm-Wild Olive evergreen sclerophyllous maquis (*Chamaeropo humilis-Oleatum sylvestris acanthetosum mollis*). Relevés 1–5 (L. Gianguzzi, 24.04.2005, in Gianguzzi & Bazan 2019a, Tab. S4 n. 74 and 82–85); 6–9 (L. Gianguzzi, 9.05.2007, in Gianguzzi & Bazan 2019a, Tab. S4 n. 77–78 and 88–87); 10–14 (L. Gianguzzi, 12.05.2014, unpubl.).

Relevé (n°)	1	2	3	4	5	6	7	8	9	10	11	12	13	14
Altitude (m)	80	80	86	84	80	76	70	67	65	78	79	81	81	86
Slope (°)	20	30	15	15	15	20	30	25	25	15	5	5	5	7
Aspect	SW	S	E	S	N	E	E	SE	S	E	E	E	E	N
Area (m <sup>2</sup> )	120	150	150	150	150	100	100	100	100	100	100	100	100	100
Total cover (%)	100	100	100	95	95	100	100	90	100	100	100	100	100	100
Tree cover (%)	100	100	100	90	90	100	100	20	100	100	95	100	100	40
Shrub cover (%)	100	100	100	90	90	100	100	80	100	100	95	100	100	70
Herbaceous cover (%)	40	40	40	40	40	40	40	30	20	30	25	20	20	15
Average height of the dominant layer (m)	6	6	5	3,5	4	5	7	3,5	5	6	6	5	5	7
Species per relevé	17	17	18	20	17	21	23	22	21	20	22	22	19	21
<b>Char. and diff. of association and subassociation</b>														
<i>Olea europaea</i> L. var. <i>syvestris</i> (Mill.) Lehr	5	5	4	5	4	5	5	5	5	5	5	5	4	5
<i>Acanthus mollis</i> L. subsp. <i>mollis</i>	2	3	2	1	+	3	3	1	2	1	+	1	1	1
<i>Chamaerops humilis</i> L.	+	1	1	1	1	1	1	1	1	.	.	1	1	+
<i>Rhamnus alaternus</i> L. subsp. <i>alaternus</i>	.	.	1	.	+	.	.	.	1	+	1	2	2	.
<b>Char. of alliance and order</b>														
<i>Pistacia lentiscus</i> L.	2	2	3	3	3	3	3	2	2	3	3	2	3	3
<i>Stachys major</i> (L.) Bartolucci & Peruzzi	+	+	1	1	2	1	+	1	1	.	1	+	1	1
<i>Allium subhirsutum</i> L. subsp. <i>subhirsutum</i>	1	1	1	1	.	1	1	+	.	+	+	.	1	.
<i>Asparagus albus</i> L.	1	1	+	.	1	.	.	1	1	1	1	+	+	.
<i>Clematis cirrhosa</i> L.	2	1	1	+	+	.	+	.	.	+	.	.	.	.
<i>Anagyris foetida</i> L.	.	.	1	1	.	2	.	2	.	.	.	1	2	1
<i>Cytisus infestus</i> (C.Presl) Guss. subsp. <i>infestus</i>	.	.	1	.	+	.	1	.	.	1	+	1	1	1
<i>Teucrium fruticans</i> L. subsp. <i>fruticans</i>	.	.	.	1	2	.	.	.	.	+	2	.	.	.
<i>Myrtus communis</i> L.	.	.	.	1	.	.	.	.	.	.	.	.	.	.
<i>Ceratonia siliqua</i> L.	.	.	.	.	.	.	.	.	.	.	.	1	.	.
<i>Rhamnus lycioides</i> subsp. <i>oleoides</i> (L.) Jahand. & Maire	.	.	.	.	.	.	.	.	.	.	.	.	.	1
<b>Char. of class <i>Quercetea ilicis</i></b>														
<i>Smilax aspera</i> L.	2	2	3	+	1	3	3	+	2	1	1	3	2	1
<i>Asparagus acutifolius</i> L.	1	1	1	1	+	1	1	1	1	1	1	1	.	+
<i>Rubia peregrina</i> L.	1	1	1	1	1	.	+	1	1	1	1	1	1	1
<i>Arisarum vulgare</i> O.Targ.Tozz. subsp. <i>vulgare</i>	+	1	1	1	+	1	.	+	1	+	+	.	1	+
<i>Daphne gnidium</i> L.	+	.	1	.	+	.	+	.	.	.	+	+	.	+
<i>Ampelodesmos mauritanicus</i> (Poir.) T.Durand & Schinz	.	.	+	1	.	+	1	+	.	.	.	.	+	.
<i>Rosa sempervirens</i> L.	.	.	.	.	.	+	2	+	+	+	.	.	.	.
<i>Quercus suber</i> L.	.	.	.	.	.	.	.	.	1	1	1	.	.	.
<i>Osyris alba</i> L.	.	.	.	.	.	.	.	.	.	1	.	+	.	+
<i>Celtis australis</i> L. subsp. <i>australis</i>	1	1	.	.	.	.	.	.	.	.	.	.	.	.
<i>Pyrus spinosa</i> Forssk.	.	.	.	.	.	+	1	.	.	.	.	.	.	.
<i>Phillyrea latifolia</i> L.	.	.	.	.	.	.	.	.	.	.	+	.	.	.
<b>Other species</b>														
<i>Oxalis pes-caprae</i> L.	3	2	2	3	3	2	+	1	+	2	+	+	1	.
<i>Oloptum miliaceum</i> (L.) Röser & H.R.Hamasha	+	.	.	1	.	+	+	1	+	+	+	+	.	+
<i>Geranium purpureum</i> L.	.	+	.	+	2	+	.	.	.	.	2	1	1	+
<i>Hyparrhenia hirta</i> (L.) Stapf subsp. <i>hirta</i>	+	+	.	.	.	.	+	1	+	+	+	.	.	.
<i>Capparis spinosa</i> L.	.	+	.	+	1	.	.	+	.	.	.	.	+	+
<i>Asphodelus ramosus</i> L. subsp. <i>ramosus</i>	.	+	.	+	.	.	.	.	+	+	.	+	.	+
<i>Mercurialis annua</i> L.	+	.	.	.	.	+	.	.	+	.	1	+	.	.
<i>Ferula communis</i> L. subsp. <i>communis</i>	.	.	.	.	.	.	+	1	.	1	.	.	.	+
<i>Charybdis pancration</i> (Steinh.) Speta	.	.	.	.	.	.	.	+	.	.	+	+	+	.
<i>Smyrniolum olusatrum</i> L.	.	.	+	1	.	.	.	.	+	.	.	.	.	.
<i>Brachypodium retusum</i> (Pers.) P.Beauv.	.	.	.	.	+	+	.	.	.	.	.	.	.	.
<i>Silene latifolia</i> Poir.	.	.	.	.	.	+	+	.	1	.	.	.	.	.
<i>Rubus ulmifolius</i> Schott	.	.	.	.	.	1	1	.	.	.	.	.	.	.
<i>Kundmannia sicula</i> (L.) DC.	.	.	.	.	.	.	+	+	.	.	.	.	.	.
<i>Umbilicus horizontalis</i> (Guss.) DC.	.	.	.	.	.	.	+	.	+	.	.	.	.	.
<i>Magydaris pastinacea</i> (Lam.) Paol.	.	.	.	.	.	.	.	.	.	+	.	.	+	.
<i>Cistus creticus</i> subsp. <i>creticus</i>	.	.	.	.	.	.	.	.	.	.	.	+	.	1
<i>Bituminaria bituminosa</i> (L.) E.H.Stirt.	.	.	.	.	.	+	.	.	.	.	.	.	.	.
<i>Crataegus monogyna</i> Jacq.	.	.	.	.	.	.	1	.	.	.	.	.	.	.



Although severely degraded and floristically impoverished, some cork oak wood nuclei still survive in the study area, forming linear features along dry stone walls and hedges which probably correspond to the past borders of the *fief*. These forest fragments host some majestic (more than 1 m of diameter at breast height: Fig. 5C) and very old individuals of *Quercus suber* L., and may be interpreted as an impoverished feature (Fig. 5D and Table 5) of *Stipa bromoidis-Quercetum suberis*, an open forest com-

munity described from the Erei Mounts (southeastern Sicily: Barbagallo 1983; Brullo et al. 2009 and references therein). Even if the main characteristic species of this association, i.e. *Aristella bromoides* (L.) Bertol. – also known as *Stipa bromoides* L. – was not observed in the study area, it has been recently reported to occur at Menfi and Castelvetrano (Scuderi and Pasta 2009) in other remnant fragments of open forestland once covering the Basso Belice lowland (see Table 6, relevé 3, from Scuderi 2006).



**Figure 5.** A, Evergreen maquis-forest nucleus referred to the *Chamaeropo humilis-Oleetum sylvestris acanthetosum mollis* (Habitat 9320) (photo: L. Gianguzzi, May 9, 2007); B, Forest nucleus dominated by *Quercus ilex* (Habitat 9340) (photo: L. Gianguzzi, May 9, 2007); C, Century-old tree of *Quercus suber* (photo: L. Gianguzzi, May 9, 2007); D, Forest strip dominated by *Quercus suber* (Habitat 9330) (photo: L. Gianguzzi, May 9, 2007).



Local fragments of *Quercus suber* woods probably represented the final stage on local well-drained and mostly sandy-loamy soils on palaeodunes. This head of series has been wiped out little by little to give room to vineyards and olive groves over wide areas of the surrounding territory. The degradation of cork oak woods usually leads to the spread of a patchy landscape made of broom-dominated shrublands referred to *Pyro amygdaliformis-Calicotometum infestae* – locally very rare due to the century-lasting removal of fuelwood – and garrigues (*Rosmarino-Thymetum capitati*) intermingled with perennial grasslands (*Hyparrhenietum hirta-pubescentis*) and annual therophytic swards belonging to *Helianthemetea guttati* Rivas Goday & Rivas-Mart. 1963). Like in the Erei territory (e.g., Niscemi, Caltagirone), on the deeper soils of the innermost part of southwestern Sicily such cork oak forests might have occurred in close contact with thermophilous woodlands dominated by *Quercus pubescens* s.l.. Downy oaks have completely disappeared from the whole territory, with the only exception of few remarkable isolated individuals growing in the bottom of the canyon of Castello di Pietra (Pasta et al. 2008) and of some little forest nuclei located in the bottom of Vallone San Vincenzo near Menfi (M. Buffa and S. Pasta, April 1994, unpubl.) and on a north-west-facing slope near the River Modione along the road connecting Partanna and Castelvetro (Lat. 37°41'00.9" N, Long. 12°49'49.1" E, A. Troia, pers. comm.).

With regards to the taxonomic value of *Q. virgiliana* as well as of various other taxa belonging to the collective group of *Q. pubescens* Willd., widely reported for Sicily and southern Italy in the last decades, a lively scientific debate is still going on (see Wellstein & Spada 2015; Pasta et al. 2016; Pignatti et al. 2017; Di Pietro et al. 2020, 2021).

Ancient floras suggest that until few centuries ago *Q. pubescens* s.l. was more common than today in the lowlands and near the coastline, while the few available data on vegetation dynamics after the abandonment of cork production (Di Benedetto et al. 1985; Di Pasquale and Garfi, 1998; Rühl et al. 2005; La Mela et al. 2007) and several co-occurrence data support the hypothesis that mixed cork oak/downy oak woodlands may have formed large forests in the thermo- and meso-Mediterranean belts of Sicily. Indeed, *Q. pubescens* s.l.-dominated forests probably represented a more common feature on clayey acid soils (e.g., Cullotta & Pasta 2004; Gianguzzi 2004; Gianguzzi et al. 2015), but underwent deep shrinkage as a result of century-long agropastoral activities.

At present, very few data were available on the occurrence of cork oak in the north-western and western Sicily (Gianguzzi et al. 2008; Badalamenti et al. 2020). As for the Tyrrhenian sector of the province of Trapani, *Q. suber* has been reported to occur only in few scattered forest/shrubland stands, such as the ‘macchie’ of Alcamo (Ponzo 1903; Borzi 1911), in locality Bosco Scorage/Arco-daci in the municipality of Buseto Palizzolo (Adragna 1976; Raimondo 1979), in the nature reserve of Zingaro in the municipalities of Castellamare del Golfo and San

**Table 4.** Holm oak wood stands (*Pistacio lentisci-Quercetum ilicis*). Relevés 1–2 (L. Gianguzzi, 9.05.2007, unpubl.).

Relevé (n°)	1	2
Altitude (m)	73	75
Slope (°)	20	15
Aspect	N	N
Area (m <sup>2</sup> )	100	100
Total cover (%)	100	80
Tree cover (%)	100	100
Scrub cover (%)	30	60
Herbaceous cover (%)	20	5
Average height of the dominant layer (m)	7.5	7.0
Species per relevé	26	21
<b>Char. association</b>		
<i>Quercus ilex</i> L.	5	5
<i>Pistacia lentiscus</i> L.	3	4
<i>Chamaerops humilis</i> L.	+	1
<i>Asparagus albus</i> L.	+	.
<b>Char. alliance (<i>Fraxino-Quercion ilicis</i>), order (<i>Quercetalia ilicis</i>) and class <i>Quercetea ilicis</i></b>		
<i>Smilax aspera</i> L.	2	3
<i>Olea europaea</i> L. var. <i>sylvestris</i> (Mill.) Lehr	2	1
<i>Phillyrea latifolia</i> L.	2	1
<i>Clematis cirrhosa</i> L.	2	1
<i>Rhamnus alaternus</i> L. subsp. <i>alaternus</i>	1	2
<i>Arisarum vulgare</i> O.Targ.Tozz. subsp. <i>vulgare</i>	1	2
<i>Rubia peregrina</i> L.	1	2
<i>Quercus suber</i> L.	1	2
<i>Pistacia lentiscus</i> L.	1	1
<i>Asparagus acutifolius</i> L.	1	1
<i>Allium subhirsutum</i> L. subsp. <i>subhirsutum</i>	1	+
<i>Stachys major</i> (L.) Bartolucci & Peruzzi	+	.
<i>Osyris alba</i> L.	+	.
<i>Daphne gnidium</i> L.	+	.
<i>Ampeledosmos mauritanicus</i> (Poir.) T.Durand & Schinz	+	.
<i>Teucrium fruticans</i> L. subsp. <i>fruticans</i>	.	+
<b>Other species</b>		
<i>Acanthus mollis</i> L. subsp. <i>mollis</i>	2	3
<i>Smyrniolum olusatrum</i> L.	2	+
<i>Oxalis pes-caprae</i> L.	1	1
<i>Crataegus monogyna</i> Jacq.	1	+
<i>Oloptum miliaceum</i> (L.) Röser & H.R.Hamasha	1	+
<i>Asphodelus ramosus</i> L. subsp. <i>ramosus</i>	+	+
<i>Stellaria neglecta</i> Weihe subsp. <i>cupaniana</i> (Jord. & Fourr.) Gutermann	+	.
<i>Silene latifolia</i> Poir.	+	.

Vito Lo Capo (Scuderi et al. 1994), at Bosco di Angimbé in the municipality of Calatafimi (Marguglio 1973; Scuderi 2006), on the slopes of Mt. Ferricini, west of Alcamo, (Scuderi 2006) and near the top of Mt. Inici in the municipality of Castellamare del Golfo (Scuderi 2006), where it was introduced after the Second World War and is now successfully established (Colomela et al. 2012).

Although south-western Sicily is correctly included in both the potential (Vessella and Schirone 2013) and the current (Pasta et al. 2000; EUFORGEN; Agrillo et al. 2018) distribution range of *Q. suber*, the information available about the distribution of local cork oak stands is extremely poor and need to be clarified. As for the area corresponding to the ancient forest of Birribaida, several neglected or unpublished occurrence data should be

**Table 5.** Degraded cork oak stands (*Stipo bromoidis-Quercetum suberis*) of Parche di Bilello (Relevés 1–2: L. Gianguzzi, 9.05.2007, unpubl.) and adjacent areas (Relevé 3: L. Scuderi, Locality Bresciana, municipality of Castelvetrano, 7.06.2005; cf. Scuderi, 2006).

Relevé (n°)	1	2	3
Altitude (m)	77	80	48
Slope (°)	5	2	2
Aspect	N	N	S
Area (m <sup>2</sup> )	100	100	200
Total cover (%)	100	90	80
Tree cover (%)	95	90	80
Scrub cover (%)	70	60	60
Herbaceous cover (%)	40	45	40
Average height of the dominant layer (m)	7.5	8.0	7.0
Species per relevé	19	15	19
<b>Char. of association and alliance <i>Erico arboreae-Quercion ilicis</i></b>			
<i>Quercus suber</i> L.	5	5	4
<i>Myrtus communis</i> L.	.	.	1
<i>Aristella bromoides</i> (L.) Bertol.	.	.	+
<b>Char. order <i>Quercetalia ilicis</i> and class <i>Quercetea ilicis</i></b>			
<i>Smilax aspera</i> L.	3	2	2
<i>Pistacia lentiscus</i> L.	2	2	2
<i>Rosa sempervirens</i> L.	1	2	2
<i>Olea europaea</i> L. var. <i>sylvestris</i> (Mill.) Lehr	2	1	1
<i>Rubia peregrina</i> L.	2	1	1
<i>Allium subhirsutum</i> L. subsp. <i>subhirsutum</i>	1	+	1
<i>Stachys major</i> (L.) Bartolucci & Peruzzi	1	+	+
<i>Asparagus acutifolius</i> L.	1	2	+
<i>Arisarum vulgare</i> O.Targ.Tozz. subsp. <i>vulgare</i>	+	1	.
<i>Rhamnus alaternus</i> L. subsp. <i>alaternus</i>	1	.	.
<i>Asparagus albus</i> L.	1	.	.
<i>Quercus ilex</i> L.	+	.	.
<i>Chamaerops humilis</i> L.	.	+	.
<i>Carex divulsa</i> Stokes	.	.	+
<b>Other species</b>			
<i>Acanthus mollis</i> L. subsp. <i>mollis</i>	2	1	.
<i>Geranium purpureum</i> L.	+	+	.
<i>Euphorbia ceratocarpa</i> Ten.	1	.	1
<i>Hyparrhenia hirta</i> (L.) Stapf subsp. <i>hirta</i>	1	.	1
<i>Oloptum miliaceum</i> (L.) Röser & H.R.Hamasha	+	.	2
<i>Rubus ulmifolius</i> Schott	.	1	1
<i>Silene latifolia</i> Poir.	+	.	.
<i>Smyrniium olusatrum</i> L.	.	+	.
<i>Crataegus monogyna</i> Jacq.	.	.	1
<i>Cistus creticus</i> L. subsp. <i>eriocephalus</i> (Viv.) Greuter & Burdet	.	.	+
<i>Solanum villosum</i> Mill.	.	.	+

added to those concerning localities Parche di Bilello and Bresciana (Table 6). In fact, cork oaks were noticed growing near Sciacca by J. W. Goethe in April 1787 during his travel across Sicily (Goethe 1813–1817), and the occurrence of the toponym ‘Suvarito’ on the map ‘Sciacca’ (IGMI 1973) suggests that *Q. suber* may have only recently gone extinct. In fact, cork oaks old records were also reported for Castelvetrano (De Stefani-Perez 1898). An additional indirect hint about the presence and the exploitation of *Q. suber* in the study area is the fact that cork dishes were part of the daily equipment of the countrymen of the Belice area until 1970s (Cusumano 1978).

Additionally, small stands and isolated trees of *Q. suber* still occur along the roadsides of the ancient Feudo (=

fiefdom) of Dimina between Castelvetrano and Partanna (Pasta et al. 2008), between Castelvetrano and Menfi (S. Pasta and S. Mattana, August 1998, unpubl.), as well as near Marinella di Selinunte (37°37'03" N, 12°51'10" E) and between Triscina and Parche di Bilello (Lat. 37°36'44.3" N, Long. 12°47'50.6" E, A. Troia, pers. comm.).

## Biogeographic remarks

The discovery of *Linaria multicaulis* subsp. *humilis* and *Serapias orientalis* subsp. *siciliensis*, discussed in the present paper, together with the recent finding of *Retama raetam* (Forssk.) Webb subsp. *gussonei* (Webb) W. Greuter (Troia and Spallino 2009), *Aristella bromoides* and *Tube-raria villosissima* (Pomel) Grosser subsp. *sicula* (Grosser) Bartolo, Pulvirenti et Salmeri (Scuderi and Pasta 2009) underlines the strong floristic similarity between the coastal lowlands (sciare) of southwestern Sicily, stretching from Marsala to Sciacca, and those of southeastern Sicily (including the territories of Comiso, Gela, Vittoria, Camarina, Caltagirone, Niscemi, Mazzarino, etc.). This latter area roughly coincides with the ‘Kamarino-Pachynense’ district, identified by Brullo et al. (1995), which probably does not deserve to be treated as a separate biogeographic unit, considering the very recent emersion of these lands (Mid to Upper Pleistocene) and the few plant taxa which result to grow exclusively there.

## Conclusions

As already highlighted by Raimondo et al. (2005) and Scuderi and Pasta (2009), the Province of Trapani is well, yet unevenly studied. Recent field surveys carried out in this territory allowed to find species new to science (e.g., Troia and Raimondo 2010) and to confirm the regional occurrence of very rare and globally endangered ones (e.g., Troia and Lansdown 2016). The improvement of the knowledge involves the need for more appropriate conservation measures to prevent the last populations of several vulnerable and legally protected species and habitats from being wiped out due to severe land-use intensification. For these reasons, taking into account not only the occurrence of a rare lichen of conservation interest (*Teloschistes chrysophthalmus*), but also considering the presence of a vascular plant of Community priority interest (*Ophrys lunulata*) and of several habitats (6220\*, 9320, 9330, 9340) of Community interest according to 92/43 EU ‘Habitats’ Directive (Table 6), we strongly recommend the institution of a new Site of Community Importance (SCI) to protect the site of Parche di Bilello, or to re-shape the ones already existing in the nearby, a strategy which has already been adopted during last years in the Province of Trapani (Troia et al. 2011, 2016).

On this purpose, we should underline that our study area is not far from the River Modione. This watercourse which still hosts several fragments of riparian forest re-

**Table 6.** Lichens, Vascular plants and habitat types of high biogeographic/conservation interest occurring at Parche di Bilello. Since 1995 all the European Orchids are included within the Appendix I of CITES. \* = species listed in the Appendixes or habitat of priority interest according to 92/43 EU Directive.

Taxon	Endemic	Narrow Range	Protected/Red-Listed	Other Criteria
<i>Acis autumnalis</i>				x
<i>Allium obtusiflorum</i>	x			
<i>Ambrosina bassii</i>			x	
<i>Anacamptis papilionacea</i> subsp. <i>grandiflora</i>			x	
<i>Barlia robertiana</i>			x	
<i>Carlina sicula</i> subsp. <i>sicula</i>	x			
<i>Desmazeria sicula</i>				x
<i>Echium italicum</i> subsp. <i>siculum</i>	x			
<i>Euphorbia ceratocarpa</i> .		x		
<i>Linaria multicaulis</i> subsp. <i>humilis</i>	x			x
<i>Ophioglossum lusitanicum</i>			x	x
<i>Ophrys apifera</i>			x	
<i>Ophrys lunulata</i>	x		*	x
<i>Ophrys lutea</i> subsp. <i>lutea</i>			x	
<i>Ophrys sicula</i>			x	
<i>Ophrys speculum</i>			x	
<i>Ophrys tenthredinifera</i> subsp. <i>grandiflora</i>		x	x	
<i>Orobanche balsensis</i>				x
<i>Pimpinella anisoides</i>		x		
<i>Serapias orientalis</i> subsp. <i>siciliensis</i>	x		x	x
<i>Tragopogon cupanii</i>		x		
<b>Habitats of the 92/43 EU Directive occurring in the study area</b>	<b>Endemic</b>	<b>Narrow Range</b>	<b>Protected/Red-Listed</b>	<b>Other Criteria</b>
6220			*	
9320			x	x
9330			x	x
9340			x	
<b>Habitats of the 92/43 EU Directive occurring in the nearby territory</b>	<b>Endemic</b>	<b>Narrow Range</b>	<b>Protected/Red-Listed</b>	<b>Other Criteria</b>
92A0			x	
92D0			x	

ferred to *Salicetum albo-pedicellatae* (*Salicion albae*). Moreover, several spots of thermo-hygrophilous scrub (*Tamarix africana*-dominated community) occur on the seasonally dry beds and the riversides of this river, while reed meadows (*Phragmitetum communis*) colonize its thalweg (data not shown). All these data suggest to promote a more comprehensive investigation of the size, importance and distribution of other patches of semi-natural habitats testifying for the previous extension of local woodlands and riparian forests. For instance, taking into account a wider portion of the territory may allow to produce a more articulated proposal for the institution of a new site of Community interest or to create the ecological corridors needed to connect the site of Parche di Bilello with an already existing hub of the Sicilian Natura 2000 network, i.e. ITA010005 'Laghetti di Preola e Gorghi Tondi e Sciare di Mazara', located within a 15 km radius from our study area.

The study area worths to be protected and valorized for the high symbolic value of the last remnant nuclei of forest vegetation, especially considering that the Parche di Bilello host the best conserved nuclei of wild olive forests occurring in the entire province of Trapani and the

last fragments of cork oak woods referred to *Stipo bromoidis-Quercetum suberis*, completely wiped out from the whole south-western Sicily. Since 2014 the management of the study area is under the responsibility of the ONG 'Cooperativa Sociale Onlus Rita Atria-Libera Terra', aiming at promoting legality and traditional extensive agricultural practices. A best practice could be to assign to this ONG the role of guardian of this area, representing a precious tile survived to the destruction of the past coastal woodland/shrubland/grassland mosaic.

The presence of bifacial pebble tools, probably dating back to Mesolithic and testifying for the long-lasting history of local human presence, has been reported for the study area (Leighton 1999; Lo Vetro and Martini 2012). As a consequence, Parche di Bilello also features among the sites of archaeological interest (Regione Siciliana 2018) according to article 142 (letter 'm') of the Decree 42 of year 2004 published by the Sicilian Regional Authority. The co-occurrence of cultural and natural heritage in the same area, a frequent feature in Mediterranean areas, should be viewed - here as elsewhere - as a chance to support the proposal of strengthening the conservation measures for this site.



## Syntaxonomic scheme

QUERCETEA ILICIS Br.-Bl. ex A. Bolòs et O. de Bolòs in A. Bolòs y Vayreda 1950

QUERCETALIA ILICIS Br.-Bl. ex Molinier 1934

**Fraxino orni-Quercion ilicis** Biondi, Casavecchia et Gigante in Biondi et al. 2013

*Pistacio lentisci-Quercetum ilicis* Brullo et Marcenò 1985

**Erico arboreae-Quercion ilicis** S. Brullo, De Martino et Marcenò 1977

*Stipo bromoidis-Quercetum suberis* Barbagallo 1983

PISTACIO LENTISCI-RHAMNETALIA ALATERNI Rivas-Martínez 1975

**Oleo sylvestris-Ceratonion siliquae** Br.-Bl. 1936 em. Rivas-Martínez 1975

*Pistacio lentisci-Chamaeropetum humilis* Brullo et Marcenò 1985

*Chamaeropo humilis-Oleetum sylvestris acanthetosum mollis* Gianguzzi et Bazan 2019

*Pyro amygdaliformis-Calicotometum infestae* Gianguzzi et La Mantia 2008

ONONIDO-ROSMARINETEA Br.-Bl. in A. Bolos y Vayreda 1950

ROSMARINETALIA OFFICIALIS Br.-Bl. ex Molinier 1934

**Cisto eriocephali-Ericion multiflorae** Biondi 2000

*Rosmarino officinalis-Thymetum capitati* Furnari 1965

LYGEO SPARTI-STIPETEA TENACISSIMAE Rivas-Martínez 1978

CYMBOPOGONO-BRACHYPODIETALIA RAMOSI Horvatic 1963

**Hyparrhenion hirtae** Br.-Bl., P. Silva et Rozeira 1956

*Hyparrhenietum hirta-pubescentis* A. et O. de Bolòs et Br.-Bl. in A. et O. Bolòs 1950

STIPO-TRACHYNIETEA DISTACHYAE Brullo in Brullo, Scelsi et Spampinato 2001

BRACHYPODIETALIA DISTACHYI Rivas-Martínez 1978

**Trachynion distachyae** Rivas-Martínez 1978

*Sedetum caerulei* Brullo 1975

STIPO-BUPLEURETALIA SEMICOMPOSITI Brullo in Brullo, Scelsi et Spampinato 2001

**Plantagini-Catapodion marini** Brullo 1985

*Ononido breviflorae-Stipetum capensis* Brullo, Guarino et Ronsisvalle 1998

SALICETEA PURPUREAE Moor 1958

SALICETALIA PURPUREAE Moor 1958

**Salicion albae** Soo 1951

*Salicetum albo-pedicellatae* Brullo et Spampinato 1991

NERIO-TAMARICETEA Br.-Bl. et O. de Bolòs 1958

TAMARICETALIA AFRICANAE Br.-Bl. et O. de Bolòs 1958

**Tamaricion africanae** Br.-Bl. et O. de Bolòs 1958

*Tamarix africana*-dominated community

PHRAGMITO-MAGNOCARICETEA Klika in Klika et V. Novák 1941

PHRAGMITETALIA KOCH 1926

**Phragmition communis** Koch 1926

*Phragmitetum communis* Koch 1926

## Other syntaxa quoted in the text

*Ammophiletea* Br.-Bl. & Tx. ex Westhoff, Dijk & Passchier 1946; *Chenopodietea* Br.-Bl. in Br.-Bl. et al. 1952; *Crithmo-Staticetea* Br.-Bl. in Br.-Bl., Roussine & Nègre 1952; *Diplotaxion eruroidis* Br.-Bl. in Br.-Bl., Gajewski, Wraber & Walas 1936; *Eragrostietalia* J. Tx. ex Poli 1966

## Acknowledgements

We are indebted to Angelo Troia (Dept. STEBICEE, University of Palermo) for sharing unpublished personal data. We also thank two anonymous reviewers and the editorial board of Plant Sociology, whose critical comments and suggestions significantly improved the quality of the manuscript.

## Bibliography

- Adragna V (1976) Il grande bosco di Arcodaci. Un'oasi poco nota nel comune di Buseto Palizzolo. "Trapani"- Rassegna della Provincia XXI (217): 19–25.
- Agrillo E, Alessi N, Jiménez-Alfaro B, Casella L, Angelini P, Argagno O, Crespo G, Fernández-González F, Monteiro-Henriques T, Silva Neto C, Attorre F (2018) The use of large databases to characterize habitat types: The case of *Quercus suber* woodlands in Europe. *Rendiconti Scienze Fisiche e Naturali dell'Accademia dei Lincei* 29: 283–293. <https://doi.org/10.1007/s12210-018-0703-x>
- Aleo M, Ottonello D (2000) La flora fanerogamica. In: Gianguzzi L, Ottonello D (eds) La Riserva di Monte Cofano (Sicilia nord-occidentale). Aspetti geomorfologici, naturalistici ed etnoantropologici, Collana Sicilia Foreste n° 8, Azienda Foreste Demaniali della Regione Siciliana, Palermo: 67–86.
- Amico Statella VM (1759) *Lexicon Topographicum Siculum*. In quo Siciliae Urbes, Oppida, cum diruta, tum extantia, Montes, Flumina, Portus, adjacentes Insulae, ac singula loca describuntur, illustrantur. In: G. Di Marzo (ed, 1855) *Dizionario topografico della Sicilia*, vol. 1, Tip. P. Morvillo, Palermo.
- Badalamenti E, Barone E, Pasta S, Sala G, La Mantia T (2012) *Ailanthus altissima* (Mill.) Swingle (fam. *Simaroubaceae*) in Sicilia e cenni storici sulla sua introduzione in Italia. *Il Naturalista siciliano* (4) 36(1): 117–164. <https://www.ssn.it/PDF/PDF%20NS%2036/117-164.pdf>
- Badalamenti E, Scalenghe R, da Silveira Bueno R, Sala G, Pizzurro GM, Giaimo A, Pasta S (2020) The cork oak in the Mountains of Palermo (Italy): Ecological insights from the south-eastern edge of its distribution range. *iForest* 13(4): 336–344. <https://doi.org/10.3832/ifer3360-013>

- Barbagallo C (1984) Vegetazione di alcuni boschi di sughera (*Quercus suber* L.) della Sicilia meridionale-orientale. Bollettino dell'Accademia Gioenia di Scienze naturali 16(321)(1983): 289–296.
- Barbagallo C, Brullo S, Guglielmo A (1980) Lineamenti della vegetazione di Monte Cofano (Sicilia occidentale). Pubblicazioni dell'Istituto di Botanica dell'Università di Catania (2) 14 pp., 6 tables.
- Bartolo G, Brullo S (1993) La classe *Crithmo-Limonietea* in Sicilia. Bollettino dell'Accademia Gioenia di Scienze naturali (4) 26 (342): 5–47.
- Bartolo G, Brullo S, Minissale P, Spampinato G (1992) Contributo alla conoscenza dei boschi a *Quercus ilex* della Sicilia. Acta Botanica Malacitana 15 (1990): 203–215. <https://doi.org/10.24310/abm.v15i.9303>
- Bartolo G, Pulvirenti S (1993) *Serapias orientalis* subsp. *siciliensis* (Orchidaceae): a new subspecies from Sicily. Candollea 48(1): 231–236.
- Bartolucci F, Peruzzi L, Galasso G, Albano A, Alessandrini A, Ardenghi NMG, et al. (2018) An updated checklist of the vascular flora native to Italy. Plant Biosystems 152 (2): 179–303. <https://doi.org/10.1080/11263504.2017.1419996>
- Bazan G, Marino P, Guarino R, Domina G, Schicchi R (2015) Bioclimatology and vegetation series in Sicily: a geostatistical approach. Annales Botanici Fennici 52: 1–18. <https://doi.org/10.5735/085.052.0202>
- Bernhardt KG (1986) *Oxalis pes-caprae* L., ein anpassungsfähiger Neophyt in Sizilien. Bauhinia 8(3): 141–148. <https://doi.org/10.5735/085.052.0202>
- Biondi E (2011) Phytosociology today: Methodological and conceptual evolution. Plant Biosystems 145(Suppl. 1): 19–29. <https://doi.org/10.1080/11263504.2011.602748>
- Biondi E, Blasi C, Allegranza M, Anzellotti I, Azzella MM, Carli E, Casavecchia S, Copiz R, Del Vico E, Facioni L, Galdenzi D (2014) Plant communities of Italy: the vegetation prodrome. Plant Biosystems 148(4): 728–814. <https://doi.org/10.1080/11263504.2014.948527>
- Bivona-Bernardi A (1845) Miglioramenti dei boschi dello Stato in Sicilia. Stamperia F. Solli, Palermo, 138 pp.
- Blasi C, Marignani M, Copiz R, Fipaldini M, Bonacquisti S, Del Vico E, Rosati L, Zavattero L, 2010. Important plant areas in Italy: from data to mapping. Biological Conservation 144(1): 220–226. <https://doi.org/10.1016/j.biocon.2010.08.019>
- Bonanno A, Ciabatti P, Liguori V, Provenzano MC, Sortino G (2000) Studio idrogeologico e idrogeochimico dell'acquifero multifalda della Piana di Castelvetro e Campobello di Mazara (Sicilia occidentale). Quaderni di Geologia applicata 7 (4): 45–59.
- Borzi A (1911) Le quercie della flora italiana. Bollettino del reale Orto botanico e Giardino coloniale di Palermo 10: 41–66.
- Braun-Blanquet J (1964) Pflanzensoziologie. Springer Verlag, Wien - New York. <https://doi.org/10.1007/978-3-7091-8110-2>
- Bresc H (1975) Una flotta mercantile periferica: la marina siciliana medievale. In: Bresc H, Gatti L, Grendi E, Borzone P (eds) Studi di Storia Navale. Centro per la Storia della Tecnica in Italia del Consiglio Nazionale delle Ricerche, Giunti, Firenze, 7–24.
- Bresc H (1980) La chasse en Sicile (XIIe-XVe siècles). In: Publications de la Faculté de Lettres de Nice n° 20 - Actes du Colloque 'La Chasse au Moyen âge' (Nice, 22–24 Juin 1979). Les Belles Lettres, Paris, 201–211.
- Bresc H (1986) Un monde méditerranéen. Economie et société en Sicile (1300–1450). Ecole française de Rome, Accademia di Scienze Lettere e Arti di Palermo, Bibliothèque des écoles françaises d'Athènes et de Rome n° 262, 981 pp. <https://doi.org/10.3406/befar.1986.1245>
- Bresc H, Sciascia L (2016) All'ombra del grande Federico: Riccardo da Lentini architetto. Edizioni Torre del Vento, Palermo, 91 pp.
- Bresc-Bautier G (1983) "Disfari et perdiri li fructi et li aglandi": Economie e risorse boschive nella Sicilia medievale (XIII–XV secolo). Quaderni storici 54(3): 941–969.
- Brullo S (1985) Sur la syntaxonomie des pelouses thérophytiques des territoires steppiques de l'Europe sud-occidentale. Documents Phytosociologiques (n.s.)9: 1–24.
- Brullo C, Brullo S, Giusso del Galdo G, Guarino R, Minissale P, Scuderi L, Siracusa G, Sciandrello S, Spampinato G (2010) The *Lygeo-Stipetea* class in Sicily. Annali di Botanica (4) 0: 57–84 + 21 tables. <https://annalidibotanica.uniroma1.it/index.php/Annalidibotanica/article/view/9119>
- Brullo S, Di Martino A, Marcenò C (1974) Osservazioni sulla vegetazione psammofila tra Capo Granitola e Selinunte (Sicilia occidentale). Bollettino di Studi e Informazioni del reale Giardino coloniale di Palermo 26: 103–110.
- Brullo S, Furnari F (1976) Le associazioni vegetali degli ambienti palustri costieri della Sicilia. Notiziario Fitosociologico 11: 1–43.
- Brullo S, Furnari F (1978) La vegetazione palustre in Sicilia. Atti II Convegno siciliano di Ecologia (Noto, SR, 23–25 ottobre 1977): 29–39.
- Brullo S, Gianguzzi L, La Mantia A, Siracusa G (2009) La classe *Quercetea ilicis* in Sicilia. Bollettino dell'Accademia Gioenia di Scienze naturali 41 (369)(2008): 1–77.
- Brullo S, Marcenò C (1980) Il *Diplotaxion eruroidis* in Sicilia, con considerazioni sulla sintassonomia e distribuzione. Notiziario Fitosociologico, 15 (1978): 27–44.
- Brullo S, Marcenò C (1985) Contributo alla conoscenza della classe *Quercetea ilicis* in Sicilia. Notiziario fitosociologico 19(1) (1984): 183–229.
- Brullo S, Minissale P, Spampinato G (1995) Considerazioni fitogeografiche sulla flora della Sicilia. Ecologia Mediterranea 21(1–2): 99–117, 1 figure. <https://doi.org/10.3406/ecmed.1995.1759>
- Brullo S, Minissale P, Spampinato G (1997) La classe *Cisto-Micromerietea* nel Mediterraneo centrale e orientale. Fitosociologia 32: 29–60.
- Brullo S, Pavone P (1985) Taxonomic considerations on the genus *Desmazeria* (Gramineae) with description of a new species: *Desmazeria pignattii*. Willdenowia 15(1): 99–106.
- Brullo S, Ronsisvalle GA (1975) La vegetazione dei Gorgi Tondi e del Lago Preola presso Mazara del Vallo (Sicilia occidentale). Notiziario fitosociologico 10: 45–67.
- Brullo S, Scelsi F, Siracusa G (1994) Contributo alla conoscenza della vegetazione terofitica della Sicilia occidentale. Bollettino dell'Accademia Gioenia di Scienze naturali (4) 27(346): 341–365.
- Calamia P, La Barbera M, Salluzzo G (2004) Bellumvider. La Reggia di Federico II di Svevia a Castelvetro. Grafill s.r.l., Palermo, 118 pp.
- Calò C, Henne PD, Curry BB, Magny M, van Leeuwen JFN, Vescovi E, La Mantia T, Pasta S, Vannièr B, Tinner W (2012) Spatio-temporal patterns of Holocene vegetation change in southern Sicily. Palaeogeography, Palaeoclimatology, Palaeoecology 323–325: 110–122. <https://doi.org/10.1016/j.palaeo.2012.01.038>
- Campo G, Romano S (2000) Le Orchidacee: 177–193. In: Gianguzzi L, Ottonello D (eds), "La Riserva di Monte Cofano (Sicilia nord-occidentale). Aspetti geomorfologici, naturalistici ed etnoantropologici", Collana Sicilia Foreste n° 8, Azienda Foreste Demaniali della Regione Siciliana. Palermo.
- Capozzi F, Panepinto F, Tretiach M (2013) Notulae Cryptogamicae 7 (Lichenes), Notula 43: *Teloschistes chrysophthalmus* (L.) Th. Fr. (*Teloschistaceae*). Informatore Botanico Italiano 45(2): 316.

- Carlón L, Laínz M, Moreno Moral G, Sánchez Pedraja O (2015) The overlooked *Orobanche balsensis* (J.A. Guim.), comb. nov., and some remarks on *O. subbaetica* Triano & A. Pujadas. *Flora Montiberica* 60: 38–53.
- Caruso E, Nobili A (eds.) (2001) Le mappe del Catasto Borbonico di Sicilia. Territori comunali e centri urbani nell'Archivio Cartografico Mortillaro di Villarena (1837–1853). Regione Siciliana, Assessorato ai Beni Culturali e Ambientali. e Pubblica Istruzione, Palermo, 569 pp.
- Catanzaro F (1983) *Cardopatum corymbosum* (L.) Pers., pianta nuova per la flora siciliana. *Il Naturalista siciliano* (4) 7(1–4): 82–83.
- Colomela D, Pasta S, Scarselli D (2012, unpublished) Relazione sugli aspetti agro-forestali, botanici (flora vascolare e habitat) ed idrogeologici della ZPS ITA010029 'Monte Cofano, Capo San Vito e Monte Sparagio'. Progetto LIFE09 NAT/IT/000099 "SICALECONS - Azioni urgenti per la conservazione della Coturnice di Sicilia, Alectoris graeca whitakeri Schiebel, 1934', Azione A3: Stesura di una cartografia dell'habitat ed idrografica georiferita della ZPS ITA010029 "Monte Cofano, Capo San Vito e Monte Sparagio". Palermo: Assessorato Risorse Agricole e Alimentari della Regione Siciliana, 54 pp.
- Cullotta S, Pasta S (2004) Vegetazione mediterranea: Sicilia, Sardegna, Calabria. In: Blasi C, Bovio G, Corona P, Marchetti M, Maturani A (eds), Incendi e complessità ecosistemica. Dalla pianificazione forestale al recupero ambientale.- Ministero dell'Ambiente e della Tutela del Territorio, Direzione per la Protezione della Natura - Società Botanica Italiana - Commissione per la Promozione della ricerca botanica, Palombi Ed., Roma, 291–307.
- Cusumano A (1978) Mestieri e lavoro contadino nella Valle del Belice. Associazione per la Conservazione delle Tradizioni popolari, Studi e materiali per la storia della cultura popolare vol. 5, Palermo, 3–34.
- Cusumano A (1983) Arti e mestieri nella Valle del Belice: il cuoio, il legno, il ferro. Associazione per la Conservazione delle Tradizioni popolari, Studi e materiali per la storia della cultura popolare vol. 13, Palermo, 70 pp.
- De Leonardis W, De Santis C, Fichera G, Giardina G., Zizza A (2003) *Linaria multicaulis* (L.) Miller (*Scrophulariaceae*) in Sicily: An investigation within its subspecific and varietal ranks. *Bocconea* 16(2): 585–595. <https://www.herbmedit.org/bocconea/16-0585.pdf>
- De Leonardis W, Giardina G, Zizza A (1999) *Linaria multicaulis* (L.) Miller subsp. *humilis* (Guss.) De Leonardis, Giardina & Zizza, comb. et stat. nov., a taxon growing in Sicily. *Flora Mediterranea* 9: 97–111. <https://www.herbmedit.org/flora/9-097.pdf>
- De Stefani-Perez T (1898) Note intorno ad alcuni Zoocedidii del *Quercus robur* e del *Q. suber* raccolti nel territorio di Castelvetro (Sicilia). *Il Naturalista siciliano* (2) 2(5–8): 156–174.
- Di Benedetto G, Maugeri G, Poli Marchese E (1985) Principali tappe del dinamismo della vegetazione nelle sugherete della Sicilia Sud-Orientale. *Notiziario Fitosociologico* 19(1)(1984): 5–12.
- Di Martino A, Perrone C (1970) Flora delle isole dello Stagnone (Marsala). I. Isola Grande. *Lav. Ist. Bot. Giard. colon. Palermo* 24: 109–166, 18 figures, 2 tables.
- Di Pasquale G, Garfi G (1998) Analyse comparée de l'évolution de la régénération de *Quercus suber* et *Quercus pubescens* après élimination du pâturage en forêt de Pisano (Sicile sud-orientale). *Ecologia Mediterranea* 24(1): 15–25. <https://doi.org/10.3406/ecmed.1998.1843>
- Di Pietro R, Conte AL, Di Marzio P, Fortini P, Farris E, Gianguzzi L, Muller M, Rosati L, Spampinato G, Gailing O (2021) Does the genetic diversity among pubescent white oaks in southern Italy, Sicily and Sardinia islands support the current taxonomic classification? *European Journal of Forest Research* 140(2): 1–17. <https://doi.org/10.1007/s10342-020-01334-z>
- Di Pietro R, Conte AL, Di Marzio P, Gianguzzi L, Spampinato G, Caldarella O, Fortini P. (2020) A multivariate morphometric analysis of diagnostic traits in southern Italy and Sicily pubescent oaks. *Folia Geobotanica* 55(3): 163–183. <https://doi.org/10.1007/s12224-020-09378-0>
- Duro A, Piccione V, Scalia C, Zampino S (1996) Precipitazioni e temperature medie mensili in Sicilia relative al sessantennio 1926–1985. *Atti 5° Workshop Progr. Strat. C.N.R. 'Clima Ambiente e Territorio nel Mezzogiorno d'Italia'* 1: 17–109.
- EUFORGEN (= European Forest Genetic Resources Programme), *Quercus suber* (<http://www.euforgen.org/species/quercus-suber/>) (last accessed: November 2, 2020)
- Fanales F (1899) Contributo alla conoscenza della flora delle sciere di Marsala. *Bollettino del reale Orto botanico e Giardino coloniale di Palermo* 3 (1–2): 1–65.
- Fazello T (1560) *De rebus siculis decades duae*. Typis excudebant Ioannes Matthaeus Mayda, et Franciscus Carrara, Panormi.
- Ferrigno GB (1931, unpublished) Teatro e giuochi a Castelvetro. Reprint in 2001 by Arti grafiche Corrao, Trapani.
- Ferrigno GB (1933) Guida di Selinunte. Società Tipografica Ospizio e Beneficenza, Palermo, 52 pp.
- Fierotti G (1988) Carta dei Suoli della Sicilia scala 1:250.000 e note illustrative. Istituto di Agronomia Generale, Università degli Studi di Palermo, Palermo.
- Frei M (1937) Studi fitosociologici su alcune associazioni litorali in Sicilia (*Ammophiletalia* e *Salicornietalia*). *Nuovo Giornale botanico italiano* (2) 44 (2): 273–294. <https://doi.org/10.1080/11263503709437958>
- Galasso G, Conti F, Peruzzi L, Ardenghi NMG, Banfi E, Celesti-Grapow L, et al. (2018) An updated checklist of the vascular flora alien to Italy. *Plant Biosystems* 152 (3): 556–592. <https://doi.org/10.1080/11263504.2017.1419996>
- Galesi R (1996) Le Orchidee del Comprensorio niscemese. Comune di Niscemi, Assessorato Territorio e Ambiente, 48 pp.
- Géhu JM, Rivas-Martínez S (1981) Notions fondamentales de phytosociologie. In: Dierschke H (ed) *Syntaxonomie*. *Berichte International Symposium IV-V*. J. Cramer, Vaduz: 5–33.
- Gennai M, Fenu G, Bacchetta G, Foggi B, Lombardi L, Pasta S, Scuderi L, Domina G, Wagensommer RP (2015) Schede per una Lista Rossa della Flora vascolare e crittogamica Italiana. *Pallenis maritima* (L.) Greuter. *Informatore botanico italiano* 47(1): 99–137.
- Geraci A, Raimondo FM, Troia A (2010) Genetic diversity and local population structure in *Ambrosina bassii* (*Araceae*, *Ambrosineae*), a Mediterranean relict species. *Biochemical Systematics and Ecology* 37(6): 737–746. <https://doi.org/10.1016/j.bse.2009.12.002>
- Gianguzzi L (2004) L'evoluzione del paesaggio vegetale del comprensorio in una prospettiva storica. In: Gianguzzi L (ed), *Il paesaggio vegetale della Riserva Naturale Orientata "Bosco della Ficuzza, Rocca Busambra, Bosco del Cappelliere, Gorgo del Drago"*. Palermo: Azienda Foreste Demaniali della Regione Siciliana, Collana "Sicilia Foreste" 22: 21–46.
- Gianguzzi L, Bazan G (2019a) A phytosociological analysis of the *Olea europaea* L. var. *sylvestris* (Mill.) Lehr. forests in Sicily. *Plant Biosystems* 154(5): 705–725. <https://doi.org/10.1080/11263504.2019.1681532>
- Gianguzzi L, Bazan G (2019b) The *Olea europaea* L. var. *sylvestris* (Mill.) Lehr. forests in the Mediterranean area. *Plant Sociology* 56(2): 3. <https://doi.org/10.1080/11263504.2019.1681532>



- Gianguzzi L, Cusimano D, Bonventre V, Romano S, Ilardi V (2012) Bio-ecological, phytosociological and conservation aspects of relictual and disjointed populations of *Simethis mattiazzii* (Vandelli) Sacc. (*Xanthorrhoeaceae*) in the Channel of Sicily. *Acta Botanica Gallica* 159(3): 303–318. <https://doi.org/10.1080/12538078.2012.737141>
- Gianguzzi L, Cusimano D, Cuttonaro P, Romano S (2016) Contribution to the phytosociological characterisation of the forest vegetation of the Sicani Mountains (inland of the north-western Sicily). *Plant Sociology* 53(1): 5–42. <http://dx.doi.org/10.7338/pls2016531/02>
- Gianguzzi L, D'Amico A, Romano S (2010) Phytosociological remarks on residual woodlands of *Laurus nobilis* in Sicily. *Lazaroa* 31: 67–84. [https://doi.org/10.5209/rev\\_LAZA.2010.v31.4](https://doi.org/10.5209/rev_LAZA.2010.v31.4)
- Gianguzzi L, Ilardi V, Raimondo FM (1996) La vegetazione del promontorio di Monte Pellegrino (Palermo). *Quaderni di Botanica ambientale e applicata* 4 (1993): 79–137.
- Gianguzzi L, La Mantia A (2009) Contributo alla conoscenza della vegetazione e del paesaggio vegetale della Riserva Naturale "Monte Cofano" (Sicilia occidentale) (con allegata Carta sinfitosociologica della vegetazione, scala 1:20.000). *Fitosociologia* 45 (1, suppl. 1): 1–55.
- Gianguzzi L, La Mantia A, Ottonello D, Romano S (2005) La flora vascolare della Riserva Naturale Monte Cofano (Sicilia Occidentale). *Il Naturalista siciliano* (4) 29(3–4): 107–152.
- Gianguzzi L, Papini F, Cusimano D (2016) Phytosociological survey vegetation map of Sicily (Mediterranean region). *Journal of Maps* 12 (5): 845–851. <https://doi.org/10.1080/17445647.2015.1094969>
- Gianguzzi L, Romano S, Caldarella O, La Rosa A. (2008) Osservazioni fitosociologiche ed ecologiche su una formazione forestale a *Quercus suber* dei Monti di Palermo (Sicilia centro-occidentale) percorsa dal fuoco nell'estate del 2007. 103° Congr. Soc. Bot. Ital. (Reggio Calabria, 17–19 Settembre 2008), riassunti: 327.
- Gianguzzi L, Scuderi L, Pasta S (2006) La flora vascolare dell'Isola di Marettimo (Arcipelago delle Egadi, Canale di Sicilia): aggiornamento ed analisi fitogeografica. *Webbia* 61 (2): 359–402. <https://doi.org/10.1080/00837792.2006.10670810>
- Goethe (von) JW (1813–1817) *Reise-Tagebuch 1786. Tagebuch der Italienischen Reise für Frau von Stein*. 2 vols.
- Grandis M. (2016) Sulla presenza di *Chasmanthe floribunda* e *Chasmanthe aethiopica* in Italia. *Acta Plantarum Notes* 4: 137–139. [https://www.academia.edu/40184939/ActaPlantarum\\_Notes\\_4\\_maggio](https://www.academia.edu/40184939/ActaPlantarum_Notes_4_maggio)
- Guarino R, Pasta S (2017) Botanical Excursions in Central and Western Sicily. *Field Guide for the 60th IAVS Symposium* (Palermo, 20–24 June 2017). Palermo University Press, Palermo, 606 pp. [https://www.academia.edu/35613890/Botanical\\_Escursions\\_in\\_Central\\_and\\_Western\\_Sicily](https://www.academia.edu/35613890/Botanical_Escursions_in_Central_and_Western_Sicily)
- Guarino R, Pasta S (2018) Sicily: the island that didn't know to be an archipelago. *Berichte der Reinhold Tüxen Gesellschaft* 30: 133–148.
- Gussone G (1827) *Florae Siculae Prodromus sive plantarum in Siciliae ulteriori nascentium enumeratio secundum systema Linneanum disposita*. Vol. 1. Ex Regia Typographia, Neapoli. <https://doi.org/10.5962/bhl.title.6700>
- Gussone G (1832–1834) *Supplementum ad Florae Siculae Prodromum, quod et specimen florae insularum Siciliae ulteriori adjacentium*. Fasciculi 2 Ex Regia Typographia, Neapoli.
- Gussone G (1842–1845) *Florae Siculae Synopsis exhibens plantas vasculares in Sicilia insulisque adjacentibus hucusque detectas secundum systema Linneanum dispositas*. 3 vols. Typ. Tramater, Neapoli. <https://doi.org/10.5962/bhl.title.50455>
- Huillard-Bréholles JLA (1860) *Historia diplomatica Friderici Secundi. Sive Constitutiones, Privilegia, Mandata, Instrumenta quae supersunt istius Imperatoris et Filiorum ejus; Accedunt Epistolae Papparum et Documenta varia*. Tomus VI. H. Plon, Parisiis.
- IGMI (Istituto Geografico Italiano) (1973) Foglio 628 'Sciacca' (scala 1:50.000). Istituto Geografico Militare, Firenze.
- ITMI (Istituto Topografico Italiano) (1872) *Carta della Sicilia* (scala 1:100.000), Tavola n. 150 'Castelvetrano'. Istituto Topografico Italiano, Firenze.
- La Mantia A, Gianguzzi L (2003) Considerations on protection and forestal restoring of the *Quercus calliprinos* vegetation in Sicily. *Bocconea* 16(2): 823–829. <https://www.herbmedit.org/bocconea/16-0823.pdf>
- La Mela Veca DS, Maetcke F, Pasta S, Rühl J (eds) (2007) *La Gestione Forestale Sostenibile nelle Aree Protette: il caso di studio della Riserva Naturale Orientata "Sugherete di Niscemi"* (CL). Dipartimento di Colture Arboree dell'Università degli Studi di Palermo - Azienda Foreste Demaniali della Regione Siciliana, Collana "Sicilia Foreste" n° 31.
- La Rosa A, Gianguzzi L, Ottonello D (2007) Primi dati sulla flora vascolare del SIC "Sistema dunale Capo Granitola, Porto Palo e Foce del Belice" (Sicilia sud-occidentale). 102° Congresso Nazionale della Società Botanica Italiana (Palermo, 26–29 settembre 2007), riassunti: 302.
- Leighton R (1999) *Sicily before History: An Archaeological Survey from the Palaeolithic to the Iron Age*. Cornell University Press, Ithaca (NY).
- Lentini F, Carbone S (2014) *Carta geologica della Sicilia* (1: 250.000). SEICa (= Società Elaborazioni Cartografiche), Firenze.
- Lo Vetro D, Martini F (2012) Il Paleolitico e il Mesolitico in Sicilia. In: R. Giglio, S. Tusa (eds), *Atti della XLI Riunione Scientifica "Dai Ciclopi agli Ecasti: Società e territorio nella Sicilia preistorica e protostorica"* (San Cipirello, PA, 16–19 novembre 2006): 19–47. Istituto Italiano di Preistoria e Protostoria, Firenze.
- Lojacono-Pojero M (1909) *Flora Sicula o descrizione delle piante spontanee o indigenate in Sicilia*. Vol. 3 (*Monocotyledones, Cryptogamae*). Palermo, 448 + xvi pp. + 20 plates.
- Lopriore G (1901) *Studi comparativi sulla flora lacustre della Sicilia*. Tip. Sicula di Monaco & Mollica, Catania, 116 pp.
- Magini GA (1620) *Regno di Sicilia*. Bologna. In: Bucchieri F. (ed), *Catalogo della mostra "La Sicilia di carta. Le carte della Sicilia. La rappresentazione cartografica della Sicilia dal Rinascimento all'età dei Lumi"*. Museo d'Arte e Archeologia "Ignazio Mormino", Fondazione del Banco di Sicilia, Palermo.
- Marguglio T (1973) Caratteristiche ecologiche e vegetazione naturale potenziale in provincia di Trapani. *Itinerari Trapanesi* 1(1): 9–15.
- Maurici F (1995) *La Sicilia di Federico II*. Città, castelli e casali. Accademia Nazionale di Scienze Lettere ed Arti, Palermo, 56 pp.
- Médail F, Pavon D, Lo Cascio P, Pasta S (2016) Sulla presenza di *Ophioglossum lusitanicum* L. sull'isola di Vulcano e aggiunte alla flora vascolare delle Isole Eolie (Sicilia nord-orientale). *Il Naturalista siciliano* (4) 40(1): 51–66. <https://www.academia.edu/38876117/Sulla-presenza-di-Ophioglossum-lusitanicum-nellisola-di-Vulcano-e-aggiunte-alla-flora-vascolare-delle-Isole-Eolie>
- Minissale P (1995) Studio fitosociologico delle praterie ad *Ampelodesmos mauritanicus* della Sicilia. *Colloques phytosociologiques* 21 (1993): 615–652.
- Mucina L, Bültmann H, Dierßen K, Theurillat J, Raus T, Čarni A, et al. (2016) *Vegetation of Europe: hierarchical floristic classification system of vascular plant, bryophyte, lichen, and algal communities*. *Applied Vegetation Sciences* 19: 3–264. <https://doi.org/10.1111/avsc.12257>

- Musarella CM, Mendoza-Fernández AJ, Mota JF, Alessandrini A, Bacchetta G, Brullo S, et al. (2018) Checklist of the gypsophilous vascular flora in Italy. *PhytoKeys* 103: 61–82. <https://doi.org/10.3897/phytokeys.103.25690>
- Napoli F (1932) Storia della città di Mazara. Stabilimento tipografico Hopps, Mazara del Vallo, 271 pp.
- Nicotra L, Campagna G (1908) Addenda ad *Floram Siculam nonnulla*. *Malpighia* 22: 3–14, 537–548.
- Nimis PL (1987) I macrolicheni d'Italia. Chiavi analitiche per la determinazione. *Gortania* 8: 101–220.
- Nimis PL, Martellos S (2017) ITALIC - The Information System on Italian Lichens. Version 5.0. University of Trieste, Dept. of Biology (<http://dryades.units.it/italic>).
- Noto GB (1732, unpubl.). Platea della Palmosa Città di Castelvetro, Suo stato, Giurisdizione, Baronie e Contea del Borgetto aggregati. Manuscript, Municipal Library of Castelvetro, 53 pp.
- Ottonello D, Catanzaro F (1986) Contributo alla flora del Trapanese. *Il Naturalista siciliano* (4) 9(1–4)(1985): 89–99.
- Ottonello D, La Mantia A, Calvaruso I (2003) Indagini preliminari finalizzate allo studio della vegetazione della Riserva Naturale Integrale Lago Preola e Gorgi Tondi (Sicilia sud-occidentale). 98° Congresso Nazionale della Società Botanica Italiana (Catania, 24–26 settembre 2003), riassunti: 300.
- Pasta S, Bambina A, Colonna Romano L, Giancontieri G, Messina G, La Mantia T, Ottonello D, Scuderi L (2008) Il sito di "Castello della Pietra" e "Riserva Zangara" (Castelvetro, Sicilia sud-occidentale: Indagine multidisciplinare e proposte di tutela. *Il Naturalista siciliano* (4) 32(1–2): 3–60. [https://www.sssn.it/PDF/PDF%20Nat.%20Sic.%201-2%202008/Pastaetalii\\_3-60.pdf](https://www.sssn.it/PDF/PDF%20Nat.%20Sic.%201-2%202008/Pastaetalii_3-60.pdf)
- Pasta S, Cullotta S, La Mantia T (2000) Ecogeografia e ruolo fisionomico-strutturale delle querce sempreverdi in Sicilia. Pp. 65–71 in: G. Bucci, G. Minotta, M. Borghetti (eds.), "Applicazioni e Prospettive per la Ricerca Forestale Italiana", Atti II Congr. S.I.S.E.F. (Bologna, 20–22 ottobre 1999).
- Pasta S, de Rigo D, Caudullo G (2016) *Quercus pubescens* in Europe: distribution, habitat, usage and threats. In: San-MiguelAyaz J, de Rigo D, Caudullo G, Houston Durrant T, Mauri A (eds), *European Atlas of Forest Tree Species*. Publ. Off. EU, Luxembourg, pp. e019e5c+
- Pignatti S, Guarino R, La Rosa M (2017–2019) *Flora d'Italia*, 2 ed. Media Business - Edagricole, Milano - Bologna, 4 vols.
- Ponzo A (1900) *La flora trapanese*. Tipografia Puccio, Palermo, 140 pp.
- Ponzo A. (1903) *La flora nei dintorni di Alcamo*. Nota I. *Bullettino della Società Botanica Italiana*: 200–212, 318–330.
- Pyšek P, Richardson DM, Rejmanek M, Webster GL, Williamson M, Kirschner J (2004) Alien plants in checklists and floras: towards better communication between taxonomists and ecologists. *Taxon* 53(1): 131–143. <https://doi.org/10.2307/4135498>
- Raimondo FM (ed), Fici S, Gianguzzi L, Lentini F, Mazzola P, Miceli G, Not R, Ottonello D, Romano S, Schicchi R (coll.) (1986) *Atlante iconografico delle piante endemiche o rare della Riserva naturale orientata dello Zingaro (Sicilia)*. Dipartimento di Scienze Botaniche, Azienda Freste Demaniali della Regione Siciliana, STASS, Palermo, pp. 84, 37 figures, 1 map.
- Raimondo FM (1979) Scheda XIX. Sughereta di Contrada Bosco Scorce. Pp. 497–498 in: Gruppo di Lavoro per la Conservazione della Natura della Società Botanica Italiana (a cura di), "Censimento dei biotopi di rilevante interesse vegetazionale meritevoli di conservazione in Italia", Vol 2. Savini-Mercuri, Camerino.
- Raimondo FM (1987) Aspetti del patrimonio vegetale spontaneo nella campagna di Buseto. In: *La flora spontanea del Busetano: un patrimonio da salvaguardare*. Quaderni della Scuola Media "A. Manzoni", Buseto Palizzolo, 6, Tipografia Abate, Paceco, 12–26.
- Raimondo FM, Bazan G, Troia A (2011) Taxa a rischio nella flora vascolare della Sicilia. *Biogeografia* (2) 30: 229–239. <https://doi.org/10.21426/B630110586>
- Raimondo FM, Castiglia G, Schicchi R (1991) La macchia insediata sulle rovine dell'antica città di Selinunte (Trapani). *Giornale botanico italiano* 12 5(3): 413 (abstract).
- Raimondo FM, Domina G, Bazan G (2005) Carta dello stato delle conoscenze floristiche della Sicilia. In: Scoppola A, Blasi C (eds), *Stato delle conoscenze sulla flora vascolare d'Italia*. Palombi Editore, Roma, pp. 203–206 + tav. XXVIII.
- Raimondo FM, Mazzola P, Domina G (2004) Check-list of the vascular plants collected during Iter Mediterraneum III. *Bocconea* 17: 65–232.
- Raimondo FM, Schicchi R (eds) (2000) *Il popolamento vegetale della Riserva Naturale dello Zingaro*. Azienda Foreste Demaniali della Regione Siciliana, Collana "Sicilia Foreste" 3 (suppl.) (1998), Palermo, 205 + i pp.
- Ravera S, Azara C, Baragatti E, Paoli L, Genovesi V, Tretiach M (2011) *Notulae Cryptogamicae* 3 (Lichenes), Notula 11: *Teloschistes chrysophthalmis* (L.) Th. Fr. (*Teloschistaceae*). *Informatore botanico italiano* 43(1): 152–153.
- Regione Siciliana (2018) Piano Paesaggistico. Ambiti Regionali 2 e 3 ricadenti in Provincia di Trapani - Carte di Analisi Sistema Antropico - Sistema Storico-Culturale. Tavola 8.7 (scala 1:25.000). Assessorato dei Beni Culturali e dell'Identità Siciliana, Dipartimento dei Beni Culturali e dell'Identità Siciliana, Soprintendenza per i Beni Culturali ed Ambientali di Trapani, Trapani. [http://www.regione.sicilia.it/beniculturali/dirbenicult/bca/ptpr/documentazione%20tecnica%20trapani/CARTOGRAFIA%201/Analisi%201/8\\_7\\_beni%20storici%20\(1\).pdf](http://www.regione.sicilia.it/beniculturali/dirbenicult/bca/ptpr/documentazione%20tecnica%20trapani/CARTOGRAFIA%201/Analisi%201/8_7_beni%20storici%20(1).pdf)
- Rivas-Martínez S (2005) Notions on dynamic-catenal phytosociology as a basis of landscape science. *Plant Biosystems* 139: 135–144. <https://doi.org/10.1080/11263500500193790>
- Rivas-Martínez S, Peñas A, Díaz TE (2004) Bioclimatic and biogeographic maps of Europe. [www.globalclimatics.org/form/maps.htm](http://www.globalclimatics.org/form/maps.htm) (last accessed: May 28, 2020)
- Rodrigues ASL, Andelman SJ, Bakarr MI, Boitani L, Brooks TM, Cowling RM, et al. (2004) Effectiveness of the global protected area network in representing species diversity. *Nature* 428(6983): 640–643. <https://doi.org/10.1038/nature02422>
- Romano S, Tobia G, Gianguzzi L (2006) Rassegna della flora vascolare dell'Isola di Levanzo (Arcipelago delle Egadi, Canale di Sicilia). *Informatore botanico italiano* 38 (2): 481–502.
- Ronsisvalle GA (1979) Vegetazione psammofila del litorale tra Gela e Mazara del Vallo (Sicilia Meridionale). *Bollettino dell'Accademia Gioenia di Scienze naturali* (4) 13(10): 9–25.
- Rosati L, Marignani M, Blasi C (2007) Vegetazione Naturale Potenziale e Gap analysis della Rete Natura 2000 in Italia. *Fitosociologia* 44(2, Suppl. 1): 61–65.
- Rosati L, Marignani M, Blasi C (2008) A gap analysis comparing Natura 2000 vs. National Protected Area network with potential natural vegetation. *Community Ecology* 9(2): 147–154. <https://doi.org/10.1556/ComEc.9.2008.2.3>
- Rühl J, Chiavetta U, La Mantia T, La Mela Veca DS, Pasta S (2005) Land cover change in the Nature Reserve "Sughereta di Niscemi" (SE Sicily) in the 20th century. In: Erasmí S., Cyffka B., Kappas M. (eds.),

- Remote Sensing & GIS for Environmental Studies: Applications in Geography, Proceedings of the 1st GGRS (Göttingen GIS & Remote Sensing Days), Environmental Studies (Göttingen, Germany, 7–8 October 2004), Göttinger Geographische Abhandlungen 113: 54–62.
- Sciascia L (2002) Nobili navarresi nella Sicilia di Federico III: Asiain, Simen de Aibar, Olleta e Caparroso. *Principe de Viana* 63: 157–166.
- Scott JM, Davis F, Csuti B, Noss RF, Butterfield B, Groves C, Anderson H, Caicco S, D'Erchia F, Edwards TC, Uliman J, Wright RG (1993) Gap analysis: a geographical approach to protection of biological diversity. *Wildlife Monographs* 123: 1–41. <https://www.jstor.org/stable/3830788>
- Scuderi G, Ilardi V, Raimondo FM (1994) La sughera nella vegetazione arborea del Trapanese. *Quaderni di Botanica ambientale e applicata* 3 (1992): 223–233. [https://www.quadernibotanicambientaleappl.it/quaderni/03\\_223-233.pdf](https://www.quadernibotanicambientaleappl.it/quaderni/03_223-233.pdf)
- Scuderi L (2006) Flora e vegetazione della provincia di Trapani (Sicilia). Tesi di Dottorato in Scienze Ambientali I - Fitogeografia dei Territori Mediterranei (XIX Ciclo), Università degli Studi di Catania, Catania, 541 pp.
- Scuderi L, Pasta S (2009) Contributi alla conoscenza della flora vascolare della provincia di Trapani (Sicilia occidentale). I. Taxa autoctoni inediti. *Il Naturalista siciliano* (4) 33 (3–4): 97–112. <http://www.sssn.it/PDF/PDF%20NS%2033/329-344.pdf>
- Serasiaux E (1989) Liste rouge des Macrolichens dans la Communauté Européenne. Centre de Recherches sur les Lichens. Sart-Tilman, Liège, 238 pp.
- Simonsohn Š (2007) The Jews in Sicily. Vol. 10: Notaries from Palermo. Part One. Brill, Leiden - Boston. <https://doi.org/10.1163/ej.9789004157620.i-6808>
- Sommier S (1907) Materiali per una florula di Pantelleria. *Bullettino della Società Botanica Italiana* 1907 14: 48–60.
- Tinner W, van Leeuwen JFN, Colombaroli D, Vescovi E, van der Knaap WO, Henne PD, Pasta S, D'Angelo S, La Mantia T (2009) Holocene environmental and climatic changes at Gorgo Basso, a coastal lake in southern Sicily, Italy. *Quaternary Science Review* 28 (15–16): 1498–1510. <https://doi.org/10.1016/j.quascirev.2009.02.001>
- Titone V (1961) Le origini della questione meridionale. I. Riveli e platee del Regno di Sicilia. Feltrinelli, Milano, 298 pp.
- Troia A (2006) Segnalazione di *Zizyphus lotus* (L.) Lam. (*Rhamnaceae*) nei pressi di Mazara del Vallo (Trapani, Sicilia). *Il Naturalista siciliano* (4) 30 (3–4): 393–400. [http://www.sssn.it/PDF/PDF%20Nat.%20Sic.%203-4%202006/Troia\\_393-400.pdf](http://www.sssn.it/PDF/PDF%20Nat.%20Sic.%203-4%202006/Troia_393-400.pdf)
- Troia A, Adragna F, Campisi P, Campo G, Dia MG, Ilardi V, La Mantia T, La Rosa A, Lo Valvo M, Muscarella C, Pasta S, Pieri V, Scuderi L, Sparacio I, Stoch F, Marrone F (2016). I pantani di Anguillara (Comune di Calatafimi-Segesta, provincia di Trapani): dati preliminari sulla biodiversità a supporto della tutela del biotopo. *Il Naturalista siciliano* (4) 40(2): 171–200. <https://www.sssn.it/PDF/PDF%20NS%2040/171-200.pdf>
- Troia A, Bazan G, Schicchi R (2011) Nuove aree di rilevante interesse naturalistico nella Sicilia centro-occidentale: proposta di tutela. *Il Naturalista siciliano* (4) 35(2): 257–293. <https://www.sssn.it/PDF/PDF%20NS%2035/257-294.pdf>
- Troia A, Lansdown R (2016) The first confirmed population of the globally endangered *Pilularia minuta* (*Marsileaceae*) in Sicily. *Webbia* 71 (2): 283–286. <https://doi.org/10.1080/00837792.2016.1195105>
- Troia A, Napolitano T (2017) Segnalazioni floristiche e vegetazionali per le zone umide costiere del territorio di Petrosino (Sicilia occidentale). *Il Naturalista siciliano* (4) 41(1): 25–34. <http://www.sssn.it/PDF/PDF%20NS%2041/25-34.pdf>
- Troia A, Raimondo FM (2010) *Isoëtes todaroana* (*Isoëtaceae*, *Lycopodiophyta*), a new species from Sicily (Italy). *American Fern Journal* 99(4) (2009): 238–243. <https://doi.org/10.1640/0002-8444-99.4.238>
- Troia A, Spallino RE (2009) Conferma della presenza nella Sicilia occidentale di *Retama raetam* (Forssk.) Webb subsp. *gussonei* (Webb) W. Greuter (*Fabaceae Cytiseae*), specie a rischio della flora italiana. *Il Naturalista siciliano* (4) 33 (3–4): 305–314. <https://www.sssn.it/PDF/PDF%20NS%2033/305-314.pdf>
- UK BAP (UK Biodiversity Group: Tranche 2 Action Plans), 1999. Volume III: Plants and Fungi. Peterborough: English Nature, 104–106.
- Vessella F, Schirone B (2013) Predicting potential distribution of *Quercus suber* in Italy based on ecological niche models: Conservation insights and reforestation involvements. *Forest Ecology and Management* 304: 150–161. <https://doi.org/10.1016/j.foreco.2013.05.006>
- Viciani D, Lastrucci L, Geri F, Foggi B (2016) Gap analysis comparing protected areas with potential natural vegetation in Tuscany (Italy) and a GIS procedure to bridge the gaps. *Plant Biosystems* 150(1): 62–72. <https://doi.org/10.1080/11263504.2014.950623>
- Vimal R, Rodrigues ASL, Mathevet R, Thompson JD (2011) The sensitivity of gap analysis to conservation targets. *Biodiversity Conservation* 20: 531–543. <https://doi.org/10.1007/s10531-010-9963-1>
- Wellstein C, Spada F (2015) The status of *Quercus pubescens* Willd. in Europe. In: Box EO, Fujiwara K (eds.) *Warm-Temperate Deciduous Forests around the Northern Hemisphere*, Springer, Cham, pp. 153–163. [https://doi.org/10.1007/978-3-319-01261-2\\_8](https://doi.org/10.1007/978-3-319-01261-2_8)

## Supplementary material 1

### Table S1

Authors: Alfonso La Rosa, Lorenzo Gianguzzi, Giuseppe Salluzzo, Leonardo Scuderi, Salvatore Pasta

Data type: table

Explanation note: Checklist of the Vascular plants observed growing in Parche di Bilello (south-western Sicily).

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/PlantSociology.58.62437.suppl1>