



UNIVERSITÀ
DEGLI STUDI
DI PADOVA

Università degli Studi di Padova

Padua Research Archive - Institutional Repository

Convergence of nonlocal geometric flows to anisotropic mean curvature motion

Original Citation:

Availability:

This version is available at: 11577/3395381 since: 2021-07-09T10:31:18Z

Publisher:

American Institute of Mathematical Sciences

Published version:

DOI: 10.3934/dcds.2021065

Terms of use:

Open Access

This article is made available under terms and conditions applicable to Open Access Guidelines, as described at <http://www.unipd.it/download/file/fid/55401> (Italian only)

(Article begins on next page)

CONVERGENCE OF NONLOCAL GEOMETRIC FLOWS TO ANISOTROPIC MEAN CURVATURE MOTION

ANNALISA CESARONI *

Department of Statistical Sciences, Università di Padova
Via Battisti 241/243
35121 Padova, Italy

VALERIO PAGLIARI

Institute of Analysis and Scientific Computing, TU Wien
Wiedner Hauptstraße 8 - 10
1040 Vienna, Austria

(Communicated by the associate editor name)

ABSTRACT. We consider nonlocal curvature functionals associated with positive interaction kernels, and we show that local anisotropic mean curvature functionals can be retrieved in a blow-up limit from them. As a consequence, we prove that the viscosity solutions to the rescaled nonlocal geometric flows locally uniformly converge to the viscosity solution to the anisotropic mean curvature motion. The result is achieved by combining a compactness argument and a set-theoretic approach related to the theory of De Giorgi's barriers for evolution equations.

1. Introduction. In this paper we prove convergence of a class of rescaled nonlocal curvature flows to local anisotropic mean curvature evolutions.

We fix an interaction kernel $K: \mathbb{R}^d \setminus \{0\} \rightarrow [0, +\infty)$, possibly singular at 0, modeling interactions between points in the space, and we define the *nonlocal curvature* associated with K of a measurable set $E \subseteq \mathbb{R}^d$ at $x \in \partial E$ as

$$H_K(E, x) := - \lim_{r \rightarrow 0^+} \int_{B(x, r)^c} K(y - x) \tilde{\chi}_E(y) dy. \quad (1)$$

Here and in the sequel, $B(x, r)$ is the open ball with center x and radius r , $E^c = \mathbb{R}^d \setminus E$ for any $E \subseteq \mathbb{R}^d$, and $\tilde{\chi}_E(x)$ is equal to 1 when $x \in E$ and it is equal to -1 otherwise.

Note that if $K \in L^1(\mathbb{R}^d)$, then the nonlocal curvature coincides with $H_K(E, x) = -(K * \tilde{\chi}_E)(x)$. More generally, we will impose conditions on K so that $C^{1,1}$ sets have bounded nonlocal curvature, see Section 2.

By using the nonlocal curvature operator, we define a nonlocal flow as follows: for a family of evolving sets $\{E(t)\}_{t \geq 0}$, we prescribe the geometric law

$$\partial_t x(t) \cdot \hat{n} = -H_K(E(t), x), \quad (2)$$

2020 *Mathematics Subject Classification.* Primary: 53E10; Secondary: 35D40, 35K93, 35R11.
Key words and phrases. Nonlocal curvature flow, anisotropic mean curvature flow, geometric equations, De Giorgi's barriers for geometric evolutions, level-set method, viscosity solutions.

The authors are members and were supported by the INDAM/GNAMPA.

* Corresponding author: Annalisa Cesaroni.

where \hat{n} is the outer unit normal to $\partial E(t)$ at the point $x(t)$.

Geometric nonlocal evolutions as (2) emerged as models for dislocations dynamics in the description of plastic behavior of metallic crystals. Dislocations are linear misalignments in the microscopic crystalline lattice, and whose normal velocity is determined by the so called Peach-Koehler force. In [3], Alvarez, Hoch, Le Bouar, and Monneau proposed a mathematical description of dislocation dynamics in terms of a nonlocal eikonal equation, where the Peach-Koehler force is encoded by a convolution kernel c_0 . The explicit expression of the kernel might be complicated, because it has to capture the physical features of the system, e.g. in general it can change sign. By then, their model has been simplified in a series of papers, in which well-posedness of the geometric evolution law was obtained, see [2, 7, 30, 27, 24].

Another interesting aspect of the nonlocal curvature (1) is that it is the first variation of the nonlocal perimeter functional

$$\text{Per}_K(E) := \int_E \int_{E^c} K(y-x) dy dx$$

(see e.g. [19]), and the geometric evolution law (2) is then understood as the L^2 gradient flow of this kind of perimeter.

When K belongs to an appropriate class of fractional kernels, existence and uniqueness of solutions in the viscosity sense to the geometric flow (2) were investigated in [29]. More recently, Chambolle, Morini, and Ponsiglione have proved in [19] well posedness of the level-set formulation of a wide class of local and nonlocal translation-invariant geometric flows. They also have exploited the minimizing movement scheme to construct solutions to flows driven by variational curvatures.

The analysis of nonlocal curvature flows as (2) has lately been carried out from various perspectives, especially in fractional case; for instance, conservation of convexity, formation of neckpinch singularities, and fattening phenomena have been considered, see [17, 23, 21].

As we anticipated, we are interested in the asymptotic behaviour of a family of nonlocal curvature flows, obtained by rescaling the kernel K . Explicitly, for any $\varepsilon > 0$ and $x \in \mathbb{R}^d$, we put

$$K_\varepsilon(x) := \frac{1}{\varepsilon^d} K\left(\frac{x}{\varepsilon}\right) \quad (3)$$

and, for a measurable set $E \subset \mathbb{R}^d$ and $x \in \partial E$, we define

$$H_\varepsilon(E, x) := \frac{1}{\varepsilon} H_{K_\varepsilon}(E, x). \quad (4)$$

We remark that this scaling is mass preserving, in the sense that, at least formally, $\|K\|_{L^1(\mathbb{R}^d)} = \|K_\varepsilon\|_{L^1(\mathbb{R}^d)}$. At the same time, we expect a localization effect in the limit.

Our main assumptions on the kernel K are listed in Section 2. In particular, we will require that K is sufficiently regular and has at most a singularity in the origin, that is $K \in W^{1,1}(\mathbb{R}^d \setminus B(0, r))$ for all $r > 0$. In addition, we assume that there exist $m > 0$ and $s \in (0, 1)$ such that

$$0 \leq K(x) \leq \frac{m}{|x|^{d+1+s}} \quad \text{if } x \in B(0, 1)^c,$$

and that for all $\lambda > 0$ and all $e \in \mathbb{S}^{d-1} := \partial B(0, 1)$ there holds

$$K, |x| |\nabla K(x)| \in L^1 \left(\left\{ x \in \mathbb{R}^d : |x \cdot e| \leq \frac{\lambda}{2} |\pi_{e^\perp}(x)|^2 \right\} \right),$$

where e^\perp is the hyperplane of vectors that are orthogonal to e , and π_{e^\perp} is the orthogonal projection operator on e^\perp . Actually, in order to exploit these properties in our proofs, we will need to make them quantitative. We refer the reader to Section 2 for a detailed presentation of the assumptions.

We point out that in [24] a similar problem was studied, but there the assumptions on the interaction kernel, and thus the choice of the rescaling, are different from ours. Indeed, the authors of [24] assume the kernel K to be bounded near the origin (hence nonsingular) and to decay as $|x|^{-(d+1)}$ at infinity. The rescaled curvature is defined as

$$\frac{1}{\varepsilon \log \varepsilon} H_{K_\varepsilon}(E, x),$$

and the authors prove that, as $\varepsilon \rightarrow 0$, it converges to an anisotropic, local curvature functional. They also show that the rescaled geometric motion approaches the flow driven by the limiting curvature.

In the last years, other results related to the asymptotic behavior of rescaled non-local functionals have appeared in the literature, mainly in the stationary setting. For radial, nonsingular kernels, it is proved in [33] that the rescaled perimeters $\varepsilon^{-1} \text{Per}_{K_\varepsilon}(E)$ converge pointwise to the local perimeter functional. In the same paper, pointwise convergence of the rescaled curvature to the local mean curvature is obtained as well. An improvement concerning the convergence of perimeters has recently been obtained in [13, 34], where Γ -convergence of the functionals $\varepsilon^{-1} \text{Per}_{K_\varepsilon}(E)$ to De Giorgi's perimeter is established for a class of singular kernels. Results in the same spirit addressing specifically the fractional case can be found in [5, 14, 16], see also [35] for Γ -convergence of nonlocal phase transitions. Finally, we recall the recent preprint [18], where stability results for nonlocal geometric evolutions are studied by using viscosity solutions arguments. In the present paper, we propose a different, more geometric, approach to the problem, as we will detail in the following.

Our first main result is the uniform convergence of the rescaled curvature functionals to a local, anisotropic mean curvature functional, when they are computed for smooth, compact sets. We fix some notations needed to formulate the precise statement.

As before, p^\perp is the hyperplane of the vectors that are orthogonal to p , and π_{p^\perp} is the orthogonal projection operator on p^\perp . We denote by $\text{Sym}(d)$ the space of $d \times d$ real symmetric matrices and by \mathcal{H}^{d-1} the $(d-1)$ -dimensional Hausdorff measure. For a C^2 hypersurface in \mathbb{R}^d Σ , we define the anisotropic mean curvature functional

$$H_0(\Sigma, x) := -\frac{1}{|\nabla \phi(x)|} \text{tr} \left(M_K(\hat{n}) \pi_{\hat{n}^\perp} \nabla^2 \phi(x) \pi_{\hat{n}^\perp} \right), \quad (5)$$

where $\phi \in C^2(\mathbb{R}^d)$ is a function such that $\Sigma \cap U = \{y \in \mathbb{R}^d : \phi(y) = 0\} \cap U$ in some open neighbourhood U of x , $\nabla \phi(x) \neq 0$, \hat{n} is the outer unit normal to Σ at x , and finally

$$\begin{aligned} M_K : \mathbb{S}^{d-1} &\longrightarrow \text{Sym}(d) \\ e &\longmapsto \int_{e^\perp} K(z) z \otimes z d\mathcal{H}^{d-1}(z). \end{aligned} \quad (6)$$

Then, we show the following:

Theorem 1.1. *Let K satisfy all the assumptions in Section 2. Let $E \subset \mathbb{R}^d$ be a set whose boundary Σ is compact and of class C^2 . Then,*

$$\lim_{\varepsilon \rightarrow 0^+} H_\varepsilon(E, x) = H_0(\Sigma, x) \quad \text{uniformly in } x \in \Sigma.$$

We recall that analog results to ours for nonsingular kernels are found in [33] and in [24], respectively for the isotropic and the anisotropic case.

Our second main result deals with the convergence of the rescaled nonlocal geometric flows

$$\partial_t x(t) \cdot \hat{n} = -H_\varepsilon(E(t), x(t)) \quad (7)$$

to the anisotropic mean curvature flow

$$\partial_t x(t) \cdot \hat{n} = -H_0(\Sigma(t), x(t)), \quad (8)$$

where $\Sigma(t) := \partial E(t)$. We develop our analysis in the framework of the level-set method. This amounts to defining the evolving set $E(t)$ and its boundary $\Sigma(t)$ as the 0 superlevel set and 0 level set of some function $\phi(t, \cdot)$, which turns out to be a viscosity solution of the nonlocal parabolic partial differential equation

$$\partial_t \phi(t, x) + |\nabla \phi(t, x)| H_\varepsilon(\{y : \phi(t, y) \geq \phi(t, x)\}, x) = 0 \quad (9)$$

if $E(t)$ solves the rescaled nonlocal geometric flow (7), or of the local parabolic partial differential equation

$$\partial_t \phi(t, x) + |\nabla \phi(t, x)| H_0(\{y : \phi(t, y) = \phi(t, x)\}, x) = 0 \quad (10)$$

if $\Sigma(t)$ solve the anisotropic mean curvature flow (8). We can state our second major result.

Theorem 1.2. *Let K satisfy all the assumptions in Section 2. Let $u_0: \mathbb{R}^d \rightarrow \mathbb{R}$ be a Lipschitz continuous function that is constant outside a compact set. Let $u_\varepsilon, u: [0, +\infty) \times \mathbb{R}^d \rightarrow \mathbb{R}$ be respectively the unique continuous viscosity solution to (9) and (10), with initial datum u_0 . Then*

$$\lim_{\varepsilon \rightarrow 0} u_\varepsilon(t, x) = u(t, x) \quad \text{locally uniformly in } [0, +\infty) \times \mathbb{R}^d.$$

The proof of Theorem 1.2 is based on the convergence of curvatures obtained in Theorem 1.1. We propose a proof based on the concept of geometric barrier, introduced by De Giorgi in [25] as a weak solution to a wide range of evolution problems. The study of barriers in relation to geometric parabolic PDEs, such as (10), was developed by Bellettini, Novaga, and Paolini in the late 90's [11, 8, 10, 9]. It turns out that, for the class of problems under consideration, viscosity theory and barriers can be compared, and this is the key point that we will exploit in our analysis.

We remark that isotropic fractional kernels such as $K(y-x) = |y-x|^{-d-s}$ for $s \in (0, 1)$ are not directly included in the class of kernels we are considering, see Example 1. Nevertheless the same kind of result as Theorem 1.1 for the fractional mean curvature as $s \rightarrow 1$ was obtained in [1, 16, 18], whereas the convergence of the level set flow has been proved in [18] by using viscosity solution methods.

Finally, we recall that there is a large literature concerning approximation results for mean curvature motions, either with local or nonlocal operators. One of the most renowned algorithmeorems is the threshold dynamics type one introduced in [12] by Bence, Merriman, and Osher. This approach was rigorously settled in [6] and [26]; then, the analysis was extended to more general diffusion operators in [31], [32], and [20] (for anisotropic and crystalline evolutions). In [15] Caffarelli and

Souganidis established the convergence of an analogous threshold dynamics scheme to the (isotropic) motion by fractional mean curvature, and this result was adapted to the anisotropic case, also in presence of a driving force, in [21].

Structure of the paper. In Section 2 we describe the class of interaction kernels that we consider in this work. In Section 3 and 4 we discuss some basic properties of the curvatures functionals, and we recall the level-set formulation for geometric flows, the notion of geometric barriers, and the main results about them. Section 5 is devoted to the proof of Theorem 1.1. In Section 6, we provide a compactness result for the family of solutions to the rescaled nonlocal problems. Eventually, Section 7 contains the proof of Theorem 1.2.

Acknowledgement. The authors warmly thank Matteo Novaga for inspiring discussions on this problem.

2. Standing assumptions on the kernel. Throughout this work, $K: \mathbb{R}^d \setminus \{0\} \rightarrow [0, +\infty)$ is a measurable function such that

$$K(y) = K(-y) \quad \text{for all } y \in \mathbb{R}^d \setminus \{0\} \quad (11)$$

and

$$K \in W^{1,1}(B(0,r)^c) \quad \text{for all } r > 0. \quad (12)$$

Note that (12) allows both K and ∇K to be singular around the origin, and it implies convergence of their integrals at infinity; however, we need to make these information quantitative.

Firstly, we require that

$$\lim_{r \rightarrow 0^+} r \int_{B(0,r)^c} K(y) dy = 0. \quad (13)$$

Then, for any $e \in \mathbb{S}^{d-1}$ and $\lambda > 0$, we set

$$Q_\lambda(e) := \left\{ y \in \mathbb{R}^d : |y \cdot e| \leq \frac{\lambda}{2} |\pi_{e^\perp}(y)|^2 \right\},$$

and we assume that

$$y \mapsto K(y), y \mapsto |y| |\nabla K(y)| \in L^1(Q_\lambda(e)) \quad \text{for all } e \in \mathbb{S}^{d-1} \text{ and } \lambda > 0. \quad (14)$$

This will imply that sets with $C^{1,1}$ compact boundary have finite curvature, see Proposition 1. We stress that we make no isotropy hypothesis on K ; still, we have to suppose some control on the mass of K in $Q_\lambda(e)$, uniformly in e . We therefore suppose that for all $\lambda > 0$ there exists $a_\lambda > 0$ such that for all $e \in \mathbb{S}^{d-1}$

$$\int_{Q_\lambda(e)} K(y) dy \leq a_\lambda. \quad (15)$$

In addition, we require that there exist $a_0, b_0 > 0$ such that for all $e \in \mathbb{S}^{d-1}$

$$\limsup_{\lambda \rightarrow 0^+} \frac{1}{\lambda} \int_{Q_\lambda(e)} K(y) dy \leq a_0, \quad (16)$$

$$\limsup_{\lambda \rightarrow 0^+} \frac{1}{\lambda} \int_{Q_\lambda(e)} |\nabla K(y)| |y| dy \leq b_0. \quad (17)$$

We assume as well that for all $e \in \mathbb{S}^{d-1}$

$$\lim_{\lambda \rightarrow +\infty} \frac{1}{\lambda} \int_{Q_\lambda(e)} K(y) dy = 0. \quad (18)$$

Finally, we suppose that, far from the origin, K is bounded above by a fractional kernel; that is, there exist $m > 0$ and $s \in (0, 1)$ such that

$$K(y) \leq \frac{m}{|y|^{d+1+s}} \quad \text{if } y \in B(0, 1)^c. \quad (19)$$

Remark 1. Inequality (19) entails that for all $\alpha < s$

$$\lim_{r \rightarrow +\infty} r^{1+\alpha} \int_{B(0,r)^c} K(y) dy = 0. \quad (20)$$

Actually, most of the results in the paper are not affected if the weaker assumption (20) replaces (19). However, for the sake of simplicity, we decided not to pursue this direction.

As a concluding comment about our assumptions on K , we describe a class of singular kernels that fits in our analysis.

Example 1 (Fractional kernels). Let us suppose that $K: \mathbb{R}^d \setminus \{0\} \rightarrow [0, +\infty)$ satisfies (11) and that there exist constants $m, \mu > 0$ and $s, \sigma \in (0, 1)$ such that

$$K(y), |y| |\nabla K(y)| \leq \frac{\mu}{|y|^{d+\sigma}} \quad \text{for all } y \in B(0, 1)$$

and

$$K(y), |y| |\nabla K(y)| \leq \frac{m}{|y|^{d+1+s}} \quad \text{for all } y \in B(0, 1)^c.$$

Then, all the assumptions above are satisfied.

Also fractional kernels with exponential decay at infinity fit in our framework; namely, these are the kernels $K: \mathbb{R}^d \setminus \{0\} \rightarrow [0, +\infty)$ that satisfy (11) and for which there exist constants $m, \mu > 0$ and $s \in (0, 1)$ such that

$$K(y), |y| |\nabla K(y)| \leq \frac{\mu e^{-m|y|}}{|y|^{d+s}}, \quad \forall y \in \mathbb{R}^d.$$

3. Preliminaries about curvature functionals. In this section we discuss some basic results about the local and nonlocal curvature functionals H_0 and H_K defined in (5) and (1).

First of all, we show that the nonlocal curvature is finite on sets with $C^{1,1}$ boundaries. Similar results are already available in [29] and [19]. Nonetheless, we detail the argument for the sake of completeness, and to recover estimate (22), which will come in handy later. We will use the following notation: for $e \in \mathbb{S}^{d-1}$, $x \in \mathbb{R}^d$ and $\delta > 0$, we denote the cylinder of center x and axis e as

$$C_e(x, \delta) := \{y \in \mathbb{R}^d : y = x + z + te, \text{ with } z \in e^\perp \cap B(0, \delta), t \in (-\delta, \delta)\}. \quad (21)$$

Proposition 1. *Let $E \subset \mathbb{R}^d$ be an open set such that ∂E is a $C^{1,1}$ -hypersurface. Then, for all $x \in \partial E$ there exist $\bar{\delta}, \lambda > 0$ such that*

$$|H_K(E, x)| \leq \int_{Q_{\lambda, \bar{\delta}}(\hat{n})} K(y) dy + \int_{B(0, \bar{\delta})^c} K(y) dy, \quad (22)$$

where $Q_{\lambda, \bar{\delta}}(\hat{n}) := \{y \in Q_\lambda(\hat{n}) : |\pi_{\hat{n}^\perp}(y)| < \bar{\delta}\}$. In particular, $H_K(E, x)$ is finite.

Proof. Let $\Sigma := \partial E$ and \hat{n} be the outer unit normal to Σ at x . By the regularity of Σ , there exist $\bar{\delta} := \bar{\delta}(x)$ and a function $f: \hat{n}^\perp \cap B(0, \bar{\delta}) \rightarrow (-\bar{\delta}, \bar{\delta})$ of class $C^{1,1}$ such that

$$\Sigma \cap C_{\hat{n}}(x, \bar{\delta}) = \{y = x + z - f(z)\hat{n} : z \in \hat{n}^\perp \cap B(0, \bar{\delta})\}, \quad (23)$$

$$E \cap C_{\hat{n}}(x, \bar{\delta}) = \{y = x + z - t\hat{n} : z \in \hat{n}^\perp \cap B(0, \bar{\delta}), t \in (f(z), \bar{\delta})\}, \quad (24)$$

$$|f(z)| \leq \frac{\lambda}{2} |z|^2 \quad \text{for some } \lambda > 0. \quad (25)$$

It is not restrictive to assume $r < \bar{\delta}$; hence, we can split the integral in (1) into the sum

$$\int_C K(y-x) \tilde{\chi}_E(y) \chi_{B(x,r)^c}(y) dy + \int_{C^c} K(y-x) \tilde{\chi}_E(y) dy,$$

where we set $C := C_{\hat{n}}(x, \bar{\delta})$. The second term above is finite as a consequence of (12); indeed, since $B(x, \bar{\delta}) \subset C$, we have that

$$\left| \int_{C^c} K(y-x) \tilde{\chi}_E(y) dy \right| \leq \int_{B(0, \bar{\delta})^c} K(y) dy. \quad (26)$$

So, we are left to show that the integral

$$I_r := \int_C K(y-x) \tilde{\chi}_E(y) \chi_{B(x,r)^c}(y) dy$$

is bounded by a constant that does not depend on r . Taking into account (24) and recalling that K belongs to $L^1(B(0, r)^c)$ for any $r > 0$, we can write

$$I_r = \int_{\hat{n}^\perp \cap B(0, \bar{\delta})} \left[\int_{f(z)}^{\bar{\delta}} K(z - t\hat{n}) b_r(z, t) dt - \int_{-\bar{\delta}}^{f(z)} K(z - t\hat{n}) b_r(z, t) dt \right] d\mathcal{H}^{d-1}(z),$$

where, for $(z, t) \in [\hat{n}^\perp \cap B(0, \bar{\delta})] \times (-\bar{\delta}, \bar{\delta})$,

$$b_r(z, t) := \begin{cases} 0 & \text{if } |z| < r \text{ and } |t| < \sqrt{r^2 - |z|^2} \\ 1 & \text{otherwise} \end{cases}. \quad (27)$$

Since K is even, we get

$$\begin{aligned} I_r &= \int_{\hat{n}^\perp \cap B(0, \bar{\delta})} \left[\int_{f(z)}^{\bar{\delta}} K(z - t\hat{n}) b_r(z, t) dt - \int_{-f(-z)}^{\bar{\delta}} K(z - t\hat{n}) b_r(z, t) dt \right] d\mathcal{H}^{d-1}(z) \\ &= - \int_{\hat{n}^\perp \cap B(0, \bar{\delta})} \int_{-f(-z)}^{f(z)} K(z - t\hat{n}) b_r(z, t) dt d\mathcal{H}^{d-1}(z) \end{aligned}$$

In view of (25) we infer

$$\begin{aligned} |I_r| &\leq \int_{\hat{n}^\perp \cap B(0, \bar{\delta})} \int_{-\frac{\lambda}{2}|z|^2}^{\frac{\lambda}{2}|z|^2} K(z - t\hat{n}) b_r(z, t) dt d\mathcal{H}^{d-1}(z) \\ &= \int_{Q_{\lambda, \bar{\delta}}(\hat{n})} K(y) \chi_{B(0, r)^c}(y) dy. \end{aligned}$$

Assumption (14) allows to take the limit in the last inequality, and we conclude that (22) holds. \square

Remark 2. We point out that (22) has been obtained just exploiting the facts that K is even, $K \in L^1(B(0, r)^c)$ for all $r > 0$, and that $K \in L^1(Q_\lambda(e))$ for all $e \in \mathbb{S}^{d-1}$ and $\lambda > 0$.

We next observe that in (22) the second integral takes into account the “tails” of the kernel K , while the first one is related to the second fundamental form of Σ . We will prove in the sequel that, under our standing assumptions, the second term is negligible in the large scale limit.

The next lemma collects two fundamental properties of H_K . We omit the proofs, which can be derived easily from the definition of H_K .

Lemma 3.1. *Let $E \subset \mathbb{R}^d$ be an open set such that $H_K(E, x)$ is finite for some $x \in \partial E$.*

1. *For any $h \in \mathbb{R}^d$ and any orthogonal matrix R , if $T(y) := Ry + h$, then*

$$H_K(E, x) = H_{\tilde{K}}(T(E), T(x)), \quad (28)$$

where $\tilde{K} := K \circ R^t$. In particular, H_K is invariant under translation.

2. *If $F \subset E$ and $x \in \partial E \cap \partial F$, then $H_K(E, x) \leq H_K(F, x)$.*

We focus now on the functional H_0 defined in (5), which is a local anisotropic mean curvature functional, the anisotropy being encoded by M_K . As a first step, we establish the well-posedness of M_K and to this aim we recall the characterization of Sobolev functions in terms of absolute continuity on lines, whose definition we include here:

Definition 3.2. Let $\Omega \subset \mathbb{R}^d$ be an open set. A function $u: \Omega \rightarrow \mathbb{R}$ is absolutely continuous on lines if u is Borel measurable in Ω and locally absolutely continuous on almost all lines parallel to coordinate axes, that is, if $\{e_1, \dots, e_d\}$ is the canonical basis, for all $i = 1, \dots, d$ there exists $N_i \subset e_i^\perp$ such that $\mathcal{H}^{d-1}(N_i) = 0$ and for all $z \in e_i^\perp \cap N_i^c$ the function $I \ni t \mapsto u(z + te_i)$ is absolutely continuous on any compact interval I such that $z + te_i \in \Omega$ when $t \in I$.

Since absolutely continuous functions are differentiable a.e., we highlight that if u is absolutely continuous on lines, then it admits partial derivatives a.e. and hence the vector ∇u is a.e. defined. On the other hand, if a function has Sobolev regularity, then it has a representative which is absolutely continuous on lines. That is the content of the following result, whose proof can be found in [28, Theorem 2.3].

Theorem 3.3. *Let $\Omega \subset \mathbb{R}^d$ be an open set. For any $p \in [1, +\infty)$, $u: \Omega \rightarrow \mathbb{R}$ belongs to the Sobolev space $W^{1,p}(\Omega)$ if and only if it coincides a.e. with a function $\tilde{u} \in L^p(\Omega)$ that is absolutely continuous on lines and whose gradient $\nabla \tilde{u}$ belongs to $L^p(\Omega; \mathbb{R}^d)$.*

Thanks to (12) and to the theorem above, we may without loss of generality suppose that the kernel K is absolutely continuous on lines in $B(0, r)^c$ for all $r > 0$. We exploit this fact to prove boundedness and continuity of M_K .

Lemma 3.4. *Let a_0 be the constant in (16). Then, for all $e \in \mathbb{S}^{d-1}$,*

$$\int_{e^\perp} K(z) |z|^2 d\mathcal{H}^{d-1}(z) \leq a_0, \quad (29)$$

and M_K is continuous on \mathbb{S}^{d-1} .

Moreover, for any $e \in \mathbb{S}^{d-1}$, there holds

$$\lim_{r \rightarrow +\infty} r^\beta \int_{e^\perp \cap B(0, r)^c} K(z) |z|^2 d\mathcal{H}^{d-1}(z) = 0 \quad \text{for all } \beta < s. \quad (30)$$

Proof. By (a slight adaptation of) Theorem 3.3, for any $e \in \mathbb{S}^{d-1}$ and any $j \in \mathbb{N}$, there exists a \mathcal{H}^{d-1} -negligible $N_j \subset \{z \in e^\perp : j|z| \geq 1\}$ such that, for all $z \in e^\perp \cap N_j^c$ with $j|z| \geq 1$, the function $t \mapsto K(z + te)$ is absolutely continuous when t belongs to closed, bounded intervals. By the arbitrariness of $j \in \mathbb{N}$, we conclude that for \mathcal{H}^{d-1} -a.e. $z \in e^\perp$, $[a, b] \ni t \mapsto K(z + te)$ is absolutely continuous for any $a, b \in \mathbb{R}$.

Hence, by the Mean Value Theorem, for \mathcal{H}^{d-1} -almost every $z \in e^\perp$ we find

$$\lim_{\lambda \rightarrow 0^+} \frac{1}{\lambda} \int_{-\frac{\lambda}{2}|z|^2}^{\frac{\lambda}{2}|z|^2} K(z + te) dt = K(z) |z|^2. \quad (31)$$

Now, for any $\lambda > 0$, (14) guarantees that

$$a_\lambda(e) := \int_{Q_\lambda(e)} K(y) dy \in (0, +\infty).$$

Moreover, we have

$$\frac{1}{\lambda} \int_{e^\perp} \int_{-\frac{\lambda}{2}|z|^2}^{\frac{\lambda}{2}|z|^2} K(z + te) dt d\mathcal{H}^{d-1}(z) = \frac{a_\lambda(e)}{\lambda}.$$

In view of (31) and (16), we can take the limit $\lambda \rightarrow 0^+$ on both sides of the last equality and this yields (29), as desired.

Now we prove that M_K is continuous. We fix $e \in \mathbb{S}^{d-1}$ and we consider a sequence of rotations R_n such that $R_n \rightarrow \text{id}$. We have

$$\begin{aligned} |M_K(R_n e) - M_K(e)| &= \left| \int_{e^\perp} K(R_n z) R_n z \otimes R_n z d\mathcal{H}^{d-1} - \int_{e^\perp} K(z) z \otimes z d\mathcal{H}^{d-1} \right| \\ &\leq \left| \int_{e^\perp} K(R_n z) [R_n z \otimes R_n z - z \otimes z] d\mathcal{H}^{d-1} \right| \\ &\quad + \left| \int_{e^\perp} [K(R_n z) - K(z)] z \otimes z d\mathcal{H}^{d-1} \right|. \end{aligned}$$

Since $K \in L^1(B(0, r)^c)$ for all $r > 0$, it holds

$$\lim_{n \rightarrow +\infty} \|K \circ R_n - K\|_{L^1(B(0, r)^c)} = 0;$$

hence, we deduce that $K(R_n z) \rightarrow K(z)$ for \mathcal{H}^{d-1} -a.e. $z \in e^\perp$ and this, together with (29), gets that the upper bound we have on $|M_K(R_n e) - M_K(e)|$ vanishes as $n \rightarrow +\infty$.

Estimate (30) is an easy consequence of assumption (19). \square

From the very definition of M_K , we notice that $\pi_{\hat{n}^\perp} M_K(\hat{n}) \pi_{\hat{n}^\perp} = M_K(\hat{n})$. Using this, we observe that if Σ, x, ϕ , and \hat{n} are the same as in (5), we have

$$\begin{aligned} H_0(\Sigma, x) &= -\frac{1}{|\nabla \phi(x)|} \text{tr} (M_K(\hat{n}) \nabla^2 \phi(x)) \\ &= -\frac{1}{|\nabla \phi(x)|} \int_{\hat{n}^\perp} K(z) \nabla^2 \phi(x) z \cdot z d\mathcal{H}^{d-1}(z). \end{aligned} \quad (32)$$

Remark 3. Let us consider a smooth hypersurface Σ whose outer unit normal at a given point x is \hat{n} , and the map $T(y) := Ry + h$, where R is an orthogonal matrix and $h \in \mathbb{R}^d$. Then, it is easy to check by using (32) that it holds

$$H_0(\Sigma, x) = \tilde{H}_0(T(\Sigma), T(x)), \quad (33)$$

where \tilde{H}_0 is the anisotropic mean curvature functional associated with the kernel $\tilde{K} := K \circ R^t$. To prove our claim, we observe that if $\Sigma = \{y \in \mathbb{R}^d : \phi(y) = 0\}$

for some smooth $\phi: \mathbb{R}^d \rightarrow \mathbb{R}$, then $T(\Sigma) = \{y \in \mathbb{R}^d : \psi(y) = 0\}$ with $\psi(y) := \phi(R^t(y-x))$. We have

$$\nabla\psi(T(y)) = R\nabla\phi(y) \quad \text{and} \quad \nabla^2\psi(T(y)) = R\nabla^2\phi(y)R^t,$$

and, therefore,

$$\begin{aligned} \tilde{H}_0(T(\Sigma), T(x)) &= -\frac{1}{|R\nabla\phi(x)|} \int_{R(\hat{n}^\perp)} \tilde{K}(z) (R\nabla^2\phi(x)R^t) z \cdot z d\mathcal{H}^{d-1}(z) \\ &= -\frac{1}{|\nabla\phi(x)|} \int_{\hat{n}^\perp} K(z) \nabla^2\phi(x) z \cdot z d\mathcal{H}^{d-1}(z). \end{aligned}$$

Remark 4 (Connection with standard mean curvature). When K is radial, that is, $K(x) = K_0(|x|)$ for some $K_0: (0, +\infty) \rightarrow [0, +\infty)$, then H_0 coincides with the standard mean curvature, up to a multiplicative constant. Indeed, let Σ be a C^2 hypersurface such that $0 \in \Sigma$ and $\Sigma \cap U = \{y \in U : \phi(y) = 0\}$ for some neighbourhood U of 0 and some smooth function $\phi: U \rightarrow \mathbb{R}$. We suppose also that $\nabla\phi(0) \neq 0$ and that the outer unit normal to Σ at 0 is e_d . We recall the expression of the mean curvature H of Σ at 0:

$$H(\Sigma, 0) = -\frac{1}{\omega_{d-1} |\nabla\phi(0)|} \int_{\mathbb{S}^{d-2}} \nabla^2\phi(0) e \cdot e d\mathcal{H}^{d-2}(e),$$

with $\omega_{d-1} := \mathcal{H}^{d-1}(\mathbb{S}^{d-1})$.

If $K(x) = K_0(|x|)$, then formula (29) reads

$$c_K := \int_0^{+\infty} r^d K_0(r) dr < +\infty,$$

and, consequently, we have

$$\begin{aligned} H_0(\Sigma, 0) &= -\frac{1}{|\nabla\phi(0)|} \int_0^{+\infty} r^d K_0(r) dr \int_{\mathbb{S}^{d-2}} \nabla^2\phi(0) e \cdot e d\mathcal{H}^{d-2}(e) \\ &= \omega_{d-1} c_K H(\Sigma, 0). \end{aligned}$$

4. Barriers and level-set flow for geometric evolutions. We devote this section to some basics about level-set formulations and barriers for the geometric flows driven by the curvatures H_K and H_0 . In particular, we recall existence and uniqueness results for the level-set flow, and we revise its connections with the notion of geometric barriers.

We consider the following geometric evolutions for the family of sets $\{E(t)\}_{t \geq 0}$:

$$\partial_t x(t) \cdot \hat{n} = -H_\varepsilon(E(t), x), \quad \partial_t x(t) \cdot \hat{n} = -H_0(E(t), x), \quad (34)$$

where \hat{n} is the outer unit normal to $\partial E(t)$ at the point $x(t)$ and $H_\varepsilon \varepsilon$ is the rescaled version of H_K defined in (4). In addition, we accompany these equations with an initial datum E_0 , which we assume to be a bounded set.

Let us begin with the level-set formulations of the geometric flows (34). First of all, we interpret the initial datum E_0 as the superlevel set of a suitable function $u_0: \mathbb{R}^d \rightarrow \mathbb{R}$. Explicitly, we suppose that $E_0 = \{x : u_0(x) \geq 0\}$ and $\partial E_0 = \{x : u_0(x) = 0\}$; moreover, throughout the paper we assume that

$$u_0: \mathbb{R}^d \rightarrow \mathbb{R} \text{ is Lipschitz and constant outside a compact } C. \quad (35)$$

Then, we consider the nonlocal and local Cauchy problems:

$$\begin{cases} \partial_t u(t, x) + |\nabla u(t, x)| H_\varepsilon(\{y : u(t, y) \geq u(t, x)\}, x) = 0 \\ u(0, x) = u_0(x) \end{cases} \quad \begin{matrix} (t, x) \in [0, +\infty) \times \mathbb{R}^d \\ x \in \mathbb{R}^d \end{matrix}, \quad (36)$$

$$\begin{cases} \partial_t u(t, x) - \operatorname{tr} \left(M_K \left(\widehat{\nabla u(t, x)} \right) \nabla^2 u(t, x) \right) = 0 \\ u(0, x) = u_0(x) \end{cases} \quad \begin{matrix} (t, x) \in [0, \infty) \times \mathbb{R}^d \\ x \in \mathbb{R}^d \end{matrix}. \quad (37)$$

Observe that

$$|\nabla u(x)| H_0(\{y : u(y) = u(x)\}, x) = -\operatorname{tr} \left(M_K \left(\widehat{\nabla u(x)} \right) \nabla^2 u(x) \right)$$

(recall that $\hat{p} := p/|p|$ if $p \neq 0$).

We remind the definition of viscosity solution for nonlocal equations, which goes back to the work [36], see also [29, 24, 19, 17].

Definition 4.1 (Solution to the rescaled problems). A locally bounded, upper semicontinuous function (resp. lower semicontinuous) $u_\varepsilon : [0, +\infty) \times \mathbb{R}^d \rightarrow \mathbb{R}$ is a viscosity subsolution (resp. supersolution) to the problem (36) if

1. $u_\varepsilon(0, x) \leq u_0(x)$ for all $x \in \mathbb{R}^d$ (resp. $u_\varepsilon(0, x) \geq u_0(x)$);
2. for all $(t, x) \in (0, +\infty) \times \mathbb{R}^d$ and for all $\phi \in C^2([0, +\infty) \times \mathbb{R}^d)$ such that $u_\varepsilon - \phi$ has a maximum at (t, x) (resp. has a minimum at (t, x)), it holds

$$\partial_t \phi(t, x) \leq 0 \quad (\text{resp. } \partial_t \phi(t, x) \geq 0) \quad \text{when } \nabla \phi(t, x) = 0,$$

or

$$\partial_t \phi(t, x) + |\phi(t, x)| H_\varepsilon(\{y : \phi(t, y) \geq \phi(t, x)\}, x) \leq 0$$

$$(\text{resp. } \partial_t \phi(t, x) + |\phi(t, x)| H_\varepsilon(\{y : \phi(t, y) > \phi(t, x)\}, x) \geq 0) \quad \text{otherwise.}$$

A continuous function $u_\varepsilon : [0, +\infty) \times \mathbb{R}^d \rightarrow \mathbb{R}$ is a viscosity solution to (36) if it is both a viscosity sub- and supersolution.

Existence and uniqueness of a viscosity solution to (36) were proved in [19], in a very general setting. A similar result can also be found in [29].

Theorem 4.2 (Comparison principle and existence of solutions to the nonlocal problem). *If the standing assumptions on the kernel and (35) hold, for all $\varepsilon > 0$, if $v_\varepsilon, w_\varepsilon : [0, +\infty) \times \mathbb{R}^d \rightarrow \mathbb{R}$ are respectively a sub- and a supersolution to (36), then $v_\varepsilon(t, x) \leq w_\varepsilon(t, x)$ for all $(t, x) \in [0, +\infty) \times \mathbb{R}^d$.*

Moreover, (36) admits a unique bounded, Lipschitz continuous viscosity solution in $[0, +\infty) \times \mathbb{R}^d$, which is constant in $\mathbb{R}^d \setminus C$, for some compact set $C \subset \mathbb{R}^d$.

We recall also the definition of solution to the limit problem (37), see [24].

Definition 4.3 (Solution to the limit problem). A locally bounded, upper semicontinuous function (resp. lower semicontinuous function) $u : [0, \infty) \times \mathbb{R}^d \rightarrow \mathbb{R}$ is a viscosity subsolution (resp. supersolution) to the Cauchy's problem (37) if

1. $u(0, x) \leq u_0(x)$ for all $x \in \mathbb{R}^d$, (resp. $u(0, x) \geq u_0(x)$);
2. for all $(t, x) \in (0, +\infty) \times \mathbb{R}^d$ and for all $\phi \in C^2([0, +\infty) \times \mathbb{R}^d)$ such that $u - \phi$ has a maximum at (t, x) (resp. a minimum at (t, x)) it holds

$$\partial_t \phi(t, x) \leq 0 \quad (\text{resp. } \partial_t \phi(t, x) \geq 0) \quad \text{when } \nabla \phi(t, x) = 0 \text{ and } \nabla^2 \phi(t, x) = 0$$

or

$$\partial_t \phi(t, x) - \operatorname{tr} \left(M_K \left(\widehat{\nabla \phi(t, x)} \right) \nabla^2 \phi(t, x) \right) \leq 0 \quad (\text{resp. } \geq 0) \quad \text{otherwise.}$$

A continuous function $u : [0, +\infty) \times \mathbb{R}^d \rightarrow \mathbb{R}$ is a viscosity solution to (37) if it is both a viscosity sub- and supersolution.

As for existence of solutions, we observe that the function

$$F_0 : \begin{array}{ccc} \mathbb{R}^d \setminus \{0\} \times \text{Sym}(d) & \longrightarrow & \mathbb{R} \\ (p, X) & \longmapsto & -\text{tr}(M_K(\hat{p})X) \end{array}$$

that defines the problem (37) has the three following properties:

1. it is continuous;
2. it is geometric, that is, for all $\lambda > 0$, $\sigma \in \mathbb{R}$, $p \in \mathbb{R}^d \setminus \{0\}$ and $X \in \text{Sym}(d)$ it holds $F_0(\lambda p, \lambda X + \sigma p \otimes p) = \lambda F_0(p, X)$.
3. it is degenerate elliptic, that is, $F_0(p, X) \geq F_0(p, Y)$ for all $p \in \mathbb{R}^d \setminus \{0\}$ and $X, Y \in \text{Sym}(d)$ such that $X \leq Y$.

It is well known [9, 22] that these conditions grant existence and uniqueness of a viscosity solution:

Theorem 4.4. *Let us suppose that (35) holds. Then, the Cauchy's problem (37) admits a unique bounded, Lipschitz continuous viscosity solution in $[0, +\infty) \times \mathbb{R}^d$, which is constant in $\mathbb{R}^d \setminus C$, for some compact set $C \subset \mathbb{R}^d$.*

Summing up, owing to Theorems 4.2 and 4.4, we get that, for every initial datum u_0 as in (35), there exist a unique viscosity solution u_ε to (37) and a unique viscosity solution u to (37). We define the level-set flows associated with these solutions. For every $\lambda \in \mathbb{R}$, we set

$$\begin{aligned} E_{\varepsilon, \lambda}^+(t) &= \{x \in \mathbb{R}^d : u_\varepsilon(t, x) \geq \lambda\}, & E_{\varepsilon, \lambda}^-(t) &= \{x \in \mathbb{R}^d : u_\varepsilon(t, x) > \lambda\}, \\ E_\lambda^+(t) &= \{x \in \mathbb{R}^d : u(t, x) \geq \lambda\}, & E_\lambda^-(t) &= \{x \in \mathbb{R}^d : u(t, x) > \lambda\}. \end{aligned} \quad (39)$$

It is well known that, as long as they are smooth, these families are solutions to the geometric flows (34) resp. with H_ε and H_0 and initial datum $E_\lambda = \{x \in \mathbb{R}^d : u_0(x) \geq \lambda\}$.

Geometric evolutions may be formulated as PDEs involving distance functions from the moving front, see for instance the survey [4] by Ambrosio; in the following definitions, we use them to express a regularity property both in time and space for a class of evolving sets (see 2 below) w.r.t. a generic geometric law.

Definition 4.5. Let $0 \leq t_0 < t_1 < +\infty$. We say that the evolutions of sets $[t_0, t_1] \ni t \mapsto D(t)$ is a *geometric subsolution* (resp. *supersolution*) to the flow associated with the curvature functional H if

1. $D(t)$ is closed and $\partial D(t)$ is compact for all $t \in [t_0, t_1]$;
2. there exists an open set $U \subset \mathbb{R}^d$ such that the distance function $(t, x) \mapsto d_{D(t)}(x)$ is of class C^∞ in $[t_0, t_1] \times U$ and $\partial D(t) \subset U$ for all $t \in [t_0, t_1]$;
3. for all $t \in (t_0, t_1)$ and $x(t) \in \partial D(t)$, it holds

$$\partial_t x(t) \cdot \hat{n} \leq -H(D(t), x(t)) \quad (\text{resp. } \partial_t x(t) \cdot \hat{n} \geq -H(D(t), x(t))), \quad (40)$$

where \hat{n} is the outer unit normal to $D(t)$ at x .

When strict inequalities hold, $D(t)$ is called *strict geometric subsolution* (resp. *strict geometric supersolution*).

Remark 5. We notice that, for any $p \in \mathbb{R}^d \setminus \{0\}$ and $X \in \text{Sym}(d)$, by (29) we get that

$$|\text{tr}(M_K(\hat{p})X)| = \frac{1}{2} \left| \int_{\hat{p}^\perp} K(z) Xz \cdot z d\mathcal{H}^{d-1}(z) \right| \leq \frac{a_0}{2} |X|,$$

This ensures that geometric sub- and supersolution for the flow associated with H_0 exist (see [9]).

Next, we remind the notion of geometric barriers w.r.t. these smooth evolutions:

Definition 4.6. Let $T > 0$ and \mathcal{F}^- and \mathcal{F}^+ be, respectively, the families of strict geometric sub- and supersolution to the flow associated with some curvature functional H , as introduced in Definition 4.5.

1. We say that the evolution of sets $[0, T] \ni t \mapsto E(t)$ is an *outer barrier* w.r.t. \mathcal{F}^- (resp. \mathcal{F}^+) if whenever $[t_0, t_1] \subset [0, T]$ and $[t_0, t_1] \ni t \mapsto D(t)$ is a smooth strict subsolution (resp. $F(t)$ is a smooth strict supersolution) such that $D(t_0) \subset E(t_0)$, then we get $D(t_1) \subset E(t_1)$ (resp. such that $F(t_0) \subset E(t_0)$, then we get $F(t_1) \subset E(t_1)$).
2. Analogously, $[0, T] \ni t \mapsto E(t)$ is an *inner barrier* w.r.t. the family \mathcal{F}^- (resp. \mathcal{F}^+) if whenever $[t_0, t_1] \subset [0, T]$ and $[t_0, t_1] \ni t \mapsto D(t)$ is a smooth strict subsolution (resp. supersolution) such that $E(t_0) \subset \text{int}(D(t_0))$, then $E(t_1) \subset \text{int}(D(t_1))$ (resp. $E(t_0) \subset \text{int}(F(t_0))$, then $E(t_1) \subset \text{int}(F(t_1))$).

We are interested in barriers for the anisotropic mean curvature motion (34) because they are comparable with level-sets flows, as the next theorem shows. Its proof can be found in [9, theorem 3.2]. For further reading about barriers for general geometric, local evolution problems, we refer to that paper and to [10].

Theorem 4.7. Let u be the unique solution to (37) with initial datum u_0 as in (35). Let E_λ^\pm the sets defined in (39).

1. The map $[0, T] \ni t \mapsto E_\lambda^-(t)$ is the minimal outer barrier for the family of strict geometric subsolutions associated with H_0 , that is $E_\lambda^-(t)$ is an outer barrier and $E_\lambda^-(t) \subset E(t)$ for any other outer barrier $E(t)$.
2. The map $[0, T] \ni t \mapsto E_\lambda^+(t)$ is the maximal inner barrier for the family of geometric strict supersolutions associated with H_0 , that is $E_\lambda^+(t)$ is an inner barrier and $E(t) \subset E_\lambda^+(t)$ for any other inner barrier $E(t)$.

Lastly, we mention a comparison principle concerning the level-set flow and strict geometric sub- and supersolutions for the nonlocal problems, see [17, Proposition A.10].

Proposition 2. Let $u_\varepsilon : [0, +\infty) \times \mathbb{R}^d \rightarrow \mathbb{R}$ be the viscosity solution to (36) with initial datum u_0 as in (35). Let $E_{\varepsilon, \lambda}^\pm(t)$ be as in (38). Then, the evolutions $t \mapsto E_{\varepsilon, \lambda}^-(t)$ and $t \mapsto E_{\varepsilon, \lambda}^+(t)$ are, respectively, an outer barrier w.r.t geometric strict subsolutions to (36) and an inner barrier w.r.t geometric strict supersolutions to (36).

5. Convergence of the rescaled nonlocal curvatures. This section is devoted to the proof of Theorem 1.1, the first main result of the paper. The argument consists of two steps: firstly, we deal in Lemma 5.1 with the pointwise convergence of the curvatures, providing a precise estimate on the error; then, in Proposition 3, we show that it is possible to make the estimate uniform when smooth, compact hypersurfaces are considered.

We fix the notations that we are going to use in the current section. Let $E \subset \mathbb{R}^d$ be a set of class C^2 . Then for all $x \in \Sigma := \partial E$, there exist an open neighborhood U of x and $\phi \in C^2(U)$ such that

$$\Sigma \cap U = \{y \in U : \phi(y) = 0\}, \quad E \cap U = \{y \in U : \phi(y) > 0\},$$

and $\nabla\phi(y) \neq 0$ for all $y \in \Sigma \cap U$. We write \hat{n} for the outer unit normal to Σ at x . Lastly, by the Implicit Function Theorem, there exist $\bar{\delta} := \bar{\delta}(x) > 0$ and $f: \hat{n}^\perp \cap B(0, \bar{\delta}) \rightarrow (-\bar{\delta}, \bar{\delta})$ such that (23) and (24) hold, and $\inf_{y \in C_{\hat{n}}(x, \bar{\delta})} |\nabla\phi(y)| > 0$.

Lemma 5.1. *Let $E \subseteq \mathbb{R}^d$ be such that $\Sigma := \partial E$ is of class C^2 . Let $x \in \Sigma$, $\bar{\delta}$, and f be as above, and let $s \in (0, 1)$ be the exponent in (19). Then, for all $\alpha, \beta \in (0, s)$, there exist $q > 1$ and $\bar{\varepsilon} \in (0, 1)$ such that $q\bar{\varepsilon} \leq \bar{\delta}$ and that, for all $\varepsilon \in (0, \bar{\varepsilon})$ and all $\delta \in (q\varepsilon, \bar{\delta})$, it holds*

$$|H_\varepsilon(E, x) - H_0(\Sigma, x)| \leq \mathcal{E}(\varepsilon, \delta),$$

where

$$\mathcal{E}(\varepsilon, \delta) := \frac{1}{\bar{\delta}} \left(\frac{\varepsilon}{\bar{\delta}} \right)^\alpha + (b_0 + 1) \|\nabla^2 f\|_{L^\infty(D)}^2 \delta + a_0 \omega_f(\delta) + |\nabla^2 f(0)| \left(\frac{\varepsilon}{\bar{\delta}} \right)^\beta, \quad (41)$$

with $D := \hat{n}^\perp \cap B(0, \bar{\delta})$ and

$$\omega_f(\delta) := \sup_{z \in B(0, \delta)} |\nabla^2 f(z) - \nabla^2 f(0)|. \quad (42)$$

Proof. We start by observing that, without loss of generality, we may assume that $x = 0$ and $\hat{n} = e_d := (0, \dots, 0, 1)$.

The argument is similar to the one followed to prove estimate (22). There exists $f: D \rightarrow (-\bar{\delta}, \bar{\delta})$ of class C^2 such that $f(0) = 0$, $\nabla f(0) = 0$, and (23) and (24) hold. Moreover,

$$\partial_i f = \frac{\partial_i \phi}{\partial_d \phi}, \quad (43)$$

$$\partial_{i,j}^2 f = \frac{1}{\partial_d \phi} (\partial_{i,j}^2 \phi + \partial_i f \partial_{j,d}^2 \phi + \partial_j f \partial_{i,d}^2 \phi + \partial_i f \partial_j f \partial_{d,d}^2 \phi) \quad (44)$$

for $i, j = 1, \dots, d-1$. Let us introduce the function

$$f_\varepsilon(z) := \frac{f(\varepsilon z)}{\varepsilon}.$$

Since f is of class C^2 , for all $z \in D$ there exists z' such that $f_\varepsilon(z) = (\varepsilon \nabla^2 f(\varepsilon z') \cdot z \cdot z)/2$. When t ranges between $-f_\varepsilon(-z)$ and $f_\varepsilon(z)$, we thus see that

$$|t| \leq \frac{\varepsilon}{2} \|\nabla^2 f\|_{L^\infty(D)} |z|^2. \quad (45)$$

Let us fix $0 < \varepsilon < \delta < \bar{\delta}$. We split H_ε into two different contributions:

$$H_\varepsilon(E, 0) = I_\varepsilon^0 + I_\varepsilon^1 := -\frac{1}{\varepsilon} \int_C K_\varepsilon(y) \tilde{\chi}_E(y) dy - \frac{1}{\varepsilon} \int_{C^c} K_\varepsilon(y) \tilde{\chi}_E(y) dy,$$

where $C := C_{e_d}(0, \delta)$. The first integral takes into account the interactions with points that are close to 0, and it approximates the anisotropic mean curvature at 0 when ε is small; the second term encodes the energy stored far away from the origin. Observe that

$$I_\varepsilon^0 = \frac{1}{\varepsilon} \int_{e_d^\perp \cap B(0, \frac{\delta}{\varepsilon})} \int_{-f_\varepsilon(-z)}^{f_\varepsilon(z)} K(z + te_d) dt d\mathcal{H}^{d-1}(z).$$

Let us define

$$J_\varepsilon := \frac{1}{\varepsilon} \int_{e_d^\perp \cap B(0, \frac{\delta}{\varepsilon})} K(z) [f_\varepsilon(z) + f_\varepsilon(-z)] d\mathcal{H}^{d-1}(z),$$

and recall that, in view of (32),

$$H_0(\Sigma, 0) = \int_{e_d^\perp} K(z) \nabla^2 f(0) z \cdot z d\mathcal{H}^{d-1}(z).$$

We consider the chain of inequalities

$$|H_\varepsilon(E, 0) - H_0(\Sigma, 0)| = |I_\varepsilon^0 + I_\varepsilon^1 - H_0(\Sigma, 0)| \leq |I_\varepsilon^0 - J_\varepsilon| + |J_\varepsilon - H_0(\Sigma, 0)| + |I_\varepsilon^1|,$$

and we estimate each term separately.

We start with I_ε^1 . We remark that, as a consequence of (20), for all $\alpha < s$ there exists $q_1 > 1$ such that

$$|I_\varepsilon^1| = \frac{1}{\varepsilon} \int_{B(0, \frac{\delta}{\varepsilon})^c} K(y) dy \leq \frac{1}{\delta} \left(\frac{\varepsilon}{\delta}\right)^\alpha \quad \text{whenever } q_1 \varepsilon < \delta. \quad (46)$$

We proceed with the other terms. We observe that

$$|I_\varepsilon^0 - J_\varepsilon| \leq \frac{1}{\varepsilon} \int_{e_d^\perp \cap B(0, \frac{\delta}{\varepsilon})} \left| \int_{-f_\varepsilon(-z)}^{f_\varepsilon(z)} [K(z + te_d) - K(z)] dt \right| d\mathcal{H}^{d-1}(z). \quad (47)$$

By Theorem 3.3, for \mathcal{H}^{d-1} -a.e. $z \in e_d^\perp$, it holds

$$K(z + te_d) - K(z) = \int_0^t \partial_d K(z + se_d) ds,$$

and this, combined with (45), implies that

$$|K(z + te_d) - K(z)| \leq \int_{-\frac{\varepsilon}{2} \|\nabla^2 f\|_{L^\infty(D)} |z|^2}^{\frac{\varepsilon}{2} \|\nabla^2 f\|_{L^\infty(D)} |z|^2} |\nabla K(z + se_d)| ds.$$

We plug this inequality in (47) and we obtain

$$\begin{aligned} & |I_\varepsilon^0 - J_\varepsilon| \\ & \leq \|\nabla^2 f\|_{L^\infty(D)} \int_{e_d^\perp \cap B(0, \frac{\delta}{\varepsilon})} |z|^2 \int_{-\frac{\varepsilon}{2} \|\nabla^2 f\|_{L^\infty(D)} |z|^2}^{\frac{\varepsilon}{2} \|\nabla^2 f\|_{L^\infty(D)} |z|^2} |\nabla K(z + se_d)| ds d\mathcal{H}^{d-1}(z) \\ & \leq \|\nabla^2 f\|_{L^\infty(D)} \frac{\delta}{\varepsilon} \int_{Q(\varepsilon)} |y| |\nabla K(y)| dy, \end{aligned}$$

where $Q(\varepsilon) := Q_{\varepsilon \|\nabla^2 f\|_{L^\infty(D)}}(e_d)$. By using (17) we get that there exists $\eta \in (0, \bar{\delta})$ such that

$$|I_\varepsilon^0 - J_\varepsilon| \leq (b_0 + 1) \|\nabla^2 f\|_{L^\infty(D)}^2 \delta \quad \text{whenever } \varepsilon < \eta. \quad (48)$$

Finally, we have

$$\begin{aligned} |J_\varepsilon - H_0(\Sigma, 0)| & \leq \omega_f(\delta) \int_{e_d^\perp \cap B(0, \frac{\delta}{\varepsilon})} K(z) |z|^2 d\mathcal{H}^{d-1}(z) \\ & \quad + |\nabla^2 f(0)| \int_{e_d^\perp \cap B(0, \frac{\delta}{\varepsilon})^c} K(z) |z|^2 d\mathcal{H}^{d-1}(z), \end{aligned}$$

ω_f being defined in (42). Thanks to (30), for all $\beta < s$, there exists $q_2 > 0$ such that, if $q_2 \varepsilon < \delta$, then

$$\left(\frac{\delta}{\varepsilon}\right)^\beta \int_{e_d^\perp \cap B(0, \frac{\delta}{\varepsilon})^c} K(z) |z|^2 d\mathcal{H}^{d-1}(z) \leq 1.$$

Recalling (29), we thus find

$$|J_\varepsilon - H_0(\Sigma, 0)| \leq a_0 \omega_f(\delta) + |\nabla^2 f(0)| \left(\frac{\varepsilon}{\delta}\right)^\beta \quad \text{whenever } q_2 \varepsilon < \delta. \quad (49)$$

Now, if we set $q := \max\{q_1, q_2\} > 1$ with q_1 and q_2 as above, both (46) and (49) hold for all $\varepsilon, \delta > 0$ such that $q\varepsilon < \delta < \bar{\delta}$. Besides, if we pick $\bar{\varepsilon} := \min\{\eta, \bar{\delta}/q\}$, (48) is satisfied as well whenever $\varepsilon < \bar{\varepsilon}$. This yields the conclusion. \square

Remark 6. In the proof of Lemma 5.1, we did not exploit assumptions (13) and (18). These will be useful in the proof of Proposition 4.

By applying the estimate on the error term given in Lemma 5.1, we deduce the desired uniform convergence.

Proposition 3. *Under the same notation and assumptions of Lemma 5.1, there exists a constant $c := c(\alpha, \beta, a_0, b_0) > 0$ such that for all $\gamma \in \left(0, \frac{\alpha}{1+\alpha}\right)$, it holds*

$$|H_\varepsilon(E, x) - H_0(\Sigma, x)| \leq c \left(\varepsilon^{\alpha-\gamma(1+\alpha)} + \|\nabla^2 f\|_{L^\infty(D)} \varepsilon^\gamma + \omega_f(q\varepsilon^\gamma) + |\nabla^2 f(0)| \varepsilon^{(1-\gamma)\beta} \right).$$

In particular, if Σ is compact, the conclusion of Theorem 1.1 holds.

Proof. We start by proving that pointwise convergence holds. We choose $\gamma \in (0, \alpha/(1+\alpha))$ and we observe that, for any $\varepsilon < \bar{\varepsilon} < 1$, we have $q\varepsilon < q\varepsilon^\gamma$. We may therefore pick $\delta = q\varepsilon^\gamma$ in (41) and check that $\mathcal{E}(\varepsilon, q\varepsilon^\gamma) \rightarrow 0$ when $\varepsilon \rightarrow 0^+$. The pointwise convergence follows.

Now, we turn to the case when Σ is compact and of class C^2 . We denote by \hat{n}_x the outer unit normal to Σ at x and by \hat{n}_x^\perp the tangent plane at the same point. Let us also define

$$V_\Sigma(\delta) := \{y \in \mathbb{R}^d : \inf_{z \in \Sigma} |y - z| < \delta\},$$

and

$$\bar{\delta} := \sup\{\delta > 0 : \text{the boundary of } V_\Sigma(\delta) \text{ is of class } C^2\} > 0.$$

This ensures that, for any $x \in \Sigma$, the implicit function f defined on \hat{n}_x^\perp ranges in $(-\bar{\delta}, \bar{\delta})$. Let us denote this function by f_x to stress that it depends on x . There exists $\bar{\varepsilon} < 1$ such that for all $\varepsilon \in (0, \bar{\varepsilon})$, for all $\gamma \in (0, \alpha/(1+\alpha))$, and for all $x \in \Sigma$ it holds

$$|H_\varepsilon(E, x) - H_0(\Sigma, x)| \leq c \left(\varepsilon^{\alpha-\gamma(1+\alpha)} + \|\nabla^2 f_x\|_{L^\infty(\hat{n}_x^\perp \cap B(0, \bar{\delta}))} \varepsilon^\gamma + \omega_{f_x}(q\varepsilon^\gamma) + |\nabla^2 f_x(0)| \varepsilon^{(1-\gamma)\beta} \right).$$

Since Σ is compact, $|\nabla^2 f_x(0)|$ and $\|\nabla^2 f_x\|_{L^\infty(\hat{n}_x^\perp \cap B(0, \bar{\delta}))}$ are bounded above by the $L^\infty(\Sigma)$ -norm of the second fundamental form of Σ ; also, there exists a function ω_Σ that vanishes in 0, that is decreasing and that satisfies $\omega_{f_x}(\delta) \leq \omega_\Sigma(\delta)$ whenever δ is sufficiently small. In conclusion, we obtain an estimate on $|H_\varepsilon(E, x) - H_0(\Sigma, x)|$ that is uniform in x , and the thesis holds. \square

6. A priori estimates for the rescaled problems. In this section we establish a compactness property for the family of solutions to the Cauchy's problems (36). Even though the result is known, we sketch its proof, because it is not explicitly stated in the literature for our setting.

Proposition 4. *Assume that $u_0: \mathbb{R}^d \rightarrow \mathbb{R}$ is as in (35), and let u_ε be the unique continuous viscosity solution to (36). Then,*

$$|u_\varepsilon(t, x) - u_\varepsilon(t, y)| \leq \|\nabla u_0\|_{L^\infty(\mathbb{R}^d)} |x - y| \quad \text{for all } t \in [0, T] \text{ and } x, y \in \mathbb{R}^d, \quad (50)$$

and there exists a constant $c > 0$ independent of ε such that

$$|u_\varepsilon(t, x) - u_\varepsilon(s, x)| \leq \|\nabla u_0\|_{L^\infty(\mathbb{R}^d)} \sqrt{c|t - s|} \quad \text{for all } t, s \in [0, T] \text{ and } x \in \mathbb{R}^d. \quad (51)$$

Proof. The equi-Lipschitz property (50) is a consequence of the Lipschitz continuity of the datum and of the comparison principle. We skip the proof, since it is completely standard and can be found, for instance, in [17, 24].

For the proof of equi-Hölder continuity, we follow the strategy of Section 5 in [24]. We point out that, however, the case that we treat differs from the one in the reference, mainly because of the possible singularity of our interaction kernel.

We fix $\eta > 0$ and $x \in \mathbb{R}^d$ and we consider

$$\phi(t, y) = Lt + A\sqrt{|y - x|^2 + \eta^2} + u_0(x), \quad (52)$$

where $A := \|\nabla u_0\|_{L^\infty(\mathbb{R}^d)}$. We claim that, for $L > 0$ sufficiently large, ϕ is a supersolution to (36) for any $\varepsilon \in (0, 1)$.

To prove the claim, we remark first of all that $\phi(0, y) \geq u_0(y)$ as a consequence of the Lipschitz continuity of u_0 . Also, we observe that, for any $y \in \mathbb{R}^d$,

$$\{z \in \mathbb{R}^d : \phi(t, z) \geq \phi(t, y)\} = B(x, |y - x|)^c.$$

Hence, to show that ϕ is a supersolution, it is sufficient to choose L so large that

$$\frac{L}{A} \geq \frac{|y - x|}{\sqrt{|y - x|^2 + \eta^2}} H_\varepsilon(B(x, |y - x|), y) \quad \text{for all } y \in \mathbb{R}^d \text{ and } \varepsilon \in (0, 1).$$

Recalling that the nonlocal curvature is invariant under translations, if we set $e := \frac{y - x}{|y - x|}$ and $r := |y - x|$, we have that the last inequality holds if and only if

$$\frac{L}{A} \geq \frac{r}{\sqrt{r^2 + \eta^2}} H_\varepsilon(B(-re, r), 0) \quad \text{for all } r > 0, e \in \mathbb{S}^{d-1} \text{ and } \varepsilon \in (0, 1). \quad (53)$$

So, we are left to prove that there exists $L_0 := L_0(\eta) > 0$ such that

$$\sup_{r > 0, e \in \mathbb{S}^{d-1}} \sup_{\varepsilon \in (0, 1)} \frac{r}{\sqrt{r^2 + \eta^2}} H_\varepsilon(B(-re, r), 0) \leq L_0; \quad (54)$$

this clearly yields (53) for $L = AL_0$.

To recover estimate (54), we use inequality (22). We get

$$0 \leq H_\varepsilon(B(-re, r), 0) \leq \int_{Q_{\frac{\varepsilon}{r}}(e)} K(y) dy + \int_{B(0, \frac{r}{2\varepsilon})^c} K(y) dy,$$

and hence

$$\frac{r}{\sqrt{r^2 + \eta^2}} H_\varepsilon(B(-re, r), 0) \leq \frac{r}{\varepsilon\eta} \left[\int_{Q_{\frac{\varepsilon}{r}}(e)} K(y) dy + \int_{B(0, \frac{r}{2\varepsilon})^c} K(y) dy \right].$$

By assumptions (13), (18), (16), and (20), there exist $\lambda, \Lambda > 0$ with the following properties:

1. $\lambda < \Lambda$;
2. if $r < \lambda\varepsilon$, then

$$\frac{r}{\varepsilon} \int_{Q_{\frac{\varepsilon}{r}}(e)} K(y) dy \leq \frac{1}{2} \quad \text{and} \quad \frac{r}{\varepsilon} \int_{B(0, \frac{\varepsilon}{2\varepsilon})^c} K(y) dy \leq \frac{1}{2}$$

and, consequently,

$$\frac{r}{\sqrt{r^2 + \eta^2}} H_\varepsilon(B(-re, r), 0) \leq \frac{1}{\eta}; \quad (55)$$

3. if $r > \Lambda\varepsilon$, then

$$\frac{r}{\varepsilon} \int_{Q_{\frac{\varepsilon}{r}}(e)} K(y) dy \leq a_0 + \frac{1}{2} \quad \text{and} \quad \frac{r}{\varepsilon} \int_{B(0, \frac{\varepsilon}{2\varepsilon})^c} K(y) dy \leq \frac{1}{2}$$

and, consequently,

$$\frac{r}{\sqrt{r^2 + \eta^2}} H_\varepsilon(B(-re, r), 0) \leq \frac{a_0 + 1}{\eta}. \quad (56)$$

Now, only the case $\lambda\varepsilon \leq r \leq \Lambda\varepsilon$ is left to discuss. In this intermediate regime, recalling (15), we easily obtain

$$\frac{r}{\sqrt{r^2 + \eta^2}} H_\varepsilon(B(-re, r), 0) \leq \frac{\Lambda}{\eta} \left(c + \int_{B(0, \frac{\lambda}{2})^c} K(y) dy \right), \quad (57)$$

with $c > 0$ depending only on λ .

In view of (55), (56), and (57), there exists a constant $c := c(a_0, \lambda, \Lambda) > 0$ such that

$$\sup_{r>0, e \in \mathbb{S}^{d-1}} \sup_{\varepsilon \in (0, 1)} \frac{r}{\sqrt{r^2 + \eta^2}} H_\varepsilon(B(-re, r), 0) \leq \frac{c}{\eta},$$

and (53) thus holds for the choice $L = Ac/\eta$.

Summing up, we proved that, for any fixed $x \in \mathbb{R}^d$, the function

$$\phi(t, y) = A \left(\frac{c}{\eta} t + \sqrt{|y - x|^2 + \eta^2} \right) + u_0(x)$$

is a supersolution to (36) for any $\varepsilon > 0$.

By means of an analogous argument we can prove that, for all $x \in \mathbb{R}^d$, the function

$$\psi(y) := -A \left(\frac{c}{\eta} t + \sqrt{|y - x|^2 + \eta^2} \right) + u_0(x),$$

is a subsolution to (36) for any $\varepsilon > 0$ and some $c = c(a_0, \lambda, \Lambda)$.

All in all, thanks to the comparison principle in Theorem 4.2, we infer that for all $(t, x) \in [0, T] \times \mathbb{R}^d$ and all $\eta > 0$,

$$|u_\varepsilon(t, x) - u_0(x)| \leq \|\nabla u_0\|_{L^\infty(\mathbb{R}^d)} \left(\frac{c}{\eta} t + \eta \right).$$

The previous estimates holds for every η , and hence, by choosing $\eta = \sqrt{ct}$, we get

$$|u_\varepsilon(t, x) - u_0(x)| \leq 2 \|\nabla u_0\|_{L^\infty(\mathbb{R}^d)} \sqrt{ct}. \quad (58)$$

Eventually, we deduce (51) from (58) by combining the facts that the problem (36) is invariant w.r.t. translations in time, that it admits a unique solution, and that $\|\nabla u_\varepsilon(t, \cdot)\|_{L^\infty(\mathbb{R}^d)} \leq \|\nabla u_0\|_{L^\infty(\mathbb{R}^d)}$ for all $t \in [0, T]$. \square

7. Convergence to the solution of the limit problem. This section is devoted to the proof of the second main result of the paper, Theorem 1.2. Theorem 1.1 establishes an asymptotic link between the rescaled nonlocal curvatures and the anisotropic mean curvature. In what follows, we take advantage of this relationship to deduce locally uniform convergence of the viscosity solutions u_ε of (36) to the viscosity solution u of (37).

To achieve the result, we compare any limit point v of $\{u_\varepsilon\}$ (which Proposition 4 proves to be a relatively compact family) with the viscosity solution u to (37). More precisely, we focus on the respective superlevel sets, and, by using the theory of geometric barriers and their relations with the level-set flows, we establish the inclusions (60) and (61). In turn, these are sufficient to conclude that $v = u$, thanks to the next lemma.

Lemma 7.1. *Let $f, g : \mathbb{R}^d \rightarrow \mathbb{R}$ be two continuous functions such that for all $\lambda \in \mathbb{R}$ there hold*

$$\{x \in \mathbb{R}^d : f(x) > \lambda\} \subseteq \{x \in \mathbb{R}^d : g(x) \geq \lambda\}$$

and

$$\{x \in \mathbb{R}^d : g(x) > \lambda\} \subseteq \{x \in \mathbb{R}^d : f(x) \geq \lambda\}.$$

Then, $f(x) = g(x)$ for all $x \in \mathbb{R}^d$.

Proof. Let $\bar{x} \in \mathbb{R}^d$ and assume that $g(\bar{x}) = \lambda$. Then, for all $\mu > 0$, we get $\bar{x} \in \{x : g(x) > \lambda - \mu\} \subseteq \{x : f(x) \geq \lambda - \mu\}$, which in particular implies $f(\bar{x}) \geq \lambda$. If $f(\bar{x}) > \lambda$, then for some $\mu_0 > 0$, we would get $\bar{x} \in \{x : f(x) > \lambda + \mu_0\} \subseteq \{x : g(x) \geq \lambda + \mu_0 > \lambda\}$, in contradiction with the fact that $g(\bar{x}) = \lambda$. So $f(\bar{x}) = \lambda$. By reversing the role of f and g , we get the conclusion. \square

Let $\lambda \in \mathbb{R}$ and $E_{\varepsilon, \lambda}^\pm(t)$ be the level-set flows associated with the solutions u_ε to (36) defined in (38). We introduce the families $\tilde{E}_\lambda^\pm(t)$, which are the set-theoretic upper limits of $E_{\varepsilon, \lambda}^\pm(t)$:

$$\tilde{E}_\lambda^-(t) := \bigcap_{\varepsilon < 1} \bigcup_{\eta < \varepsilon} E_{\eta, \lambda}^-(t) \quad \text{and} \quad \tilde{E}_\lambda^+(t) := \bigcap_{\varepsilon < 1} \bigcup_{\eta < \varepsilon} E_{\eta, \lambda}^+(t). \quad (59)$$

Remark 7. It is an immediate consequence of the definition that, for any $\bar{\varepsilon} < 1$,

$$\tilde{E}_\lambda^-(t) = \bigcap_{\varepsilon < \bar{\varepsilon}} \bigcup_{\eta < \varepsilon} E_{\eta, \lambda}^-(t) \quad \text{and} \quad \tilde{E}_\lambda^+(t) = \bigcap_{\varepsilon < \bar{\varepsilon}} \bigcup_{\eta < \varepsilon} E_{\eta, \lambda}^+(t).$$

We are ready to discuss the proof of our convergence result:

Proof of Theorem 1.2. We divide the proof in three steps, starting with a preliminary observation. By Proposition 4, we know that the family u_ε is relatively compact in $C([0, T] \times \mathbb{R}^d)$ and, consequently, there exist a subsequence $\{u_{\varepsilon_n}\}$ and a function $v \in C([0, T] \times \mathbb{R}^d)$ such that $u_{\varepsilon_n} \rightarrow v$ locally uniformly as $\varepsilon \rightarrow 0^+$. We remark that the conclusion is achieved if we show that $v = u$. Indeed, since the argument applies to any converging subsequence of $\{u_\varepsilon\}$, it follows that the whole family $\{u_\varepsilon\}$ locally uniformly converges to u , as desired.

From now on we reason on a subsequence that we still denote $\{u_\varepsilon\}$ and that we suppose to be locally uniformly converging to v .

Step 1: we claim that for every $\lambda \in \mathbb{R}$,

$$\{x \in \mathbb{R}^d : v(t, x) > \lambda\} \subseteq \tilde{E}_\lambda^-(t) \subseteq \tilde{E}_\lambda^+(t) \subseteq \{x \in \mathbb{R}^d : v(t, x) \geq \lambda\} \quad (60)$$

with $\tilde{E}_\lambda^\pm(t)$ as in (59).

In this part of the proof we exploit only the pointwise convergence of $\{u_\varepsilon\}$. Without loss of generality, we discuss just the case $\lambda = 0$.

Let us fix $\bar{x} \in \mathbb{R}^d$ such that $v(t, \bar{x}) > 0$, that is, $v(t, \bar{x}) = \mu$ for some $\mu > 0$. Since v is the limit of $\{u_\varepsilon\}$, there exists $\bar{\varepsilon} > 0$ such that

$$u_\varepsilon(t, \bar{x}) \geq \frac{\mu}{2} > 0 \quad \text{for all } \varepsilon < \bar{\varepsilon},$$

and hence $\bar{x} \in \tilde{E}_0^-(t)$. This shows that $\{x \in \mathbb{R}^d : v(t, x) > 0\} \subseteq \tilde{E}_0^-(t)$.

Let us now turn to the inclusion $\tilde{E}_0^+(t) \subseteq \{x \in \mathbb{R}^d : v(t, x) \geq 0\}$. By definition, if $\bar{x} \in \tilde{E}_0^+(t)$, then for all $\varepsilon < 1$ there exists $\eta_\varepsilon < \varepsilon$ such that $u_{\eta_\varepsilon}(t, \bar{x}) \geq 0$. Taking the limit $\varepsilon \rightarrow 0$, we get

$$v(t, \bar{x}) = \lim_{\varepsilon \rightarrow 0} u_{\eta_\varepsilon}(t, \bar{x}) \geq 0.$$

Step 2: we claim that, for all $\lambda \in \mathbb{R}$,

$$\{x \in \mathbb{R}^d : u(t, x) > \lambda\} \subseteq \tilde{E}_\lambda^-(t) \subseteq \tilde{E}_\lambda^+(t) \subseteq \{x \in \mathbb{R}^d : u(t, x) \geq \lambda\} \quad (61)$$

where u is the viscosity solution to (37).

We will firstly show that $\tilde{E}_\lambda^-(t)$ and $\tilde{E}_\lambda^+(t)$ are, respectively, an outer barrier for the family of strict geometric subsolutions and an inner barrier associated with the flow of H_0 . If these assertions hold true, then Theorem 4.7 immediately entails the conclusion, because it states that $\{x \in \mathbb{R}^d : u(t, x) > \lambda\}$ is the minimal outer barrier for the family of strict geometric subsolutions, and that $\{x \in \mathbb{R}^d : u(t, x) \geq \lambda\}$ is the maximal inner barrier for the family of strict geometric supersolutions.

We prove just that $\tilde{E}_0^-(t)$ is an outer barrier for the family of strict geometric subsolutions, since the arguments for $\lambda \neq 0$ and $\tilde{E}_0^+(t)$ are the same.

Let us consider, for some $0 \leq t_0 < t_1 \leq T$, a family of evolving sets $t \mapsto D(t)$ which is a strict geometric subsolution to the anisotropic mean curvature motion when $t \in [t_0, t_1]$. Explicitly, we suppose that there exists $\ell > 0$ such that

$$\partial_t x(t) \cdot \hat{n}_D(t, x(t)) \leq -H_0(\partial D(t), x(t)) - \ell \quad \text{for all } t \in (t_0, t_1] \text{ and } x(t) \in \partial D(t), \quad (62)$$

where \hat{n}_D is the outer unit normal to $D(t)$; we assume as well that

$$D(t_0) \subset \tilde{E}_0^-(t_0). \quad (63)$$

We want to show that $D(t_1) \subset \tilde{E}_0^-(t_1)$.

Recalling definition (59), we get from (63) that for all $\varepsilon < 1$ there exists $\eta_\varepsilon \leq \varepsilon$ such that

$$D(t_0) \subseteq E_{\eta_\varepsilon, 0}^-(t_0). \quad (64)$$

Since for $t \in [t_0, t_1]$ the second fundamental forms of $\partial D(t)$ are uniformly bounded, we can apply Theorem 1.1 and we deduce that

$$\lim_{\varepsilon \rightarrow 0} H_\varepsilon(D(t), x) = H_0(D(t), x) \quad \text{uniformly in } t \in [t_0, t_1] \text{ and } x \in \partial D(t).$$

Consequently, there exists $\bar{\varepsilon} := \bar{\varepsilon}(\ell)$ such that, for all $\varepsilon < \bar{\varepsilon}$,

$$\partial_t x(t) \cdot \hat{n}_D(t, x(t)) \leq -H_\varepsilon(D(t), x(t)) - \frac{\ell}{2} \quad \text{for all } t \in (t_0, t_1] \text{ and } x(t) \in \partial D(t),$$

or, in other words, $t \mapsto D(t)$ is a strict geometric subsolution to all the rescaled problems of parameter $\varepsilon \in (0, \bar{\varepsilon})$. By (64) and Proposition 2, we obtain that for all $\varepsilon < \bar{\varepsilon}$ there exists $\eta_\varepsilon \leq \varepsilon$ such that

$$D(t) \subset E_{\eta_\varepsilon, 0}^-(t) \quad \text{for all } t \in [t_0, t_1].$$

We take advantage of Remark 7 to deduce from the previous inclusion that

$$D(t) \subseteq \tilde{E}_0^-(t) \quad \text{for all } t \in [t_0, t_1].$$

In particular, we conclude that $D(t_1) \subseteq \tilde{E}_0^-(t_1)$, as desired.

Step 3: we conclude $v = u$.

By (60) and (61), we deduce that, for every $\lambda \in \mathbb{R}$ and $t \in [0, T]$,

$$\begin{aligned} \{x \in \mathbb{R}^d : v(t, x) > \lambda\} &\subseteq \{x \in \mathbb{R}^d : u(t, x) \geq \lambda\}, \\ \{x \in \mathbb{R}^d : u(t, x) > \lambda\} &\subseteq \{x \in \mathbb{R}^d : v(t, x) \geq \lambda\}. \end{aligned}$$

The proof is thus accomplished by applying Lemma 7.1. \square

REFERENCES

- [1] N. Abatangelo and E. Valdinoci. A notion of nonlocal curvature. *Numerical Functional Analysis and Optimization*, 35, no 7-9:793–815, 2014.
- [2] O. Alvarez, P. Cardaliaguet, and R. Monneau. Existence and uniqueness for dislocation dynamics with nonnegative velocity. *Interfaces Free Bound.*, 7(4):415–434, 2005.
- [3] O. Alvarez, P. Hoch, Y. Le Bouar, and R. Monneau. Dislocation dynamics: Short-time existence and uniqueness of the solution. *Arch. Ration. Mech. Anal.*, 181(3):449–504, 2006.
- [4] L. Ambrosio. *Geometric evolution problems, distance function and viscosity solutions*, pages 5–93. Springer Berlin Heidelberg, Berlin, Heidelberg, 2000.
- [5] L. Ambrosio, G. De Philippis, and L. Martinazzi. Γ -convergence of nonlocal perimeter functionals. *Manuscripta Math.*, 134(3-4):377–403, 2011.
- [6] G. Barles and C. Georgelin. A simple proof of convergence for an approximation scheme for computing motions by mean curvature. *SIAM J. Numer. Anal.*, 32(2):484–500, 1995.
- [7] G. Barles and O. Ley. Nonlocal first-order Hamilton-Jacobi equations modelling dislocations dynamics. *Commun. Partial Differ. Equations*, 31(8):1191–1208, 2006.
- [8] G. Bellettini. Alcuni risultati sulle minime barriere per movimenti geometrici di insiemi. *Bollettino UMI*, 7:485–512, 1997.
- [9] G. Bellettini and M. Novaga. Comparison results between minimal barriers and viscosity solutions for geometric evolutions. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.*, 26(1):97–131, 1998.
- [10] G. Bellettini and M. Novaga. *Some aspects of De Giorgi's barriers for geometric evolutions*, pages 115–151. Springer Berlin Heidelberg, Berlin, Heidelberg, 2000.
- [11] G. Bellettini and M. Paolini. Some results on minimal barriers in the sense of De Giorgi applied to driven motion by mean curvature. *Rend. Accad. Naz. Sci. XL Mem. Mat. Appl. (5)*, (19):43–67, 1995.
- [12] J. K. Bence, B. Merriman, and S. Osher. *Diffusion generated motion by mean curvature*. Amer. Math. Soc., Providence, RI, 1992.
- [13] J. Berendsen and V. Pagliari. On the asymptotic behaviour of nonlocal perimeters. *ESAIM Control Optim. Calc. Var.*, 25: Paper No. 48, 27, 2019.
- [14] J. Bourgain, H. Brezis, and P. Mironescu. Another look at Sobolev spaces. In *Optimal control and partial differential equations*, pages 439–455. IOS, Amsterdam, 2001.
- [15] L. A. Caffarelli and P. E. Souganidis. Convergence of nonlocal threshold dynamics approximations to front propagation. *Arch. Ration. Mech. Anal.*, 195, no 1:1–23, 2010.
- [16] L.A. Caffarelli and E. Valdinoci. Uniform estimates and limiting arguments for nonlocal minimal surfaces. *Calc. Var. Partial Differential Equations*, 41(1-2):203–240, 2011.
- [17] A. Cesaroni, S. Dipierro, M. Novaga, and E. Valdinoci. Fattening and nonfattening phenomena for planar nonlocal curvature flows. *Math. Ann.*, 375(1-2):687–736, 2019.
- [18] A. Cesaroni, L. De Luca, M. Novaga, and M. Ponsiglione. Stability results for nonlocal geometric evolutions and limit cases for fractional mean curvature flows. *Comm. Partial Differential Equations, to appear*, arxiv preprint <https://arxiv.org/abs/2003.02248>, 2020.
- [19] A. Chambolle, M. Morini, and M. Ponsiglione. Nonlocal curvature flows. *Arch. Ration. Mech. Anal.*, 218(3):1263–1329, 2015.
- [20] A. Chambolle and M. Novaga. Convergence of an algorithm for the anisotropic and crystalline mean curvature flow. *SIAM J. Math. Anal.*, 37(6):1978–1987, 2006.
- [21] A. Chambolle, M. Novaga, and B. Ruffini. Some results on anisotropic fractional mean curvature flows. *Interfaces Free Bound.*, 19(3):393–415, 2017.

- [22] Y.-G. Chen, Y. Giga, and S. Goto. Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations. *J. Differential Geom.*, 33:749–786, 1991.
- [23] E. Cinti, C. Sinestrari, and E. Valdinoci. Neckpinch singularities in fractional mean curvature flows. *Proc. Amer. Math. Soc.*, 146(6):2637–2646.
- [24] F. Da Lio, N. Forcadel, and R. Monneau. Convergence of a non-local eikonal equation to anisotropic mean curvature motion. application to dislocations dynamics. *J. Eur. Math. Soc. (JEMS)*, 10(4):1105–1119, 2008.
- [25] E. De Giorgi. Barriers, boundaries, motion of manifolds, March 18, 1994. Conference held at Dipartimento di Matematica, Univ. of Pavia.
- [26] L.C. Evans. Convergence of an algorithm for mean curvature motion. *Indiana Univ. Math. J.*, 42(2):533–557, 1993.
- [27] N. Forcadel, C. Imbert, and R. Monneau. Homogenization of some particle systems with two-body interactions and of the dislocation dynamics. *DCDS-A*, 23(3):785–826, 2008.
- [28] P. Hajlasz. *Sobolev spaces on metric-measure spaces*, volume 338 of *Contemp. Math.* Amer. Math. Soc., Providence, RI, 2003.
- [29] C. Imbert. Level set approach for fractional mean curvature flows. *Interfaces Free Bound.*, 11(1):153–176, 2009.
- [30] C. Imbert, R. Monneau, and E. Rouy-Mironescu. Homogenization of first order equations with u/ε -periodic Hamiltonians. part ii: application to dislocations dynamics. *Comm. in PDEs*, 33(1):479–516, 2008.
- [31] H. Ishii. A generalization of the Bence, Merriman and Osher algorithm for motion by mean curvature. *Proceedings of the international conference on curvature flows and related topics held in Levico, Italy, June 27-July 2nd, 1994*, pages 111–127, 1995.
- [32] H. Ishii, G.E. Pires, and P. E. Souganidis. Threshold dynamics type approximation schemes for propagating fronts. *J. Math. Soc. Japan*, 51(2):267–308, 1999.
- [33] J. M. Mazon, J. D. Rossi, and J. Toledo. Nonlocal perimeter, curvature and minimal surfaces for measurable sets. *J. Anal. Math.*, 138(1):235–279, 2019.
- [34] V. Pagliari. Halfspaces minimise nonlocal perimeter: a proof via calibrations. *Ann. Mat. Pura Appl.*, 199(4):1685–1696, 2020.
- [35] O. Savin and E. Valdinoci. Γ -convergence for nonlocal phase transitions. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 29(4):479–500, 2012.
- [36] D. Slepčev. Approximation schemes for propagation of fronts with nonlocal velocities and Neumann boundary conditions. *Nonlinear Anal.*, 52:79–115, 2003.

Received xxxx 20xx; revised xxxx 20xx.

E-mail address: annalisa.cesaroni@unipd.it

E-mail address: valerio.pagliari@tuwien.ac.at