



The anti-inflammatory effect of bacterial short chain fatty acids is partially mediated by endocannabinoids

Amrita Vijay, Afroditi Kouraki, Sameer Gohir, James Turnbull, Anthony Kelly, Vicky Chapman, David A Barrett, William J Bultiewicz & Ana M Valdes

To cite this article: Amrita Vijay, Afroditi Kouraki, Sameer Gohir, James Turnbull, Anthony Kelly, Vicky Chapman, David A Barrett, William J Bultiewicz & Ana M Valdes (2021) The anti-inflammatory effect of bacterial short chain fatty acids is partially mediated by endocannabinoids, Gut Microbes, 13:1, 1997559, DOI: [10.1080/19490976.2021.1997559](https://doi.org/10.1080/19490976.2021.1997559)

To link to this article: <https://doi.org/10.1080/19490976.2021.1997559>



© 2021 The Author(s). Published with license by Taylor & Francis Group, LLC.



[View supplementary material](#)



Published online: 17 Nov 2021.



[Submit your article to this journal](#)



Article views: 2718




[View related articles](#)



[View Crossmark data](#)

RESEARCH PAPER



The anti-inflammatory effect of bacterial short chain fatty acids is partially mediated by endocannabinoids

Amrita Vijay^a, Afroditi Kouraki^{a,b}, Sameer Gohir^{a,b}, James Turnbull^{b,c}, Anthony Kelly^{a,b}, Vicky Chapman^{a,c}, David A Barrett^{b,d}, William J Bultiewicz^e, and Ana M Valdes^{b,d}

^aDivision of Rheumatology, Orthopedics and Dermatology, School of Medicine, University of Nottingham, Nottingham, UK; ^bNihr Nottingham Biomedical Research Centre, University of Nottingham, Nottingham, UK; ^cArthritis Research UK Pain Centre, University of Nottingham, Medical School, Queen's Medical Centre, Nottingham, UK; ^dDAB-Centre for Analytical Bioscience, Advanced Materials and Healthcare Technologies Division, School of Pharmacy, University of Nottingham, Nottingham, UK; ^eLowcountry Gastroenterology Associates, Mount Pleasant, South Carolina, USA

ABSTRACT

The endocannabinoid (EC) system has pleiotropic functions in the body. It plays a key role in energy homeostasis and the development of metabolic disorders being a mediator in the relationship between the gut microbiota and host metabolism. In the current study we explore the functional interactions between the endocannabinoid system and the gut microbiome in modulating inflammatory markers. Using data from a 6 week exercise intervention (treatment $n = 38$ control $n = 40$) and a cross sectional validation cohort ($n = 35$), we measured the associations of 2-arachidonoylglycerol (2-AG), anandamide (AEA), *N*-oleoylethanolamine (OEA) and *N*-palmitoylethanolamine (PEA) with gut microbiome composition, gut derived metabolites (SCFAs) and inflammatory markers both cross-sectionally and longitudinally. At baseline AEA and OEA were positively associated with alpha diversity ($\beta(\text{SE}) = .32 (.06), P = .002; .44 (.04), P < .001$) and with SCFA producing bacteria such as *Bifidobacterium* (2-AG $\beta(\text{SE}) = .21 (.10), P < .01$; PEA $\beta(\text{SE}) = .23 (.08), P < .01$), *Coprococcus 3* and *Faecalibacterium* (PEA $\beta(\text{SE}) = .29 (.11), P = .01; .25 (.09), P < .01$) and negatively associated with *Collinsella* (AEA $\beta(\text{SE}) = -.31 (.12), P = .004$). Additionally, we found AEA to be positively associated with SCFA Butyrate ($\beta(\text{SE}) = .34 (.15), P = .01$). AEA, OEA and PEA all increased significantly with the exercise intervention but remained constant in the control group. Changes in AEA correlated with SCFA butyrate and increases in AEA and PEA correlated with decreases in TNF- α and IL-6 statistically mediating one third of the effect of SCFAs on these cytokines. Our data show that the anti-inflammatory effects of SCFAs are partly mediated by the EC system suggesting that there may be other pathways involved in the modulation of the immune system via the gut microbiome.

ARTICLE HISTORY

Received 4 August 2021
Revised 13 October 2021
Accepted 15 October 2021

KEYWORDS

Endocannabinoids; exercise; gut microbes; SCFA; inflammatory markers; intervention



Introduction


Exercise is known to elicit a feeling of euphoria, referred to as a “runner’s high”, which recent studies indicate is the result of activation of the endocannabinoid system.^{1,2} Endocannabinoids (ECs), such as anandamide (AEA), 2-arachidonoylglycerol (2-AG), *N*-palmitoylethanolamine (PEA), and *N*-oleoylethanolamine (OEA) are lipid mediators that bind to specific receptors and elicit cell signaling. The EC system modulates systemic energy metabolism, inflammation, pain, and brain biology³ and is comprised of ECs, its receptors, most notably the G-protein coupled receptors CB1, CB2, and the enzymes that produce and degrade ECs.⁴ The role of this system in modulating

inflammation,⁵ muscle strength^{6,7} and energy metabolism^{8,9} is now widely documented in humans and in other mammals.¹⁰

In addition, there is a vast body of evidence suggesting that the gut microbiome and exercise are interconnected to regulate metabolism and homeostasis, independent of diet.¹¹ Specifically, exercise has been shown in both animal model and human studies to increase the relative abundance of butyrate-producing microbes and thereby increase the production of butyrate, a short chain fatty acid with systemic anti-inflammatory benefits.^{12–17}

Separately, the gut microbiome and the EC system have also been connected to metabolic regulation and homeostasis.¹⁸ For over a decade now it

CONTACT Amrita Vijay  amrita.vijay@nottingham.ac.uk  Division of Rheumatology, Orthopedics and Dermatology, School of Medicine, University of Nottingham, Nottingham, UK

 Supplemental data for this article can be accessed on the [publisher's website](#).

© 2021 The Author(s). Published with license by Taylor & Francis Group, LLC.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

has been known that specific gut microbial strains modulate the expression of cannabinoid and μ -opioid receptors in intestinal cells.¹⁹ Extensive work in animal models has shown that gut microbes also counteract obesity-induced overactivity of the EC system in the mouse colon, with subsequent reduction of gut permeability to lipopolysaccharide

(LPS) (i.e. decreased metabolic endotoxaemia) and increased adipogenesis.^{20,21} Accordingly, prebiotics, probiotics and antibiotics affect the intestinal EC system. These effects of the microbiota appear to be mediated in part by the modulation of EC inactivating enzymes, which also metabolize EC-related mediators with activity at non-cannabinoid receptors.²² Furthermore, dysregulation of the EC system has been connected to digestive disorders such as inflammatory bowel disease, irritable bowel syndrome, as well as obesity.^{23–25} These conditions involve both a dysregulated microbiota (dysbiosis) and altered short chain fatty acids (SCFAs) levels.^{26–28}

To date the mechanisms and the extent to which the anti-inflammatory effects of gut microbial production of (SCFAs) are mediated or induced by changes in the EC system have not been explored. Moreover, the links between changes in specific bacterial strains and EC levels in response to dietary or other interventions are lacking in humans. In this study we have investigated the cross-sectional links between ECs and gut microbiome composition in two cohorts and further investigated the relationship between changes in ECs and gut microbiome in response to an exercise intervention.

Methods

Study population

The longitudinal cohort belonged to community dwelling individuals (Age: >45 y) as part of the iBEATO study.²⁹ The cross-sectional cohort was an independent cohort consisting of healthy individuals aged >18 y. The associations of ECs with microbiome and metabolomic data was carried out cross-sectionally first in the longitudinal cohort and then validated in the independent cohort consisting

of healthy volunteers. Longitudinal analysis was performed using baseline and follow-up data in the webex cohort.²⁹

All participants provided written informed consent. For the longitudinal cohort, ethical approval was obtained from the Research Ethics Committee (ref:18/EM/0154) and the Health Research Authority (protocol no: 18021) and the trial is registered under the clinicaltrials.gov database (NCT03545048). For the cross-sectional (validation cohort), ethical approval was sought by the West Midlands Black Country Research Ethics Committee (18/WM/0066).

Sample collection

Baseline blood, stool, and anthropometric measures (such as, height and weight) were collected in both cohort studies. Blood samples were collected from participants between 8:30am and 10am during each visit. Participants were instructed to come in a fasted state at least since 9 pm the night before (i.e. minimum fasting time was 11.5 hours). Blood samples were collected using Serum Separator Tubes (SST) and were processed within 2–3 hours of collection for separating serum and aliquoted for storage at -80 C until the end of the intervention period.

Metabolomic analysis

Lipidomic measurements

2-AG, AEA, PEA and OEA measurements were extracted from .5 ml serum samples and quantified against a fully extracted calibration line using targeted liquid chromatography tandem mass spectrometry (LC-MS/MS) based on the method described previously.³⁰

Serum short-chain fatty acids

Serum SCFAs in the Webex study cohort were measured using the standardized procedures by Metabolon Inc., Durham, USA.³¹ Serum SCFA in the validation study cohort were measured by the Mass Spectrometry Department, King's College London using in-situ pentafluorobenzoylation of the free acid species, followed by gas chromatography-

negative-chemical-ionization mass spectrometry (GC-NCI-MS) determination of the resulting derivatives as described previously.^{32,33}

Inflammatory markers

Pro and anti-inflammatory serum markers were measured by Affinity Biomarkers, London using the standardized Human Proinflammatory panel 1 assay kit (cat number K151A0H-1), distributed by Meso Scale Discovery as described previously.³³

Gut microbiome sequencing

Fecal sample collection and processing methods were the same in both cohort studies. Fecal samples were collected by the participant at home using previously provided collection kits and frozen immediately at -80°C until further processing. Stool DNA extraction was carried out according to Goodrich et al.³⁴ using 100 mg of the stool sample. There was no homogenization prior to this step. Gut microbiome composition was determined by 16S rRNA gene sequencing carried out as previously described.^{32,33} Briefly, the V4 region of the 16S rRNA gene was amplified using universal primers 355 F (CCAGACTCCTACGGGAGGCAGC) and 806 R (GGACTACHVGGGTWTCTAAT). Amplified DNA was sequenced on the MiSeq platform (Illumina, 300bp paired-end reads). Read filtering and clustering were carried out using the MYcrobiota pipeline. Briefly, chimeric sequences were filtered using the VSEARCH algorithm within Mothur, and reads were clustered into operational taxonomic units (OTUs) using closed-reference clustering against the SILVA database v132 based on a 97% similarity. Diversity metrics (Shannon index observed OTUs and Unweighted UniFrac) were calculated by rarefying the OTU table down to 7000 sequences per sample 50 times and taking the average. These analyses were carried out in QIIME2 (v2018.11).

Gene expression

Barcoded libraries for RNA-seq were prepared with 5ng of RNA using TruSeq Stranded Total RNA HT Sample Prep Kit with Ribo-Zero Gold kit (Illumina) per manufacturer's protocol. Paired-

end sequencing (100 bp \times 2) was performed on HiSeq 4000 sequencers (Illumina) at Genewiz (UK).³⁵

Statistical analysis

All statistical analyses were carried out in R v4.0.3. OTUs with a relative abundance of $<.1\%$ in every sample were removed, and zero inflated relative OTU abundances were inverse normal transformed before further analyses. All analyses were adjusted for age, gender and BMI and multiple testing using false discovery rate (FDR $q < .05$). Linear regressions were first carried out independently in both cohorts to identify significant associations between endocannabinoids, gut microbiota composition, SCFAs and cytokines along with adjusting for covariates such as age, sex, BMI and multiple testing (FDR $q < .05$). The standardized estimates were meta-analyzed to produce a combined effect after adjusting for covariates and multiple testing. Meta-analysis takes the effect size, standard error and sample size into account to give an overall effect from the different groups studied. For the cross-sectional analysis, we used fixed-effects inverse-variance models since our cohorts were homogeneous having adjusted for age, sex and BMI.

Results

The descriptive characteristics of the cohorts are shown in Table 1.

The associations of endocannabinoids with gut microbiome composition, SCFAs and inflammatory cytokines were assessed both cross-sectionally and longitudinally using two independent cohorts that were matched for age and gender as described in Figure 1.

Cross-sectional and reproducible associations of endocannabinoids with gut microbiome composition, SCFAs and inflammatory markers

We first carried a cross-sectional analysis to investigate the associations of endocannabinoid levels with gut microbiome composition, short chain fatty acids and inflammatory markers. We first tested this using the baseline data from the longitudinal cohort wherein we found all four

Table 1.

	Longitudinal cohort (Webex) (N = 78)				Cross-sectional cohort (N = 35)
	Control group (n = 40)		Exercise group (n = 38)		Mean (± SD)
	Baseline Mean (± SD)	Follow-up Mean (± SD)	Baseline Mean (± SD)	Follow-up Mean (± SD)	
<i>Demographics</i>					
Age (y)	67.59 (9.71)		65.32 (10.08)		69.51 (8.77)
Men/Women (%)	8/32 (20/80)		10/28 (26/73)		10/25 (32/68)
BMI (kg/m ²)	32.86 (7.80)		32.58 (7.50)		29.81 (5.12)
					30.31 (7.26)
<i>Endocannabinoids</i>					
2-AG	52.40 (42.12)	49.32 (32.13)	42.50 (28.85)	44.61 (30.48)	41.91 (36.81)
AEA	1.68 (.38)	1.53 (.26)	1.61 (.35)	1.79 (.44)**	.94 (.34)
OEA	7.18 (1.49)	7.15 (2.03)	6.71 (1.87)	7.37 (2.13)**	4.48 (1.78)
PEA	21.99 (33.22)	19.19 (16.99)	26.41 (24.14)	42.07 (31.18)**	26.03 (21.92)
<i>Cytokines</i>					
INF γ	18.46 (17.91)	17.15 (15.63)	14.29 (10.80)	11.25 (7.30)	12.71 (8.12)
IL-10	1.02 (1.69)	.79 (.76)	.55 (.36)	.42 (.31) *	.67 (.53)
IL-13	3.44 (2.49)	2.91 (1.15)	3.26 (1.54)	1.81 (.98)**	3.18 (2.71)
IL-1	.34 (.45)	.22 (.15)	.28 (.27)	.18 (.09)	.41 (.39)
IL-4	.34 (.62)	.32 (.21)	.30 (.57)	.10 (.03)**	.15 (.09)
IL-6	3.16 (2.93)	2.82 (2.74)	2.67 (.86)	2.02 (.71)	1.42 (1.22)
IL-8	41.75 (35.62)	40.04 (35.84)	42.50 (2.21)	32.80 (2.20)	31.79 (17.19)
TNF α	7.05 (6.46)	6.44 (6.03)	7.78 (9.01)	5.91 (3.52)*	3.37 (.89)
<i>SCFAs</i>					
Acetic acid	28.98 (17.89)	26.92 (16.45)	33.61 (18.43)	32.38 (17.12)	43.43 (68.69)
Butyric acid	11.04 (2.53)	10.63 (2.74)	10.86 (2.67)	17.23 (4.10)*	8.39 (.73)
Propionic acid	2.28 (5.98)	2.28 (6.17)	2.42 (6.57)	6.91 (2.03)*	10.13 (.79)
Valeric acid	3.21 (7.33)	2.92 (7.7)	3.20 (4.31)	3.38 (4.63)	1.25 (.52)
Iso-butyric acid	10.39 (2.90)	10.65 (3.09)	9.67 (2.35)	11.79 (3.47)*	10.69 (5.70)
Iso-valeric acid	5.69 (3.17)	5.95 (3.02)	5.55 (2.52)	5.61 (2.30)	7.26 (2.45)

BMI (Body Mass Index); SCFA (Short Chain Fatty Acids)

* $p < .05$; ** $p < .001$. p values are FDR corrected obtained from paired matched t test between baseline and follow-up.

endocannabinoids were positively associated with Shannon diversity, SCFA producing bacteria such as *Bifidobacteria* and *Coprococcus 3* and negatively associated with *Collinsella* and *Escherichia Shigella*, a pathogenic bacterium (Figure 2). We found the endocannabinoids mainly AEA, 2AG and the endocannabinoid like compound OEA to be significantly associated with butyrate (β (SE) = .38 (.10), $P = .01$), propionate (β (SE) = .31 (.08) $P = .01$) and iso-butyrate (β (SE) = .34 (.08), $P = .02$), respectively. Furthermore, EC levels were positively associated with anti-inflammatory markers such as IL-10 but negatively associated with a range of pro-inflammatory cytokines as shown in Figure 2. These findings were then validated in an independent cohort consisting of healthy individuals matched for age and gender wherein we found similar significant associations (supplementary figure S1). We tested reproducibility of the significant associations of ECs with the above mentioned traits by carrying out a meta-analysis by combining the direction of effects observed in

both independent cohorts and found that the associations of endocannabinoids with Shannon diversity, specific OTUs, SCFAs and pro and anti-inflammatory markers were significantly reproducible as shown in Figure 3.

Proportional variance explained by the ECs on the associations of the gut microbiome with inflammatory markers

We then explored the proportional effect of the gut microbiota and ECs on inflammatory markers that were significantly associated (FDR $p < .05$) with both these parameters. We explored these effects by formal mediation where ECs were fitted as mediator of the effect of SCFAs on inflammatory markers. Overall, we found that ECs partially mediated the association between SCFAs and inflammatory markers. AEA mediated 33% ($P < .001$) of the effect of SCFA (butyrate) on TNF α and 27% ($P = .001$) of the effect of SCFA on IL-6. We also tested how much of the effect of ECs on inflammatory markers is mediated by SCFA and found that 56% ($P = .02$)




Figure 1. CONSORT flow diagram.




Figure 2. Matrix plot showing the associations of: (a) OTU abundance and short chain fatty acids with endocannabinoids; and (b) pro and anti-inflammatory markers with endocannabinoids. Associations are based on data from the longitudinal cohort. Squares represent beta coefficients with size and color varying based on size and direction of association. (FDR adjusted $*p < .05$).




Figure 3. Meta-analysis of beta coefficients of with 95% CIs from cross-sectional analysis derived from both cohorts. The figure represents the strongest associations of ECs with (a) Microbiome composition; (b) Short chain fatty acids and (c) Pro and anti-inflammatory cytokines.




Figure 4. Mediator effects of ECs and SCFAs on pro-inflammatory markers: TNF α and IL-6. (a) SCFA: butyrate, iso-butyrate (b) EC: AEA, PEA.

and 48% ($P = .001$) of SCFA mediated the effects on TNF α AND IL-6 respectively, as shown in Figure 4.

Longitudinal association of endocannabinoids with gut microbiome composition, SCFAs and inflammatory markers and response to exercise

We found that levels of AEA, OEA and PEA were all significantly increased in the exercise group but not in the control group. Similarly, we observed significant decreases in proinflammatory cytokines and increases in specific SCFAs in the exercise but not in the control group (Table 1). We then

assessed the correlations between changes in ECs with change in microbiome composition, SCFAs and inflammatory cytokines. The associations in the intervention arm alone did not show changes in ECs to be associated with changes in SCFAs (Supplementary Figure S2) after FDR correction. Therefore, we looked at changes from baseline to follow up using data from both arms looking at the overall change effect of ECs with SCFAs and other markers. We found that amongst the ECs, changes in AEA and OEA were positively associated with gut microbiome diversity ($\beta(\text{SE}) = .32 (.06)$, $P = .002$; $.44 (.04)$, $P < .001$). Increases in ECs were also associated with increased abundance of

SCFA producing bacteria such as *Bifidobacterium* (2-AG β (SE) = .21 (.10), $P < .01$; PEA β (SE) = .23 (.08), $P < .01$), *Coprococcus 3* and *Faecalibacterium* (PEA β (SE) = .29 (.11), $P = .01$; .25 (.09), $P < .01$) and negatively associated with *Collinsella* (AEA β (SE) = -.31 (.12), $P = .004$). Additionally, we found AEA to be positively associated with SCFA Butyrate (β (SE) = .34 (.15), $P = .01$). When we associated ECs with cytokines, we found positive associations of 2AG and OEA with anti-inflammatory markers such as IL-10 and negative associations of most ECs with some of the pro-inflammatory cytokines, TNF α and IL-6 (Figure 5).

Association of endocannabinoids with gene expression levels

The association of endocannabinoids with gene expression levels of specific cannabinoid receptors as well as SCFA receptors from the longitudinal

cohort was tested. We found most ECs to be positively associated with one of the main cannabinoid receptors CNR2 (all $P < .05$). Interestingly, AEA and OEA were found to be positively associated with FFAR2 which is one of the main SCFA fatty acid receptors (Figure 6).

Discussion

In this study we find that approximately one third of the anti-inflammatory downstream effects of SCFAs are statistically mediated by endocannabinoids but approximately two thirds of the effect of SCFAs on cytokines appears not to be related to ECs. Using an exercise intervention, we report that increases in SCFA-producing bacteria and decreases in the pro-inflammatory genus *Collinsella* are correlated with increases in endocannabinoid circulating levels.




Figure 5. Matrix plot showing the associations of (a) change in OTU abundance and short chain fatty acids with change endocannabinoids; (b) change in pro- and anti-inflammatory markers with change endocannabinoids. Squares represent beta coefficients with size and color varying based on size and direction of association. (FDR adjusted $*p < .05$).




Figure 6. Heat map indication positive (light blue) and negative (dark blue) associations of ECs with gene expression levels of specific EC and SCFA receptor proteins. (* $p < .05$).

Previous research has shown increased abundance of *Collinsella* to be strongly associated with increased intake of processed food and low vegetable intake (Menni et al., 2021; Wolf et al., 2019), whereas decreased abundance of *Collinsella* was observed following a Mediterranean diet intervention (Ghosh et al., 2020). Furthermore, it was found that this genus significantly increased the risk of nonalcoholic steatohepatitis, the most serious form of nonalcoholic fatty liver disease whereby inflammation causes liver damage that can progress to cirrhosis (Astbury et al., 2020). Moreover, increased levels of *Collinsella* have been detected in individuals with type 2 diabetes³⁶ and were noted to decrease in a weight loss study where insulin sensitivity improved during the course of the study.³⁷ *Collinsella* was also positively correlated with heightened inflammatory cell count profiles (i.e. lymphocytes).³⁸ The concomitant changes observed in EC levels, *Collinsella* and cytokines suggest that this genus might exert some of its pro-inflammatory effects via modulating the EC system and it is possible that some of the effects of this

genus on insulin resistance may also be linked to ECs given the tight relationship between EC modulation and insulin resistance.³⁹

So far, no previous study examined the potential link between the EC system, exercise, and the gut microbiome. In a study with obese and normal weight women, high moderate to vigorous activity levels were not only higher in normal weight women but were also associated with higher OEA and AEA levels.⁴⁰ Studies have shown that acute physical exercise increases circulating AEA, but not 2-AG, levels in humans.^{41,42} Two studies on elite rugby players showed that depending on diet and BMI the athletes had higher microbial diversity⁴³ and increased fecal SCFA levels⁴⁴ compared to non-active controls, whereas another study found cardiorespiratory fitness associated with higher levels of microbiome diversity, especially butyrate-producing bacteria independent of diet.⁴⁵ While using small numbers of participants, a longitudinal study using 6 weeks of progressive endurance exercise without changing diet in

overweight women found an increase in SCFA-producing bacteria,⁴⁶ independent of age, weight, fat % as well as energy and fiber intake.⁴⁷ Therefore, our data confirm findings from previous studies that have shown that ECs and SCFAs increase with exercise and further reveal the strong correlation between increases in ECs and decreases in pro-inflammatory cytokines. It is possible therefore that improved EC tone induced by exercise may be mediating the shift in the gut microbiota to increased SCFA producers, thereby increasing the SCFA production without a dietary change. This is a hypothesis that needs testing in a controlled experimental setting.

Using formal mediation analysis, we find that ECs are statistically mediating up to a third of the effect of SCFAs on the circulating levels of pro and anti-inflammatory cytokines. We further show that EC levels, specifically AEA and OEA, are positively correlated not just with EC system genes such as cannabinoid receptor 2 (CNR2) but also with higher expression levels of the SCFA receptors *FFAR2* but not with long fatty acid receptor like *FFAR4*. These data complement results from animal model studies showing that ECs can attenuate central and peripheral inflammation, can modulate gut microbiota composition and can reduce markers of gut permeability.⁴⁸ CNR2 deficient mice had higher serum levels of the anti-inflammatory cytokine IL-10 compared to controls following a bacterial lipopolysaccharide challenge.⁴⁹ The anti-inflammatory role of CNR2 was further demonstrated both in a chronic murine animal model and in IBD patients.⁵⁰ *FFAR2* was found to be necessary for the inulin-induced reduction of food-intake and protection against diet-induced obesity.⁵¹ Overall, our data confirm that ECs and SCFAs are crucial modulators of the effects of the gut microbiome on human metabolism and physiology.

Consistent with previous studies,^{52–54} we find anti-inflammatory effects of endocannabinoids, including positive associations of 2AG and OEA with anti-inflammatory markers (IL-10) and negative associations of most ECs with pro-inflammatory cytokines (TNF α and IL-6). At the same time, we show that the anti-inflammatory effects of ECs are partly mediated

by increased levels of SCFAs and specifically butyrate. Elevated levels of butyrate have been shown to decrease mucosal permeability by increasing the secretion of mucins.^{55,56} Furthermore, ECs have emerged as important players in modulating gut permeability via enhancing the production of the tight-junction protein, occludin-1, as well as decreasing the expression of claudin-1, that serves as a paracellular barrier.^{57,58} Although further *in-vitro* and *in-vivo* studies are required to unravel the specific pathways involved, we think that the EC system and the gut microbiome play a role jointly in regulating an inflammatory status.

Our study has a number of strengths. Firstly, the findings from our study reinforce previous knowledge of an interaction between the EC system and the microbiome and add to previous literature by revealing an interaction between ECs, SCFAs and inflammatory system markers. Secondly, we have validated the cross-sectional associations between ECs and bacterial abundances in an independent cohort, and we have used gene expression data to annotate the pathways involved. Lastly, we have assessed the relationship between ECs, SCFAs and cytokines both cross-sectionally and longitudinally and have shown that simple lifestyle interventions such as exercise can modulate inflammatory markers via SCFAs and ECs.

We also note some study limitations. The transcriptomic assay we used did not include probes for *FFAR1* and *FFAR3* nor did it include *CNR1* making our gene expression data only partially informative. The exercise intervention we carried out was performed in individuals with pain in knee osteoarthritis and may not be directly relevant to other groups. However, the associations between ECs, SCFAs and cytokines were validated in a healthy age and sex matched smaller cohort suggesting that our data are generalizable. The findings presented from the formal mediation analysis are purely statistical and do not indicate causality.

In conclusion, in this study we show that circulating levels of ECs are consistently associated with higher levels of SCFAs, with higher microbiome diversity and with lower levels of the pro-inflammatory genus *Collinsella*. We also

show statistically that the anti-inflammatory effects of SCFAs are up to one third mediated by the EC system.

Acknowledgments

We would like to thank Dr Rakesh Jha, Medical School, University of Nottingham for sample preparations as part of the endocannabinoid measurements.

Author contributions

Study conceptualization: A.M.V., AV; Analyzed the data: A.V.; Contributed materials/analysis tools: A.K., J.T., T.K., wrote the manuscript: A.V., A.M.V. Reviewed and edited the manuscript: A.M.V., A.K., J.T., T.K., D.A.B., W.J.B.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

This work was funded by Versus Arthritis [Grant Number: 20777], a Pfizer Global medical grant “Global Awards for Advancing Chronic Pain Research (ADVANCE) 2018” [Grant Number: WI243608], and by the NIHR Nottingham Biomedical Research Centre. AMV and AV are supported by MRC grant MR/V027883/1.

ORCID

Amrita Vijay  <http://orcid.org/0000-0002-9595-5680>

Ana M Valdes  <http://orcid.org/0000-0003-1141-4471>

Data availability statement

The data that support the findings of this study are available at www.ebi.ac.uk/metabolights/MTBLS3203.

References

- Hicks SD, Jacob P, Perez O, Baffuto M, Gagnon Z, Middleton FA. The transcriptional signature of a runner's high. *Med Sci Sports Exerc.* 2019;51(5):970–978. doi:10.1249/MSS.0000000000001865.
- Siebers M, Biedermann SV, Bindila L, Lutz B, Fuss J. Exercise-induced euphoria and anxiolysis do not depend on endogenous opioids in humans. *Psychoneuroendocrinology.* 2021;126:105173. doi:10.1016/j.psyneuen.2021.105173.
- Watkins BA. Diet, endocannabinoids, and health. *Nutr Res.* 2019;70:32–39. doi:10.1016/j.nutres.2019.06.003.
- Iannotti FA, Di Marzo V. The gut microbiome, endocannabinoids and metabolic disorders. *J Endocrinol.* 2021;248(2):R83–97. doi:10.1530/JOE-20-0444.
- Donvito G, Nass SR, Wilkerson JL, Curry ZA, Schurman LD, Kinsey SG, Lichtman AH. The endogenous cannabinoid system: a budding source of targets for treating inflammatory and neuropathic pain. *Neuropsychopharmacology.* 2018;43:52–79.
- Galdino G, Romero T, Silva JFPD, Aguiar D, Paula AMD, Cruz J, Parrella C, Piscitelli F, Duarte I, Di Marzo V, et al. Acute resistance exercise induces antinociception by activation of the endocannabinoid system in rats. *Anesth Analg.* 2014;119(3):702–715. doi:10.1213/ANE.0000000000000340.
- Schönke M, Martinez-Tellez B, Rensen PC. Role of the endocannabinoid system in the regulation of the skeletal muscle response to exercise. *Curr Opin Pharmacol.* 2020;52:52–60. doi:10.1016/j.coph.2020.05.003.
- Rakotoarivelo V, Sihag J, Flamand N. Role of the endocannabinoid system in the adipose tissue with focus on energy metabolism. *Cells [Internet].* 2021;10(6):1279. doi:10.3390/cells10061279.
- Gatta-Cherifi B, Cota D. New insights on the role of the endocannabinoid system in the regulation of energy balance. *Int J Obes.* 2016;40(2):210–219. doi:10.1038/ijo.2015.179.
- Silver RJ. The endocannabinoid system of animals. *Animals (Basel) [Internet].* 2019;9. <http://dx.doi.org/10.3390/ani9090686>
- Mailing LJ, Allen JM, Buford TW, Fields CJ, Woods JA. Exercise and the gut microbiome: a review of the evidence, potential mechanisms, and implications for human health. *Exerc Sport Sci Rev.* 2019;47(2):75–85. doi:10.1249/JES.0000000000000183.
- Matsumoto M, Inoue R, Tsukahara T, Ushida K, Chiji H, Matsubara N, Hara H. Voluntary running exercise alters microbiota composition and increases n-butyrate concentration in the rat cecum. *Biosci Biotechnol Biochem.* 2008;72(2):572–576. doi:10.1271/bbb.70474.
- Kang SS, Jeraldo PR, Kurti A, Miller MEB, Cook MD, Whitlock K, Goldenfeld N, Woods JA, White BA, Chia N, et al. Diet and exercise orthogonally alter the gut microbiome and reveal independent associations with anxiety and cognition. *Mol Neurodegener.* 2014;9(1):36. doi:10.1186/1750-1326-9-36.

14. Evans CC, LePard KJ, Kwak JW, Stancukas MC, Laskowski S, Dougherty J, Moulton L, Glawe A, Wang Y, Leone V, et al. Exercise prevents weight gain and alters the gut microbiota in a mouse model of high fat diet-induced obesity. *PLoS One*. 2014;9(3):e92193. doi:10.1371/journal.pone.0092193.
15. Allen JM, Mailing LJ, Niemiro GM, Moore R, Cook MD, White BA, Holscher HD, Woods JA. Exercise alters gut microbiota composition and function in lean and obese humans. *Med Sci Sports Exerc*. 2018;50(4):747–757. doi:10.1249/MSS.0000000000001495.
16. Bressa C, Bailén-Andrino M, Pérez-Santiago J, González-Soltero R, Pérez M, Montalvo-Lominchar MG, Maté-Muñoz JL, Domínguez R, Moreno D, Larrosa M. Differences in gut microbiota profile between women with active lifestyle and sedentary women. *PLoS One*. 2017;12(2):e0171352. doi:10.1371/journal.pone.0171352.
17. Haen E. The peripheral lymphocyte as clinical model for receptor disturbances: asthmatic diseases. *Bull Eur Physiopathol Respir*. 1987;23:539–541.
18. Gioacchini G, Rossi G, Carnevali O. Host-probiotic interaction: new insight into the role of the endocannabinoid system by in vivo and ex vivo approaches. *Sci Rep*. 2017;7(1):1261. doi:10.1038/s41598-017-01322-1.
19. Rousseaux C, Thuru X, Gelot A, Barnich N, Neut C, Dubuquoy L, Dubuquoy C, Merour E, Geboes K, Chamaillard M, et al. *Lactobacillus acidophilus* modulates intestinal pain and induces opioid and cannabinoid receptors. *Nat Med*. 2007;13(1):35–37. doi:10.1038/nm1521.
20. Mehrpouya-Bahrami P, Chitralla KN, Ganewatta MS, Tang C, Murphy EA, Enos RT, Velazquez KT, McCellan J, Nagarkatti M, Nagarkatti P. Blockade of CB1 cannabinoid receptor alters gut microbiota and attenuates inflammation and diet-induced obesity. *Sci Rep*. 2017;7(1):15645. doi:10.1038/s41598-017-15154-6.
21. Liu J, Bátkai S, Pacher P, Harvey-White J, Wagner JA, Cravatt BF, Gao B, Kunos G. Lipopolysaccharide induces anandamide synthesis in macrophages via CD14/MAPK/phosphoinositide 3-kinase/NF- κ B independently of platelet-activating factor *. *J Biol Chem*. 2003;278(45):45034–45039. doi:10.1074/jbc.M306062200.
22. Geurts L, Everard A, Van Hul M, Essaghir A, Duparc T, Matamoros S, Plovier H, Castel J, Denis RGP, Bergiers M, et al. Adipose tissue NAPE-PLD controls fat mass development by altering the browning process and gut microbiota. *Nat Commun*. 2015;6(1):6495. doi:10.1038/ncomms7495.
23. Izzo AA, Sharkey KA. Cannabinoids and the gut: new developments and emerging concepts. *Pharmacol Ther*. 2010;126(1):21–38. doi:10.1016/j.pharmthera.2009.12.005.
24. Izzo AA, Piscitelli F, Capasso R, Aviello G, Romano B, Borrelli F, Petrosino S, Di Marzo V. Peripheral endocannabinoid dysregulation in obesity: relation to intestinal motility and energy processing induced by food deprivation and re-feeding. *Br J Pharmacol*. 2009;158(2):451–461. doi:10.1111/j.1476-5381.2009.00183.x.
25. Camilleri M, Kolar GJ, Vazquez-Roque MI, Carlson P, Burton DD, Zinsmeister AR. Cannabinoid receptor 1 gene and irritable bowel syndrome: phenotype and quantitative traits. *Am J Physiol Gastrointest Liver Physiol*. 2013;304(5):G553–60. doi:10.1152/ajpgi.00376.2012.
26. Undseth R, Jakobsdottir G, Nyman M, Berstad A, Valeur J. Low serum levels of short-chain fatty acids after lactulose ingestion may indicate impaired colonic fermentation in patients with irritable bowel syndrome. *Clin Exp Gastroenterol*. 2015;8:303–308.
27. Parada Venegas D, De la Fuente MK, Landskron G, González MJ, Quera R, Dijkstra G, Harmsen HJM, Faber KN, Hermoso MA. Short Chain Fatty Acids (SCFAs)-mediated gut epithelial and immune regulation and its relevance for inflammatory bowel diseases. *Front Immunol*. 2019;10:277. doi:10.3389/fimmu.2019.00277.
28. Chambers ES, Preston T, Frost G, Morrison DJ. Role of gut microbiota-generated short-chain fatty acids in metabolic and cardiovascular health. *Curr Nutr Rep*. 2018;7(4):198–206. doi:10.1007/s13668-018-0248-8.
29. Gohir SA, Eek F, Kelly A, Abhishek A, Valdes AM. Effectiveness of internet-based exercises aimed at treating knee osteoarthritis: the iBEAT-OA randomized clinical trial. *JAMA Netw Open*. 2021;4(2):e210012. doi:10.1001/jamanetworkopen.2021.0012.
30. Wong A, Sagar DR, Ortori CA, Kendall DA, Chapman V, Barrett DA. Simultaneous tissue profiling of eicosanoid and endocannabinoid lipid families in a rat model of osteoarthritis. *J Lipid Res*. 2014;55(9):1902–1913. doi:10.1194/jlr.M048694.
31. Bridgewater BR EAM. High resolution mass spectrometry improves data quantity and quality as compared to unit mass resolution mass spectrometry in high-throughput profiling metabolomics. *Metabolomics* [Internet]. 2014;4. <https://www.omicsonline.org/open-access/high-resolution-mass-spectrometry-improves-data-quantity-and-quality-as-compared-to-unit-mass-resolution-mass-spectrometry-in-high-throughput-profiling-metabolomics-2153-0769-4-132.php?aid=32765>
32. Vijay A, Astbury S, Le Roy C, Spector TD, Valdes AM. The prebiotic effects of omega-3 fatty acid supplementation: a six-week randomised intervention trial. *Gut Microbes*. 2021;13(1):1–11. doi:10.1080/19490976.2020.1863133.
33. Vijay A, Astbury S, Panayiotis L, Marques FZ, Spector TD, Menni C, Valdes AM. Dietary interventions reduce traditional and novel cardiovascular risk markers by altering the gut microbiome and their metabolites. *Front Cardiovasc Med*. 2021;8:691564. doi:10.3389/fcvm.2021.691564.

34. Goodrich JK, Waters JL, Poole AC, Sutter JL, Koren O, Blekhman R, Beaumont M, Van Treuren W, Knight R, Bell JT, et al. Human genetics shape the gut microbiome. *Cell*. 2014;159(4):789–799. doi:10.1016/j.cell.2014.09.053.
35. Werling DM, Pochareddy S, Choi J, An J-Y, Sheppard B, Peng M, Li Z, Dastmalchi C, Santpere G, Sousa AMM, et al. Whole-genome and RNA sequencing reveal variation and transcriptomic coordination in the developing human prefrontal cortex. *Cell Rep*. 2020;31(1):107489. doi:10.1016/j.celrep.2020.03.053.
36. Lambeth SM, Carson T, Lowe J, Ramaraj T, Leff JW, Luo L, Bell CJ, Shah VO. Composition, diversity and abundance of gut microbiome in prediabetes and type 2 diabetes. *J Diabetes Obes*. 2015;2:1–7.
37. Frost F, Storck LJ, Kacprowski T, Gärtner S, Rühlemann M, Bang C, Franke A, Völker U, Aghdassi AA, Steveling A, et al. A structured weight loss program increases gut microbiota phylogenetic diversity and reduces levels of *Collinsella* in obese type 2 diabetics: a pilot study. *PLoS One*. 2019;14(7):e0219489. doi:10.1371/journal.pone.0219489.
38. Menni C, Louca P, Berry SE, Vijay A, Astbury S, Leeming ER, Gibson R, Asnicar F, Piccinno G, Wolf J, et al. High intake of vegetables is linked to lower white blood cell profile and the effect is mediated by the gut microbiome. *BMC Med*. 2021;19(1):37. doi:10.1186/s12916-021-01913-w.
39. Vilches-Flores A, Garcia-Luna GM. Endocannabinoid receptors activity modifies insulin secretion in response to glucose exposition. *Diabetes* [Internet]. 2018 [cited 2021 Jul 8];67(Supplement 1):2173–P. doi:10.2337/db18-2173-P
40. Fernández-Aranda F, Sauchelli S, Pastor A, Gonzalez ML, de la Torre R, Granero R, Jiménez-Murcia S, Baños R, Botella C, Fernández-Real JM, et al. Moderate-vigorous physical activity across body mass index in females: moderating effect of endocannabinoids and temperament. *PLoS One*. 2014;9(8):e104534. doi:10.1371/journal.pone.0104534.
41. Heyman E, Gamelin F-X, Goekint M, Piscitelli F, Roelands B, Leclair E, Di Marzo V, Meeusen R. Intense exercise increases circulating endocannabinoid and BDNF levels in humans—possible implications for reward and depression. *Psychoneuroendocrinology*. 2012;37(6):844–851. doi:10.1016/j.psyneuen.2011.09.017.
42. Raichlen DA, Foster AD, Seillier A, Giuffrida A, Gerdeman GL. Exercise-induced endocannabinoid signaling is modulated by intensity. *Eur J Appl Physiol*. 2013;113:869–875. doi:10.1007/s00421-012-2495-5.
43. Clarke SF, Murphy EF, O’Sullivan O, Lucey AJ, Humphreys M, Hogan A, Hayes P, O’Reilly M, Jeffery IB, Wood-Martin R, et al. Exercise and associated dietary extremes impact on gut microbial diversity. *Gut*. 2014;63(12):1913–1920. doi:10.1136/gutjnl-2013-306541.
44. Barton W, Penney NC, Cronin O, Garcia-Perez I, Molloy MG, Holmes E, Shanahan F, Cotter PD, O’Sullivan O. The microbiome of professional athletes differs from that of more sedentary subjects in composition and particularly at the functional metabolic level. *Gut*. 2018;67:625–633.
45. Estaki M, Pither J, Baumeister P, Little JP, Gill SK, Ghosh S, Ahmadi-Vand Z, Marsden KR, Gibson DL. Cardiorespiratory fitness as a predictor of intestinal microbial diversity and distinct metagenomic functions. *Microbiome*. 2016;4(1):42. doi:10.1186/s40168-016-0189-7.
46. Louis P, Flint HJ. Formation of propionate and butyrate by the human colonic microbiota. *Environ Microbiol*. 2017;19(1):29–41. doi:10.1111/1462-2920.13589.
47. Munukka E, Ahtiainen JP, Puigbó P, Jalkanen S, Pahkala K, Keskitalo A, Kujala UM, Pietilä S, Hollmén M, Elo L, et al. Six-week endurance exercise alters gut metagenome that is not reflected in systemic metabolism in over-weight women. *Front Microbiol*. 2018;9:2323. doi:10.3389/fmicb.2018.02323.
48. Cristiano C, Pirozzi C, Coretti L, Cavaliere G, Lama A, Russo R, Lembo F, Mollica MP, Meli R, Calignano A, et al. Palmitoylethanolamide counteracts autistic-like behaviours in BTBR T+tf/J mice: contribution of central and peripheral mechanisms. *Brain Behav Immun*. 2018;74:166–175. doi:10.1016/j.bbi.2018.09.003.
49. Sun X, Cappelletti M, Li Y, Karp CL, Divanovic S, Dey SK. *Cnr2* deficiency confers resistance to inflammation-induced preterm birth in mice. *Endocrinology*. 2014;155(10):4006–4014. doi:10.1210/en.2014-1387.
50. Leinwand KL, Jones AA, Huang RH, Jedlicka P, Kao DJ, de Zoeten EF, Ghosh S, Moaddel R, Wehkamp J, Ostaff MJ, et al. Cannabinoid receptor-2 ameliorates inflammation in murine model of Crohn’s disease. *J Crohns Colitis*. 2017;11(11):1369–1380. doi:10.1093/ecco-jcc/jjx096.
51. Brooks L, Viardot A, Tsakmaki A, Stolarczyk E, Howard JK, Cani PD, Everard A, Sleeth ML, Psychas A, Anastasovskaj J, et al. Fermentable carbohydrate stimulates FFAR2-dependent colonic PYY cell expansion to increase satiety. *Mol Metab*. 2017;6(1):48–60. doi:10.1016/j.molmet.2016.10.011.
52. Kerr DM, Harhen B, Okine BN, Egan LJ, Finn DP, Roche M. The monoacylglycerol lipase inhibitor JZL184 attenuates LPS-induced increases in cytokine expression in the rat frontal cortex and plasma: differential mechanisms of action. *Br J Pharmacol*. 2013;169(4):808–819. doi:10.1111/j.1476-5381.2012.02237.x.

53. Khan RN, Maner-Smith K, Owens AJ, Barbian ME, Jones RM, Naudin RC. At the heart of microbial conversations: endocannabinoids and the microbiome in cardiometabolic risk. *Gut Microbes*. 2021;13(1):1911572. doi:10.1080/19490976.2021.1911572.
54. Naidu PS, Kinsey SG, Guo TL, Cravatt BF, Lichtman AH. Regulation of inflammatory pain by inhibition of fatty acid amide hydrolase. *J Pharmacol Exp Ther*. 2010;334(1):182–190. doi:10.1124/jpet.109.164806.
55. Jung T-H, Park JH, Jeon W-M, Han K-S. Butyrate modulates bacterial adherence on LS174T human colorectal cells by stimulating mucin secretion and MAPK signaling pathway. *Nutr Res Pract*. 2015;9(4):343–349. doi:10.4162/nrp.2015.9.4.343.
56. Schilderink R, Verseijden C, Seppen J, Muncan V, Van Den Brink GR, Lambers TT, van Tol EA, de Jonge WJ. The SCFA butyrate stimulates the epithelial production of retinoic acid via inhibition of epithelial HDAC. *Am J Physiol Gastrointest Liver Physiol*. 2016;310(11):G1138–46. doi:10.1152/ajpgi.00411.2015.
57. Alhamoruni A, Lee AC, Wright KL, Larvin M, O’Sullivan SE. Pharmacological effects of cannabinoids on the Caco-2 cell culture model of intestinal permeability. *J Pharmacol Exp Ther*. 2010;335(1):92–102. doi:10.1124/jpet.110.168237.
58. Muccioli GG, Naslain D, Bäckhed F, Reigstad CS, Lambert DM, Delzenne NM, Cani PD. The endocannabinoid system links gut microbiota to adipogenesis. *Mol Syst Biol*. 2010;6(1):392. doi:10.1038/msb.2010.46.