



## Length-weight relationship of four demersal species from the San Matías Gulf, Patagonia, Argentina

MARÍA EUGENIA LÓPEZ<sup>1,2,3\*§</sup>, NAIR SOLEDAD RUIZ<sup>1,2,3,§</sup>, MARÍA ALEJANDRA ROMERO<sup>1,4,5</sup>, MATÍAS OCAMPO REINALDO<sup>1,4,5</sup> & RAÚL ALBERTO GONZÁLEZ<sup>1,4,5</sup>

<sup>1</sup> Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET)

<sup>2</sup> Instituto de Desarrollo Costero “Dr. H.C. Héctor E. Zaixso” (IDC), Universidad Nacional de la Patagonia San Juan Bosco (UNPSJB)

<sup>3</sup> Centro de Investigación y Transferencia “Golfo San Jorge” (CONICET-UNPSJB-UNPA)

<sup>4</sup> Centro de Investigación Aplicada y Transferencia Tecnológica en Recursos Marinos “Almirante Storni” (CIMAS)

<sup>5</sup> Escuela Superior de Ciencias Marinas (ESCiMar), Universidad Nacional del Comahue (UNCo)

\*Corresponding author: [melopez@live.com.ar](mailto:melopez@live.com.ar)

§ Authors contributed equally and should be considered joint first authors

**Abstract.** This study provides the length-weight relationship (LWR) for the main fish of fishery importance in northern Patagonia, Argentina. The LWR for *Genypterus brasiliensis* and *Acanthistius patachonicus* was not available for the Argentine Continental Shelf, so these results contribute to our knowledge of these species.

**Key words:** fisheries resources, northern Patagonia, trawl fishery.

**Resumen:** Relación largo-peso de cuatro especies demersales del Golfo San Matías, Patagonia, Argentina. Este estudio provee la relación largo peso (RLP) de los recursos pesqueros más importantes de la Patagonia norte, Argentina. Las RLP para *Genypterus brasiliensis* y *Acanthistius patachonicus* no estaban disponibles para la Plataforma Continental Argentina, por lo que estos resultados contribuyen al conocimiento de estas especies.

**Palabras clave:** recursos pesqueros, Patagonia norte, pesquerías de arrastre.

The length-weight relationship (LWR) indicates the average weight at a given size, usually based on the use of a potential equation that shows a relationship between both variables. LWR of fishes is of utmost importance in fishery sciences and management and has been studied since the late 19th century (Froese 1998, Garcia-Ayala *et al.* 2014). It is useful for estimating the biomass of fish stocks when only length data is known, and for understanding the life cycle of fishes since it is related to growth, energy budget, reproductive patterns, and feeding conditions of individuals (Froese 2006). Estimates of mean weight by length class are required for stock assessment models such as virtual population analysis and similar approaches (Pauly 1993). In the San Matías Gulf (SMG), a trawl

demersal fishery has been developed since 1971 which focuses on the Argentine hake *Merluccius hubbsi* Marini, 1933 (Romero *et al.* 2013). Other demersal species such as the pink cusk-eel *Genypterus brasiliensis* Regan, 1903, the Argentine seabass *Acanthistius patachonicus* (Jenyns, 1842), and the Brazilian flathead *Percophis brasiliensis* Quoy and Gaimard, 1825 are also important commercial resources. The Fishery Statistics for the last decade show that these species represent more than 50% of the total catches of trawl fishery that operates in the SMG (Millán 2017). In addition to their economic importance, these species are preyed upon by the system's most conspicuous predators, such as the South American sea lion *Otaria flavescens* (Romero *et al.* 2011), the common

dolphin *Delphinus delphis* and the dusky dolphin *Lagenorhynchus obscurus* (Romero *et al.* 2012), as well as by cartilaginous fish, such as the La Plata skate *Atlantoraja platana* (Coller 2012) and the smallnose fanskate *Symterygia bonaparti* (Estalles *et al.* 2016). Previous studies determined the LWR for these species in other sites of their geographic distribution (Haimovici & Velasco 2000, Frota *et al.* 2004, Vianna *et al.* 2004, Macchi *et al.* 2006, Vaz-Dos-Santos *et al.* 2009, Portela *et al.* 2012, Rico *et al.* 2018) and some of them were made some years ago in the SMG (Rubinich & González 2001, Ocampo Reinaldo 2010). However, knowing this relationship for individuals present in the SMG, would provide basic information to understand growth patterns that could later contribute to the differentiation, assessment and management of fish stocks. In particular, the *M. hubbsi* stock in the SMG constitutes an independent demographic unit (Sardella & Timi 2004, González *et al.* 2007, Machado Schiaffino *et al.* 2011) from the other stocks described in the Argentine Exclusive Economic Zone. Therefore, the aim of this study is to determine the LWR of *M. hubbsi*, *G. brasiliensis*, *A. patachonicus* and *P. brasiliensis* sampled in the SMG. Accurately estimating the growth condition of these species will determine if somatic growth is isometric ( $b=3$ ) or allometric ( $b \neq 3$ ), providing basic knowledge of the intrinsic characteristics of the populations analyzed.

Fish samples were obtained from two sources (Table I): a) the CIMAS (Centro de Investigación

Aplicada y Transferencia Tecnológica en Recursos Marinos “Almirante Storni”, San Antonio Oeste, Río Negro, Argentina) fish database; b) a research survey in 2016. Specimens were first sexed and the total length (TL) was measured to the nearest 1 cm with an ichthyometer, and wet weights (W) were determined to the nearest 0.01 g using an electronic scale. Error distribution (multiplicative or additive) was evaluated for each species' data set using likelihood analysis in order to decide which model (linear regression with logarithmically transformed data or non-linear regression with raw data) was appropriate for estimating the parameters of the relationship (Xiao *et al.* 2011). As the assumption of multiplicative log-normal error was better supported, the parameters of the length-weight relationship ( $W = a LT^b$ , where W: total weight in grams, LT: total length in cm) were estimated by fitting linear regressions to the log-transformed data ( $\log [W] = \log [a] + \log [TL]$ ). In addition, differences between sexes for each species were compared using ANCOVA. In order to verify if calculated  $b$  was significantly different from 3, the Student's t-test was employed. All analyses were carried out using R software (R Core Team 2018).

A total of 3008 specimens belonging to four species were used to estimate the length-weight relationships. There were no differences in LWR between sexes for *M. hubbsi*, *A. patachonicus* and

**Table I.** Origin and size of the sample.

Species	Family	Origin of the sample	N
<i>Merluccius hubbsi</i>	Merlucciidae	Research survey	1812
<i>Acanthistius patachonicus</i>	Serranidae	CIMAS database	762
<i>Genypterus brasiliensis</i>	Ophidiidae	Research survey	150
<i>Percophis brasiliensis</i>	Percophidae	Research survey	294

**Table II.** Descriptive statistics and estimated length-weight relationship parameters for four fish species from the San Matías Gulf. \* b is significantly different from 3 ( $P < 0.05$ ).

Species	Total length range (cm)	Total weight range (g)	Regression parameters (CI 95% <sub>inf</sub> – CI 95% <sub>sup</sub> )		R <sup>2</sup>	P-value ANCOVA
			a	b		
<i>Merluccius hubbsi</i>	19 - 81	41 – 3.485	0.0095 (0.0090 – 0.0100)	2.919* (2.905 – 2.934)	0.99	0.19
<i>Acanthistius patachonicus</i>	21 - 55	200 – 2.696	0.0438 (0.0387 – 0.0496)	2.723* (2.689 – 2.757)	0.97	0.17
<i>Genypterus brasiliensis</i>	30 - 62	106 - 1.616	0.0018 (0.0012 – 0.0028)	3.229 (3.122 – 3.336)	0.96	0.10
<i>Percophis brasiliensis</i>	32 - 71	121 – 1.579	0.0034 (0.0024- 0.0046)	3.067 (2.986 – 3.147)	0.95	0.004

*G. brasiliensis*, so results from pooled individuals are presented for each species in Table II. Although the LWR for *P. brasiliensis* showed significant differences between sexes ( $P < 0.05$ ), a model using pooled individuals is also presented (Table II) for comparison purposes. Based on the results of the t-test, the growth exponent of *G. brasiliensis* and *P. brasiliensis* were not significantly different ( $P > 0.05$ ) from 3, while *M. hubbsi* and *A. patachonicus* were significantly different ( $P < 0.05$ ) from 3.

In this study we provide the length-weight relationship of the main commercial species landed by trawl demersal fisheries in the San Matías Gulf. All species showed an allometric coefficient of around 3, which is the expected value for most fish species (Froese 2006). *M. hubbsi* and *A. patachonicus* present allometric growth. *Acanthistius patachonicus* had the lowest coefficient  $b$  value, with slightly negative allometry. Although this result is similar to that obtained by Rubinich & González (2001) ( $b = 2.7$ ) using samples from the SMG, this study provides the proportionality parameter  $a$  and the range of sizes in which the determined LWR could be used. There are no previous studies that report the complete LWR of *A. patachonicus* for its distribution, so this is the first record. On the other hand, the value of the coefficient  $b$  for *M. hubbsi* indicates negative allometry growth. When individuals collected in 2016 were incorporated in the analysis, similar results were obtained to those of Ocampo Reinaldo (2010) for individuals from SMG ( $b = 3.00$ ). The available information on the allometric coefficient of the LWR for *M. hubbsi* populations located in Argentina ( $b = 2.85$  Macchi *et al.* 2006), Uruguay ( $b \sim 2.7$  Portela *et al.* 2012) and Brazil ( $b = 2.87$  Haimovici & Velasco 2000,  $b = 2.93$  Frota *et al.* 2004) suggests that this species shows growth in length faster than in weight. Nevertheless, Dias *et al.* (2014) when analyzing individuals from Brazil, reported positive asymmetry ( $b = 3.503$ ) for that population, although this could be influenced by the low number of individuals used ( $n = 52$ ) to calculate the LWR.

The LWR calculated for *G. brasiliensis* in the present study represents the first report for the Argentine sea. *G. brasiliensis* present isometric growth and compared with published data, the coefficient  $b$  estimated here is higher than that obtained for the same species from the central Brazilian coast ( $b = 2.76$  Frota *et al.* 2004). This difference could be linked to sample size, since Frota *et al.* (2004) analyzed fewer individuals ( $n =$

41). The growth pattern of *P. brasiliensis* was found to be isometric which indicates that an increment in length and weight occurs at equivalent rate. The value of the allometric coefficient  $b$  estimated for pooled sexes of *P. brasiliensis* is lower than those previously determined for the southwest Atlantic coastal waters (San Román 1974, Rico *et al.* 2018) and the southern coast of Brazil (Vianna *et al.* 2004). This variation could be explained by exogenous factors, such as fish habitat, sample size, and length of specimens examined; and various endogenous factors, such as sex of individuals, stage of sex maturity, and the amount of food contained in the stomach (Froese 2006). Significant differences were found between sexes of *P. brasiliensis* have already been reported for individuals from the Buenos Aires coast (Rico *et al.* 2018). Our results provide essential information for more complex analyses of stock status and the condition of fishes. This study updates the length-weight parameters for some of the main fishery resources in northern Patagonia, and reports for the first time LWR data for Argentine populations of *Gerypterus brasiliensis* and *Acanthistius patachonicus*.

#### Acknowledgments

We acknowledge to the people who collaborated in obtaining and processing the samples. This study was carried out thanks to the logistical support of the Escuela Superior de Ciencias Marinas of the Universidad Nacional del Comahue. We also thank Mariano Malvé and Sofia Sola for correcting the English language and making suggestions to improve the manuscript.

**Funding:** This work was supported by the projects “Elaboración y ejecución de programas para definiciones de planes de manejo de recursos de la pesca en la Plataforma Continental Argentina” STAN CONICET (ST 2872), PICT-2014-1671 and PICT-2015-0534. All authors are supported by CONICET.

#### References

- Coller, N. M. 2012. Biología, ecología y explotación de la raya platana *Atlantoraja platana* (Günther, 1880) (Chondrichthyes, Rajidae) del golfo San Matías. **PhD. Thesis**. Universidad Nacional de la Plata, La plata, Argentina, 179p.
- Días, J. F., Fernandez, W. S. & Schmidt, T. C. S. 2014. Length-weight relationship of 73 fish species caught in the southeastern inner continental shelf region of Brazil. **Latin**

- American Journal of Aquatic Research**, 42(1): 127-136.
- Estalles, M. L., Perier M. R. & Di Giácomo, E. E. 2016. Trophic ecology of the smallnose fanskate *Sympterygia bonapartii* in the San Matías Gulf, northern Patagonia, Argentina. **Ichthyology Research**, 63(2): 207-217.
- Froese, R. 1998. Length–weight relationships for 18 less studied fish species. **Journal of Applied Ichthyology**, 14: 117–118.
- Froese, R. 2006. Cube law, condition factor and weight-length relationships: History, meta-analysis and recommendations. **Journal of Applied Ichthyology**, 22(4): 241–253.
- Frota L. O., Costa P. A. S. & Braga, A. C. 2004. Length-weight relationships of marine fishes from the central Brazilian coast. **ICLARM Quart**, 27(1-2): 20-26.
- Garcia-Ayala, J. R., Brambilla, E. M., Travassos, F. A., Carvalho, E. D. & David, G. S. 2014. Length-weight relationships of 29 fish species from the Tucuru Reservoir (Tocantins/Araguaia Basin, Brazil). **Journal of Applied Ichthyology**, 30: 1092–1095.
- González, R., Narvarte, M. & Caille, G. M. 2007. An assessment of the sustainability of the hake *Merluccius hubbsi* artisanal fishery in San Matías Gulf, Patagonia, Argentina. **Fisheries Research**, 87: 58–67.
- Haimovici, M. & Velasco, G. 2000. Length-Weight Relationship of Marine Fishes from Southern Brazil. Fishbyte Section Editorial Naga. **ICLARM Quart**, 23(1): 19-23.
- Machado Schiaffino, G., Juanes, F. & Garcia-Vazquez, E. 2011. Identifying unique populations in long-dispersal marine species: Gulfs as priority conservation areas. **Biological Conservation**, 144: 330–338.
- Macchi, G. J., Pájaro, M., Militelli, M. I., Radovani N. & Rivas, L. 2006. Influence of size, age and maternal condition on the oocytedry weight of Argentine hake (*Merluccius hubbsi*). **Fisheries Research**, 80: 345–349.
- Millán, D. 2017. **Estadística pesquera oficial de la provincia de Río Negro**. Dirección de Pesca, Ministerio de Agricultura Ganadería y Pesca. Provincia de Río Negro. URL: <https://minagri.rionegro.gov.ar> (Accessed 01/30/2019)
- Ocampo Reinaldo, M. 2010. Evaluación pesquera integral de la merluza común (*Merluccius hubbsi* Marini, 1933) del golfo San Matías y efectos de la explotación de esta especie sobre otros componentes de la trama trófica. **PhD Thesis**. Universidad Nacional de Córdoba, Córdoba, Argentina, 156 p.
- Pauly, D. 1993. Fishbyte Section Editorial Naga. **ICLARM Quart**, 16: 26.
- Portela, J. M., González, M. C. & Bellido, J. M. 2012. Preliminary results of an exploratory fishing targeting deep-water species off Uruguay. **FishBase** – World Wide Web Electronic publication, accessible at <http://www.fishbase.org> (Accessed 7/16/2019).
- R Core Team. 2018. **R: A language and environment for statistical computing**. Vienna: R Foundation for Statistical Computing.
- Rubinich, J. P. & González, R. A. 2001 Edad y crecimiento del mero *Acanthistius brasiliensis* (Pisces, Serranidae) en el Golfo San Matías (Rio Negro, Argentina). **Resúmenes del IX Congreso Latinoamericano de Ciencias del Mar**: 362.
- Rico, M. R., Lagos, A.N., Rodríguez, J. S. & Lorenzo, M. I. 2018. Estado de la pesquería de pez palo (*Percophis brasiliensis*) en el área del Río de la Plata, Zona Común de Pesca Argentino-Uruguaya y aguas jurisdiccionales adyacentes al norte de los 39°S. **Frente Marítimo**, 25: 117-146.
- Romero, M. A., Dans, S., González, R., Svendsen, G., García, N. & Crespo, E. 2011. Solapamiento trófico entre el lobo marino de un pelo *Otaria flavescens* (itálica) y la pesquería de arrastre demersal del Golfo San Matías, Patagonia, Argentina. **Latin american journal of aquatic research**, 39(2): 344-358.
- Romero, M. A., Dans, S. L., García, N. A., Svendsen, G. M., González, R. & Crespo E. A. 2012. Feeding habits of two sympatric dolphin species off North Patagonia, Argentina. **Marine Mammal Science**, 28(2): 364-377.
- Romero, M. A.; Ocampo Reinaldo, M.; Williams, G.; Narvarte, M.; Gagliardini, D. A. & González, R. 2013. Understanding the dynamics of an enclosed trawl demersal fishery in Patagonia (Argentina): A holistic approach combining multiple data sources. **Fisheries Research**, 140: 73-82.
- San Román, N. A. 1974. Crecimiento del pez palo, *Percophis brasiliensis* (Quoy y

- Gaimard, 1824), del Atlántico Sur (Perciformes Trachinoidei). **Physis**, 33(86): 361-369.
- Sardella, N. & Timi, J. 2004. Parasites of Argentine hake in the Argentine Sea: population and infracommunity structure as evidence for host stock discrimination. **Journal of Fish Biology**, 65: 1472–1488.
- Vaz-Dos-Santos, A. M., Rossi-Wongtschowski, C. L. D. B. & Figueiredo, J. L. 2009. Parâmetros da reprodução e relação comprimento-peso da merluza *Merluccius hubbsi* (Teleostei: Merlucciidae, Estoque Sudeste Brasileiro (21°S–29°S), Ano 2004. **Bolletim do Instituto de Pesca**, 35(1): 1–16.
- Vianna, M., Costa, F. E. S. & Ferreira, C. N. 2004. Length-weight relationship of fish caught as by-catch by shrimp fishery in the Southeastern coast of Brazil. **Bolletim do Instituto de Pesca**, 30: 81–85.
- Xiao, X., White, E. P., Hooten, M. B., & Durham, S. L. 2011. On the use of log-transformation vs. nonlinear regression for analyzing biological power laws. **Ecology**, 92: 1887–1894.

Received: June 2019

Accepted: September 2019

Published: January 2020