

R. Crescimbeni, S. Hartzstein and O. Salinas

*Two-weighted inequalities for the fractional integral
associated to the Schrödinger operator*

MATHEMATICAL INEQUALITIES & APPLICATIONS

AIMS AND SCOPE

Mathematical Inequalities & Applications (MIA, Math. Inequal. Appl.) brings together original research papers in all areas of mathematics, provided they are concerned with inequalities or their role. From time to time MIA will publish invited survey articles. Short notes with interesting results or open problems will also be accepted.

MIA is published quarterly, in January, April, July and October.

SUBMISSION

Manuscripts should be submitted electronically in PostScript or Adobe Acrobat PDF format to the MIA Editorial Office through MIA page: www.ele-math.com or by e-mail mia@ele-math.com

The author who submitted the article for publication will be denoted as a corresponding author. He/She manages all communication and correspondence with the MIA regarding the article. Authors may indicate a member of the Editorial Board whom they consider appropriate for the article. However, assignment to that particular editor is not assured.

In order to facilitate refereeing, copies of those papers (whether by the author or someone else) which are essential and referred to in the manuscript but are not conveniently accessible, should be enclosed.

COPYRIGHT

The acceptance of the article automatically implies the copyright transfer to MIA. Manuscripts are accepted for review with the understanding that the same work has not been published (except in the form of an abstract), that it is not under consideration for publication elsewhere, that it will not be submitted to another journal, and that its submission for publication has been approved by all the authors.

PREPARATION OF MANUSCRIPT

Manuscripts should be written in English.

The publisher strongly encourages submission of manuscripts written in TeX or one of its variants LaTeX, AMSTeX or AMSLaTeX. On acceptance of the paper, authors will be asked to send final version of file(s) to the Editorial Office.

TITLE PAGE

A title page should include the title of the article, author's name (no degrees) author's affiliation, e-mail addresses, mailing address of the corresponding author and running head less than 60 characters. Authors should be listed alphabetically.

ABSTRACT, KEY WORDS, SUBJECT CLASSIFICATION

The manuscript must be accompanied by a brief abstract, no longer than 100-150 words. It should make minimal use of mathematical symbols and displayed formulas. Mathematics Subject Classification (2000) with primary (and secondary) subject classification codes and a list of 4-5 key words must be given.

FIGURES

Figures should be prepared in a digital form suitable for direct reproduction, grey scale only, preferably at resolution of 300 dpi or higher, sized to fit a journal page, and in EPS, TIFF or JPEG format. Figures should have a short descriptive title and should be numbered consecutively in order of appearance in the text. The position of each figure in the text should be indicated.

REFERENCES

Bibliographic references should be listed alphabetically at the end of the article. The author should consult Mathematical Reviews for the standard abbreviations of journal names.

PROOFS

The author will receive proofs only once. Late return of the proofs will delay the article to a later issue.

PAGE CHARGES, OFFPRINTS

There are no page charges. Authors will receive PDF file of the printed article free of charge. Additional offprints may be ordered from 'MIA' prior to publication.

FORTHCOMING PAPERS

Papers accepted and prepared for publication will appear in the forthcoming section of Journal Web page. They are identical in form as final printed papers, except volume, issue and page numbers.

MIA is published by Publishing House **ELEMENT**, Zagreb, Croatia.

The publication of this journal is not supported by Ministry of science, education and sports, Croatia

All correspondence and subscription orders should be addressed to the Editorial Office:

Mathematical Inequalities & Applications
Editorial Office
Menceticeva 2, 10000 Zagreb, Croatia
www.ele-math.com
e-mail: mia@ele-math.com
Fax: +385 1 6008799

TWO-WEIGHTED INEQUALITIES FOR THE FRACTIONAL INTEGRAL ASSOCIATED TO THE SCHRÖDINGER OPERATOR

R. CRESCIMBENI, S. HARTZSTEIN AND O. SALINAS

(Communicated by L. Pick)

Abstract. In this article we prove that the fractional integral operator associated to the Schrödinger second order differential operator $\mathcal{L}^{-\alpha/2} = (-\Delta + V)^{-\alpha/2}$ maps with continuity weak Lebesgue space $L^{p,\infty}(v)$ into weighted Campanato-Hölder type spaces $BMO_{\mathcal{L}}^{\beta}(w)$, thus improving regularity under appropriate conditions on the pair of weights (v, w) and the parameters p , α and β . We also prove the continuous mapping from $BMO_{\mathcal{L}}^{\beta}(v)$ to $BMO_{\mathcal{L}}^{\gamma}(w)$ for adequate pair of weights. Our results improve those known for the same weight in both sides of the inequality and they also enlarge the families of weights known for the classical fractional integral associated to the Laplacian operator $\mathcal{L} = -\Delta$.

1. Introduction

Regularity estimates of solutions of second order differential operators are central in the study of partial differential equations. Sometimes these results are closely related to regularity estimates for negative powers of those operators. Keeping in mind this fact in this paper we focus our attention on estimates on Campanato-Hölder type spaces of fractional integrals -negative powers- of the Schrödinger differential operator

$$\mathcal{L} = -\Delta + V,$$

on \mathbb{R}^d with $d \geq 3$, where the potential $V \geq 0$ belongs to a reverse Hölder class RH_q , for some exponent $q > \frac{d}{2}$, as defined in (1.4). For a deeper insight in this direction see [14, 24, 25].

At this point we must recall that the Hölder- α continuous space of functions f such that $\|f\|_{\mathcal{C}^{\alpha}} = \sup_{x \neq y} \frac{|f(x) - f(y)|}{|x - y|^{\alpha}} < \infty$, $0 < \alpha \leq 1$, can be identified with the Campanato space BMO^{α} defined by the seminorm $\|f\|_{BMO^{\alpha}} = \sup_B \frac{1}{|B|^{1+\alpha/\lambda}} \int_B |f(x) - f_B| dx$, where the supremum is taken over all the balls $B \in \mathbb{R}^d$ and f_B is the mean value of f on B , see for example [38, 22, 26]. In the Schrödinger setting analogous result was obtained in [2] by identifying a Campanato-type space $BMO_{\mathcal{L}}^{\alpha}$ with certain type

Mathematics subject classification (2010): 42B35, 35J10.

Keywords and phrases: Fractional integral, weights, Schrödinger, *BMO*, Lipschitz.

The first author was supported by Depto. de Matemática, Fac. de Economía y Administración, Univ. Nac. del Comahue, Argentina. The other two authors were partially supported by Instituto de Matemática Aplicada del Litoral (CONICET) and Depto. de Matemática, Fac. de Ingeniería Química, Univ. Nac. del Litoral, Argentina..

Hölder- α continuous space, see next section. This identification will be the key tool to interpret the information given by our results in terms of regularity.

Let us recall that negative powers of the Schrödinger operator can be expressed in terms of the heat diffusion semigroup generated by \mathcal{L} , $e^{-t\mathcal{L}}$, as

$$\mathcal{I}_\alpha f(x) = \mathcal{L}^{-\alpha/2} f(x) = \int_0^\infty e^{-t\mathcal{L}} f(x) t^{\alpha/2} \frac{dt}{t}, \quad \alpha > 0. \tag{1.1}$$

The above operator is also named (Schrödinger) fractional integral operator of order α . When $V = 0$ then $\mathcal{L} = -\Delta$ is the Laplacian operator and we have the classical fractional integral I_α .

For each $t > 0$ the operator $e^{-t\mathcal{L}}$ is an integral operator with kernel $k_t(x, y)$ having a better behaviour far away from the diagonal $\{(x, x), x \in \mathbb{R}^d\}$ than the heat diffusion kernel $\frac{1}{(4\pi t)^{d/2}} e^{-|x-y|^2/4t}$ associated to $-\Delta$, see Lemma 2.1 below and [12], [13] and [23]. It follows from this property that $\mathcal{I}_\alpha f$ is finite a.e. for $f \in L^p$ with $p \geq 1$.

When applied on L^p -spaces the value $p = d/\alpha$ constitutes a breaking-point for the classical fractional integral I_α . More clearly, for $p < d/\alpha$ it satisfies a (p, q) -norm inequality with $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{d}$, see [18, 19, 37, 21], but for $p \geq d/\alpha$ I_α instead shows regularity properties of the form

$$\frac{1}{|B|^{1+\frac{\alpha}{d}-\frac{1}{p}}} \int_B |I_\alpha f(x) - c_{B,f}| dx \leq C \|f\|_p \tag{1.2}$$

for $f \in L^p$, any ball B and some constant $c_{B,f}$. In other words $I_\alpha f$ belongs to the Campanato space BMO^β with $\beta = \frac{\alpha}{d} - \frac{1}{p}$, see for example [38, 22, 29]. In particular, when $p = d/\alpha$ the arrival space is $BMO^0 = BMO$, the well known space of bounded mean oscillation functions of John and Nirenberg.

The Schrödinger fractional integral \mathcal{I}_α behaves similarly to I_α when $p < d/\alpha$ in the sense that it also maps L^p in L^q with $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{d}$ but for $p = d/\alpha$ the behavior of \mathcal{I}_α is better since it maps $L^{d/\alpha}$ into a space, denoted $BMO_{\mathcal{L}}$, which is in fact smaller than the classical BMO -space, see [10]. Finally for $p > d/\alpha$ the operator \mathcal{I}_α maps with continuity L^p into a Campanato-type space $BMO_{\mathcal{L}}^\beta$, see [2, 3, 24]. This space is shown to be the dual of the H^p -space introduced in [11] and [13], as it can be easily checked from the atomic decomposition given there. For definition and properties of weighted H^p -spaces see [1].

Going back again to the classical setting, Harboure, Viviani and Salinas in [17] obtained a more general estimate than (1.2) with I_α defined on a weak weighted L^p -space. That is

$$\frac{1}{v(B)|B|^{\frac{\alpha}{d}-\frac{1}{p}}} \int_B |I_\alpha f(x) - c_{B,f}| dx \leq C \left[\frac{f}{v} \right]_p \tag{1.3}$$

for an adequate class of weights $H(\alpha, p)$.

We recall that the weak weighted $L^{p,\infty}(v)$, $p > 1$, is the space of measurable functions f such that $[\frac{f}{v}]_p = (\sup_{t>0} t^p |\{x : \frac{|f(x)|}{v(x)} > t\}|)^{\frac{1}{p}} < \infty$ where v is a measurable

non-negative function and that the expression on the left side of (1.3) represents a semi-norm in a weighted $BMO^{\alpha-d/p}(v)$, see also [35] and references therein.

The class of weights $H(\alpha, p)$ introduced in [17] was later used in [2] to derive for the Schrödinger fractional integral \mathcal{I}_α estimates, one of which is of the type of (1.3), that show the continuity of the operator from $L^{p,\infty}(v)$ into a weighted Campanato-type space $BMO_{\mathcal{L}}^\beta(v)$. However, since the kernel of \mathcal{I}_α behaves away from the diagonal better than the kernel of the classical fractional integral, it is natural to wonder if there exists a wider class of weights than those used in [2] from which the same continuity result can be deduced.

The first kind of estimates in the present paper gives a positive answer to this question. Moreover, it deals with the two-weight version of the boundedness results obtained in [2].

Our two-weighted results involve hypothesis based on a “power bump” property. This type of conditions already appeared in several papers dealing with two-weighted inequalities, see for instance [32, 15]. Even a weaker “log bump” condition also appeared, for example, in [7, 8, 27, 9] and references therein. For two-weighted inequalities for classical potential operators see for instance [5, 31, 30, 6, 34, 33, 28], and for the Schrödinger fractional integral and maximal operators associated see [20].

The purpose of those hypothesis is to get simpler conditions on the weights by avoiding extra assumptions or conditions involving the operators under consideration.

In order to introduce our main results we turn our attention to the Schrödinger operator $\mathcal{L} = -\Delta + V$. We say that the function V belongs to a reverse Hölder class of order q denoted by RH_q for some $q > \frac{d}{2}$ if

$$\left(\frac{1}{|B|} \int_B V(y)^q dy\right)^{\frac{1}{q}} \leq \frac{C}{|B|} \int_B V(y) dy \tag{1.4}$$

for any ball $B \subset \mathbb{R}^d$. In the sequel we denote $q_0 = \sup\{q : V \in RH_q\}$ and $\delta_0 = \min(1, 2 - \frac{d}{q_0})$.

A “critical radii” function associated to V is defined by

$$\rho(x) = \sup \left\{ r > 0 : \frac{1}{r^{d-2}} \int_{B(x,r)} V \leq 1 \right\}, \quad x \in \mathbb{R}^d. \tag{1.5}$$

Such function is finite for all $x \in \mathbb{R}^d$ and plays an important role in the description of the spaces and, hence, in the inequalities related to regularity of the operators acting on or arriving at these spaces associated to \mathcal{L} , see [10, 12, 13, 36].

We also denote by L_{loc}^1 the set of locally integrable functions of \mathbb{R}^d . By a weight we mean a locally integrable function $w > 0$ a.e. and along this work we denote $w(E) = \int_E w(x)dx$ for any measurable subset $E \in \mathbb{R}^d$.

Given $\eta \geq 1$, a weight w belongs to the class D_η , $w \in D_\eta$, if there exists a constant C such that $w(tB) \leq Ct^{d\eta} w(B)$ for any ball $B \subset \mathbb{R}^d$ and $t \geq 1$. It is easy to see that a weight w belongs to $D = \cup_{\eta \geq 1} D_\eta$ if and only if it satisfies the doubling condition

$$w(2B) \leq Cw(B) \text{ for any ball } B \tag{1.6}$$

and some constant C .

The classes of weights defined in this work are naturally associated to the decay of the kernel of \mathcal{I}_α and the function “critical radius” ρ related. There is some connection between these classes and a two-weighted version of the classes $A_p^{\rho, \theta}$, introduced in [4] in relation with L^p -norm inequalities for several operators. As usual p' denotes the Hölder’s conjugate exponent of p .

DEFINITION 1.1. Given $1 < p \leq \infty$, δ and λ real numbers and $N_0 \geq 0$, the pair of weights (v, w) belongs to $\mathcal{S}(p, \delta, \lambda, N_0)$ if there exist $C > 0$ such that

$$\left(\frac{v^{p'}(\theta B)}{|\theta B|^{1-\frac{p'\lambda}{d}}} \right)^{\frac{1}{p'}} \leq C \left(1 + \frac{\theta R}{\rho(x)} \right)^{N_0} \frac{w(B)}{|B|^{1-\frac{\delta}{d}}}$$

for all $\theta \geq 1$, any ball $B = B(x, R)$ and $\theta B = B(x, \theta R)$. The above inequality can be rephrased as

$$\left(v^{p'}(\theta B) \right)^{\frac{1}{p'}} \leq C \left(1 + \frac{\theta R}{\rho(x)} \right)^{N_0} \theta^{\frac{d}{p'}-\lambda} \frac{w(B)}{|B|^{\frac{1}{p}-\frac{\delta-\lambda}{d}}}.$$

The class $\mathcal{S}(1, \delta, \lambda, N_0)$ is defined by

$$\sup_{\theta B} v \leq C \left(1 + \frac{\theta R}{\rho(x)} \right)^{N_0} \theta^{-\lambda} \frac{w(B)}{|B|^{1-\frac{\delta-\lambda}{d}}}. \tag{1.7}$$

We also define $\mathcal{S}(p, \delta, \lambda) = \bigcup_{N_0 \geq 0} \mathcal{S}(p, \delta, \lambda, N_0)$. If (v, v) belongs to $\mathcal{S}(p, \delta, \lambda)$ we simply say that $v \in \mathcal{S}(p, \delta, \lambda)$. Our first result is

THEOREM 1.1. Let $1 < p < \infty$, $\alpha > 0$ and $\alpha - \frac{d}{p} - \lambda < \delta_0$. If w is doubling (see (1.6)), $(v, w) \in \mathcal{S}((p'r)', \delta, \lambda)$ for some $r > 1$ and $\lambda - \delta \leq \alpha - \frac{d}{p}$ then there exists a positive constant C such that for all $f \in L^{p,\infty}(v)$

$$\frac{1}{w(B)} \int_B |\mathcal{I}_\alpha f(x) - c_B| dx \leq CR^{\alpha-\frac{d}{p}+\delta-\lambda} \left[\frac{f}{v} \right]_p, \quad R < \rho(x_B) \tag{1.8}$$

for any (subcritical) ball $B = B(x_B, R)$, and some positive constant c_B and

$$\frac{1}{w(B)} \int_B |\mathcal{I}_\alpha f(x)| dx \leq C\rho(x_B)^{\alpha-\frac{d}{p}+\delta-\lambda} \left[\frac{f}{v} \right]_p \tag{1.9}$$

for any critical ball $B = B(x_B, \rho(x_B))$.

In the one-weighted situation, the following corollary of Theorem 1.1 improves the result in [2] since the inequalities therein are deduced for a wider class of weights.

COROLLARY 1.1. Let $1 < p < \infty$, $\frac{d}{p} \leq \alpha$ and $\alpha - \frac{d}{p} - \lambda < \delta_0$. If v is doubling and belongs to $\mathcal{S}(p, \lambda, \lambda)$ then there exists a constant C such that for all $f \in L^{p,\infty}(v)$

$$\frac{1}{v(B)} \int_B |\mathcal{I}_\alpha f(x) - c_B| dx \leq CR^{\alpha-\frac{d}{p}} \left[\frac{f}{v} \right]_p, \tag{1.10}$$

for $B = B(x_B, R)$ with $R < \rho(x_B)$, and some positive constant c_B , and

$$\frac{1}{v(B)} \int_B |\mathcal{I}_\alpha f(x)| dx \leq C \rho(x_B)^{\alpha - \frac{d}{p}} \left[\frac{f}{v} \right]_p \tag{1.11}$$

for any critical ball $B = B(x_B, \rho(x_B))$.

Inequalities (1.8) and (1.9) indicate that the arrival space for \mathcal{I}_α is the following weighted Campanato-type space associated to the Schrödinger operator which was introduced in [2].

DEFINITION 1.2. Given a weight w and $\beta \geq 0$ the space $BMO_{\mathcal{L}}^\beta(w)$ is the set of functions f in L^1_{loc} satisfying for any ball $B = B(x, r)$, with $x \in \mathbb{R}^d$ and $r > 0$,

$$\frac{1}{w(B)} \int_B |f - f_B| \leq C |B|^{\frac{\beta}{d}}, \quad \text{with } f_B = \frac{1}{|B|} \int_B f, \tag{1.12}$$

and

$$\frac{1}{w(B)} \int_B |f| \leq C |B|^{\frac{\beta}{d}}, \quad \text{if } r \geq \rho(x). \tag{1.13}$$

Since (1.13) implies (1.12) for $r \geq \rho(x)$ then it is enough to consider (1.12) only for radius $r < \rho(x)$. The constants in (1.12) and (1.13) are independent of the choice of B but may depend on f . A norm (up to an identification of functions differing by a constant) in the space $BMO_{\mathcal{L}}^\beta(w)$ is given by the infima of the constants C satisfying (1.12) and (1.13). As in the classical case, the mean value f_B in (1.12) may be replaced by any positive constant c_B depending only on the ball.

In view of the above definition we are able to rephrase Theorem 1.1 and Corollary 1.1 in terms of a continuous mapping. That is,

$$\mathcal{I}_\alpha : L^{p, \infty}(v) \rightarrow BMO_{\mathcal{L}}^{\alpha - \frac{d}{p} + \delta - \lambda}(w)$$

and

$$\mathcal{I}_\alpha : L^{p, \infty}(v) \rightarrow BMO_{\mathcal{L}}^{\alpha - \frac{d}{p}}(v),$$

with continuity.

In our next theorem we obtain pointwise regularity estimates for $\mathcal{I}_\alpha f$ whenever $f \in BMO_{\mathcal{L}}^\beta(v)$ with β small. Those kind of inequalities characterize weighted Campanato-type spaces of order smaller than 1. Let us consider the function W_β defined by

$$W_\beta(x, r) = \int_{B(x, r)} \frac{w(z)}{|z - x|^{d - \beta}} dz \tag{1.14}$$

for $x \in \mathbb{R}^d$, $r > 0$, $\beta > 0$ and $w \in L^1_{loc}$.

THEOREM 1.2. *Let $\alpha > 0$, $\beta \geq 0$, $\beta > \lambda$, $\beta - \lambda + \alpha < \delta_0$ and δ and λ be real numbers. If $0 < \beta + \alpha + \delta - \lambda < 1$, w is a doubling weight (see ((1.6))) and $(v, w) \in \mathcal{S}(\infty, \delta, \lambda)$ then there exists a constant C such that for all $f \in BMO_{\mathcal{L}}^{\beta}(v)$*

$$|\mathcal{I}_{\alpha}f(x) - \mathcal{I}_{\alpha}f(y)| \leq C\|f\|_{BMO_{\mathcal{L}}^{\beta}(v)}(W_{\beta+\alpha+\delta-\lambda}(x, |x-y|) + W_{\beta+\alpha+\delta-\lambda}(y, |x-y|)) \tag{1.15}$$

if $|x - y| < \rho(x)$ and

$$\frac{1}{w(B(x, \rho(x)))} \int_{B(x, \rho(x))} |\mathcal{I}_{\alpha}f(x)| dx \leq C\|f\|_{BMO_{\mathcal{L}}^{\beta}(v)} \rho(x)^{\beta+\alpha+\delta-\lambda} \tag{1.16}$$

for all $x \in \mathbb{R}^d$ and $r > 0$.

In the case $v = w$ and $\delta - \lambda = 0$ in Theorem 1.2 we recover Theorem 2 in [2].

Notice that the function $W_{\beta}(x, r)$ defined in (1.14) is finite for all $r > 0$ for almost every $x \in \mathbb{R}^d$. It also is increasing as a function of r for any fixed x and if w is doubling (see (1.6)) then W_{β} is also doubling in the same sense.

The function W_{β} was first considered in [17] and later also used in [2] to define a Lipschitz-Hölder-type space associated to \mathcal{L} . That is, $\Lambda_{\mathcal{L}}^{\beta}(w)$ is the set of functions f such that

$$|f(x) - f(y)| \leq C(W_{\beta}(x, |x-y|) + W_{\beta}(y, |x-y|)) \tag{1.17}$$

and

$$|f(x)| \leq CW_{\beta}(x, \rho(x)). \tag{1.18}$$

for almost all x and y in \mathbb{R}^d . A (quasi) norm is defined on $\Lambda_{\mathcal{L}}^{\beta}(w)$ by taking the maximum of the two infima of the constants satisfying (1.17) and (1.18) respectively.

PROPOSITION 1.1. ([2]) *If $0 < \beta < 1$ and w satisfies the doubling condition (1.6) then $\Lambda_{\mathcal{L}}^{\beta}(w) = BMO_{\mathcal{L}}^{\beta}(w)$ and their norms are equivalent.*

REMARK 1.1. When proving Proposition 1.1, the authors showed that (1.13) jointly with (1.17) imply (1.18). That is if f satisfy (1.13) and (1.17) simultaneously then $f \in \Lambda_{\mathcal{L}}^{\beta}(w)$. The fact that w is doubling is essential to obtain the identification between Lipschitz-Hölder and Campanato type spaces in Proposition 1.1. Hence Theorem 1.2 can be rephrased by saying that

$$\mathcal{I}_{\alpha} : BMO_{\mathcal{L}}^{\beta}(v) \rightarrow \Lambda_{\mathcal{L}}^{\beta+\alpha+\delta-\lambda}(w) = BMO_{\mathcal{L}}^{\beta+\alpha+\delta-\lambda}(w)$$

is continuous, under the hypothesis of that theorem. In the one-weight situation, we obtain

$$\mathcal{I}_{\alpha} : BMO_{\mathcal{L}}^{\beta}(v) \rightarrow \Lambda_{\mathcal{L}}^{\beta+\alpha}(v) = BMO_{\mathcal{L}}^{\beta+\alpha}(v).$$

This article is organized as follows. In Section 2 we give the preliminary definitions and results. In Section 3 we study properties of the class of weights given in Definition 1.1 and show some examples. Section 4 and Section 5 are devoted to the main lemmas and the proofs of Theorems 1.1 and 1.2 respectively.

Throughout this work, we denote by C a constant that may change from one occurrence to other.

2. Preliminaries

Some important properties of the critical rati ρ given in (1.5) are shown in the following propositions.

PROPOSITION 2.1. ([36]) *There exist C and $j_0 \geq 1$ such that*

$$C^{-1}\rho(x)\left(1 + \frac{|x-y|}{\rho(x)}\right)^{-j_0} \leq \rho(y) \leq C\rho(x)\left(1 + \frac{|x-y|}{\rho(x)}\right)^{\frac{j_0}{j_0+1}}$$

for all $x, y \in \mathbb{R}^d$.

PROPOSITION 2.2. ([11]) *There exists a sequence of points $\{x_k\}_{k=1}^\infty$ in \mathbb{R}^d , so that the family $B_k = B(x_k, \rho(x_k))$, $k \geq 1$, satisfies*

1. $\cup_k B_k = \mathbb{R}^d$.
2. *There exists N such that for all $k \in \mathbb{N}$, $\text{card}\{j : 4B_j \cap 4B_k \neq \emptyset\} \leq N$.*

The doubling condition on w is a crucial point in the proof given by the authors in [2] of the following proposition.

PROPOSITION 2.3. ([2]) *Given $\beta \geq 0$, $w \in D$ (see (1.6)) and $\{x_k\}_{k=1}^\infty$ a sequence as in Proposition 2.2, a function f belongs to $BMO_{\mathcal{L}}^\beta(w)$ if, and only if, f satisfies (1.12) for any ball B , and*

$$\int_{B(x_k, \rho(x_k))} |f| \leq Cw(B(x_k, \rho(x_k)))\rho(x_k)^\beta \text{ for all } k \geq 1. \tag{2.1}$$

The previous result allows us to provide the following characterization of the space $BMO_{\mathcal{L}}^\beta(w)$ that in the sequel will be used as it definition.

COROLLARY 2.1. ([2]) *Let $\beta \geq 0$ and $w \in D$ (see (1.6)). A function f belongs to $BMO_{\mathcal{L}}^\beta(w)$ if, and only if, for some constant C*

$$\frac{1}{w(B)} \int_B |f - f_B| \leq CR^\beta, \text{ if } B = B(x, r) \text{ and } r < \rho(x) \tag{2.2}$$

and

$$\frac{1}{w(B(x, \rho(x)))} \int_{B(x, \rho(x))} |f| \leq C\rho(x)^\beta. \tag{2.3}$$

If $w \equiv 1$ the atomic decomposition given in [11] and [13] shows that $BMO_{\mathcal{L}}^\beta$, $\beta \geq 0$, is the dual space of the H^p -space defined in those works. In this setting, the $BMO_{\mathcal{L}}$ space, $\beta = 0$, was defined in [10] as a natural substitute of L^∞ in the context of the semigroup generated by the operator \mathcal{L} .

In the case that w is not doubling then inequality (2.1) implies the condition

$$\int_B |f| \leq Cw(cB) |B|^{\frac{\beta}{d}}, \quad \text{if } r \geq \rho(x), \tag{2.4}$$

for the geometric constant $c = c_0$ in Proposition 2.1 and some $C > 0$. Therefore, if w is not doubling a different space should be defined, $BMO_{c,\mathcal{L}}^\beta(w)$, c a positive number, as the one satisfying (2.4) and

$$\int_B |f - f_B| \leq Cw(cB) |B|^{\frac{\beta}{d}}, \quad \text{if } r < \rho(x), \tag{2.5}$$

for some constant C . Clearly, $BMO_{1,\mathcal{L}}^\beta(w) = BMO_{\mathcal{L}}^\beta(w)$. Thus, a different version of Corollary 2.1 can be obtained as follows,

COROLLARY 2.2. *Let $\beta \geq 0$ and w a weight. If f belongs to $BMO_{c,\mathcal{L}}^\beta(w)$ then for some constant C it satisfies*

$$\int_B |f - f_B| \leq Cw(cB) R^\beta \quad \text{if } B = B(x, r) \text{ and } r < \rho(x), \tag{2.6}$$

and

$$\int_{B(x,\rho(x))} |f| \leq Cw(B(x, c\rho(x))) \rho(x)^\beta \quad \text{for all } x \in \mathbb{R}^d. \tag{2.7}$$

Reciprocally, if f satisfies inequalities (2.6) and (2.7) then f belongs to $BMO_{\tilde{c},\mathcal{L}}^\beta(w)$ where $\tilde{c} \geq c$.

Hence, by Corollary 2.2 if w is not doubling it is still possible to obtain a weaker version of Theorem 1.1 and Corollary 1.1, with $BMO_{c,\mathcal{L}}^{\alpha-\frac{d}{p}+\delta-\lambda}(w)$ as arrival space.

In the remaining part of this section some useful lemmas related to the kernel K_α of \mathcal{J}_α will be stated and given the references to their proofs. That kernel is given by the formula

$$K_\alpha(x, y) = \int_0^\infty k_t(x, y) t^{\alpha/2} \frac{dt}{t},$$

where k_t is the kernel of the operator $e^{-t\mathcal{L}}$ ($t > 0$).

LEMMA 2.1. ([23]) *Given $N > 0$ there exists a constant $C = C_N$ such that for all x and y in \mathbb{R}^d ,*

$$k_t(x, y) \leq Ct^{-\frac{d}{2}} e^{-\frac{|x-y|^2}{ct}} \left(1 + \frac{\sqrt{t}}{\rho(x)} + \frac{\sqrt{t}}{\rho(y)} \right)^{-N}.$$

As consequence of the above inequality it follows that

$$K_\alpha(x, y) \leq \frac{C}{|x - y|^{d-\alpha}} \tag{2.8}$$

for all x and y in \mathbb{R}^d .

LEMMA 2.2. ([13]) *Given $N > 0$ and $0 < \nu < \delta_0$, there exists a constant $C = C_N$ such that*

$$|k_t(x, y) - k_t(x_0, y)| \leq C \left(\frac{|x - x_0|}{\sqrt{t}} \right)^\nu t^{-\frac{d}{2}} e^{-\frac{|x-y|^2}{ct}} \left(1 + \frac{\sqrt{t}}{\rho(x)} + \frac{\sqrt{t}}{\rho(y)} \right)^{-N}$$

for all x, y and x_0 in \mathbb{R}^d with $|x - x_0| < \sqrt{t}$.

We recall that a function ψ is said to be *rapidly decaying* (see [12]) if for each $N > 0$ there exists a constant C_N such that $|\psi(x)| \leq C_N(1 + |x|)^{-N}$. Its dilation is defined by $\psi_t(x) = \frac{1}{t^{\frac{d}{2}}} \psi\left(\frac{x}{\sqrt{t}}\right)$ for $t > 0$.

Some estimates on the function $q_t(x, y) = k_t(x, y) - \tilde{k}_t(x, y)$, where \tilde{k}_t is the kernel of the classical heat operator $e^{-t\Delta}$, will be useful later.

LEMMA 2.3. ([12]) *There exist a rapidly decaying no negative function ψ , $0 < \nu < 2 - \frac{d}{q_0}$ and $C > 0$ such that for $x, y \in \mathbb{R}^d$ and $t > 0$*

$$|q_t(x, y)| \leq C \left(\frac{\sqrt{t}}{\rho(x)} \right)^\nu \psi_t(x - y).$$

LEMMA 2.4. ([12]) *For all $0 < \nu < \delta_0$ there exists a rapidly decaying function ψ and a constant $C > 0$ such that $|q_t(x, y + h) - q_t(x, y)| \leq \left(\frac{|h|}{\rho(x)}\right)^\nu \psi_t(x - y)$ for x, y in \mathbb{R}^d , $t > 0$ and $|h| < \min(C\rho(y), \frac{|x-y|}{4})$.*

3. Properties and examples of the class of weights

LEMMA 3.1. *The classes $\mathcal{S}(p, \delta, \lambda)$ are increasing in p for $1 \leq p \leq \infty$ and δ and λ real numbers. That is, $\mathcal{S}(p, \delta, \lambda) \subseteq \mathcal{S}(\tau p, \delta, \lambda) \subseteq \mathcal{S}(\infty, \delta, \lambda)$ for all $\tau > 1$ and $1 \leq p < \infty$.*

Proof. Let $B = B(x, t)$. By Hölder’s inequality and Definition 1.1 there exists $C > 0$ and $N_0 \geq 0$ such that, if $p > 1$,

$$\left(\frac{v^{(\tau p)'}(\theta B)}{|\theta B|} \right)^{\frac{1}{(\tau p)'}} \leq \left(\frac{v^{p'}(\theta B)}{|\theta B|} \right)^{\frac{1}{p'}} \leq C \left(1 + \frac{\theta t}{\rho(x)} \right)^{N_0} \theta^{-\lambda} \frac{w(B)}{|B|^{1-\frac{\delta-\lambda}{d}}}.$$

For $p = 1$ the proof follows in the same way by replacing the second term above by $\sup_{\theta B} v$. That is,

$$\left(\frac{v^{t'}(\theta B)}{|\theta B|}\right)^{\frac{1}{t'}} \leq C \sup_{\theta B} v \leq C \left(1 + \frac{\theta t}{\rho(x)}\right)^{N_0} \theta^{-\lambda} \frac{w(B)}{|B|^{1-\frac{\delta-\lambda}{d}}}.$$

for $1 \leq p < \infty$. Analogously, $\mathcal{S}(p, \delta, \lambda) \subseteq \mathcal{S}(\infty, \delta, \lambda)$. \square

In order to show a more precise relationship between the class of weights defined in Definition 1.1 and the classes introduced in [17] and [34] we prove the next lemma.

LEMMA 3.2. *Given $1 < p < \infty$, λ, δ real numbers and $N_0 \geq 0$, if $(v, w) \in \mathcal{S}(p, \delta, \lambda, N_0)$ and $\xi + \frac{d}{p} + \lambda > 0$ then there exists a constant $C > 0$ such that*

$$|B|^{1+\frac{\xi}{d}} \left(\int_{\mathbb{R}^d-B} \frac{v^{p'}(y)}{|x-y|^{(d+\xi)p'}} \left(1 + \frac{|x-y|}{\rho(x)}\right)^{-Np'} dy\right)^{\frac{1}{p'}} \leq C \frac{w(B)}{|B|^{\frac{1}{p}-\frac{\gamma}{d}}}, \tag{3.1}$$

and

$$\left(v^{p'}(B)\right)^{\frac{1}{p'}} \left(1 + \frac{|B|^{\frac{1}{d}}}{\rho(x)}\right)^{-N} \leq C \frac{w(B)}{|B|^{\frac{1}{p}-\frac{\gamma}{d}}}, \tag{3.2}$$

where $\gamma = \delta - \lambda$, and $N \geq N_0$ and both inequalities hold for any ball $B = B(x, R)$. Reciprocally, if (3.1) and (3.2) hold for some ξ and γ real numbers and $N \geq 0$ then $(v, w) \in \mathcal{S}(p, \gamma - \frac{d}{p} - \xi, -\frac{d}{p} - \xi, N)$.

Proof. Inequality (3.2) is a direct application of Definition 1.1 when $\theta = 1$. On the other hand, given a ball B of radius t , by a dyadic decomposition and Definition 1.1 we get

$$\begin{aligned} & \left(\int_{\mathbb{R}^d-B} \frac{v^{p'}(y)}{|x-y|^{(d+\xi)p'}} \left(1 + \frac{|x-y|}{\rho(x)}\right)^{-Np'} dy\right)^{\frac{1}{p'}} \\ & \leq C \left(\sum_{k=0}^{\infty} \frac{1}{(2^k t)^{(d+\xi)p'}} v^{p'}(B(x, 2^k t)) \left(1 + \frac{2^k t}{\rho(x)}\right)^{-Np'}\right)^{\frac{1}{p'}} \\ & \leq C \frac{1}{|B|^{1+\frac{\xi}{d}}} \left(\sum_{k=0}^{\infty} 2^{-k(d+\xi)p'} 2^{k(d-\lambda p')}\right)^{\frac{1}{p'}} \frac{w(B)}{|B|^{\frac{1}{p}-\frac{\delta-\lambda}{d}}} \\ & \leq C \frac{w(B)}{|B|^{1+\frac{1}{p}+\frac{\xi}{d}-\frac{\delta-\lambda}{d}}} \left(\sum_{k=0}^{\infty} 2^{-k(\xi+\frac{d}{p}+\lambda)p'}\right)^{\frac{1}{p'}} \\ & \leq C \frac{w(B)}{|B|^{1+\frac{1}{p}+\frac{\xi}{d}-\frac{\delta-\lambda}{d}}} \end{aligned}$$

if $\xi + \frac{d}{p} + \lambda > 0$.

Reciprocally, (3.1) and (3.2) jointly are equivalent to the inequality

$$|B|^{1+\frac{\xi}{d}} \left(\int_{\mathbb{R}^d} v^{p'}(y) \left[|x-y|^{(d+\xi)p'} \left(1 + \frac{|x-y|}{\rho(x)} \right)^{Np'} + |B|^{(1+\frac{\xi}{d})p'} \left(1 + \frac{|B|^{\frac{1}{d}}}{\rho(x)} \right)^{Np'} \right]^{-1} dy \right)^{\frac{1}{p'}} \leq C \frac{w(B)}{|B|^{\frac{1}{p}-\frac{\gamma}{d}}}.$$

Moreover, we easily get that

$$|B|^{1+\frac{\xi}{d}} \left(\int_{\mathbb{R}^d} v^{p'}(y) \left[|x-y|^{(d+\xi)p'} \left(1 + \frac{|x-y|}{\rho(x)} \right)^{Np'} + |B|^{(1+\frac{\xi}{d})p'} \left(1 + \frac{|B|^{\frac{1}{d}}}{\rho(x)} \right)^{Np'} \right]^{-1} dy \right)^{\frac{1}{p'}} \geq |B|^{1+\frac{\xi}{d}} \frac{(v^{p'}(\theta B))^{\frac{1}{p'}}}{|\theta B|^{1+\frac{\xi}{d}} \left(1 + \frac{\theta R}{\rho(x)} \right)^N}.$$

Then

$$|B|^{1+\frac{\xi}{d}} \frac{(v^{p'}(\theta B))^{\frac{1}{p'}}}{|\theta B|^{1+\frac{\xi}{d}} \left(1 + \frac{\theta R}{\rho(x)} \right)^N} \leq C \frac{w(B)}{|B|^{\frac{1}{p}-\frac{\gamma}{d}}}.$$

That is, $(v^{p'}(\theta B))^{\frac{1}{p'}} \leq C \theta^{d+\xi} \left(1 + \frac{\theta R}{\rho(x)} \right)^N \frac{w(B)}{|B|^{\frac{1}{p}-\frac{\gamma}{d}}}$. Now setting $\frac{d}{p'} - \lambda = d + \xi$ and $\delta - \lambda = \gamma$, we obtain that $(v, w) \in \mathcal{S}(p, \delta, \lambda, N)$ with $\delta = \gamma - \frac{d}{p} - \xi$ and $\lambda = -\frac{d}{p} - \xi$. \square

The above lemma allows us to compare the family of weights defined in this work with other known classes. For example, taking $N_0 = 0$, $\alpha = \delta - \lambda$, $-(\frac{d}{p} + \lambda) < 1 - \alpha$ and p fixed in our class we obtain the set of weights $H(p, \alpha, 2\alpha - \frac{d}{p} - d)$ in [34] and if, in addition, we set $\delta - \lambda = 0$ and consider the family of one weights $v = w$ then we obtain the class in [17].

3.1. The case $\delta = \lambda$

This special case of classes of weights displays significant features that will be described in this section. Let us first define an extension of the class $A_p^{\rho, \theta}$ introduced in [4] to a family of pair of weights in the following way. The pair (w, v) belongs to $A_p^{\rho, N}$, $N \geq 0$, if there exists a constant C such that

$$(w(B))^{\frac{1}{p}} \left(v^{-\frac{1}{p-1}}(B) \right)^{\frac{1}{p'}} \leq C |B| \left(1 + \frac{r}{\rho(x)} \right)^N$$

for every ball $B = B(x, r)$ and $1 < p < \infty$. In the case $p = 1$ the pair (w, v) satisfies inequality

$$w(B) \leq C |B| \inf_B v \left(1 + \frac{r}{\rho(x)} \right)^N.$$

PROPOSITION 3.1. *Let $1 < q < \infty$. If $(w^{q'}, v^{q'}) \in A_1^{p,N}$ then $(w, v) \in \mathcal{S}(q, 0, 0, \frac{N}{q'})$.*

Proof. Given $\theta \geq 1$ and $B = B(x, r)$, statement $(w^{q'}, v^{q'}) \in A_1^{p,N}$ implies that

$$\begin{aligned} (w^{q'}(\theta B))^{\frac{1}{q'}} \left(1 + \frac{\theta r}{\rho(x)}\right)^{-\frac{N}{q'}} &\leq C|\theta B|^{\frac{1}{q'}} \inf_B v \leq C\theta^{\frac{d}{q'}} |B|^{\frac{1}{q'}} \inf_B v \\ &\leq C\theta^{\frac{d}{q'}} |B|^{\frac{1}{q'}} \frac{v(B)}{|B|} \leq C\theta^{\frac{d}{q'}} \frac{v(B)}{|B|^{\frac{1}{q'}}}, \end{aligned}$$

which is the desired inequality. \square

PROPOSITION 3.2. *Let $1 < q < \infty$. If $w \in D_\eta$ and $(v^{q'}, w^{q'}) \in A_{q'+1}^{p,N}$ then $(v, w) \in \mathcal{S}(q, -\eta, -\eta, N(1 + \frac{1}{q'}))$.*

Proof. The hypothesis $(v^{q'}, w^{q'}) \in A_{q'+1}^{p,N}$ implies that

$$v^{q'}(\theta B) (w^{-1}(\theta B))^{q'} \leq C|\theta B|^{q'+1} \left(1 + \frac{\theta r}{\rho(x_{\theta B})}\right)^{N(q'+1)}.$$

Then Hölder’s inequality, the doubling condition and the fact $|\theta B| \geq |B|$ show that

$$\begin{aligned} (v^{q'}(\theta B))^{\frac{1}{q'}} \left(1 + \frac{\theta r}{\rho(x_{\theta B})}\right)^{-N(1+\frac{1}{q'})} &\leq C|\theta B|^{\frac{1}{q'}} \frac{|\theta B|}{w^{-1}(\theta B)} \\ &\leq C|\theta B|^{\frac{1}{q'}} \frac{w(\theta B)}{|\theta B|} \\ &\leq C\theta^{\frac{d}{q'}+\eta} \frac{w(B)}{|B|^{\frac{1}{q'}}} \end{aligned}$$

which proves that $(v, w) \in \mathcal{S}(q, -\eta, -\eta, N(1 + \frac{1}{q'}))$. \square

The one weight case $v \in \mathcal{S}(p, \lambda, \lambda)$ satisfies special properties. The case $N_0 = 0$ was defined in [2] in connection with the Schrödinger operator and in [17] related to the Laplacian operator. One of the features of this class of weights is a Reverse-Hölder-type inequality. Thereafter, we will be able to recover the results in [2] for $N_0 = 0$.

DEFINITION 3.1. Given $1 \leq p < \infty$ a weight v satisfies the Reverse-Hölder type inequality of order p , and say $v \in RH_p(\rho)$ if there exist $C > 0$ and $N_0 \geq 0$ such that for any ball $B = B(x, R)$,

$$\left(\frac{v^p(B)}{|B|}\right)^{\frac{1}{p}} \leq C \left(1 + \frac{R}{\rho(x)}\right)^{N_0} \frac{v(B)}{|B|}. \tag{3.3}$$

We say that $v \in RH_\infty(\rho)$, if there exist $C > 0$ and $N_0 \geq 0$ such that for any ball $B = B(x, R)$,

$$\sup_B v \leq C \left(1 + \frac{R}{\rho(x)}\right)^{N_0} \frac{v(B)}{|B|}. \tag{3.4}$$

LEMMA 3.3. Let $1 \leq p \leq \infty$ and λ be a real number. If $v \in \mathcal{S}(p, \lambda, \lambda)$ then

1. $v \in RH_{p'}(\rho)$.
2. v is \mathcal{L} -doubling, that is, there exist $\eta_1 > 0$, $C > 0$ and N_0 big enough, such that

$$v(\theta B) \leq C\theta^{\eta_1} \left(1 + \frac{R}{\rho(x)}\right)^{N_0} v(B)$$

for any ball $B = B(x, R)$.

3. $v^{p'}$ (if $p' < \infty$) or $\sup v$ (if $p' = \infty$) are \mathcal{L} -doubling, that is, there exist $\eta_2 > 0$, $C > 0$ and N_0 big enough, such that

$$v^{p'}(\theta B) \leq C\theta^{\eta_2} \left(1 + \frac{R}{\rho(x)}\right)^{N_0 p'} v^{p'}(B) \quad \text{if } p' < \infty \tag{3.5}$$

or

$$\sup_{\theta B} v \leq C\theta^{\eta_2} \left(1 + \frac{R}{\rho(x)}\right)^{N_0} \sup_B v \quad \text{if } p' = \infty$$

for any ball $B = B(x, R)$.

Reciprocally, if $v \in RH_{p'}(\rho)$ and it is \mathcal{L} -doubling then there exists a real number λ such that $v \in \mathcal{S}(p, \lambda, \lambda)$.

Proof. The first item is an immediate consequence of Definition 1.1 in the case $\delta = \lambda$ and $\theta = 1$. For the second and third item we apply first Hölder’s inequality, the Definition 1.1 for some $N_0 \geq 0$ and $\delta = \lambda$ and, again, Hölder’s inequality, to get

$$\begin{aligned} \frac{v(\theta B)}{|\theta B|} &\leq \left(\frac{v^{p'}(\theta B)}{|\theta B|}\right)^{\frac{1}{p'}} \leq C\theta^{-\lambda+N_0} \left(1 + \frac{R}{\rho(x)}\right)^{N_0} \frac{v(B)}{|B|} \\ &\leq C\theta^{-\lambda+N_0} \left(1 + \frac{R}{\rho(x)}\right)^{N_0} \left(\frac{v^{p'}(B)}{|B|}\right)^{\frac{1}{p'}}. \end{aligned}$$

Hence, taking on one side the first and third term and, on the other, the second and fourth term, we obtain

$$v(\theta B) \leq C\theta^{d-\lambda+N_0} \left(1 + \frac{R}{\rho(x)}\right)^{N_0} v(B) \tag{3.6}$$

and

$$v^{p'}(\theta B) \leq C\theta^{\left(\frac{d}{p'}-\lambda+N_0\right)p'} \left(1 + \frac{R}{\rho(x)}\right)^{N_0 p'} v^{p'}(B) \quad \text{if } p' < \infty.$$

On the other hand taking $\sup v$ in (3.6) it follows the case $p' = \infty$.

Reciprocally, if $v \in RH_{p'}(\rho)$ and it is \mathcal{L} -doubling then there exists a constant C and a real number η such that

$$\begin{aligned} \left(\frac{v^{p'}(\theta B)}{|\theta B|}\right)^{\frac{1}{p'}} &\leq C\left(1 + \frac{\theta R}{\rho(x)}\right)^{N_0} \frac{v(\theta B)}{|\theta B|} \\ &\leq C\left(1 + \frac{\theta R}{\rho(x)}\right)^{N_0} \theta^{\eta-d} \left(1 + \frac{R}{\rho(x)}\right)^{N_1} \frac{v(B)}{|B|} \\ &\leq C\left(1 + \frac{\theta R}{\rho(x)}\right)^{N_0+N_1} \theta^{\eta-d} \frac{v(B)}{|B|}. \end{aligned}$$

That is,

$$\left(v^{p'}(\theta B)\right)^{\frac{1}{p'}} \leq C\theta^{\frac{d}{p'}+\eta-d} \left(1 + \frac{\theta R}{\rho(x)}\right)^{N_0+N_1} \frac{v(B)}{|B|^{\frac{1}{p'}}$$

which implies that $v \in \mathcal{S}(p, d - \eta, d - \eta)$. Analogously, if $v \in RH_\infty(\rho)$ and v is \mathcal{L} -doubling then $v \in \mathcal{S}(\infty, d - \eta, d - \eta)$. \square

The next lemma is the fundamental key for proving in the one-weight setting that the class $\mathcal{S}(p, \lambda, \lambda)$ is open to the left in p .

LEMMA 3.4. *Let $1 < q \leq \infty$ and λ be a real number. If $v \in \mathcal{S}(q, \lambda, \lambda)$ then there exist $\tau_0 > 1$ and $K_0 \geq 0$ such that*

$$\left(\frac{v^{\tau q'}(B)}{|B|}\right)^{\frac{1}{\tau q'}} \leq C\left(1 + \frac{R}{\rho(x)}\right)^{K_0} \left(\frac{v^{q'}(B)}{|B|}\right)^{\frac{1}{q'}}$$

for $1 \leq \tau \leq \tau_0$ and any ball $B = B(x, r)$.

Proof. By Lemma 3.3 and Definition 3.3 we get $\left(\frac{v^{q'}(B)}{|B|}\right)^{\frac{1}{q'}} \leq C\frac{v(B)}{|B|}$ for any ball $B = B(x, R)$ such that $R \leq \rho(x)$. In this situation, the proof in [16], page 268, shows that there is $\tau_0 > 1$ and a constant C_1 such that if $1 \leq \tau \leq \tau_0$ then

$$\left(\frac{v^{\tau q'}(B)}{|B|}\right)^{\frac{1}{\tau q'}} \leq C_1 \left(\frac{v^{q'}(B)}{|B|}\right)^{\frac{1}{q'}}. \tag{3.7}$$

On the other hand, in the case $R \geq \rho(x)$ let us denote $\mathcal{F} = \{j : B_j \cap B \neq \emptyset\}$ with $B_j = B(x_j, \rho(x_j))$ and $\{x_j\}_{j \in \mathbb{N}}$ the sequence in Proposition 2.2. Using Proposition 2.1, if $j \in \mathcal{F}$ and $R \geq \rho(x)$ then

$$\rho(x_j) \leq C\rho(x) \left(1 + \frac{R}{\rho(x)}\right)^{\frac{j_0}{j_0+1}} \leq C\rho(x) \left(1 + \frac{R}{\rho(x)}\right) \leq CR \left(1 + \frac{R}{\rho(x)}\right)$$

and, thus, $\cup_{j \in \mathcal{F}} B_j \subset cB$, with $c = 4C\left(1 + \frac{R}{\rho(x)}\right)$. By Proposition 2.1, if $j \in \mathcal{F}$ then

$$\begin{aligned} C\rho(x_j) &\geq \rho(x) \left(1 + \frac{|x_j - x|}{\rho(x)}\right)^{-j_0} \geq \rho(x) \left(1 + \frac{cR}{\rho(x)}\right)^{-j_0} \\ &\geq \frac{1}{C} \rho(x) \left(1 + \frac{R}{\rho(x)}\right)^{-j_0} \left(1 + \frac{R}{\rho(x)}\right)^{-j_0} \geq \frac{1}{\tilde{C}} \rho(x) \left(1 + \frac{R}{\rho(x)}\right)^{-2j_0}. \end{aligned} \tag{3.8}$$

Hence, by (3.7), (3.8) and (3.5), we obtain

$$\begin{aligned}
 \left(\frac{1}{|B|}v^{\tau q'}(B)\right)^{\frac{1}{\tau q'}} &\leq \left(\frac{1}{|B|}\sum_{j\in\mathcal{F}}v^{\tau q'}(B_j)\right)^{\frac{1}{\tau q'}} \leq C\left(\frac{1}{|B|}\sum_{j\in\mathcal{F}}\frac{v^{q'}(B_j)^\tau}{|B_j|^{\tau-1}}\right)^{\frac{1}{\tau q'}} \\
 &\leq C\left(\frac{1}{|B|\rho(x)^{d(\tau-1)}}\left(1+\frac{R}{\rho(x)}\right)^{2j_0d(\tau-1)}\sum_{j\in\mathcal{F}}v^{q'}(B_j)^\tau\right)^{\frac{1}{\tau q'}} \\
 &\leq C\left(\frac{1}{|B|R^{d(\tau-1)}}\left(1+\frac{R}{\rho(x)}\right)^{(2j_0+1)d(\tau)}\right)^{\frac{1}{\tau q'}}\left(\sum_{j\in\mathcal{F}}v^{q'}(B_j)^\tau\right)^{\frac{1}{\tau q'}} \\
 &\leq C\left(1+\frac{R}{\rho(x)}\right)^{3j_0\frac{d}{q'}}\left(\frac{1}{|B|}\sum_{j\in\mathcal{F}}v^{q'}(B_j)\right)^{\frac{1}{q'}} \\
 &\leq C\left(1+\frac{R}{\rho(x)}\right)^{3j_0\frac{d}{q'}}\left(\frac{1}{|B|}\int_{cB}\left(\sum_{j\in\mathcal{F}}\chi_{B_j}\right)v^{q'}\right)^{\frac{1}{q'}} \\
 &\leq C\left(1+\frac{R}{\rho(x)}\right)^{3j_0\frac{d}{q'}}\left(\frac{1}{|B|}v^{q'}(cB)\right)^{\frac{1}{q'}} \\
 &\leq C\left(1+\frac{R}{\rho(x)}\right)^{3j_0\frac{d}{q'}+N_0}c^{\eta_2}\left(\frac{1}{|B|}v^{q'}(B)\right)^{\frac{1}{q'}} \\
 &\leq C\left(1+\frac{R}{\rho(x)}\right)^{K_0}\left(\frac{1}{|B|}v^{q'}(B)\right)^{\frac{1}{q'}},
 \end{aligned}$$

with $K_0 = 3j_0\frac{d}{q'} + N_0 + \eta_2$, and N_0 and η_2 the exponents in (3.5). \square

We are now able to prove the openness result we mentioned before. This result is central in the proof of lemmas and theorems in the one-weight situation.

LEMMA 3.5. *Let $1 < p < \infty$ and λ be a real number. If $v \in \mathcal{S}(p, \lambda, \lambda)$ then there exists $1 < p_0 < p$ such that $v \in \mathcal{S}(q, \lambda, \lambda)$ for $p_0 < q \leq p$.*

Proof. Let us choose τ_0 is as in Lemma 3.4 and set $p_0 = (\tau_0 p')' < p$. Note that if $p_0 < q \leq p$ then $q = (\tau p')'$ for some $1 \leq \tau \leq \tau_0$. Now using Lemma 3.4 and that $v \in \mathcal{S}(p, \lambda, \lambda)$ it follows that

$$\left(\frac{v^{q'}(\theta B)}{|\theta B|}\right)^{\frac{1}{q'}} \leq C\left(1+\frac{\theta R}{\rho(x)}\right)^{K_0}\left(\frac{v^{p'}(\theta B)}{|\theta B|}\right)^{\frac{1}{p'}} \leq C\left(1+\frac{\theta R}{\rho(x)}\right)^{K_0+N_0}\theta^{-\lambda}\frac{v(B)}{|B|}$$

for every $\theta \geq 1$. \square

In the remaining part of this section we show some examples of pairs of weights which belong to the classes defined in this work.

EXAMPLE 3.1. Let us consider the pairs of potential weights $(|x|^{-\varepsilon}, |x|^{-\beta})$ and let us analyze the values of ε and β that allow this pair to belongs to the class of weights defined in this work. Assuming, for example, that $\rho = 1$ then the pair $v(x) = |x|^{-\varepsilon}$ and $w(x) = |x|^{-\beta}$ belongs to the class $\mathcal{S}(p, \delta, \lambda, N_0)$, $1 < p < \infty$, if and only if $\beta \leq \varepsilon < \frac{d}{p}$, $\delta \leq \min(0, \beta)$ and $N_0 \geq \lambda - \delta + \beta - \varepsilon \geq 0$.

In fact, condition $\varepsilon < \frac{\lambda}{p'}$ is equivalent to the local integrability of $v(x) = |x|^{-\varepsilon}$. Moreover,

$$\left(\frac{v^{p'}(B(x,r))}{|B(x,r)|}\right)^{\frac{1}{p'}} \approx \begin{cases} |x|^{-\varepsilon} & \text{if } |x| \geq 2r \\ r^{-\varepsilon} & \text{if } |x| < 2r. \end{cases} \tag{3.9}$$

We want to prove inequality

$$\left(\frac{v^{p'}(B(x,\theta t))}{|B(x,\theta t)|}\right)^{\frac{1}{p'}} (1 + \theta t)^{-N_0} \leq C \frac{w(B(x,t))}{|B(x,t)|^{1-\frac{\delta-\lambda}{d}}} \tag{3.10}$$

for all $\theta \geq 1$ but, by (3.9), it is equivalent to the following three inequalities

$$\begin{aligned} |x|^{\beta-\varepsilon} &\leq C\theta^\lambda t^{\delta-\lambda} (1 + \theta t)^{N_0} && \text{if } \theta t < \frac{|x|}{2}, \\ |x|^\beta &\leq C\theta^{\varepsilon+\lambda} t^{\varepsilon+\delta-\lambda} (1 + \theta t)^{N_0} && \text{if } t < \frac{|x|}{2} \leq \theta t \\ 1 &\leq C\theta^{\varepsilon+\lambda} t^{\varepsilon-\beta+\delta-\lambda} (1 + \theta t)^{N_0} && \text{if } \frac{|x|}{2} \leq t. \end{aligned}$$

However, a careful analysis of the behavior of θ , t and $|x|$ on each region leads to the above inequalities only if $\delta - \lambda \leq \beta - \varepsilon \leq 0$, $\delta \leq \min(0, \beta)$ and $N_0 \geq \beta - \varepsilon - \delta + \lambda$. Reciprocally, these conditions are sufficient to prove the above inequalities.

4. Technical Lemmas and proof of Theorem 1.1

The next lemma gives an estimate for the mean value of order $q < p$ on any ball for functions in $L^{p,\infty}(v)$ and it is a fundamental inequality used in the remaining lemmas of this section.

LEMMA 4.1. *Given $1 < q < p < \infty$ and a weight v , there exists a constant C such that*

$$\left(\int_B \left(\frac{|f(x)|}{v(x)}\right)^q dx\right)^{\frac{1}{q}} \leq C|B|^{\frac{1}{q}-\frac{1}{p}} \left[\frac{f}{v}\right]_p$$

for any ball B and $f \in L^{p,\infty}(v)$.

Proof. If $q < p$ then for $a = \frac{[\frac{f}{v}]_p}{|B|^{\frac{1}{p}}}$ we get the statement as follow

$$\begin{aligned} \int_B \left(\frac{|f(x)|}{v(x)}\right)^q dx &= q \left(\int_0^a + \int_a^\infty\right) t^{q-1} |B \cap \left\{\frac{|f|}{v} > t\right\}| dt \\ &\leq |B|a^q + q \left[\frac{f}{v}\right]_p^p \int_a^\infty t^{q-p-1} dt \leq |B|a^q + \frac{q}{p-q} \left[\frac{f}{v}\right]_p^p a^{q-p}. \quad \square \end{aligned}$$

REMARK 4.1. Condition $q < p$ is crucial to get the statement in Lemma 4.1 since it ensures integrability of the distribution function. Therefore in the foregoing lemmas it will be required that $(v, w) \in \mathcal{S}(q, \delta, \lambda)$ for some $q < p$ when $v \neq w$ or $\delta \neq \lambda$. Nevertheless, in the case $v = w$ and $\delta = \lambda$ by Lemma 3.5 it will be enough to require that $v \in \mathcal{S}(p, \lambda, \lambda)$ since in this case it follows that $v \in \mathcal{S}(q, \lambda, \lambda)$ for all $q < p$ close enough to p . Hence, in the one-weight situation Lemma 4.1 will still apply.

LEMMA 4.2. Given $1 < q < p < \infty$ and λ and δ real numbers if $(v, w) \in \mathcal{S}(q, \delta, \lambda)$ then there exist a positive constant C and $N_0 \geq 0$ such that for all $f \in L^{p, \infty}(v)$

$$\int_{\theta B} |f(x)| dx \leq C \theta^{\frac{d}{p'} - \lambda} \left(1 + \frac{\theta R}{\rho(x)}\right)^{N_0} \frac{w(B)}{|B|^{\frac{1}{p} - \frac{\delta - \lambda}{d}}} \left[\frac{f}{v}\right]_p$$

for all $\theta \geq 1$ and every ball $B = B(x_B, R)$. If $v = w \in \mathcal{S}(p, \lambda, \lambda)$, i.e. $\delta = \lambda$, then the above inequality also holds.

Proof. By Hölder’s inequality and Lemma 4.1, if $(v, w) \in \mathcal{S}(q, \delta, \lambda)$ for some $q < p$ or $v = w \in \mathcal{S}(p, \lambda, \lambda)$ and $q < p$, close enough to p then, for some $C > 0$ and $N_0 \geq 0$,

$$\begin{aligned} \int_{\theta B} |f(x)| dx &\leq \left(\int_{\theta B} v^{q'}(x) dx\right)^{\frac{1}{q'}} \left(\int_{\theta B} \left(\frac{|f(x)|}{v(x)}\right)^q dx\right)^{\frac{1}{q}} \\ &\leq C \left(1 + \frac{\theta R}{\rho(x_B)}\right)^{N_0} \theta^{\frac{d}{q'} - \lambda} \frac{w(B)}{|B|^{\frac{1}{q} - \frac{\delta - \lambda}{d}}} \left(|\theta B|^{\frac{1}{q} - \frac{1}{p}} \left[\frac{f}{v}\right]_p\right) \\ &\leq C \theta^{\frac{d}{p'} - \lambda} \frac{w(B)}{|B|^{\frac{1}{p} - \frac{\delta - \lambda}{d}}} \left(1 + \frac{\theta R}{\rho(x_B)}\right)^{N_0} \left[\frac{f}{v}\right]_p. \quad \square \end{aligned}$$

To prove Theorem 1.1 we need estimates for the fractional integral of the local and global parts of a function $f \in L^{p, \infty}(v)$. The following two lemmas give us those estimates.

LEMMA 4.3. Given $1 < q < p < \infty$, λ and δ real numbers and $\alpha > 0$, if $(v, w) \in \mathcal{S}(q, \delta, \lambda)$ then there exists $N_0 \geq 0$ and C such that for all $f \in L^{p, \infty}(v)$

$$\frac{1}{w(B)} \int_B \mathcal{I}_\alpha(|f| \chi_{2B})(x) dx \leq C \left(1 + \frac{R}{\rho(x_B)}\right)^{N_0} |B|^{\frac{\alpha}{d} - \frac{1}{p} + \frac{\delta - \lambda}{d}} \left[\frac{f}{v}\right]_p$$

for any ball $B = B(x_B, R)$. If $v = w \in \mathcal{S}(p, \lambda, \lambda)$, i.e. $\delta = \lambda$, then the above inequality also holds.

Proof. By (2.8), Fubini’s Theorem, $\alpha > 0$ and Lemma 4.2 applied to $\theta = 2$ there exists $N_0 \geq 0$ such that

$$\begin{aligned} \frac{1}{w(B)} \int_B \mathcal{I}_\alpha(|f|\chi_{2B})(x)dx &= \frac{1}{w(B)} \int_B \int_{2B} K_\alpha(x,y)|f(y)|dy dx \\ &\leq C \frac{1}{w(B)} \int_B \int_{2B} \frac{|f(y)|}{|x-y|^{d-\alpha}} dy dx \\ &\leq C \frac{1}{w(B)} \int_{2B} |f(y)| \int_{B(y,2R)} \frac{1}{|x-y|^{d-\alpha}} dx dy \\ &\leq C \frac{|B|^{\frac{\alpha}{d}}}{w(B)} \int_{2B} |f(y)| dy \\ &\leq C \left(1 + \frac{R}{\rho(x_B)}\right)^{N_0} |B|^{\frac{\alpha}{d} - \frac{1}{p} + \frac{\delta-\lambda}{d}} \left[\frac{f}{v}\right]_p. \quad \square \end{aligned}$$

LEMMA 4.4. Given $1 < q < p < \infty$ and λ and δ real numbers, if $(v,w) \in \mathcal{S}(q,\delta,\lambda)$ then there exist positive constants C and $N_0 \geq 0$ such that for all $f \in L^{p,\infty}(v)$

$$\int_{\mathbb{R}^d \setminus 2B} \frac{|f(y)|}{|x-y|^{d+m}} \left(1 + \frac{|x-y|}{\rho(x)}\right)^{-N} dy \leq C \frac{w(B)}{|B|^{1+\frac{1}{p}+\frac{m+\lambda-\delta}{d}}} \left[\frac{f}{v}\right]_p$$

for any constant m such that $m + \lambda + \frac{d}{p} > 0$, $N \geq N_0$ and any ball $B = B(x_B, R)$. Moreover, if $v = w \in \mathcal{S}(p,\lambda,\lambda)$, i.e. $\delta = \lambda$, then the above inequality also holds.

Proof. Using a dyadic decomposition and Lemma 4.2 there exist $N_0 \geq 0$ such that if $N \geq N_0$ and, also, $m + \lambda + \frac{d}{p} > 0$ then

$$\begin{aligned} &\int_{\mathbb{R}^d \setminus 2B} \frac{|f(y)|}{|x-y|^{d+m}} \left(1 + \frac{|x-y|}{\rho(x)}\right)^{-N} dy \\ &\leq C \sum_{k=1}^\infty (2^k R)^{-(d+m)} \left(1 + \frac{2^k R}{\rho(x)}\right)^{-N} \int_{2^{k+1}B} f(y) dy \\ &\leq C \sum_{k=0}^\infty (2^k R)^{-(d+m)} \left(1 + \frac{2^k R}{\rho(x)}\right)^{-(N-N_0)} 2^{k(\frac{d}{p'}-\lambda)} \frac{w(B)}{|B|^{\frac{1}{p}-\frac{\delta-\lambda}{d}}} \left[\frac{f}{v}\right]_p \\ &\leq C |B|^{-\frac{d+m}{d}} \sum_{k=0}^\infty 2^{-k(m+\lambda+\frac{d}{p})} \frac{w(B)}{|B|^{\frac{1}{p}-\frac{\delta-\lambda}{d}}} \left[\frac{f}{v}\right]_p \\ &\leq C \frac{w(B)}{|B|^{1+\frac{1}{p}+\frac{m+\lambda-\delta}{d}}} \left[\frac{f}{v}\right]_p. \quad \square \end{aligned}$$

The next auxiliary lemma is a key tool to prove Theorem 1.1.

LEMMA 4.5. 1. Given s, ρ and β positive numbers and $N \geq 0$ there exists a constant C such that

$$\int_0^\infty \frac{t^{-\frac{\beta}{2}}}{(1 + \frac{\sqrt{t}}{\rho})^N} e^{-\frac{s^2}{t}} \frac{dt}{t} \leq C \frac{1}{s^\beta (1 + \frac{s}{\rho})^N}.$$

2. Given $s > \sigma$, $\rho > 0$, β a real number, $N \geq 0$ and $M > \beta + N$ there is a constant C such that

$$\int_0^{\sigma^2} \frac{t^{-\frac{\beta}{2}}}{(1 + \frac{\sqrt{t}}{\rho})^N} e^{-\frac{s^2}{t}} \frac{dt}{t} \leq C \left(\frac{\sigma}{s}\right)^{M-N-\beta} \frac{1}{s^\beta (1 + \frac{s}{\rho})^N}.$$

Proof. (1) By the change of variable $u = \frac{s^2}{t}$ we get

$$\int_0^\infty \frac{t^{-\frac{\beta}{2}}}{(1 + \frac{\sqrt{t}}{\rho})^N} e^{-\frac{s^2}{t}} \frac{dt}{t} = \frac{1}{s^\beta} \int_0^\infty \frac{u^{\frac{\beta}{2}}}{(1 + \frac{s}{\rho\sqrt{u}})^N} e^{-u} \frac{du}{u}.$$

If $\frac{s}{\rho} < 1$ then $1 + \frac{s}{\rho} \leq 2$ and hence $\frac{u^{\frac{\beta}{2}}}{(1 + \frac{s}{\rho\sqrt{u}})^N} \leq u^{\frac{\beta}{2}} \leq 2^N \frac{u^{\frac{\beta}{2}}}{(1 + \frac{s}{\rho})^N}$. On the other hand, if $\frac{s}{\rho} > 1$ then $1 + \frac{s}{\rho} \leq 2\frac{s}{\rho}$ and thus $\frac{u^{\frac{\beta}{2}}}{(1 + \frac{s}{\rho\sqrt{u}})^N} \leq \frac{u^{\frac{\beta+N}{2}}}{(\frac{s}{\rho})^N} \leq 2^N \frac{u^{\frac{\beta+N}{2}}}{(1 + \frac{s}{\rho})^N}$. Therefore

$$\int_0^\infty \frac{t^{-\frac{\beta}{2}}}{(1 + \frac{\sqrt{t}}{\rho})^N} e^{-\frac{s^2}{t}} \frac{dt}{t} \leq \frac{2^N}{s^\beta (1 + \frac{s}{\rho})^N} \int_0^\infty (u^{\frac{\beta}{2}} + u^{\frac{\beta+N}{2}}) e^{-u} \frac{du}{u} \leq C_N \frac{1}{s^\beta (1 + \frac{s}{\rho})^N}.$$

(2) By the same change of variable as above, the fact that $\sup_{t>1} t^{M/2} e^{-t} \leq C_M$ for any $M > 0$ and some $C_M > 0$, and following the steps in the above proof, it follows that

$$\begin{aligned} \int_0^{\sigma^2} \frac{t^{-\frac{\beta}{2}}}{(1 + \frac{\sqrt{t}}{\rho})^N} e^{-\frac{s^2}{t}} \frac{dt}{t} &= \frac{1}{s^\beta} \int_{(\frac{s}{\sigma})^2}^\infty \frac{u^{\frac{\beta}{2}}}{(1 + \frac{s}{\rho\sqrt{u}})^N} e^{-u} \frac{du}{u} \\ &\leq C \frac{1}{s^\beta (1 + \frac{s}{\rho})^N} \int_{(\frac{s}{\sigma})^2}^\infty (u^{\frac{\beta-M}{2}} + u^{\frac{\beta+N-M}{2}}) \frac{du}{u} \\ &\leq C \frac{1}{s^\beta (1 + \frac{s}{\rho})^N} \left(\left(\frac{s}{\sigma}\right)^{\beta-M} + \left(\frac{s}{\sigma}\right)^{\beta+N-M} \right) \\ &\leq C \frac{1}{s^\beta (1 + \frac{s}{\rho})^N} \left(\frac{s}{\sigma}\right)^{\beta+N-M} \\ &\leq C \frac{\sigma^{M-N-\beta}}{s^{M-N} (1 + \frac{s}{\rho})^N}. \quad \square \end{aligned}$$

4.1. Proof of Theorem 1.1

Let $(v, w) \in \mathcal{S}(q, \delta, \lambda)$ for some $q < p$ or $v = w \in \mathcal{S}(p, \lambda, \lambda)$, i.e. $\delta = \lambda$ and split $f = f_1 + f_2$, with $f_1 = f\chi_{2B}$ and first assume that $R = \rho(x_B)$. Thus

$$\frac{1}{w(B)} \int_B |\mathcal{I}_\alpha f(x)| dx \leq \frac{1}{w(B)} \int_B |\mathcal{I}_\alpha f_1(x)| dx + \frac{1}{w(B)} \int_B |\mathcal{I}_\alpha f_2(x)| dx \leq \mathcal{I}_1 + \mathcal{I}_2.$$

By Lemma 4.3

$$\mathcal{I}_1 = \frac{1}{w(B)} \int_B |\mathcal{I}_\alpha f_1(x)| dx \leq C|B|^{\frac{\alpha}{d} - \frac{1}{p} + \frac{\delta - \lambda}{d}} \left[\frac{f}{v} \right]_p. \tag{4.1}$$

To estimate \mathcal{I}_2 we display

$$|\mathcal{I}_\alpha f_2(x)| = \left| \int_0^\infty e^{-t\mathcal{L}} f_2(x) t^{\frac{\alpha}{2}} \frac{dt}{t} \right| \leq \int_0^\infty \int_{(2B)^c} k_t(x, y) |f(y)| dy t^{\frac{\alpha}{2}} \frac{dt}{t}.$$

Notice that if $x \in B$ and $y \in \mathbb{R}^d \setminus 2B$ then $2R \leq |x - y| \leq 2|x_B - y|$. Moreover, by Lemma 2.1,

$$\rho(x) \leq C\rho(x_B) \left(1 + \frac{|x - x_B|}{\rho(x_B)} \right)^{\frac{k_0}{k_0+1}} \leq C\rho(x_B). \tag{4.2}$$

Hence, by Lemmas 2.1 and 4.5 given $N > 0$ and $M > 0$, to be chosen later, there is a constant C such that

$$\begin{aligned} |\mathcal{I}_\alpha f_2(x)| &\leq C \int_{(2B)^c} \int_0^\infty t^{\frac{\alpha-d}{2}} \left(1 + \frac{\sqrt{t}}{\rho(x)} \right)^{-(N+M)} e^{-\frac{|x-y|^2}{t}} \frac{dt}{t} |f(y)| dy \\ &\leq C \int_{(2B)^c} \frac{|f(y)|}{|x-y|^{d-\alpha}} \left(1 + \frac{|x-y|}{\rho(x)} \right)^{-(N+M)} dy \\ &\leq C \int_{(2B)^c} \frac{|f(y)|}{|x_B-y|^{d-\alpha}} \left(1 + \frac{|x_B-y|}{\rho(x_B)} \right)^{-(N+M)} dy \\ &\leq C\rho(x_B)^M \int_{(2B)^c} \frac{|f(y)|}{|x_B-y|^{d-\alpha+M}} \left(1 + \frac{|x_B-y|}{\rho(x_B)} \right)^{-N} dy. \end{aligned}$$

By Lemma 4.4, taking $R = \rho(x_B)$, $N \geq N_0$ and choosing M such that $M - \alpha + \lambda + \frac{d}{p} > 0$ we get

$$|\mathcal{I}_\alpha f_2(x)| \leq C\rho(x_B)^M \frac{w(B)}{|B|^{1 + \frac{M-\alpha+\lambda-\delta}{d} + \frac{1}{p}}} \left[\frac{f}{v} \right]_p \leq C \frac{w(B)}{|B|^{1 - \frac{\alpha+\delta-\lambda}{d} + \frac{1}{p}}} \left[\frac{f}{v} \right]_p.$$

Then

$$\mathcal{I}_2 = \frac{1}{w(B)} \int_B |\mathcal{I}_\alpha f_2(x)| dx \leq C|B|^{\frac{\alpha}{d} - \frac{1}{p} + \frac{\delta - \lambda}{d}} \left[\frac{f}{v} \right]_p. \tag{4.3}$$

Estimates (4.1) and (4.3) give the proof of (1.9).

To prove (1.8), we consider $R < \rho(x_B)$, define $c_B = \int_{\mathbb{R}^2} e^{-t\mathcal{L}} f_2(x_B) t^{\frac{\alpha}{2}} \frac{dt}{t}$ and split

$$\frac{1}{w(B)} \int_B |\mathcal{I}_\alpha f(x) - c_B| dx \leq \frac{1}{w(B)} \int_B |\mathcal{I}_\alpha f_1(x)| dx + \frac{1}{w(B)} \int_B |\mathcal{I}_\alpha f_2(x) - c_B| dx.$$

As in (4.1) for $\alpha > 0$ it follows that

$$\frac{1}{w(B)} \int_B |\mathcal{I}_\alpha f_1(x)| dx \leq C |B|^{\frac{\alpha}{d} - \frac{1}{p} + \frac{\delta - \lambda}{d}} \left[\frac{f}{v} \right]_p.$$

On the other hand, for $x \in B$, we have

$$\begin{aligned} |\mathcal{I}_\alpha f_2(x) - c_B| &\leq \int_0^{R^2} |e^{-t\mathcal{L}} f_2(x)| t^{\frac{\alpha}{2}} \frac{dt}{t} + \int_{R^2}^\infty |e^{-t\mathcal{L}} f_2(x) - e^{-t\mathcal{L}} f_2(x_B)| t^{\frac{\alpha}{2}} \frac{dt}{t} \\ &= J_1(x) + J_2(x). \end{aligned}$$

If $|x - x_B| < R \leq \rho(x_B)$ then, as in (4.2), $\rho(x) \leq C\rho(x_B)$. Hence we apply Lemma 2.1, Lemma 4.5 with $\beta = d - \alpha$, $s = |x - y|$, $\sigma = R$, $M + d$ in place of M , and N positive such that $M > N + d - \alpha$, and use that if $|x - x_B| < R$ and $|y - x_B| \geq 2R$ then $|y - x| > R$, to get

$$\begin{aligned} J_1(x) &= \int_0^{R^2} |e^{-t\mathcal{L}} f_2(x)| t^{\frac{\alpha}{2}} \frac{dt}{t} \\ &\leq \int_0^{R^2} \left(\int_{(2B)^c} t^{-\frac{d}{2}} \left(1 + \frac{\sqrt{t}}{\rho(x)} \right)^{-N} e^{-\frac{|x-y|^2}{t}} |f(y)| dy \right) t^{\frac{\alpha}{2}} \frac{dt}{t} \\ &\leq C \int_{(2B)^c} \left(\int_0^{R^2} t^{\frac{\alpha-d}{2}} \left(1 + \frac{\sqrt{t}}{\rho(x)} \right)^{-N} e^{-\frac{|x-y|^2}{t}} \frac{dt}{t} \right) |f(y)| dy \\ &\leq CR^{\alpha+M-N} \int_{(2B)^c} \frac{|f(y)|}{|x-y|^{d+M-N}} \left(1 + \frac{|x-y|}{\rho(x)} \right)^{-N} dy. \end{aligned}$$

By Lemma 4.4 if $M - N + \lambda + \frac{d}{p} > 0$ and $N \geq N_0$ for some N_0 big enough then

$$J_1(x) \leq C |B|^{\frac{\alpha}{d} + \frac{M-N}{d}} \frac{w(B)}{|B|^{1 + \frac{M-N}{d} + \frac{1}{p} - \frac{\delta - \lambda}{d}}} \left[\frac{f}{v} \right]_p \leq C \frac{w(B)}{|B|^{1 - \frac{\alpha + \delta - \lambda}{d} + \frac{1}{p}}} \left[\frac{f}{v} \right]_p \tag{4.4}$$

where we chose $M > \max(N + d - \alpha, N - \lambda - \frac{d}{p})$ and $N \geq N_0$.

On the other hand, we use Lemma 2.2 and Lemma 4.5 to get for any $v < \delta_0$ a constant C such that if $x \in B$ then

$$\begin{aligned} J_2(x) &= \int_{R^2}^\infty |e^{-t\mathcal{L}} f_2(x) - e^{-t\mathcal{L}} f_2(x_B)| t^{\frac{\alpha}{2}} \frac{dt}{t} \\ &= \int_{R^2}^\infty \int_{(2B)^c} |k_t(x, y) - k_t(x_B, y)| |f(y)| dy t^{\frac{\alpha}{2}} \frac{dt}{t} \\ &\leq C |x - x_B|^v \int_{R^2}^\infty \left(\int_{(2B)^c} \frac{t^{\frac{\alpha-d-v}{2}}}{\left(1 + \frac{\sqrt{t}}{\rho(x)} \right)^N} e^{-\frac{|x-y|^2}{t}} |f(y)| dy \right) \frac{dt}{t} \\ &\leq CR^v \int_{(2B)^c} \left(\int_{R^2}^\infty \frac{t^{\frac{\alpha-d-v}{2}}}{\left(1 + \frac{\sqrt{t}}{\rho(x)} \right)^N} e^{-\frac{|x-y|^2}{t}} \frac{dt}{t} \right) |f(y)| dy \\ &\leq CR^v \int_{(2B)^c} \frac{|f(y)|}{|x_B - y|^{d+v-\alpha}} \left(1 + \frac{|x_B - y|}{\rho(x_B)} \right)^{-N} dy. \end{aligned}$$

Proceeding as in (4.4), since we can choose v such that $\alpha - \lambda - \frac{d}{p} < v < \delta_0$ and apply Lemma 4.4, we get

$$J_2(x) \leq CR^v \frac{w(B)}{|B|^{\frac{d+v-\alpha-\delta+\lambda}{d} + \frac{1}{p}}} \left[\frac{f}{v} \right]_p = C \frac{w(B)}{|B|^{\frac{d-\alpha-\delta+\lambda}{d} + \frac{1}{p}}} \left[\frac{f}{v} \right]_p. \tag{4.5}$$

By integrating (4.4) and (4.5) on B we get (1.8).

The proof in the case $v = w \in \mathcal{S}(p, \lambda, \lambda)$ follows from Lemma 3.5, Remark 4.1 and the above reasoning. \square

5. Technical lemmas and proof of Theorem 1.2

In $BMO_{\mathcal{L}}^\beta(v)$ the average control is only on balls with radii greater than ρ at their centers (Corollary 2.1). However, for lower radii some kind of estimate can be proved. The following is a variation of Lemma 6 in [2].

LEMMA 5.1. *Let β, λ, δ and N_0 real numbers such that $\beta > \lambda$ and $\beta, N_0 \geq 0$. If $(v, w) \in \mathcal{S}(\infty, \delta, \lambda, N_0)$ then there exist $C > 0$ such that for any $f \in BMO_{\mathcal{L}}^\beta(v)$ and $k \in \mathbb{N} \cup \{0\}$*

$$\int_{2^k B} |f| \leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} |B|^{\frac{\beta-\lambda+\delta}{d}} w(B) \times \begin{cases} 2^{k(d+\beta-\lambda+N_0)} \left(\frac{R}{\rho(x_B)}\right)^{N_0} & \text{if } k > j_0 \\ 2^{kd} \left(\frac{\rho(x_B)}{R}\right)^{\beta-\lambda} & \text{if } k \leq j_0 \end{cases} \tag{5.1}$$

for any ball $B = B(x_B, R)$ with $R \leq \rho(x_B)$ and $j_0 \in \mathbb{N} \cup \{0\}$ such that $2^{j_0} R \leq \rho(x_B) < 2^{j_0+1} R$.

Proof. Using Definitions 1.13 and 1.1 we consider two cases. If $k \geq j_0 + 1$, then for some $C > 0$, $N_0 \geq 0$ and any $f \in BMO_{\mathcal{L}}^\beta(v)$,

$$\begin{aligned} \int_{2^k B} |f| &\leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} |2^k B|^{\frac{\beta}{d}} v(2^k B) \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} |2^k B|^{\frac{\beta-\lambda+d}{d}} \left(1 + \frac{2^k R}{\rho(x_B)}\right)^{N_0} \frac{w(B)}{|B|^{1-\frac{\delta}{d}}} \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} 2^{k(\beta-\lambda+d+N_0)} \left(\frac{R}{\rho(x_B)}\right)^{N_0} |B|^{\frac{\beta-\lambda+\delta}{d}} w(B). \end{aligned}$$

If $k \leq j_0$ and $\beta - \lambda > 0$ by a dyadic decomposition and the previous inequality, then there exists a constant C such that for $f \in BMO_{\mathcal{L}}^\beta(v)$

$$\begin{aligned}
 \frac{1}{|2^k B|} \int_{2^k B} |f| &\leq C \sum_{j=k}^{j_0+1} \frac{1}{|2^j B|} \int_{2^j B} |f(z) - f_{2^j B}| dz + \frac{1}{|2^{j_0+1} B|} \int_{2^{j_0+1} B} |f(z)| dz \\
 &\leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \left\{ \sum_{j=k}^{j_0+1} |2^j B|^{\frac{\beta}{d}} \frac{v(2^j B)}{|2^j B|} + 2^{j_0(\beta-\lambda+N_0)} \left(\frac{R}{\rho(x_B)}\right)^{N_0} |B|^{\frac{\beta-\lambda+\delta}{d}} \frac{w(B)}{|B|} \right\} \\
 &\leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \left\{ \sum_{j=k}^{j_0+1} 2^{j(\beta-\lambda)} \left(1 + \frac{2^j R}{\rho(x_B)}\right)^{N_0} + \left(\frac{\rho(x_B)}{R}\right)^{\beta-\lambda} \right\} |B|^{\frac{\beta-\lambda+\delta}{d}} \frac{w(B)}{|B|} \\
 &\leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \left\{ \sum_{j=k}^{j_0+1} 2^{j(\beta-\lambda)} + \left(\frac{\rho(x_B)}{R}\right)^{\beta-\lambda} \right\} |B|^{\frac{\beta-\lambda+\delta}{d}} \frac{w(B)}{|B|} \\
 &\leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \left(\frac{\rho(x_B)}{R}\right)^{\beta-\lambda} |B|^{\frac{\beta-\lambda+\delta}{d}} \frac{w(B)}{|B|},
 \end{aligned}$$

that is,

$$\int_{2^k B} |f| \leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} 2^{kd} \left(\frac{\rho(x_B)}{R}\right)^{\beta-\lambda} |B|^{\frac{\beta-\lambda+\delta}{d}} w(B). \quad \square$$

LEMMA 5.2. Given $\beta \geq 0$, $\beta > \lambda$ with λ, δ and α real numbers and $N_0 \geq 0$, let us assume that $(v, w) \in \mathcal{S}(\infty, \delta, \lambda, N_0)$.

1. If $M > N_0 + \beta - \lambda + \alpha$ then for some constant $C = C_M$ and any ball $B = B(x_B, R)$ with $R \leq \rho(x_B)$,

$$\int_{|x_B - y| \geq 2\rho(x_B)} \frac{|f(y)|}{|x_B - y|^{d+M-\alpha}} dy \leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \frac{R^{\delta-d}}{\rho(x_B)^{-\beta+\lambda+M-\alpha}} w(B). \quad (5.2)$$

2. If $M > \alpha$ then for some constant $C = C_M$ and any ball $B = B(x_B, R)$ with $R \leq \rho(x_B)$,

$$\int_{2R \leq |x_B - y| < 2\rho(x_B)} \frac{|f(y)|}{|x_B - y|^{d+M-\alpha}} dy \leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \frac{R^{\delta-d-M+\alpha}}{\rho(x_B)^{-\beta+\lambda}} w(B) \quad (5.3)$$

for all $f \in BMO_{\mathcal{L}}^\beta(v)$.

Proof. Let $j_0 \in \mathbb{N} \cup \{0\}$ such that $2^{j_0} R \leq \rho(x_B) < 2^{j_0+1} R$. By a dyadic decomposing and the first item in Lemma 5.1, if $M > \beta - \lambda + \alpha + N_0$ then there exists a

constant C such that

$$\begin{aligned} \int_{|x_B-y|\geq 2\rho(x_B)} \frac{|f(y)|}{|x_B-y|^{d+M-\alpha}} dy &\leq \sum_{k=j_0}^{\infty} \frac{1}{(2^k R)^{d+M-\alpha}} \int_{2^{k+1}B} f(y) dy \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \sum_{k=j_0}^{\infty} 2^{-k(-\beta-\alpha+\lambda+M-N_0)} \left(\frac{R}{\rho(x_B)}\right)^{N_0} R^{\beta-\lambda+\delta-d-M+\alpha} w(B) \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} 2^{-j_0(-\beta-\alpha+\lambda+M-N_0)} \left(\frac{R}{\rho(x_B)}\right)^{N_0} R^{\beta-\lambda+\delta-d-M+\alpha} w(B) \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \left(\frac{R}{\rho(x_B)}\right)^{-\beta-\alpha+\lambda+M} R^{\beta-\lambda+\delta-d-M+\alpha} w(B). \end{aligned}$$

On the other hand, if $M > \alpha$ we get

$$\begin{aligned} \int_{2R \leq |x_B-y| < 2\rho(x_B)} \frac{|f(y)|}{|x_B-y|^{d+M-\alpha}} dy &\leq \sum_{k=0}^{j_0} \frac{1}{(2^k R)^{d+M-\alpha}} \int_{2^{k+1}B} f(y) dy \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \sum_{k=0}^{j_0} \frac{1}{2^{k(M-\alpha)}} \left(\frac{\rho(x_B)}{R}\right)^{\beta-\lambda} R^{\beta-\lambda+\delta-d-M+\alpha} w(B) \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \left(\frac{\rho(x_B)}{R}\right)^{\beta-\lambda} R^{\beta-\lambda+\delta-d-M+\alpha} w(B). \quad \square \end{aligned}$$

In the following two lemmas we study certain kind of integrability for the oscillations of f .

LEMMA 5.3. *Let $\beta \geq 0$, λ and $\delta \in \mathbb{R}$, $\alpha \geq 0$ and $\beta > \lambda$.*

1. *If $(v, w) \in \mathcal{S}(\infty, \delta, \lambda, N_0)$, and $M > \beta + \alpha - \lambda + N_0$ then, for some $C > 0$,*

$$\int_{|x_B-y|\geq 2\rho(x_B)} \frac{|f(y) - f_B|}{|x_B-y|^{d+M-\alpha}} dy \leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \frac{R^{\delta-d}}{\rho(x_B)^{-\beta+\lambda+M-\alpha}} w(B) \quad (5.4)$$

for any ball $B := B(x_B, R)$ with $R \leq \rho(x_B)$ and $f \in BMO_{\mathcal{L}}^\beta(v)$.

2. *If $(v, w) \in \mathcal{S}(\infty, \delta, \lambda)$, and $M > \beta - \lambda + \alpha$ then, for some $C > 0$*

$$\int_{2R < |x_B-y| \leq 2\rho(x_B)} \frac{|f(y) - f_B|}{|y-x_B|^{d+M-\alpha}} dy \leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \frac{w(B)}{R^{-\beta+\lambda-\delta+M-\alpha+d}} \quad (5.5)$$

for any ball $B := B(x_B, R)$ with $R \leq \rho(x_B)$ and $f \in BMO_{\mathcal{L}}^\beta(v)$.

Proof. Let $j_0 \in \mathbb{N} \cup \{0\}$ such that $2^{j_0}R \leq \rho(x_B) < 2^{j_0+1}R$. Using a dyadic decomposition, Lemma 5.1 and Definition 1.1, and choosing M such that $M > N_0 + \beta -$

$\lambda + \alpha > \alpha$ we get

$$\begin{aligned} & \int_{|x_B-y|>2\rho(x_B)} \frac{|f(y) - f_B|}{|x_B - y|^{d+M-\alpha}} dy \leq C \sum_{j=j_0+1}^{\infty} \frac{1}{(2^j R)^{d+M-\alpha}} \int_{2^{j+1}B} |f(y) - f_B| dy \\ & \leq C \sum_{j=j_0+1}^{\infty} \frac{1}{(2^j R)^{d+M-\alpha}} \left\{ \int_{2^{j+1}B} |f(y)| dy + |f_B| (2^j R)^d \right\} \\ & \leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \frac{w(B)}{R^{d+M-\alpha-\beta-\delta+\lambda}} \\ & \quad \times \sum_{j=j_0+1}^{\infty} 2^{-j(d+M-\alpha)} \left\{ 2^{j(d+\beta-\lambda+N_0)} \left(\frac{R}{\rho(x_B)}\right)^{N_0} + 2^{jd} \left(\frac{\rho(x_B)}{R}\right)^{\beta-\lambda} \right\} \\ & \leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \frac{w(B)}{R^{d+M-\alpha-\beta-\delta+\lambda}} \\ & \quad \times \left\{ \left(\frac{R}{\rho(x_B)}\right)^{N_0} \sum_{j=j_0+1}^{\infty} 2^{j(\beta-\lambda-M+\alpha+N_0)} + \left(\frac{\rho(x_B)}{R}\right)^{\beta-\lambda} \sum_{j=j_0+1}^{\infty} 2^{-j(M-\alpha)} \right\} \\ & \leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \left(\frac{R}{\rho(x_B)}\right)^{-\beta+\lambda+M-\alpha} \frac{w(B)}{R^{d+M-\alpha-\beta-\delta+\lambda}}. \end{aligned}$$

On the other hand, by Definition 1.1 and (2.2) if $M > \beta - \lambda + \alpha > \alpha$ it follows that for some $C > 0$,

$$\begin{aligned} & \int_{2R < |x_B-y| \leq 2\rho(x_B)} \frac{|f(y) - f_B|}{|x_B - y|^{d+M-\alpha}} dy \leq C \sum_{j=1}^{j_0} \frac{1}{(2^j R)^{d+M-\alpha}} \int_{2^{j+1}B} |f(y) - f_B| dy \\ & \leq \sum_{j=1}^{j_0} \frac{1}{(2^j R)^{M-\alpha}} \left\{ \frac{1}{(2^{j+1}R)^d} \int_{2^{j+1}B} |f(y) - f_{2^{j+1}B}| dy + \sum_{k=0}^j |f_{2^{k+1}B} - f_{2^k B}| \right\} \\ & \leq C \sum_{j=1}^{j_0} \frac{1}{(2^j R)^{M-\alpha}} \sum_{k=0}^{j+1} \frac{1}{(2^k R)^d} \int_{2^k B} |f(y) - f_{2^k B}| dy \\ & \leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \sum_{j=1}^{j_0} \frac{1}{(2^j R)^{M-\alpha}} \sum_{k=1}^{j+1} (2^k R)^\beta \frac{v(2^k B)}{(2^k R)^d} \\ & \leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \sum_{j=1}^{j_0} \frac{1}{2^{j(M-\alpha)}} \sum_{k=1}^{j+1} 2^{k\beta} 2^{-k\lambda} \left(1 + \frac{2^k R}{\rho(x_B)}\right)^{N_0} \frac{w(B)}{R^{d-\beta+\lambda-\delta+M-\alpha}} \\ & \leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \sum_{k=1}^{j_0+1} 2^{k(\beta-\lambda)} \sum_{j=k-1}^{j_0+1} \frac{1}{2^{j(M-\alpha)}} \frac{w(B)}{R^{d-\beta+\lambda-\delta+M-\alpha}} \\ & \leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \sum_{k=1}^{j_0+1} 2^{k(\beta-\lambda-M+\alpha)} \frac{w(B)}{R^{d-\beta+\lambda-\delta+M-\alpha}} \\ & \leq \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \frac{w(B)}{R^{d-\beta+\lambda-\delta+M-\alpha}}. \quad \square \end{aligned}$$

Since by Proposition 1.1 the pointwise regularity conditions (1.17) and (2.3) characterize a Campanato-type space then the proof of Theorem 1.2 will follow from the

verification of those two conditions and using Remark 1.1.

LEMMA 5.4. *Let $\beta \geq 0$, $\beta > \lambda$ and $\delta \in \mathbb{R}$. If $(v, w) \in \mathcal{S}(\infty, \delta, \lambda)$ and $\alpha > 0$ then there exists a constant C such that for all $f \in BMO_{\mathcal{S}}^{\beta}(v)$,*

$$\frac{1}{w(B(x_B, \rho(x_B)))} \int_{B(x_B, \rho(x_B))} |\mathcal{I}_{\alpha} f(x)| dx \leq C \|f\|_{BMO_{\mathcal{S}}^{\beta}(v)} \rho(x_B)^{\beta - \lambda + \delta + \alpha} \tag{5.6}$$

for any $x_B \in \mathbb{R}^d$.

Proof. Let us denote $\rho := \rho(x_B)$, $B := B(x_B, \rho)$ and split, $f = f_1 + f_2$ with $f_1 = f \chi_{2B}$. Hence by inequality (2.8), Lemma 5.1 and the fact that $\alpha > 0$ it follows that

$$\begin{aligned} \frac{1}{w(B)} \int_B |\mathcal{I}_{\alpha} f_1(x)| dx &\leq \frac{1}{w(B)} \int_B \int_{2B} \frac{|f(y)|}{|x-y|^{d-\alpha}} dy dx \leq C \rho^{\alpha} \frac{1}{w(2B)} \int_{2B} |f(y)| dy \\ &\leq C \rho^{\beta - \lambda + \delta + \alpha} \|f\|_{BMO_{\mathcal{S}}^{\beta}(v)}. \end{aligned} \tag{5.7}$$

On the other hand, if $x \in B$ and $y \in (2B)^c$ then $|x-y| \geq |x_B-y|/2 \geq \rho$ and, by (4.2), $\rho(x) \leq C\rho$. Hence, by Lemma 2.1, given $N > 0$ there is a constant C such that

$$|k_t(x, y)| \leq C \frac{e^{-\frac{|x-y|^2}{t}}}{t^{\frac{d}{2}} \left(1 + \frac{\sqrt{t}}{\rho(x)}\right)^N} \leq C \rho^N \frac{e^{-\frac{|x-y|^2}{t}}}{t^{\frac{d+N}{2}}}. \tag{5.8}$$

Thus, if $N > \alpha - d$, using the change of variable $s = \frac{|x-y|^2}{t}$, we get

$$\begin{aligned} |\mathcal{I}_{\alpha} f_2(x)| &\leq \int_0^{\infty} t^{\frac{\alpha}{2}} \left(\int_{(2B)^c} |k_t(x, y)| |f(y)| dy \right) \frac{dt}{t} \\ &\leq C \rho^N \int_{(2B)^c} |f(y)| \left(\int_0^{\infty} \frac{e^{-\frac{|x-y|^2}{t}}}{t^{\frac{d+N-\alpha}{2}}} \frac{dt}{t} \right) dy \\ &\leq C \rho^N \int_0^{\infty} s^{\frac{d+N-\alpha}{2}} e^{-s} \frac{ds}{s} \int_{(2B)^c} \frac{|f(y)|}{|x-y|^{d+N-\alpha}} dy \\ &\leq C \rho^N \int_{(2B)^c} \frac{|f(y)|}{|x_B-y|^{d+N-\alpha}} dy. \end{aligned}$$

Since $(v, w) \in \mathcal{S}(\infty, \delta, \lambda, N_0)$ for some $N_0 \geq 0$, using (5.2) and taking $N > N_0 + \beta - \lambda + \alpha$, we obtain

$$|\mathcal{I}_{\alpha} f_2(x)| \leq C \rho^N \frac{w(B)}{\rho^{d-\alpha-\beta-\delta+\lambda+N}} \|f\|_{BMO_{\mathcal{S}}^{\beta}(v)} \leq C \rho^{\beta - \lambda + \alpha + \delta} \frac{w(B)}{\rho^d} \|f\|_{BMO_{\mathcal{S}}^{\beta}(v)}.$$

Hence

$$\frac{1}{w(B)} \int_B |\mathcal{I}_{\alpha} f_2(x)| dx \leq C \rho^{\beta - \lambda + \alpha + \delta} \|f\|_{BMO_{\mathcal{S}}^{\beta}(v)}. \tag{5.9}$$

With (5.7) and (5.9) we get the proof of the lemma. \square

LEMMA 5.5. Let $\beta \geq 0$, $\alpha > 0$, $\lambda < \beta$, $\alpha + \beta - \lambda < \delta_0$ and $\alpha + \beta - \lambda + \delta \leq d$. If w is a doubling weight and $(v, w) \in \mathcal{S}(\infty, \delta, \lambda)$, then there exists a constant C such that

$$|\mathcal{I}_\alpha f(x) - \mathcal{I}_\alpha f(y)| \leq C \|f\|_{BMO_{\mathcal{S}}^\beta(v)} (W_{\beta-\lambda+\alpha+\delta}(x, |x-y|) + W_{\beta-\lambda+\alpha+\delta}(y, |x-y|))$$

for $|x-y| < \rho(x)$, $f \in BMO_{\mathcal{S}}^\beta(v)$ and W defined as in (1.14).

Proof. Set $R = |x-y| < \rho(x)$ and denote $B = B(x, R)$. Then

$$\begin{aligned} |\mathcal{I}_\alpha f(x) - \mathcal{I}_\alpha f(y)| &\leq \int_{\mathbb{R}^d} \int_0^\infty |k_t(x, z) - k_t(y, z)| t^{\frac{\alpha}{2}} \frac{dt}{t} |f(z)| dz \\ &\leq \left(\int_0^{\rho(x)^2} + \int_{\rho(x)^2}^\infty \right) t^{\frac{\alpha}{2}} \int_{\mathbb{R}^d} |k_t(x, z) - k_t(y, z)| |f(z)| dz \frac{dt}{t} = I + J. \end{aligned}$$

By splitting J we get

$$J = \int_{\rho(x)^2}^\infty t^{\frac{\alpha}{2}} \left(\int_{|x-z| \leq 4\rho(x)} + \int_{|x-z| > 4\rho(x)} \right) |k_t(x, z) - k_t(y, z)| |f(z)| dz \frac{dt}{t} = J_1 + J_2.$$

Let $j_0 \in \mathbb{N} \cup \{0\}$ such that $2^{j_0}R \leq \rho(x) < 2^{j_0+1}R$. Since $\sqrt{t} > \rho(x) > R = |x-y|$ then by Lemma 2.2 and arguments similar to those used to obtain (5.8), given $N > 0$, $v < \delta_0$ and $L > 0$ there exists a constant $C = C_{v, N, L}$ such that

$$\begin{aligned} |k_t(x, z) - k_t(y, z)| &\leq C \left(\frac{R}{\sqrt{t}} \right)^v t^{-\frac{d}{2}} \left(1 + \frac{\sqrt{t}}{\rho(x)} \right)^{-N} e^{-\frac{|x-z|^2}{t}} \\ &\leq CR^v \rho(x)^N \times \begin{cases} t^{-\frac{d+v+N}{2}} & \text{if } \frac{|x-z|^2}{t} \leq 16 \\ t^{-\frac{d+v+N-L}{2}} |x-z|^{-L} & \text{if } \frac{|x-z|^2}{t} > 16. \end{cases} \end{aligned} \tag{5.10}$$

In the case $\frac{|x-z|^2}{t} \leq 16$ choosing v such that $\alpha + \beta - \lambda \leq v < \delta_0$ and $N > 0$ such that $N > d - \alpha + v$, using (5.10) and Lemma 5.1, since $\alpha + \beta - \lambda + \delta \leq d$ we can obtain

$$\begin{aligned} J_1 &\leq CR^v \rho(x)^N \int_{\rho(x)^2}^\infty t^{\frac{\alpha-d-v-N}{2}} \frac{dt}{t} \int_{|x-z| < 2^{j_0+3}R} |f(z)| dz \\ &\leq C \|f\|_{BMO_{\mathcal{S}}^\beta(v)} R^v \rho(x)^{\alpha-d-v} 2^{j_0(d+\beta-\lambda+N_0)} \left(\frac{R}{\rho(x)} \right)^{N_0} R^{\beta-\lambda+\delta} w(B) \\ &\leq C \|f\|_{BMO_{\mathcal{S}}^\beta(v)} \left(\frac{R}{\rho(x)} \right)^{-\beta+\lambda+v-\alpha} R^{\beta-\lambda+\alpha+\delta-d} w(B) \\ &\leq C \|f\|_{BMO_{\mathcal{S}}^\beta(v)} R^{\beta-\lambda+\alpha+\delta-d} w(B) \leq C \|f\|_{BMO_{\mathcal{S}}^\beta(v)} W_{\beta-\lambda+\alpha+\delta}(x, R) \end{aligned}$$

for some constants $C > 0$ and $N_0 \geq 0$.

In the case $\frac{|x-z|^2}{t} > 16$ setting $N_0 \geq 0$ such that $(v, w) \in \mathcal{S}(\infty, \delta, \lambda, N_0)$, v such that $\beta - \lambda + \alpha < v$, choosing $M_1 > N_0 + \beta - \lambda + \alpha - v$, defining $M = M_1 + v$ in

(5.2), choosing $N > M_1 + \nu$ and $L = d - \alpha + M_1 + \nu$ in (5.10) and finally assuming that $\beta - \lambda + \alpha + \delta \leq d$ there exists a constant C such that

$$\begin{aligned} J_2 &\leq C R^\nu \rho(x)^N \int_{\rho(x)^2}^{\infty} t^{\frac{\alpha-d-\nu-N}{2}} t^{\frac{d-\alpha+\nu+M_1}{2}} \frac{dt}{t} \int_{|x-z| \geq 4\rho(x)} \frac{|f(z)|}{|x-z|^{d-\alpha+\nu+M_1}} dz \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^{\beta}(\nu)} R^\nu \rho(x)^N \int_{\rho(x)^2}^{\infty} t^{\frac{-N+M_1}{2}} \frac{dt}{t} \left(\frac{R}{\rho(x)}\right)^{-\beta+\lambda-\alpha+\nu+M_1} \frac{R^{\beta-\lambda+\delta} w(B)}{R^{d-\alpha+\nu+M_1}} \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^{\beta}(\nu)} \left(\frac{R}{\rho(x)}\right)^{-\beta+\lambda+\nu-\alpha} R^{\beta-\lambda+\alpha+\delta-d} w(B) \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^{\beta}(\nu)} W_{\beta-\lambda+\alpha+\delta}(x, R). \end{aligned}$$

To deal with I we split

$$I \leq \int_0^{\rho(x)^2} t^{\frac{\alpha}{2}} \int_{\mathbb{R}^d} (|q_t(x, z) - q_t(y, z)| + |\tilde{k}_t(x, z) - \tilde{k}_t(y, z)|) |f(z)| dz \frac{dt}{t} = I_1 + I_2,$$

where \tilde{k} is the kernel of the classical heat operator $e^{-t\Delta}$ and $q_t = k_t - \tilde{k}_t$. For the first term above we get that

$$\begin{aligned} I_1 &\leq \int_0^{\rho(x)^2} t^{\frac{\alpha}{2}} \int_{\mathbb{R}^d} \left(\chi_{\{|x-z|>4R\}} |q_t(x, z) - q_t(y, z)| \right. \\ &\quad \left. + \chi_{\{|x-z|\leq 4R\}} |q_t(x, z)| + \chi_{\{|x-z|\leq 4R\}} |q_t(y, z)| \right) |f(z)| dz \frac{dt}{t} \\ &= I_{1,1} + I_{1,2} + I_{1,3}. \end{aligned}$$

By Lemma 2.4, in the case $t < \rho(x)^2$ and any $N > 0$ there exists C (independent of t)

$$|q_t(x, z) - q_t(y, z)| \leq C \left(\frac{R}{\rho(x)}\right)^\nu \frac{1}{t^{\frac{d}{2}} \left(1 + \frac{|x-z|}{\sqrt{t}}\right)^{d+N}} \leq C \left(\frac{R}{\rho(x)}\right)^\nu \frac{t^{\frac{N}{2}}}{|x-z|^{d+N}},$$

where $R = |x-y| < C\rho(x)$ and $|x-z| > 4R$.

To estimate $I_{1,1}$ we split $\{z : |x-z| > 4R\} = \{z : 4R < |x-z| \leq 4\rho(x)\} \cup \{z : |x-z| > 4\rho(x)\}$, define in the above inequality $N = \xi$ with $0 < \xi < \nu - (\alpha + \beta - \lambda)$ for the first domain and $N = N_2 > N_0 + \beta - \lambda$ for the second and use Lemma 5.2 and condition $\alpha + \beta - \lambda + \delta \leq d$ to get, for some constant C ,

$$\begin{aligned} I_{1,1} &\leq C \left(\frac{R}{\rho(x)}\right)^\nu \left(\int_0^{\rho(x)^2} t^{\frac{\alpha+\xi}{2}} \frac{dt}{t} \int_{\{4R < |x-z| \leq 4\rho(x)\}} \frac{|f(z)|}{|x-z|^{d+\xi}} dz \right. \\ &\quad \left. + \int_0^{\rho(x)^2} t^{\frac{\alpha+N_2}{2}} \frac{dt}{t} \int_{\{|x-z|>4\rho(x)\}} \frac{|f(z)|}{|x-z|^{d+N_2}} dz \right) \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^{\beta}(\nu)} \left(\left(\frac{\rho(x)}{R}\right)^{\xi+\beta-\lambda+\alpha-\nu} + \left(\frac{R}{\rho(x)}\right)^{-\beta+\lambda+\nu-\alpha} \right) \frac{R^{\beta-\lambda+\alpha+\delta} w(B)}{R^d} \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^{\beta}(\nu)} R^{\beta-\lambda+\alpha+\delta-d} w(B) \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^{\beta}(\nu)} W_{\beta-\lambda+\alpha+\delta}(x, R). \end{aligned} \tag{5.11}$$

On the other hand, by Lemma 2.3, for $\sqrt{t} < \rho$ we get

$$|q_t(x, z)| \leq C_M \left(\frac{\sqrt{t}}{\rho(x)} \right)^v t^{-\frac{d}{2}} \left(1 + \frac{|x-z|}{\sqrt{t}} \right)^{-M}.$$

If $\sqrt{t} < R/4$ then we set an integer $n_t \geq 2$ such that $2^{n_t} \sqrt{t} \leq R < 2^{n_t+1} \sqrt{t}$, use a dyadic decomposition and the above inequality to get

$$\begin{aligned} I_{1,2} &\leq C \left(\int_0^{\frac{R^2}{16}} \left(\frac{\sqrt{t}}{\rho(x)} \right)^v t^{\frac{\alpha-d}{2}} \right. \\ &\quad \times \left(\int_{|x-z| < 4\sqrt{t}} |f(z)| dz + \sum_{j=2}^{n_t} \int_{2^j \sqrt{t} \leq |x-z| < 2^{j+1} \sqrt{t}} \left(\frac{\sqrt{t}}{|x-z|} \right)^M |f(z)| dz \right) \frac{dt}{t} \\ &\quad \left. + \int_{\frac{R^2}{16}}^{\frac{\rho(x)^2}{16}} \left(\frac{\sqrt{t}}{\rho(x)} \right)^v t^{\frac{\alpha-d}{2}} \frac{dt}{t} \int_{|x-z| \leq 2R} |f(z)| dz \right) = H_1 + H_2. \end{aligned}$$

Now, since $\beta - \lambda > 0$ then we are able to apply Lemma 5.1, that w is doubling, $\alpha + \beta - \lambda + \delta \leq d$, $\beta - \lambda < \beta - \lambda + \alpha < v$ and choose $M > d$ to obtain

$$\begin{aligned} H_1 &\leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \int_0^{\frac{R^2}{16}} \left(\frac{\sqrt{t}}{\rho(x)} \right)^v t^{\frac{\alpha-d}{2}} \\ &\quad \times \left(4^d \left(\frac{\rho(x)}{\sqrt{t}} \right)^{\beta-\lambda} t^{\frac{\beta-\lambda+\delta}{2}} w(B(x, 4\sqrt{t})) + \sum_{j=2}^{n_t} 2^{-jM} \int_{|x-z| < 2^{j+1} \sqrt{t}} |f(z)| dz \right) \frac{dt}{t} \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \int_0^{\frac{R^2}{16}} \left(\frac{\sqrt{t}}{\rho(x)} \right)^v t^{\frac{\beta-\lambda+\alpha+\delta-d}{2}} \\ &\quad \times \left(\frac{\rho(x)}{\sqrt{t}} \right)^{\beta-\lambda} \left(1 + \sum_{j=2}^{n_t} 2^{-jM+jd} \right) w(B(x, 4\sqrt{t})) \frac{dt}{t} \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \int_0^{\frac{R^2}{16}} \left(\frac{\sqrt{t}}{\rho(x)} \right)^{v-\beta+\lambda} \frac{dt}{t} W_{\beta-\lambda+\alpha+\delta}(x, R) \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} W_{\beta-\lambda+\alpha+\delta}(x, R). \end{aligned}$$

Analogously, since $\beta - \lambda + \alpha + \delta \leq d$ and $\beta - \lambda < \beta - \lambda + \alpha < v \leq d$ then

$$\begin{aligned} H_2 &\leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \int_{\frac{R^2}{16}}^{\frac{\rho(x)^2}{16}} \left(\frac{\sqrt{t}}{\rho(x)} \right)^v t^{\frac{\alpha-d}{2}} \frac{dt}{t} 2^d \left(\frac{\rho(x)}{R} \right)^{\beta-\lambda} R^{\delta+\beta-\lambda} w(B(x, 4R)) \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \int_{\frac{R^2}{16}}^{\frac{\rho(x)^2}{16}} \left(\frac{\sqrt{t}}{\rho(x)} \right)^{v+\alpha-d} \frac{dt}{t} \left(\frac{\rho(x)}{R} \right)^{\beta-\lambda+\alpha-d} R^{\beta-\lambda+\delta+\alpha-d} w(B(x, R)) \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^\beta(v)} \int_{\frac{R^2}{16}}^{\frac{\rho(x)^2}{16}} \left(\frac{\sqrt{t}}{\rho(x)} \right)^{\beta-\lambda+\alpha-d} \left(\frac{\sqrt{t}}{\rho(x)} \right)^{v-\beta+\lambda} \frac{dt}{t} \\ &\quad \times \left(\frac{\rho(x)}{R} \right)^{\beta-\lambda+\alpha-d} W_{\beta-\lambda+\alpha+\delta}(x, R) \end{aligned}$$

$$\begin{aligned} &\leq C\|f\|_{BMO_{\mathcal{L}}^\beta(v)} \int_0^{\frac{\rho(x)^2}{16}} \left(\frac{\sqrt{t}}{\rho(x)}\right)^{\nu-\beta+\lambda} \frac{dt}{t} W_{\beta-\lambda+\alpha+\delta}(x, R) \\ &\leq C\|f\|_{BMO_{\mathcal{L}}^\beta(v)} W_{\beta-\lambda+\alpha+\delta}(x, R). \end{aligned}$$

Hence

$$I_{1,2} \leq C\|f\|_{BMO_{\mathcal{L}}^\beta(v)} W_{\beta-\lambda+\alpha+\delta}(x, R). \tag{5.12}$$

Since $I_{1,3}$ is similar to $I_{1,2}$ we also obtain

$$I_{1,3} \leq C\|f\|_{BMO_{\mathcal{L}}^\beta(v)} W_{\beta-\lambda+\alpha+\delta}(x, R). \tag{5.13}$$

Inequalities (5.11), (5.12) and (5.13) give the seek estimate of I_1 .

On the other hand, since $\int_{\mathbb{R}^d} (\tilde{k}_t(x, z) - \tilde{k}_t(y, z)) dz = 0$ then

$$\begin{aligned} I_2 &\leq \int_0^{\rho(x)^2} t^{\frac{\alpha}{2}} \int_{\mathbb{R}^d} |\tilde{k}_t(x, z) - \tilde{k}_t(y, z)| |f(z) - f_B| dz \frac{dt}{t} \\ &\leq \int_0^{\rho(x)^2} t^{\frac{\alpha}{2}} \left(\int_{|x-z|>2R} |\tilde{k}_t(x, z) - \tilde{k}_t(y, z)| |f(z) - f_B| dz \right. \\ &\quad \left. + \int_{|x-z|\leq 2R} |\tilde{k}_t(x, z)| |f(z) - f_B| dz + \int_{|x-z|\leq 2R} |\tilde{k}_t(y, z)| |f(z) - f_B| dz \right) \frac{dt}{t} \\ &= I_{2,1} + I_{2,2} + I_{2,3}. \end{aligned}$$

It is not difficult to see that $|\tilde{k}_t(x, z) - \tilde{k}_t(y, z)| \leq Ce^{-\frac{|x-z|^2}{ct}} t^{-\frac{d}{2}-1} |x-y||x-z|$ for $|x-z| > 2|x-y|$ and $\alpha < \beta - \lambda + \alpha < \delta_0 \leq 1$ then, choosing $M > N_0 - \lambda + \alpha + \beta - 1$ and using (5.5) and (5.4) in Lemma 5.3, it follows that

$$\begin{aligned} I_{2,1} &\leq CR \int_{|x-z|>2R} |x-z| |f(z) - f_B| \int_0^{\rho(x)^2} t^{\frac{\alpha-d-2}{2}} e^{-\frac{|x-z|^2}{ct}} \frac{dt}{t} dz \\ &\leq CR \int_{|x-z|>2R} \frac{|f(z) - f_B|}{|x-z|^{d+1-\alpha}} \int_{\frac{|x-z|^2}{c\rho(x)^2}}^\infty s^{\frac{d+2-\alpha}{2}} e^{-s} \frac{ds}{s} dz \\ &\leq CR \left(\int_{2R < |x-z| \leq 2\rho(x)} \frac{|f(z) - f_B|}{|x-z|^{d+1-\alpha}} dz \left(\int_0^\infty s^{\frac{d+2-\alpha}{2}} e^{-s} \frac{ds}{s} \right) \right. \\ &\quad \left. + \int_{|x-z|>2\rho(x)} \frac{|f(z) - f_B|}{|x-z|^{d+1-\alpha}} \int_{\frac{|x-z|^2}{c\rho(x)^2}}^\infty s^{\frac{d+2-\alpha-M}{2}} e^{-\frac{s}{c}} \frac{ds}{s} dz \right) \\ &\leq CR \left(\int_{2R < |x-z| \leq 2\rho(x)} \frac{|f(z) - f_B|}{|x-z|^{d+1-\alpha}} dz + \rho(x)^M \int_{|x-z|>2\rho(x)} \frac{|f(z) - f_B|}{|x-z|^{d-\alpha+1+M}} dz \right) \\ &\leq C\|f\|_{BMO_{\mathcal{L}}^\beta(v)} \frac{w(B)}{R^{d-\beta-\delta+\lambda-\alpha}} \leq C\|f\|_{BMO_{\mathcal{L}}^\beta(v)} W_{\beta-\lambda+\alpha+\delta}(x, R). \end{aligned}$$

For $I_{2,2}$ we use that $(v, w) \in \mathcal{S}(\infty, \delta, \lambda)$ and that w is doubling to get

$$\begin{aligned} I_{2,2} &\leq C \int_{|x-z| \leq 2R} |f(z) - f_B| \int_0^{\rho(x)^2} t^{\frac{\alpha-d}{2}} e^{-\frac{|x-z|^2}{ct}} \frac{dt}{t} dz \\ &\leq C \int_{|x-z| \leq 2R} \frac{|f(z) - f_B|}{|x-z|^{d-\alpha}} \left(\int_{\frac{|x-z|^2}{cp(x)^2}}^{\infty} s^{\frac{-\alpha+d}{2}} e^{-s} \frac{ds}{s} \right) dz \\ &\leq C \int_{|x-z| \leq 2R} \frac{|f(z) - f_B|}{|x-z|^{d-\alpha}} dz \leq C \sum_{j=0}^{\infty} (2^{-j}R)^{\alpha-d} \int_{|x-z| < 2^{-j+1}R} |f(z) - f_B| dz \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^{\beta}(v)} \sum_{j=0}^{\infty} (2^{-j}R)^{\beta+\alpha-d} v(2^{-j+1}B) \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^{\beta}(v)} \sum_{j=0}^{\infty} (2^{-j}R)^{\beta-\lambda+\alpha+\delta-d} w(2^{-j+1}B) \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^{\beta}(v)} \sum_{j=1}^{\infty} (2^{-j}R)^{\beta-\lambda+\alpha+\delta-d} w(2^{-j+1}B \setminus 2^{-j}B) \\ &\leq C \|f\|_{BMO_{\mathcal{L}}^{\beta}(v)} \int_{|x-z| \leq 2R} \frac{w(z)}{|x-z|^{d-\beta+\lambda-\alpha-\delta}} dz = C \|f\|_{BMO_{\mathcal{L}}^{\beta}(v)} W_{\beta-\lambda+\alpha+\delta}(x, R). \end{aligned}$$

The estimate for $I_{2,3}$ is obtained in the same way. \square

5.1. Proof of Theorem 1.2

Since $\beta - \lambda + \alpha + \delta < 1$ and w is doubling then to prove this Theorem it will be enough to use the pointwise characterization of the space $BMO_{\mathcal{L}}^{\alpha+\beta-\lambda+\delta}(w)$ provided by Proposition 1.1. But then Lemma 5.4 and Lemma 5.5 give the two desired inequalities for $\mathcal{I}_{\alpha}f$.

The authors are deeply grateful to the referee for his interesting suggestions to make this paper more clear and complete.

REFERENCES

- [1] B. BONGIOANNI, A. CABRAL AND E. HARBOURE, *Schrödinger type singular integrals: weighted estimates for $p = 1$* , Math. Nach., **289** (11–12): 1341–1369, 2016.
- [2] B. BONGIOANNI, E. HARBOURE AND O. SALINAS, *Weighted inequalities for negative powers of Schrödinger operators*, J. Math. Anal. Appl. **348** (1): 12–27, 2008.
- [3] B. BONGIOANNI, E. HARBOURE AND O. SALINAS, *Riesz transforms related to Schrödinger operators acting on BMO type spaces*, J. Math. Anal. Appl. **357** (1): 115–131, 2009.
- [4] B. BONGIOANNI, E. HARBOURE AND O. SALINAS, *Classes of weights related to Schrödinger operators*, J. Math. Anal. Appl. **373** (2): 563–579, 2011.
- [5] C. PÉREZ AND R. L. WHEEDEN, *Uncertainly principle estimates for vector fields*, J. Func. Anal., **141**: 146–188, 2001.
- [6] D. CRUZ-URIBE AND C. PÉREZ, *Two-weight, weak-type norm inequalities for fractional integrals, Calderón-Zygmund operators and commutators*, Indiana Math. J., **49**: 697–721, 2000.
- [7] D. CRUZ-URIBE AND C. PÉREZ, *On the two-weight problem for singular integral operators*, Ann. Scuola Norm. Sup. Pisa CL. Sci., I (5): 821–849, 2002.

- [8] D. CRUZ-URIBE, J. M. MARTELL AND C. PÉREZ, *Sharp two-weight inequalities for singular integrals, with applications to the Hilbert transform and the Sarason conjecture*, Adv. Math., **216** (2): 647–676, 2007.
- [9] D. CRUZ-URIBE, A. REZNIKOV AND A. VOLBERG, *Logarithmic bump conditions and the two-weight boundedness of Calderón-Zygmund operators*, Adv. in Math., **255**: 706–729, 2014.
- [10] J. DZIUBÁŃSKI, G. GARRIGÓS, T. MARTÍNEZ, J. TORREA AND J. ZIENKIEWICZ, *BMO spaces related to Schrödinger operators with potentials satisfying a reverse Hölder inequality*, Math. Z., **249** (2): 329–356, 2005.
- [11] J. DZIUBÁŃSKI AND J. ZIENKIEWICZ, *Hardy spaces H^1 associated to Schrödinger operators with potential satisfying reverse Hölder inequality*, Rev. Mat. Iberoamericana, **15** (2): 279–296, 1999.
- [12] J. DZIUBÁŃSKI AND J. ZIENKIEWICZ, *H^p spaces for Schrödinger operators*, Fourier Analysis and Related Topics (Bedlewo) – Banach Center of Publ., Polish Acad. Sci. Inst Math., Warsaw, **56**: 45–53, 2002.
- [13] J. DZIUBÁŃSKI AND J. ZIENKIEWICZ, *H^p spaces associated with Schrödinger operator with potential from reverse Hölder classes*, Colloq. Math., **98** (1):5–38, 2003.
- [14] T. EKHOLOM AND R. L. FRANK, *On Lieb-Thirring inequalities for Schrödinger operators with virtual level*, Commun. Math. Phys., **264**: 725–740, 2006.
- [15] J. GARCÍA-CUERVA AND J. M. MARTELL, *Two-weight norm inequalities for maximal operators and fractional integrals on non-homogeneous spaces*, Indiana Univ. Math. J., **5** (3):1241–1280, 2001.
- [16] F. W. GEHRING, *The L^p -integrability of the partial derivatives of a quasiconformal mappings*, Acta Math., **130**: 265–277, 1973.
- [17] E. HARBOURE, O. SALINAS AND B. VIVIANI, *Boundedness of the fractional integral on weighted Lebesgue and Lipschitz spaces*, Trans. Amer. Math. Soc., **349** (1): 235–255, 1997.
- [18] G. H. HARDY AND J. E. LITTLEWOOD, *Some properties of fractional integrals I*, Math. Zeit., **27**: 565–606, 1927.
- [19] G. H. HARDY AND J. E. LITTLEWOOD, *Some properties of fractional integrals II*, Math. Zeit., **34**: 403–439, 1932.
- [20] S. HARTZSTEIN, O. SALINAS AND M. A. VIGNATTI, *Two-weighted inequalities for maximal operators related to Schrödinger differential operators*, Forum Mathematicum (to appear), 2020.
- [21] L. HEDBERG, *On certain convolution inequalities*, Proc. Amer. Math. Soc., **36**: 505–510, 1972.
- [22] S. JANSON, *On functions on the mean oscillation*, Arb. Mat., **14**: 189–196, 1976.
- [23] K. KURATA, *An estimate on the heat kernel of magnetic Schrödinger operators and uniformly elliptic operators with non-negative potentials*, J. London Math. Soc.. Second Series, **62** (3): 885–903, 2000.
- [24] T. MA, P. R. STINGA, J. L. TORREA AND C. ZHANG, *Regularity estimates in Hölder spaces for Schrödinger operators via a $T1$ theorem*, J. Math. Anal. Appl., **388**: 817–837, 2012.
- [25] T. MA, P. R. STINGA, J. L. TORREA AND C. ZHANG, *Regularity properties of Schrödinger operators*, Annali di Matematica, **193**: 561–589, 2014.
- [26] N. G. MEYERS, *Mean oscillation over cubes and Hölder continuity*, Proc. Amer. Math. Soc., **15**: 717–721, 1964.
- [27] K. MOEN, *Weighted inequalities for multilinear fractional integral operators*, Collect. Math., **60** (2): 213–238, 2009.
- [28] J. PAN AND W. SUN, *Two-weight norm inequalities for fractional maximal functions and fractional integral operators on weighted Morrey spaces*, Math. Nach. (to appear), pages 1–13, 2020.
- [29] J. PEETRE, *On the theory of $\mathcal{L}_{p,\lambda}$ spaces*, J. Func. Anal., **4**: 71–87, 1969.
- [30] C. PÉREZ, *Two weighted norm inequalities for Riesz potentials and uniform L^p -weighted Sobolev inequalities*, Indiana Univ. Math. J., **39** (1): 31–44, 1990.
- [31] C. PÉREZ, *Two weighted inequalities for potential and fractional type maximal operators*, Indiana Math. J., **43**: 663–683, 1994.
- [32] C. PÉREZ, *On sufficient conditions for the boundedness of the Hardy-Littlewood maximal operator between weighted L^p -spaces with different weights*, Proc. London Math. Soc. (3), **71** (1): 135–157, 1995.
- [33] G. PRADOLINI, *A class of pair of weights related to the boundedness of the fractional integral operator between L^p and Lipschitz spaces*, Comment. Math. Univ. Carolinae, **42** (1): 133–152, 2001.
- [34] G. PRADOLINI, *Two-weighted norm inequalities for the fractional integral operator between L^p and Lipschitz spaces*, Comment. Math. (Prace Mat.), **41**: 147–169, 2001.

- [35] G. PRADOLINI AND O. SALINAS, *The fractional integral between weighted Orlicz and BMO_ϕ spaces on spaces of homogeneous type*, Comment. Math. Univ. Carolinae, **44** (3): 469–487, 2003.
- [36] Z. W. SHEN, *L^p estimates for Schrödinger operators with certain potentials*, Ann. Inst. Fourier (Grenoble), **45** (2): 513–546, 1995.
- [37] S. L. SOBOLEV, *On a theorem of Marcinkiewicz concerning interpolation of operators*, Jour. de Math. Pures et Appliquées, **35**: 223–248, 1956.
- [38] S. SPANNE, *Some function spaces defined using the mean oscillation over cubes*, Ann. Scuola Norm. Sup. Pisa CL. Sci., **19**: 593–608, 1965.

(Received October 24, 2018)

R. Crescimbeni
Dept. de Matemática
Fac. de Economía y Administración, UNCo e IITCI-CONICET
Neuquén, Argentina
e-mail: raquel.crescimbeni@comahue-conicet.gob.ar

S. Hartzstein
Dept. de Matemática
Fac. de Ingeniería Química, UNL e IMAL-CONICET
Santa Fe, Argentina
e-mail: shartzstein@gmail.com

O. Salinas
Dept. de Matemática
Fac. de Ingeniería Química, UNL e IMAL-CONICET
Santa Fe, Argentina
e-mail: salinas@santafe-conicet.gov.ar

