

Montana Tech Library

Digital Commons @ Montana Tech

Safety Health & Industrial Hygiene

Faculty Scholarship

2021

Hazardous Drug Wipe Sampling in Healthcare Facilities

Brandi Gruenewald

David P. Gilkey

Follow this and additional works at: <https://digitalcommons.mtech.edu/shih>



Part of the [Other Pharmacy and Pharmaceutical Sciences Commons](#), and the [Pharmacy Administration, Policy and Regulation Commons](#)



2021

ISSN 1015-5589, Vol. XXX, No. 2

World Safety Journal

© Copyright 2021 | WSO



Traffic Congestion in Beirut, Lebanon: Environmental Impact, Financial Consequences, and Possible Solutions!
by E.M. Choueiri

Hazardous Drug Wipe Sampling in Healthcare Facilities
by B. Gruenewald and D. Gilkey

Prevalence and Determinants of Frailty: A Cross-Sectional Study from Rural and Urban Communities in South Lebanon
by F. al-Lababidi, A. Khalil, N. Bassil, E.M. Choueiri, and H. Akoum

Why the Industry is Losing Faith in HSE Awards? 250 Professionals Rethink!
by H. Lal

Trip Attraction Models, Using Regression Analysis, for Shopping Malls in Lebanon
by R. Wehbe, E. Otayek, and E.M. Choueiri

WSO's Certification Program is
accredited in compliance with
ISO/IEC 17024:2012



World Safety Organization

Statement of Purpose and Objective

WSO's purpose is to internationalize all safety fields, including occupational and environmental safety and health, accident prevention movement, etc., and to disseminate throughout the world the practices skills, arts, and technologies of safety and accident prevention.

WSO's objective is to protect people, property, resources, and the environment on local, regional, national, and international levels. WSO membership is open to all individuals and entities involved in the safety and accident prevention field, regardless of race, color, creed, ideology, religion, social status, sex, or political beliefs.

WSO is in Consultative Category II Status (Non-Governmental Organization - NGO) to the Economic and Social Council of the United Nations.

WSO is a non-profit, non-sectarian, non-political organization, dedicated to:

“Making Safety a Way of Life ... Worldwide.”



Editor-in-Chief

Prof. Dr. Elias M. Choueiri

*Member of the WSO
Board of Directors
and WSO Liaison Officer
to the United Nations*

Associate Editor

Dr. Janis K. Jansz

*Curtin University, Western Australia
Director of the WSO National Office for Australia, and
Member of the WSO Board of Directors*

Editorial Board

Prof. Michael Daramola

*University of Pretoria
South Africa*

Dr. Richard Franklin

*James Cook University
Queensland, Australia*

Dr. Charles H. Baker

*World Safety Organization
USA*

Dr. Milos Nedved

*Central Queensland University
Australia*

Dr. David P. Gilkey

*Montana Tech University
USA*

Ms. Zahra Etemadi

*Fremantle Ports
Australia*

Cover Photo

On Tuesday 4th August 2020, at 6:09 p.m. local time, a large explosion occurred at a warehouse in the Port of Beirut, Lebanon, which resulted not only in the destruction of buildings and infrastructure in a large section of Beirut, but also devastated the lives of many people who were just going about their daily lives, leaving some 200 deaths, 6,000 injured, 300,000 people homeless and an unknown number unaccounted for in its wake. Reconstruction cost of the various buildings and infrastructure is currently estimated at some \$15 billion, whereas the rebuilding of the lives of many people in Beirut and beyond will take much more than just money. Lebanon cannot do this on its own, as it is highly in debt and its finances are nearly depleted – the country itself is on the brink of collapse. Assistance from the international community and financial institutions is very much needed, in order to aid this country in its endeavors to pick up the rubble, dust down the pieces and start rebuilding the various buildings and infrastructure, as well as the lives and livelihoods of its people – so that there may be hope for both Lebanon and its people!

Table of Contents

Traffic Congestion in Beirut, Lebanon: Environmental Impact, Financial Consequences, and Possible Solutions! by E.M. Choueiri.....	1-8
Hazardous Drug Wipe Sampling in Healthcare Facilities by B. Gruenewald and D. Gilkey	9-13
Prevalence and Determinants of Frailty: A Cross-Sectional Study from Rural and Urban Communities in South Lebanon by F. al-Lababidi, A. Khalil, N. Bassil, E.M. Choueiri, and H. Akoum	14-26
Why the Industry is Losing Faith in HSE Awards? 250 Professionals Rethink! by H. Lal	27-36
Trip Attraction Models, Using Regression Analysis, for Shopping Malls in Lebanon by R. Wehbe, E. Otayek, and E.M. Choueiri	37-47

Article Submission

The World Safety Journal (WSJ) is a quarterly refereed journal (ISSN: 1015-5589) that includes original full length papers, review articles and case studies on the most recent outcomes of advanced research in any occupational safety and health fields.

All articles shall be written in concise English and typed with a minimum font size of 12 point. Articles should have an abstract of not more than 300 words. Articles shall be submitted as Times New Roman print and presented in the form the writer wants published. On a separate page, the author should supply the author's name, contact details, professional qualifications, current employment position, a brief bio, and a photo of the author. This should be submitted with the article.

Writers should include all references and acknowledgments. Authors are responsible for ensuring that their works do not infringe on any copyright. Failure to do so can result in the writer being accountable for breach of copyright. The accuracy of the references is the author's responsibility.

References

Articles should be referenced according to the Publication Manual of the American Psychological Association, 7th ed.

Books are referenced as follows:

Author. (Year of publication). *Title of publication*. Publisher.

Articles are referenced as follows:

Author (Year). Title of article. *Name of Journal*. Volume (Issue), Page numbers of article.

Internet information is referenced as follows:

Name of author. (Year of publication). *Title*. DOI or web address if no DOI.

Submissions should be mailed to:

World Safety Organization

Attn: Editorial Staff /

Prof. Dr. Elias M. Choueiri

World Management Center,

106 W Young St #F

Warrensburg, MO 64093, USA

or emailed to: elias.choueiri@gmail.com

Articles, wherever possible, must be up-to-date and relevant to the Safety Industry. ***All articles are Blind Peer Reviewed by at least two referees before being accepted for publication***

Disclaimer

Opinions expressed by contributors in articles or reproduced articles are the individual opinions of such contributors or the authors and not necessarily those of the World Safety Organization. Reproduction of articles or abstracts contained in this journal is approved providing the source is acknowledged.



World Safety Journal

A peer-reviewed journal,
published by the World Safety Organization



Hazardous Drug Wipe Sampling in Healthcare Facilities

Brandi Gruenewald and David Gilkey^(*)

Montana Technological University, 1300 West Park, Butte, MT, USA

Keywords

Pharmaceutical hazards
Antineoplastic drugs
Wipe samples
Industrial hygiene.

Abstract

Hazardous drugs are associated with causing acute and chronic side effects to healthcare workers that experience occupational exposures. Antineoplastic drugs are known to cause headaches, nausea, vomiting, hair loss, mutagenic activity, spontaneous abortions, infertility, and congenital malformations. Currently, there are no acceptable thresholds for exposures to this type of hazardous drugs. The use of as low as reasonably acceptable (ALARA) is used for exposures to these types of drugs. Occupational exposure risk should be evaluated within facilities where they are used. Performing hazardous drug wipe sampling in areas that are high risk for contamination can provide information to facilities on how to protect their employees.

Introduction and Background

Antineoplastic drugs are a workplace hazard. These drugs are known to be toxic to cells that are non-cancerous (Vyas, Yiannakis, Turner and Sewell, 2013). These types of drugs are associated with adverse side effects for employees with both acute and chronic exposures. Some of the earliest reports of these drugs posing occupational risks was in 1979 (Soteriades et al., 2020). The levels were quantifiable in the urine of nurses handling these mutagenic drugs. Previously, the worker exposures were higher levels (e.g., mg/mL) and currently the exposures are much lower. Most exposures recently are nanograms per milliliter (ng/mL) (Soteriades et al., 2020). Since the 1940's, the toxicity of cancer treatments has been known to cause side-effects to both patients and to the healthcare workers handling these drugs while performing their daily duties (Soteriades et al., 2020). Because healthcare workers handle these toxic drugs, the occupational risk should be evaluated.

The symptoms associated with occupational exposure to Antineoplastic drugs include headaches, nausea, vomiting, hair loss, hypersensitivity, mutagenic activity, spontaneous abortions, infertility, and congenital malformations (Vyas, Yiannakis, Turner and Sewell, 2013). These symptoms have been reported in healthcare workers that are being exposed to these cytotoxic drugs at much lower doses than patients (Dugheri et al, 2018). The drugs are also known to cause irritation and/or damage to the skin, eyes, and mucous membranes. In another study, it was been shown that the compounds were

^(*) *Corresponding Author*: Dgilkey@mtech.edu

mutagenic to mammalian cells in cell cultures (Harrison, Peters, and Bing, 2006). Antineoplastics have no therapeutic relevance to individuals that do not require these types of drug therapies.

By 2020, it was expected that there will be a rise in yearly cancer diagnoses to 16 million, globally (Dugheri et al., 2018). The market for cancer treatment drugs is expected to generate approximately US \$161.3B by the end of 2021. Treatments included in the estimate are chemotherapy, hormone therapy, immunotherapy, and targeted therapy, with chemotherapy projected to be 50% of the revenue. Antineoplastic drugs are classified as hazardous chemicals by National Institute for Occupational Safety and Health (NIOSH) (Dugheri et al., 2018).

NIOSH published the Preventing Occupational Exposures to Antineoplastic and Other Hazardous Drugs in Health Care Settings Alert in 2004. The list was recently updated in May 2020 to include newly approved drugs by the Food and Drug Administration (NIOSH, 2020). Of the drugs on this list from NIOSH, roughly half of them are antineoplastics. The purpose of this alert was to bring awareness to employees the risk involved with the handling of these drugs and outline protective measures they could implement for their facility (Fuller, Bain, Sperrazza, and Mazzuckelli, 2007).

Currently, there is no safe occupational threshold, such as a Permissible Exposure Limit (PEL), for hazardous drugs, however, there have been proposals for possible threshold limits for single drugs only (NIOSH, 2020). Healthcare facilities are trying to minimize the occupational exposure by utilizing the NIOSH hierarchy of controls, by implementing engineering controls, administrative controls and personal protective equipment, as well as environmental monitoring (i.e. wipe sampling) and biological monitoring (laboratory testing) as a method to identify problems and create more worker awareness (Dugheri et al., 2018).

The occupational exposure to hazardous drugs by healthcare workers is proposed to happen most commonly by dermal contact. It is not likely that healthcare workers are exposed via inhalation of the hazardous drugs. Hazardous drugs that have a low molecular weight (< 500 Daltons) are of concern because these drugs are easily absorbed through the skin, whereas some of the current hazardous drugs have a molecular weight of >40,000 Daltons (Conner and Smith, 2016; Connor, Zock and Snow, 2016).

The larger molecular weight limits the dermal uptake from contaminated surfaces. However, nurses have a higher risk of exposure and possible dermal uptake of these higher molecular weight drugs due to constant hand-washing practices, which damages their skin and causes cracks that these drugs can penetrate through (Conner and Smith, 2016; Connor, Zock and Snow, 2016).

Wipe sampling is one of the most common practices for hazardous drug contamination assessment (Conner and Smith, 2016; Connor, Zock and Snow, 2016). Environmental monitoring has shown that hazardous drugs can be found in the air and on work surfaces in sterile compounding rooms, manufacturing and packaging areas for the compounded sterile products and clinical administration areas (Harrison, Peters, and Bing, 2006). Performing hazardous drug wipe sampling in areas at risk for exposure can provide information for environmental monitoring on cleaning processes and handling of hazardous materials.

Environmental wipe sampling has been used for the last 20 years in healthcare facilities to evaluate contamination within the workplace. Conner, Zock and Snow (2016) stated that other reasons for surface sampling is as follows: hazard identification and evaluation, exposure assessment, facility characterization, housekeeping, selection of engineering controls, evaluation of engineering and administrative/work practice controls, evaluation of exposure pathways, selection of personal protective equipment, compliance with regulations and standards, source identification, education and training, and investigation of complaints. The samples that were collected in the facilities of were not

used for worker exposure but to look at the environmental contamination as a possibility for worker exposure (Conner and Smith, 2016; Connor, Zock and Snow, 2016).

Recommended Wipe Sampling Methods

Wipe sampling was originally developed to evaluate other agents such as lead, asbestos, methamphetamine, and antibiotics (Connor and Smith, 2016; Connor, Zock and Snow, 2016). This methodology was evaluated and applied to hazardous drugs. However, not all hazardous drugs can be analyzed because not all drugs have methods designed to analyze them in a laboratory. Additional methods can be developed for other drugs that do not currently have a testing method determined, as long as there are antibodies available for the drug (Conner and Smith, 2016; Connor, Zock and Snow, 2016).

When performing hazardous wipe sampling, there is a need to have a strategy in mind for which factors are to be assessed and what variables may be present in the sampling evaluation. Some factors to consider are the types of hazardous drugs that are being used and quantities stored and used within the facility. Once a sampling plan has been completed, a certified laboratory and/or industrial hygiene professional that conducts this type of sampling should be identified. Currently, there are no standards for sampling and analysis for these drugs, therefore it is essential to identify a laboratory that is experienced with hazardous drug wipe sampling analysis (Power, Sessink, Gesy, and Charbonneau, 2014). The laboratory should have a validation process for the drugs their facility evaluates. These validation methods should include how samples are stored for stability, medium desorption efficiency, limits of detection and quantitation, calibration curves and quality control methods. Their method should give greater than 90% extraction efficiency, which is preferred, however greater than 75% extraction is acceptable (Conner & Smith, 2016).

When hazardous drug wipe sampling is performed, the sample size should be no less than 100 cm², however if a smaller sample size is used, more samples must be taken which can increase the cost of this testing since the cost is per wipe rather than per drug being tested for. A more acceptable size would be 400 cm², which would give a larger sampling area and reduce the cost by not needing as many samples for one location. A sampling plan should be devised so that the facility knows exactly what locations were sampled and make note of what type of activity takes place in those locations (i.e., surface of the biological safety cabinet – admixing of hazardous drugs). Common locations for hazardous drug wipe sampling would be the geometric center of the engineering controls (biological safety cabinet or compounding aseptic containment isolator), where the direct compounding area is located. The floors directly below the engineering controls, pass-throughs from the hazardous drug storage into the negative pressure hazardous drug ISO Class 7 buffer areas and the pass through from the hazardous buffer areas to the general pharmacy, equipment, counters, storage containers, door handles, high touch areas, and computer keyboards (Conner and Smith, 2016; Connor, Zock and Snow, 2016).

Once the sampling has been completed, samples are sent overnight to a laboratory where the samples are processed and analyzed. Typical methods used for specimen recovery and analysis are gas chromatography, liquid chromatography, high-performance liquid chromatography, ultra-high-performance liquid chromatography along with mass spectrometry, tandem mass spectrometry or inductively coupled plasma mass spectrometry. These methods determine the concentration of hazardous drugs present on the wipe samples that have been collected within the healthcare facility (Conner and Smith, 2016; Connor, Zock and Snow, 2016).

Summary and Conclusions

Currently, no standards or regulations exist for an acceptable level of exposure to hazardous drug, and nothing is known about the synergistic effects multiple drugs could elicit in human systems. The only allowable standard for exposure is as low as reasonably achievable (ALARA). A common approach is put in place workplace controls, as well as implementation of personnel training on the handling of hazardous drugs, cleaning, deactivation and decontamination of work surfaces and surveys given to healthcare staff involved with the processes of admixture and administration of these therapies could work to lower potential contamination and the exposure of employees to these toxic drugs in the workplace (Soteriades, et al., 2020). Despite the usual healthcare protocols, workers may not be fully protected in their facilities. The overwhelming evidence is that an occupational risk persists for those handling antineoplastic drugs. The highest risk groups include pharmacists and their team that compounds antineoplastic drugs and the nursing staff that administers these drugs to patients (Vyas, Yiannakis, Turner and Sewell, 2013). Surface sampling should be used as part of an environmental monitoring program. In doing so, results will inform staff the larger picture of possible contamination and exposures on within the facility (Conner and Smith, 2016; Connor, Zock and Snow, 2016).

References

- Connor, T. & Smith, J. (2016). New Approaches to Wipe Sampling Methods for Antineoplastic and Other Hazardous Drugs in Healthcare Settings. *Pharmaceutical Technology in Hospital Pharmacy*, 1(3), 107-114.
- Connor, T., Zock, M. & Snow, A. (2016). Surface wipe sampling for antineoplastic (chemotherapy) and other hazardous drug residue in healthcare settings: Methodology and recommendations. *Journal of Occupational and Environmental Hygiene*, 13(9), 658-667.
- Dugheri, S., Bonari, A., Pompilio, I., Boccalon, P, Tognoni, D., Cecchi, M. & Arcangeli, G. (2018). Analytical strategies for assessing occupational exposure to antineoplastic drugs in healthcare workplaces. *Medycyna Pracy*, 69(6), 589-603.
- Fuller, T., Bain, E., Sperrazza, K. & Mazzuckelli, L. (2007). A Survey of the Status of Hazardous Drug Awareness and Control in a Sample Massachusetts Nursing Population. *Journal of Occupational and Environmental Hygiene*, 4(11), D113-D119
- Harrison, B., Peters, B. & Bing, M. (2006). Comparison of surface contamination with cyclophosphamide and fluorouracil using a closed-system drug transfer device versus standard preparation techniques. *American Journal of Health-System Pharmacy*, 63(18), 1736-1744.
- NIOSH. (2020). *Hazardous drug exposures in healthcare*. Retrieved from <https://www.cdc.gov/niosh/topics/hazdrug/default.html>
- Power, L., Sessink, P., Gesy, K. & Charbonneau, F. (2014). Hazardous Drug Residue on Exterior Vial Surfaces: Evaluation of a Commercial Manufacturing Process. *Hospital Pharmacy*, 49(4), 355-362.
- Soteriades, E., Economidou, S., Tsivitanidou, A., Polyviou, P., Lorimer, A., Katodritis, N. & Theophanous-K. (2020). Environmental assessment of cytotoxic drugs in the Oncology Center of Cyprus. *PloS One*, 15(3), E0216098.
- Vyas, N., Yiannakis, D., Turner, A. & Sewell, G. (2013). Occupational exposure to anti-cancer drugs: A review of effects of new technology. *Journal of Oncology Pharmacy Practice*, 20(4), 278-287.

Authors

Brandi GRUENEWALD is Operations Manager of Advanced Infusion Therapeutics, LLC, dba Advanced 797, a pharmacy cleanroom certification, consulting, and education company. She is currently a master's degree graduate student at Montana Technological University in the Industrial Hygiene program. She received her B.S. in Molecular Biology from Northeastern State University, in Broken Arrow, Oklahoma. She is a nationally registered CETA Certification Professional in Sterile Compounding Facilities (RCP-SCF). Her professional focus is working with pharmacy professionals to develop, test and maintain high-quality pharmacy compounding environments and achieve regulatory compliance.



David P. GILKEY, D.C., Ph.D., CPE, CSP, REHS is an Associate Professor at Montana Technological University, 1300 West Park St. Butte, MT 59701. He is an Associate Professor at Montana Technological University in Butte, MT. David has 40 years' experience in occupational and environmental health with expertise in ergonomics, safety, and workplace wellness. Dr. Gilkey earned his Doctor of Chiropractic degree from Southern California Health Sciences University and Ph.D. from Colorado State University with a focus in occupational health, safety, industrial hygiene and ergonomics. He is a Certified Professional Ergonomist (CPE), Certified Safety Professional (CSP) and Registered Environmental Health Specialist (REHS). Dr. Gilkey has authored and/or co-authored 40 articles in peer reviewed scientific journals, 60 articles in trade journals and has provided four book chapter contributions in the areas of ergonomics, occupational safety, and environmental health. He has taught both undergraduate and graduate level courses in environmental and public health, safety and ergonomics. His research has focused in translational (R2P) research looking at methods to enhance safe work practices in agriculture where ATVs are used in farm and ranch operations. He was actively involved in construction safety climate research with an emphasis on evaluating differences between Latino and Non-Latinos workers. His most recent work is focused on safety climate in mining.