

2021

Computerized tumor multinucleation index (MuNI) is prognostic in p16+ oropharyngeal carcinoma

Can F Koyuncu
Case Western Reserve University

Wade L Thorstad
Washington University School of Medicine in St. Louis

R D Chernock
Washington University School of Medicine in St. Louis

Jingqin Luo
Washington University School of Medicine in St. Louis

et al

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Recommended Citation

Koyuncu, Can F; Thorstad, Wade L; Chernock, R D; Luo, Jingqin; and et al, "Computerized tumor multinucleation index (MuNI) is prognostic in p16+ oropharyngeal carcinoma." *The Journal of Clinical Investigation*, . (2021).
https://digitalcommons.wustl.edu/open_access_pubs/10387

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact scales@wustl.edu.

Computerized tumor multinucleation index (MuNI) is prognostic in p16⁺ oropharyngeal carcinoma

Can F. Koyuncu, ... , James S. Lewis Jr., Anant Madabhushi

J Clin Invest. 2021;131(8):e145488. <https://doi.org/10.1172/JCI145488>.

Clinical Medicine

Oncology

Patients with p16⁺ oropharyngeal squamous cell carcinoma (OPSCC) are potentially cured with definitive treatment. However, there are currently no reliable biomarkers of treatment failure for p16⁺ OPSCC. Pathologist-based visual assessment of tumor cell multinucleation (MN) has been shown to be independently prognostic of disease-free survival (DFS) in p16⁺ OPSCC. However, its quantification is time intensive, subjective, and at risk of interobserver variability.

We present a deep-learning–based metric, the multinucleation index (MuNI), for prognostication in p16⁺ OPSCC. This approach quantifies tumor MN from digitally scanned H&E-stained slides. Representative H&E-stained whole-slide images from 1094 patients with previously untreated p16⁺ OPSCC were acquired from 6 institutions for optimization and validation of the MuNI.

The MuNI was prognostic for DFS, overall survival (OS), or distant metastasis–free survival (DMFS) in p16⁺ OPSCC, with HRs of 1.78 (95% CI: 1.37–2.30), 1.94 (1.44–2.60), and 1.88 (1.43–2.47), respectively, independent of age, smoking status, treatment type, or tumor and lymph node (T/N) categories in multivariable analyses. The MuNI was also prognostic for DFS, OS, and DMFS in patients with stage I and stage III OPSCC, separately.

MuNI holds promise as a low-cost, tissue-nondestructive, H&E stain–based digital biomarker test [...]

Computerized tumor multinucleation index (MuNI) is prognostic in p16⁺ oropharyngeal carcinoma

Can F. Koyuncu,^{1,2} Cheng Lu,¹ Kaustav Bera,¹ Zelin Zhang,³ Jun Xu,³ Paula Toro,¹ German Corredor,^{1,2} Deborah Chute,⁴ Pingfu Fu,⁵ Wade L. Thorstad,⁶ Farhoud Faraji,⁷ Justin A. Bishop,⁸ Mitra Mehrad,⁹ Patricia D. Castro,¹⁰ Andrew G. Sikora,^{10,11} Lester D.R. Thompson,¹² R.D. Chernock,⁶ Krystle A. Lang Kuhs,⁹ Jingqin Luo,⁶ Vlad Sandulache,^{10,11} David J. Adelstein,⁴ Shlomo Koyfman,⁴ James S. Lewis Jr.,⁹ and Anant Madabhushi^{1,2}

¹Department of Biomedical Engineering, Case Western Reserve University, Cleveland, Ohio, USA. ²Louis Stokes Cleveland Veterans Affairs (VA) Medical Center, Cleveland, Ohio, USA. ³Nanjing University of Information Science and Technology, Nanjing, China. ⁴Cleveland Clinic Foundation, Cleveland, Ohio, USA. ⁵Department of Population and Quantitative Health Sciences, Case Western Reserve University, Cleveland, Ohio, USA. ⁶Washington University in St. Louis, St. Louis, Missouri, USA. ⁷UCSD, San Diego, California, USA. ⁸University of Texas (UT) Southwestern Medical Center, Dallas, Texas, USA. ⁹Vanderbilt University Medical Center, Nashville, Tennessee, USA. ¹⁰Department of Otolaryngology, Head and Neck Surgery, Baylor College of Medicine, Houston, Texas, USA. ¹¹ENT Section, Operative Care Line, Michael E. DeBakey VA Medical Center, Houston, Texas, USA. ¹²Southern California Permanente Medical Group, Pasadena, California, USA.

BACKGROUND. Patients with p16⁺ oropharyngeal squamous cell carcinoma (OPSCC) are potentially cured with definitive treatment. However, there are currently no reliable biomarkers of treatment failure for p16⁺ OPSCC. Pathologist-based visual assessment of tumor cell multinucleation (MN) has been shown to be independently prognostic of disease-free survival (DFS) in p16⁺ OPSCC. However, its quantification is time intensive, subjective, and at risk of interobserver variability.

METHODS. We present a deep-learning-based metric, the multinucleation index (MuNI), for prognostication in p16⁺ OPSCC. This approach quantifies tumor MN from digitally scanned H&E-stained slides. Representative H&E-stained whole-slide images from 1094 patients with previously untreated p16⁺ OPSCC were acquired from 6 institutions for optimization and validation of the MuNI.

RESULTS. The MuNI was prognostic for DFS, overall survival (OS), or distant metastasis-free survival (DMFS) in p16⁺ OPSCC, with HRs of 1.78 (95% CI: 1.37–2.30), 1.94 (1.44–2.60), and 1.88 (1.43–2.47), respectively, independent of age, smoking status, treatment type, or tumor and lymph node (T/N) categories in multivariable analyses. The MuNI was also prognostic for DFS, OS, and DMFS in patients with stage I and stage III OPSCC, separately.

CONCLUSION. MuNI holds promise as a low-cost, tissue-nondestructive, H&E stain-based digital biomarker test for counseling, treatment, and surveillance of patients with p16⁺ OPSCC. These data support further confirmation of the MuNI in prospective trials.

FUNDING. National Cancer Institute (NCI), NIH; National Institute for Biomedical Imaging and Bioengineering, NIH; National Center for Research Resources, NIH; VA Merit Review Award from the US Department of VA Biomedical Laboratory Research and Development Service; US Department of Defense (DOD) Breast Cancer Research Program Breakthrough Level 1 Award; DOD Prostate Cancer Idea Development Award; DOD Lung Cancer Investigator-Initiated Translational Research Award; DOD Peer-Reviewed Cancer Research Program; Ohio Third Frontier Technology Validation Fund; Wallace H. Coulter Foundation Program in the Department of Biomedical Engineering; Clinical and Translational Science Award (CTSA) program, Case Western Reserve University; NCI Cancer Center Support Grant, NIH; Career Development Award from the US Department of VA Clinical Sciences Research and Development Program; Dan L. Duncan Comprehensive Cancer Center Support Grant, NIH; and Computational Genomic Epidemiology of Cancer Program, Case Comprehensive Cancer Center. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH, the US Department of VA, the DOD, or the US Government.




Figure 1. Example of tumor cell MN. (A) Typical nonkeratinizing SSC, for comparison, which has fusiform, high nuclear/cytoplasmic ratios with inconspicuous nucleoli and brisk mitotic activity. The nuclei are relatively consistent in size across the whole tumor (original magnification, $\times 20$). (B) Tumor with an area of MN, in which cells have 3 or more nuclei (original magnification, $\times 40$).

Introduction

The continued increase in the incidence of oropharyngeal squamous cell carcinoma (OPSCC) in the setting of declining rates of tobacco use has been attributed to HPV, with almost 70% of all OPSCCs being HPV⁺ (1–3). HPV-related OPSCC has now overtaken cervical cancer as the most common HPV-related malignancy in the US (1, 2). HPV positivity has been well demonstrated to confer favorable survival for patients with OPSCC (4, 5). This has led to separate tumor, node, metastasis (TNM) staging systems for HPV-related (p16⁺) and HPV-unrelated OPSCC patients in the new 8th edition of the American Joint Commission on Cancer/Union for International Cancer Control (AJCC/UICC) guidelines (3), with p16 IHC currently recognized as a suitable surrogate marker for HPV in these patients (4–6).

Clinically, tumor and lymph node (T/N) status along with smoking have previously been shown to influence the risk of recurrence and/or death for patients with p16⁺ OPSCC (4, 5). However, these clinical parameters fail to capture intrinsic biological characteristics of p16⁺ OPSCC that may be relevant to treatment sensitivity and thus a clinical response. Machczynski et al. nicely discuss the limitations of even the new 8th edition of the AJCC guidelines regarding staging for p16⁺ OPSCC (7). Wuerdemann et al. found that the 8th edition's staging system did not discriminate well between patients with HPV⁺ stage II OPSCC and those with HPV⁺ stage III OPSCC (8). Similarly, no significant differences in OS have been found between the 8th edition's guidelines for (9) stage I versus stage II and (10) stage II versus stage III OPSCC.

Currently, the biomarker landscape for HPV-related OPSCC

validation cohorts and very small sample sizes. In addition, these strategies rely on tissue-destructive sampling or require expensive methods for gene expression profiling. To date, however, no single signature has been validated as a practical predictor of DFS, OS, or distant metastasis-free survival (DMFS) in multiple cohorts, a critical step toward translating the signature into a clinical test. Reliable and sensitive prognostic tools for p16⁺ OPSCC could support clinical decision making. For instance, in high-risk patients, a safe de-escalation of treatment may not be possible, whereas, conversely, very low-risk patients could benefit from de-escalation and thus avoid

adverse effects related to unnecessary intensive therapy (5).

Traditionally, tissue morphology and architecture within the tumor microenvironment (TME) have been shown to be reflective of tumor characteristics and to carry rich prognostic and predictive information across myriad histologic types (15–17). For head and neck squamous cell carcinoma (HNSCC), Hartman et al. reported that high numbers of tumor-infiltrating CD8⁺ T cells in the TME are associated with oropharyngeal localization and limited tumor growth and that the patients with high infiltration of CD8⁺ T cells also have significantly better outcomes (18). Lewis et al. previously visually identified anaplasia and multinucleation (MN) in the TME as novel prognostic and independently associated features in patients with p16⁺ OPSCC (Figure 1 and ref. 19). However, the recognition and quantification of morphologic features is time intensive and requires human interpretation, leading to interobserver variability and bias.

In this work, we present a computerized metric called the multinucleation index (MuNI), a quantification of MN density in epithelial (EP) regions, to risk-stratify patients with p16⁺ OPSCC for DFS, OS, and DMFS. Two machine-learning networks, specifically, conditional generative adversarial networks (cGANs) (20), were used for the MuNI calculations: (a) GAN_{MN} to segment MN events and (b) GAN_{EP} to segment cancer nuclei in EP regions in digitized H&E-stained whole-slide images (WSIs). By calculating the ratio of the MN events to the total number of EP nuclei identified on the slide image, we calculated a MuNI for every slide and every patient with p16⁺ OPSCC (Figure 2). A large cohort of 1094 previously untreated patients with p16⁺ OPSCC obtained from




Figure 2. Overall flowchart of the image analysis pipeline. The training step involved building MN and EP segregators. The same deep-learning architecture, cGAN, was used to build the segregators. Each block in the cGAN models consisted of convolution, BatchNorm, and ReLU layers. The MuNI calculation phase started with the extraction of tiles from tissue regions. The tiles were then input into the MN and EP models separately. Finally, the MuNI was calculated automatically, which was the ratio of MNs within EP regions to EP cells.

<https://doi.org/10.1172/JCI145488DS1>). Patients were followed for a mean and median of 63 and 59 months, respectively (range, 1–200 months). Briefly, the median age for the entire set was 58 years; 66% of patients were current or former smokers; and 93.4% of patients were White. The Non-White races consisted of Black (5.3%) and Asian (0.4%). Supplemental Table 1 provides statistical differences in the clinical variables and MuNI across the cohorts.

Experiment 1: association between the MuNI and OS, DFS, and DMFS in all patients. Table 1 provides the results of the univariate analysis for the major clinical and pathologic features and for the computerized detection of MN in different cohorts. Kaplan-Meier (KM) survival curves for MuNIs in the training and validation cohorts are presented in Figures 3, 4, and 5. Cohort-specific KM curves are provided in the supplemental material (Supplemen-

0.001), and DMFS (HR: 1.88, 95% CI: 1.43–2.47, $P < 0.001$) (Table 2). Visual examples of 3 high-risk and low-risk samples identified by MuNI are shown in Supplemental Figure 10.

Experiment 2: association between the MuNI and OS, DFS, and DMFS in the AJCC 8th edition's defined stage groups. The prognostic ability of the MuNI was evaluated for patients within the AJCC 8th edition's defined stage groups. On univariate and multivariable analyses, we found that the MuNI was prognostic for DFS, OS, and DMFS for the S_{VA} patients with stage I and stage III tumors, respectively. The HRs for predicting DFS, OS, and DMFS on univariate analysis were 1.93 (95% CI: 1.32–2.82, $P < 0.001$), 1.76 (95% CI: 1.13–2.74, $P < 0.01$), and 1.88 (CI: 1.26–2.83, $P < 0.02$), respectively, for stage I tumors, and 2.29 (CI: 1.47–3.56, $P < 0.001$), 2.31 (CI: 1.42–3.77, $P < 0.02$), and 2.11 (CI: 1.33–3.35, $P < 0.01$), respectively,

Table 1. HRs and P values from univariate Cox proportional hazards model analysis of DFS across 6 institutions

Variable	Univariate Cox proportional hazards model analysis											
	Validation cohort (S_{VA})		D_2		D_3		D_4		D_5		D_6	
	HR (95% CI)	P	HR (95% CI)	P	HR (95% CI)	P	HR (95% CI)	P	HR (95% CI)	P	HR (95% CI)	P
Age (≤ 56 vs. > 56 yr)	1.41 (1.11–1.78)	0.01	1.87 (0.77–4.54)	0.17	0.90 (0.46–1.75)	0.76	0.90 (0.54–1.49)	0.67	1.21 (0.80–1.83)	0.37	2.04 (1.25–3.33)	0.01
Sex	1.48 (0.99–2.21)	0.06	1.77 (0.55–5.67)	0.44	1.00 (0.30–3.26)	0.99	0.36 (0.03–4.61)	0.43	1.15 (0.6–2.21)	0.69	1.94 (0.81–4.62)	0.26
Race (White vs. Black or Asian)	0.83 (0.45–1.52)	0.58	4.3 (0.26–71.7)	0.03	2.16 (0.41–11.4)	0.19	0.48 (0.18–2.20)	0.42	0.57 (0.23–1.38)	0.33	0.52 (0.12–2.18)	0.50
Smoking status (0 vs. 1)	1.31 (1.03–1.67)	0.04	1.99 (0.85–4.68)	0.16	1.36 (0.68–2.7)	0.39	1.08 (0.57–2.02)	0.81	1.44 (0.93–2.23)	0.13	0.89 (0.54–1.46)	0.64
Overall stage (stage I/II vs. stage III)	1.79 (1.32–2.43)	< 0.001	1.88 (0.66–5.33)	0.16	1.88 (0.82–4.32)	0.08	1.35 (0.79–2.32)	0.22	1.56 (0.98–2.49)	0.04	2.44 (0.77–7.66)	0.02
T stage (T1/2 vs. T3/4)	1.87 (1.45–2.43)	< 0.001	1.57 (0.66–3.76)	0.28	2.15 (1.05–4.41)	0.02	1.04 (0.64–1.67)	0.88	1.92 (1.25–2.95)	0.01	2.37 (1.13–5)	0.01
N stage (N0/1 vs. N2/3)	1.72 (1.34–2.2)	< 0.001	1.3 (0.53–3.21)	0.54	2.73 (1.25–5.96)	0.01	0.97 (0.52–1.63)	0.76	1.15 (0.75–1.77)	0.50	1.61 (0.87–2.98)	0.08
MuNI (high vs. low)	1.79 (1.42–2.26)	< 0.001	2.64 (1.15–6.04)	0.05	2.15 (1.07–4.3)	0.06	1.45 (0.89–2.34)	0.14	1.53 (1.00–2.35)	0.04	1.96 (1.2–3.21)	0.01

HRs and P values from univariate Cox proportional hazards model analysis of DFS across 6 institutions. Bolded values indicate significant HR or P values. Data for the other races are in Supplemental Table 1.

so was trained on S_{TR} using age, sex, smoking status, TNM stage, and the MuNI to predict OS. We calculated the risk score for each patient for risk stratification. The median of the risk scores in the S_{TR} group was defined as the cutoff to dichotomize low- and high-risk patients, and the same cutoff was used in the S_{VA} group. The HR value for predicting OS on univariate analysis was 2.42 (95% CI: 1.86–3.15, $P < 0.001$) for S_{VA} (Figure 5). We conducted the same experiments using the clinical variables only, without the MuNI, and the corresponding HR was 1.43 (CI: 1.09–1.87, $P < 0.01$).

Experiment 4: evaluating the resilience of the MuNI against batch effects to account for site-specific preanalytic variations. We performed qualitative analysis of the MuNI resilience against batch effects. As a baseline, for each WSI coming from any of the 6 sites, image metrics related to image brightness and contrast were calculated using HistoQC. The features were then embedded into 2D space for visualization using the t -distributed stochastic neighbor embedding (t -SNE) algorithm (Figure 6A). Likewise, we performed t -SNE mapping to calculate MuNI-specific statistics (Figure 6, B and C). We assessed a total of 8 different statistics (MuNI of the entire WSI, mean, median, SD, minimum, maximum, and 33rd–66th percentiles) for the MuNI from across every tile of the WSI.

stantial toxicity and morbidity from these therapies. Consequently, there is a clear need to develop a quantifiable and reproducible biomarker to stratify high- and low-risk patients with p16⁺ OPSCC (23). Low-risk patients might then potentially benefit from therapy de-intensification, whereas high-risk patients would continue standard or intensified management.

Lewis et al. previously identified anaplasia and MN as novel prognostic features in patients with p16⁺ OPSCC (19). These were strongly and independently associated with disease recurrence and death from the disease and also correlated with DFS in a cohort of surgically treated patients with OPSCC ($n = 149$). However, identification of the above-mentioned morphologic features is pathologist dependent, and, although no specific study in the literature documents it, implies subjectivity and potential bias (24). We found specific examples difficult to discern and quantify, such as overlapping nuclei from separate cells that were not truly multinucleated and large, anaplastic, irregular nuclei that were also not truly multinucleated. Additionally, the study was performed at a single institution in a cohort of only surgically treated patients, for whom all slides of resected tumor, including lymph node metastases, were reviewed. Interestingly, in this study, the pathologist's quantification of MNs on the single H&E-stained slides for 478 patients with p16⁺ OPSCC





Figure 3. KM DMFS curves. KM DMFS curves for all patients in the training/validation cohorts and groups of patients at different cancer stages in the validation cohort according to the AJCC 8th edition's definition.

DFS, as well as DMFS among the AJCC 8th edition's defined stage I and stage III patients in both univariate and multivariable analyses. If confirmed in a prospective clinical trial setting, we believe this finding could have major implications for clinical practice. Patients with stage I disease, currently the target of de-escalation treatment strategies, could be further stratified using the MuNI

to exclude those who might have a high chance of treatment failure resulting in recurrence (25). Although multiple clinical trials are currently exploring therapeutic de-intensification strategies, they are limited by a dependence on clinical parameters to identify appropriate patients at low risk of disease recurrence (12). The identification of biologically meaningful markers of a good prog-






Figure 5. KM OS curves. KM OS curves for all patients in the training/validation cohorts and groups of patients at different cancer stages in the validation cohort according to the AJCC 8th edition's definition. The last row presents KM OS curves for all patients in the validation cohort using an integrated classifier comprising clinical variables for predicting OS.

nosis is of critical importance. Similarly, patients with stage III disease who are further categorized as high risk by the MuNI may merit the maintenance of treatment intensity by incorporating surgical resection, consistently utilizing concurrent chemotherapy, or intensifying chemoradiotherapy. Taken together, this would represent a novel, viable precision oncology approach to treating patients with p16⁺ OPSCC in the modern era.

In spite of the differences in clinical and pathological data between the sites (Supplemental Table 1), the MuNI was prognostic across the different sites using a single threshold cutoff learned from a single site, although with modest HRs for death. In Figure 6, the

Quantitative histomorphometric (QH) approaches for the prognostication of disease outcomes have been previously proposed for many cancers. These approaches fall into 2 major categories: hand-crafted (or domain-inspired) and deep-learning- or neural network-based approaches. We have previously introduced 2 hand-crafted-based approaches, OHbIC (26) and QuHbIC (27), to stratify the risk of patients with head and neck carcinomas using H&E-stained tumor microarrays (TMAs). The first study showed the independent prognostic value of OHbIC, which utilizes nuclear shape and texture features for predicting disease-specific survival, in a cohort ($n = 115$) of patients with oral cavity squamous cell

Table 2. HRs and P values from multivariable Cox proportional hazards model analysis of DFS across 6 institutions

Variable	Multivariable Cox proportional hazards model analysis controlling for other variables											
	Validation cohort, S _{va}		D ₂		D ₃		D ₄		D ₅		D ₆	
	HR (95% CI)	P	HR (95% CI)	P	HR (95% CI)	P	HR (95% CI)	P	HR (95% CI)	P	HR (95% CI)	P
Age	1.04 (1.03–1.06)	< 0.001	1.10 (1.04–1.16)	< 0.001	1.01 (0.97–1.06)	0.68	1.03 (1.00–1.06)	0.07	1.03 (1.01–1.06)	0.02	1.07 (1.03–1.10)	< 0.001
Sex	1.45 (0.88–2.40)	0.14	1.92 (0.40–9.31)	0.42	0.54 (0.12–2.32)	0.40	(0.00 – inf.)	1.00	1.27 (0.61–2.64)	0.52	1.32 (0.39–4.46)	0.66
Race (White vs. Black or Asian)	0.92 (0.58–1.45)	0.71	0.37 (0.07–1.82)	0.22	0.98 (0.24–4.08)	0.98	0.67 (0.34–1.30)	0.24	2.70 (0.84–8.63)	0.10	0.62 (0.08–4.83)	0.30
Smoking status (0 vs. 1)	1.32 (0.99–1.78)	0.06	1.52 (0.54–4.28)	0.43	1.41 (0.66–3.04)	0.36	0.89 (0.47–1.69)	0.72	1.18 (0.72–1.95)	0.51	1.10 (0.60–2.00)	0.78
Overall stage	0.92 (0.67–1.25)	0.58	0.75 (0.24–2.42)	0.64	0.80 (0.33–1.90)	0.61	1.32 (0.65–2.39)	0.52	0.78 (0.44–1.37)	0.38	0.80 (0.40–1.65)	0.56
T stage	1.33 (1.08–1.64)	0.01	1.51 (0.68–3.38)	0.31	1.41 (0.76–2.64)	0.28	0.88 (0.62–1.32)	0.59	1.59 (1.07–2.36)	0.02	1.77 (1.07–2.94)	0.03
N stage	1.28 (1.03–1.58)	0.02	0.94 (0.44–1.97)	0.86	1.70 (0.93–3.09)	0.08	1.00 (0.68–1.46)	0.98	1.14 (0.78–1.65)	0.50	1.52 (0.92–2.53)	0.10
MuNI (low vs. high)	1.78 (1.37–2.30)	< 0.001	2.22 (0.76–6.51)	0.15	2.15 (0.92–5.03)	0.08	1.46 (0.86–2.49)	0.16	1.45 (0.95–2.21)	0.08	1.90 (0.95–3.79)	0.07

HRs and P values from multivariable Cox proportional hazards model analysis of DFS across 6 institutions. Bolded values indicate significant HR or P values. Data for the other races are in Supplemental Table 1. inf., infinitive.

to learn best representations for predicting prognosis categories of interest without requiring a pathologist's input. However, because of the multilayered, nonlinear structure of deep-learning models, they are considered black boxes, and their output is not interpretable by pathologists or translatable into any directly visual form. Interestingly, unlike these models, the method presented here utilizes the power of deep learning with the interpretability of hand-crafted (i.e., visually identified) features. In other words, deep learning was used not to make a direct prognostic prediction but rather to quantify MNs in WSIs and derive a prognostic metric based on the number of MNs identified on the WSIs. As demonstrated in experiment 4, the hybrid approach to identify a computational pathology-based biomarker was found to be resilient against batch effects.

Our study has some limitations. The cohorts from different institutions were found to have significant differences in their MuNIs, as well as in certain clinical and pathologic parameters (Supplemental Table 1 and Supplemental Figure 9). Nonetheless, we found that the MuNI was prognostic for the entire set of validation cohorts in both univariate and multivariable analyses, although it was not consistently prognostic for each of the separate

since there were 471 patients, whereas in the group with stage II disease, which included 245 patients, that difference was not apparent. In the group of patients with stage III disease, whose overall prognosis was much worse, we could detect the difference, even though there were only 169 patients. Finally, this study was based on retrospectively collected data. Analyses of slides from completed multi-institutional, prospective clinical trials, or better yet, a prospective clinical trial with the MuNI embedded within it, are required to validate the findings, minimize the potential for bias, and determine whether the MuNI can specifically predict a patient's response to treatment.

In conclusion, the MuNI is a tissue-nondestructive, reproducible, rapid, and cost-efficient artificial intelligence-enabled (AI-enabled) biomarker with the potential to risk-stratify patients with p16⁺ OPSCC. The MuNI only relies on the quantitative measurement of MN tumor cells in digitized H&E-stained tissue from primary tumors, without the need for visual or manual segmentation of tumor versus nontumor regions. These specimens are already consistently obtained from patients with OPSCC in routine practice. This makes the MuNI potentially widely introducible into clinical practice at US institutions and useful in low- and




Figure 6. Patients in the entire set were embedded into 2D feature space and then plotted using *t*-SNE. The x and y axes refer to *t*-SNE dimensions 1 and 2, respectively. Embedment was performed using (A) low-level image features such as color and texture extracted via HistoQC and (B) the MuNI and (C) MuNI-derived metrics. Each dot represents a patient, and the dots are colored according to (A and B) the site they originated from or (C) their progressor or nonprogressor label. (A) Each site clusters separately because of the differences in patient demographics between the sites. Despite the differences, (B) no site was clustered away from the others, indicating that the MuNI-derived metrics were reproducible across sites. (C) Separation between progressor and nonprogressor patients using the MuNI metrics.

diagnosis of OPSCC. Immunohistochemistry for p16 had been performed at the respective institutions in routine clinical practice, and only those tumor specimens that were classified as p16⁺ by nationally accepted pathologic standards (extensive nuclear and cytoplasmic staining present in 70% or more of the tumor specimen with at least moderate to strong intensity) were included (32). H&E-stained glass slides from the primary tumors of each patient were re-reviewed by the collaborating pathologists for the selection of the most representative tumor

sent to Case Western Reserve University, where they underwent WSI scanning at $\times 40$ resolution (0.25 $\mu\text{m}/\text{pixel}$ resolution) using a Ventana iScan HT slide scanner. WSI quality checking was performed using HistoQC (33), an automatic, rapid, and quantifiable quality control tool for computational pathology that excluded an additional 61 patients' specimens because of poor image quality. Finally, 1094 specimens remained for the analysis. The patients were divided into training and validation sets, S_{TR} and S_{VA} , respectively. D_1 comprised




Figure 7. Inclusion and exclusion criteria for the study. Data on a total of 1485 patients with OPSCC were gathered from the 6 institutions. After reviewing the clinical data and WSI image quality, 391 of the patients were excluded. A total of 1094 specimens remained for the analysis. SCPMG, Southern California Permanente Medical Group; WUSTL, Washington University in St. Louis; JHU, Johns Hopkins University; MEDVAMC, Michael E. DeBakey VA Medical Center; CCF, Cleveland Clinic Foundation; VUMC, Vanderbilt University Medical Center.

of 6 cases from S_{TR}. A total of 153 image patches, each corresponding to 512 × 512 pixels, were cropped at ×10 magnification and then annotated by a pathologist. Of these, 102 were used for training GAN_{EP}. Its performance was then evaluated quantitatively on the remaining 51 images and yielded a pixel-level F1 score of 0.88.

Automated detection and segmentation of MN

MNs were segmented using another cGAN model (GAN_{MN}) that involved 2 steps. Training of the GAN_{MN} for nuclei segmentation was done using 30 images from a public data set corresponding to multiple organs (34). Patches of 256 × 256 pixels were extracted from these images at ×40 magnification and fed into the model during training. Segmentation performance was quantitatively and qualitatively verified to be suitable for MN segmentation. For independent validation of GAN_{MN}, another publicly available data

to identify and eliminate false-positive MN exemplars, given that MNs are typically larger compared with EP cells and lymphocytes. A detailed description of the network architecture and validation of GAN_{MN} is provided in the supplemental material (Supplemental Method 1).

MuNI

Utilizing GAN_{EP} and GAN_{MN}, EP and MN masks were extracted for each WSI. For a WSI, m was used to denote the number of tiles extracted from the WSI and M_{MN}^i and M_{EP}^i corresponded to the number of detected MNs and EP cells in tile i extracted from the WSI, respectively. The normalized MuNI for the WSI was then defined as the ratio of total MNs to EP cells.

$$\text{MuNI} = \frac{\sum_{i=1}^m M_{MN}^i}{\sum_{i=1}^m M_{EP}^i}$$

Statistics

The similarity of clinical and pathologic variables between different institutional cohorts was calculated using ANOVA. Associations between the dichotomized MuNIs and the other categorical clinical and pathologic variables were determined by a 2-sided Fisher's exact test. KM survival curves and univariate log-rank tests were applied to correlate DFS, OS, and DMFS with the MuNI. Multivariable Cox proportional hazards models were also used to investigate the independent prognostic ability of MuNI for DFS, OS, and DMFS after accounting for age, sex, race, smoking status, treatment type, and T/N categories. The same univariate and multivariable analyses were further performed for each cancer stage group defined by the AJCC's 8th edition. The prognostic power of the MuNI, together with other clinical variables, was analyzed using Cox regression via Lasso. HRs associated with 95% CIs and *P* values from Wald tests were reported. All tests were 2 tailed, with the significance level set at 0.05.

Study approval

The present studies in humans were reviewed and approved by the IRBs of Washington University in St. Louis, Missouri, USA, Johns Hopkins University in Baltimore, Maryland, USA, Vanderbilt University Medical Center in Nashville, Tennessee, USA, the Cleveland Clinic Foundation in Cleveland, Ohio, USA, Case Western Reserve University in Cleveland, Ohio, USA, the Southern California Permanente Medical Group in Los Angeles, California, USA, and the Michael E. DeBakey VA Medical Center in Houston, Texas, USA.

Author contributions

CFK and CL contributed to the methodology, experiments, and writing of the manuscript. ZZ and JX contributed to the methodology and reviewed the manuscript. KB, PAT, GC, and PF helped review the manuscript. DC, WLT, FF, JAB, MM, PDC, AGS, LDRT, RDC, KALK, JL, VS, DJA, and SK contributed to data collection and reviewed the manuscript. JSL and AM contributed to the methodology, experiments, and writing of the manuscript.

Acknowledgments

This work was supported by the NCI, NIH (1U24CA199374-01, R01CA202752-01A, R01CA208236-01A1, R01CA216579-01A1, R01CA220581-01A1, 1U01CA239055-01); the National Institute for Biomedical Imaging and Bioengineering, NIH (1R43EB028736-01); the National Center for Research Resources, NIH (1CO6RR12463-01); a VA Merit Review Award (IBX004121A) from the US Department of VA Biomedical Laboratory Research and Development Service; a US DOD Breast Cancer Research Program Breakthrough Level 1 Award (W81XWH-19-1-0668); a US DOD Prostate Cancer Idea Development Award (W81XWH-15-1-0558); a US DOD Lung Cancer Investigator-Initiated Translational Research Award (W81XWH-18-1-0440); the US DOD Peer Reviewed Cancer Research Program (W81XWH-16-1-0329); the Ohio Third Frontier Technology Validation Fund; the Wallace H. Coulter Foundation Program in the Department of Biomedical Engineering; a CTSA award to Case Western Reserve University; NCI Cancer Center Support Grant P30CA125123, NIH; Career Development Award 1 IK2 CX001953 from the US Department of VA Clinical Sciences Research and Development Program; Dan L. Duncan Comprehensive Cancer Center Support Grant (NCI-CA125123), NIH; and the Computational Genomic Epidemiology of Cancer Program at Case Comprehensive Cancer Center (T32CA094186). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH, the US Department of VA, the DOD, or the US Government.

Address correspondence to: James S. Lewis Jr., Vanderbilt University Medical Center, 1211 Medical Center Dr., Nashville, Tennessee, 37232 USA. Phone: 615.343.0233; Email: james.lewis@vumc.org. Or to: Anant Madabhushi, Louis Stokes Cleveland VA Medical Center, 10701 East Blvd., Cleveland, Ohio, 44106 USA. Phone: 216.368.8519; Email: anant.madabhushi@case.edu.

- Gillison ML, et al. Epidemiology of human papillomavirus-positive head and neck squamous cell carcinoma. *J Clin Oncol.* 2015;33(29):3235–3242.
- Chaturvedi AK, et al. Human papillomavirus and rising oropharyngeal cancer incidence in the United States. *J Clin Oncol.* 2011;29(32):4294–4301.
- O'Sullivan B. HPV-mediated (p16+) oropharyngeal cancer. In: Greene FL, et al., eds. *AJCC Cancer Staging Manual.* Springer; 2017:113–121.
- Ang KK, et al. Human papillomavirus and survival of patients with oropharyngeal cancer. *N Engl J Med.* 2010;363(1):24–35.
- nalaryngol. 2020;277(9):2407–2412.
- Wuerdemann N, et al. Risk factors for overall survival outcome in surgically treated human papillomavirus-negative and positive patients with oropharyngeal cancer. *Oncol Res Treat.* 2017;40(6):320–327.
- Gupta P, et al. Validation and assessment of discordance of the 8th edition AJCC (American Joint Committee on Cancer) clinical and pathologic staging systems in patients with p16+ oropharyngeal cancer treated with surgery and adjuvant radiation at a single institution. *Oral Oncol.* 2020;27(1):1–10.
- of head and neck cancer. *Oncoimmunology.* 2014;3(1):e27403.
- Mizumachi T, et al. Confirmation of the eighth edition of the AJCC/UICC TNM staging system for HPV-mediated oropharyngeal cancer in Japan. *Int J Clin Oncol.* 2017;22(4):682–689.
- Würdemann N, et al. Prognostic impact of AJCC/UICC 8th edition new staging rules in oropharyngeal squamous cell carcinoma. *Front Oncol.* 2017;7:129.
- Ali S, et al. Selective invocation of shape priors for deformable segmentation and morphologic classification of head and neck cancer. *Oncoimmunology.* 2014;3(1):e27403.

- 2018;86:278–287.
19. Lewis JS, et al. Tumor cell anaplasia and multinucleation are predictors of disease recurrence in oropharyngeal squamous cell carcinoma, including among just the human papillomavirus-related cancers. *Am J Surg Pathol*. 2012;36(7):1036–1046.
20. Mahmood F, et al. Deep adversarial training for multi-organ nuclei segmentation in histopathology images. *IEEE Trans Med Imaging*. 2020;39(11):3257–3267.
21. Adelstein DJ, et al. Role of treatment deintensification in the management of p16+ oropharyngeal cancer: ASCO provisional clinical opinion. *J Clin Oncol*. 2019;37(18):1578–1589.
22. Contreras JA, et al. Eliminating postoperative radiation to the pathologically node-negative neck: long-term results of a prospective phase II study. *J Clin Oncol*. 2019;37(28):2548–2555.
23. Eze N, et al. Biomarker driven treatment of head and neck squamous cell cancer. *Cancers Head Neck*. 2017;2:6.
24. Ranganathan K, et al. Intra-observer and inter-observer variability in two grading systems for oral epithelial dysplasia: a multi-centre study in India. *J Oral Pathol Med*. 2020;49(9):984–955.
25. Ma DJ, et al. Phase II evaluation of aggressive dose de-escalation for adjuvant chemoradiotherapy in human papillomavirus-associated oropharynx squamous cell carcinoma. *J Clin Oncol*. 2019;37(22):1909–1918.
26. Lu C, et al. An oral cavity squamous cell carcinoma quantitative histomorphometric-based image classifier of nuclear morphology can risk stratify patients for disease-specific survival. *Mod Pathol*. 2017;30(12):1655–1665.
27. Lewis JS, et al. A quantitative histomorphometric classifier (QuHbIC) identifies aggressive versus indolent p16-positive oropharyngeal squamous cell carcinoma. *Am J Surg Pathol*. 2014;38(1):128–137.
28. Bera K, et al. Artificial intelligence in digital pathology—new tools for diagnosis and precision oncology. *Nat Rev Clin Oncol*. 2019;16(11):703–715.
29. Bulten W, et al. Automated deep-learning system for Gleason grading of prostate cancer using biopsies: a diagnostic study. *Lancet Oncol*. 2020;21(2):233–241.
30. Ström P, et al. Artificial intelligence for diagnosis and grading of prostate cancer in biopsies: a population-based, diagnostic study. *Lancet Oncol*. 2020;21(2):222–232.
31. Skrede OJ, et al. Deep learning for prediction of colorectal cancer outcome: a discovery and validation study. *Lancet*. 2020;395(10221):350–360.
32. Begum S, et al. Detection of human papillomavirus in cervical lymph nodes: a highly effective strategy for localizing site of tumor origin. *Clin Cancer Res Off J Am Assoc Cancer Res*. 2003;9(17):6469–6475.
33. Janowczyk A, et al. HistoQC: an open-source quality control tool for digital pathology slides. *JCO Clin Cancer Inform*. 2019;3:1–7.
34. Kumar N, et al. A dataset and a technique for generalized nuclear segmentation for computational pathology. *IEEE Trans Med Imaging*. 2017;36(7):1550–1560.
35. Naylor P, et al. Segmentation of nuclei in histopathology images by deep regression of the distance map. *IEEE Trans Med Imaging*. 2019;38(2):448–459.