

Platelet FcγRIIA-induced serotonin release exacerbates the severity of Transfusion-Related Acute Lung Injury in mice

Tracking no: ADV-2021-004336R2

Marie-Belle El Mdawar (Etablissement Français du Sang-Grand Est, France) Blandine Maître (EFS-Grand Est, France) Stephanie Magnenat (EFS Grand Est, INSERM, France) Florian Tupin (Etablissement Français du Sang-Grand Est, France) Friederike Jonsson (Institut Pasteur/ INSERM, France) Christian Gachet (EFS, Inserm, France) Henri de la Salle (EFS-Alsace, France) Béatrice Hechler (Etablissement Français du Sang-Grand Est, France)

Abstract:

Transfusion-related acute lung injury (TRALI) remains a major cause of transfusion-related fatalities. The mechanism of human antibody-mediated TRALI, especially the involvement of the Fcγ receptors, is not clearly established. Contrary to mice, human platelets are unique in their expression of the FcγRIIA/CD32A receptor, suggesting that our understanding of the pathogenesis of antibody-mediated TRALI is partial, since the current murine models incompletely recapitulate the human immunology. We evaluated the role of FcγRIIA/CD32A in TRALI using a humanized mouse model expressing the FcγRIIA/CD32A receptor. When challenged with a recombinant chimeric human IgG1/mouse anti-MHC I monoclonal antibody, these mice showed exacerbated alveolar edema and higher mortality compared to WT mice. Unlike in WT mice, monocytes/macrophages were accessory for TRALI initiation in CD32A+ mice pointing to the decisive contribution of another cell type. Platelet activation was dramatically increased in CD32A+ animals resulting in their increased consumption and massive release of their granule contents. Platelet depletion prevented the exacerbation of TRALI in CD32A+ mice but did not affect TRALI in WT animals. By blocking platelet serotonin uptake with fluoxetine, we showed that the severity of TRALI in CD32A+ mice resulted from the serotonin released by activated platelets. Furthermore, inhibition of 5-hydroxytryptamine 2A serotonin receptor with sarpgrelate, before or after the induction of TRALI, abolished the aggravation of lung edema in CD32A+ mice. Our findings demonstrate that platelet FcγRIIA/CD32A activation exacerbates antibody-mediated TRALI and provide a rationale for designing prophylactic and therapeutic strategies targeting the serotonin pathway to attenuate TRALI in patients.

Conflict of interest: No COI declared

COI notes:

Preprint server: No;

Author contributions and disclosures: M.-B.E.M., B.M., S.M., F.T., F.J. and B.H. performed experiments and analyzed data; F.J. contributed essential material; H.d.l.S., B.M. and B.H. designed the research and wrote the manuscript; C.G. and F.J. critically reviewed the manuscript; all authors contributed to the final manuscript.

Non-author contributions and disclosures: No;

Agreement to Share Publication-Related Data and Data Sharing Statement: For original data, please contact beatrice.hechler@efs.sante.fr

Clinical trial registration information (if any):

Platelet FcγRIIA-induced serotonin release exacerbates the severity of Transfusion-Related Acute Lung Injury in mice

Short title: FcγRIIA/CD32A expression in mice exacerbates TRALI.

Marie-Belle El Mdawar^{1*}, Blandine Maître^{1*}, Stéphanie Magnenat¹, Florian Tupin¹, Friederike Jönsson², Christian Gachet¹, Henri de la Salle^{1#} and Beatrice Hechler^{1#}

*M.-B.E.M. and B.M co-first authors. #H.d.S. and B.H co-last authors.

¹ Université de Strasbourg, INSERM, Etablissement Français du Sang (EFS)-Grand Est, BPPS UMR_S 1255, Fédération de Médecine Translationnelle de Strasbourg (FMTS), F-67000 Strasbourg, France; ² Institut Pasteur, Department of Immunology, Unit of Antibodies in Therapy and Pathology, UMR INSERM U1222, Paris, France.

Corresponding author : Beatrice Hechler, UMR_S1255 INSERM, Université de Strasbourg, Etablissement Français du Sang-Grand Est, 10 rue Spielmann, BP 36, F-67065 Strasbourg Cedex, France. Tel. +33 3 88 21 25 25, Fax +33 3 88 21 25 21, email: beatrice.hechler@efs.sante.fr

Word counts:

- Text: 4127 words
- Abstract: 246 words
- Figure number: 6
- Reference count: 47

Abstract

Transfusion-related acute lung injury (TRALI) remains a major cause of transfusion-related fatalities. The mechanism of human antibody-mediated TRALI, especially the involvement of the Fc γ receptors, is not clearly established. Contrary to mice, human platelets are unique in their expression of the Fc γ RIIA/CD32A receptor, suggesting that our understanding of the pathogenesis of antibody-mediated TRALI is partial, since the current murine models incompletely recapitulate the human immunology. We evaluated the role of Fc γ RIIA/CD32A in TRALI using a humanized mouse model expressing the Fc γ RIIA/CD32A receptor. When challenged with a recombinant chimeric human IgG1/mouse anti-MHC I monoclonal antibody, these mice showed exacerbated alveolar edema and higher mortality compared to WT mice. Unlike in WT mice, monocytes/macrophages were accessory for TRALI initiation in CD32A⁺ mice pointing to the decisive contribution of another cell type. Platelet activation was dramatically increased in CD32A⁺ animals resulting in their increased consumption and massive release of their granule contents. Platelet depletion prevented the exacerbation of TRALI in CD32A⁺ mice but did not affect TRALI in WT animals. By blocking platelet serotonin uptake with fluoxetine, we showed that the severity of TRALI in CD32A⁺ mice resulted from the serotonin released by activated platelets. Furthermore, inhibition of 5-hydroxytryptamine 2A serotonin receptor with sarpogrelate, before or after the induction of TRALI, abolished the aggravation of lung edema in CD32A⁺ mice. Our findings demonstrate that platelet Fc γ RIIA/CD32A activation exacerbates antibody-mediated TRALI and provide a rationale for designing prophylactic and therapeutic strategies targeting the serotonin pathway to attenuate TRALI in patients.

Key points

- Platelet Fc γ RIIA/CD32A activation exacerbates the severity of antibody-mediated TRALI.
- Platelet serotonin release and activation of 5-HT_{2A} serotonin receptors is responsible for the Fc γ RIIA/CD32A-dependent exacerbation of TRALI.

Introduction

Transfusion-related acute lung injury (TRALI) is a syndrome characterized by the development of non-cardiogenic pulmonary edema. TRALI contributes significantly to the morbidity of transfusion, or can even be fatal ¹⁻³. TRALI can result from the presence in the transfused product of alloantibodies specific for polymorphic HLA/MHC class I or class II molecules or for neutrophil antigens, the latter two constituting the major causes of mortality in this syndrome. The *in situ* detection of platelets and neutrophils in the lungs of patients who died from TRALI has led to the proposition that these cells participate in its pathogenesis ⁴.

In order to analyze the mechanisms of alloantibody-mediated TRALI, several models have been developed in H-2^d BALB/c mice, all of which use the anti-H-2D^d/H-2K^d major histocompatibility complex class I (MHC I) mouse monoclonal antibody (mAb) 34-1-2S. In a two-hit model, an inflammatory priming step, induced by intraperitoneal administration of a low dose of lipopolysaccharides (LPS, 0.1 mg/kg), is followed 24 h later by intravenous injection of 34-1-2S (0.5 mg/kg) ⁵. In one-hit models, the animals are not sensitized with LPS, but at least ten-fold higher amounts of the antibody need to be injected to induce TRALI ^{6,7}. These models have allowed one to assess the respective contributions of the different blood cell populations and have resulted in various conclusions. Monocytes/macrophages were found to be essential, whereas CD4⁺ T cells and dendritic cells behaved as negative regulators of TRALI; depending on the studies, neutrophils and platelets were described to be either required or dispensable for lung edema formation ⁵⁻¹⁵. Recently, endothelial MHC I has been identified as the critical molecular target for initiation of anti-MHC I mAb-mediated lung injury ¹⁶.

One challenge with respect to the mechanisms of antibody-mediated TRALI is to elucidate the biological features of the interactions between the Fc part of immunoglobulins G (IgGs) and their specific Fc γ receptors (Fc γ Rs) present on immune cells. In different studies, Fc γ Rs were found to be major ^{9,13,17} or accessory ⁷ mediators of anti-MHC I-induced lung injury. Concerning the repertoire of Fc γ Rs, one major difference between man and mouse is the

presence in the human genome of the *FCGR2A* gene, encoding an activating low affinity receptor for IgGs called Fc γ RIIA, also known as CD32A¹⁸. This receptor is expressed on various immune cells of myeloid origin susceptible to participate in antibody-mediated TRALI responses, namely neutrophils, monocytes, macrophages, dendritic cells and platelets. It bears an intracellular immunoreceptor tyrosine-based activation motif (ITAM) that mediates cell activation upon its crosslinking by IgG-immune complexes (IgG-ICs)¹⁹. Unlike leukocytes, which express various types of Fc γ R, human platelets possess only Fc γ RIIA/CD32A.

It is well established that IgG-ICs can strongly activate human platelets, resulting in their aggregation and the release of platelet granule contents²⁰⁻²². In contrast, Fc γ RIIA/CD32A is absent from mice and mouse platelets, devoid of any Fc γ R, cannot be activated by IgG-ICs. Thus, in the current murine models of antibody-mediated TRALI, Fc γ R-dependent responses would appear to be biased towards the involvement of circulating leukocytes and/or macrophages, while ignoring the potential contribution of Fc γ R-mediated platelet responses to the pathogenesis of TRALI in man. Accordingly, we have reported that in WT animals, Fc γ R⁺ patrolling monocytes and/or macrophages are required, while Fc γ R⁻ platelets are dispensable for the early steps of experimental antibody-induced TRALI leading to the formation of lung edema^{8,12}. These concerns prompted us to make use of transgenic mice expressing human Fc γ RIIA/CD32A, especially as these mice have been shown to reliably mimic several human physiological and pathological responses. Indeed, in these animals, platelet Fc γ RIIA/CD32A participates not only in protective immune responses²⁰, but also in severe IgG-IC-dependent anaphylaxis^{21,23}, heparin-induced thrombocytopenia²⁴ and sepsis²⁵.

In this work, we explored the role of Fc γ RIIA/CD32A in the pathogenesis of TRALI using a new model of antibody-mediated TRALI induced by LPS sensitization and challenge with a recombinant chimeric human/mouse anti-mouse MHC I (H-2D^d/H-2K^d) mAb in mice expressing a human Fc γ RIIA/CD32A transgene. This strategy revealed a crucial contribution of platelet Fc γ RIIA/CD32A to antibody-induced TRALI through platelet activation and serotonin release,

with exacerbation of TRALI responses, thereby opening up a new dimension in our understanding of the pathogenesis of TRALI in humans.

Methods

Mice

Transgenic mice expressing human Fc γ RIIA/CD32A (C57BL/6, H-2^b)²⁶ were kindly provided by P. Bruhns (Institut Pasteur, France) and bred in our animal facilities. These animals were crossed with BALB/cByJ mice (H-2^d) (JAXTM, Charles River). The presence of the hCD32A transgene was assessed by PCR analysis of genomic DNA using the primers 5'-CAATTTTGCTGCTATGGGC-3' and 5'-CTGGTCAAGGTCACATTCTTC-3'. TRALI experiments were performed on the F1 progeny (C57BL/6 x BALB/c) using 6-7 week-old male mice expressing the human Fc γ RIIA/CD32A transgene (CD32A⁺) and their wild-type littermates (CD32A⁻). Mice were housed in the animal facilities of the EFS-Grand Est (agreement N°E67-482-10). Experiments were approved by the Regional Committee for Ethics in Animal Experimentation of Strasbourg (CREMEAS, CEEA-35) and authorized by the French Ministry of Research, according to the regulations of the European Community (APAFIS#2018100815412773).

Two-event TRALI model

A previously described two-hit model⁵ was used with a few modifications. Male mice were primed with an intraperitoneal (i.p.) injection of LPS (0.1 mg/kg body weight) and 24 h later, anesthetized with xylazine (20 mg/kg body weight) and ketamine (100 mg/kg body weight). TRALI was induced by retro-orbital injection of the recombinant chimeric human IgG1/mouse 34-1-2S anti-MHC I mAb (anti-H-2D^d/H-2K^d, 1.5 mg/kg). The animals were maintained on warming plate heated to 37°C and sacrificed 10-15 min or 2 h after injection of the mAb. TRALI responses were assessed by measuring the protein content of bronchoalveolar lavages (BALs) in right lungs and by analyzing the pulmonary lesions in left lungs by histochemistry⁸. Animals used as negative controls received saline. To visualize vascular leakage, 1% Evans Blue (50 μ L) was injected intravenously (i.v.) prior to injection of the anti-MHC I mAb. All experiments were performed by operators blind to the genotype of the mice.

Statistical analyses

Statistical analyses were performed with GraphPad software (GraphPad Prism 5.02). Survival was analyzed using the log-rank test. All other values were reported as the mean \pm SEM and analyzed using 1- or 2-way ANOVA followed by a Dunn's test or a Bonferroni post hoc analysis, depending on the normal distribution of the data established using the Shapiro-Wilk test. For figures 3A,B, 4 and 5A,E,G, statistical analyses were performed without considering the negative controls, i.e., mice injected with saline instead of the anti-MHC I mAb, because the n-values of the negative controls were lower than those of the other conditions. These low n-values of the negative control animals could be justified by the high reproducibility of the measurements between independent experiments and the ethical requirement to reduce the number of animals.

An extended methods section is available in the online Supplemental Material.

Results

Fc γ RIIA/CD32A expression severely aggravates the alveolar edema and increases mortality of antibody-mediated TRALI.

To set up a humanized model of TRALI allowing evaluation of the biological activity of CD32A, we first cloned the variable regions of the heavy and light chains of the mouse mAb 34-1-2S and then fused them to the constant regions of the human IgG1 (hIgG1) heavy chain and kappa light chain, respectively, thus generating a chimeric human/mouse anti-MHC I mAb, called hIgG1-34-1-2S. Human IgG1 was preferred to other IgG subclasses on account of its biological activities which better resemble those of mouse IgG2a, together with its higher affinity for CD32A as compared to other human IgG subclasses ²⁷. Indeed, both the recombinant mAb hIgG1-34-1-2S and the original mouse mAb IgG2a-34-1-2S bound to H-2K^d MHC I with comparable affinity and stained circulating CD45⁺ leukocytes from BALB/c mice with the same efficacy (**Supplemental Figure 1A, B**). In addition, hIgG1-34-1-2S- and mIgG2a-34-1-2S-containing immune complexes readily bound to Fc γ RIIA/CD32A and all mouse Fc γ Rs-expressing CHO transfectants, whereas monomeric hIgG1-34-1-2S or mIgG2a-34-1-2S only bound efficiently to mFc γ RI- and Fc γ RIV-expressing cells (**Supplemental Figure 1C**). Therefore, we may anticipate that hIgG1-34-1-2S can activate all endogenous mouse Fc γ Rs, in addition to Fc γ RIIA/CD32A.

hIgG1-34-1-2S was administered to LPS-sensitized (0.1 mg/kg, i.p.) CD32A⁻ and CD32A⁺ mice ⁵ and lung alveolar edema, a typical feature of experimental TRALI, was evaluated by quantifying the proteins in bronchoalveolar lavages (BALs) 15 min later. The protein content of BALs increased significantly in CD32A⁻ and CD32A⁺ mice as compared to negative control animals. The increase was greater in CD32A⁺ mice, reaching twice the levels in CD32A⁻ animals (1.11±0.14 vs 0.48±0.09 mg/mL, CD32A⁺ vs CD32A⁻, respectively, **p<0.01), indicating that the alveolar edema was more severe in CD32A⁺ than in CD32A⁻ mice (**Figure 1A**). Histological evaluation indicated that acute lung injury was also significantly

increased in CD32A⁺ as compared to CD32A⁻ mice (**Figure 1B**). The enhanced pulmonary vascular permeability in CD32A⁺ animals was confirmed by measurement of the extravasation of Evans Blue dye into the lung tissues and BALs 15 min after injection of hlgG1-34-1-2S (**Figure 1C**). This increased vascular permeability appeared to be specific to the lungs, since no extravasation of the dye could be observed in other vascular territories of CD32A⁺ mice such as the glabrous skin of the ears, snout and paws (**Figure 1D**). Within 2 h following hlgG1-34-1-2S injection, 40% of the CD32A⁻ mice died, while this percentage increased to 90% in CD32A⁺ mice (**p=0.0027, log-rank test) (**Figure 1E, left panel**). At death, or at the 2 h time point, alveolar edema was also more severe in CD32A⁺ than in CD32A⁻ mice (**Figure 1E, right panel**). Altogether, these results indicated that Fc γ RIIA/CD32A expression exacerbates the disruption of the pulmonary endothelial barrier provoked by antibody-mediated TRALI, leading to reduced survival.

Fc γ RIIA/CD32A expression increases the activation of circulating neutrophils and platelets during antibody-mediated TRALI

Since all blood cell populations expressing CD32A could potentially contribute to this effect, we first evaluated the activation state of circulating leukocytes and platelets during TRALI. Baseline blood cell counts were similar between CD32A⁻ and CD32A⁺ mice (**Supplemental Table 1**). Within 15 min of injection of hlgG1-34-1-2S, plasma levels of the granular protein myeloperoxidase (MPO) were significantly higher in CD32A⁺ as compared to CD32A⁻ mice (**Figure 2A**), suggesting an increased activation of blood neutrophils in CD32A⁺ mice. Immunohistology using an anti-mouse neutrophil antibody showed increased neutrophil sequestration in the lungs of CD32A⁺ as compared to CD32A⁻ mice (**Figure 2B**). In addition, the drop of circulating platelet count was more severe in CD32A⁺ than in CD32A⁻ mice (**Figure 2C**), suggesting enhanced platelet activation in CD32A⁺ animals. Indeed, the circulating platelets of CD32A⁺ mice exhibited higher CD62P expression as compared to those of CD32A⁻ mice (**Figure 2D**). Immunohistology using an anti-mouse platelet antibody provided evidence for increased platelet sequestration in the lungs of CD32A⁺ as compared to CD32A⁻ animals

(**Figure 2E**). Histology also revealed the presence of a few occlusive thrombi in the pulmonary microcirculation, but in similar proportions in CD32A⁺ and CD32A⁻ mice (**Supplemental Figure 2**), ruling out a major contribution of thrombi to the increased edema formation. Thus, the activation of platelets and neutrophils was quantitatively more important in CD32A⁺ animals experiencing antibody-mediated TRALI.

Monocytes/macrophages are required in WT mice but only accessory in CD32A⁺ mice for the initiation of antibody-mediated TRALI

Given the increase in the activation state and pulmonary sequestration of neutrophils in CD32A⁺ mice during TRALI, we looked at whether these cells contributed to the aggravation of lung injury. Neutrophil depletion (**Supplemental Figure 3A, B**) had no significant impact on TRALI in CD32A⁻ mice (**Figure 3A**), in agreement with our previous observations in the context of TRALI induced by the mIgG2a mAb 34-1-2S in WT mice¹². Depletion of neutrophils in CD32A⁺ animals did not reduce but rather slightly increased the protein content of BALs (**Figure 3A**), an effect which cannot be easily explained. In any case, neutrophils did not appear to be responsible for the aggravation of alveolar edema in this humanized model of antibody-mediated TRALI.

Next, we investigated whether monocytes/macrophages contributed to lung injury in CD32A⁺ mice. These cells were eliminated using clodronate liposomes, delivered 18 h after LPS, i.e., 6 h before anti-MHC I mAb injection to allow inflammation to proceed. Under these conditions, TRALI was totally abolished in CD32A⁻ mice (**Figure 3B**), in agreement with our previous observations in the context of TRALI induced by the mIgG2a mAb 34-1-2S in WT mice¹². In contrast, depletion of monocytes/macrophages in CD32A⁺ animals only partially reduced the protein content of BALs (**Figure 3B**), indicating that these cells played an accessory role and thereby pointing to the decisive intervention of another cell population.

Platelets are required for the Fc γ RIIA/CD32A-induced aggravation of antibody-mediated TRALI

The enhanced activation of platelets during TRALI in CD32A⁺ mice raised the question as to whether platelet activation or consumption was a marker of disease severity or actively contributed to the aggravation of TRALI. To explore the role of platelets in TRALI, we depleted circulating platelets using the anti-GPIb α mAb cocktail R300, 48 h before inducing TRALI (**Supplemental Figure 4**). Under these conditions, the alveolar edema associated with IgG1-34-1-2S-mediated TRALI was not modified in CD32A⁻ mice (**Figure 4**), consistent with our previous published work demonstrating that platelets are dispensable for TRALI induced by the mIgG2a mAb 34-1-2S in WT mice⁸. In contrast, the protein content of BALs was significantly reduced in thrombocytopenic CD32A⁺ mice, falling to the level observed in CD32A⁻ animals, although remaining significantly above resting levels (**Figure 4**). These results showed that platelet Fc γ RIIA/CD32A activation exacerbates antibody-mediated TRALI.

Platelet-released serotonin, but not platelet-activating factor, is responsible for the Fc γ RIIA/CD32A-induced aggravation of antibody-mediated TRALI

We next wondered whether a mediator derived from platelets or released from their granules might be responsible for this effect. Platelet-activating factor (PAF), which is produced by activated platelets and plays a key role during anaphylaxis²⁸, was not involved in platelet Fc γ RIIA/CD32A-mediated aggravation of TRALI, since the PAF receptor antagonist ABT-491 had no impact on the protein content of BALs in either CD32A⁻ or CD32A⁺ mice (**Figure 5A**).

We next evaluated whether platelet granule contents were released during TRALI. Within 15 min of the onset of TRALI, platelet factor 4 (PF4) (**Figure 5B**) and serotonin (**Figure 5C**), stored in alpha and dense granules respectively and released during activation-induced platelet degranulation, were detected at high concentrations in the plasma of CD32A⁺ but not of CD32A⁻ animals. Serotonin, a known inducer of vascular permeability²⁹⁻³¹, has been recently identified by us and others as playing a critical part in platelet-dependent IgG-mediated anaphylaxis in CD32A⁺ transgenic animals^{21,25}. The vast majority of the total serotonin pool of the body is stored in platelet dense granules³². Platelets do not synthesize serotonin but use

a serotonin transporter (SERT) to capture plasma serotonin ³². Blocking platelet serotonin uptake with the SERT inhibitor fluoxetine, resulted in almost complete depletion of the serotonin content of dense granules (**Figure 5D**). Whereas fluoxetine treatment had no observable impact on the lung edema in CD32A⁻ mice, it abolished its aggravation in CD32A⁺ mice (**Figure 5E**). These findings indicated that platelet-released serotonin was responsible for the Fc γ RIIA/CD32A-dependent exacerbation of TRALI. One may note that the exacerbation of TRALI-associated thrombocytopenia still occurred in serotonin-depleted CD32A⁺ mice (**Figure 5F**), suggesting that serotonin, although a platelet activator, was not responsible for this effect.

Interestingly, release of platelet serotonin occurred normally during the residual TRALI in clodronate-treated CD32A⁺ mice (**Figure 5G**), suggesting that Fc γ RIIA/CD32A-mediated platelet activation and serotonin release was sufficient to initiate TRALI in transgenic animals. Conversely, monocyte/macrophage depletion completely abolished TRALI in fluoxetine-treated CD32A⁺ mice (**Figure 5H**), strongly suggesting that release of serotonin from platelets was responsible for the low grade TRALI responses in clodronate-treated CD32A⁺ animals. The TRALI-associated thrombocytopenia remained exacerbated in CD32A⁺ mice with or without clodronate treatment (**Figure 5I**). In contrast, TRALI-associated platelet consumption was inhibited in CD32A⁻ mice treated with clodronate (**Figure 5I**), indicating that in WT animals, but not in transgenic ones, platelet consumption partly relied on monocytes and/or macrophages.

Blockade of the 5-hydroxytryptamine 2A serotonin receptor prevents the exacerbation of antibody-mediated TRALI in CD32A⁺ mice

Finally, we investigated how the serotonin released by platelets expressing Fc γ RIIA/CD32A could aggravate the development of TRALI-associated lung edema. Among the numerous serotonin receptor subtypes present on circulating blood cells and cells of the vessel wall, the 5-hydroxytryptamine (5-HT) 2A receptor has been shown to mediate the increase in vascular

permeability occurring during IgG-IC-induced CD32A-mediated shock ²⁵. To explore the involvement of this receptor in the serotonin-induced exacerbation of lung edema in CD32A⁺ mice, we used the selective 5-HT_{2A} antagonist sarpogrelate ³³. Intravenous injection of sarpogrelate 5 min before injection of hlgG1-34-1-2S had no significant effect on lung edema in CD32A⁻ mice, but prevented its aggravation in CD32A⁺ animals (**Figure 6A**). The 5-HT_{2A} serotonin receptor therefore appeared to be responsible for the serotonin-induced exacerbation of Fc_γRIIA/CD32A-dependent TRALI.

These results suggested that the 5-HT_{2A} receptor could represent a pharmacological target to attenuate TRALI in humans. To test this hypothesis, we first checked the kinetics of TRALI development by measuring lung respiration parameters. Five min following challenge with hlgG1-34-1-2S, CD32A⁺ animals showed increased signs of respiratory distress, as evidenced by their enhanced pulmonary elastance as compared to CD32A⁻ mice (**Figure 6B**). Thus, when sarpogrelate was injected at this time point, it effectively abolished the aggravation of lung edema occurring in CD32A⁺ mice, but not in CD32A⁻ animals (**Figure 6C**). These findings showed that sarpogrelate prevents the exacerbation of TRALI mediated by platelet Fc_γRIIA/CD32A, which would provide a rationale for targeting serotonin receptors as a therapeutic strategy for TRALI in humans.

Discussion

The pathophysiology of antibody-mediated TRALI involves complex interactions between various blood cells, IgGs and molecular mediators, rendering difficult the identification of the main molecular and cellular effectors of this pathology. Hence the availability of murine models of TRALI has permitted a better understanding of the mechanisms implicated in its pathogenesis. In the present study, we describe a new mouse model of antibody-mediated TRALI, based on the expression of a human Fc γ RIIA/CD32A transgene and the use of a recombinant chimeric human/mouse mAb derived from the anti-MHC I mouse mAb 34-1-2S. Remarkably, the TRALI reactions induced by this recombinant antibody - namely, the early increases in pulmonary elastance, the protein content of BALs and mortality - were more severe in transgenic than in WT animals. Our results indicate that platelet Fc γ RIIA/CD32A activation exacerbates antibody-mediated TRALI through its involvement in platelet serotonin release and the subsequent activation of 5-HT_{2A} serotonin receptors. These findings provide a rationale for targeting serotonin receptors to attenuate TRALI in humans.

Our data identify Fc γ RIIA/CD32A as an important determinant of the severity of antibody-mediated TRALI. Nevertheless, a contribution of Fc γ RI and Fc γ RIV to TRALI cannot be excluded, given that these receptors have previously been shown to contribute to IgG2a-mediated anaphylaxis³⁴. In addition, a potential contribution of complement cannot be excluded as recently highlighted in the case of TRALI-inducing *versus* TRALI-resistant antibodies^{16,35}.

We and others have previously reported that, in WT mice, monocytes/macrophages are essential to induce TRALI with the mouse IgG2a anti-MHC I mAb 34-1-2S^{7,12,13,15}, while neutrophils and platelets are dispensable^{7,8,11,12}. Controversies have long existed, however, especially regarding the contribution of the recipient platelets to the pathology, notwithstanding the recent consensus that platelets may not be critically required for the onset and development of TRALI in WT mice^{36,37}. Here, similar cell-dependent features were observed

in CD32A⁻ animals when TRALI was induced by the chimera hIgG1-34-1-2S. In CD32A⁺ mice, the assessed TRALI responses induced by hIgG1-34-1-2S did not appear to require the presence of circulating neutrophils, demonstrating that expression of Fc γ R1IA/CD32A in mice does not confer a major additional function on neutrophils during TRALI. Remarkably, liposome clodronate sensitive cells (monocytes, macrophages and/or dendritic cells) were not required for the initiation of lung edema in CD32A⁺ mice, although they contributed to it, pointing to a new mechanism and to the decisive contribution of another cell type.

The administration of hIgG1-34-1-2S induced thrombocytopenia in WT and CD32A⁺ mice probably by two different mechanisms. In WT animals, monocyte/macrophage depletion reduced thrombocytopenia, suggesting that some of the anti-MHC I-opsonized platelets were captured by liver and/or splenic macrophages. In contrast, in CD32A⁺ mice, thrombocytopenia was not affected by macrophage depletion. This effect could result firstly from an increased adsorption of platelets on the opsonized endothelium, most likely through an Fc γ R1IA/CD32A-dependent mechanism, as suggested by the enhanced immunohistochemical staining of platelets on lung sections of CD32A⁺ mice. Secondly, the increased activation state of these platelets, as revealed by the large amounts of granule-specific molecules released into the blood of CD32A⁺ mice during TRALI, might stabilize this adsorption. Consequently, in transgenic animals, platelets would be efficiently sequestered on the vessel wall.

The massive release of platelet granule contents also suggests that the amplification of TRALI responses in these animals is platelet-dependent. In agreement, prior depletion of circulating platelets abolished the aggravation of TRALI observed in CD32A⁺ mice. In fact, our analysis identified platelet-derived serotonin as the mediator responsible for Fc γ R1IA/CD32A-induced aggravation of TRALI, since the latter was abrogated when platelet serotonin stores were pharmacologically depleted *in vivo* prior to induction of TRALI. Notably, in CD32A⁺ mice, low grade lung edema occurred in the absence of monocytes and macrophages and was completely abolished by fluoxetine treatment, strongly suggesting that Fc γ R1IA/CD32A-

mediated platelet activation and serotonin release are sufficient to initiate the pathogenesis of TRALI.

A major finding was that the 5-HT_{2A} serotonin receptor subtype is responsible for Fc γ RIIIA/CD32A-induced exacerbation of lung edema formation. This property could result from the particular physiology of lung circulation. By targeting 5-HT_{2A}⁺ vascular smooth muscle cells (VSMCs), serotonin provokes pulmonary arterial vasoconstriction and subsequent increase in pulmonary arterial pressure ^{38,39}. We previously proposed that the transient increase of pulmonary arterial pressure was responsible for lung edema formation during TRALI ¹². This hypothesis was based on a bundle of observations: (i) alveolar edema occurs when interstitial edema exceeds a threshold due to plasma leakage through the vessel wall before flooding into the alveoli ⁴⁰, (ii) during anti-MHC I-induced TRALI, the pre-capillary arterioles, but not the post-capillary venules, are the exclusive site of edema formation ¹², (iii) NF449, a selective P2X₁ receptor antagonist, which blocks VSMC constriction, also inhibits TRALI ¹². Similarly, the contraction of pulmonary VSMCs is positively regulated by 5-HT_{2A} serotonin receptors present on their surface ^{38,39} and inhibited by specific 5-HT₂ receptor antagonists ^{41,42}. Hence it is tempting to speculate that the abrogation of the increase in pulmonary edema by sarpogrelate might be due to the inhibition of serotonin-induced pulmonary periarteriolar VSMC contraction.

Whether the transient increase of pulmonary arterial pressure does occur during the very early steps of TRALI in humans is a question difficult to address because pulmonary arterial pressure is rarely measured during transfusion of a patient. Nevertheless, in two cases of human TRALI observed after the transfusion of blood products containing anti-HLA class I antibodies, the first manifestation of the reaction was an increase in pulmonary arterial blood pressure, which occurred before the onset of hypoxemia and acute respiratory distress, while at the same time systemic arterial blood pressure decreased ^{43,44}. These cases point to the possibility that serotonin-induced pulmonary arterial contraction is an early event in the development of TRALI. Of note, whereas serotonin acts as a powerful vasoconstrictor in human pulmonary arteries, it acts as a vasodilator in peripheral vessels ^{45,46}. One may

speculate that the vascular territory-dependent effects of serotonin could explain the specific localization of TRALI lesions in pulmonary tissue. Whether the use of selective serotonin reuptake inhibitors in patients with depression is associated with fewer cases of antibody-mediated TRALI is currently unknown.

TRALI is an ARDS, the mainstay treatment of which remains general supportive care, including fluid resuscitation or mechanical ventilation ⁴⁷. No pharmacological agents able to improve the outcome of ARDS have yet been identified. Nevertheless, experiments in the mouse have suggested that administration of IL-10 or CRP inhibitors could represent promising therapeutic or prophylactic therapies against TRALI ¹⁴. Our findings now indicate that administration of sarpogrelate after the onset of TRALI efficiently inhibits edema formation in CD32A⁺ mice. These results therefore provide a rationale for targeting serotonin receptors to block the pathological progression of TRALI.

Overall, the present humanized model of antibody-mediated TRALI offers new insights into the mechanisms underlying TRALI as it may occur in humans, while identifying a crucial contribution of platelet Fc γ R1IA/CD32A and serotonin release to the severity of the syndrome. The same mechanism also underlines the role of platelet serotonin in lung pathophysiology as recently highlighted in IgG-dependent anaphylaxis studies ^{21,25}. This would suggest a similar crucial contribution of platelet Fc γ R1IA/CD32A to antibody-induced TRALI and IgG-dependent anaphylaxis, through platelet activation and serotonin release, leading to increased vascular permeability and edema formation. Future clinical investigations will aim at determining how our observations can be translated into the human health care setting. In addition to blood transfusion therapy, these findings could have implications in other fields relevant to antibody-mediated damage where platelets are active players such as organ transplantation and acute rejection.

Data Sharing Statement

For original data, please contact beatrice.hechler@efs.sante.fr

Acknowledgements

This work was supported by a project grant from the EFS (APR 2014) and by INSERM, EFS and ARMESA (Association de Recherche et Développement en Médecine et Santé Publique). F.J. is an employee of the CNRS (Centre National de la Recherche Scientifique). The authors would like to thank M. Freund and her collaborators for animal care, Laurie Ruch and Oceane Schlienger for expert technical assistance and the NIH Tetramer Core Facility (tetramer.yerkes.emory.edu) for provision of MHC-peptide complexes. We are particularly grateful to Patrick England and the Plateforme de Biophysique Moléculaire (PFBMI), Institut Pasteur (Paris, France), for help with BLI measurements and to J. Mulvihill for revising the English of the manuscript. The visual abstract was created using <https://biorender.com>.

Authors' Contributions

M.-B.E.M., B.M., S.M., F.T., F.J. and B.H. performed experiments and analyzed data; F.J. contributed essential material; H.dS., B.M. and B.H. designed the research and wrote the manuscript; C.G. and F.J. critically reviewed the manuscript; all authors contributed to the final manuscript.

Conflict-of-Interest Statement

The authors declare no competing financial interest.

References

1. Hendrickson JE, Roubinian NH, Chowdhury D, et al. Incidence of transfusion reactions: a multicenter study utilizing systematic active surveillance and expert adjudication. *Transfusion*. 2016;56(10):2587-2596.
2. Vlaar APJ, Kleinman S. An Update of the Transfusion-Related Acute Lung Injury (TRALI): A Proposed Modified Definition and Classification Scheme Definition. *Indian J Hematol Blood Transfus*. 2020;36(3):556-558.
3. Semple JW, Rebetz J, Kapur R. Transfusion-associated circulatory overload and transfusion-related acute lung injury. *Blood*. 2019;133(17):1840-1853.
4. Dry SM, Bechard KM, Milford EL, Churchill WH, Benjamin RJ. The pathology of transfusion-related acute lung injury. *Am J Clin Pathol*. 1999;112(2):216-221.
5. Looney MR, Nguyen JX, Hu Y, Van Ziffle JA, Lowell CA, Matthay MA. Platelet depletion and aspirin treatment protect mice in a two-event model of transfusion-related acute lung injury. *J Clin Invest*. 2009;119(11):3450-3461.
6. Fung YL, Kim M, Tabuchi A, et al. Recipient T lymphocytes modulate the severity of antibody-mediated transfusion-related acute lung injury. *Blood*. 2010;116(16):3073-3079.
7. Strait RT, Hicks W, Barasa N, et al. MHC class I-specific antibody binding to nonhematopoietic cells drives complement activation to induce transfusion-related acute lung injury in mice. *J Exp Med*. 2011;208(12):2525-2544.
8. Hechler B, Maitre B, Magnenat S, et al. Platelets are dispensable for antibody-mediated transfusion-related acute lung injury in the mouse. *J Thromb Haemost*. 2016;14(6):1255-1267.
9. Looney MR, Su X, Van Ziffle JA, Lowell CA, Matthay MA. Neutrophils and their Fc gamma receptors are essential in a mouse model of transfusion-related acute lung injury. *J Clin Invest*. 2006;116(6):1615-1623.
10. Peters AL, Van Stein D, Vlaar AP. Antibody-mediated transfusion-related acute lung injury; from discovery to prevention. *Br J Haematol*. 2015;170(5):597-614.
11. Cognasse F, Tariket S, Hamzeh-Cognasse H, et al. Platelet depletion limits the severity but does not prevent the occurrence of experimental transfusion-related acute lung injury. *Transfusion*. 2020;60(4):713-723.

12. El Mdawar MB, Maitre B, Magnenat S, Gachet C, Hechler B, de la Salle H. The ATP-gated P2X1 ion channel contributes to the severity of antibody-mediated Transfusion-Related Acute Lung Injury in mice. *Sci Rep.* 2019;9(1):5159.
13. McKenzie CG, Kim M, Singh TK, Milev Y, Freedman J, Semple JW. Peripheral blood monocyte-derived chemokine blockade prevents murine transfusion-related acute lung injury (TRALI). *Blood.* 2014;123(22):3496-3503.
14. Kapur R, Kim M, Aslam R, et al. T regulatory cells and dendritic cells protect against transfusion-related acute lung injury via IL-10. *Blood.* 2017;129(18):2557-2569.
15. Kapur R, Kasetty G, Rebetz J, Egesten A, Semple JW. Osteopontin mediates murine transfusion-related acute lung injury via stimulation of pulmonary neutrophil accumulation. *Blood.* 2019;134(1):74-84.
16. Cleary SJ, Kwaan N, Tian JJ, et al. Complement activation on endothelium initiates antibody-mediated acute lung injury. *J Clin Invest.* 2020.
17. Semple JW, Kim M, Hou J, et al. Intravenous immunoglobulin prevents murine antibody-mediated acute lung injury at the level of neutrophil reactive oxygen species (ROS) production. *PLoS One.* 2012;7(2):e31357.
18. Qiao J, Al-Tamimi M, Baker RI, Andrews RK, Gardiner EE. The platelet Fc receptor, FcγRIIIa. *Immunol Rev.* 2015;268(1):241-252.
19. Daeron M. Fc receptor biology. *Annu Rev Immunol.* 1997;15:203-234.
20. Arman M, Krauel K. Human platelet IgG Fc receptor FcγRIIIA in immunity and thrombosis. *J Thromb Haemost.* 2015;13(6):893-908.
21. Beutier H, Hechler B, Godon O, et al. Platelets expressing IgG receptor FcγRIIIA/CD32A determine the severity of experimental anaphylaxis. *Sci Immunol.* 2018;3(22).
22. Duffau P, Seneschal J, Nicco C, et al. Platelet CD154 potentiates interferon-alpha secretion by plasmacytoid dendritic cells in systemic lupus erythematosus. *Sci Transl Med.* 2010;2(47):47ra63.
23. Jonsson F, Mancardi DA, Zhao W, et al. Human FcγRIIIA induces anaphylactic and allergic reactions. *Blood.* 2012;119(11):2533-2544.
24. Reilly MP, Taylor SM, Hartman NK, et al. Heparin-induced thrombocytopenia/thrombosis in a transgenic mouse model requires human platelet factor 4 and platelet activation through FcγRIIIA. *Blood.* 2001;98(8):2442-2447.

25. Cloutier N, Allaey I, Marcoux G, et al. Platelets release pathogenic serotonin and return to circulation after immune complex-mediated sequestration. *Proc Natl Acad Sci U S A*. 2018;115(7):E1550-E1559.
26. McKenzie SE, Taylor SM, Malladi P, et al. The role of the human Fc receptor Fc gamma RIIA in the immune clearance of platelets: a transgenic mouse model. *J Immunol*. 1999;162(7):4311-4318.
27. Overdijk MB, Verploegen S, Ortiz Buijsse A, et al. Crosstalk between human IgG isotypes and murine effector cells. *J Immunol*. 2012;189(7):3430-3438.
28. Ishii S, Kuwaki T, Nagase T, et al. Impaired anaphylactic responses with intact sensitivity to endotoxin in mice lacking a platelet-activating factor receptor. *J Exp Med*. 1998;187(11):1779-1788.
29. Cloutier N, Pare A, Farndale RW, et al. Platelets can enhance vascular permeability. *Blood*. 2012;120(6):1334-1343.
30. Kubes P, Gaboury JP. Rapid mast cell activation causes leukocyte-dependent and -independent permeability alterations. *Am J Physiol*. 1996;271(6 Pt 2):H2438-2446.
31. Majno G, Palade GE. Studies on inflammation. 1. The effect of histamine and serotonin on vascular permeability: an electron microscopic study. *J Biophys Biochem Cytol*. 1961;11:571-605.
32. Adnot S, Houssaini A, Abid S, Marcos E, Amsellem V. Serotonin transporter and serotonin receptors. *Handb Exp Pharmacol*. 2013;218:365-380.
33. Saini HK, Takeda N, Goyal RK, Kumamoto H, Arneja AS, Dhalla NS. Therapeutic potentials of sarpogrelate in cardiovascular disease. *Cardiovasc Drug Rev*. 2004;22(1):27-54.
34. Khodoun MV, Kucuk ZY, Strait RT, et al. Rapid desensitization of mice with anti-Fc gammaRIIb/Fc gammaRIII mAb safely prevents IgG-mediated anaphylaxis. *J Allergy Clin Immunol*. 2013;132(6):1375-1387.
35. Zeeuw van der Laan EAN, van der Velden S, Bentlage AEH, et al. Biological and structural characterization of murine TRALI antibody reveals increased Fc-mediated complement activation. *Blood Adv*. 2020;4(16):3875-3885.
36. Semple JW, Kapur R. The contribution of recipient platelets in TRALI: has the jury reached a verdict? *Transfusion*. 2020;60(5):886-888.

37. Zeeuw van der Laan EAN, van der Velden S, Porcelijn L, Semple JW, van der Schoot CE, Kapur R. Evaluation of Platelet Responses in Transfusion-Related Acute Lung Injury (TRALI). *Transfus Med Rev.* 2020.
38. MacLean MR, Sweeney G, Baird M, McCulloch KM, Houslay M, Morecroft I. 5-Hydroxytryptamine receptors mediating vasoconstriction in pulmonary arteries from control and pulmonary hypertensive rats. *Br J Pharmacol.* 1996;119(5):917-930.
39. MacLean MR, Herve P, Eddahibi S, Adnot S. 5-hydroxytryptamine and the pulmonary circulation: receptors, transporters and relevance to pulmonary arterial hypertension. *Br J Pharmacol.* 2000;131(2):161-168.
40. Staub NC. Pulmonary edema. *Physiol Rev.* 1974;54(3):678-811.
41. Desmecht D, Linden A, Amory H, Lekeux P. Hemodynamic responses to *Pasteurella haemolytica* inoculation in calves given type 2 serotonergic antagonist. *Can J Physiol Pharmacol.* 1996;74(5):572-579.
42. Selig WM, Bloomquist MA, Cohen ML, Fleisch JH. Serotonin-induced pulmonary responses in the perfused guinea pig lung: evidence for 5HT₂ receptor-mediated pulmonary vascular and airway smooth muscle constriction. *Pulm Pharmacol.* 1988;1(2):93-99.
43. Kojima T, Nishisako R, Sato H. A patient with possible TRALI who developed pulmonary hypertensive crisis and acute pulmonary edema during cardiac surgery. *J Anesth.* 2012;26(3):460-463.
44. Lin Y, Kanani N, Naughton F, Pendergrast J, Karkouti K. Case report: transfusion-related acute lung injury (TRALI) - a clear and present danger. *Can J Anaesth.* 2007;54(12):1011-1016.
45. Egermayer P, Town GI, Peacock AJ. Role of serotonin in the pathogenesis of acute and chronic pulmonary hypertension. *Thorax.* 1999;54(2):161-168.
46. Rudolph AM, Paul MH. Pulmonary and systemic vascular response to continuous infusion of 5-hydroxytryptamine (serotonin) in the dog. *Am J Physiol.* 1957;189(2):263-268.
47. Vlaar AP, Juffermans NP. Transfusion-related acute lung injury: a clinical review. *Lancet.* 2013;382(9896):984-994.

Figure Legends

Figure 1: Fc γ RIIA/CD32A expression severely aggravates the alveolar edema and increases the mortality of antibody-mediated TRALI.

The pathogenesis of TRALI was analyzed 15 min (**A, B, C, D**) or 2 h (**E**) after injection of the chimeric human/mouse anti-MHC I mAb hlgG1-34-1-2S (1.5 mg/kg, i.v.) into LPS-sensitized (0.1 mg/kg, i.p.) CD32A⁻ or CD32A⁺ mice. **A.** Protein concentrations in BALs were measured 15 min after injection of hlgG1-34-1-2S or in negative controls (CD32A⁻: n=9-20, CD32A⁺: n=3-20). **B.** The histological features of lung injury were determined 15 min after injection of hlgG1-34-1-2S or in negative controls (CD32A⁻: n=7-17, CD32A⁺: n=7-19) (Scale bar, 200 μ m). **C.** Exemplary photographs showing increased pulmonary vascular permeability in CD32A⁺ as compared to CD32A⁻ mice revealed by extravasation of Evans Blue dye into the lung tissue and BALs 15 min after administration of hlgG1-34-1-2S. **D.** Extravasation of Evans Blue dye was not observed in the vascular territories of the glabrous skin of the ears, snout or paws 15 min after injection of hlgG1-34-1-2S or saline into CD32A⁺ or CD32A⁻ mice. **E.** Left: Kaplan-Meier survival plots for CD32A⁻ and CD32A⁺ mice (n=10, p=0.0027, log-rank test). Right: Protein concentrations in BALs of CD32A⁻ and CD32A⁺ mice at death or at the 2 h endpoint. Results are presented as the mean \pm SEM. *p<0.05, **p<0.01, ***p<0.001 by 1-way ANOVA (**A, B**) or a Student's t-test (**E**, right).

Figure 2: Fc γ RIIA/CD32A expression increases the activation of circulating neutrophils and platelets during antibody-mediated TRALI.

LPS-sensitized (0.1 mg/kg, i.p.) CD32A⁻ or CD32A⁺ mice were injected with the mAb hlgG1-34-1-2S (1.5 mg/kg, i.v.) or vehicle and examined 15 min later. **A.** Blood plasma levels of myeloperoxidase (MPO) (CD32A⁻: n=5-11, CD32A⁺: n=5-12). **B.** Left: Neutrophil areas in the lungs of CD32A⁻ and CD32A⁺ mice (n=3). Right: Representative lung sections from CD32A⁻ and CD32A⁺ mice immunolabeled with the anti-neutrophil mAb 1A8 (Scale bar, 50 μ m). **C.**

Circulating platelet counts before administration of LPS, 24 h later and 15 min after hlgG1-34-1-2S challenge (n=8). **D.** Percentage of CD62P-positive platelets (n=6). **E.** Left: Platelet areas in the lungs of CD32A⁻ and CD32A⁺ mice (n=3). Right: Representative lung sections from CD32A⁻ and CD32A⁺ mice immunolabeled with the anti-platelet mAb RAM.1 (Scale bar, 50 μ m). Results are presented as the mean \pm SEM. ns p>0.05, *p<0.05, **p<0.01, ***p<0.001 by 1-way ANOVA (**A,D**), 2-way ANOVA (**C**), or a Student's t-test (**B,E**).

Figure 3: Monocytes/macrophages are required in WT mice but only accessory in CD32A⁺ mice for the initiation of antibody-mediated TRALI

A. The Gr-1 mAb RB6-8C5 or control rlgG2bk (0.5 mg/kg, i.v.) was administered 24 h before LPS (0.1 mg/kg, i.p.) and protein concentrations in BALs were analyzed 15 min after injection of the mAb hlgG1-34-1-2S (1.5 mg/kg, i.v.) or vehicle into CD32A⁺ (n=3-16) or CD32A⁻ (n=3-16) mice. **B.** Clodronate liposomes (2 mL/kg, i.v.), or vehicle, were injected 6 h before hlgG1-34-1-2S challenge (1.5 mg/kg, i.v.) into LPS-sensitized (0.1 mg/kg, i.p.) CD32A⁻ or CD32A⁺ mice. Protein concentrations in BALs were evaluated 15 min later (CD32A⁻: n=7-12, CD32A⁺: n=4-13). Results are presented as the mean \pm SEM. ns p>0.05, *p<0.05, ***p<0.001 by 1-way ANOVA (**A,B**).

Figure 4: Platelets are required for the Fc γ RIIA/CD32A-induced aggravation of antibody-mediated TRALI.

The mAb R300 or control rlgG2a (0.5 mg/kg, i.v.) was administered 24 h before LPS (0.1 mg/kg, i.p.) and protein concentrations in BALs were analyzed 15 min after injection of the mAb hlgG1-34-1-2S (1.5 mg/kg, i.v.) or vehicle into CD32A⁻ or CD32A⁺ mice (n \geq 9). Results are presented as the mean \pm SEM. ns p>0.05, *p<0.05, **p<0.01, ***p<0.001 by 1-way ANOVA.


Figure 5: Platelet-released serotonin, but not platelet-activating factor, is responsible for the Fc γ RIIA/CD32A-induced aggravation of antibody-mediated TRALI.

A. The platelet-activating factor (PAF) receptor antagonist ABT-491 (1.25 mg/kg, i.v.) or saline was administered 5 min before injection of the mAb hlgG1-34-1-2S (1.5 mg/kg, i.v.) or vehicle into LPS-sensitized (0.1 mg/kg, i.p.) CD32A⁻ or CD32A⁺ mice and protein concentrations in BALs were analyzed 15 min later (CD32A⁻: n=3-8, CD32A⁺: n=3-7). Plasma concentrations of platelet factor 4 (PF4) (CD32A⁻: n=3-12, CD32A⁺: n=5-12) (**B**) and serotonin (CD32A⁻: n=3-12, CD32A⁺: n=5-12) (**C**) 15 min after injection of the mAb hlgG1-34-1-2S (1.5 mg/kg, i.v.) or vehicle into LPS-sensitized (0.1 mg/kg, i.p.) CD32A⁻ or CD32A⁺ mice. **D.** Serotonin concentrations in the serum of CD32A⁻ or CD32A⁺ mice treated or not with the SERT inhibitor fluoxetine (160 mg/L) in drinking water for 3 weeks (CD32A⁻: n=11-19, CD32A⁺: n=5-18). **E.** CD32A⁻ or CD32A⁺ mice, treated or not with fluoxetine, were sensitized with LPS (0.1 mg/kg, i.p.) and 24 h later, injected with the mAb hlgG1-34-1-2S (1.5 mg/kg, i.v.) or vehicle. Protein concentrations in BALs were evaluated 15 min afterwards (CD32A⁻: n=3-16, CD32A⁺: n=3-16). **F.** Circulating platelet counts before and 15 min after injection of the mAb hlgG1-34-1-2S (1.5 mg/kg, i.v.) or vehicle into LPS-sensitized (0.1 mg/kg, i.p.) CD32A⁻ or CD32A⁺ mice treated or not with fluoxetine (n \geq 10). **G.** Plasma concentrations of serotonin before and 15 min after injection of the mAb hlgG1-34-1-2S (1.5 mg/kg, i.v.) or vehicle into LPS-sensitized (0.1 mg/kg, i.p.) CD32A⁻ or CD32A⁺ mice treated or not with clodronate (CD32A⁻: n=4-6, CD32A⁺: n=4-6). **H.** Protein concentrations in BALs were evaluated 15 min after injection of the mAb hlgG1-34-1-2S (1.5 mg/kg, i.v.) or vehicle into LPS-sensitized (0.1 mg/kg, i.p.) CD32A⁻ or CD32A⁺ mice treated or not with fluoxetine or clodronate (CD32A⁻: n=5-9, CD32A⁺: n=3-9). **I.** Circulating platelet counts before and 15 min after injection of the mAb hlgG1-34-1-2S (1.5 mg/kg, i.v.) or vehicle into LPS-sensitized (0.1 mg/kg, i.p.) CD32A⁻ or CD32A⁺ mice treated or not with clodronate (CD32A⁻: n=7-12, CD32A⁺: n=4-12). Results are presented as the mean \pm SEM. ns p>0.05, *p<0.05, **p<0.01, ***p<0.001 by 1-way ANOVA (**A-I**).


Figure 6: Blockade of the 5-hydroxytryptamine 2A serotonin receptor prevents the exacerbation of antibody-mediated TRALI in CD32A⁺ mice.

A. The selective 5-HT_{2A} serotonin receptor antagonist sarpogrelate (1 mg/kg, i.v.) or saline was administered 5 min before hlgG1-34-1-2S challenge (1.5 mg/kg, i.v.) in LPS-sensitized (0.1 mg/kg, i.p.) CD32A⁻ or CD32A⁺ mice. Protein concentrations in BALs were evaluated 15 min later (n=6). **B.** Pulmonary elastance (Ers) in LPS-sensitized (0.1 mg/kg, i.p.) CD32A⁻ or CD32A⁺ mice following injection of the mAb hlgG1-34-1-2S (0.75 mg/kg, i.v., n=4) or vehicle (n=3). **C.** Sarpogrelate (1 mg/kg, i.v.) or saline was administered 5 min after hlgG1-34-1-2S challenge (1.5 mg/kg, i.v.) in LPS-sensitized (0.1 mg/kg, i.p.) CD32A⁻ or CD32A⁺ mice. Protein concentrations in BALs were evaluated 15 min later (n≥22). Results are presented as the mean ± SEM. ns p>0.05, **p<0.01 by 1-way ANOVA (**A,C**).


A.



B.



C.



D.




Figure 1A-D

E.






Figure 1E


A.



B.



C.



D.




Figure 2A-D

E.





Figure 2E

A.



B.




Figure 3A,B






Figure 4


A.



B.



C.



D.




Figure 5A-D




Figure 5E-H

I.






Figure 5I

A.



B.



C.




Figure 6A-C