

Research Articles: Development/Plasticity/Repair

Central nervous system hypomyelination disrupts axonal conduction and behaviour in larval zebrafish

<https://doi.org/10.1523/JNEUROSCI.0842-21.2021>

Cite as: J. Neurosci 2021; 10.1523/JNEUROSCI.0842-21.2021

Received: 19 April 2021

Revised: 11 August 2021

Accepted: 16 August 2021

This Early Release article has been peer-reviewed and accepted, but has not been through the composition and copyediting processes. The final version may differ slightly in style or formatting and will contain links to any extended data.

Alerts: Sign up at www.jneurosci.org/alerts to receive customized email alerts when the fully formatted version of this article is published.

Copyright © 2021 Madden et al.

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license, which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.

1 **Central nervous system hypomyelination disrupts axonal conduction and behaviour in larval zebrafish**

2

3 **Abbreviated title:** CNS myelination in zebrafish circuit function

4

5 M.E. Madden¹, D. Suminaite¹, E. Ortiz², J.E. Early¹, S. Koudelka¹, M.R. Livesey^{1,3}, I.H. Bianco⁴, M. Granato²,

6 D.A. Lyons¹

7

8 1. Centre for Discovery Brain Sciences, University of Edinburgh, Chancellor's Building, 49 Little France
9 Crescent, Edinburgh BioQuarter, Edinburgh, EH16 4SB

10 2. Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104

11 3. Sheffield Institute for Translational Neuroscience, Department of Neuroscience, The University of
12 Sheffield, Sheffield, S10 2HQ

13 4. Neuroscience, Physiology and Pharmacology, University College London, Gower Street, London,
14 WC1E 6BT

15 Corresponding author email address: David A Lyons (David.Lyons@ed.ac.uk)

16

17 Figures: 5

18 Abstract: 238 words

19 Introduction: 619 words

20 Discussion: 1294 words

21

22 Acknowledgements: This work was supported by Wellcome Trust Senior Research Fellowships (102836/Z/13/Z
23 and 214244/Z/18/Z) to DAL, a Wellcome Trust Edinburgh Clinical Academic Track PhD studentship
24 (205042/Z/16/Z) to MEM, National Institutes of Health awards (MH109498 and NS118921) to MG, a National
25 Institutes of Health NIDCD award (5T32DC016903) to EO, and a Sir Henry Dale Fellowship from the Royal
26 Society & Wellcome Trust (101195/Z/13/Z) and a UCL Excellence Fellowship to IHB.

27

28

29

30 **Abstract**

31

32 Myelination is essential for central nervous system (CNS) formation, health and function. As a model organism,
33 larval zebrafish have been extensively employed to investigate the molecular and cellular basis of CNS
34 myelination, due to their genetic tractability and suitability for non-invasive live cell imaging. However, it has
35 not been assessed to what extent CNS myelination affects neural circuit function in zebrafish larvae, prohibiting
36 the integration of molecular and cellular analyses of myelination with concomitant network maturation. To test
37 whether larval zebrafish might serve as a suitable platform with which to study the effects of CNS myelination
38 and its dysregulation on circuit function, we generated zebrafish myelin regulatory factor (*myrf*) mutants with
39 CNS-specific hypomyelination and investigated how this affected their axonal conduction properties and
40 behaviour. We found that *myrf* mutant larvae exhibited increased latency to perform startle responses following
41 defined acoustic stimuli. Furthermore, we found that hypomyelinated animals often selected an impaired
42 response to acoustic stimuli, exhibiting a bias towards reorientation behaviour instead of the stimulus-
43 appropriate startle response. To begin to study how myelination affected the underlying circuitry, we established
44 electrophysiological protocols to assess various conduction properties along single axons. We found that the
45 hypomyelinated *myrf* mutants exhibited reduced action potential conduction velocity and an impaired ability to
46 sustain high frequency action potential firing. This study indicates that larval zebrafish can be used to bridge
47 molecular and cellular investigation of CNS myelination with multiscale assessment of neural circuit function.

48

49 **Significance statement**

50 Myelination of central nervous system axons is essential for their health and function, and it is now clear that
51 myelination is a dynamic life-long process subject to modulation by neuronal activity. However, it remains
52 unclear precisely how changes to myelination affects animal behaviour and underlying action potential
53 conduction along axons in intact neural circuits. In recent years, zebrafish have been employed to study cellular
54 and molecular mechanisms of myelination, due to their relatively simple, optically transparent, experimentally
55 tractable vertebrate nervous system. Here we find that changes to myelination alter the behaviour of young
56 zebrafish and action potential conduction along individual axons, providing a platform to integrate molecular,
57 cellular and circuit level analyses of myelination using this model.

58

59

60 **Introduction**

61

62 Myelination is a well-characterised regulator of axonal health and function. In recent years it has become clear
63 that myelination in the central nervous system (CNS) is dynamically regulated over time, including by neuronal
64 activity, leading to the view that activity-regulated myelination might represent a form of functional plasticity
65 (Fields, 2015). Furthermore, disruption to myelin is observed in numerous diseases of the CNS, and its
66 regulation may represent a viable therapeutic strategy. Indeed, major insights have emerged from studies in
67 multiple systems into the cellular and molecular mechanisms of CNS myelination, its regulation by neuronal
68 activity, and its disruption in disease (Almeida, 2018; Gibson et al., 2018; Mount and Monje, 2017; Nave and
69 Werner, 2014). In parallel, an increasing number of studies indicate that the generation of new oligodendrocytes
70 (Geraghty et al., 2019; McKenzie et al., 2014; Pan et al., 2020; Steadman et al., 2020; Wang et al., 2020), and
71 the degree of myelination (Bonnefil et al., 2019; Liu et al., 2016; Makinodan et al., 2012; Sampaio-Baptista
72 et al., 2013), are important for distinct behaviours. However, how dynamic regulation of myelination, or disruption
73 to myelin per se, actually affects the activity of circuits underlying these behaviours remains much less clear.
74 This is partly due to the difficulty in visualising changes to myelination along single axons over time in the
75 mammalian brain while concomitantly assessing their conduction properties and, in turn, evaluating how
76 alteration to conduction affects neural circuit function.

77 Zebrafish are well established as a model organism for the study of myelination. The small size and
78 transparency of their larvae, in combination with their genetic tractability and established transgenic tools,
79 allows the assessment of myelin made by individual oligodendrocytes and along single axons in vivo e.g. (Auer
80 et al., 2018; Koudelka et al., 2016). Together, these features have facilitated innumerable discoveries into the
81 molecular and cellular mechanisms of myelination in this model (Preston and Macklin, 2015). Despite this
82 progress, it remains unknown how CNS myelination affects the function of individual axons, neural circuits, or
83 the behaviour of larval zebrafish, and thus it is not clear whether integrated multiscale assessments of CNS
84 myelination from molecule through circuit can be performed in this model. However, it is now clear that larval
85 zebrafish exhibit a diverse repertoire of experimentally tractable innate and stereotypical locomotor behaviours
86 (Marques et al., 2018), many of which are mediated by reticulospinal (RS) neurons – a diverse set of neurons of
87 the midbrain and hindbrain that process multimodal sensory information, and project descending axons to the
88 spinal cord to coordinate specific motor outputs (Gahtan and O'Malley, 2003; Metcalfe et al., 1986).
89 Intriguingly, RS axons are first to be myelinated in the zebrafish CNS and exhibit activity-regulated myelination

90 at larval stages (Almeida et al., 2011; Hines et al., 2015; Koudelka et al., 2016), implying that regulation of their
91 myelination might influence circuit function in early larvae. In vivo electrophysiological recordings from
92 subsets of individual RS neurons are feasible (Roy and Ali, 2013; Saint-Amant and Drapeau, 2003; Tanimoto et
93 al., 2009), which in principle permits direct measurement of myelinated axon conduction properties underlying
94 behaviour. However, how disruption to CNS myelination affects the behaviour or axonal conduction properties
95 of larval zebrafish remains to be investigated.

96 In this study, we set out to investigate whether changes to CNS myelination can be detected in
97 behaviour and in the conduction properties of single axons in zebrafish larvae. To achieve this, we created a
98 myelin gene regulatory factor (*myrf*) mutant line, which exhibits severe CNS hypomyelination. Using this
99 mutant, we demonstrate that both behavioural and electrophysiological consequences of hypomyelination are
100 indeed detectable in the relevant circuitry in vivo, providing proof of principle that integrated analysis is feasible
101 in this model organism, offering a framework for future investigations.

102

103

104 **Materials and Methods**

105

106 *Zebrafish maintenance*

107 Zebrafish were raised and maintained under standard conditions in the BVS Aquatics Facility in the Queen's
108 Medical Research Institute, University of Edinburgh. Adult and larval animals were maintained on a 14 hours
109 light and 10 hours dark cycle. Embryos were stored in 10mM HEPES-buffered E3 embryo medium or
110 conditioned aquarium water with 0.000001% methylene blue at 28.5°C. All experiments were performed under
111 the project license 70/8436 with approval from the UK Home Office. The *myrf^{me70}* line was maintained in a
112 Tupfel Long Fin (TL) wildtype background. Within this manuscript, 'Tg' denotes a stable, germline inserted
113 transgenic line.

114

115 *Transgenic and mutant lines*

116 The *myrf^{me70}* mutant line was established during this study is described in this manuscript. The following
117 transgenic lines were also used in this study: Tg(mbp:eGFP-CAAX) (Almeida et al., 2011; Mensch et al., 2015),
118 Tg(mbp:nls-eGFP) (Karttunen et al., 2017).

119

120 *Generation of *myrf^{me70}* mutants*

121 A freely available guide selection tool (<http://crispr.mit.edu>) was used to select sgRNA sequences against the
122 second exon of the zebrafish *myrf* gene. sgRNA (target sequence CATTGACACCAGTATCCTGG) was
123 synthesised using DNA template oligomers (5'-
124 AAAGCACCGACTCGGTGCCACTTTTTCAAGTTGATAACGGACTAGCCTATTTTAACTTGCTATTC
125 TAGCTCTAAAACCCAGGATACTGGTGTCAATGCTATAGTGAGTCGTATTACGC-3') (Integrated
126 DNA Technologies, Belgium) consisting of DNA coding for the T7 promotor, DNA recognition sequence
127 (sgRNA variable region) and the sgRNA scaffold. sgRNA synthesis was performed using Ambion
128 MEGAshortscript T7 Transcription Kit (Thermo Fisher Scientific) and the synthesised DNA oligomers as
129 template. Transcribed sgRNA was purified using Ambion MEGAclean kit (Thermo Fisher Scientific). The
130 expression vector for Cas9 protein, pCS2-nCas9n (Addgene plasmid #47929) (Jao et al., 2013), was used to
131 transcribe Cas9 mRNA using the mMMESSAGE mMachine SP6 kit (Thermo fisher Scientific) and purified using
132 an RNeasy mini kit (Qiagen). Injection solutions were prepared with a final concentration of 300ng/μl nCas9n
133 mRNA and 10ng/μl sgrNA in nuclease free water and 0.05% phenol red (Sigma Aldrich). Wildtype embryos

134 were injected at the single or two cell stage with 1.5nL injection solution. Injected F0 animals were raised to
135 adulthood and outcrossed to wildtype animals to create F1 offspring. Clutches of F1 offspring were raised to
136 adulthood and genotyped to identify heterozygous carriers of function disrupting mutant alleles. *myrf^{me70}* refers
137 to a specific allelic mutation consisting of the deletion of two cytosine nucleotides and insertion of a single
138 adenine nucleotide (wildtype sequence: 5'-CCAGTATCCTGGAGGAATA-3'; *myrf^{me70}* mutant allele: 5'-
139 CCAGTATATGGAGGAATA-3').

140

141 *Genotyping*

142 Tissue was genotyped using primers *myrf-f* (5' AACTGTGCGTAGGAACACGATA-3') and *myrf-r* (5'-
143 TGGACCTCCGTGAAACAACACTG-3') in a standard PCR reaction. The PCR product was digested using
144 restriction enzyme PspGI (New England Biolabs), which cleaves wildtype product into 131bp and 157bp
145 fragments. The mutant product remains uncut as the *myrf^{me70}* allele contains a frameshifting indel which
146 abolishes the PspGI cutting site. PCR products were visualised on a 2% gel following gel electrophoresis. All
147 analyses were performed blinded to genotype.

148

149 *Quantitative RT-PCR*

150 Total RNA was extracted from whole brains of adult *myrf^{me70}* wildtype and homozygous siblings using a
151 modified Trizol RNA extraction protocol (TRIzolTM Reagent, Thermo Fisher Scientific). RNA concentration
152 and integrity were assessed using a nanodrop spectrophotometer (NanoDrop One^c, Thermo Fisher Scientific).
153 RNA clean-up was performed if necessary. cDNA synthesis was performed using Accuscript Hi Fidelity First
154 Strand Synthesis kit (Agilent). The amount of RNA entered into the reaction was normalised between samples.
155 Primers *mbp-f* (5'-ACAGAGACCCCACTCTT-3') and *mbp-r* (5'-TCCCAGGCCCAATAGTTCTC-3')
156 were used to amplify *mbp* transcripts within a qPCR reaction (Brilliant III Ultra-fast SYBR Green qPCR Master
157 Mix, Agilent). Transcript levels were detected using Roche Light Cycler 96 (Roche Life Science) with the
158 following amplification protocol: preincubation 95° for 180s, two step amplification 40 cycles: 95° for 10s then
159 60° for 20s, followed by high-resolution melting. Each sample was run in triplicate. Housekeeping gene *ef1a*
160 was used as a reference gene, using primers *ef1a-f* (5'-TGGTACTTCTCAGGCTGACT-3') and *ef1a-r*
161 (5'TGACTCCAACGATCAGCTGT-3'). The delta-delta CT method was used to quantify expression levels. All
162 values were normalised to wildtypes to provide the relative expression of the gene of interest.

163

164

165 *Transmission electron microscopy*

166 Larval tissue was prepared for TEM using the microwave fixation protocol as previously described (Czopka and
167 Lyons, 2011; Karttunen et al., 2017). For adult tissue, adult zebrafish were terminally anaesthetised in tricaine
168 and perfused intracardially with PBS followed by primary fixative solution (4% paraformaldehyde, 2.5%
169 glutaraldehyde, 0.1M sodium cacodylate) (Sigma Aldrich). Adults were subsequently incubated in fresh primary
170 fixative solution for 24 hours at 4°C. Spinal cords were dissected and processed using the microwave fixation
171 protocol described for larval tissue. TEM images were obtained using a JEOL JEM-1400 Plus Electron
172 Microscope. Image magnification ranged from 11.2-17kx magnification for larval spinal cords, and 1.7kx for
173 adult spinal cord.

174

175 *Single cell labelling*

176 Fertilised eggs from *myr^{me70}* heterozygous adult in-crosses were microinjected between the single and four-cell
177 stage with 10ng/μl plasmid DNA encoding mbp:mCherry-CAAX (Mensch et al., 2015) and 25ng/μl tol2
178 transposase mRNA in nuclease free water with 10% phenol red. Animals were screened at 4 dpf for mosaically
179 labelled oligodendrocytes and subsequently imaged. Isolated single cells from any level in the dorsal spinal cord
180 were imaged. Images were obtained in 4 and 6 dpf larvae.

181

182 *Live imaging*

183 Larvae were anaesthetised in tricaine/MS-222 (ethyl3-aminobenzoate methanesulfonate salt, Sigma Aldrich) in
184 HEPES buffered E3 embryo medium and embedded in 1.3-1.5% low melting point agarose (Invitrogen). All
185 fluorescent images were acquired using a Zeiss LSM 880 confocal microscope with a 20x objective (Zeiss Plan-
186 Apochromat 20x dry, NA = 0.8, Carl Zeiss Microscopy). Z-stacks were obtained through the entire single cell,
187 axon or spinal cord according to each experiment. For time course imaging, a single oligodendrocyte was
188 imaged as at 4 dpf. Larvae were then extracted from agarose gel, recovered in embryo medium and maintained
189 with daily feeds and water exchange until imaging of the same cell was repeated at 6 dpf. For automated
190 imaging of the entire spinal cord and peripheral nervous system, Vertebrate Automated Screening Technology
191 (VAST) was utilised as described previously (Early et al., 2018). Briefly, larvae are arrayed into individual
192 wells of a 96-well plate containing MS-222 treated HEPES buffered E3 embryo media. Fish are loaded and
193 oriented for imaging using a Large Particle (LP) Sampler and VAST BioImager system (Union Biometrica Inc)

194 fitted with a 600 μ m capillary tube. Embryos are automatically loaded into the capillary, positioned and imaged
195 using an AxioCam 506m CCD Camera, a CSU-X1 spinning disk confocal scanner, a 527/54+645/60nm double
196 bandpass emission filter, 1.6x C-Mount adapter, a PIFOC P-725.4CD piezo objective scanner, W-Plan-
197 Apochromat 10x 0.5NA objective and an Axio Examiner D1. Z-stacks covering the depth of the capillary were
198 acquired using a 4 μ m z-interval, 3x3 binning and 60ms exposure. Images were acquired using brightfield and
199 the appropriate fluorescent channel. Following imaging, larvae were dispensed into a corresponding well of a
200 96-well collection plate and whole tissue retained for genotyping. Unless otherwise stated, all confocal images
201 presented in this manuscript represent a lateral view of the zebrafish spinal cord, with anterior to the left and
202 posterior to the right, and dorsal/ventral at the top/bottom of the image respectively. Within experiments, images
203 were obtained using similar laser intensity and optical gain settings. All imaging was performed blinded to
204 genotype.

205

206 *Transmission electron microscopy*

207 TEM images were tiled using the automated photo merge tool in Adobe Photoshop 2019. The number of
208 ensheathed axons was counted in one hemi-spinal cord section per larva using the cell counter tool in FIJI
209 ImageJ. Axon caliber is defined as the area of the axon within this manuscript. Axonal area was calculated using
210 the freehand line and measure tool in FIJI ImageJ. A g-ratio represents the ratio between the inner and outer
211 diameter of the myelin sheath (i.e. a larger g-ratio values denotes a thinner myelin sheath). This calculation
212 assumed perfect circularity of axons, which is not true to larval zebrafish TEM preparations. Thus, within these
213 experiments g-ratio was calculated by dividing axonal area by the axonal + associated myelin area.

214

215 *VAST*

216 Images obtained using VAST were stitched and processed using FIJI ImageJ software (Schindelin et al., 2012)
217 and custom macros (Early et al., 2018). Semi-automated oligodendrocyte counts were performed on the
218 maximum intensity projection images (Early et al., 2018). Cell count values represent all oligodendrocytes in
219 the spinal cord (dorsal and ventral tracts). Morphometric analysis of larval developmental features was
220 performed on brightfield images. Measurements of ocular diameter, body length and swim bladder height were
221 performed using the line and measure tool in FIJI ImageJ (National Institutes of Health).

222

223 *Single cell imaging*

224 Confocal z-stack images were airyscan processed using Zen software (ZEISS). Images were opened in FIJI
225 ImageJ. Cells were included for analysis only if all myelin sheaths were distinguishable. Myelin sheath lengths
226 were measured using the freehand line and measure tools. Myelin sheath number was equivalent to the number
227 of measurements performed. Total myelin per cell was calculated as sum of all myelin sheaths lengths per cell.
228 Abnormal sheaths were defined at sheaths with abnormal elongation profiles, incomplete wrapping or myelin
229 blebbing. For time course experiments, net growth or shrinkage of myelin sheaths was calculated as the average
230 myelin sheath length at 6 dpf minus the average myelin sheath length at 4 dpf. Where possible, all myelin
231 sheaths per cell were measured. In instances where measurement of all myelin sheaths was not possible (due to
232 other cells coming into close proximity), only isolated myelin sheaths were analysed at each time point. The
233 number of retracted sheaths was recorded, and these sheaths were excluded from sheath growth analysis.

234

235 *Electrophysiology*

236 Zebrafish were dissected as described previously in Roy & Ali (2013) to access the Mauthner neuron. In short, 6
237 dpf anaesthetised zebrafish were laid on their sides on a Sylgard dish and secured using tungsten pins through
238 their notochords in a dissection solution containing the following (in mM): 134 NaCl, 2.9 KCl, 2.1 CaCl₂, 1.2
239 MgCl₂, 10 HEPES, 10 glucose and 160mg/ml tricaine, adjusted to pH 7.8 with NaOH. Their eyes as well as
240 lower and upper jaws were removed using forceps to expose the ventral surface of the hindbrain, which was
241 secured with an additional tungsten pin. The motor neurons in the anterior spinal cord were exposed as
242 described by Wen et al (2005). A dissecting tungsten pin was used to remove the skin and the muscle overlaying
243 the motor neurons in a single segment. Following the dissection, zebrafish together with their recording
244 chamber were moved to the rig and washed with extracellular solution containing the following (in mM): 134
245 NaCl, 2.9 KCl, 2.1 CaCl₂, 1.2 MgCl₂, 10 HEPES and 10 glucose with 15 μ M tubocurarine. The cells were
246 visualised using Olympus microscope capable of DIC using 60X water immersion NA = 1 objective lens and
247 Rolera Bolt Scientific camera with Q-Capture Pro 7 software. The stimulating electrode filled with extracellular
248 solution was then positioned in the mid spinal cord lightly touching the exposed neurons underneath. Mauthner
249 whole-cell recordings were performed with thick-walled borosilicate glass pipettes pulled to 6–10 M Ω . The
250 internal solution contained the following (in mM): 25 K-gluconate, 15 KCl, 10 HEPES, 5 EGTA, 2 MgCl₂, 0.4
251 NaGTP, 2 Na₂ATP, and 10 Na-phosphocreatine, adjusted to pH 7.4 with KOH. Upon formation of whole-cell
252 patch clamp, 270s – long recording was performed in the current - clamp configuration. Cell resting membrane
253 potential was established as an average of the first 5 seconds of the recording if the cell did not fire during that

254 time. To measure the conduction velocity along the Mauthner axon, the zebrafish were washed with recording
255 solution containing the following (in mM): 134 NaCl, 2.9 KCl, 2.1 CaCl₂, 1.2 MgCl₂, 10 HEPES and 10 glucose
256 with addition of (in μM) 50 AP5, 20 strychnine, 100 picrotoxin and 50 CNQX. The antidromic Mauthner action
257 potentials were recorded following the field stimulation by the stimulating electrode connected to DS2A
258 Isolated Voltage Stimulator (Digitimer) in the spinal cord. 30 consecutive action potentials were recorded every
259 5 seconds using Clampex 10.7 at 100kHz sampling rate and filtered at 2 kHz using MultiClamp 700B. At the
260 end of the recording, images of the zebrafish were obtained with 4X objective and stitched using Adobe
261 Photoshop. The resulting image was then transferred to FIJI and the distance between stimulating and the
262 recording electrode was measured. The conduction velocity of action potential was calculated dividing the
263 distance between the electrodes by the latency from the stimulus artefact to the peak of action potential. Action
264 potential latency and half-width were measured using homebuilt MATLAB script. For the analysis of action
265 potential fidelity consecutive trains of 10 stimuli were delivered at 1, 10, 100, 300, 500 and 1000Hz every 30s.
266 Recordings were made at 20kHz sampling rate and filtered at 2 kHz. The number of action potentials were
267 calculated using Clampfit 10.7 software and the action potential success rate was calculated as a number of
268 action potentials fired out of 10 and multiplied by 100.

269

270 *Behavioural assay*

271 Analysis of startle behaviour in 5-6 dpf *myrf^{me70}* mutant and wildtype larvae was performed as previously
272 described (Burgess and Granato, 2007a, Wolman et al, 2011). Briefly, larvae were placed into individual wells
273 of a 6 x 6 custom made acrylic testing plate containing E3 embryo media. A series of 10 acoustic stimuli (40.6
274 dB, 1000 Hz, 3ms duration) were delivered to the plate with an interstimulus interval of 20 seconds. Behaviour
275 was recorded using a high-speed camera (Photron Fastcam Mini UX) at 1000frames/s. Analysis of recorded
276 video footage was performed using FLOTE v2.0 tracking software (Burgess and Granato 2007a). Larvae that
277 responded to less than 70% of the stimuli were excluded from further kinematic analysis. Average behavioural
278 latency was calculated as an average per larva over all behavioural responses. Short latency C-starts (SLC) and
279 long latency C-starts (LLC) were defined by identifying a latency value (16ms) separating the two peaks of the
280 latency bimodal distribution in wildtype *myrf^{me70}* larvae. Behavioural latency, c-bend duration, initial turn angle,
281 and angular velocity for SLC and LLC events were defined and analysed as previously described (Burgess and
282 Granato, 2007a).

283

284 *Experimental design and statistical analysis*

285 Unless stated otherwise, all experiments were performed on 6 dpf larvae from adult heterozygous in-crosses. All
286 subjects were the offspring of third generation, or younger, adults. The experimenter was blinded to the
287 genotype of the larvae during all experiments and analysis. The sex of the animals was unknown as sex
288 specification has not occurred at this stage of larval development. All graphs and statistical testing were
289 performed using GraphPad Prism. All data was assessed for Gaussian distribution using a D'agostino Pearson
290 omnibus normality test. Parametric continuous data was analysed using a two-tailed unpaired student's t test, or
291 two-way ANOVA, according to the number of variables being compared. Non-parametric continuous data was
292 analysed using a Mann-Whitney test. If the number of values were too small to assess for normality, it was
293 assumed that data was non-parametric. Results were considered statistically significant when $p < 0.05$. Within
294 figures, p values are denoted as follows: non-significant i.e. $p > 0.05$ 'ns', $p < 0.05$ '*', $p < 0.01$ '***', $p < 0.001$
295 '****', $p < 0.0001$ '*****'. Unless otherwise stated, all data was averaged per biological replicate (N represents
296 number of larvae). Throughout the figures, error bars represent mean \pm standard deviation for parametric data, or
297 median and interquartile range for non-parametric data. Details of statistical tests, precise p and n values for
298 each experiment are provided in the appropriate figure legends.

299

300 *Code accessibility*

301 Custom written code to perform automated cell counts is available in a previous publication (Early et al., 2018).

302 Code to interpret electrophysiological data is available at https://github.com/skotuke/Mauthner_analysis.

303

304 **Results**

305

306 **Targeting myelin gene regulatory factor to create a larval zebrafish model of CNS-specific**
307 **hypomyelination**

308 To begin our investigations into the role of CNS myelination in neural circuit function, we sought to establish a
309 larval zebrafish model with CNS-specific hypomyelination. Mammalian studies have identified myelin gene
310 regulatory factor (*myrf*) as a transcription factor vital for CNS myelin formation and maintenance (Bujalka et
311 al., 2013; Emery et al., 2009). Zebrafish possess a single ortholog of *myrf* and, similar to mammals, *myrf*
312 expression in the CNS appears to be restricted to oligodendrocytes (Klingseisen et al., 2019; Treichel and Hines,
313 2018). We used CRISPR/Cas9 technology to target a guide RNA to exon 2 of the zebrafish *myrf* gene, the first
314 conserved exon across all predicted splice variant isoforms, and in doing so created the *myrf*^{me70} mutant
315 (Methods and **Figure 1A**). Morphometric analysis of larval body features of *myrf*^{me70} mutants at larval stages
316 showed them to be indistinguishable from siblings (data not shown), and in contrast to mammalian *myrf*
317 mutants (Emery et al., 2009), homozygous *myrf*^{me70} mutants remain viable through to adulthood. Adult *myrf*^{me70}
318 mutants exhibited an almost complete absence of *mbp* mRNA (**Figure 1B**), and transmission electron
319 microscopy (TEM) assessment indicated effectively no myelin in the adult spinal cord (**Figure 1C and D**). In
320 addition, and unlike larvae, homozygous adult *myrf*^{me70} were grossly identifiable from their siblings by their
321 smaller size. Adult *myrf*^{me70} mutants were also infertile, due to the absence of detectable gonadal tissue in
322 females, confirmed via histopathology, which also revealed evidence of cardiomyopathy (data not shown) -
323 findings consistent with proposed roles of *myrf* outside the CNS (Hamanaka et al., 2019; Pinz et al., 2018;
324 Rossetti et al., 2019).

325 Given the potential to study myelination of well-defined circuits at high resolution over time at larval
326 stages when *myrf*^{me70} mutants are morphologically indistinguishable from siblings, we next analysed our
327 transgenic reporter of myelination Tg(*mbp:eGFP-CAAX*) at 6 days post fertilisation (dpf). This indicated that
328 the gross level of CNS myelination was also reduced in *myrf*^{me70} mutant larvae relative to wildtype siblings
329 (**Figure 1E**). To quantify myelination in larvae, TEM was performed on transverse sections of the spinal cord
330 (CNS) and posterior lateral line nerve of the peripheral nervous system (PNS) at 6 dpf (**Figure 2A-C**). At this
331 timepoint, we observed a 66% reduction in the number of myelinated axons in the spinal cord of *myrf*^{me70}
332 mutants relative to wildtype siblings (35.29 ± 7.83 myelinated axons in wildtypes, 12.00 ± 4.34 myelinated
333 axons in mutants, $p \leq 0.0001$, unpaired t-test) (**Figure 2D**). In contrast, and demonstrating specificity of

334 hypomyelination to the CNS, similar numbers of myelinated axons were observed in the PNS of mutant and
335 wildtype siblings (7.33 ± 1.53 myelinated axons in wildtypes, 9.00 ± 3.83 myelinated axons in mutants, $p =$
336 0.52 , unpaired t-test) (**Figure 2E**).

337 Despite the large number of unmyelinated axons in *myrf^{me70}* mutants, our TEM analyses indicated that
338 some axons remained ensheathed in the larval CNS, including the very large diameter Mauthner axons, the first
339 reticulospinal axons to be myelinated in the zebrafish CNS (Almeida et al., 2011). Although Mauthner axons
340 were ensheathed in *myrf^{me70}* mutants at 6 dpf, they had significantly thinner myelin sheaths compared to
341 wildtype siblings (average g-ratio: 0.48 ± 0.009 in wildtypes, 0.80 ± 0.08 in homozygous mutants, $p = 0.0009$,
342 unpaired t-test, **Figure 2F**). A similar finding was observed in the other axons that were ensheathed in *myrf^{me70}*
343 mutants at this stage, with greater g-ratio values (denoting thinner myelin) for other large caliber ($>0.3\mu\text{m}^2$)
344 axons in mutants than in wildtype siblings (average g-ratio: 0.60 ± 0.08 wildtypes, 0.71 ± 0.08 mutants, p
345 ≤ 0.0001 , unpaired t test, **Figure 2G**). Despite the generally severe hypomyelination phenotype, the presence of
346 some large caliber myelinated axons in zebrafish *myrf^{me70}* mutants at larval stages contrasts with our analysis of
347 adult zebrafish mutants and *myrf* mutant mice which both have essentially a complete absence of CNS
348 myelination (Emery et al., 2009). This suggests that the full effects of *myrf* knockout may be masked at early
349 stages, either by maternal gene expression or genetic compensatory mechanisms (Rossi et al., 2015).

350 To examine the cellular basis of CNS hypomyelination in *myrf^{me70}* mutant larvae, we first assessed
351 myelinating oligodendrocyte number using the transgenic reporter Tg(*mbp:nls-eGFP*) (Karttunen et al., 2017)
352 (**Figure 3A**). At 6 dpf, the timepoint at which TEM was performed, the number of detectable oligodendrocytes
353 was reduced by 21% in *myrf^{me70}* mutants relative to wildtype siblings ($p = 0.0002$, unpaired t-test, **Figure 3B**). In
354 addition, the fluorescent intensity of *myrf^{me70}* mutant oligodendrocyte nuclei was reduced, consistent with
355 reduced *mbp* expression. Because, the reduction in cell number was not sufficient to explain the reduction in
356 myelin observed using TEM, we assessed the morphology of individual myelinating oligodendrocytes using
357 mosaic cell labelling with the *mbp:mCherry-CAAX* reporter construct (Almeida et al., 2011) (**Figure 3C**). We
358 found that both myelin sheath number ($p = 0.02$, Mann Whitney test, **Figure 3D**) and length ($p = 0.002$,
359 unpaired t test, **Figure 3E**) were reduced in *myrf^{me70}* mutants by 33% and 25% respectively at 6 dpf, with total
360 myelin (sum of sheath lengths) per individual oligodendrocyte reduced by 47% in mutants relative to wildtypes
361 ($p \leq 0.0001$, unpaired t test, **Figure 3F**). In addition to being required for the initiation of myelination, previous
362 studies in rodents indicate that *myrf* is also essential for myelin sheath maintenance (Koenning et al., 2012).
363 Having observed that adult *myrf^{me70}* mutants have a much more severe hypomyelination phenotype than larvae

364 (Figure 1D), we wanted to assess how the morphology of single oligodendrocytes changed over time. To do so,
365 we imaged single oligodendrocytes at 4 dpf and again at 6 dpf (Figure 3G). We found that between these
366 timepoints mutant oligodendrocytes demonstrated a net shrinkage in myelin sheath length, while wildtype
367 oligodendrocytes showed a net growth ($p = 0.009$, Mann Whitney test, Figure 3H). Furthermore, the number of
368 myelin sheaths that were completely retracted during this timeframe was significantly higher in *myrf^{ae70}* mutant
369 oligodendrocytes ($p = 0.003$, unpaired t test, Figure 3I). Also consistent with a failure to maintain healthy
370 myelin sheaths, the number of myelin sheaths exhibiting an abnormal morphology (i.e. incomplete wrapping,
371 abnormal elongation profiles or myelin blebs) was significantly higher in mutant versus wildtype
372 oligodendrocytes at 6 dpf (Figure 3J).

373 In summary, disrupting *myrf* leads to a CNS-specific hypomyelination phenotype in larval zebrafish,
374 caused by a reduction in the number of oligodendrocytes, with those that remain having fewer and shorter
375 sheaths. The majority of sheaths that are made are thinner, and, based on our documentation of almost complete
376 absence of myelin in adults, not maintained long-term. Therefore, the phenotype in the *myrf^{ae70}* mutant fulfilled
377 our aim to generate a CNS-specific model of hypomyelination to study the effects on neural circuit function at
378 larval stages.

379

380 *myrf^{ae70}* mutants exhibit an increase in the latency to perform startle responses and an impaired 381 behavioural choice in response to a defined auditory stimulus

382 Given that many larval zebrafish sensorimotor behaviours are mediated by RS neurons, whose axons are
383 myelinated early and exhibit activity-regulated myelination (Koudelka et al., 2016), we hypothesised that
384 *myrf^{ae70}* mutants would display detectable differences in the performance of RS-mediated behaviours. To test
385 this, we chose to first examine acoustic-startle behaviour, for which the underlying circuit is relatively well
386 described (Hale et al., 2016). Briefly, a high-intensity acoustic stimulus activates the auditory (VIIIth) nerve,
387 which courses into the hindbrain to synapse onto the Mauthner cell at its lateral dendrite. Once the threshold
388 potential is exceeded, an action potential is elicited and rapidly propagated along the Mauthner axon, which
389 crosses into, and extends along, the contralateral tract of the spinal cord. Along its length, collateral branches
390 make synapses with interneurons and primary motor neurons that coordinate motor output. Activation of a
391 Mauthner axon results in a stereotypical, high-velocity ‘c-bend’ away from the stimulus, followed by a fast burst
392 swim (Kimmel et al., 1974) (Figure 4A). The latency to perform such a response is defined as the time taken
393 from stimulus presentation to the onset of a c-bend (Figure 4J). Given that myelin increases conduction

394 velocity along a single axon (Waxman, 1980), we made the prediction that the latency to execute the motor
395 responses following an acoustic stimulus would be delayed in *myrf^{ae70}* mutants.

396 Motor behaviour was assessed using an established high-throughput assay (Burgess and Granato,
397 2007a). *myrf^{ae70}* larvae were arrayed into individual wells of a 6x6 custom made plate attached to an amplifier
398 delivering a series of acoustic stimuli at 20 second intervals (**Figure 4B**). Using a high-speed (1000Hz) camera,
399 behavioural responses were recorded and subsequently analysed using FLOTE software (Burgess and Granato,
400 2007a, 2007b). Overall, the frequency of responses to acoustic stimuli was similar between groups (**Figure 4E**).
401 However, on average, *myrf^{ae70}* mutants exhibited a 66% increase in their average latency to elicit a response
402 compared to wildtype siblings (wildtypes: 10.55ms (9.6-16.16ms), mutants: 17.60ms (12.90-21.88ms), $p =$
403 0.003, Mann Whitney Test, **Figure 4F**).

404 Interestingly, larval behavioural responses to acoustic stimuli can be modulated across variable
405 stimulus properties, exhibiting decision-making capabilities of the underlying circuitry (Burgess and Granato,
406 2007a; Jain et al., 2018). For example, in larval zebrafish, while high intensity threatening stimuli induce the
407 short-latency c-bend startle response, also known as the ‘short latency c-start’ (SLC), lower stimulus intensities
408 induce a distinct longer latency reorientation-like behaviour, initially defined as a ‘long latency c-start’ (LLC).
409 These kinematically and behaviourally distinct responses are executed by activity in partially overlapping
410 circuitry, with the crucial difference that SLCs are driven by recruitment of Mauthner neurons, while LLCs
411 appear to be driven by alternative pathways e.g. preoptine neurons (Burgess and Granato, 2007a; Marquart et
412 al., 2019) (**Figure 4A**). Given that hypomyelination in *myrf^{ae70}* is widespread within the CNS, we anticipated
413 that the large overall increase in latency to respond to acoustic stimuli might be due to significant delays in the
414 performance of both SLC and LLC responses. However, when data was segmented into SLCs or LLCs, the
415 latency to perform an SLC was increased by 6.4% (10.03 ± 0.85 ms in wildtypes, 10.67 ± 0.83 ms in mutants, p
416 $= 0.006$, unpaired t test **Figure 4G and I**), but the latency to perform LLCs remained unaffected (**Figure 4H**
417 **and I**), begging the question as to what caused the much larger overall increase in latencies to respond to
418 acoustic stimuli.

419 We reasoned that if the latency to perform SLCs was only affected to a small degree and LLCs not at
420 all, the overall large increase in latency to perform all responses might be due to a biased selection of the longer
421 latency LLCs over the much shorter latency SLCs. Indeed, when we compared their relative frequency, we saw
422 that LLCs represented a significantly increased proportion of behavioural responses in *myrf^{ae70}* mutants relative
423 to wildtypes (SLC:LLC ratio: 10:1 in wildtypes, 2.9:1 in mutants, $p \leq 0.0001$, Kolmogorov-Smirnov test,

424 **Figure 4C and D**). To ensure that this apparent bias in behavioural selection was not due to SLCs simply being
425 so slow as to be detected as LLCs, we analysed additional kinematic parameters (**Figure 4J-N**), which have
426 specific values associated with each type of response (Burgess and Granato, 2007a). No differences were found
427 in the duration, maximum angular velocity or initial turning angle of SLCs or LLCs between wildtype and
428 mutant larvae (**Figure 4K-M**), consistent with the conclusion that the increased frequency of LLCs represents
429 true LLC events, rather than delayed and inappropriately classified SLCs.

430 In summary, we have shown that *myrf^{me70}* mutants exhibited delayed latency to perform Mauthner-
431 mediated startle responses (SLCs), and an unexpected bias towards performing Mauthner-independent
432 reorientation behaviours (LLCs) in response to the same acoustic stimuli. This shows that hypomyelination in
433 the larval zebrafish can be detected in overt changes to behaviour and highlights the complexity of how
434 dysregulation of myelination impacts circuit function, even when executing relatively simple sensorimotor
435 transformations.

436

437 **Action potential conduction is impaired along the Mauthner axon in *myrf^{me70}* mutants**

438 In order to investigate how myelination affects conduction along larval zebrafish axons, we set out to establish
439 an electrophysiological platform that would allow us to measure and compare multiple aspects of axonal
440 conduction in vivo. We focussed our analysis on the Mauthner neuron and axon, due to its characteristic
441 morphology and anatomical location, and given its established role in mediating the SLC. To begin with, we
442 performed whole-cell current-clamp recordings of the Mauthner neuron cell body while stimulating its axon in
443 the spinal cord with an extracellular electrode (**Figure 5A**). We first tested whether loss of *myrf* function
444 affected intrinsic properties of the Mauthner neuron, by assessing its resting membrane potential: we found that
445 this remained stable in mutants (siblings: $-70.82 \pm 2.76\text{mV}$, mutants: $-70.68 \pm 1.25\text{mV}$, $p = 0.9077$, unpaired t-
446 test, **Figure 5B**). Our experimental configuration allowed us to record antidromic action potentials propagating
447 along the Mauthner axon. Therefore, we next assessed whether the shape of action potentials was disrupted by
448 hypomyelination, by measuring the width of the action potential at its half-height (action potential half-width) at
449 6 dpf, which we found to be similar in control and *myrf^{me70}* mutant animals (siblings: $0.64 \pm 0.09\text{ms}$, *myrf^{me70}*
450 mutants: $0.60 \pm 0.06\text{ms}$, $p = 0.2610$, unpaired t-test, **Figure 5C and D**). These data indicate that the degree of
451 hypomyelination along Mauthner axons in *myrf^{me70}* mutants at these stages does not affect the Mauthner resting
452 membrane potential or greatly affect the shape of the action potentials.

453 Given the well-defined role for myelin in speeding-up action potential conduction, and the evidence of
454 an increased latency to perform the Mauthner-dependent SLC response, we next measured the latency of action
455 potential conduction along the Mauthner axon in controls and *myrf^{me70}* mutants. This analysis showed that the
456 normalised latency of action potentials was significantly increased in *myrf^{me70}* mutants when compared to
457 siblings (siblings: 0.80 ± 0.11 ms/mm, mutants: 0.97 ± 0.07 ms/mm, $p = 0.0003$, **Figure 5F**) resulting in an 18%
458 reduction in conduction velocity (siblings: 1.27 ± 0.17 m/s, mutants: 1.04 ± 0.08 m/s, $p = 0.0005$, **Figure 5G**).
459 This reduction in conduction velocity supports our finding of a delayed execution of SLCs in *myrf^{me70}* mutants.
460 We next assessed whether the precision of action potential propagation might be impaired due to
461 hypomyelination, which might interfere with synaptic signalling in the circuit. To do so, we measured the
462 ‘jitter’, or imprecision, in the timing of action potential arrival following stimulation, as the standard deviation
463 of 30 action potential peak times aligned to the stimulus artefact (**Figure 5H**). No differences were observed in
464 the precision of action potential arrival in *myrf^{me70}* mutants at 6 dpf (siblings: 0.006 ± 0.002 ms, mutants: $0.006 \pm$
465 0.0009 ms, $p = 0.8166$, unpaired t-test, **Figure 5I**). These data suggest that hypomyelination leads to slower, but
466 nonetheless precise, action potential propagation.

467 Given that the action potentials conducted along Mauthner axons in *myrf* mutants are likely to be
468 sufficient to trigger downstream motor output, albeit with a longer delay, we next asked whether the
469 hypomyelination of Mauthner axon might lead to an increased failure to reliably propagate action potentials.
470 Therefore, we implemented a strategy to robustly test the ability of the myelinated axon to faithfully transmit
471 action potentials. With our preparation, we observed that the Mauthner cell could spontaneously fire short trains
472 of action potentials (1-10) at high frequency (~300Hz) while in the resting state, prior to our adding
473 pharmacological reagents to block network-level input on to Mauthner and ahead of taking control of
474 stimulating activity in the preparation, for the analyses noted above (data not shown). On the basis of this
475 observation, and given the evidence from studies in rats that dysmyelination can influence firing frequency
476 (Kim et al., 2013) we established a high-frequency stimulation paradigm to assess how hypomyelination
477 affected the ability of the Mauthner axon to sustain high frequency firing of action potentials. To do so, we used
478 our field stimulation procedure and delivered 10 stimuli at various frequencies via the stimulating electrode and
479 recorded the number of action potentials fired by the Mauthner cell, which allowed us to assess action potential
480 success rate (**Figure 5J**). Given that myelination reduces axonal current leakage, we predicted that our high
481 frequency stimulation protocol may reveal failed action potential propagation. When we analysed the success
482 rate of action potential firing, we found that this was indistinguishable between siblings and mutants at 300Hz,

483 insignificantly different at 500Hz, but significantly impaired at 1000Hz stimulation, where we found that
484 Mauthner cells from mutants fired with a significantly lower success rate (siblings: $55.79 \pm 10.17\%$, mutants:
485 $38.89 \pm 17.64\%$, $p = 0.0014$, two-way ANOVA, **Figure 5K**). This assay suggests that hypomyelination impairs
486 the ability of axons to propagate action potentials faithfully, which could contribute to the behavioural shift
487 away from Mauthner-mediated responses to auditory stimuli.

488 In conclusion, we have established an electrophysiology platform that allows direct measurement of
489 single cell (i.e. Mauthner) conduction properties in vivo. In doing so, we have demonstrated that
490 hypomyelination of the Mauthner axon leads to slowed conduction velocity, and with a high frequency
491 stimulation paradigm we reveal a loss of fidelity of action potential propagation along the hypomyelinated
492 Mauthner axon.

493

494

495 **Discussion**

496 We have demonstrated that CNS hypomyelination leads to behavioural alterations and impaired conduction
497 along axons in larval zebrafish. We found that *myrf^{me70}* mutant zebrafish larvae exhibit CNS-specific
498 hypomyelination, representing the first model with which one can study the role of CNS myelin in behaviour.
499 These mutants exhibited an increased latency to execute the stereotypical rapid acoustic startle responses (SLCs)
500 and were also biased towards performing longer latency reorientation behaviours (LLCs) in response to startle-
501 inducing acoustic stimuli. The fact that our analysis revealed phenotypes in both the speed of executing a
502 specific behaviour and in the selection of the correct behavioural response to a sensory stimulus indicates the
503 complex roles that myelination plays in regulating circuit function. These findings provide encouragement that
504 studying additional behaviours will offer further entry-points into studying how alterations to myelination affect
505 the function of other neural circuits. Indeed, there are now a large number of behavioural paradigms that allow
506 analysis of larval zebrafish circuit function, from various sensorimotor transformations (Dunn et al., 2016;
507 Henriques et al., 2019; Naumann et al., 2016), behaviours regulated by sensory experience over time (Burgess
508 and Granato, 2007a; Wolman et al., 2011) and those driven by inter-individual interactions, such as sociability
509 (Dreosti et al., 2015; Larsch and Baier, 2018).

510 In addition to studying behaviour, we established electrophysiological protocols to assess the
511 conduction properties of single neurons and axons, focusing on the Mauthner neuron due to the ease of its
512 identification and its involvement in the acoustic startle response. We found that conduction along the

513 hypomyelinated Mauthner axon was reduced, and that Mauthner axons in *myrf^{ue70}* mutants exhibited an
514 increased failure to propagate action potentials in response to high-frequency stimulation. It remains to be
515 determined precisely how disruption to the conduction properties of neurons and axons caused by
516 hypomyelination affects circuit function and behavioural outputs. For example, the slowed execution of the SLC
517 may be due to more than the slower conduction along the hypomyelinated Mauthner axon of *myrf* mutants,
518 including slower conduction elsewhere in the circuit. Precisely how hypomyelination leads to a biased
519 recruitment of LLCs over SLCs in response to the same auditory cue in *myrf^{ue70}* mutants also remains to be
520 elucidated, but could be influenced by the impaired ability to sustain high frequency firing along the
521 hypomyelinated Mauthner axons, and dysregulated recruitment of downstream motor pools. However, with our
522 antidromic preparation, we cannot rule out the possibility that the reduced success rate of high frequency action
523 potential conduction was influenced by impaired generation of action potentials in the axon. Therefore,
524 establishing methods to record orthodromic action potentials remains an important challenge for the future. In
525 addition, to study how dysregulation of myelination influences synaptic signalling, electrophysiological
526 analyses through the paired recordings of neurons known to communicate within circuits will be required. These
527 studies, alongside the ability to assess the conduction properties of additional neurons, will be required to
528 generate complete circuit models of how myelin influences even simple behaviours. Our study documented
529 behavioural alteration and disruption to conduction in larvae with CNS-specific hypomyelination, but many
530 challenges remain in integrating our understanding of circuit function across scales from conduction and
531 synaptic communication through population-level neuronal activity and the execution of specific behaviours.
532 However, we believe that the zebrafish represents a model in which such a multi-scale analyses of myelination
533 on neural circuit function is feasible.

534

535 The larval zebrafish has numerous advantages that facilitate analyses of circuit function across scales.
536 The larval CNS is relatively simple compared to mammalian models; with approximately one hundred thousand
537 neurons by 6 dpf, only a relatively small proportion (on the order of a few hundred neurons) have myelinated
538 axons at this stage (Hildebrand et al., 2017). The myelination of those axons is generally very stereotyped, with
539 myelination of certain neuronal subtypes (e.g. reticulospinal neurons) adaptable and responsive to neuronal
540 activity (Koudelka et al., 2016). With the aim of studying myelination from the perspective of neural circuits,
541 we previously developed tools to study patterns of myelination along single axons in vivo (Koudelka et al.,
542 2016). These tools, together with increasing availability of neuron-specific drivers coupled with circuit maps of

543 the larval fish brain provide a great opportunity to map myelination at single cell resolution across the larval
544 zebrafish CNS, and to do so over time. Even with myelination patterns mapped, a corresponding challenge will
545 be to manipulate myelin from the point of view of specific neurons/axons and circuits. As noted above, it
546 remains unclear whether the longer latency to execute the startle response is simply due to hypomyelination of
547 Mauthner axons, or elsewhere in the circuit, and it may even be influenced by complex integrative functions that
548 affect timing across the circuit. Therefore, it will be important to develop methods to regulate myelination in a
549 neuron/axon and circuit-specific manner. One possibility might be to selectively ablate oligodendrocytes in
550 specific circuits. Although oligodendrocyte ablation can be carried out at single cell resolution in zebrafish
551 larvae (Auer et al., 2018), it leads to inflammatory reactions by cells such as microglia (Karttunen et al., 2017),
552 which may be relevant to disease contexts, but would confound the disentangling of the role of myelin per se in
553 healthy circuits. Therefore, an additional approach might be to express cell surface proteins that inhibit
554 myelination (Redmond et al., 2016) along the axons of specific neuronal cell types (Burgess et al., 2009; Tabor
555 et al., 2018; Yamanaka et al., 2013), selectively preventing their myelination. Furthermore, as signals and
556 receptors that influence adaptive activity-regulated myelination are identified, yet more strategies to influence
557 myelination in localised manners may emerge.

558 In addition to needing more refined methods to map and manipulate myelination of specific circuits,
559 additional tools to assess function across scales from single axon to behaving animal will be required. Given the
560 challenges of integrating complex electrophysiological protocols with behavioural observation in small
561 zebrafish larvae, it is possible that optical methods to assess function across scales provides a better opportunity
562 to bridge analyses across scales. Indeed, optical imaging approaches have already proven hugely powerful in the
563 study of larval zebrafish brain function. For example, two-photon and light-sheet microscopy-based imaging
564 studies allow the analysis of the activity of individual neurons (Abdelfattah et al., 2019) through to sampling the
565 activity of effectively all neurons the entire larval zebrafish brain, at multiple volumes per second with sub-
566 cellular resolution (Ahrens et al., 2013, 2012; Chen et al., 2018). In fact, sophisticated imaging platforms that
567 allow monitoring of neuronal activity in the brain during the execution of behaviours have been developed,
568 including during acoustic stimulus-driven responses (Jain et al., 2018; Lacoste et al., 2015). Furthermore, the
569 coordinated activity of ensembles of neurons have been investigated in the larval brain, which provides an
570 opportunity to investigate how potentially even subtle alterations to myelination in development, health or
571 disease might influence relatively high-order network activity (Diana et al., 2018; Romano et al., 2015; Sumbre
572 et al., 2008; Wolf et al., 2017). To date, most optical analyses of neuronal activity in zebrafish have been carried

573 out using genetically encoded Ca^{2+} reporters, but the limited temporal kinetics of even the fastest Ca^{2+} reporters
574 may preclude the analysis of millisecond-scale changes to conduction properties, which our data indicate can be
575 expected with disruption to larval myelination. However, ongoing development and refinement of voltage
576 indicators appear to exhibit photodynamic properties with the sensitivity to detect functional changes to
577 conduction and synaptic properties at the appropriate temporal resolution, including in larval zebrafish
578 (Abdelfattah et al., 2019). Employing indicators that allow bona fide assessment of conduction in the intact
579 brain, during the execution of behaviours has the potential to provide a transformative capacity to interrogate
580 how myelin influences circuit function.

581 In summary, our study presents larval zebrafish as a viable model to study myelination across scales
582 from molecular and cellular analyses of how myelin organises and supports axons through to functional
583 assessments of conduction, synaptic communication, network function and behaviour over time.

584 **References:**

- 585 Abdelfattah, A.S., Kawashima, T., Singh, A., Novak, O., Liu, H., Shuai, Y., Huang, Y.-C., Campagnola, L.,
586 Seeman, S.C., Yu, J., Zheng, J., Grimm, J.B., Patel, R., Friedrich, J., Mensh, B.D., Paninski, L.,
587 Macklin, J.J., Murphy, G.J., Podgorski, K., Lin, B.-J., Chen, T.-W., Turner, G.C., Liu, Z., Koyama, M.,
588 Svoboda, K., Ahrens, M.B., Lavis, L.D., Schreiter, E.R., 2019. Bright and photostable chemigenetic
589 indicators for extended in vivo voltage imaging. *Science* eaav6416.
590 <https://doi.org/10.1126/science.aav6416>
- 591 Ahrens, M.B., Li, J.M., Orger, M.B., Robson, D.N., Schier, A.F., Engert, F., Portugues, R., 2012. Brain-wide
592 neuronal dynamics during motor adaptation in zebrafish. *Nature* 485, 471–477.
593 <https://doi.org/10.1038/nature11057>
- 594 Ahrens, M.B., Orger, M.B., Robson, D.N., Li, J.M., Keller, P.J., 2013. Whole-brain functional imaging at
595 cellular resolution using light-sheet microscopy. *Nat. Methods* 10, 413–420.
596 <https://doi.org/10.1038/nmeth.2434>
- 597 Almeida, R.G., 2018. The Rules of Attraction in Central Nervous System Myelination. *Front. Cell. Neurosci.*
598 12, 367. <https://doi.org/10.3389/fncel.2018.00367>
- 599 Almeida, R.G., Czopka, T., French-Constant, C., Lyons, D.A., 2011. Individual axons regulate the myelinating
600 potential of single oligodendrocytes in vivo. *Development* 138, 4443–4450.
601 <https://doi.org/10.1242/dev.071001>
- 602 Auer, F., Vagionitis, S., Czopka, T., 2018. Evidence for Myelin Sheath Remodeling in the CNS Revealed by In
603 Vivo Imaging. *Curr. Biol.* 28, 549–559.e3. <https://doi.org/10.1016/j.cub.2018.01.017>
- 604 Bonnefil, V., Dietz, K., Amatruda, M., Wentling, M., Aubry, A.V., Dupree, J.L., Temple, G., Park, H.-J.,
605 Burghardt, N.S., Casaccia, P., Liu, J., 2019. Region-specific myelin differences define behavioral
606 consequences of chronic social defeat stress in mice. *eLife* 8, e40855.
607 <https://doi.org/10.7554/eLife.40855>
- 608 Bujalka, H., Koenning, M., Jackson, S., Perreau, V.M., Pope, B., Hay, C.M., Mitew, S., Hill, A.F., Lu, Q.R.,
609 Wegner, M., Srinivasan, R., Svaren, J., Willingham, M., Barres, B.A., Emery, B., 2013. MYRF Is a
610 Membrane-Associated Transcription Factor That Autoproteolytically Cleaves to Directly Activate
611 Myelin Genes. *PLoS Biol.* 11, e1001625. <https://doi.org/10.1371/journal.pbio.1001625>
- 612 Burgess, H.A., Granato, M., 2007a. Sensorimotor Gating in Larval Zebrafish. *J. Neurosci.* 27, 4984–4994.
613 <https://doi.org/10.1523/JNEUROSCI.0615-07.2007>

- 614 Burgess, H.A., Granato, M., 2007b. Modulation of locomotor activity in larval zebrafish during light adaptation.
615 J. Exp. Biol. 210, 2526–2539. <https://doi.org/10.1242/jeb.003939>
- 616 Burgess, H.A., Johnson, S.L., Granato, M., 2009. Unidirectional startle responses and disrupted left-right co-
617 ordination of motor behaviors in *robo3* mutant zebrafish. Genes Brain Behav. 8, 500–511.
618 <https://doi.org/10.1111/j.1601-183X.2009.00499.x>
- 619 Chen, X., Mu, Y., Hu, Y., Kuan, A.T., Nikitchenko, M., Randlett, O., Chen, A.B., Gavornik, J.P., Sompolinsky,
620 H., Engert, F., Ahrens, M.B., 2018. Brain-wide Organization of Neuronal Activity and Convergent
621 Sensorimotor Transformations in Larval Zebrafish. Neuron 100, 876–890.e5.
622 <https://doi.org/10.1016/j.neuron.2018.09.042>
- 623 Czopka, T., Lyons, D.A., 2011. Dissecting Mechanisms of Myelinated Axon Formation Using Zebrafish, in:
624 Methods in Cell Biology. Elsevier, pp. 25–62. <https://doi.org/10.1016/B978-0-12-381320-6.00002-3>
- 625 Diana, G., Sainsbury, T.T.J., Meyer, M.P., 2018. Bayesian inference of neuronal ensembles. Biorxiv.
626 <https://doi.org/10.1101/452557>
- 627 Dreosti, E., Lopes, G., Kampff, A.R., Wilson, S.W., 2015. Development of social behavior in young zebrafish.
628 Front. Neural Circuits 9. <https://doi.org/10.3389/fncir.2015.00039>
- 629 Dunn, T.W., Gebhardt, C., Naumann, E.A., Riegler, C., Ahrens, M.B., Engert, F., Del Bene, F., 2016. Neural
630 Circuits Underlying Visually Evoked Escapes in Larval Zebrafish. Neuron 89, 613–628.
631 <https://doi.org/10.1016/j.neuron.2015.12.021>
- 632 Early, J.J., Cole, K.L., Williamson, J.M., Swire, M., Kamadurai, H., Muskavitch, M., Lyons, D.A., 2018. An
633 automated high-resolution in vivo screen in zebrafish to identify chemical regulators of myelination.
634 eLife 7, e35136. <https://doi.org/10.7554/eLife.35136>
- 635 Emery, B., Agalliu, D., Cahoy, J.D., Watkins, T.A., Dugas, J.C., Mulinyawe, S.B., Ibrahim, A., Ligon, K.L.,
636 Rowitch, D.H., Barres, B.A., 2009. Myelin Gene Regulatory Factor Is a Critical Transcriptional
637 Regulator Required for CNS Myelination. Cell 138, 172–185.
638 <https://doi.org/10.1016/j.cell.2009.04.031>
- 639 Fields, R.D., 2015. A new mechanism of nervous system plasticity: activity-dependent myelination. Nat. Rev.
640 Neurosci. 16, 756–767. <https://doi.org/10.1038/nrn4023>
- 641 Gahtan, E., O'Malley, D.M., 2003. Visually guided injection of identified reticulospinal neurons in zebrafish: A
642 survey of spinal arborization patterns. J. Comp. Neurol. 459, 186–200.
643 <https://doi.org/10.1002/cne.10621>

- 644 Geraghty, A.C., Gibson, E.M., Ghanem, R.A., Greene, J.J., Ocampo, A., Goldstein, A.K., Ni, L., Yang, T.,
645 Marton, R.M., Paşca, S.P., Greenberg, M.E., Longo, F.M., Monje, M., 2019. Loss of Adaptive
646 Myelination Contributes to Methotrexate Chemotherapy-Related Cognitive Impairment. *Neuron* 103,
647 250-265.e8. <https://doi.org/10.1016/j.neuron.2019.04.032>
- 648 Gibson, E.M., Geraghty, A.C., Monje, M., 2018. Bad wrap: Myelin and myelin plasticity in health and disease:
649 Myelin and Myelin Plasticity in Health and Disease. *Dev. Neurobiol.* 78, 123–135.
650 <https://doi.org/10.1002/dneu.22541>
- 651 Hale, M.E., Katz, H.R., Peek, M.Y., Fremont, R.T., 2016. Neural circuits that drive startle behavior, with a
652 focus on the Mauthner cells and spiral fiber neurons of fishes. *J. Neurogenet.* 30, 89–100.
653 <https://doi.org/10.1080/01677063.2016.1182526>
- 654 Hamanaka, K., Takata, A., Uchiyama, Y., Miyatake, S., Miyake, N., Mitsuhashi, S., Iwama, K., Fujita, A.,
655 Imagawa, E., Alkanaq, A.N., Koshimizu, E., Azuma, Y., Nakashima, M., Mizuguchi, T., Saitsu, H.,
656 Wada, Y., Minami, S., Katoh-Fukui, Y., Masunaga, Y., Fukami, M., Hasegawa, T., Ogata, T.,
657 Matsumoto, N., 2019. MYRF haploinsufficiency causes 46,XY and 46,XX disorders of sex
658 development: bioinformatics consideration. *Hum. Mol. Genet.* 28, 2319–2329.
659 <https://doi.org/10.1093/hmg/ddz066>
- 660 Henriques, P.M., Rahman, N., Jackson, S.E., Bianco, I.H., 2019. Nucleus Isthmi Is Required to Sustain Target
661 Pursuit during Visually Guided Prey-Catching. *Curr. Biol.* 29, 1771-1786.e5.
662 <https://doi.org/10.1016/j.cub.2019.04.064>
- 663 Hildebrand, D.G.C., Cicconet, M., Torres, R.M., Choi, W., Quan, T.M., Moon, J., Wetzel, A.W., Scott
664 Champion, A., Graham, B.J., Randlett, O., Plummer, G.S., Portugues, R., Bianco, I.H., Saalfeld, S.,
665 Baden, A.D., Lillaney, K., Burns, R., Vogelstein, J.T., Schier, A.F., Lee, W.-C.A., Jeong, W.-K.,
666 Lichtman, J.W., Engert, F., 2017. Whole-brain serial-section electron microscopy in larval zebrafish.
667 *Nature* 545, 345–349. <https://doi.org/10.1038/nature22356>
- 668 Hines, J.H., Ravanelli, A.M., Schwindt, R., Scott, E.K., Appel, B., 2015. Neuronal activity biases axon selection
669 for myelination in vivo. *Nat. Neurosci.* 18, 683–689. <https://doi.org/10.1038/nn.3992>
- 670 Jain, R.A., Wolman, M.A., Marsden, K.C., Nelson, J.C., Shoenhard, H., Echeverry, F.A., Szi, C., Bell, H.,
671 Skinner, J., Cobbs, E.N., Sawada, K., Zamora, A.D., Pereda, A.E., Granato, M., 2018. A Forward
672 Genetic Screen in Zebrafish Identifies the G-Protein-Coupled Receptor CaSR as a Modulator of

- 673 Sensorimotor Decision Making. Curr. Biol. 28, 1357-1369.e5.
674 <https://doi.org/10.1016/j.cub.2018.03.025>
- 675 Jao, L.-E., Wente, S.R., Chen, W., 2013. Efficient multiplex biallelic zebrafish genome editing using a CRISPR
676 nuclease system. *Proc. Natl. Acad. Sci.* 110, 13904–13909. <https://doi.org/10.1073/pnas.1308335110>
- 677 Karttunen, M.J., Czopka, T., Goedhart, M., Early, J.J., Lyons, D.A., 2017. Regeneration of myelin sheaths of
678 normal length and thickness in the zebrafish CNS correlates with growth of axons in caliber. *PLOS*
679 ONE 12, e0178058. <https://doi.org/10.1371/journal.pone.0178058>
- 680 Kim, J.H., Renden, R., von Gersdorff, H., 2013. Dysmyelination of Auditory Afferent Axons Increases the Jitter
681 of Action Potential Timing during High-Frequency Firing. *J. Neurosci.* 33, 9402–9407.
682 <https://doi.org/10.1523/JNEUROSCI.3389-12.2013>
- 683 Kimmel, C.B., Patterson, J., Kimmel, R.O., 1974. The development and behavioral characteristics of the startle
684 response in the zebra fish. *Dev. Psychobiol.* 7, 47–60. <https://doi.org/10.1002/dev.420070109>
- 685 Klingseisen, A., Ristoiu, A.-M., Kegel, L., Sherman, D.L., Rubio-Brotons, M., Almeida, R.G., Koudelka, S.,
686 Benito-Kwiecinski, S.K., Poole, R.J., Brophy, P.J., Lyons, D.A., 2019. Oligodendrocyte Neurofascin
687 Independently Regulates Both Myelin Targeting and Sheath Growth in the CNS. *Dev. Cell* 51, 730-
688 744.e6. <https://doi.org/10.1016/j.devcel.2019.10.016>
- 689 Koenning, M., Jackson, S., Hay, C.M., Faux, C., Kilpatrick, T.J., Willingham, M., Emery, B., 2012. Myelin
690 Gene Regulatory Factor Is Required for Maintenance of Myelin and Mature Oligodendrocyte Identity
691 in the Adult CNS. *J. Neurosci.* 32, 12528–12542. <https://doi.org/10.1523/JNEUROSCI.1069-12.2012>
- 692 Koudelka, S., Voas, M.G., Almeida, R.G., Baraban, M., Soetaert, J., Meyer, M.P., Talbot, W.S., Lyons, D.A.,
693 2016. Individual Neuronal Subtypes Exhibit Diversity in CNS Myelination Mediated by Synaptic
694 Vesicle Release. *Curr. Biol.* 26, 1447–1455. <https://doi.org/10.1016/j.cub.2016.03.070>
- 695 Lacoste, A.M.B., Schoppik, D., Robson, D.N., Haesemeyer, M., Portugues, R., Li, J.M., Randlett, O., Wee,
696 C.L., Engert, F., Schier, A.F., 2015. A Convergent and Essential Interneuron Pathway for Mauthner-
697 Cell-Mediated Escapes. *Curr. Biol.* 25, 1526–1534. <https://doi.org/10.1016/j.cub.2015.04.025>
- 698 Larsch, J., Baier, H., 2018. Biological Motion as an Innate Perceptual Mechanism Driving Social Affiliation.
699 *Curr. Biol.* 28, 3523-3532.e4. <https://doi.org/10.1016/j.cub.2018.09.014>
- 700 Liu, J., Dupree, J.L., Gacias, M., Frawley, R., Sikder, T., Naik, P., Casaccia, P., 2016. Clemastine Enhances
701 Myelination in the Prefrontal Cortex and Rescues Behavioral Changes in Socially Isolated Mice. *J.*
702 *Neurosci.* 36, 957–962. <https://doi.org/10.1523/JNEUROSCI.3608-15.2016>

- 703 Makinodan, M., Rosen, K.M., Ito, S., Corfas, G., 2012. A Critical Period for Social Experience-Dependent
704 Oligodendrocyte Maturation and Myelination. *Science* 337, 1357–1360.
705 <https://doi.org/10.1126/science.1220845>
- 706 Marquart, G.D., Tabor, K.M., Bergeron, S.A., Briggman, K.L., Burgess, H.A., 2019. Preoptine non-giant
707 neurons drive flexible escape behavior in zebrafish (preprint). *Neuroscience*.
708 <https://doi.org/10.1101/668517>
- 709 Marques, J.C., Lackner, S., Félix, R., Orger, M.B., 2018. Structure of the Zebrafish Locomotor Repertoire
710 Revealed with Unsupervised Behavioral Clustering. *Curr. Biol.* 28, 181-195.e5.
711 <https://doi.org/10.1016/j.cub.2017.12.002>
- 712 McKenzie, I.A., Ohayon, D., Li, H., Paes de Faria, J., Emery, B., Tohyama, K., Richardson, W.D., 2014. Motor
713 skill learning requires active central myelination. *Science* 346, 318–322.
714 <https://doi.org/10.1126/science.1254960>
- 715 Mensch, S., Baraban, M., Almeida, R., Czopka, T., Ausborn, J., El Manira, A., Lyons, D.A., 2015. Synaptic
716 vesicle release regulates myelin sheath number of individual oligodendrocytes in vivo. *Nat. Neurosci.*
717 18, 628–630. <https://doi.org/10.1038/nn.3991>
- 718 Metcalfe, W.K., Mendelson, B., Kimmel, C.B., 1986. Segmental homologies among reticulospinal neurons in
719 the hindbrain of the zebrafish larva. *J. Comp. Neurol.* 251, 147–159.
720 <https://doi.org/10.1002/cne.902510202>
- 721 Mount, C.W., Monje, M., 2017. Wrapped to Adapt: Experience-Dependent Myelination. *Neuron* 95, 743–756.
722 <https://doi.org/10.1016/j.neuron.2017.07.009>
- 723 Naumann, E.A., Fitzgerald, J.E., Dunn, T.W., Rihel, J., Sompolinsky, H., Engert, F., 2016. From Whole-Brain
724 Data to Functional Circuit Models: The Zebrafish Optomotor Response. *Cell* 167, 947-960.e20.
725 <https://doi.org/10.1016/j.cell.2016.10.019>
- 726 Nave, K.-A., Werner, H.B., 2014. Myelination of the Nervous System: Mechanisms and Functions. *Annu. Rev.*
727 *Cell Dev. Biol.* 30, 503–533. <https://doi.org/10.1146/annurev-cellbio-100913-013101>
- 728 Pan, S., Mayoral, S.R., Choi, H.S., Chan, J.R., Kheirbek, M.A., 2020. Preservation of a remote fear memory
729 requires new myelin formation. *Nat. Neurosci.* 23, 487–499. [https://doi.org/10.1038/s41593-019-0582-](https://doi.org/10.1038/s41593-019-0582-1)
730 1
- 731 Pinz, H., Pyle, L.C., Li, D., Izumi, K., Skraban, C., Tarpinian, J., Braddock, S.R., Telegrafi, A., Monaghan,
732 K.G., Zackai, E., Bhoj, E.J., 2018. De novo variants in Myelin regulatory factor (MYRF) as candidates

- 733 of a new syndrome of cardiac and urogenital anomalies. *Am. J. Med. Genet. A.* 176, 969–972.
734 <https://doi.org/10.1002/ajmg.a.38620>
- 735 Preston, M.A., Macklin, W.B., 2015. Zebrafish as a model to investigate CNS myelination: Zebrafish
736 Myelination. *Glia* 63, 177–193. <https://doi.org/10.1002/glia.22755>
- 737 Redmond, S.A., Mei, F., Eshed-Eisenbach, Y., Osso, L.A., Leshkowitz, D., Shen, Y.-A.A., Kay, J.N., Aurrand-
738 Lions, M., Lyons, D.A., Peles, E., Chan, J.R., 2016. Somatodendritic Expression of JAM2 Inhibits
739 Oligodendrocyte Myelination. *Neuron* 91, 824–836. <https://doi.org/10.1016/j.neuron.2016.07.021>
- 740 Romano, S.A., Pietri, T., Pérez-Schuster, V., Jouary, A., Haudrechy, M., Sumbre, G., 2015. Spontaneous
741 Neuronal Network Dynamics Reveal Circuit's Functional Adaptations for Behavior. *Neuron* 85, 1070–
742 1085. <https://doi.org/10.1016/j.neuron.2015.01.027>
- 743 Rossetti, L.Z., Glington, K., Yuan, B., Liu, P., Pillai, N., Mizerik, E., Magoulas, P., Rosenfeld, J.A., Karaviti, L.,
744 Sutton, V.R., Lalani, S.R., Scott, D.A., 2019. Review of the phenotypic spectrum associated with
745 haploinsufficiency of *MYRF*. *Am. J. Med. Genet. A.* *ajmg.a.61182*.
746 <https://doi.org/10.1002/ajmg.a.61182>
- 747 Rossi, A., Kontarakis, Z., Gerri, C., Nolte, H., Hölper, S., Krüger, M., Stainier, D.Y.R., 2015. Genetic
748 compensation induced by deleterious mutations but not gene knockdowns. *Nature* 524, 230–233.
749 <https://doi.org/10.1038/nature14580>
- 750 Roy, B., Ali, D.W., 2013. Patch Clamp Recordings from Embryonic Zebrafish Mauthner Cells. *J. Vis. Exp.*
751 50551. <https://doi.org/10.3791/50551>
- 752 Saint-Amant, L., Drapeau, P., 2003. Whole-cell patch-clamp recordings from identified spinal neurons in the
753 zebrafish embryo. *Methods Cell Sci.* 25, 59–64. <https://doi.org/10.1023/B:MICS.0000006896.02938.49>
- 754 Sampaio-Baptista, C., Khrapitchev, A.A., Foxley, S., Schlagheck, T., Scholz, J., Jbabdi, S., DeLuca, G.C.,
755 Miller, K.L., Taylor, A., Thomas, N., Kleim, J., Sibson, N.R., Bannerman, D., Johansen-Berg, H.,
756 2013. Motor Skill Learning Induces Changes in White Matter Microstructure and Myelination. *J.*
757 *Neurosci.* 33, 19499–19503. <https://doi.org/10.1523/JNEUROSCI.3048-13.2013>
- 758 Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C.,
759 Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J., Hartenstein, V., Eliceiri, K., Tomancak, P.,
760 Cardona, A., 2012. Fiji: an open-source platform for biological-image analysis. *Nat. Methods* 9, 676–
761 682. <https://doi.org/10.1038/nmeth.2019>

- 762 Steadman, P.E., Xia, F., Ahmed, M., Mocle, A.J., Penning, A.R.A., Geraghty, A.C., Steenland, H.W., Monje,
763 M., Josselyn, S.A., Frankland, P.W., 2020. Disruption of Oligodendrogenesis Impairs Memory
764 Consolidation in Adult Mice. *Neuron* 105, 150-164.e6. <https://doi.org/10.1016/j.neuron.2019.10.013>
- 765 Sumbre, G., Muto, A., Baier, H., Poo, M., 2008. Entrained rhythmic activities of neuronal ensembles as
766 perceptual memory of time interval. *Nature* 456, 102–106. <https://doi.org/10.1038/nature07351>
- 767 Tabor, K.M., Smith, T.S., Brown, M., Bergeron, S.A., Briggman, K.L., Burgess, H.A., 2018. Presynaptic
768 Inhibition Selectively Gates Auditory Transmission to the Brainstem Startle Circuit. *Curr. Biol.* 28,
769 2527-2535.e8. <https://doi.org/10.1016/j.cub.2018.06.020>
- 770 Tanimoto, M., Ota, Y., Horikawa, K., Oda, Y., 2009. Auditory Input to CNS Is Acquired Coincidentally with
771 Development of Inner Ear after Formation of Functional Afferent Pathway in Zebrafish. *J. Neurosci.*
772 29, 2762–2767. <https://doi.org/10.1523/JNEUROSCI.5530-08.2009>
- 773 Treichel, A.J., Hines, J.H., 2018. Development of an Embryonic Zebrafish Oligodendrocyte–Neuron Mixed
774 Coculture System. *Zebrafish* 15, 586–596. <https://doi.org/10.1089/zeb.2018.1625>
- 775 Wang, F., Ren, S.-Y., Chen, J.-F., Liu, K., Li, R.-X., Li, Z.-F., Hu, B., Niu, J.-Q., Xiao, L., Chan, J.R., Mei, F.,
776 2020. Myelin degeneration and diminished myelin renewal contribute to age-related deficits in
777 memory. *Nat. Neurosci.* 23, 481–486. <https://doi.org/10.1038/s41593-020-0588-8>
- 778 Waxman, S.G., 1980. Determinants of conduction velocity in myelinated nerve fibers. *Muscle Nerve* 3, 141–
779 150. <https://doi.org/10.1002/mus.880030207>
- 780 Wen, H., 2005. Paired Motor Neuron-Muscle Recordings in Zebrafish Test the Receptor Blockade Model for
781 Shaping Synaptic Current. *J. Neurosci.* 25, 8104–8111. <https://doi.org/10.1523/JNEUROSCI.2611-05.2005>
- 782
- 783 Wolf, S., Dubreuil, A.M., Bertoni, T., Böhm, U.L., Bormuth, V., Candelier, R., Karpenko, S., Hildebrand,
784 D.G.C., Bianco, I.H., Monasson, R., Debrégeas, G., 2017. Sensorimotor computation underlying
785 phototaxis in zebrafish. *Nat. Commun.* 8, 651. <https://doi.org/10.1038/s41467-017-00310-3>
- 786 Wolman, M.A., Jain, R.A., Liss, L., Granato, M., 2011. Chemical modulation of memory formation in larval
787 zebrafish. *Proc. Natl. Acad. Sci.* 108, 15468–15473. <https://doi.org/10.1073/pnas.1107156108>
- 788 Yamanaka, I., Miki, M., Asakawa, K., Kawakami, K., Oda, Y., Hirata, H., 2013. Glycinergic transmission and
789 postsynaptic activation of CaMKII are required for glycine receptor clustering *in vivo*. *Genes Cells* 18,
790 211–224. <https://doi.org/10.1111/gtc.12032>

791 Zottoli, S.J., Bentley, A.P., Prendergast, B.J., Rieff, H., I., 1995. Comparative Studies on the Mauthner Cell of
792 Teleost Fish in Relation to Sensory Input. *Brain Behav. Evol.* 46, 151–164.
793
794

795 **Figure Legends**

796

797 **Figure 1: *myrf^{me70}* mutants display a gross reduction in the level of CNS myelination at the adult and**
798 **larval stages.**

799 **A, Top:** *myrf* gene structure composed of 27 exons. Red arrowhead marks the location of the mutation in exon
800 2. Scale bar equates to 1000bp. Schematic created using <http://wormweb.org>. **Middle:** Wildtype and mutant
801 nucleotide sequences spanning the mutagenesis site. The guide RNA (gRNA) target site (red line) and
802 restriction enzyme (RE) recognition site (green line) are labelled. **Bottom:** Amino acid sequence indicating that
803 the *myrf^{me70}* mutation results in shift in the open reading frame leading to downstream coding for a premature
804 stop codon (*).

805 **B,** The relative concentration of *mbp* mRNA is reduced by 95% in mutants (0.04 ± 0.03 au) compared to
806 wildtypes (1.003 ± 0.13 au, $p = 0.0002$, unpaired t test, $N = 3$ adult brains per genotype).

807 **C,** Transverse section of the spinal cord in an adult *myrf^{me70}* sibling showing extensive myelination of ventral
808 spinal cord (dashed box). 20x objective. Scale bar = 100 μ m.

809 **D,** TEM images of the spinal cord in the region of the ventral spinal tract (outlined in C) in *myrf^{me70}* adult
810 siblings (top) and mutants (bottom). Panels i-iv display different fields of view within the region of interest.
811 Thick myelin sheaths are clearly visible in siblings, particularly surrounding the Mauthner axon. There is a lack
812 of myelin surrounding the Mauthner axon in the mutant sample, and distinct reduction in the level of
813 myelination in the remainder of surrounding spinal cord. Occasional hypomyelinated and dysmyelinated axons
814 can be observed in the mutant samples. Scale bar = 5 microns for panels i-iii. Scale bar = 1 micron for panel iv.
815 'm' denotes the Mauthner axon.

816 **E, Top:** Brightfield images of *myrf^{me70}* wildtype and mutant larvae at 6 dpf. Black box defines the anatomical
817 region imaged across animals. Scale bar = 0.5mm. **Bottom:** Confocal microscopy images of the spinal cord at 6
818 dpf in *myrf^{me70}* Tg(*mbp:eGFP-CAAX*) larvae. Scale bar = 20 μ m.

819

820

821

822 **Figure 2: *myrf^{me70}* mutants display CNS-specific hypomyelination at 6 dpf.**

823 **A**, TEM images of the myelinated tracts in the dorsal (top row) and ventral spinal cord (bottom rows). Scale
824 bars = 1 μ m.

825 **B**, Schematic of the transverse section of a 6 dpf larval zebrafish at the level of the urogenital opening. **Inset**:
826 transverse section of the spinal cord at the same level. Myelinated (green) axons are located in the ventral and
827 dorsal spinal tracts of the spinal cord (CNS) as well as the posterior lateral line (PNS). m = Mauthner axons.

828 **C**, TEM images of the posterior lateral line at 6 dpf. Scale bar = 1 μ m.

829 **D**, The average number of myelinated axons in one hemi-spinal cord is reduced by 66% in mutants (wildtypes:
830 35.29 ± 7.83 myelinated axons, mutants: 12.00 ± 4.34 myelinated axons, $p \leq 0.0001$, unpaired t-test, $N = 7$
831 wildtypes, $N = 8$ mutants).

832 **E**, The number of myelinated axons in the PNS is similar between genotypes (wildtypes: 7.33 ± 1.53
833 myelinated axons, mutants: 9.00 ± 3.83 myelinated axons, $p = 0.52$, unpaired t-test, $N = 3$ wildtypes, $N = 4$
834 mutants). Values represent mean \pm standard deviation.

835 **F**, G-ratio of Mauthner axons in wildtype and mutant siblings (wildtypes: 0.48 ± 0.009 , mutants: 0.80 ± 0.08 , p
836 $= 0.0009$, unpaired t-test).

837 **G**, G-ratios for myelinated axons for small caliber (area $<0.3\mu\text{m}^2$) and large caliber (area $>0.3\mu\text{m}^2$) myelinated
838 axons. The g-ratio of small caliber axons is similar between groups (wildtypes: 0.57 (0.52 to 0.62), mutants:
839 0.59 (0.52 to 0.70), $p = 0.51$, Mann Whitney test, $n = 53$ myelinated axons in wildtypes, $n = 17$ myelinated
840 axons in mutants). The g-ratios for large caliber axons are significantly higher in mutants than wildtype siblings
841 (wildtypes: 0.60 ± 0.08 , mutants: 0.71 ± 0.08 , $p \leq 0.0001$, unpaired t-test, $n = 33$ myelinated axons in
842 wildtypes, $n = 19$ myelinated axons in mutants).

843

844

845 **Figure 3: *myrf^{μe70}* mutants have fewer oligodendrocytes which produce less myelin and fail to maintain**
846 **myelin sheaths over time.**

847 **A**, Confocal images of the spinal cord at 6 dpf in sibling control and *myrf^{μe70}* Tg(mbp:nls-eGFP) larvae. Scale
848 bar = 100μm.

849 **B**, Oligodendrocyte numbers in the spinal cord at 6 dpf (wildtype: 304.8 ± 39.07, mutants: 239.3 ± 50.48, p =
850 0.0002, unpaired t test, N = 15 wildtypes, N = 22 mutants). Error bars represent mean ± standard deviation.

851 **C**, Representative confocal images of single oligodendrocytes mosaically labelled with mbp:mCherry-CAAX
852 reporter construct in a wildtype (top) and mutant (bottom) at 6 dpf. Scale bar = 15μm.

853 **D**, Average myelin sheath number was reduced in *myrf^{μe70}* mutants relative to wildtype siblings at 6 dpf
854 (wildtypes: 10.50 (7.00 to 14.00) sheaths per cell, mutants: 7.00 (5.00 to 10.50) sheaths per cell, p = 0.02, Mann
855 Whitney test). Values and error bars represent median and IQR.

856 **E**, Average myelin sheath length was reduced from 41.83 ± 9.68μm in wildtypes to 31.35 ± 11.49μm in
857 mutants at 6 dpf (p = 0.002, unpaired t test). Error bars represent mean ± standard deviation.

858 **F**, Total myelin produced per oligodendrocyte was reduced from 458.2 ± 156.4μm in wildtypes to 241.1
859 ± 138.6μm in mutants at 6 dpf (p ≤ 0.0001, unpaired t test). Error bars represent mean ± standard deviation.

860 **D-F**: N = 20 wildtypes, N = 27 mutants.

861 **G**, Confocal images of a single mutant oligodendrocyte labelled with mbp:mCherry-CAAX at 4 and 6 dpf. A
862 myelin sheath (*) and myelinated neuronal cell body (#) are observed at 4 dpf and subsequently retracted by 6
863 dpf. Arrowheads label myelin sheaths which are observed to shrink between 4 and 6 dpf. Scale bar = 15μm.

864 **H**, Myelin sheaths belonging to wildtype oligodendrocytes demonstrated a net growth of 6.24 ± 3.43μm
865 between 4 and 6 dpf, while mutants display net shrinkage of myelin sheaths by -0.31 ± 4.79μm (p = 0.003,
866 unpaired t test). Error bars represent mean ± standard deviation.

867 **I**, Between 4 and 6 dpf, wildtype oligodendrocytes retracted 0 (0 to 0) myelin sheaths, while mutants retracted 2
868 (1 to 3) myelin sheaths (p = 0.009, Mann Whitney test). Error bars represent median and IQR.

869 **J**, Number of abnormal myelin sheaths at 6 dpf (wildtypes: 0.00 (0.00-0.00); mutants: 2 (0.00-3.00), p ≤ 0.0001,
870 Mann Whitney test). Error bars represent median and IQR.

871 **H – I**: N = 11 wildtypes, N = 7 mutants. **J**: N = 20 wildtypes, N = 27 mutants.

872

873

874 **Figure 4: *myrf^{ue70}* mutants exhibit increased latency to perform startle responses, and a tendency to**
875 **perform avoidance behaviour, in response to defined acoustic stimuli.**

876 **A**, Overview of the neuronal circuitry involved in motor response to auditory stimuli. **Startle response (SLC):**
877 sensory input from the ear, via the auditory nerve (red), is received at the lateral dendrite of the Mauthner cell
878 body (black). The axon of the Mauthner cell crosses into the contralateral aspect of the spinal cord where it
879 extends along the ventral tract to recruit motor neurons directly along the length of the larvae. Recruitment of
880 motor neurons allows muscle contraction on the side of the body contralateral to the stimulus, allowing a rapid,
881 high-velocity c-bend (motor response) away from the stimulus (inset). **Avoidance behaviour (LLC):** sensory
882 input is detected by prepontine neurons (purple) in the hindbrain, which recruit ipsilateral motor neurons
883 indirectly, resulting in a low-velocity, longer latency, c-bend away from the stimulus.

884 **B**, Schematic of the behavioural rig.

885 **C**, Relative frequency histogram displaying the distribution of latencies for behavioural responses in response to
886 acoustic stimuli in wildtype and mutant larvae (N = 24 wildtype larvae, n = 220 events; N = 35 mutant larvae, n
887 = 299 events; Kolmogorov-Smirnov test, $p \leq 0.0001$).

888 **D**, Number and proportion of events (SLC vs LLC) per genotype.

889 **E**, React rate per fish (median react rate = 100% in both wildtypes and mutants, $p = 0.24$, Mann-Whitney test, N
890 = 25 wildtype larvae, N = 38 mutant larvae). Larvae are excluded from subsequent analysis if they exhibit a
891 react rate <70%.

892 **F**, Average latency values per fish (wildtype: 10.55ms (9.6-16.15ms), mutants: 17.6ms (12.9-21.88ms), $p =$
893 0.003, Mann-Whitney test).

894 **G**, Average latency of short latency c-starts (<16ms) (wildtypes: 10.03 ± 0.85 ms, mutants: 10.67 ± 0.83 ms, $p =$
895 0.006, unpaired t test).

896 **H**, Average latency of long latency c-starts (>16ms) (wildtypes: 43.20 ± 8.95 ms; mutants: 38.91 ± 10.15 ms, $p =$
897 0.28, unpaired t test).

898 **I**, Mean and standard deviations values for SLC and LLC responses per genotype.

899 J-M analysis of c-bend kinematics:

900 **J**, Example trace of orientation over time during a behavioural response to an acoustic stimulus. C-bend
901 kinematics are calculated from individual traces for each response per fish. Latency is the time from stimulus
902 onset to behavioural onset (red star). C-bend duration (A) is time from behaviour onset to initial turn angle (blue

903 star). Maximum angular velocity is defined as the change in orientation over time (B/A). Turning angle equates
904 to the initial turn angle.

905 **K**, Initial turn duration (SLC: wildtypes: 10.06 ± 0.70 ms, mutants: 9.81 ± 0.72 ms, $p = 0.20$, unpaired t-test;
906 LLC: wildtypes: 14.30 ± 3.65 ms, mutants: 13.25 ± 3.10 , $p = 0.42$, unpaired t test).

907 **L**, Maximum angular velocity (SLC: wildtypes: 24° /ms (22.78-28.68 $^\circ$ /ms), mutants: 25° /ms (23.10-26.60 $^\circ$ /ms),
908 $p = 0.73$, Mann-Whitney test; LLC: wildtypes: $16.16 \pm 6.61^\circ$ /ms, mutants: $13.67 \pm 4.95^\circ$ /ms, $p = 0.24$, unpaired
909 t test).

910 **M**, Initial turn angle (SLC: wildtypes: $121.9 \pm 10.80^\circ$, mutants: $127.4 \pm 9.86^\circ$, $p = 0.051$, unpaired t test; LLC:
911 wildtypes: $85.11 \pm 35.78^\circ$, mutants: $83.78 \pm 29.07^\circ$, $p = 0.91$, unpaired t test).

912 **N**, Descriptive statistics (mean \pm standard deviation) for c-bend kinematics.

913 For **Figures E-G and K-M**, $N = 23$ wildtypes, $N = 35$ mutant larvae. For **Figures D & E** values represent
914 median and interquartile range, for **Figures F-K**, values represent mean \pm standard deviation.

915

916

917 **Figure 5: Whole cell current-clamp recordings from Mauthner cells demonstrate slower conduction**
918 **velocity times and abnormal spiking profiles in *myrf^{ue70}* mutants.**

919 **A**, Electrophysiological preparation for recording from Mauthner neuron in a whole-cell current clamp
920 configuration while stimulating with an extracellular monopolar field electrode midway through the spinal cord.

921 **B**, Resting membrane potential is unchanged in (siblings (n = 18 cells): $-70.82 \pm 2.76\text{mV}$, mutants (n = 6 cells):
922 $-70.68 \pm 1.25\text{mV}$, $p = 0.9077$ at 6 dpf).

923 **C**, Sample trace of an action potential recorded at 6 dpf in a wildtype fish illustrating the measurement of half-
924 width. Half-width is described as width of action potential (ms) at its half height.

925 **D**, Half-width of action potential is unchanged (siblings (n = 18 cells): $0.64 \pm 0.09\text{ms}$, mutants (n = 9 cells):
926 $0.60 \pm 0.06\text{ms}$, $p = 0.2610$ at 6 dpf).

927 **E**, An example of current – clamp recording from Mauthner neuron in a 6 dpf wildtype and mutant following
928 field stimulation (stimulus artefact is indicated by a grey dashed line). Latency is described as time from the
929 onset of stimulus artefact to the peak of action potential.

930 **F**, Normalised action potential latency is increased in mutants at 6 dpf (siblings (n = 19 cells): $0.80 \pm$
931 0.11ms/mm , mutants (n = 9 cells): $0.97 \pm 0.07\text{ms/mm}$, $p = 0.0003$ at 6 dpf).

932 **G**, Conduction velocity of Mauthner action potentials is significantly decreased in mutant larvae (siblings (n =
933 19 cells): $1.27 \pm 0.17\text{m/s}$, mutants (n = 9 cells): $1.04 \pm 0.08\text{m/s}$, $p = 0.0005$ at 6 dpf).

934 **H**, Sample traces of three subsequent action potentials recorded from the same wildtype Mauthner cell at 6 dpf
935 superimposed and aligned to the peak of stimulus artefact. The area outlined by the rectangle is magnified in the
936 inset and demonstrates slight imprecision of action potential arrival.

937 **I**, Precision of action potential arrival is comparable in siblings and mutants (siblings (n = 16 cells): $0.0064 \pm$
938 0.0019ms , mutants (n = 8 cells): $0.0062 \pm 0.0009\text{ms}$, $p = 0.8166$ at 6 dpf).

939 **J**, Sample trace of a train of action potentials fired following 10 stimuli at 1000 Hz at 6 dpf in a *myrf^{ue70}* mutant
940 and sibling.

941 **K**, Mauthner neurons in mutant larvae do not sustain prolonged action potential trains of high frequency
942 stimulation (siblings (n = 19 cells): $55.79 \pm 10.17\%$ mutants (n = 9 cells): $38.89 \pm 17.64\%$ at 6 dpf, $p = 0.0014$
943 at 6 dpf).

944 For **Figures 5 B, D, F, G, I, K** error bars represent mean \pm standard deviation. Unpaired t-test for **Figures 5 B,**
945 **D, F, G, I** and a two-way ANOVA for **5K**. Scale bars are 10mV and 1ms for **Figures 5C, E, H, J** and 5 mV and
946 200 μs for **Figure 5H inset**.









