



6-2021

Four Decades of Understanding Martian Geomorphology: Revisiting Baker's "The Geomorphology of Mars"

Anshuman Bhardwaj
University of Aberdeen

Lydia Sam
University of Aberdeen

Saeideh Gharehchahi
Jacksonville State University, sgharehchahi@jsu.edu

Follow this and additional works at: https://digitalcommons.jsu.edu/fac_res



Part of the [Geomorphology Commons](#)

Recommended Citation

Bhardwaj, A., Sam, L. and Gharehchahi, S. (2021). Four decades of understanding Martian geomorphology: Revisiting Baker's 'The geomorphology of Mars'. *Progress in Physical Geography*, <https://doi.org/10.1177/03091333211026215>

This Article is brought to you for free and open access by the Faculty Scholarship & Creative Work at JSU Digital Commons. It has been accepted for inclusion in Research, Publications & Creative Work by an authorized administrator of JSU Digital Commons. For more information, please contact digitalcommons@jsu.edu.

1 **Four decades of understanding Martian geomorphology: Revisiting Baker’s “The**
2 **geomorphology of Mars”**

3 Anshuman Bhardwaj^{1*}, Lydia Sam¹, Saeideh Gharehchahi²

4 ¹School of Geosciences, University of Aberdeen, Meston Building, King’s College, Aberdeen
5 AB24 3UE, UK

6 ²Department of Chemistry and Geosciences, Jacksonville State University, Jacksonville, AL
7 36265, USA

8 *Corresponding author

9 **Abstract:** Owing to multiple successful orbiter and rover missions in the past two and half
10 decades, our understanding of the Martian atmosphere, terrain, and subsurface has
11 continuously evolved. This prompts the need to revisit the first holistic review of Martian
12 geomorphology based on useful images from Viking Mission orbiters, authored by Prof. Victor
13 R. Baker. Several of the remote sensing-based interpretations and recommendations in Baker’s
14 (1981) paper are as valid even today as they were four decades back. With an unprecedented
15 focus on Mars exploration in the coming decades, it is important to briefly revisit the advances
16 and prospects in Martian geomorphology research.

17 **Keywords:** Mars, geomorphology, Viking Mission, planetary exploration, remote sensing

18 **I Introduction**

19 In our solar system, Mars is the planet with the highest Earth similarity and relative planetary
20 habitability indices, based on various physical and physicochemical determinants (Schulze-
21 Makuch et al., 2011). Being terrestrial planets, the structural and compositional similarities
22 between Earth and Mars are further apparent from the relative geological and
23 geomorphological interpretations (Baker, 1981). The Martian regolith contains minerals and
24 the temperatures are within an acceptable range for the existence of life (as we know it). The

25 moderate Martian gravity can enable future colonization, and the Martian obliquity and the day
26 length are also comparable to Earth, giving the red planet its distinct seasons. Thus, it is not a
27 surprise that the leading space agencies and space companies are investing significant resources
28 in enabling further Mars exploration within the next couple of decades. However, while
29 technological advancements in engineering and computing have certainly bolstered this
30 confidence, the contribution from the vast influx of orbiter and rover observations, in the past
31 two and half decades in facilitating our understanding of the Martian atmosphere, terrain, and
32 subsurface, cannot be ignored (Bhardwaj et al., 2019a).

33 This prompts the need to revisit the first holistic and comprehensive account of Viking
34 Mission-based interpretations of Martian geomorphology, titled “The geomorphology of Mars”
35 and authored by Prof. Victor R. Baker in 1981. Undoubtedly there have been considerable
36 developments in the discipline since the publication of Baker’s (1981) paper; thus, revisiting
37 this work will clearly highlight the impacts of evolving techniques and tools on planetary
38 geomorphological interpretations. Although the short format of this “classics revisited” paper
39 does not allow for a detailed analysis of all the advances made in Martian geomorphology
40 research, we have provided key references throughout this article which the interested readers
41 can further explore. Instead, here we focus on key facets of Baker’s (1981) work to highlight
42 the status of our understanding of the Martian geomorphology in the Viking Mission era and
43 the considerable advancements since then. It is interesting to identify and suitable to
44 acknowledge, how many of Baker’s (1981) viewpoints still hold relevance, across several
45 themes within the discipline of Martian geomorphology.

46 **II. Planetary geomorphology through terrestrial analogy**

47 Baker starts his paper with an interesting example of the Chief Geologist for the US Geological
48 Survey, G.K. Gilbert, who had to abandon planned fieldworks after 1892 congressional budget

49 cuts to the Survey. Gilbert channelised this to an opportunity by utilising his time in studying
50 the moon through the naval observatory telescope in Washington, and thus, publishing one of
51 the most detailed accounts of contemporary lunar geomorphology (Gilbert, 1893). This
52 highlights how a scientist trained in terrestrial geomorphology can contribute significantly to
53 planetary geomorphology. Baker gives some more contemporary examples (e.g., Mutch, 1979;
54 Sharp 1980) discussing the relevance of analogy-based planetary exploration using remote
55 sensing images. Starting from the best available spatial resolution of ~10 m/pixel for the Viking
56 images to as high as ~25 cm/pixel High Resolution Imaging Science Experiment (HiRISE)
57 camera resolution for Mars today, over these past decades, we have seen numerous similar
58 analogous interpretations (e.g., Edgett et al., 2003; Irwin et al., 2004; Tsibulskaya et al., 2020;
59 Wood, 2006) advancing our knowledge of Martian landforms. Interestingly, while mentioning
60 the ~10 m/pixel resolution Viking images, Baker (1981) writes, “this is better resolution than
61 is available for portions of Earth”. Coming to the present scenario, the ~25 cm/pixel resolution
62 provided by the HiRISE camera for Mars is in public domain and we cannot expect such freely
63 available dataset for parts of Earth. Although the advent of unmanned aerial vehicles (UAVs)
64 (Bhardwaj et al., 2016; Gaffey and Bhardwaj, 2020) has led to an option of acquiring images,
65 comparable to HiRISE-resolutions, for Mars analogue research on Earth (e.g., Bhardwaj et al.,
66 2019b; Sam et al., 2020a; Sam et al., 2020b), the applications of this technique are still limited,
67 owing to the generally inaccessible nature of the analogue sites. In the subsequent sections,
68 various relevant references are provided which can be taken as examples of some remarkable
69 approaches where the knowledge of terrestrial geomorphology was comprehensively
70 extrapolated to Mars.

71 **III. Geomorphic map of Mars**

72 As an important contribution, Baker’s (1981) Figure 1 provides a holistic geomorphic map of
73 Mars, representing global distribution of various physiographic features. This map was

74 modified from the geological map by Scott and Carr (1978), issued as a US Geological Survey
75 publication. Baker (1981) classified the heavily cratered equatorial and southern highlands as
76 cratered uplands, and ejecta and uplifted blocks of ancient terrain caused by large impacts as
77 mountainous terrain. He further characterised the fretted uplands and isolated mesas along the
78 boundary between heavily cratered uplands and northern plains as knobby terrain. He also
79 highlighted three chronological volcanic plains, with Tharsis being the younger lava flows, the
80 rolling plains constituting the majority of Elysium Planitia as the intermediate age lava flows,
81 and ridged plains as the older lava plains. Baker (1981) further classified the northern plains as
82 a “complex lowland showing extensive evidence of ice-contact volcanism, permafrost features,
83 and aeolian modification”. Chaotic terrain or fractured terrain and valleys were also key
84 components of Baker’s (1981) Martian geomorphic map. Although with volumes of new
85 multisensory and higher resolution datasets, the mapping scale has improved severalfold,
86 undoubtedly, all the major geomorphic units, as presented by Baker (1981), are equally relevant
87 even today.




Figure 1. Global geomorphic map of Mars. This map is modified from the geological map by Tanaka et al. (2014). Data Source: <http://pubs.usgs.gov/sim/3292>.

90 As an interesting exercise, we adopted the similar approach as Baker's (1981) and compiled a
91 geomorphic map of major physiographic regions on Mars (Figure 1) using the data modified
92 from the most recent and complete geological map of Mars (Tanaka et al., 2014), available
93 from the US Geological Survey. This global dataset is a derived product of unprecedented
94 diversity (spectral, topographic, thermophysical, and subsurface), quality (high spatial and
95 spectral resolutions), and volume of remotely sensed data acquired since the Viking Orbiters.
96 In particular, the inclusion of precise topographic data such as Mars Global Surveyor (MGS)
97 Mars Orbiter Laser Altimeter (MOLA) digital elevation model (DEM) (463 m/pixel resolution
98 at lower latitudes to 115 m/pixel near the poles) (Smith and others, 2001) and the Mars Odyssey
99 (ODY) Thermal Emission Imaging System (THEMIS) daytime infrared (IR) image mosaic
100 (100 m/pixel) (Christensen and others, 2004) aided the visual interpretations greatly by
101 providing 3D terrain and infrared multispectral information (Tanaka et al., 2014). As
102 mentioned above, the map in Figure 1 is more detailed, understandably owing to the vastness
103 of the input data, but the close resemblance of the major physiographic classes with the ones
104 presented in Figure 1 of Baker (1981) are undeniable.

105 **IV. Involvement of geomorphologists in planetary sciences**

106 Baker's (1981) paper was the first review that discussed a very vital contemporary issue (on
107 Page 476 of Baker, 1981); "Why then have so few geomorphologists become involved in the
108 study of the fascinating landscapes of Mars?" With inclusion of the first two tables in his article,
109 Baker (1981) provided two main reasons for this: (1) the limited distribution of journals and
110 government documents publishing the planetary science papers, and (2) the lack of access to
111 the planetary surface images which could enable geomorphic interpretations. Baker (1981)
112 argued how increased involvement of geomorphologists could be great in promoting planetary
113 sciences as a discipline. There is no denying that in the past four decades, comparative
114 planetology has advanced as a science and the role of the geomorphological approaches in this

115 advancement has been undisputed. Not only the important themes mentioned by Baker (1981),
116 such as catastrophic flooding (e.g., Rodríguez et al., 2014), volcanism (e.g., Brož et al., 2020),
117 impact cratering (e.g., Palumbo and Head, 2018), aeolian erosion (e.g., Williams et al., 2020),
118 sapping (e.g., Goldspiel and Squyres, 2000), thermokarst (e.g., Dundas et al., 2015), and large-
119 scale landslides (e.g., Magnarini et al., 2019), have been greatly explored, but also, our
120 understanding of the effects of reduced gravity (e.g., Jacobsen and Burr, 2016), catastrophism
121 (Pacifci et al., 2009), and atmospheric processes (e.g., Matsubara et al., 2018) on Martian
122 landscape evolution has considerably improved. Nevertheless, the internet revolution leading
123 to digital information dissemination (Feldman, 2002) played a significant role in solving the
124 two aforementioned issues highlighted by Baker (1981). The online platform not only led to
125 the increase in the number of planetary journals, but also to the distribution of articles to an
126 interested reader. Moreover, the World Wide Web and the evolution in computing systems and
127 geographic information system software enabled easy and efficient data transfers, processing,
128 and interpretations.

129 **V. Major geomorphic features on Mars and conceptual advances**

130 Baker (1981) compiled the information on the geomorphic features on Mars and in the
131 following sub-sections, we revisit each of them briefly, providing some recent references to
132 assess the major conceptual advances.

133 *1. Impact craters*

134 Baker (1981) highlights how active resurfacing processes on Earth have erased the majority of
135 the ancient impact craters, making them relatively rare terrestrial landforms to perform
136 comparative planetology. Nevertheless, in past decades, several terrestrial craters have been
137 proposed and explored as Mars analogue sites around the globe. For example, Australia's arid
138 climate, coupled with the low-relief and tectonically stable terrain, has ensured best

139 preservation of several of the craters (West et al., 2010). Undeniably, the majority of these
140 craters have been identified on constantly improving mid-to-high resolution satellite images;
141 for example, the 260 m diameter Hickman Crater in Western Australia (Glikson et al., 2008).
142 Moving further, Baker (1981) provides perspectives on crater densities and ways to enable
143 relative and absolute dating of Martian surface. He further mentions the morphological
144 uniqueness of several of the Martian craters, in terms of being surrounded by layered debris
145 instead of ejecta of ballistic origin, and cites Mouginis-Mark (1979) to imply that the layered
146 debris might be a consequence of the entrainment of permafrost-melt water into the ejecta.
147 Now, with years of observational data, numerical modelling, and laboratory experiments, we
148 know that in addition to the subsurface volatiles, particle size and density, and atmospheric
149 density and pressure also contribute to the morphology of ejecta blankets (Barlow, 2005).
150 Moreover, we have further learnt about the astrobiological potential of the numerous small-to-
151 medium sized impact craters on Mars with clearly defined flat floors containing a possible
152 sedimentary record (Cockell and Lee, 2002; Lindsay and Brasier, 2006).

153 *2. Volcanic landforms*

154 Baker (1981) starts the discussion on volcanic structures by providing dimensional and
155 contextual information about the highest, i.e., Olympus Mons, and the widest, i.e., Alba Patera,
156 volcanoes on Mars. He further mentions the possible phreatic (explosive) phases in the early
157 eruptive history of large Martian volcanoes, owing to the eruptions through water-saturated (or
158 ice-rich) megaregolith materials (Greeley and Spudis, 1981). These explosive eruptions
159 probably changed to effusive lava production, constructing prominent shields and domes once
160 the megaregolith was depleted in water (Baker, 1981). Baker (1981) also cites Reimers and
161 Komar's (1979) hypothesis on the pyroclastic activity to be a result of volcanic interaction with
162 an ice-rich permafrost. However, the most recent and comprehensive review on this topic (Brož
163 et al., 2020) concludes several major points in this discussion of explosive volcanism on Mars.

164 Brož et al. (2020) concur that although, the indications of explosive volcanism have been
165 identified at various locations on Mars, the evidence is still less common than for effusive
166 activity. Brož et al. (2020) also infer that the possible explosive eruptions on Mars would have
167 behaved differently from those on Earth, since the observed edifices are often different in
168 shapes from their terrestrial counterparts. Baker (1981) mentions the presence of pseudocraters
169 and pedestal craters in the northern plains of Mars. There still is an uncertainty on the exact
170 formation mechanism of these landforms. The pseudocraters are now categorised as rootless
171 cones and both igneous and mud volcano hypotheses are proposed as their formation
172 mechanism (Czechowski et al., 2020; Dapremont and Wray, 2021). Similarly, the pedestal
173 craters, even today, are believed to be a result of presence of ice-rich layers during their
174 formation, when the ejecta formed an erosion-resistant layer shielding the surroundings
175 (Kenkmann and Wulf, 2018). Baker (1981) cites Hodges and Moore (1979) in proposing the
176 table mountains of Iceland as a possible analogue for the Martian pedestal craters.

177 *3. Aeolian landforms*

178 Aeolian landforms are one of those surficial features on Mars which highlight the effect of
179 improvement in spatial resolution of the imaging camera on advancing our geomorphic
180 knowledge. During the 1970s, available coarser resolution images, captured by Mariner 9
181 (McCauley et al., 1972) and Viking Orbiters (Cutts et al., 1976), revealed large, low albedo
182 dune masses, now known as large dark dunes (LDDs). Baker (1981) mentions these huge dark-
183 coloured dunes surrounding the northern polar cap of Mars. With the advent of high-resolution
184 imagers, such as the Mars Orbiter Camera (MOC) (Malin and Edgett, 2001), High Resolution
185 Stereo Camera (Neukum and Jaumann, 2004), Context (CTX) camera (Malin et al., 2007), and
186 HiRISE (McEwen et al., 2007), during the 1990s and 2000s, the captured m-to-cm resolution
187 images made it possible to observe and study smaller aeolian landforms such as wind ripples,
188 granule ripples, yardangs, dust devils, ventifacts, and transverse aeolian ridges (TARs)

189 (Bhardwaj et al., 2019c). An updated account of aeolian landforms on Mars can be read in
190 Bridges et al. (2013).

191 *4. Hillslopes and mass movement*

192 In his comprehensive and interesting review, Brunsten (1993) appropriately highlighted the
193 complexity involved in discussing mass movement as an isolated discipline. Studying mass
194 movements mandatorily needs an interdisciplinary approach involving geomorphology,
195 geology, hydrology, geophysics, and soil mechanics (Brunsten, 1993). This makes
196 understanding the nature of mass movements even more complex for the places (e.g., Mars)
197 where sufficient multidisciplinary data are unavailable. Nevertheless, Baker (1981) relatively
198 effectively uses published contemporary examples of the Valles Marineris system (e.g.,
199 Lucchitta, 1979; Sharp, 1973) to highlight the typical “spur-and-gully topography” (Lucchitta,
200 1978) defining numerous steep escarpments and hillslopes on Mars. Such complex topography
201 is prone to produce an immense array of mass movement features (Baker, 1981), some of which
202 are difficult to interpret and characterise even today (e.g., Bhardwaj et al., 2019a; Bhardwaj et
203 al., 2019d). Undeniably, the unavailability of high-resolution multisource datasets for the
204 majority of the Martian terrain makes the interpretations even more speculative. Baker (1981)
205 also compiled published examples to emphasise the relatively massive dimensions of mass
206 movements on Mars, undetected on Earth.

207 Interestingly, one of the possible mass movement features, which Baker (1981) refers to as
208 lobate debris deposits and discusses towards the end of this section, are extensively investigated
209 in the past two decades. These lobate debris deposits resemble terrestrial glaciers, with valleys
210 filled with debris, in some instances originating from cirque-like heads, and locally marked by
211 prominent longitudinal ridges. Squyres (1978; 1979) correlated the lobate debris deposits with
212 regions of probable high frost deposition and proposed their possible analogy with terrestrial

213 rock glaciers (Baker, 1981). All these glacial-type formations on Mars, displaying evidence of
214 viscous flow, are now characterised within an umbrella term called Viscous flow feature (VFF)
215 (Souness et al., 2012). Four major types of VFFs are identified and studied (Hubbard et al.,
216 2011; Souness et al., 2012; Squyres, 1978, Squyres, 1979; Squyres and Carr, 1986): (1) lobate
217 debris aprons (LDA), (2) lineated valley fill (LVF), (3) concentric crater fill (CCF), and (4)
218 glacier-like forms (GLF). Interested reader can find an updated account of VFFs in Berman et
219 al. (2021). Koutnik and Pathare (2021) recently presented an informative and updated account
220 of LDA and GLF in terms of their analogy with terrestrial debris-covered glaciers. Their review
221 can be helpful in providing a holistic account of analogy between debris-covered glaciers on
222 Earth and dust and debris-covered ice on Mars.

223 *5. Periglacial and permafrost features*

224 Although morphologically, VFFs qualify to be characterised as permafrost landforms often
225 observed in periglacial landscape on Earth, Baker (1981) at the very onset of this section,
226 defines “periglacial” as a geomorphic environment categorised by very low annual
227 temperatures, freeze-thaw episodes, and strong wind action. Baker (1981) further defines the
228 term “permafrost” as used in comparative planetology. Although “permafrost” refers to frozen
229 ground, irrespective of its water content, in planetology, the term is often used as a synonym
230 for “ground ice” (Baker, 1981). Baker (1981) starts by discussing polygonal fracture patterns
231 which are typical of permafrost terrain. For the initially observed fractures in Martian northern
232 plains, the two most probable proposed mechanisms were permafrost ice wedging, and cooling-
233 contraction cracking in lava flows. However, the massive dimensions of these cracks (hundreds
234 of metres wide with average spacings of 5-10 km) put constraint on both these hypotheses.
235 Terrestrial ice-wedge polygons generally vary from 1-100 m in diameter (Baker, 1981), and
236 with HiRISE images, today many regions on Mars have been identified with polygons
237 comparable in dimensions to their terrestrial counterparts (Soare et al., 2021). Baker (1981)

238 further discusses thermokarst landforms and scalloped terrain on Mars and highlights the
239 contemporary views of thermokarst landforms being a result of melting ground ice. However,
240 in the past couple of decades, both sublimation (e.g., Dundas et al., 2015) and melting (e.g.,
241 Soare et al., 2008), have been investigated as the formation mechanisms for these thermokarst
242 landforms on Mars.

243 *6. Polar terrains*

244 Baker (1981) starts this section with describing Mariner and Viking observations of Martian
245 polar caps. Based on the temperatures observed by the infrared radiometers of the Viking
246 orbiters, Baker (1981) asserts that the “northern cap must be water ice”. Subsequent
247 multisensory observations have validated the residual ice caps to be primarily consisting of
248 water ice. A recent paper (Ojha et al., 2019) placed compositional constraint on the polar
249 silicate-rich basal unit below the ice-rich north polar layered deposit, by modelling its gravity
250 signature in both spatial and spectral domains. These estimates suggest that even the silicate-
251 rich basal unit below the polar layered deposits may contain $55\pm 25\%$ water ice, corresponding
252 to ~ 1.5 m global water equivalent, making it one of the largest reservoirs of water-ice on Mars
253 (Ojha et al., 2019).

254 *7. Channels and valleys*

255 Baker (1981) starts this section by highlighting the excitement that was linked with the
256 discovery of channels, valleys, and related features of possible aqueous origin on Mars.
257 However, like any other hypothesis of possible liquid water on Mars, this discovery was also
258 not without controversies, and soon, in addition to water, lava flow, wind, glacial ice,
259 liquefaction of crustal materials, debris flows, liquid carbon dioxide, and liquid alkanes were
260 suggested as other possible channel-carving agents (Baker, 1981). To characterise the widely
261 variable channel morphologies, Masursky (1973) adopted a broader context and proposed four

262 classes: (1) broad large-sized channels originating from chaotic terrain, (2) narrow
263 intermediate-sized channels, (3) small valleys across the heavily cratered terrain, and (4)
264 volcanic channels (Baker, 1981). Sharp and Malin (1975) proposed an additional category
265 called fretted channels, in addition to Masursky's categories (Baker, 1981). Using high-
266 resolution MOC images of channels and valleys, Malin and Edgett (2003) provided
267 geomorphic evidence for aqueous sedimentation on early Mars. Mangold et al. (2004)
268 interpreted the geomorphic characteristics, especially the high degree of branching, of the
269 valleys in Valles Marineris region, to propose atmospheric precipitation during 2.9 to 3.4
270 billion years as their formation mechanism. However, a recent paper (Galofre et al., 2020), puts
271 a constraint on entirely precipitation and surface water runoff-based hypotheses for valley
272 formations on Mars and proposes subglacial and fluvial erosion as the predominant
273 mechanisms.

274 **VI. Summary**

275 As evident from revisiting Baker's (1981) paper, the relevance and impact of geomorphology
276 as a discipline in progressing comparative planetology in general, and Mars landscape research
277 in particular, are indisputable. The timely compilation and survey of contemporary literature
278 following Viking Missions, and raising the awareness of geomorphology community on Mars
279 exploration, are the highpoints of this first holistic review of Martian geomorphology presented
280 by Baker (1981). With nearly the entire Martian terrain covered at ~6 m/pixel CTX resolution,
281 and continuously increasing volume of submeter HiRISE data, undoubtedly, the prospects for
282 performing comprehensive local-scale geomorphic analyses have considerably improved.
283 Moreover, the operational rovers on Mars are also providing stereoscopic and multispectral
284 images. In fact, the perceived success of the first unmanned aerial vehicle on Mars, in the form
285 of a mini-helicopter named Ingenuity onboard the recently landed Perseverance Rover, as an

286 image capture platform can really transform the next stage of exploring Martian
287 geomorphology.

288 **Declaration of conflicting interests**

289 The author(s) declared no potential conflicts of interest with respect to the research, authorship,
290 and/or publication of this article.

291 **Acknowledgement**

292 The authors acknowledge the encouragement, helpful suggestions, and support from Prof.
293 David R. Butler.

294 **References**

295 Baker, V. R. (1981). The geomorphology of Mars. *Progress in Physical Geography*, 5(4), 473-
296 513.

297 Barlow, N. G. (2005). A review of Martian impact crater ejecta structures and their implications
298 for target properties. *Large meteorite impacts III*, Geological Society of America Special Paper,
299 384, 433-442.

300 Berman, D. C., Chuang, F. C., Smith, I. B., & Crown, D. A. (2021). Ice-rich landforms of the
301 southern mid-latitudes of Mars: A case study in Nereidum Montes. *Icarus*, 355, 114170.

302 Bhardwaj, A., Sam, L., Akanksha, Martín-Torres, F. J., & Kumar, R. (2016). UAVs as remote
303 sensing platform in glaciology: Present applications and future prospects. *Remote sensing of*
304 *environment*, 175, 196-204.

305 Bhardwaj, A., Sam, L., Martín-Torres, F. J., & Zorzano, M. P. (2019a). Are Slope Streaks
306 Indicative of Global-Scale Aqueous Processes on Contemporary Mars?. *Reviews of*
307 *Geophysics*, 57(1), 48-77.

308 Bhardwaj, A., Sam, L., Martin-Torres, F. J., & Zorzano, M. P. (2019c). Distribution and
309 morphologies of transverse Aeolian ridges in ExoMars 2020 Rover landing site. *Remote*
310 *Sensing*, 11(8), 912.

311 Bhardwaj, A., Sam, L., Martin-Torres, J., & Mier, M. P. Z. (2019d). Revisiting Enigmatic
312 Martian Slope Streaks. *Earth and Space Science*, 100, 100.

313 Bhardwaj, A., Sam, L., Martín-Torres, F. J., Zorzano, M. P., & Ramírez Luque, J. A. (2019b).
314 UAV imaging of a Martian brine analogue environment in a fluvio-aeolian setting. *Remote*
315 *Sensing*, 11(18), 2104.

316 Bridges, N., Geissler, P., Silvestro, S., & Banks, M. (2013). Bedform migration on Mars:
317 Current results and future plans. *Aeolian Research*, 9, 133-151.

318 Brož, P., Bernhardt, H., Conway, S. J., & Parekh, R. (2020). An overview of explosive
319 volcanism on Mars. *Journal of Volcanology and Geothermal Research*, 107125.

320 Brunsten, D. (1993). Mass movement; the research frontier and beyond: a geomorphological
321 approach. *Geomorphology*, 7(1-3), 85-128.

322 Christensen, P.R., Jakosky, B.M., Kieffer, H.H., and 8 others, 2004, The Thermal Emission
323 Imaging System (THEMIS) for the Mars 2001 Odyssey mission: *Space Science Reviews*, v.
324 110, p. 85–130.

325 Cockell, C. S., & Lee, P. (2002). The biology of impact craters—a review. *Biological Reviews*,
326 77(3), 279-310.

327 Cutts, J.A.; Blasius, K.R.; Briggs, G.A.; Carr, M.H.; Greeley, R.; Masursky, H. North polar
328 region of Mars: Imaging results from Viking 2. *Science* 1976,194, 1329–1337

329 Czechowski, L., Zalewska, N., Zambrowska, A., Ciazela, M., Witek, P., & Kotlarz, J. (2020,
330 September). Mechanism of Origin of Chains of Cones in Crise PLanitia. In European Planetary
331 Science Congress (pp. EPSC2020-895).

332 Dapremont, A. M., & Wray, J. J. (2021). Igneous or Mud Volcanism on Mars? The Case Study
333 of Hephaestus Fossae. *Journal of Geophysical Research: Planets*, 126(2), e2020JE006390.

334 Dundas, C. M., Byrne, S., & McEwen, A. S. (2015). Modeling the development of Martian
335 sublimation thermokarst landforms. *Icarus*, 262, 154-169.

336 Edgett, K. S., Williams, R. M., Malin, M. C., Cantor, B. A., & Thomas, P. C. (2003). Mars
337 landscape evolution: Influence of stratigraphy on geomorphology in the north polar region.
338 *Geomorphology*, 52(3-4), 289-297.

339 Feldman, M. P. (2002). The Internet revolution and the geography of innovation. *International*
340 *Social Science Journal*, 54(171), 47-56.

341 Gaffey, C., & Bhardwaj, A. (2020). Applications of unmanned aerial vehicles in cryosphere:
342 Latest advances and prospects. *Remote Sensing*, 12(6), 948.

343 Galofre, A. G., Jellinek, A. M., & Osinski, G. R. (2020). Valley formation on early Mars by
344 subglacial and fluvial erosion. *Nature Geoscience*, 13(10), 663-668.

345 Gilbert, G.K. (1893) The moon's face: a study of the origin of its features. *Philosophical*
346 *Society of Washington Bulletin* 12, 241-92.

347 Glikson, A. Y., Hickman, A. H., & Vickers, J. (2008). Hickman Crater, Ophthalmia Range,
348 Western Australia: evidence supporting a meteorite impact origin. *Australian Journal of Earth*
349 *Sciences*, 55(8), 1107-1117.

350 Goldspiel, J. M., & Squyres, S. W. (2000). Groundwater sapping and valley formation on Mars.
351 *Icarus*, 148(1), 176-192.

352 Greeley, R. and Spudis, P.D. 1981: Volcanism on Mars. *Reviews of Geophysics and Space*
353 *Physics* 19, 13-41.

354 Hodges, C.A. and Moore, H.J. 1979: The subglacial birth of Olympus Mons and its aureoles.
355 *Journal of Geophysical Research* 84, 8061-74.

356 Hubbard, B., Milliken, R.E., Kargel, J.S., Limaye, A., Souness, C., 2011. Geomorphologi-cal
357 characterisation and interpretation of a mid-latitude glacier-like form: Hellas Planitia, Mars.
358 *Icarus* 211 (1), 330–346.

359 Irwin III, R. P., Howard, A. D., & Maxwell, T. A. (2004). Geomorphology of Ma'adim Vallis,
360 Mars, and associated paleolake basins. *Journal of Geophysical Research: Planets*, 109(E12).

361 Jacobsen, R. E., & Burr, D. M. (2016). Greater contrast in Martian hydrological history from
362 more accurate estimates of paleodischarge. *Geophysical Research Letters*, 43(17), 8903-8911.

363 Kenkmann, T., & Wulf, G. (2018). Impact Cratering. In *Planetary Geology* (pp. 123-145).
364 Springer, Cham.

365 Koutnik, M.R. and Pathare, A.V., 2021. Contextualizing lobate debris aprons and glacier-like
366 forms on Mars with debris-covered glaciers on Earth. *Progress in Physical Geography: Earth*
367 *and Environment* 45: 130-186.

368 Lindsay, J., & Brasier, M. (2006). Impact craters as biospheric microenvironments, Lawn Hill
369 structure, northern Australia. *Astrobiology*, 6(2), 348-363.

370 Lucchitta, B.K. (1978). Morphology of chasma walls, Mars. *Journal of Research of the United*
371 *States Geological Survey* 6, 651-62.

372 Lucchitta, B.K. (1979). Landslides in Valles Marineris, Mars. *Journal of Geophysical Research*
373 84, 8097-8113.

374 Magnarini, G., Mitchell, T. M., Grindrod, P. M., Goren, L., & Schmitt, H. H. (2019).
375 Longitudinal ridges imparted by high-speed granular flow mechanisms in martian landslides.
376 Nature communications, 10(1), 1-7.

377 Malin, M.C.; Bell, J.F.; Cantor, B.A.; Caplinger, M.A.; Calvin, W.M.; Clancy, R.T.; Edgett,
378 K.S.; Edwards, L.; Haberle, R.M.; James, P.B.; et al. Context Camera Investigation on board
379 the Mars Reconnaissance Orbiter. J. Geophys. Res. Planets 2007,112.

380 Malin, M.C.; Edgett, K.S. Mars Global Surveyor Mars Orbiter Camera: Interplanetary cruise
381 through primary mission. J. Geophys. Res. 2001,106, 23429–23570.

382 Malin, M. C., & Edgett, K. S. (2003). Evidence for persistent flow and aqueous sedimentation
383 on early Mars. Science, 302(5652), 1931-1934.

384 Mangold, N., Quantin, C., Ansan, V., Delacourt, C., & Allemand, P. (2004). Evidence for
385 precipitation on Mars from dendritic valleys in the Valles Marineris area. Science, 305(5680),
386 78-81.

387 Masursky, H. 1973: An overview of geological results from Mariner 9. Journal of Geophysical
388 Research 78, 4009-30.

389 Matsubara, Y., Howard, A. D., & Irwin III, R. P. (2018). Constraints on the Noachian
390 paleoclimate of the Martian highlands from landscape evolution modeling. Journal of
391 Geophysical Research: Planets, 123(11), 2958-2979.

392 McCauley, J.F.; Carr, M.H.; Cutts, J.A.; Hartmann, W.K.; Masursky, H.; Milton, D.J.; Sharp,
393 R.P.; Wilhelms, D.E. Preliminary Mariner 9 report on the geology of Mars. Icarus1972,17,
394 289–327.

395 McEwen, A.S.; Eliason, E.M.; Bergstrom, J.W.; Bridges, N.T.; Hansen, C.J.; Delamere, W.A.;
396 Grant, J.A.; Gulick, V.C.; Herkenhoff, K.E.; Keszthelyi, L.; et al. Mars reconnaissance

397 orbiter's high resolution imaging science experiment (HiRISE). *J. Geophys. Res. Planets*
398 2007,112.

399 Mouginis-Mark, P.J. 1979: Martian fluidized crater morphology: variations with crater size,
400 latitude, altitude, and target material. *Journal of Geophysical Research* 84, 8011-22.

401 Mutch, T.A. (1979). Planetary surfaces. *Reviews of Geophysics and Space Physics* 17, 1694-
402 1722.

403 Neukum, G.; Jaumann, R. HRSC: The High Resolution Stereo Camera of Mars Express, Mars
404 Express: The Scientific Payload; Wilson, A., Chicarro, A., Eds.; ESA Publications Division:
405 Noordwijk, The Netherlands, 2004; pp. 17–35, ISBN 92-9092-556-6.

406 Ojha, L., Nerozzi, S., & Lewis, K. (2019). Compositional constraints on the north polar cap of
407 Mars from gravity and topography. *Geophysical Research Letters*, 46(15), 8671-8679.

408 Pacifici, A., Komatsu, G., & Pondrelli, M. (2009). Geological evolution of Ares Vallis on Mars:
409 Formation by multiple events of catastrophic flooding, glacial and periglacial processes. *Icarus*,
410 202(1), 60-77.

411 Palumbo, A. M., & Head, J. W. (2018). Impact cratering as a cause of climate change, surface
412 alteration, and resurfacing during the early history of Mars. *Meteoritics & Planetary Science*,
413 53(4), 687-725.

414 Riemers, C.E. and Komar, P.D. 1979: Evidence for explosive volcanic density currents on
415 certain Martian volcanoes. *Icarus* 39, 88-110.

416 Rodríguez, J.A.P., Gulick, V.C., Baker, V.R., Platz, T., Fairén, A.G., Miyamoto, H., Kargel,
417 J.S., Rice, J.W. and Glines, N., 2014. Evidence for Middle Amazonian catastrophic flooding
418 and glaciation on Mars. *Icarus*, 242, pp.202-210.

419 Sam, L., Bhardwaj, A., Singh, S., & Martin-Torres, F. J. (2020a). UAV Imaging of Small Caves
420 in Icelandic Lava Field as Possible Mars Analogues. In 3rd International Planetary Caves
421 Conference (Vol. 2197, p. 1053).

422 Sam, L., Bhardwaj, A., Singh, S., Martin-Torres, F. J., Zorzano, M. P., & Ramírez Luque, J.
423 A. (2020b). Small lava caves as possible exploratory targets on Mars: Analogies drawn from
424 UAV imaging of an Icelandic lava field. *Remote Sensing*, 12(12), 1970.

425 Schulze-Makuch, D., Méndez, A., Fairén, A. G., von Paris, P., Turse, C., Boyer, G., Davila, A.
426 F., de Sousa Antonio, M. R., Catling, D., & Irwin, L. N. (2011). A two-tiered approach to
427 assessing the habitability of exoplanets. *Astrobiology*, 11(10), 1041–1052.

428 Scott, D.H. and Carr, M.H. 1978: Geologic map of Mars. United States Geological Survey
429 Miscellaneous Geologic Investigations Map I-1083.

430 Sharp, R.P. (1973): Mars: troughed terrain. *Journal of Geophysical Research* 78, 4063-72.

431 Sharp, R.P. (1980). Geomorphological processes on terrestrial planetary surfaces. *Annual*
432 *Reviews of Earth and Planetary Sciences* 8, 231-61.

433 Sharp, R.P. and Malin, M.C. 1975: Channels on Mars. *Geological Society of America Bulletin*
434 86, 593-609.

435 Smith, D.E., Zuber, M.T., Frey, H.V., and 21 others, 2001, Mars Orbiter Laser Altimeter—
436 Experiment summary after the first year of global mapping of Mars: *Journal of Geophysical*
437 *Research*, v. 106, no. E10, p.23,689–23,722.

438 Soare, R. J., Conway, S. J., Williams, J. P., Philippe, M., Mc Keown, L. E., Godin, E., &
439 Hawksell, J. (2021). Possible ice-wedge polygonisation in Utopia Planitia, Mars and its
440 latitudinal gradient of distribution. *Icarus*, 358, 114208.

441 Soare, R. J., Osinski, G. R., & Roehm, C. L. (2008). Thermokarst lakes and ponds on Mars in
442 the very recent (late Amazonian) past. *Earth and Planetary Science Letters*, 272(1-2), 382-393.

443 Souness, C., Hubbard, B., Milliken, R. E., & Quincey, D. (2012). An inventory and population-
444 scale analysis of martian glacier-like forms. *Icarus*, 217(1), 243-255.

445 Squyres, S.W. 1978: Martian fretted terrain: flow of erosional debris. *Icarus* 34, 600-613.

446 Squyres, S.W. 1979: The distribution of lobate debris aprons and similar flows on Mars.
447 *Journal of Geophysical Research* 84, 8087-96.

448 Squyres S. W. and Carr M. H. (1986) Geomorphic evidence for the distribution of ground ice
449 on Mars. *Science*, 231, 249-252.

450 Tanaka, K.L., J.A. Skinner, Jr., J.M. Dohm, R.P. Irwin, III, E.J. Kolb, C.M. Fortezzo, Thomas
451 Platz, G.G. Michael, and T.M. Hare, 2014, Geologic Map of Mars, Scale 1:20,000,000, U.S.
452 Geological Survey Scientific Investigations Map SIM 3292. URL:
453 <http://pubs.usgs.gov/sim/3292>

454 Tsibulskaya, V., Hepburn, A. J., Hubbard, B., & Holt, T. (2020). Surficial geology and
455 geomorphology of Greg crater, Promethei Terra, Mars. *Journal of Maps*, 16(2), 524-533.

456 West, M. D., Clarke, J. D., Thomas, M., Pain, C. F., & Walter, M. R. (2010). The geology of
457 Australian Mars analogue sites. *Planetary and Space Science*, 58(4), 447-458.

458 Williams, J., Day, M., Chojnacki, M., & Rice, M. (2020). Scarp orientation in regions of active
459 aeolian erosion on Mars. *Icarus*, 335, 113384.

460 Wood, L. J. (2006). Quantitative geomorphology of the Mars Eberswalde delta. *Geological*
461 *Society of America Bulletin*, 118(5-6), 557-566.