

2021

Hepatotoxicity reports in the FDA adverse event reporting system database: A comparison of drugs that cause injury via mitochondrial or other mechanisms

Payal Rana

Michael D. Aleo

Xuerong Wen

University of Rhode Island, xuerongwen@uri.edu

Stephen Jon Kogut

University of Rhode Island, skogut@uri.edu

Follow this and additional works at: https://digitalcommons.uri.edu/php_facpubs

Citation/Publisher Attribution

Rana, P., Aleo, M. D., Wen, X., & Kogut, S. (2021). Hepatotoxicity reports in the FDA adverse event reporting system database: A comparison of drugs that cause injury via mitochondrial or other mechanisms. *Acta Pharmaceutica Sinica B*. <https://doi.org/10.1016/j.apsb.2021.05.028>


Available at: <https://doi.org/10.1016/j.apsb.2021.05.028>

This Article is brought to you for free and open access by the Pharmacy Practice at DigitalCommons@URI. It has been accepted for inclusion in Pharmacy Practice Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons@etal.uri.edu.

Journal Pre-proof

Hepatotoxicity reports in the FDA adverse event reporting system database: A comparison of drugs that cause injury via mitochondrial or other mechanisms

Payal Rana, Michael D. Aleo, Xuerong Wen, Stephen Kogut



PII: S2211-3835(21)00207-0

DOI: <https://doi.org/10.1016/j.apsb.2021.05.028>

Reference: APSB 1116

To appear in: *Acta Pharmaceutica Sinica B*

Received Date: 24 December 2020

Revised Date: 10 May 2021

Accepted Date: 24 May 2021


Please cite this article as: Rana P, Aleo MD, Wen X, Kogut S, Hepatotoxicity reports in the FDA adverse event reporting system database: A comparison of drugs that cause injury via mitochondrial or other mechanisms, *Acta Pharmaceutica Sinica B*, <https://doi.org/10.1016/j.apsb.2021.05.028>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Chinese Pharmaceutical Association and Institute of Materia Medica, Chinese Academy of Medical Sciences. Production and hosting by Elsevier B.V. All rights reserved.

GA

In this study, we investigated liver injury reports submitted to the FAERS database and compared the frequency of reports between drugs that can cause hepatotoxicity *via* mitochondrial mechanisms and those without mitochondrial mechanisms of toxicity.



Original article**Hepatotoxicity reports in the FDA adverse event reporting system database: A comparison of drugs that cause injury *via* mitochondrial or other mechanisms****Payal Rana^{a,*}, Michael D. Aleo^a, Xuerong Wen^b, Stephen Kogut^b**^a*Drug Safety Research & Development, Pfizer, Groton, CT 06340, USA*^b*University of Rhode Island, College of Pharmacy, Kingston, RI 02881, USA*

Received 24 December 2020; received 2 March 2021; accepted 19 May 2021

*Corresponding author. Tel.: +1 0 715 6154.

E-mail address: payal.m.rana@pfizer.com (Payal Rana).

Running title: Hepatotoxicity reports in the FDA adverse event reporting system database

Abstract Drug-induced liver injury (DILI) is a leading reason for preclinical safety attrition and post-market drug withdrawals. Drug-induced mitochondrial toxicity has been shown to play an essential role in various forms of DILI, especially in idiosyncratic liver injury. This study examined liver injury reports submitted to the Food and Drug Administration (FDA) Adverse Event Reporting System (FAERS) for drugs associated with hepatotoxicity *via* mitochondrial mechanisms compared with non-mitochondrial mechanisms of toxicity. The frequency of hepatotoxicity was determined at a group level and individual drug level. A reporting odds ratio (ROR) was calculated as the measure of effect. Between the two DILI groups, reports for DILI involving mitochondrial mechanisms of toxicity had a 1.43 (95% CI 1.42–1.45; $P < 0.0001$) times higher odds compared to drugs associated with non-mitochondrial mechanisms of toxicity. Antineoplastic, antiviral, analgesic, antibiotic, and antimycobacterial drugs were the top 5 drug classes with the highest ROR values. Although the top 20 drugs with the highest ROR values included drugs with both mitochondrial and non-mitochondrial injury mechanisms, the top 4 drugs (ROR values >18 : benzbromarone, troglitazone, isoniazid, rifampin) were associated with mitochondrial mechanisms of toxicity. The major demographic influence for DILI risk was also examined. There was a higher mean patient age among reports for drugs that were associated with mitochondrial mechanisms of toxicity [56.1 ± 18.33 (SD)] compared to non-mitochondrial mechanisms [48 ± 19.53 (SD)] ($P < 0.0001$), suggesting that age may play a role in susceptibility

to DILI *via* mitochondrial mechanisms of toxicity. Univariate logistic regression analysis showed that reports of liver injury were 2.2 (odds ratio: 2.2, 95% CI 2.12–2.26) times more likely to be associated with older patient age, as compared with reports involving patients less than 65 years of age. Compared to males, female patients were 37% less likely (odds ratio: 0.63, 95% CI 0.61–0.64) to be subjects of liver injury reports for drugs associated with mitochondrial toxicity mechanisms. Given the higher proportion of severe liver injury reports among drugs associated with mitochondrial mechanisms of toxicity, it is essential to understand if a drug causes mitochondrial toxicity during preclinical drug development when drug design alternatives, more clinically relevant animal models, and better clinical biomarkers may provide a better translation of drug-induced mitochondrial toxicity risk assessment from animals to humans. Our findings from this study align with mitochondrial mechanisms of toxicity being an important cause of DILI, and this should be further investigated in real-world studies with robust designs.

KEY WORDS Mitochondrial toxicity; FAERS database; Adverse event reporting; Drug-induced liver injury; Hepatotoxicity

Abbreviations: AE, adverse event; CI, confidence interval; CNS, center nervous system; DILI, drug-induced liver injury; DNA, deoxyribonucleic acid; FAERS, FDA's Adverse Event Reporting System; FDA, US Food and Drug Administration; MedDRA, Medical Dictionary for Regulatory Activities; NCTR-LTKB, National Center for Toxicological Research-Liver Toxicity Knowledge Base; NSAID, nonsteroidal anti-inflammatory drugs; ROR, Reporting Odds Ratio

1. Introduction

Severe drug-induced liver injury (DILI) is a rare, potentially life-threatening adverse event with an incidence of 10–15 cases per 10,000 to 100,000 patients per year¹⁻⁴. The clinical profile of DILI is challenging to diagnose as it can mimic almost any type of liver disease^{5,6} and mild, asymptomatic transaminase elevations may mimic those caused by diet⁷. Prediction of liver injury remains a challenge for the pharmaceutical industry, regulators, and clinicians⁵. DILI symptoms range from non-specific mild elevations in liver enzymes (aminotransferases) to severe liver illnesses such as cirrhosis or acute hepatitis^{5,6} and there is no specific biomarker that links drug exposure as the contributing cause of liver injury⁸.

There are two types of DILI: intrinsic (*i.e.*, dose-dependent) and idiosyncratic (*i.e.*, dose-independent)⁶. A well-known example of intrinsic DILI is acetaminophen-induced liver injury, as it is dose-dependent, the onset is hours to days, and predictable⁹. On the other hand, idiosyncratic DILI is not dose-dependent, occurs in a small portion of drug-exposed individuals, its onset is days to weeks, and unpredictable⁹. Idiosyncratic DILI is highly dependent on environmental and host factors that alter the susceptibility of individual patient responses to the drug. Hamilton et al.¹⁰ suggested that DILI is the convergence of three influencing risk factors: host factors, environmental factors, and drug-specific factors. Host-related risk factors include genetics, ethnicity, gender, comorbidities, alterations in drug transport, drug clearance capabilities, age, and mitochondrial function variability¹⁰. Environmental risk factors include lifestyle, viral co-infection, co-prescriptions, diet, and alcohol consumption¹⁰. Finally, drug-specific risk factors include the relationship of applied dose (exposure) and chemical structure with reactive metabolite formation, mitochondrial dysfunction, and lipophilicity¹⁰⁻¹². Therefore, mechanisms of DILI, whether intrinsic or idiosyncratic, are a multivariable, highly complex process that varies from patient to patient and is influenced by host, environmental, and drug-specific factors that influence the liver's ability to adapt and recover from an injury caused by a drug⁸.

In recent years, drug-induced mitochondrial toxicity has been shown to play an essential role in intrinsic and idiosyncratic DILI. Many medications from different drug classes, such as antidiabetic, antilipidemic, antiviral, antibiotic, anti-inflammatory, and antipsychotic agents have toxicities mediated by mitochondrial mechanisms, which may contribute to DILI¹³. Mitochondrial toxicants affect mitochondrial homeostasis by numerous mechanisms such as oxidative stress, inhibition or uncoupling of respiratory complexes of electron transport chain, impairment of mitochondrial replication or promoting mitochondrial DNA damage¹⁴. Drug-induced mitochondrial toxicity is difficult to be detected in standard preclinical animal testing models and requires specific studies to examine disruptions in liver energy status¹⁵. Only recently, there has been the development of clinical biomarkers specific for mitochondrial dysfunction in disease¹⁶ and DILI¹⁷ beyond changes in blood lactate. With these inadequacies, a drug candidate can enter human clinical trials only to fail for evidence of mitochondrial toxicity^{18,19}. Examples of non-mitochondrial toxicity mechanisms that drive DILI are generation of reactive metabolites, activation of cell death pathways, activation of innate or adaptive

immune response pathways, or disruption of cellular homeostasis²⁰. This study evaluates the frequency of reports of hepatotoxicity injury in drugs that cause DILI with mitochondrial and non-mitochondrial mechanisms.

Patient demographics influence risk or susceptibility for DILI. Boelsterli and Lim²¹ indicated that older age and female gender were important susceptibility factors for DILI; however, the reasons were still unknown. There are no clinical studies that link the sensitivity of the female gender to DILI caused by mitochondrial dysfunction. Amacher et al.²² indicated that women are more susceptible to DILI than men. Several hypotheses were proposed to explain gender differences in susceptibility, including pharmacokinetic or pharmacodynamic differences, interactions of sex hormones with signaling molecules, and a difference in immune system responses²². Similarly, it is believed that older adults are more susceptible to DILI caused by mitochondrial dysfunction. The review published by Will et al.¹³ indicated that the most commonly used prescription and over-the-counter medications for geriatric patients, such as antilipidemic, pain, and heartburn medications, had published reports of toxicities linked to mitochondrial dysfunction¹³. As the United States' elderly population is growing rapidly, identifying and addressing risk factors of DILI, where mitochondrial dysfunction may play a substantial role in adverse events, will be beneficial to this vulnerable patient population. Therefore, in this study, we evaluated the patient age and gender associated with DILI reports (measured by reporting odds ratio) for hepatotoxicants with mitochondrial and non-mitochondrial injury mechanisms.

Given that mitochondrial dysfunction is a common characteristic of drugs that cause liver injury, a better understanding of the association between the probability of liver injury induced by drugs that are mitochondrial toxicants and the influence of patient's age and gender would be beneficial for clinicians and drug developers. If a drug is associated with mitochondrial mechanisms of liver injury, clinicians could incorporate mitochondrial injury-specific biomarkers into clinical trials²³⁻²⁵. Additionally, the development of clinically relevant animal models or study designs may provide drug-induced mitochondrial toxicity risk translation from animals to humans²⁴.

This study investigated liver injury reports submitted to the US Food and Drug Administration (FDA) Adverse Event Reporting System (FAERS) and compared the frequency

of reports between drugs that can cause hepatotoxicity *via* mitochondrial mechanisms and those without mitochondrial mechanisms of toxicity. Additionally, we determined if there were age and gender differences associated with DILI reports involving drugs with or without mitochondrial toxicities.

2. Methods

2.1. Study design

2.1.1. Data source

FAERS database is the largest national repository of spontaneous drug event reports, having accumulated over 28 million adverse events reports. Healthcare professionals, patients, manufacturers, and lawyers can submit potential drug-induced adverse events of small and large therapeutic (biologics) classes and medical devices to the FAERS database. The FAERS database has a vital role in post-market drug surveillance in terms of detection and characterization of drug and device-related adverse events.

We extracted adverse event reports from the FAERS database for the timeframe from January 1998 to May 2019. In this study, the reports included severe adverse events, such as hospitalization, disability, or death. The types of reports were classified by FAERS as direct, expedited, or periodic. Direct reports were submitted to FDA from consumer or health care professionals; whereas, expedited reports were sent from the manufacturer within 15 days of severe adverse events occurrence not included in the product label²⁶. Adverse drug event reports considered periodic were submitted from manufacturers, included in the label, and sent to the FDA quarterly or annually²⁶. The main selection criterion was “primary suspect” drugs. “secondary suspect” drugs were excluded because of the greater uncertainty of the association between the drug and the reported adverse events. FAERS reports were coded using the MedDRA (Medical Dictionary for Regulatory Activities) terms for DILI²⁷. Although DILI has complex clinical symptoms, there has been documentation for the utilization of the FAERS database to investigate emerging DILI adverse events for newly marketed drugs⁸.

2.1.2. Inclusion/exclusion criteria

Drugs that cause liver injury have been annotated using the United States National Center for Toxicological Research Liver Toxicity Knowledge Base (NCTR-LTKB), which utilizes

hepatotoxic descriptions from the FDA-approved drug labeling regulatory documents as well as evaluating causality evidence in the literature²⁸. This database was created by the FDA to help clinicians, toxicologists, and researchers access information on DILI annotation of various drugs^{28,29}. NCTR-LTKB serves as a centralized source to study the mechanism of DILI and the development or validation of emerging biomarkers and predictive models²⁹. This is the largest publicly available annotated DILI dataset containing three groups based on their potential to cause liver toxicity [Most DILI concern-(192 drugs), Lesser DILI concern-(278 drugs), and No-DILI concern (312 drugs)] with confirmed causal evidence connecting a drug to liver injury²⁸. The FAERS database uses FDA drug labeling information for the classification of drugs according to their potential to cause DILI. This study utilized drugs with “most-DILI concern”, which were defined based on hepatotoxicity resulting in market withdrawal (in US and ex-US), black box warning, or high severity of liver injury noted as part of the warning and precautions label²⁸⁻³⁰. Therefore, both mitochondrial and non-mitochondrial mechanisms of toxicity groups are associated with severe hepatic injury.

The study drugs represented various drug classes such as analgesic, anti-inflammatory, antidepressant, antibiotic, antidiabetic, and antineoplastic agents. Most of these drugs had been withdrawn, have boxed warnings, or have warnings and precautions for liver injury in their prescribing labels. The details of DILI severity categories based on the DILI description are included in the drug labeling: severity level 1; steatosis, level 2; cholestasis and steatohepatitis, level 3; liver aminotransferases increase, level 4; hyperbilirubinemia, level 5; jaundice, level 6; liver necrosis, level 7; acute liver failure, and level 8; hepatotoxicity²⁸. Examples of withdrawn drugs include bromfenac, chlorzoxazone, troglitazone, and trovafloxacin, which have been assigned a severity level of 8, suggesting evidence of fatal hepatotoxicity. Drugs such as bosentan, danazol, ketoconazole, nefazodone, tolcapone, and valproic acid have box warning in their product labeling and have severity categories ranging from 3 to 8.

2.1.3. Classification of drugs as mitochondrial toxicants

Drugs with mitochondrial mechanisms of toxicity were defined by literature evidence of mitochondrial injury mechanisms (yes or no) of *in vitro* (e.g., cellular production of reactive oxygen species *via* oxidative stress, inhibition or uncoupling of respiratory complexes of electron transport chain, induction of mitochondrial membrane permeability transition pore,

inhibition of mitochondrial fatty acid oxidation or mitochondrial DNA damage, etc.)³¹⁻⁴² or *in vivo* mitochondrial toxicity from animal studies (evidence of impairment of oxidative phosphorylation complexes or histopathological alterations of mitochondria in *in vivo* animal models, etc.)^{43,44}. Our classification was based on the parent drug-induced toxicity (direct impact on mitochondria) and not the metabolite. Possible drug effects on mitochondrial biogenesis or respiratory capacity were not considered. Drugs with the non-mitochondrial mechanisms of toxicity were defined by literature evidence of the alternative mechanisms of injury or lack of evidence. It is important to note that 8.2% of drugs had no literature evidence of the type of toxicity mechanism, meaning it could be a mitochondrial or non-mitochondrial mechanism. Furthermore, the non-mitochondrial mechanisms of toxicity drugs are not proven to have non-mitochondrial mechanisms. For these drugs, there is simply no evidence of mitochondrial mechanisms of toxicity information that is publicly available yet.

2.2. Outcome

We determined the number of reports for hepatotoxicity at a group level and an individual drug level using the Reporting Odds Ratio (ROR). As shown in Table 1, we calculated total hepatotoxicity and all other adverse events for both the DILI groups. For ROR calculations, numerators are derived by multiplying the hepatotoxicity reports for mitochondrial mechanisms of toxicity drug group with all other adverse event reports of non-mitochondrial mechanisms of toxicity per drug group. The denominator is calculated by multiplying hepatotoxicity adverse event reports of non-mitochondrial mechanisms of toxicity with all the adverse events reported for mitochondrial mechanisms of toxicity per drug group⁴⁵. Therefore, the ROR for drugs associated with mitochondrial mechanism of toxicity was 1.43 [ROR=(40,343×1,342,486)/(586,989×64,358)=1.43].

Insert Table 1

We also examined the RORs at the individual drug level, as shown in Table 2. A case (hepatotoxicity reports) or non-case (all other adverse event reports) disproportionality approach was utilized by creating a two-by-two contingency table, as demonstrated below using acetaminophen as an example⁴⁵. During this timeframe, a total of 383,540 hepatotoxicity reports and a total of 27,852,908 adverse event counts of any drug type were collected. For ROR

calculations, numerators are derived by multiplying the hepatotoxicity reports for a drug of interest with all other adverse events reports. The denominator is calculated by multiplying hepatotoxicity adverse event reports of all other drugs (excluding acetaminophen) with all the adverse events reported with a drug of interest⁴⁵ (ROR for acetaminophen= $[8,509 \times 27,852,908] / (51,732 \times 383,540) = 11.94$). Within the timeframe, a ROR higher than 1 for a drug indicates a higher proportion of severe liver injury reports for a drug of interest than all the other drug reports in the database. In this case, acetaminophen was associated with proportionally more reports for serious liver adverse events than other drugs in the database.

Insert Table 2

2.3. Association of age, gender, and other factors in two groups of DILI (mitochondrial and non-mitochondrial mechanisms of toxicity)

Mitochondrial function declines with aging¹³; therefore, we evaluated if reports of severe DILI were disproportionately associated with older patient age, indicating potential susceptibility to DILI from mitochondrial mechanisms of toxicity. Furthermore, since gender may play an important role in the sensitivity of DILI, we also evaluated the frequency of reports according to patient gender. The mean and standard deviation (SD) of patient age were calculated and compared between DILI reports caused by drugs associated with mitochondrial mechanisms of toxicity and DILI reports associated with non-mitochondrial mechanisms. Patient age was dichotomized into ≤ 65 years or > 65 years for comparison. Other factors, including drug severity class, patient weight, report type, and label section, were examined in a descriptive analysis.

2.4. Statistical analysis

Descriptive statistics were used to compare the gender and age of reports for the DILI drug groups associated with mitochondrial mechanisms of toxicity and associated with non-mitochondrial mechanisms. The statistical significance of differences in categorical variables such as age group, DILI severity, drug label, and report type between two categories of DILI drugs was examined using the chi-square test. Whereas, differences in continuous variables such as mean patient age between the two categories of hepatotoxic drugs were compared with the two-tailed Student's *t*-test for independent samples. The unadjusted association of age and

gender with mitochondrial mechanisms of toxicity DILI group (against non-mitochondrial mechanisms of toxicity DILI group) was determined using univariate logistic regression analysis. ROR calculations were carried out using a two-by-two contingency table using OpenEpi (version 3.01; Centers for Disease Control and Prevention), which calculates 95% CI and *P*-values *via* Taylor series⁴⁶. Chi-square tests were performed with SAS version 9.4 (SAS Institute, Cary, NC, USA), and two-sided *t*-tests were performed at the 0.05 significance level *via* GraphPad Prism version 8 (La Jolla, CA, USA). All statistical tests were two-sided with a significance level at 0.05.

3. Results

We included 192 drugs classified as having the highest DILI risk (“Most-DILI concern”) *via* the NCTR-LTKB database. Out of these 192 drugs, 134 drugs had searchable FAERS reports, while the remaining 58 drugs were either withdrawn before our study period or were withdrawn from the European market before US approval. Therefore, the final data set contained 134 drugs, which were categorized as 56 drugs causing hepatotoxicity *via* mitochondrial mechanisms, and 78 drugs were classified as causing hepatotoxicity *via* non-mitochondrial mechanisms.

Table 3 indicates the characteristics of the event reports included in the study. A total of 104,701 adverse event reports were extracted from FAERS for the period spanning January 1998 to May 2019. Of these, 40,343 (38.5%) reports of hepatotoxicity were for drugs that were associated with mitochondrial mechanisms of toxicity, whereas 64,358 (61.5%) reports of hepatotoxicity were for drugs associated with non-mitochondrial mechanisms of toxicity. Furthermore, drugs were categorized based on the NCTR-LTKB severity classification. There was a statistically significant difference in DILI severity ($P < 0.0001$) between the two groups of DILI drugs (mitochondrial mechanisms compared to non-mitochondrial mechanisms). There was a 5.5 percentage point difference in reports for more severe DILI (liver failure/hepatotoxicity) for drugs associated with mitochondrial mechanisms of toxicity compared to non-mitochondrial mechanisms (76.3% compared to 70.8%, respectively, $P < 0.0001$). As shown in Table 3, the FAERS reports were classified based on the drug label section for liver injury; there was a statistically significant difference in drug labels ($P < 0.0001$) between the two groups of DILI drugs (mitochondrial mechanisms compared to non-mitochondrial mechanisms). Additionally, 24.6% of mitochondrial mechanisms of toxicity

drugs had a box warning label as compared to 19.8% of non-mitochondrial mechanisms of toxicity drugs, and 64.9% of mitochondrial mechanisms of toxicity drugs had warning and precautions label as compared to 79.1% of non-mitochondrial mechanisms of toxicity drugs. For drugs withdrawn due to hepatotoxicity, there were high numbers of reports ($n = 4227$, 10.5%) for drugs that are associated with mitochondrial toxicity mechanisms, compared to a lower number of reports ($n = 747$, 1.2%) for drugs with non-mitochondrial mechanisms of toxicity ($P < 0.0001$). Over 88% of reports were expedited, while the rest of the reports were either direct or periodic. In summary, there was a statistically significant difference between drug severity classification, label, and liver injury severity according to the drug's ability to cause toxicity through mitochondrial mechanisms.

We also examined patient bodyweight, but 79%–81% of the reports did not have this information documented. Among the 19%–21% of reports where the bodyweight data was present, the average difference between the two groups of DILI drugs (mitochondrial compared to non-mitochondrial mechanisms) was only 1.6 kg (68.6 ± 20.8 compared to 70.2 ± 23.4 ; $P < 0.0001$). In this case, the small P -value may be because the large sample size overpowered the comparison. As large numbers of the reports were missing bodyweight, further analysis was not performed. As shown in Table 3, a higher percentage of males were the subjects of hepatotoxicity reports *via* mitochondrial mechanisms compared to the subjects of reports involving hepatotoxicity *via* the non-mitochondrial mechanisms (49.1% compared to 37.8%, $P < 0.0001$). About 7% to 8.3% of reports were missing information about the patient's gender.

Insert Table 3

Table 3 presents the difference in the mean and distribution of age among the two groups. The patient's age was recorded in more than 71% of the reports from both the groups. As shown in Table 3, there was a statistically significant difference ($P < 0.0001$) between the mean age of patients with hepatotoxicity in drugs that are associated with mitochondrial mechanisms [56.1 ± 18.33 (SD)] compared to non-mitochondrial mechanisms of toxicity [48 ± 19.53 (SD)]. In other words, reports involving drugs associated with mitochondrial mechanisms of hepatic toxicity displayed a higher mean age than reports for drugs associated with non-mitochondrial mechanisms of hepatic toxicity.

Fig. 1 indicates the ROR values of drugs associated with mitochondrial mechanisms of toxicity; benzbromarone, troglitazone, isoniazid, rifampin, and nimesulide had the highest ROR values in the group. Fig. 2 indicates the ROR values of drugs associated with non-mitochondrial mechanisms of toxicity; telithromycin, gemtuzumab ozogamicin, mexiletine, dactomycin, and disulfiram had the highest ROR values in the group. Table 4 indicates the top 20 drugs with the highest ROR values in both groups of hepatotoxicants. The top 20 drugs with the highest ROR values included drugs with either mitochondrial or non-mitochondrial injury mechanisms. The top 4 drugs, which had ROR values higher than 18 (benzbromarone, troglitazone, isoniazid, rifampin), were associated with mitochondrial mechanisms of toxicity. Furthermore, the top two drugs, benzbromarone, and troglitazone were withdrawn from the market.

Insert Table 4

Insert Figs. 1 and 2

Table 5 indicates the RORs between the two groups of drugs that caused liver injury *via* mitochondrial compared to non-mitochondrial mechanisms. Between the two DILI groups, reports for DILI involving mitochondrial mechanisms of toxicity had a 1.43 (95% CI 1.42–1.45; $P < 0.0001$) times higher odds compared to drugs associated with non-mitochondrial mechanisms of toxicity. The univariate logistic regression model was used after dichotomizing age and gender. Table 6 indicates a statistically significant risk association of age or gender with hepatotoxic drugs with mitochondrial toxicity mechanisms. Reports of liver injury were 2.2 (odds ratio: 2.2, 95% CI 2.12–2.26) times more likely to be associated with older patient age, as compared with reports involving patients under 65 years of age. On the other hand, female patients were 37% less likely to be subjects of liver injury reports for drugs associated with mitochondrial mechanisms of toxicity compared to males (Odds Ratio 0.63, 95% CI 0.61–0.64). Supporting Information Tables S1 to S7 contain DILI reports, all adverse event reports, ROR, and 95% Confidence Interval (CI) for all the drugs evaluated in the study.

Insert Tables 5 and 6

Fig. 3 indicates the totality of all ROR scores of DILI drugs with mitochondrial or non-mitochondrial mechanisms of toxicity. Drugs from the antineoplastic, antiviral, analgesic, antibiotic, and antimycobacterial classes were the top 5 drugs classes associated with higher

ROR scores. Drugs from the antidiabetic, antiretroviral, anti-inflammatory, anti-Parkinson, vasoactive, neuroprotective, and antihyperlipidemic drug classes were primarily associated with mitochondrial mechanisms. Alternatively, leukotriene pathway modulators, alcohol antagonists, CNS stimulants, and platelet inhibitor drug classes were the drugs with non-mitochondrial mechanisms having higher RORs. Figs. 4 and 5 categorize these two groups of drugs based on the drug label section and severity class. We did not observe any notable trend between mitochondrial and non-mitochondrial mechanisms and drug label section, or severity class.

Insert Fig. 3–5

4. Discussion

Prediction and characterization for DILI during preclinical drug development and post-approval remains a challenge for the pharmaceutical industry, toxicologists, clinicians, physicians, health authorities, and regulators⁵. Characterizing DILI has been a challenge due to its unpredictability, lack of accurate biomarkers, poorly defined pathogenesis, and its potential to cause fatal liver failure⁵. In the past two decades, drug-induced mitochondrial dysfunction has been established as an important contributing mechanism associated with liver, muscle, heart, kidney, and central nervous system toxicity¹³. Mitochondrial dysfunction is one of the reasons known to cause muscle toxicity by HMG-CoA (3-hydroxy-3-methylglutaryl coenzyme A) reductase inhibitor (cerivastatin), cardiovascular toxicity by anthracyclines (daunorubicin, doxorubicin, idarubicin), and DILI by an antidepressant (nefazodone), antibiotics [isoniazid, ketoconazole(oral)], and anxiolytic (panadiplon) drugs^{19,47-50}.

We calculated the ROR for reports of severe hepatotoxicity adverse events among drugs with the highest risk for DILI, for drugs having mitochondrial or non-mitochondrial mechanisms of toxicity. Brinker et al.⁸ indicated that various measures of disproportionate reporting of adverse events such as Proportional Reporting Ratio, Multi-item Gamma Poisson Shrinker, and the Bayesian Confidence Propagation Neutral Network had been used in analyses of surveillance databases. Each of these methods may have different strengths and limitations and may lead to different sensitivity and specificity for a drug's risk reporting⁸. Various health regulatory authorities use different statistical measures for reporting. For example, the European Medicines Agency uses Proportional Reporting Ratio; FDA and UK's Medicines and Healthcare products

Regulatory Agency uses Multi-item Gamma Poisson Shrinker. Whereas, the World Health Organization has utilized the Bayesian Confidence Propagation Neural Network method for reporting⁸. These reporting measures have been used to generate hypotheses and do not infer adverse event-causal associations. It has been suggested that there is not one single measure of effect that is superior to the others^{8,51,52}. Our study utilized reporting odds ratios to characterize the frequency of liver injury reports as it is a straightforward and frequently used measure for the analysis of FAERS data^{45,53-57}.

The review published by Will et al.¹³ indicated that the most commonly used prescriptions and over the counter medications for geriatric patients had published reports of various toxicities linked to mitochondrial dysfunction. Our study reported that reports for DILI involving mitochondrial mechanisms of toxicity had a 1.43 (95% CI 1.42–1.45; $P < 0.0001$) times higher odds compared to drugs associated with non-mitochondrial mechanisms of toxicity. The ROR scores with the highest risk of liver injury based on mitochondrial or non-mitochondrial mechanisms were highest for drugs from the antineoplastic, antiviral, analgesics, antibiotics, and antimycobacterial classes. This finding agreed with the published literature. Sonawane et al.²⁶ indicated that antineoplastic, analgesics, and antibiotics were among the top 10 drugs that reported severe adverse events in the FAERS database from 2006 to 2014. Additionally, our study observed that over 88% of adverse reports were expedited, while the rest of the reports were either direct or periodic in both drug categories. This observation also agreed with the published literature. Sonawane et al.²⁶ also reported that expedited reports were the most common and over 72% of all serious adverse events with available data on the report type.

Antidiabetic, antiretroviral, anti-inflammatory, anti-Parkinson, vasoactive, neuroprotective, antihyperlipidemic drug classes were primarily associated with mitochondrial toxicity mechanisms. In recent years, an impaired mitochondrial function has been documented as one of the critical factors in inflammation, sarcopenia, metabolic (obesity, type 2 diabetes, non-alcoholic fatty liver disease), and neurodegenerative diseases (Parkinson's, Alzheimer's, Huntington's diseases)^{13,58-60}. Patients with reduced mitochondrial function occurring as a manifestation of their underlying disease state may be more vulnerable to drugs that cause toxicity *via* mitochondrial mechanisms. Alternatively, antiadrenergic, leukotriene pathway

modulators, alcohol antagonist, CNS stimulants, and platelet inhibitors were drug classes that were primarily associated with non-mitochondrial toxicity mechanisms.

We identified statistically significant differences ($P < 0.0001$) in drug severity classification, label section for liver injury, and report type between these two mechanisms for DILI. For drugs withdrawn for liver injury, there were a higher number of hepatotoxicity reports (10.5%) associated with mitochondrial than non-mitochondrial mechanisms (1.2%). Dykens and Will (2007) noted that 38 marketed drugs withdrew from the market between 1994 and 2006. Among these, for cerivastatin, nefazodone, troglitazone, and tolcapone, there was substantial evidence of mitochondrial-induced organ toxicity⁴⁸. Therefore, our observations agreed with reports in the medical literature of drug-induced mitochondrial dysfunction playing an important role in drug withdrawal. Furthermore, Boelsterli and Lim²¹, in 2007, suggested that several drugs, such as amiodarone, dantrolene, diclofenac, isoniazid, lamivudine, leflunomide, mefenamic acid, nimesulide, perhexiline, simvastatin, stavudine, sulindac, tolcapone, troglitazone, trovafloxacin, and valproic acid, are associated with idiosyncratic DILI with a clear link to mitochondrial toxicity. Many of these drugs reported a relatively higher ROR in our study.

Our study reported an older mean patient age [56.1 ± 18.33 (SD)] associated with reports for drugs that cause DILI *via* mitochondrial mechanisms compared to mean age [48 ± 19.53 (SD)] associated with reports for drugs that cause injury *via* non-mitochondrial mechanisms ($P < 0.0001$). This was further substantiated in a univariate logistic regression analysis where reports of liver injury were 2.2 (odds ratio: 2.2, 95% CI 2.12–2.26) times more likely to be associated with older patient age, as compared with reports involving patient ages under 65 years. This finding is consistent with physiological information indicating age as a risk factor for both mitochondrial DNA abnormality and increased oxidative stress-related injury⁵⁹. There is evidence that mitochondrial function declines with age, including the role of mitochondrial DNA mutation, increased production of reactive oxygen species, and the dysfunction in oxidative phosphorylation pathways⁵⁸. The hallmark of mitochondrial aging includes a decreased mitochondrial number, reduced mitochondrial function, and individual electron transport chain activities¹³. Mitochondrial function deteriorates progressively with age.

Therefore, older age populations may be more vulnerable to hepatotoxic drugs associated with mitochondrial mechanisms of toxicity.

Our study indicated that female patients were 37% less likely to report liver injury adverse events for drugs associated with mitochondrial mechanisms of toxicity compared to males. There are conflicting reports associating male gender as a susceptibility factor for DILI, and a clear link for this association is absent in the literature^{21,61}. Several articles allude to the potential involvement of a reactive metabolite, and differences in pharmacokinetics, pharmacodynamics, sex hormones, and immune system response between males and females^{21,22}.

5. Conclusions

Mitochondria play an important part in DILI, including idiosyncratic liver injury. There have been various proposed mechanisms for mitochondrial involvement in DILI⁵⁹. There is a gap in the literature describing the differences in clinical outcomes for patients who experienced DILI from mitochondrial mechanisms of toxicity as compared with non-mitochondrial mechanisms of toxicity drugs. There are limitations in detecting drugs that have mitochondrial liability in the drug development phase of the discovery. For the most part, drug-induced mitochondrial toxicity does not reveal itself in animal models due to the young age, lack of genetic divergence, health status, and lack of concomitant drug exposure¹³. Therefore, drug-induced mitochondrial toxicity is often idiosyncratic, meaning it is not predictable until a large population is exposed^{21,60}. Based on this study, we provide evidence of a higher proportion of reports of severe liver injury adverse events among drugs associated with mitochondrial mechanisms of toxicity as compared with non-mitochondrial mechanisms of toxicity. Furthermore, we found that reports of liver injury were 2.2 (odds ratio: 2.2, 95% CI 2.12–2.26) times more likely to be associated with older patient age, as compared with reports involving patients ages under 65 years. This finding aligns with the theory that age is a susceptibility factor in liver injury *via* the mitochondrial mechanisms of toxicity.

6. Limitation

The FAERS database describes adverse event reports but does not include information about the number of patients treated with a drug. Therefore, incidence rates, prevalence rates, and causal

relationships between drugs and safety adverse events cannot be determined for drugs according to mitochondrial or non-mitochondrial mechanisms of toxicity. For example, the population incidence of DILI may be higher for drugs associated with non-mitochondrial mechanisms than mitochondrial mechanisms of toxicity. Nevertheless, researchers and health authorities have used the FAERS database for adverse event signal identification, developing ideas, and hypothesis generation despite this limitation. The hypothesis and ideas generated using this database could serve as a foundation for more robust study designs, and for *in vitro* or *in vivo* studies investigating the causal relationship of a drug with liver injury. The FAERS database provides a suitable source to evaluate the volume and characteristics of adverse event reports for marketed medications. Furthermore, factors such as age, gender, weight, drug severity class, and label section of FAERS reports can provide valuable insights to health authorities during the post-market surveillance of marketed medications.

The FAERS database is a spontaneous reporting system with limitations when used for drug safety research, including the potential for under or over-reporting events, duplicate reports, influence of media, and uncertainty of reported events^{8,62}. For example, troglitazone received significant media attention due to a class-action lawsuit which called attention to its DILI risk. Therefore, troglitazone may have a higher number of hepatotoxicity reports than some drugs that did not receive media attention. Moreover, the FAERS database could be associated with the “Weber effect”, where adverse event reports are higher in initial marketing stages following a gradual decline⁸.

Mitochondria have a diverse role in the pathophysiology of DILI. In current literature, most of the mitochondrial-induced toxicity is derived from *in vitro* studies. *In vitro* assays using immortalized cell lines or primary human hepatocytes have their limitations as they generally lack competent metabolic function, xenobiotic biotransformation capacity, appropriate drug receptors and transporters, and cellular architecture. Therefore, it is unclear how mitochondrial *in vitro* mechanisms truly translate to liver injury outcomes in humans; there appear to be strong associations as outlined here.

Characterizing DILI drugs based on mitochondrial dysfunction *versus* other mechanisms may have limitations. For acetaminophen, mitochondrial dysfunction plays an essential role in liver injury. This was demonstrated in primary human hepatocytes and preclinical models.

Mechanistically, acetaminophen produces a reactive metabolite leading to disruption of cellular homeostasis. However, acetaminophen as a parent drug does not directly affect the mitochondrial respiratory chain or cause direct toxicity to mitochondria. Therefore, we included acetaminophen in non-mitochondrial DILI drugs while considering that mitochondrial dysfunction plays an important role in acetaminophen-induced liver injury. Moreover, given the limitations of the data source we were unable to discern intrinsic from idiosyncratic DILI.

Patients with an underlying condition such as obesity may be more vulnerable to drugs that cause toxicity *via* mitochondrial mechanisms; thus, we attempted to include patient weight in our study. However, about 79%–81% of the reports missed the bodyweight information; therefore, the effect of patient weight was not examined. The study may also have several unmeasured confounding factors as patient comorbidities, pre-existing liver disease, and concomitant drug use is not captured in FAERS reports. Additionally, the findings regarding age and gender are unadjusted; therefore, it should be used merely for hypothesis generation. Moreover, gender bias may be due to disease demographics. Some of the DILI drugs with mitochondrial toxicity mechanisms are prescribed for diseases with a higher male predisposition. For example, benzbromarone is prescribed for gout, which has six times higher occurrence in males⁶³. Similarly, isoniazid and rifampin are prescribed for the treatment of tuberculosis, which has two times higher occurrence in males⁶⁴.

ROR depends on the reporting rates of liver injury adverse events and all other adverse events reports in compared drug classes. DILI drugs associated with non-mitochondrial mechanisms of toxicity have a significantly higher number of non-hepatic adverse events reports. Therefore, we are not sure if larger ROR values are due to the higher reporting of hepatotoxicity in the drugs with mitochondrial mechanisms of toxicity, or higher reporting of non-hepatic adverse events reported for the drugs with non-mitochondrial mechanisms of toxicity.

For this analysis, we utilized ROR, which is a disproportionality measurement of spontaneous reports and not a method to measure drug-related risks quantitatively. Regulatory actions in response to safety concerns related to age and gender using the FAERS database must be determined *via* individual cases to determine causality. Despite these database limitations, we were able to show that drugs that cause hepatotoxicity *via* mitochondrial mechanisms were

associated with a higher proportion of adverse event reports than drugs having non-mitochondrial mechanisms of toxicity. Additionally, age may play a role in susceptibility to DILI *via* mitochondrial mechanisms of toxicity. Our findings from this study align with mitochondrial mechanisms of toxicity being an important cause of DILI, and this should be further investigated in real-world studies with robust designs.

Acknowledgments

We thank Dr. Michael Leach and Dr. K. Nasir Khan for their support of our research. We also thank Dr. Mark Gosink, Dr. Seda Arat, and Lisa Marroquin for their review and assistance on this article.

Author contributions

Payal Rana was responsible for hypothesis generation, predictive analysis, manuscript outline, and generation of figures & tables. Payal Rana, Dr. Stephen Kogut and Dr. Michael Aleo were accountable for creating the primary manuscript. Dr. Xuerong Wen was responsible for guiding statistical analysis on univariate and multivariable logistic regression models. All authors reviewed, edited, and refined the final manuscript and have given approval to the final version.

Conflicts of interest

The authors declare that there was no conflict of interest.

References

- 1 Sgro C, Clinard F, Ouazir K, Chanay H, Allard C, Guilleminet C, et al. Incidence of drug-induced hepatic injuries: a French population-based study. *Hepatology (Baltimore)* 2002; **36**: 451-5.
- 2 Shapiro MA, Lewis JH. Causality assessment of drug-induced hepatotoxicity: promises and pitfalls. *Clin Liver Dis* 2007; **11**: 477-505, v.
- 3 Friis H, Andreasen PB. Drug-induced hepatic injury: an analysis of 1100 cases reported to the Danish Committee on Adverse Drug Reactions between 1978 and 1987. *J Int Med* 1992; **232**: 133-8.
- 4 Holt M, Ju C. Drug-induced liver injury. *Handbook of experimental pharmacology*. Berlin: Springer; 2010. p.3-27.

- 5 Alempijevic T, Zec S, Milosavljevic T. Drug-induced liver injury: do we know everything?. *World J Hepatol* 2017; **9**: 491-502.
- 6 Raschi E, Poluzzi E, Koci A, Caraceni P, Ponti FD. Assessing liver injury associated with antimycotics: concise literature review and clues from data mining of the FAERS database. *World J Hepatol* 2014; **6**: 601-12.
- 7 Purkins L, Love ER, Eve MD, Wooldridge CL, Cowan C, Smart TS, et al. The influence of diet upon liver function tests and serum lipids in healthy male volunteers resident in a Phase I unit. *Br J Clin Pharmacol* 2004; **57**: 199-208.
- 8 Brinker AD, Lyndly J, Tønning J, Moeny D, Levine JG, Avigan MI. Profiling cumulative proportional reporting ratios of drug-induced liver injury in the FDA Adverse Event Reporting System (FAERS) database. *Drug saf* 2013; **36**: 1169-78.
- 9 EASL Clinical Practice Guidelines: drug-induced liver injury. *J Hepatol* 2019; **70**: 1222-61.
- 10 Hamilton LA, Collins-Yoder A, Collins RE. Drug-induced liver injury. *AACN Adv Crit Care* 2016;**27**:430-440.
- 11 Aleo MD, Luo Y, Swiss R, Bonin PD, Potter DM, Will Y. Human drug-induced liver injury severity is highly associated with dual inhibition of liver mitochondrial function and bile salt export pump. *Hepatology (Baltimore)* 2014; **60**: 1015-22.
- 12 Aleo MD, Shah F, Allen S, Barton HA, Costales C, Lazzaro S, et al. Moving beyond binary predictions of human drug-induced liver injury (DILI) toward contrasting relative risk potential. *Chem Res Toxicol* 2020; **33**: 223-38.
- 13 Will Y, Shields JE, Wallace KB. Drug-induced mitochondrial toxicity in the geriatric population: challenges and future directions. *Biology (Basel)* 2019;**8**:32.
- 14 Vuda M, Kamath A. Drug induced mitochondrial dysfunction: mechanisms and adverse clinical consequences. *Mitochondrion* 2016; **31**: 63-74.
- 15 Haasio K, Nissinen E, Söpanen L, Heinonen EH. Different toxicological profile of two COMT inhibitors *in vivo*: the role of uncoupling effects. *J Neural Transm (Vienna)* 2002;**109**:1391-401.
- 16 Boenzi S, Diodato D. Biomarkers for mitochondrial energy metabolism diseases. *Essays Biochem* 2018; **62**: 443-54.

- 17 Kakisaka K, Yoshida Y, Suzuki Y, Sato T, Kuroda H, Miyasaka A, et al. Serum markers for mitochondrial dysfunction and cell death are possible predictive indicators for drug-induced liver injury by direct acting antivirals. *Hepatol Res* 2018; **48**: 78-86.
- 18 Low Wang CC, Galinkin JL, Hiatt WR. Toxicity of a novel therapeutic agent targeting mitochondrial complex I. *Clin Pharm Ther* 2015; **98**: 551-9.
- 19 Brass EP, Hoppel CL. Mitochondria as targets of drug toxicity: lessons from the R118 phase I experience. *Clin Pharmacol Ther* 2015; **98**: 464-6.
- 20 Stirnimann G, Kessebohm K, Lauterburg B. Liver injury caused by drugs: an update. *Swiss Med Wkly* 2010; **140**: w13080.
- 21 Boelsterli UA, Lim PL. Mitochondrial abnormalities—a link to idiosyncratic drug hepatotoxicity?. *Toxicol Appl Pharmacol* 2007; **220**: 92-107.
- 22 Amacher DE. Female gender as a susceptibility factor for drug-induced liver injury. *Hum Exp Toxicol* 2014; **33**: 928-39.
- 23 Shi Q, Yang X, Mattes WB, Mendrick DL, Harrill AH, Beger RD. Circulating mitochondrial biomarkers for drug-induced liver injury. *Biomark Med* 2015; **9**: 1215-23.
- 24 Ramachandran A, Duan L, Akakpo JY, Jaeschke H. Mitochondrial dysfunction as a mechanism of drug-induced hepatotoxicity: current understanding and future perspectives. *J Clin Transl Res* 2018; **4**:75-100.
- 25 Hunt CM. Mitochondrial and immunoallergic injury increase risk of positive drug rechallenge after drug-induced liver injury: a systematic review. *Hepatology (Baltimore)* 2010; **52**: 2216-22.
- 26 Sonawane KB, Cheng N, Hansen RA. Serious adverse drug events reported to the FDA: analysis of the FDA Adverse Event Reporting System 2006–2014 Database. *J Manag Care Spec Pharm* 2018; **24**: 682-90.
- 27 Sarntivijai S, Zhang S, Jagannathan DG, Zaman S, Burkhart KK, Omenn GS, et al. Linking MedDRA[®]-coded clinical phenotypes to biological mechanisms by the ontology of adverse events: a pilot study on tyrosine kinase inhibitors. *Drug saf* 2016; **39**: 697-707.
- 28 Chen M, Vijay V, Shi Q, Liu Z, Fang H, Tong W. FDA-approved drug labeling for the study of drug-induced liver injury. *Drug discov Today* 2011; **16**: 697-703.

- 29 Thakkar S, Chen M, Fang H, Liu Z, Roberts R, Tong W. The Liver Toxicity Knowledge Base (LKTb) and drug-induced liver injury (DILI) classification for assessment of human liver injury. *Expert Rev Gastroenterol Hepatol* 2018; **12**: 31-8.
- 30 Hong H, Thakkar S, Chen M, Tong W. Development of decision forest models for prediction of drug-induced liver injury in humans using a large set of FDA-approved drugs. *Sci Rep* 2017; **7**: 17311.
- 31 Hynes J, Swiss RL, Will Y. High-throughput analysis of mitochondrial oxygen consumption. *Methods Mol Biol* 2012; **810**: 59-72.
- 32 Lin Z, Will Y. Evaluation of drugs with specific organ toxicities in organ-specific cell lines. *Toxicol Sci* 2012; **126**: 114-27.
- 33 Luo Y, Rana P, Will Y. Palmitate increases the susceptibility of cells to drug-induced toxicity: an *in vitro* method to identify drugs with potential contraindications in patients with metabolic disease. *Toxicol Sci* 2012; **129**: 346-62.
- 34 Marroquin LD, Hynes J, Dykens JA, Jamieson JD, Will Y. Circumventing the Crabtree effect: replacing media glucose with galactose increases susceptibility of HepG2 cells to mitochondrial toxicants. *Toxicol Sci* 2007; **97**: 539-47.
- 35 Nadanaciva S, Dykens JA, Bernal A, Capaldi RA, Will Y. Mitochondrial impairment by PPAR agonists and statins identified *via* immunocaptured OXPHOS complex activities and respiration. *Toxicol Appl Pharmacol* 2007; **223**: 277-87.
- 36 Nadanaciva S, Rana P, Beeson GC, Chen D, Ferrick DA, Beeson CC, et al. Assessment of drug-induced mitochondrial dysfunction *via* altered cellular respiration and acidification measured in a 96-well platform. *J Bioenerg Biomembr* 2012; **44**: 421-37.
- 37 Nadanaciva S, Will Y. New insights in drug-induced mitochondrial toxicity. *Curr Pharm Des* 2011; **17**: 2100-12.
- 38 Nadanaciva S, Will Y. Investigating mitochondrial dysfunction to increase drug safety in the pharmaceutical industry. *Curr Drug Targets* 2011; **12**: 774-82.
- 39 Nadanaciva S, Will Y. The role of mitochondrial dysfunction and drug safety. *IDrugs* 2009; **12**: 706-10.
- 40 Rana P, Aleo MD, Gosink M, Will Y. Evaluation of *in vitro* mitochondrial toxicity assays and physicochemical properties for prediction of organ toxicity using 228 pharmaceutical drugs. *Chem Res Toxicol* 2019; **32**: 156-67.

- 41 Shah F, Leung L, Barton HA, Will Y, Rodrigues AD, Greene N, et al. Setting clinical exposure levels of concern for drug-induced liver injury (DILI) using mechanistic *in vitro* assays. *Toxicol Sci* 2015; **147**: 500-14.
- 42 Shah F, Louise-May S, Greene N. Chemotypes sensitivity and predictivity of *in vivo* outcomes for cytotoxic assays in THLE and HepG2 cell lines. *Bioorg Med Chem Lett* 2014; **24**: 2753-7.
- 43 Porceddu M, Buron N, Roussel C, Labbe G, Fromenty B, Borgne-Sanchez A. Prediction of liver injury induced by chemicals in human with a multiparametric assay on isolated mouse liver mitochondria. *Toxicol Sci* 2012; **129**: 332-45.
- 44 Begriche K, Massart J, Robin MA, Borgne-Sanchez A, Fromenty B. Drug-induced toxicity on mitochondria and lipid metabolism: mechanistic diversity and deleterious consequences for the liver. *J Hepatol* 2011; **54**: 773-94.
- 45 Sakaeda T, Tamon A, Kadoyama K, Okuno Y. Data mining of the public version of the FDA Adverse Event Reporting System. *Int J Med Sci* 2013; **10**: 796-803.
- 46 Sullivan KM, Dean A, Soe MM. OpenEpi: a web-based epidemiologic and statistical calculator for public health. *Public Health Rep* 2009; **124**: 471-4.
- 47 Dykens JA, Jamieson JD, Marroquin LD, Nadanaciva S, Xu JJ, Dunn MC, et al. *In vitro* assessment of mitochondrial dysfunction and cytotoxicity of nefazodone, trazodone, and buspirone. *Toxicol Sci* 2008; **103**: 335-45.
- 48 Dykens JA, Will Y. The significance of mitochondrial toxicity testing in drug development. *Drug Discov Today* 2007; **12**: 777-85.
- 49 Ulrich RG, Bacon JA, Brass EP, Cramer CT, Petrella DK, Sun EL. Metabolic, idiosyncratic toxicity of drugs: overview of the hepatic toxicity induced by the anxiolytic, panadipilon. *Chem Biol Interact* 2001; **134**: 251-70.
- 50 Dykens JA, Jamieson J, Marroquin L, Nadanaciva S, Billis PA, Will Y. Biguanide-induced mitochondrial dysfunction yields increased lactate production and cytotoxicity of aerobically-poised HepG2 cells and human hepatocytes *in vitro*. *Toxicol Appl Pharmacol* 2008; **233**: 203-10.
- 51 Deshpande G, Gogolak V, Smith SW. Data mining in drug safety. *Pharm Med* 2010; **24**: 37-43.

- 52 Hauben M, Madigan D, Gerrits CM, Walsh L, Van Puijenbroek EP. The role of data mining in pharmacovigilance. *Expert Opin Drug Saf* 2005; **4**: 929-48.
- 53 Shimada K, Hasegawa S, Nakao S, Mukai R, Sasaoka S, Ueda N, et al. Adverse reaction profiles of hemorrhagic adverse reactions caused by direct oral anticoagulants analyzed using the Food and Drug Administration Adverse Event Reporting System (FAERS) database and the Japanese Adverse Drug Event Report (JADER) database. *Int J Med Sci* 2019; **16**: 1295-303.
- 54 Fujimoto M, Hosomi K, Takada M. Statin-associated lower urinary tract symptoms: data mining of the public version of the FDA adverse event reporting system, FAERS. *Int J Clin Pharmacol Ther* 2014; **52**: 259-66.
- 55 Kaur K, Fayad R, Saxena A, Frizzell N, Chanda A, Das S, et al. Fluoroquinolone-related neuropsychiatric and mitochondrial toxicity: a collaborative investigation by scientists and members of a social network. *J Community Support Oncol* 2016; **14**: 54-65.
- 56 Ohyama K, Hori Y, Sugiura M. Evaluation of syncope association with α_1 -adrenoceptor blockers in males using the FAERS database: impact of concomitant hypertension. *Pharmazie* 2019; **74**: 755-9.
- 57 Zhai Y, Ye X, Hu F, Xu J, Guo X, Zhuang Y, et al. Endocrine toxicity of immune checkpoint inhibitors: a real-world study leveraging US Food and Drug Administration adverse events reporting system. *J Immunother Cancer* 2019; **7**: 286.
- 58 Srivastava S. The mitochondrial basis of aging and age-related disorders. *Genes (Basel)* 2017;**8**:398.
- 59 Boelsterli UA, Ho HK, Zhou S, Leow KY. Bioactivation and hepatotoxicity of nitroaromatic drugs. *Curr Drug Metab* 2006; **7**: 715-27.
- 60 Boelsterli UA. Disease-related determinants of susceptibility to drug-induced idiosyncratic hepatotoxicity. *Curr Opin Drug Discov Dev* 2003; **6**: 81-91.
- 61 Worland T, Chin KL, Rodrigues B, Nicoll A. A retrospective case-controlled cohort study of inpatient drug induced liver injury: the RIDDLE study. *Transl Gastroenterol Hepatol* 2020; **5**: 33.
- 62 Johnson K, Guo C, Gosink M, Wang V, Hauben M. Multinomial modeling and an evaluation of common data-mining algorithms for identifying signals of disproportionate reporting in pharmacovigilance databases. *Bioinformatics (Oxford)* 2012; **28**: 3123-30.

- 63 Singh JA. Racial and gender disparities among patients with gout. *Curr Rheumatol Rep* 2013; **15**: 307.
- 64 Horton KC, MacPherson P, Houben RM, White RG, Corbett EL. Sex differences in tuberculosis burden and notifications in low- and middle-income countries: a systematic review and meta-analysis. *PLoS Med* 2016; **13**: e1002119.
- 65 Xu JJ, Henstock PV, Dunn MC, Smith AR, Chabot JR, de Graaf D. Cellular imaging predictions of clinical drug-induced liver injury. *Toxicol Sci* 2008; **105**: 97-105.
- 66 Ghosh R, Goswami SK, Feitoza LF, Hammock B, Gomes AV. Diclofenac induces proteasome and mitochondrial dysfunction in murine cardiomyocytes and hearts. *Int J Cardiol* 2016; **223**: 923-35.
- 67 Nadanaciva S, Aleo MD, Strock CJ, Stedman DB, Wang H, Will Y. Toxicity assessments of nonsteroidal anti-inflammatory drugs in isolated mitochondria, rat hepatocytes, and zebrafish show good concordance across chemical classes. *Toxicol Appl Pharmacol* 2013; **272**: 272-80.
- 68 Mahmud T, Rafi SS, Scott DL, Wrigglesworth JM, Bjarnason I. Nonsteroidal antiinflammatory drugs and uncoupling of mitochondrial oxidative phosphorylation. *Arthritis Rheum* 1996; **39**: 1998-2003.
- 69 Seo SK, Lee HC, Woo SH, Jin HO, Yoo DH, Lee SJ, et al. Sulindac-derived reactive oxygen species induce apoptosis of human multiple myeloma cells *via* p38 mitogen activated protein kinase-induced mitochondrial dysfunction. *Apoptosis* 2007; **12**: 195-209.
- 70 Sorensen EM, Acosta D. Relative toxicities of several nonsteroidal antiinflammatory compounds in primary cultures of rat hepatocytes. *J Toxicol Environ Health* 1985; **16**: 425-40.
- 71 Hynes J, Nadanaciva S, Swiss R, Carey C, Kirwan S, Will Y. A high-throughput dual parameter assay for assessing drug-induced mitochondrial dysfunction provides additional predictivity over two established mitochondrial toxicity assays. *Toxicol In Vitro* 2013; **27**: 560-9.
- 72 Murray MD, Brater DC. Renal toxicity of the nonsteroidal anti-inflammatory drugs. *Annu Revi Pharmacol Toxicol* 1993; **33**: 435-65.

- 73 Dybing E, Nelson SD, Mitchell JR, Sasame HA, Gillette JR. Oxidation of alpha-methyldopa and other catechols by cytochrome P-450-generated superoxide anion: possible mechanism of methyldopa hepatitis. *Mol Pharmacol* 1976; **12**: 911-20.
- 74 Jatoe SD, Lauriault V, McGirr LG, O'Brien PJ. The toxicity of disulphides to isolated hepatocytes and mitochondria. *Drug Metabol Drug Interact* 1988; **6**: 395-412.
- 75 Felser A, Stoller A, Morand R, Schnell D, Donzelli M, Terracciano L, et al. Hepatic toxicity of dronedarone in mice: role of mitochondrial beta-oxidation. *Toxicology* 2014; **323**: 1-9.
- 76 Gomes MP, Goncalves CA, de Brito JCM, Souza AM, da Silva Cruz FV, Bicalho EM, et al. Ciprofloxacin induces oxidative stress in duckweed (*Lemna minor* L.): implications for energy metabolism and antibiotic-uptake ability. *J Hazard Mater* 2017; **328**: 140-9.
- 77 Salimi A, Eybagi S, Seydi E, Naserzadeh P, Kazerouni NP, Pourahmad J. Toxicity of macrolide antibiotics on isolated heart mitochondria: a justification for their cardiotoxic adverse effect. *Xenobiotica* 2016; **46**: 82-93.
- 78 Liguori MJ, Anderson MG, Bukofzer S, McKim J, Pregonzer JF, Retief J, et al. Microarray analysis in human hepatocytes suggests a mechanism for hepatotoxicity induced by trovafloxacin. *Hepatology (Baltimore)* 2005; **41**: 177-86.
- 79 Leiva M, Ruiz-Bravo A, Moreno E, Jimenez-Valera M. Telithromycin inhibits the production of proinflammatory mediators and the activation of NF-kappaB in *in vitro*-stimulated murine cells. *FEMS Immunol Med Microbiol* 2008; **53**: 343-50.
- 80 Santos NA, Medina WS, Martins NM, Mingatto FE, Curti C, Santos AC. Aromatic antiepileptic drugs and mitochondrial toxicity: effects on mitochondria isolated from rat liver. *Toxicol In Vitro* 2008; **22**: 1143-52.
- 81 Finsterer J. Toxicity of antiepileptic drugs to mitochondria. *Handb Exp Pharmacol* 2017; **240**: 473-88.
- 82 Finsterer J, Scorza FA. Effects of antiepileptic drugs on mitochondrial functions, morphology, kinetics, biogenesis, and survival. *Epilepsy Res* 2017; **136**: 5-11.
- 83 Dieckhaus CM, Thompson CD, Roller SG, Macdonald TL. Mechanisms of idiosyncratic drug reactions: the case of felbamate. *Chem Biol Interact* 2002; **142**: 99-117.
- 84 Poplawska M, Borowicz KK, Czuczwar SJ. The safety and efficacy of fosphenytoin for the treatment of status epilepticus. *Expert Rev Neurother* 2015; **15**: 983-92.

- 85 Higgins SC, Pilkington GJ. The *in vitro* effects of tricyclic drugs and dexamethasone on cellular respiration of malignant glioma. *Anticancer Res* 2010; **30**: 391-7.
- 86 Choi HS, Park JH, Ahn JH, Hong S, Cho JH, Won MH, et al. The anti-inflammatory activity of duloxetine, a serotonin/norepinephrine reuptake inhibitor, prevents kainic acid-induced hippocampal neuronal death in mice. *J Neurol Sci* 2015; **358**: 390-7.
- 87 Yuan SY, Cheng CL, Ho HC, Wang SS, Chiu KY, Su CK, et al. Nortriptyline induces mitochondria and death receptor-mediated apoptosis in bladder cancer cells and inhibits bladder tumor growth *in vivo*. *Eur J Pharmacol* 2015; **761**: 309-20.
- 88 Sahdeo S, Tomilov A, Komachi K, Iwahashi C, Datta S, Hughes O, et al. High-throughput screening of FDA-approved drugs using oxygen biosensor plates reveals secondary mitofunctional effects. *Mitochondrion* 2014; **17**: 116-25.
- 89 Somchit N, Hassim SM, Samsudin SH. Itraconazole- and fluconazole-induced toxicity in rat hepatocytes: a comparative *in vitro* study. *Hum Exp Toxicol* 2002; **21**: 43-8.
- 90 Rodriguez RJ, Acosta D, Jr. Inhibition of mitochondrial function in isolated rat liver mitochondria by azole antifungals. *J Biochem Toxicol* 1996; **11**: 127-31.
- 91 Knasmuller S, Parzefall W, Helma C, Kassie F, Ecker S, Schulte-Hermann R. Toxic effects of griseofulvin: disease models, mechanisms, and risk assessment. *Crit Rev Toxicol* 1997; **27**: 495-537.
- 92 Yanni SB, Augustijns PF, Benjamin DK, Jr., Brouwer KL, Thakker DR, Annaert PP. In vitro investigation of the hepatobiliary disposition mechanisms of the antifungal agent micafungin in humans and rats. *Drug Metab Dispos* 2010; **38**: 1848-56.
- 93 Iverson SL, Utrecht JP. Identification of a reactive metabolite of terbinafine: insights into terbinafine-induced hepatotoxicity. *Chem Res Toxicol* 2001; **14**: 175-81.
- 94 Haegler P, Joerin L, Krahenbuhl S, Bouitbir J. Hepatocellular toxicity of imidazole and triazole antimycotic agents. *Toxicol Sci* 2017; **157**: 183-95.
- 95 Williams CD, McGill MR, Lebofsky M, Bajt ML, Jaeschke H. Protection against acetaminophen-induced liver injury by allopurinol is dependent on aldehyde oxidase-mediated liver preconditioning. *Toxicol Appl Pharmacol* 2014; **274**: 417-24.
- 96 Kim HS, Lim SW, Jin L, Jin J, Chung BH, Yang CW. The protective effect of febuxostat on chronic tacrolimus-induced nephrotoxicity in rats. *Nephron* 2017; **135**: 61-71.

- 97 Yoshikawa Y, Hosomi H, Fukami T, Nakajima M, Yokoi T. Establishment of knockdown of superoxide dismutase 2 and expression of CYP3A4 cell system to evaluate drug-induced cytotoxicity. *Toxicology In Vitro* 2009; **23**: 1179-87.
- 98 Jamieson JD, Smith EB, Dalvie DK, Stevens GJ, Yanochko GM. Myeloperoxidase-mediated bioactivation of 5-hydroxythiabendazole: a possible mechanism of thiabendazole toxicity. *Toxicol In Vitro* 2011; **25**: 1061-6.
- 99 Leung K, Quezada M, Chen Z, Kanel G, Kaplowitz N. Niacin-induced anicteric microvesicular steatotic acute liver failure. *Hepatol Commun* 2018; **2**: 1293-8.
- 100 Orlando R, Piccoli P, De Martin S, Padrini R, Palatini P. Effect of the CYP3A4 inhibitor erythromycin on the pharmacokinetics of lignocaine and its pharmacologically active metabolites in subjects with normal and impaired liver function. *Br J Clin Pharmacol* 2003; **55**: 86-93.
- 101 Yang SD, Bai ZL, Zhang F, Ma L, Yang DL, Ding WY. Levofloxacin increases the effect of serum deprivation on anoikis of rat nucleus pulposus cells *via* Bax/Bcl-2/caspase-3 pathway. *Toxicol Mech Methods* 2014; **24**: 688-96.
- 102 Huang SP, Chien JY, Tsai RK. Ethambutol induces impaired autophagic flux and apoptosis in the rat retina. *Dis Model Mech* 2015; **8**: 977-87.
- 103 Alachkar H, Fulton N, Sanford B, Malnassy G, Mutonga M, Larson RA, et al. Expression and polymorphism (rs4880) of mitochondrial superoxide dismutase (SOD2) and asparaginase induced hepatotoxicity in adult patients with acute lymphoblastic leukemia. *Pharmacogenomics J* 2017; **17**: 274-9.
- 104 Waseem M, Parvez S, Tabassum H. Mitochondria as the target for the modulatory effect of curcumin in oxaliplatin-induced toxicity in isolated rat liver mitochondria. *Arch Med Res* 2017; **48**: 55-63.
- 105 Zhang J, Salminen A, Yang X, Luo Y, Wu Q, White M, et al. Effects of 31 FDA approved small-molecule kinase inhibitors on isolated rat liver mitochondria. *Arch Toxicol* 2017; **91**: 2921-38.
- 106 Rana P, Nadanaciva S, Will Y. Mitochondrial membrane potential measurement of H9c2 cells grown in high-glucose and galactose-containing media does not provide additional predictivity towards mitochondrial assessment. *Toxicol In Vitro* 2011; **25**: 580-7.

- 107 Huuskonen MT, Loppi S, Dhungana H, Keksa-Goldsteine V, Lemarchant S, Korhonen P, et al. Bexarotene targets autophagy and is protective against thromboembolic stroke in aged mice with tauopathy. *Sci Rep* 2016; **6**: 33176.
- 108 Kim SY, Song X, Zhang L, Bartlett DL, Lee YJ. Role of Bcl-xL/Beclin-1 in interplay between apoptosis and autophagy in oxaliplatin and bortezomib-induced cell death. *Biochem Pharmacol* 2014; **88**: 178-88.
- 109 Megias-Vericat JE, Montesinos P, Herrero MJ, Moscardo F, Boso V, Martinez-Cuadron D, et al. Impact of novel polymorphisms related to cytotoxicity of cytarabine in the induction treatment of acute myeloid leukemia. *Pharmacogenet Genomics* 2017; **27**: 270-4.
- 110 Zhu P, Zhu JJ. Tumor treating fields: a novel and effective therapy for glioblastoma: mechanism, efficacy, safety and future perspectives. *Chin Clin Oncol* 2017; **6**: 41.
- 111 Arndt C, Hawkins D, Anderson JR, Breitfeld P, Womer R, Meyer W. Age is a risk factor for chemotherapy-induced hepatopathy with vincristine, dactinomycin, and cyclophosphamide. *J Clin Oncol* 2004; **22**: 1894-901.
- 112 Gutierrez-Escobar AJ, Mendez-Callejas G. Interactome analysis of microtubule-targeting agents reveals cytotoxicity bases in normal cells. *Genomics Proteomics Bioinformatics* 2017; **15**: 352-60.
- 113 Nuvoli B, Camera E, Mastrofrancesco A, Briganti S, Galati R. Modulation of reactive oxygen species via ERK and STAT3 dependent signalling are involved in the response of mesothelioma cells to exemestane. *Free Radic Biol Med* 2018; **115**: 266-77.
- 114 Wang DD, Liu Y, Li N, Zhang Y, Jin Q, Hao DC, et al. Induction of CYP1A1 increases gefitinib-induced oxidative stress and apoptosis in A549 cells. *Toxicol In Vitro* 2017; **44**: 36-43.
- 115 van Hell AJ, Haimovitz-Friedman A, Fuks Z, Tap WD, Kolesnick R. Gemcitabine kills proliferating endothelial cells exclusively via acid sphingomyelinase activation. *Cell Signal* 2017; **34**: 86-91.
- 116 Yamauchi T, Matsuda Y, Tasaki T, Negoro E, Ikegaya S, Takagi K, et al. Induction of DNA strand breaks is critical to predict the cytotoxicity of gemtuzumab ozogamicin against leukemic cells. *Cancer Sci* 2012; **103**: 1722-9.

- 117 Kirkwood JM, Bender C, Agarwala S, Tarhini A, Shipe-Spotloe J, Smelko B, et al. Mechanisms and management of toxicities associated with high-dose interferon alfa-2b therapy. *J Clin Oncol* 2002; **20**: 3703-18.
- 118 Doherty KR, Wappel RL, Talbert DR, Trusk PB, Moran DM, Kramer JW, et al. Multi-parameter *in vitro* toxicity testing of crizotinib, sunitinib, erlotinib, and nilotinib in human cardiomyocytes. *Toxicol Appl Pharmacol* 2013; **272**: 245-55.
- 119 Gustafsson F, Foster AJ, Sarda S, Bridgland-Taylor MH, Kenna JG. A correlation between the *in vitro* drug toxicity of drugs to cell lines that express human P450s and their propensity to cause liver injury in humans. *Toxicol Sci* 2014; **137**: 189-211.
- 120 Paemane A, Sornjai W, Kittisenachai S, Sirinonthanawech N, Roytrakul S, Wongtrakul J, et al. Nevirapine induced mitochondrial dysfunction in HepG2 cells. *Sci Rep* 2017; **7**: 9194.
- 121 Viengchareun S, Caron M, Auclair M, Kim MJ, Frachon P, Capeau J, et al. Mitochondrial toxicity of indinavir, stavudine and zidovudine involves multiple cellular targets in white and brown adipocytes. *Antivir Ther* 2007; **12**: 919-29.
- 122 Heidari R, Babaei H, Eghbal M. Mechanisms of methimazole cytotoxicity in isolated rat hepatocytes. *Drug Chem Toxicol* 2013; **36**: 403-11.
- 123 Banerjee A, Linscheer WG, Chiji H, Murthy UK, Cho C, Nandi J, et al. Induction of an ATPase inhibitor protein by propylthiouracil and protection against paracetamol (acetaminophen) hepatotoxicity in the rat. *Br J Pharmacol* 1998; **124**: 1041-7.
- 124 Kohler JJ, Hosseini SH, Green E, Fields E, Abuin A, Ludaway T, et al. Absence of mitochondrial toxicity in hearts of transgenic mice treated with abacavir. *Cardiovasc Toxicol* 2010; **10**: 146-51.
- 125 Blas-Garcia A, Marti-Rodrigo A, Victor VM, Polo M, Alegre F, Funes HA, et al. The purine analogues abacavir and didanosine increase acetaminophen-induced hepatotoxicity by enhancing mitochondrial dysfunction. *J Antimicrob Chemother* 2016; **71**: 916-26.
- 126 Purnell PR, Fox HS. Efavirenz induces neuronal autophagy and mitochondrial alterations. *J Pharmacol Exp Ther* 2014; **351**: 250-8.

- 127 Dhillon R, Rossi S, Herrine SK. Pegylated interferon 2a and 2b in combination with ribavirin for the treatment of chronic hepatitis C in HIV infected patients. *Ther Clin Risk Manag* 2008; **4**: 789-96.
- 128 Blas-Garcia A, Polo M, Alegre F, Funes HA, Martinez E, Apostolova N, et al. Lack of mitochondrial toxicity of darunavir, raltegravir and rilpivirine in neurons and hepatocytes: a comparison with efavirenz. *J Antimicrob Chemother* 2014; **69**: 2995-3000.
- 129 Robertson K, Liner J, Meeker RB. Antiretroviral neurotoxicity. *J Neurovirol* 2012; **18**: 388-99.
- 130 Latronico T, Pati I, Ciavarella R, Fasano A, Mengoni F, Lichtner M, et al. *In vitro* effect of antiretroviral drugs on cultured primary astrocytes: analysis of neurotoxicity and matrix metalloproteinase inhibition. *J Neurochem* 2018; **144**: 271-84.
- 131 Nunez M, Soriano V. Hepatotoxicity of antiretrovirals: incidence, mechanisms and management. *Drug Saf* 2005; **28**: 53-66.
- 132 Isidori A, Borin L, Elli E, Latagliata R, Martino B, Palumbo G, et al. Iron toxicity—its effect on the bone marrow. *Blood Rev* 2018; **32**: 473-9.
- 133 Lee AU, Farrell GC. Mechanism of azathioprine-induced injury to hepatocytes: roles of glutathione depletion and mitochondrial injury. *J Hepatol* 2001; **35**: 756-64.
- 134 Xuan J, Ren Z, Qing T, Couch L, Shi L, Tolleson WH, et al. Mitochondrial dysfunction induced by leflunomide and its active metabolite. *Toxicology* 2018; **396-397**: 33-45.
- 135 Widemann BC, Adamson PC. Understanding and managing methotrexate nephrotoxicity. *Oncologist* 2006; **11**: 694-703.
- 136 Linares V, Alonso V, Domingo JL. Oxidative stress as a mechanism underlying sulfasalazine-induced toxicity. *Exp Opin Drug Saf* 2011; **10**: 253-63.
- 137 Wu Y, Beland FA, Chen S, Liu F, Guo L, Fang JL. Mechanisms of tolvaptan-induced toxicity in HepG2 cells. *Biochem Pharmacol* 2015; **95**: 324-36.
- 138 Chakrabarti M, Ghosh I, Jana A, Ghosh M, Mukherjee A. Genotoxicity of antiobesity drug orlistat and effect of caffeine intervention: an *in vitro* study. *Drug And Chem Toxicol* 2017; **40**: 339-43.
- 139 Kalota A, Selak MA, Garcia-Cid LA, Carroll M. Eltrombopag modulates reactive oxygen species and decreases acute myeloid leukemia cell survival. *PLoS One* 2015; **10**: e0126691.

- 140 Neri M, Bello S, Bonsignore A, Cantatore S, Riezzo I, Turillazzi E, et al. Anabolic androgenic steroids abuse and liver toxicity. *Mini Rev Med Chem* 2011; **11**: 430-7.
- 141 Pozzi R, Fernandes KR, de Moura CF, Ferrari RA, Fernandes KP, Renno AC, et al. Nandrolone decanoate induces genetic damage in multiple organs of rats. *Arch Environ Contam Toxicol* 2013; **64**: 514-8.
- 142 Welder AA, Robertson JW, Fugate RD, Melchert RB. Anabolic-androgenic steroid-induced toxicity in primary neonatal rat myocardial cell cultures. *Toxicol Appl Pharmacol* 1995; **133**: 328-42.
- 143 Mu X, LeBlanc GA. Developmental toxicity of testosterone in the crustacean *Daphnia magna* involves anti-ecdysteroidal activity. *Gen Comp Endocrinol* 2002; **129**: 127-33.
- 144 Graham RM. Cyclosporine: mechanisms of action and toxicity. *Cleve Clin J Med* 1994; **61**: 308-13.
- 145 Joshi EM, Heasley BH, Macdonald TL. 2-ABT-S-oxide detoxification by glutathione S-transferases A1-1, M1-1 and P1-1: implications for toxicity associated with zileuton. *Xenobiotica* 2009; **39**: 197-204.
- 146 Durham JA, Gandolfi AJ, Bentley JB. Hepatotoxicological evaluation of dantrolene sodium. *Drug Chem Toxicol* 1984; **7**: 23-40.
- 147 Moreno-Ortega AJ, Al-Achbili LM, Alonso E, de Los Rios C, Garcia AG, Ruiz-Nuno A, et al. Neuroprotective effect of the novel compound ITH33/IQM9.21 against oxidative stress and Na⁺ and Ca²⁺ overload in motor neuron-like NSC-34 cells. *Neurotox Res* 2016; **30**: 380-91.
- 148 Lee JS, Kim YH, Park YM. The toxicity of nonsteroidal anti-inflammatory eye drops against human corneal epithelial cells *in vitro*. *J Korean Med Sci* 2015; **30**: 1856-64.
- 149 Shimizu S, Atsumi R, Nakazawa T, Izumi T, Sudo K, Okazaki O, et al. Ticlopidine-induced hepatotoxicity in a GSH-depleted rat model. *Arch Toxicol* 2011; **85**: 347-53.
- 150 Shevell M, Schreiber R. Pemoline-associated hepatic failure: a critical analysis of the literature. *Pediatr Neurol* 1997; **16**: 14-6.
- 151 Huang H, Li LJ, Zhang HB, Wei AY. Papaverine selectively inhibits human prostate cancer cell (PC-3) growth by inducing mitochondrial mediated apoptosis, cell cycle arrest and downregulation of NF-kappaB/PI3K/Akt signalling pathway. *J BUON* 2017; **22**: 112-8.

Tables**Table 1** Reporting odds ratio estimates for DILI drug groups (FAERS reports).

DILI groups	Hepatotoxicity	All other adverse events	Total
Drugs associated with mitochondrial mechanisms of toxicity	40,343	586,989	627,332
Drugs associated with non-mitochondrial mechanisms of toxicity	64,358	1,342,486	1,406,844
Total	104,701	1,929,475	2,034,176

Table 2 Example reporting odds ratio estimate for an individual drug: acetaminophen (FAERS Reports).

Drug	Hepatotoxicity	All other adverse events	Total
Acetaminophen	8,509	51,732	60,241
All other drugs of any type	383,540	27,852,908	28,236,448
Total	392,049	27,904,640	28,296,689

Table 3 Characteristics of patients and hepatotoxic drugs associated with mitochondrial mechanisms of toxicity compared to non-mitochondrial mechanisms.

Characteristics	Hepatotoxicity <i>via</i> mitochondrial mechanism (56 drugs)	Hepatotoxicity <i>via</i> non-mitochondrial mechanism (78 drugs)
FAERS report counts (<i>n</i>)	40,343 (38.5%)	64,358 (61.5%)
Reports based on NKTR drug severity classification		
3 - Liver aminotransferases increase	0 (0%)	3,048 (4.7%)
4 - Hyperbilirubinemia	1,958 (4.9%)	2,292 (3.6%)
5 - Jaundice	7,526 (18.7%)	13,392 (20.8)
6 - Liver necrosis	0 (0%)	35 (0.05%)
7 - Acute liver failure	4,581 (11.3%)	17,207 (26.7%)
8 - Fatal hepatotoxicity	26,278 (65%)	28,384 (44.1)
Reports combined based on less and severe DILI		
Less severe injury	9484 (23.5%)	18,767 (29.2%)
Liver failure/hepatotoxicity	30,859 (76.5%)	45,591 (70.8%)
Reports based on drug label section		
Warning & precautions	26,177 (64.9%)	50,898 (79.1%)
Box warning	9,939 (24.6%)	12,713 (19.8%)
Withdrawn	4227 (10.5%)	747 (1.2%)
Report type		
Direct	1,992 (4.9%)	2,393 (3.7%)
Expedited	35,569 (88.2%)	57,119 (88.8%)
Periodic	2,782 (6.9%)	4,846 (7.5%)
Patient characteristics		
Weight (kg)		
FAERS report counts (<i>n</i>)	7666 (19%)	13532 (21%)
Weight missing	32677 (81%)	50826 (79%)
Weight Mean \pm SD	68.6 \pm 20.8	70.2 \pm 23.4
Gender		
Male	19,818 (49.1%)	24,353 (37.8%)
Female	17,711 (43.9%)	34,690 (53.9%)
Gender missing	2,814 (7%)	5315 (8.3%)
Age (year)		
FAERS report counts (<i>n</i>)	30,324 (75.2%)	46310 (71.9%)
Age missing	10019 (24.8%)	18046 (28%)

Age Mean \pm SD 56.1 \pm 18.33 48 \pm 19.53

A statistical difference between two DILI groups across categorical variables was performed using a chi-square test. Comparisons of continuous variables were performed using *t*-tests; *P* values were < 0.0001 for all the variables ($P < 0.05$ was considered significant); *n* is number of instances.

Table 4 Top 20 drugs with the highest reporting odds ratio in both DILI groups.

Drug class	Drug name	Severity class	Label section	Mitochondrial toxicity	ROR
Antigout agent	Benzbromarone	8	Withdrawn	Yes	36.31
Antidiabetic agent	Troglitazone	8	Withdrawn	Yes	31.02
Antimycobacterial	Isoniazid	8	Box warning	Yes	20.79
Antimycobacterial	Rifampin	8	Warnings and precautions	Yes	18.64
Antibiotics	Telithromycin	8	Warnings and precautions	No	18.33
Antineoplastics	Gemtuzumab ozogamicin	8	Box warning	No	18.08
Antiarrhythmics	Mexiletine	3	Box warning	No	17.8
Antineoplastics	Dactinomycin	8	Warnings and precautions	No	17.25
Anti-inflammatory agent	Nimesulide	8	Withdrawn	Yes	15.07
Antialcoholics	Disulfiram	8	Warnings and precautions	No	14.82
Antivirals	Didanosine	8	Warnings and precautions	Yes	14.38
Stimulants; central nervous system	Pemoline	8	Withdrawn	No	14.24
Platelet inhibitors	Ticlopidine	4	Warnings and precautions	No	13.51
Antibiotics	Trovafloxacin mesylate	8	Withdrawn	Yes	13.48
Antithyroid agents	Propylthiouracil	8	Box warning	No	13.33
NSAID	Bromfenac	8	Withdrawn	No	13.01
Antiretroviral drugs	Stavudine	8	Box warning	Yes	12.83
Hormone modifiers	Danazol	8	Box warning	Yes	12.82
Antiparkinson agents	Tolcapone	8	Box warning	Yes	12.25
Antivirals	Tipranavir	8	Box warning	No	12.04

Table 5 Reporting odds ratio estimate for hepatotoxic drugs associated with mitochondrial mechanisms of toxicity compared to non-mitochondrial mechanisms.

DILI group	Odds ratio	95% CI	<i>P</i> -value
Drugs associated with mitochondrial mechanisms of toxicity	1.43	1.42-1.45	< 0.0001

Table 6 Association of age and gender with hepatotoxic drugs with mitochondrial toxicity mechanisms as compared with non-mitochondrial mechanisms, using a univariate logistic regression model.

Independent variable	Odds ratio	95% CI	<i>P</i> -value
Age			
<65 years	Reference		
>65 years	2.2	2.12-2.26	< 0.0001
Gender			
Male	Reference		
Female	0.63	0.61-0.64	< 0.0001

Figures




Figure 1 Reporting odds ratios (RORs) for hepatotoxic drugs associated with mitochondrial mechanisms of toxicity. Benzbromarone, troglitazone, isoniazid, rifampin, and nimesulide had the highest ROR values in this group.




Figure 2 Reporting odds ratios (RORs) for hepatotoxic drugs associated with non-mitochondrial mechanisms of toxicity. Telithromycin, gemtuzumab ozogamicin, mexiletine, dactinomycin, and disulfiram had the highest ROR values in the group.




Figure 3 Sum of all ROR of “most-DILI-concern” drugs associated with mitochondrial and non-mitochondrial mechanisms of toxicity per therapeutic class. Drugs from the antineoplastic, antiviral, analgesic, antibiotic, and antimycobacterial classes were the top 5 drug classes associated with higher ROR scores. Drugs from the antidiabetic, antiretroviral, anti-inflammatory, anti-Parkinson, vasoactive, neuroprotective, and antihyperlipidemic drug classes were primarily associated with mitochondrial mechanisms. Alternatively, leukotriene pathway modulators, alcohol antagonists, CNS stimulants, and platelet inhibitor drug classes were the drugs with non-mitochondrial mechanisms having higher RORs.




Figure 4 Categorization based on liver injury drug label for “most-DILI concern” drugs based on their association with mitochondrial and non-mitochondrial mechanisms of toxicity. There was not any notable trend between two groups based on drug label.




Figure 5 Categorization based on liver injury severity class for “most-DILI concern” drugs based on their association with mitochondrial and non-mitochondrial

mechanisms of toxicity. There was not any notable trend between two groups based on severity class.

Journal Pre-proof