

## The clinical diagnosis of Achilles tendinopathy: a scoping review

Matthews, Wesley; Ellis, Richard; Furness, James; Hing, Wayne A

*Published in:*  
PEERJ

*DOI:*  
[10.7717/peerj.12166](https://doi.org/10.7717/peerj.12166)

*Licence:*  
CC BY

[Link to output in Bond University research repository.](#)

*Recommended citation(APA):*  
Matthews, W., Ellis, R., Furness, J., & Hing, W. A. (2021). The clinical diagnosis of Achilles tendinopathy: a scoping review. *PEERJ*, 9, [e12166]. <https://doi.org/10.7717/peerj.12166>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

# The clinical diagnosis of Achilles tendinopathy: a scoping review

Wesley Matthews<sup>1</sup>, Richard Ellis<sup>2,3</sup>, James Furness<sup>1</sup> and Wayne A. Hing<sup>1</sup>

<sup>1</sup> Bond Institute of Health and Sport, Faculty of Health Sciences and Medicine, Bond University, Gold Coast, Queensland, Australia

<sup>2</sup> Active Living and Rehabilitation: Aotearoa New Zealand, Health and Rehabilitation Research Institute, Faculty of Health and Environmental Sciences, Auckland University of Technology, Auckland, New Zealand

<sup>3</sup> Department of Physiotherapy, School of Clinical Sciences, Faculty of Health and Environmental Sciences, Auckland University of Technology, Auckland, New Zealand

## ABSTRACT

**Background:** Achilles tendinopathy describes the clinical presentation of pain localised to the Achilles tendon and associated loss of function with tendon loading activities. However, clinicians display differing approaches to the diagnosis of Achilles tendinopathy due to inconsistency in the clinical terminology, an evolving understanding of the pathophysiology, and the lack of consensus on clinical tests which could be considered the gold standard for diagnosing Achilles tendinopathy. The primary aim of this scoping review is to provide a method for clinically diagnosing Achilles tendinopathy that aligns with the nine core health domains.

**Methodology:** A scoping review was conducted to synthesise available evidence on the clinical diagnosis and clinical outcome measures of Achilles tendinopathy. Extracted data included author, year of publication, participant characteristics, methods for diagnosing Achilles tendinopathy and outcome measures.

**Results:** A total of 159 articles were included in this scoping review. The most commonly used subjective measure was self-reported location of pain, while additional measures included pain with tendon loading activity, duration of symptoms and tendon stiffness. The most commonly identified objective clinical test for Achilles tendinopathy was tendon palpation (including pain on palpation, localised tendon thickening or localised swelling). Further objective tests used to assess Achilles tendinopathy included tendon pain during loading activities (single-leg heel raises and hopping) and the Royal London Hospital Test and the Painful Arc Sign. The VISA-A questionnaire as the most commonly used outcome measure to monitor Achilles tendinopathy. However, psychological factors (PES, TKS and PCS) and overall quality of life (SF-12, SF-36 and EQ-5D-5L) were less frequently measured.

**Conclusions:** There is significant variation in the methodology and outcome measures used to diagnose Achilles tendinopathy. A method for diagnosing Achilles tendinopathy is proposed, that includes both results from the scoping review and recent recommendations for reporting results in tendinopathy.

Submitted 6 May 2021  
Accepted 25 August 2021  
Published 28 September 2021

Corresponding author  
Wesley Matthews,  
wesley.matthews@student.bond.edu.au

Academic editor  
Charles Okpala

Additional Information and  
Declarations can be found on  
page 39

DOI 10.7717/peerj.12166

© Copyright  
2021 Matthews et al.

Distributed under  
Creative Commons CC-BY 4.0

OPEN ACCESS

**Subjects** Evidence Based Medicine, Kinesiology, Orthopedics, Sports Injury, Sports Medicine

**Keywords** Tendinopathy, Diagnosis, Achilles, Tendon, Tendinosis, Tendinitis

## INTRODUCTION

Achilles tendinopathy describes the clinical presentation of pain localised to the Achilles tendon and associated loss of function with tendon loading activities ([De Vos et al., 2021](#); [Millar et al., 2021](#)). However, clinicians display differing approaches to the diagnosis of Achilles tendinopathy due to inconsistency in the clinical terminology, an evolving understanding of the pathophysiology, and the lack of consensus on clinical tests which could be considered the gold standard for diagnosing Achilles tendinopathy ([De Vos et al., 2021](#); [Millar et al., 2021](#); [Docking, Ooi & Connell, 2015](#); [Cook et al., 2016](#)). Conversely, when describing the clinical condition of persistent pain and dysfunction of the Achilles tendon in relation to mechanical loading, consensus agreement has identified the preferred terminology to be ‘tendinopathy’ rather than other common terms such as ‘tendinitis’ and ‘tendinosis’ ([Scott et al., 2020](#)). However, the consensus agreement for terminology does not provide a clear criteria with which to diagnose Achilles tendinopathy ([De Vos et al., 2021](#)).

Additionally, when considering the diagnosis of Achilles tendinopathy, distinctions can be made between the diagnosis of tendinopathy and clinical diagnosis of Achilles tendinopathy. As described by [Aggarwal et al. \(2015\)](#), a diagnosis is based off a broad set of signs and symptoms to reflect all the potential features and severity of a pathology. Whereas, a clinical diagnosis of Achilles tendinopathy requires a specific set of signs, symptoms and tests to define a homogenous group of patients across studies and geographical regions ([Aggarwal et al., 2015](#)). In the case of Achilles tendinopathy, the diagnosis of Achilles tendinopathy is determined by the presentation of pain localised to the Achilles tendon and associated loss of function with tendon loading activities ([De Vos et al., 2021](#); [Millar et al., 2021](#)). However, this broad description may include other pathological disease processes such as retrocalcaneal bursitis, complete or partial rupture of the Achilles, tarsal tunnel syndrome, neuroma/neuritis of the sural nerve, rupture posterior tibial tendon, or arthritic conditions of the ankle that need to be differentially diagnosed ([Hutchison et al., 2013](#)). Thus, it becomes relevant to understand the process to determine a clinical diagnosis of Achilles tendinopathy.

The clinical diagnosis of Achilles tendinopathy is predominantly derived from patient history, patient reported load related pain, and pain provocation tests ([Millar et al., 2021](#)). Patient history, localised Achilles tendon pain and pain on palpation are considered key to diagnosing Achilles tendinopathy ([De Vos et al., 2021](#); [Millar et al., 2021](#)) and can all be assessed reliably ([Hutchison et al., 2013](#)). Additional pain provoking tests; such as the single leg heel raise, hop test, Royal London Hospital Test or Painful Arc Sign; have been suggested as useful to confirm a clinical diagnosis of Achilles tendinopathy ([Millar et al., 2021](#); [Hutchison et al., 2013](#); [Reiman et al., 2014](#)). However, many leading researchers disagree on the which clinical tests are essential to diagnose Achilles tendinopathy ([De Vos et al., 2021](#)). Conversely, it is agreed that uniform diagnostic criteria would be useful in identifying possible subclassifications of Achilles tendinopathy and thus improving tailored individual treatment programmes or monitoring patient progress ([De Vos et al., 2021](#)).

**Table 1** The nine core health domains of tendinopathy as recommended by [Vicenzino et al. \(2020\)](#).

Domain	Description	Example
Patient rating of overall condition	A single assessment numerical evaluation	0–100%
Pain on activity or loading	Patient reported intensity of pain during a tendon loading activity.	VAS, NRS
Participation	Patient rating of participation levels in sport or engagement across other areas.	Tegner Activity Scale
Function	Patient rating of function and not referring to the intensity of their pain.	Patient Specific Function Scale
Psychological factors	Patient rating of psychological impact (e.g. Pain self efficacy, kinesiophobia, catastrophisation).	PCS
Disability	Scores from a combination of patient rated pain and disability due to pain in relation to tendon specific loading activities	VISA-A
Physical function capacity	The quantitative measures of physical tasks such as number of hops, number of squats and dynamometry.	Single leg heel raise
Quality of life	Patient rating of general wellbeing	EQ-5D
Pain over a specified time	Patient reported intensity of pain over a specified time period (e.g. morning, night, 24 h).	VAS, NRS

**Note:**

VAS, visual analogue scale; NRS, numerical rating scale; PCS, pain catastrophisation scale; VISA, Victorian Institute of Sport Assessment; EQ-5D, EuroQol-5 dimension.


Recently, [Vicenzino et al. \(2020\)](#) identified nine core health domains in tendinopathy following consensus agreement from both health care practitioners and patients. These included patient rating of overall condition, pain on activity or loading, participation, function, psychological factors, disability, physical function capacity, quality of life, and pain over a specified timeframe ([Vicenzino et al., 2020](#)). An overview of the nine core health domains of tendinopathy ([Vicenzino et al., 2020](#)) are presented in [Table 1](#). Using the determined core health domains, specific measures will need to be identified specific to Achilles tendinopathy ([Vicenzino et al., 2020](#)). The introduction of the nine core health domains in tendinopathy ([Vicenzino et al., 2020](#)) in addition to previously identified gaps in the literature, including; a lack of consistency in terminology used to diagnose Achilles tendinopathy ([De Vos et al., 2021; Scott et al., 2020](#)), lack of a consensus on the clinical diagnosis of Achilles tendinopathy ([De Vos et al., 2021](#)), and the need for a uniform method with which to clinically diagnose Achilles tendinopathy ([De Vos et al., 2021](#)). Thus there is a requirement to identify the methods with which these gaps can be addressed and allow for greater consistency in the clinical diagnosis of Achilles tendinopathy in both research and clinical practice.

Therefore, the primary aim of this scoping review is to provide a method for clinically diagnosing Achilles tendinopathy that aligns with the nine core health domains. In order to achieve this, specific objectives have been determined that include: (1) identifying the most common clinical tests used to diagnose Achilles tendinopathy, (2) identifying the most common outcome measures used to assess Achilles tendinopathy, and (3) summarising the studies to date.

## METHODOLOGY

### Study design

A scoping review was conducted to synthesise available evidence on the clinical diagnosis and clinical outcome measures of Achilles tendinopathy. Due to the wide-ranging nature

**Figure 1** Overall study design.

Full-size DOI: 10.7717/peerj.12166/fig-1

of the topic, a scoping review was used to facilitate the collection and charting of evidence with the aim of identifying key themes, knowledge gaps and types of evidence currently available. [Figure 1](#) provides an overview of the overall study design and process to answer the primary aim and specific objectives.

### Search strategy

A single researcher (WM) completed a literature search to identify, screen and select studies in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analysis Extension for Scoping Reviews (PRISMA-ScR) ([Tricco et al., 2018](#)). A detailed, multistep search of PubMed, CINAHL, ProQuest and SPORTDiscus was conducted between May 2020 and July 2020, before being updated in April 2021. In addition to the electronic database search, reference lists from included articles were reviewed for additional articles. To ensure a broad search, key words were truncated to allow for variations in spelling and combined using Boolean operators in addition to the use of MeSH terms to allow for review of all relevant articles. The full electronic search for the PubMed database is provided in [Table 2](#).

**Table 2** Electronic database search strategy example.

Database	Search strategy	Results
PubMed	(“tendineous”[All Fields] OR “tendinopathy”[MeSH Terms] OR “tendinopathy”[All Fields] OR “tendinitis”[All Fields] OR “tendons”[MeSH Terms] OR “tendons”[All Fields] OR “tendinous”[All Fields] OR (“tendinopathy”[MeSH Terms] OR “tendinopathy”[All Fields] OR “tendinopathy”[All Fields] OR “tendinosis”[All Fields]) OR (“tendinopathy”[MeSH Terms] OR “tendinopathy”[All Fields] OR “tendinopathies”[All Fields]) OR (“tendinopathy”[MeSH Terms] OR “tendinopathy”[All Fields] OR “tendonopathy”[All Fields]) OR (“tendinopathy”[MeSH Terms] OR “tendinopathy”[All Fields] OR “tendonitis”[All Fields] OR “tendon s”[All Fields] OR “tendonous”[All Fields] OR “tendons”[MeSH Terms] OR “tendons”[All Fields] OR “tendon”[All Fields] OR (“tendinopathy”[MeSH Terms] OR “tendinopathy”[All Fields] OR “tendonosis”[All Fields])) AND (“diagnosable”[All Fields] OR “diagnosi”[All Fields] OR “diagnosis”[MeSH Terms] OR “diagnosis”[All Fields] OR “diagnose”[All Fields] OR “diagnosed”[All Fields] OR “diagnoses”[All Fields] OR “diagnosing”[All Fields] OR “diagnosis”[MeSH Subheading]) AND (“achiles”[All Fields] OR “achille”[All Fields] OR “achille s”[All Fields] OR “achilles tendon”[MeSH Terms] OR (“achilles”[All Fields] AND “tendon”[All Fields]) OR “achilles tendon”[All Fields] OR “achilles”[All Fields])	7,162

## Eligibility criteria


Methods for data extraction specific to scoping reviews were informed by the Population-Concept-Context framework as recommended by the Joanna Briggs Institute (JBI) Reviewer's Manual ([Peters et al., 2020](#)). *Population* was defined as any person clinically diagnosed with Achilles tendinopathy regardless of location (insertional or midportion). *Concept* included any study reporting on the methods used to clinically diagnose Achilles tendinopathy including subjective measures, objective measures and outcome measures. *Context* included all periods of time, outcomes, comparators, follow-up, rehabilitation settings and duration and type of intervention.

Eligible articles were full-text and included original research, reviews, scoping reviews, systematic reviews, meta-analyses, case-series and clinical commentaries. Studies were included if they provided adequate information on the method of clinical diagnosis (either subjective measures, objective measures or both subjective and objective measures), and clinical outcome measures used. Studies were excluded if they were non-English, had no description of clinical diagnosis, not specific to Achilles tendinopathy or included asymptomatic Achilles tendon states only.

## Data extraction and synthesis

WM extracted data from publications meeting the inclusion criteria into an Excel spreadsheet. Data extraction, grouping and plotting were performed by WM in line with previously published recommendations ([Peters et al., 2020](#)), where extracted data included author, year of publication, participant characteristics, methods for diagnosing Achilles tendinopathy and outcome measures. Data was extracted in tabular and graphical forms with results grouped by study design and categorised according to the hierarchy of evidence ([Daly et al., 2006](#); [Evans, 2003](#); [Merlin, Weston & Tooher, 2009](#)). Diagnostic criteria were presented in tabular form including year of publication, population, subjective and objective measures. Terminology and outcome measures were presented in graphical form with terminology grouped by publication year and outcome measures grouped by purpose of measure (disability, pain, psychological, quality of life).

Following data extraction, data synthesis was performed according to a previously published methodological framework ([Thomas & Harden, 2008](#)). Data was synthesised



**Figure 2** Preferred Reporting Items for Systematic Reviews and Meta-analysis flow diagram.

Full-size DOI: [10.7717/peerj.12166/fig-2](https://doi.org/10.7717/peerj.12166/fig-2)

into the following categories: (1) subjective measures, (2) objective measures and (3) outcome measures. Results were plotted according to publication date, terminology, study design and clinical diagnostic measures. Results were then compared to the nine core health domains of tendinopathy (*Vicenzino et al., 2020*) to identify areas of overlap and gaps in the current evidence. Studies could be allocated to multiple groups. Quality appraisal was not required as per recommended methodology for scoping reviews (*Peters et al., 2020; Arksey & O'Malley, 2005*).

## RESULTS

### Selection of sources of evidence

The search results are displayed in the PRISMA Flow Diagram (Fig. 2). The search strategy generated 11,561 results with two further results identified via reference list searching. Following duplicate removal and title and abstract screening, 554 full-text articles were

reviewed for inclusion in the study. Of these, 395 were excluded for the following reasons: 240 provided insufficient information on the method of diagnosing Achilles tendinopathy, 11 assessed asymptomatic Achilles tendons only, 46 did not have access to the full text, 52 were not in English and 47 were not specific to Achilles tendinopathy. Thus, 159 articles (*Millar et al., 2021; Hutchison et al., 2013; Reiman et al., 2014; Abate & Salini, 2019; Aiyegbusi, Tella & Sanusi, 2020; Aldridge, 2004; Alfredson, 2003; Alfredson & Cook, 2007; Alfredson & Spang, 2018; Aronow, 2005; Asplund & Best, 2013; Azevedo et al., 2009; Bains & Porter, 2006; Barge-Caballero et al., 2008; Barker-Davies et al., 2017; Baskerville et al., 2018; Benazzo, Todesca & Ceciliani, 1997; Benito, 2016; Bhatty, Khan & Zubairy, 2019; Bjordal, Lopes-Martins & Iversen, 2006; Boesen et al., 2017; Borda & Selhorst, 2017; Brown et al., 2006; Garcia et al., 2010; Cassel et al., 2018; Chazan, 1998; Cheng, Zhang & Cai, 2016; Chester et al., 2008; Chimenti et al., 2016; Chimenti et al., 2017; Chimenti et al., 2020; Cook, Khan & Purdam, 2002; Coombes et al., 2018; Courville, Coe & Hecht, 2009; Creaby et al., 2017; Crill, Berlet & Hyer, 2014; De Jonge et al., 2011; De Marchi et al., 2018; Den Hartog, 2009; Divani et al., 2010; Docking et al., 2015; Duthon et al., 2011; Ebbesen et al., 2018; Eckenrode, Kietrys & Stackhouse, 2019; Feilmeier, 2017; Finnamore et al., 2019; Florit et al., 2019; Fredericson, 1996; Furia & Rompe, 2007; Gärdin et al., 2016; Gatz et al., 2020; Habets et al., 2017; Hasani et al., 2020; Hernández-Sánchez et al., 2018; Holmes & Lin, 2006; Horn & McCollum, 2015; Hu & Flemister, 2008; Hutchison et al., 2011; Irwin, 2010; Järvinen et al., 2001; Jayaseelan, Weber & Jonely, 2019; Jewson et al., 2017; Jowett, Richmond & Bedi, 2018; Jukes, Scott & Solan, 2020; Kader et al., 2002; Karjalainen et al., 2000; Khan et al., 2003; Knobloch, 2007; Knobloch et al., 2007; Knobloch et al., 2008; Kragsnaes et al., 2014; Krogh et al., 2016; Lakshmanan & O'Doherty, 2004; Leung & Griffith, 2008; Lohrer & Nauck, 2009; Longo et al., 2009; Longo, Ronga & Maffulli, 2009; Maffulli, Giai Via & Oliva, 2015; Maffulli, Giuseppe Longo & Denaro, 2012; Maffulli & Kader, 2002; Maffulli et al., 2003; Maffulli et al., 2011; Maffulli et al., 2020; Maffulli et al., 2012; Maffulli et al., 2008; Maffulli et al., 2019; Maffulli, Sharma & Luscombe, 2004; Maffulli, Via & Oliva, 2014; Maffulli et al., 2008; Mafi, Lorentzon & Alfredson, 2001; Magnussen, Dunn & Thomson, 2009; Mansur et al., 2019; Mansur et al., 2017; Mantovani et al., 2020; Martin et al., 2018; Mayer et al., 2007; McCormack et al., 2015; McShane, Ostick & McCabe, 2007; Murawski et al., 2014; Nadeau et al., 2016; Neeter et al., 2003; Nichols, 1989; De Mesquita et al., 2018; O'Neill et al., 2019; Oloff et al., 2015; Ooi et al., 2015; Paavola et al., 2002; Paavola et al., 2002; Paavola et al., 2000; Paoloni et al., 2004; Papa, 2012; Pedowitz & Beck, 2017; Petersen, Welp & Rosenbaum, 2007; Pingel et al., 2013; Post et al., 2020; Praet et al., 2018; Rabello et al., 2020; Rasmussen et al., 2008; Reid et al., 2012; Reiter et al., 2004; Romero-Morales et al., 2019a; Rompe, Furia & Maffulli, 2009; Rompe et al., 2008; Rompe et al., 2007; Roos et al., 2004; Ryan et al., 2009; Saini et al., 2015; Santamato et al., 2019; Sayana & Maffulli, 2007; Scholes et al., 2018; Scott, Huisman & Khan, 2011; Sengkerij et al., 2009; Sharma & Maffulli, 2006; Silbernagel et al., 2007; Silbernagel et al., 2001; Simpson & Howard, 2009; Solomons et al., 2020; Sorosky et al., 2004; Stenson et al., 2018; Stergioulas et al., 2008; Syvertson et al., 2017; Tan & Chan, 2008; Thomas et al., 2010; Turner et al., 2020; Vallance et al., 2020; Van der Vlist et al., 2020; Van der Vlist et al., 2020; Van Sterkenburg et al., 2011;*

*Verrall, Schofield & Brustad, 2011; Von Wehren et al., 2019; Wang et al., 2012; Wei et al., 2017; Welsh & Clodman, 1980; Xu et al., 2019; Zellers et al., 2019; Zhang et al., 2017; Zhang et al., 2020; Zhuang et al., 2019; Romero-Morales et al., 2019b*) were included in this scoping review.

### Characteristics of sources of evidence

In grouping the included articles by publication type, narrative reviews were the most common (27.2%) followed by cohort studies (19.6%), case control studies (18.8%), randomised controlled trials (12.7%), cross-sectional studies (10.8%), case reports (3.8%), protocols (3.2%), systematic reviews (1.9%), clinical guidelines (1.9%) and one consensus statement (0.6%). The years of publication of included studies ranged from 1980 to 2021, with 2017 to 2020 producing the most publications. [Table 3](#) provides the general characteristics of the reviewed studies, including year of publication, type of publication, terminology and tendinopathy location.

As highlighted in [Fig. 3](#), the terminology used to describe tendon pain varied, with ‘tendinopathy’ being the most prevalent term used to describe tendon pain. Thus, during this scoping review, tendinopathy, will be used to describe pain located in the Achilles tendon that impairs function.

### Results of individual sources of evidence

#### Clinical guidelines and consensus statements

Two of the included clinical guidelines ([Garcia et al., 2010](#); [Martin et al., 2018](#)) discussed midportion Achilles tendinopathy, with one clinical guideline ([Thomas et al., 2010](#)) and one consensus statement ([Xu et al., 2019](#)) discussing insertional Achilles tendinopathy ([Table 4](#)). Clinical measures used to diagnose Achilles tendinopathy was consistent across the clinical guidelines and consensus statement, with location of pain being the main differentiating factor between diagnosing midportion or insertional tendinopathy. Common methods with which midportion tendinopathy was diagnosed included subjective reporting of pain located in the Achilles tendon 2–6 cm above the calcaneal insertion that is increased with tendon loading and reported tendon stiffness. Similarly, insertional tendinopathy was diagnosed *via* subjective reporting of pain and swelling at the calcaneal insertion of the Achilles tendon. Pain on palpation was utilised to confirm clinical diagnosis in both midportion and insertional tendinopathy. While additional objective tests for midportion tendinopathy included the ‘Painful Arc Sign’ and ‘Royal London Hospital Test’.

#### Systematic reviews

All three included systematic reviews assessed midportion Achilles tendinopathy ([Table 5](#)) ([Reiman et al., 2014](#); [Hutchison et al., 2011](#); [Magnussen, Dunn & Thomson, 2009](#)). Subjective reporting of pain with tendon loading was included as a diagnostic feature of midportion Achilles tendinopathy in all three systematic reviews ([Leung & Griffith, 2008](#); [Hutchison et al., 2011](#); [Magnussen, Dunn & Thomson, 2009](#)). Two of the systematic reviews ([Reiman et al., 2014](#); [Hutchison et al., 2011](#)) identified the location of tendon pain

**Table 3** Characteristics of included studies.

Characteristics	No. of studies (n)	References
<b>Year of publication</b>		
Before 1990	2	(Nichols, 1989; Welsh & Clodman, 1980)
1990–1999	3	(Benazzo, Tedesca & Ceciliani, 1997; Chazan, 1998; Frederickson, 1996)
2000–2009	59	(Aldridge, 2004; Alfredson, 2003; Alfredson & Cook, 2007; Aronow, 2005; Azeredo et al., 2009; Bains & Porter, 2006; Barge-Caballero et al., 2008; Bjordal, Lopes-Martins & Iversen, 2006; Brown et al., 2006; Chester et al., 2008; Cook, Khan & Purdam, 2002; Courville, Coe & Hecht, 2009; Den Hartog, 2009; Furia & Rompe, 2007; Holmes & Lin, 2006; Hu & Flemister, 2008; Järvinen et al., 2001; Kader et al., 2002; Karjalainen et al., 2000; Khan et al., 2003; Knobloch, 2007; Knobloch et al., 2007; Knobloch, 2008; Lakshmanan & O'Doherty, 2004; Leung & Griffith, 2008; Lohrer & Nauck, 2009; Longo et al., 2009; Longo, Ronga & Maffulli, 2009; Maffulli et al., 2003; Maffulli, Sharma & Luscombe, 2004; Maffulli et al., 2008; Maffulli, Sharma & McCabe, 2007; Neeter et al., 2003; Mafi, Lorentzon & Alfredson, 2001; Magnussen, Dunn & Thomson, 2009; Mayer et al., 2007; McShane, Ostick & McCabe, 2007; Rasmussen et al., 2008; Reiter et al., 2004; Paavola et al., 2002; Paavola et al., 2000; Paoloni et al., 2004; Petersen, Welp & Rosenbaum, 2007; Rasmussen et al., 2008; Reiter et al., 2004; Rompe, Furia & Maffulli, 2009; Rompe et al., 2008; Rompe et al., 2007; Roos et al., 2004; Ryan et al., 2009; Sayana & Maffulli, 2007; Sengheraj et al., 2009; Sharma & Maffulli, 2006; Silbernagel et al., 2007; Silbernagel et al., 2001; Simpson & Howard, 2009; Sorosky et al., 2004; Stergioulas et al., 2008; Tan & Chan, 2008)
2010–2019	79	(Hutchison et al., 2013; Reiman et al., 2014; Abate & Salini, 2019; Alfredson & Spang, 2018; Asplund & Best, 2013; Barker-Davies et al., 2017; Baskerville et al., 2018; Benito, 2016; Bhatty, Khan & Zubairy, 2019; Boesen et al., 2017; Borda & Sellhorst, 2017; Carića et al., 2010; Cassel et al., 2018; Cheng, Zhang & Cai, 2016; Chimenti et al., 2016; Chimenti et al., 2017; Coombes et al., 2018; Creaby et al., 2017; Grill, Berlet & Hyer, 2014; De Jonge et al., 2011; De Marchi et al., 2018; Divani et al., 2010; Docking et al., 2015; Duthon et al., 2011; Ebkesen et al., 2018; Ekenrode, Kietrys & Stackhouse, 2019; Feilmeyer, 2017; Finnimore et al., 2019; Florit et al., 2019; Gärdin et al., 2017; Habets et al., 2016; Hernández-Sánchez et al., 2018; Horn & McCollum, 2015; Hutchison et al., 2011; Irwin, 2010; Jayaseelan, Weber & Jones, 2019; Jewson et al., 2017; Jowett, Richmond & Bedi, 2018; Kragsnaes et al., 2014; Krogh et al., 2016; Maffulli, Giai Via & Oliva, 2015; Maffulli, Giuseppe Longo & Denaro, 2012; Maffulli et al., 2012; Maffulli et al., 2013; Maffulli, Via & Oliva, 2014; Mansur et al., 2019; Mansur et al., 2017; Martin et al., 2018; McCormack et al., 2015; Murawski et al., 2014; Nadeau et al., 2016; De Mesquita et al., 2018; O'Neill et al., 2019; Oloff et al., 2015; Ooi et al., 2015; Papa, 2012; Pedowitz & Beck, 2017; Pingel et al., 2013; Praet et al., 2018; Reid et al., 2012; Romero-Morales et al., 2019a; Saini et al., 2015; Santamato et al., 2019; Scholes et al., 2019; Stenson et al., 2018; Syverton et al., 2017; Thomas et al., 2010; Van Sterkenburg et al., 2011; Verrell, Schofield & Brustad, 2011; Von Wehren et al., 2019; Wang et al., 2012; Wei et al., 2017; Xu et al., 2019; Zellers et al., 2019; Zhang et al., 2017; Zhuang et al., 2019; Romero-Morales et al., 2019b)
2020–2021	16	(Millar et al., 2021; Ajegbusi, Tella & Sanusi, 2020; Chimenti et al., 2020; Gatz et al., 2020; Hasani et al., 2020; Jukes, Scott & Solan, 2020; Maffulli et al., 2020; Mantovani et al., 2020; Post et al., 2020; Rabello et al., 2020; Solomonis et al., 2020; Turner et al., 2020; Vallance et al., 2020; Van der Vlist et al., 2020; Van der Vlist et al., 2020; Zhang et al., 2020)
<b>Type of publication</b>		
Clinical guidelines	3	(Garcia et al., 2010; Martin et al., 2018; Thomas et al., 2010)
Consensus statement	1	(Xu et al., 2019)
Systematic reviews	3	(Reiman et al., 2014; Hutchison et al., 2011; Magnusson, Dunn & Thomson, 2009)
RCT	20	(Bjordal, Lopes-Martins & Iversen, 2006; Boesen et al., 2017; Brown et al., 2006; Ebbesen et al., 2018; Gatz et al., 2020; Knobloch et al., 2007; Krogh et al., 2016; Mafi, Lorentzon & Alfredson, 2001; Mayer et al., 2007; Paoloni et al., 2004; Petersen, Welp & Rosenbaum, 2007; Rasmussen et al., 2008; Rompe, Furia & Maffulli, 2009; Rompe et al., 2008; Rompe et al., 2007; Roos et al., 2004; Silbernagel et al., 2007; Stenson et al., 2018; Syverton et al., 2017; Von Wehren et al., 2019; Wei et al., 2019)
Cohort studies	31	(Alfredson & Spang, 2018; Barge-Caballero et al., 2008; Cheng, Zhang & Cai, 2016; Chester et al., 2008; Crill, Berlet & Hyer, 2014; Duthon et al., 2011; Florit et al., 2019; Jowett, Richmond & Bedi, 2018; Karjalainen et al., 2000; Khan et al., 2003; Knobloch, 2007; Knobloch et al., 2008; Maffulli et al., 2008; Mansur et al., 2019; McCormack et al., 2015; Murawski et al., 2014; O'Neill et al., 2019; Oloff et al., 2015; Paavola et al., 2002; Paavola et al., 2000; Sayana & Maffulli, 2007; Silbernagel et al., 2007; Stenson et al., 2007; Syverton et al., 2018; Von Wehren et al., 2017; Wei et al., 2019; Wei et al., 2019; Zhang et al., 2020; Zhuang et al., 2019)

(Continued)


Table 3 (continued)

Characteristics	No. of studies (n)	References
Case control studies	30	(Hutchison et al., 2013; Abate & Salini, 2019; Azevedo et al., 2009; Cassel et al., 2018; Chimenti et al., 2016; Coombes et al., 2020; Coombes et al., 2018; Creaby et al., 2017; Eckenrode, Kietrys & Stackhouse, 2019; Gärdin et al., 2016; Hernández-Sánchez et al., 2018; Holmes & Lin, 2006; Jewson et al., 2017; Leung & Griffith, 2008; Lohrer & Nauk, 2009; Maffulli et al., 2003; Nadeau et al., 2016; Neeter et al., 2003; De Mesquita et al., 2018; Ooi et al., 2015; Pingel et al., 2013; Rabello et al., 2020; Reid et al., 2012; Reiter et al., 2004; Romero-Morales et al., 2019a; Ryan et al., 2009; Sengkerij et al., 2009; Verrall, Schofield & Brustad, 2011; Zhang et al., 2017; Romero-Morales et al., 2019b)
Cross-sectional studies	17	(Aiyegbusi, Tella & Samusi, 2020; De Jonge et al., 2011; De Marchi et al., 2018; Divani et al., 2010; Docking et al., 2015; Finnimore et al., 2019; Kragsnaes et al., 2014; Longo et al., 2009; Maffulli et al., 2008; Mantovani et al., 2020; Praet et al., 2018; Santamato et al., 2019; Scholes et al., 2018; Turner et al., 2020; Vallance et al., 2020; Van der Vlist et al., 2020; Wang et al., 2012)
Narrative reviews	43	(Millar et al., 2021; Alldridge, 2004; Alfredson, 2003; Alfredson & Cook, 2007; Aronow, 2005; Asplund & Best, 2013; Bains & Porter, 2006; Baskerville et al., 2018; Benazzo, Todessa & Cecilliani, 1997; Blatty, Khan & Zukairy, 2019; Chimenti, 1998; Chimenti et al., 2017; Cook, Khan & Purdam, 2002; Courville, Coe & Hecht, 2009; Den Hartog, 2009; Feilheimer, 2017; Fredericson, 1996; Faria & Rompe, 2007; Horn & McCollum, 2015; Hu & Flemister, 2008; Irwin, 2010; Järvinen et al., 2001; Jukes, Scott & Solan, 2020; Kader et al., 2002; Longo, Ronga & Maffulli, 2009; Maffulli, Gai & Oliva, 2015; Maffulli, Giuseppe Longo & Denaro, 2012; Maffulli et al., 2012; Maffulli, Sharma & Luscombe, 2004; Maffulli, Via & Oliva, 2004; Pedowitz & Beck, 2017; Saini et al., 2015; Scott, Huisman & Khan, 2011; Sharma & Maffulli, 2006; Simpson & McCabe, 2007; Nichols, 1989; Paavola et al., 2002; Tan & Chan, 2008) (Benito, 2016; Borda & Sellhorst, 2017; Jayaseelan, Weber & Jones, 2019; Maffulli et al., 2011; Papa, 2012; Van Sterkenburg et al., 2011) (Barker-Davies et al., 2017; Habets et al., 2017; Hasani et al., 2020; Mansur et al., 2017; Post et al., 2020)
Case reports	6	
Protocols	5	
Terminology		
Tendon pain	2	(Neeter et al., 2003; Silbernagel et al., 2001)
Tendinitis	3	(Alldridge, 2004; Paavola et al., 2000; Welsh & Clodman, 1980)
Tendinosis	3	(Gärdin et al., 2016; Karjalainen et al., 2000; Wei et al., 2017)
Tendinopathy	144	(Millar et al., 2021; Hutchison et al., 2013; Reiman et al., 2014; Abate & Salini, 2019; Ayiegbusi, Tella & Sanusi, 2020; Alfredson, 2003; Alfredson & Cook, 2007; Alfredson & Spang, 2018; Aronow, 2005; Asplund & Best, 2013; Azevedo et al., 2009; Bains & Porter, 2006; Barge-Caballero et al., 2008; Barker-Davies et al., 2017; Baskerville et al., 2018; Benazzo, Todessa & Cecilliani, 1997; Benito, 2016; Bhatty, Khan & Zubairy, 2019; Bjordal, Lopes-Martins & Iversen, 2006; Boesen et al., 2017; Borda & Sellhorst, 2017; Brown et al., 2006; Carcia et al., 2010; Casel et al., 2010; Cook, Khan & Purdam, 2002; Coombes et al., 2018; Courville, Coe & Hecht, 2009; Creaby et al., 2016; Chimenti et al., 2017; Chimenti et al., 2020; Cook, Khan & Purdam, 2002; Coombes et al., 2018; Courville, Coe & Hecht, 2009; Creaby et al., 2017; Grill, Berlet & Hyer, 2014; De Jonge et al., 2011; De Marchi et al., 2018; Divani et al., 2010; Docking et al., 2015; Duthon et al., 2011; Ebbesen et al., 2018; Eckenrode, Kietrys & Stackhouse, 2019; Feilheimer, 2017; Finnimore et al., 2019; Florit et al., 2019; Furia & Rompe, 2007; Gatz et al., 2020; Habets et al., 2017; Hasani et al., 2020; Hernández-Sánchez et al., 2018; Holmes & Lin, 2006; Horn & McCollum, 2015; Hu & Flemister, 2008; Hutchison et al., 2011; Irwin, 2010; Järvinen et al., 2001; Jayasedan, Weber & Jones, 2019; Jewson et al., 2017; Jowett, Richmond & Bedi, 2018; Jukes, Scott & Solan, 2020; Kader et al., 2002; Khan et al., 2003; Knoblock et al., 2007; Knoblock et al., 2008; Kragsnaes et al., 2014; Krogh et al., 2016; Lakshmanan & O'Doherty, 2004; Leung & Griffith, 2008; Lohrer & Nuack, 2009; Longo et al., 2009; Maffulli, Gai & Oliva, 2015; Maffulli, Giuseppe Longo & Denaro, 2012; Maffulli & Kader, 2002; Maffulli et al., 2003; Maffulli et al., 2011; Maffulli et al., 2020; Maffulli et al., 2012; Maffulli et al., 2013; Post et al., 2020; Praet et al., 2018; Rabello et al., 2020; Rasmussen et al., 2008; Reid et al., 2012; Reiter et al., 2004; Romero-Morales et al., 2019a; Rompe, Faria & Maffulli, 2009; Rompe et al., 2008; Rompe et al., 2007; Roos et al., 2004; Ryan et al., 2009; Saini et al., 2015; Santamato et al., 2019; Sayana & Maffulli, 2007; Scholes et al., 2018; Sengkerij et al., 2009; Sharma & Maffulli, 2006; Silbernagel et al., 2007; Simpson & Howard, 2009; Solomons et al., 2020; Soroky et al., 2004; Stenson et al., 2008; Stergioulas et al., 2008; Syvertsen et al., 2017; Tan & Chan, 2008; Thomas et al., 2010; Turner et al., 2020; Vallance et al., 2020; Van der Vlist et al., 2020; Van Sterkenburg et al., 2011; Verrall, Schofield & Brustad, 2011; Von Wehren et al., 2019; Xu et al., 2019; Zellers et al., 2019; Zhang et al., 2017; Zhang et al., 2020; Zhuang et al., 2019; Romero-Morales et al., 2019b)

**Table 3** (continued)

Characteristics	No. of studies (n)	References
Combined terminology	7	(Chazan, 1998; Den Hartog, 2009; Fredericson, 1996; Nichols, 1989; Pedowitz & Beck, 2017; Scott, Huisman & Khan, 2011; Wang et al., 2012)
Tendinopathy location		
Insertional	21	(Aldridge, 2004; Benazzo, Tedesca & Cecilianni, 1997; Benito, 2016; Cheng, Zhang & Cai, 2016; Chimenti et al., 2016; Chimenti et al., 2017; Den Hartog, 2009; Hu & Flemister, 2008; Irwin, 2010; Mafulli et al., 2019; Mansur et al., 2019; Mansur et al., 2017; McCormack et al., 2015; Rompe et al., 2008; Stenson et al., 2018; Thomas et al., 2010; Wei et al., 2017; Xu et al., 2019; Zellers et al., 2019; Zhang et al., 2020; Zhuang et al., 2019)
Midportion	83	(Hutchison et al., 2013; Reiman et al., 2014; Abate & Salini, 2019; Alfredson, 2003; Alfredson & Cook, 2007; Azevedo et al., 2009; Bains & Porter, 2006; Boesen et al., 2017; Borda & Sellhorst, 2017; Brown et al., 2006; Garcia et al., 2010; Chester et al., 2008; Courville, Coe & Hecht, 2009; Creaby et al., 2017; Crill, Berlet & Hyer, 2014; De Jonge et al., 2011; De Marchi et al., 2018; Divani et al., 2010; Duthon et al., 2011; Feltmeier, 2017; Finnimore et al., 2019; Gärden et al., 2016; Habets et al., 2017; Hasani et al., 2020; Hutchison et al., 2011; Järvinen et al., 2001; Jayaseelan, Weber & Jones, 2019; Jowett, Richmond & Bedi, 2018; Kader et al., 2002; Knobloch et al., 2008; Krogh et al., 2016; Lakshmanan & O'Doherty, 2004; Lohrer & Nauck, 2009; Longo, Ronga & Maffulli, 2009; Maffulli, Giuseppe Longo & Denaro, 2012; Maffulli & Kader, 2002; Maffulli et al., 2003; Maffulli et al., 2011; Maffulli et al., 2008; Maffulli, Sharma & Luscombe, 2004; Maffulli, Via & Oliva, 2014; Maffulli et al., 2008; Mafi, Lorenzon & Alfredson, 2001; Magnussen, Dunn & Thomson, 2009; Martin et al., 2018; Mayer et al., 2007; McShane, Ostik & McCabe, 2007; Murawski et al., 2014; Nadeau et al., 2016; Neeter et al., 2019; Paavola et al., 2002; Paavola et al., 2002; Paavola et al., 2000; Paoloni et al., 2004; Papa, 2012; Petersen, Welp & Rosenbaum, 2007; Pingel et al., 2013; Praet et al., 2018; Reid et al., 2012; Romero-Morales et al., 2019a; Rompe, Furia & Maffulli, 2009; Rompe et al., 2007; Roos et al., 2004; Ryan et al., 2009; Santamato et al., 2019; Sayana & Maffulli, 2007; Scholes et al., 2018; Scott, Huisman & Khan, 2011; Sengenjir et al., 2009; Sharma & Maffulli, 2006; Silbernagel et al., 2007; Simpson & Howard, 2009; Solomons et al., 2020; Stergioulas et al., 2008; Syverson et al., 2017; Tan & Chan, 2008; Van der Vlist et al., 2020; Van der Vlist et al., 2020; Van Sterkenburg et al., 2011; Von Wehren et al., 2019; Wang et al., 2012; Romero-Morales et al., 2019b)
Both	35	(Alfredson & Spang, 2018; Antonow, 2005; Asplund & Best, 2013; Barker-Davies et al., 2017; Bhatty, Khan & Zubairy, 2019; Chimenti et al., 2020; Cook, Khan & Purdam, 2002; Combes et al., 2018; Docking et al., 2015; Eckenrode, Kietrys & Stackhouse, 2019; Fredericson, 1996; Furia & Rompe, 2007; Gatz et al., 2020; Hernández-Sánchez et al., 2018; Horn & McCollum, 2015; Jukes, Scott & Solan, 2020; Karjalainen et al., 2000; Khan et al., 2003; Knobloch, 2007; Knobloch et al., 2007; Krasnues et al., 2014; Maffulli, Giai Via & Oliva, 2015; Maffulli et al., 2020; Ooi et al., 2015; Pedowitz & Beck, 2017; Post et al., 2020; Rabello et al., 2020; Reiter et al., 2004; Saini et al., 2015; Turner et al., 2020; Vallance et al., 2020; Verrall, Schofield & Brustad, 2011; Welsh & Clohnan, 1980; Zhang et al., 2017)
Not specified	20	(Millar et al., 2021; Aiyegbusi, Tella & Sanusi, 2020; Baskerville et al., 2008; Barge-Caballero et al., 2018; Bjordal, Lopes-Martins & Iversen, 2006; Cassel et al., 2018; Ebbesen et al., 2018; Florit et al., 2019; Holmes & Lin, 2006; Jewson et al., 2017; Leung & Griffith, 2008; Longo et al., 2009; Maffulli et al., 2012; Mantovani et al., 2020; Nichols, 1989; De Mesquita et al., 2018; Oloff et al., 2015; Rasmussen et al., 2001; Sorosky et al., 2004)

**Note:***n*, number; RCT, randomised controlled trial.



**Figure 3** Terminology used to describe the clinical presentation of Achilles tendon pain and impaired function.  
[Full-size](#) DOI: 10.7717/peerj.12166/fig-3

**Table 4** Clinical guidelines and consensus statement.

Author	Year	Location	Subjective history	Clinical tests
<i>Garcia et al. (2010)</i>	2010	Midportion	Location of pain (2–6 cm above calcaneal insertion) Pain with tendon loading Tendon stiffness	Pain on palpation Painful Arc Sign Royal London Hospital Test Single-leg heel Raise Hopping
<i>Thomas et al. (2010)</i>	2010	Insertional	Location of pain (insertion) Pain with tendon loading Swelling	Pain on palpation Localised tendon thickening on palpation
<i>Martin et al. (2018)</i>	2018	Midportion	Location of pain (2–6 cm above calcaneal insertion) Pain with tendon loading Tendon stiffness	Pain on palpation Painful Arc Sign Royal London Hospital Test
<i>Xu et al. (2019)</i>	2019	Insertional	Location of pain (insertion) Pain with tendon loading	Pain on palpation Localised swelling on palpation Pain with active dorsiflexion Silverskiold Test

**Note:**

cm, centimetres.

as 2–6 cm above the calcaneal insertion, with one (*Magnussen, Dunn & Thomson, 2009*) defining the location of tendon pain as 2–7 cm above the calcaneal insertion. Palpation of the Achilles tendon, passive dorsiflexion, pain with single-leg heel raise and pain hopping or jumping were included as clinical tests in all included systematic reviews (*Reiman et al., 2014; Hutchison et al., 2011; Magnussen, Dunn & Thomson, 2009*). Two of the systematic reviews (*Reiman et al., 2014; Hutchison et al., 2011*) included the ‘Painful Arc Sign’ and ‘Royal London Hospital Test’ as diagnostic measures for midportion Achilles tendinopathy.

#### Randomised controlled trials

**Table 6** highlights the characteristics of the included randomised controlled trials. Thirteen of the included studies (*Boesen et al., 2017; Brown et al., 2006; Krogh et al., 2016*;

**Table 5** Systematic reviews.

Author	Year	Sample size	Location	Subjective history	Clinical tests
Magnussen, Dunn & Thomson (2009)	2009	677 (M/F = 347/330)	Midportion	Location of pain (2–7 cm above calcaneal insertion) Pain with tendon loading	Pain on palpation Localised swelling on palpation Localised tendon thickening on palpation Pain with passive dorsiflexion Single-leg heel raise Hopping
Hutchison et al. (2011)	2011	578 (M/F = not specified)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Pain with tendon loading Tendon stiffness	Pain on palpation Localised swelling on palpation Localised tendon thickening on palpation Painful Arc Sign Royal London Hospital Test Reduced dorsiflexion Single-leg heel Raise Jump test
Reiman et al. (2014)	2014	31 (M/F = 27/4)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Pain with tendon loading Tendon stiffness	Pain on palpation Localised swelling on palpation Localised tendon thickening on palpation Painful Arc Sign Royal London Hospital Test Pain with dorsiflexion Single-leg heel Raise Hopping

**Note:**

cm, centimetres; M, male; F, female.

*Mafi, Lorentzon & Alfredson, 2001; Mayer et al., 2007; Paoloni et al., 2004; Petersen, Welp & Rosenbaum, 2007; Rompe, Furia & Maffulli, 2009; Rompe et al., 2007; Roos et al., 2004; Solomons et al., 2020; Stergioulas et al., 2008; Van der Vlist et al., 2020*) investigated midportion Achilles tendinopathy, one study (*Rompe et al., 2008*) investigated insertional Achilles tendinopathy, two studies (*Gatz et al., 2020; Knobloch et al., 2007*) investigated both insertional and midportion Achilles tendinopathy, and four studies (*Bjordal, Lopes-Martins & Iversen, 2006; Ebbesen et al., 2018; Rasmussen et al., 2008; Silbernagel et al., 2001*) did not specify a location of interest. All of the included randomised controlled trials used location of pain as a diagnostic feature of Achilles tendinopathy. Eight of the studies (*Knobloch et al., 2007; Mafi, Lorentzon & Alfredson, 2001; Paoloni et al., 2004; Petersen, Welp & Rosenbaum, 2007; Rompe, Furia & Maffulli, 2009; Rompe et al., 2008; Roos et al., 2004; Stergioulas et al., 2008*) which assessed midportion tendinopathy defined the location of pain as 2–6 cm above the calcaneal insertion, with three studies (*Boesen et al., 2017; Krogh et al., 2016; Van der Vlist et al., 2020*) defining midportion tendinopathy as 2–7 cm above the calcaneal insertion. Of the included studies, 17 included symptom duration as part of their diagnostic criteria, with various durations including four weeks (*Roos et al., 2004*), six weeks (*Brown et al., 2006*), two months (*Gatz et al., 2020; Van der Vlist et al., 2020*), three months (*Boesen et al., 2017; Ebbesen et al., 2018; Mafi, Lorentzon & Alfredson, 2001; Paoloni et al., 2004; Petersen, Welp & Rosenbaum, 2007*);

**Table 6** Randomised controlled trials.

Author	Year	Sample size	Location	Subjective history	Clinical tests
Mafi, Lorentzon & Alfredson (2001)	2001	44 (M/F = 24/20)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Duration of symptoms (>3 months)	Pain on palpation
Silbernagel et al. (2001)	2001	49 (M/F = 36/13)	Not specified	Location of pain Duration of symptoms (>3 months)	Pain on palpation Single leg heel raise Hopping Range of motion
Paoloni et al. (2004)	2004	65 (M/F = 40/25)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Duration of symptoms (>3 months) Gradual onset of pain	Pain on palpation Localised tendon thickening on palpation Hopping
Roos et al. (2004)	2004	44 (M/F = 21/23)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Duration of symptoms (>4 weeks) Pain with tendon loading	Pain on palpation
Bjordal, Lopes-Martins & Iversen (2006)	2006	7 (M/F = not specified)	Not specified	Location of pain Pain with tendon loading	Pain on palpation Hopping
Brown et al. (2006)	2006	26 (M/F = 17/9)	Midportion	Location of pain Duration of symptoms (>6 weeks) Gradual onset of pain Pain with tendon loading	Pain on palpation Single-leg heel raise Hopping
Knobloch et al. (2007)	2007	20 (M/F = 11/9)	Insertional Midportion	Location of pain (2–6 cm above calcaneal insertion) Location of pain (insertion) Pain with tendon loading Swelling	Not specified
Mayer et al. (2007)	2007	31 (M/F = 31/0)	Midportion	Location of pain Duration of symptoms (>6 months) Pain with tendon loading	Pain on palpation Localised tendon thickening on palpation
Petersen, Welp & Rosenbaum (2007)	2007	100 (M/F = 60/40)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Duration of symptoms (>3 months) Gradual onset of pain Pain with tendon loading	Pain on palpation Localised tendon thickening on palpation
Rompe et al. (2007)	2007	75 (M/F = 29/46)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Duration of symptoms (>6 months) Pain with tendon loading Swelling	Not specified
Rasmussen et al. (2008)	2008	48 (M/F = 28/20)	Not specified	Location of pain Duration of symptoms (>3 months)	Pain on palpation Localised swelling on palpation Pain with dorsiflexion
Rompe et al. (2008)	2008	50 (M/F = 20/30)	Insertional	Location of pain Pain with tendon loading Duration of symptoms (>6 months)	Pain on palpation Painful Arc Sign Royal London Hospital Test
Stergioulas et al. (2008)	2008	40 (M/F = 25/15)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Duration of symptoms (>6 months) Pain with tendon loading	Pain on palpation Reduced active dorsiflexion

**Table 6** (continued)

Author	Year	Sample size	Location	Subjective history	Clinical tests
Rompe, Furia & Maffulli (2009)	2009	68 (M/F = 30/38)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Duration of symptoms (>6 months) Pain with tendon loading Swelling	Not specified
Krogh et al. (2016)	2016	24 (M/F = 13/11)	Midportion	Location of pain (2–7 cm above calcaneal insertion) Pain with tendon loading	Pain on palpation Localised tendon thickening on palpation
Boesen et al. (2017)	2017	60 (M/F = 60/0)	Midportion	Location of pain (2–7 cm above calcaneal insertion) Duration of symptoms (>3 months)	Pain on palpation Localised tendon thickening on palpation Single-leg heel raise
Ebbesen et al. (2018)	2018	44 (M/F = 25/19)	Not specified	Location of pain Duration of symptoms (>3 months)	Not specified
Gatz et al. (2020)	2020	42 (M/F = 20/22)	Insertional Midportion	Pain with tendon loading Duration of symptoms (>2 months) Tendon stiffness	Pain on palpation
Solomons et al. (2020)	2020	52 (M/F = 24/28)	Midportion	Location of pain Duration of symptoms (>3 months) Pain with tendon loading	Double leg heel raise Single leg heel raise Jump Hopping
Van der Vlist et al. (2020)	2020	91 (M/F = 45/46)	Midportion	Location of pain (2–7 cm above calcaneal insertion) Duration of symptoms (>2 months) Pain with tendon loading	Pain on palpation Localised swelling on palpation

**Note:**

cm, centimetres; M, male; F, female.

*Rasmussen et al., 2008; Silbernagel et al., 2001; Solomons et al., 2020* and six months (*Mayer et al., 2007; Rompe, Furia & Maffulli, 2009; Rompe et al., 2008; Rompe et al., 2007; Stergioulas et al., 2008*). Palpation was the most commonly used objective test, with 15 of the included studies (*Bjordal, Lopes-Martins & Iversen, 2006; Boesen et al., 2017; Brown et al., 2006; Gatz et al., 2020; Krogh et al., 2016; Mafi, Lorentzon & Alfredson, 2001; Mayer et al., 2007; Paoloni et al., 2004; Petersen, Welp & Rosenbaum, 2007; Rasmussen et al., 2008; Rompe et al., 2008; Roos et al., 2004; Silbernagel et al., 2001; Stergioulas et al., 2008; Van der Vlist et al., 2020*) using palpation to assess pain, localised tendon thickening or localised swelling. Four studies (*Ebbesen et al., 2018; Knobloch et al., 2007; Rompe, Furia & Maffulli, 2009; Rompe et al., 2007*) used solely subjective history to diagnose Achilles tendinopathy.

**Cohort studies**

Of the included cohort studies, 21 were prospective cohort studies (*Cheng, Zhang & Cai, 2016; Chester et al., 2008; Crill, Berlet & Hyer, 2014; Duthon et al., 2011; Jowett, Richmond & Bedi, 2018; Karjalainen et al., 2000; Khan et al., 2003; Knobloch, 2007; Knobloch et al., 2008; Maffulli et al., 2008; Mansur et al., 2019; McCormack et al., 2015; O'Neill et al., 2019; Oloff et al., 2015; Paavola et al., 2002; Paavola et al., 2000; Sayana & Maffulli, 2007; Silbernagel et al., 2007; Syvertson et al., 2017; Zhuang et al., 2019*) and 10 were retrospective cohort studies (*Alfredson & Spang, 2018; Barge-Caballero et al., 2008; Florit et al., 2019; Murawski et al., 2014; Stenson et al., 2018; Von Wehren et al., 2019*;

*Wei et al., 2017; Welsh & Clodman, 1980; Zellers et al., 2019; Zhang et al., 2020).* Midportion Achilles tendinopathy was investigated in 15 studies (*Chester et al., 2008; Crill, Berlet & Hyer, 2014; Duthon et al., 2011; Jowett, Richmond & Bedi, 2018; Knobloch et al., 2008; Lakshmanan & O'Doherty, 2004; Maffulli et al., 2008; Murawski et al., 2014; O'Neill et al., 2019; Paavola et al., 2002; Paavola et al., 2000; Sayana & Maffulli, 2007; Silbernagel et al., 2007; Syvertson et al., 2017; Von Wehren et al., 2019*), insertional tendinopathy was investigated in eight studies (*Cheng, Zhang & Cai, 2016; Mansur et al., 2019; McCormack et al., 2015; Stenson et al., 2018; Wei et al., 2017; Zellers et al., 2019; Zhang et al., 2020; Zhuang et al., 2019*), both insertional and midportion tendinopathy was investigated in five studies (*Alfredson & Spang, 2018; Karjalainen et al., 2000; Khan et al., 2003; Knobloch, 2007; Welsh & Clodman, 1980*), and three studies did not specify tendinopathy location (Table 7) (*Barge-Caballero et al., 2008; Florit et al., 2019; Oloff et al., 2015*).

Location of pain was the most prominent diagnostic feature, with 26 studies (*Cheng, Zhang & Cai, 2016; Chester et al., 2008; Crill, Berlet & Hyer, 2014; Duthon et al., 2011; Florit et al., 2019; Jowett, Richmond & Bedi, 2018; Karjalainen et al., 2000; Knobloch, 2007; Knobloch et al., 2008; Lakshmanan & O'Doherty, 2004; Maffulli et al., 2008; Mansur et al., 2019; McCormack et al., 2015; Murawski et al., 2014; O'Neill et al., 2019; Oloff et al., 2015; Sayana & Maffulli, 2007; Silbernagel et al., 2007; Stenson et al., 2018; Syvertson et al., 2017; Von Wehren et al., 2019; Wei et al., 2017; Welsh & Clodman, 1980; Zellers et al., 2019; Zhang et al., 2020; Zhuang et al., 2019*) using it as a criteria to diagnose both midportion and insertional Achilles tendinopathy. Midportion tendinopathy was defined as an area 2–4 cm above the calcaneal insertion in two studies (*Crill, Berlet & Hyer, 2014; Welsh & Clodman, 1980*), 2–6 cm above the calcaneal insertion in six studies (*Chester et al., 2008; Knobloch, 2007; Knobloch et al., 2008; Maffulli et al., 2008; Sayana & Maffulli, 2007; Syvertson et al., 2017*), and 2–7 cm above the calcaneal insertion in one study (*Murawski et al., 2014*). Insertional tendinopathy was defined as the distal 2 cm in three studies (*Cheng, Zhang & Cai, 2016; Mansur et al., 2017; McCormack et al., 2015*), and the Achilles ‘insertion’ in five studies (*Knobloch, 2007; Stenson et al., 2018; Zellers et al., 2019; Zhang et al., 2020; Zhuang et al., 2019*). Pain with tendon loading was utilised as a diagnostic criteria in 18 studies (*Alfredson & Spang, 2018; Barge-Caballero et al., 2008; Cheng, Zhang & Cai, 2016; Florit et al., 2019; Khan et al., 2003; Knobloch, 2007; Maffulli et al., 2008; McCormack et al., 2015; Murawski et al., 2014; O'Neill et al., 2019; Paavola et al., 2002; Paavola et al., 2000; Sayana & Maffulli, 2007; Silbernagel et al., 2007; Syvertson et al., 2017; Von Wehren et al., 2019; Wei et al., 2017; Welsh & Clodman, 1980*), and duration of symptoms was utilised in 16 studies (*Cheng, Zhang & Cai, 2016; Chester et al., 2008; Duthon et al., 2011; Jowett, Richmond & Bedi, 2018; Karjalainen et al., 2000; Knobloch et al., 2008; Lakshmanan & O'Doherty, 2004; McCormack et al., 2015; O'Neill et al., 2019; Oloff et al., 2015; Paavola et al., 2000; Silbernagel et al., 2007; Stenson et al., 2018; Von Wehren et al., 2019; Wei et al., 2017; Zhang et al., 2020*). Duration of symptoms varied significantly with studies defining tendinopathy as symptoms lasting less than 6 months (*Paavola et al., 2000*), more than 6 weeks (*McCormack et al., 2015; Von Wehren et al., 2019*), more than 2 months (*Silbernagel et al., 2007*), more than

**Table 7** Cohort studies.

Author	Year	Study design	Sample size	Location	Subjective history	Clinical tests
Welsh & Clodman (1980)	1980	Retrospective	50 (M/F = 28/22)	Insertional Midportion	Location of pain (2–4 cm above calcaneal insertion) Pain with tendon loading	Pain on palpation
Karjalainen et al. (2000)	2000	Prospective	100 (M/F = 75/25)	Insertional Midportion	Location of pain Duration of symptoms (not specified)	Pain on palpation Localised tendon thickening on palpation
Paavola et al. (2000)	2000	Prospective	107 (M/F = 78/29)	Midportion	Pain with tendon loading Duration of symptoms (<6 months)	Pain on palpation Range of motion Single-leg heel raise Single-leg stance
Paavola et al. (2002)	2002	Prospective	42 (M/F = 29/13)	Midportion	Pain with tendon loading	Pain on palpation Localised tendon thickening on palpation Localised swelling on palpation Range of Motion Single-leg heel raise
Khan et al. (2003)	2003	Prospective	45 (M/F = 27/18)	Insertional Midportion	Pain with tendon loading Tendon stiffness VISA-A	Pain on palpation
Lakshmanan & O'Doherty (2004)	2004	Prospective	15 (M/F = 12/3)	Midportion	Location of pain Duration of symptoms (>6 months)	Not specified
Knobloch (2007)	2007	Prospective	64 (M/F = 39/25)	Insertional Midportion	Location of pain (2–6 cm above calcaneal insertion)-midportion Location of pain (insertion) Pain with tendon loading Swelling	Not specified
Sayana & Maffulli (2007)	2007	Prospective	34 (M/F = 18/16)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Pain with tendon loading	Pain on palpation Painful Arc Sign Royal London Hospital Test
Silbernagel et al. (2007)	2007	Prospective	37 (M/F = 20/17)	Midportion	Location of pain Duration of symptoms (>2 months) Pain with tendon loading Swelling	Counter movement Jump Hopping Heel raise
Barge-Caballero et al. (2008)	2008	Retrospective	242 (M/F = 191/51)	Not specified	Pain with tendon loading	Pain on palpation
Chester et al. (2008)	2008	Prospective	16 (M/F = 11/5)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Duration of symptoms (>3 months)	Pain on palpation Localised swelling on palpation
Knobloch et al. (2008)	2008	Prospective	121 (M/F = 74/47)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Duration of symptoms (>3 months)	Pain on palpation Localised swelling on palpation
Maffulli et al. (2008)	2008	Prospective	45 (M/F = 29/16)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Pain with tendon loading	Pain on palpation Painful Arc Sign Royal London Hospital Test
Duthon et al. (2011)	2011	Prospective	14 (M/F = 11/3)	Midportion	Location of pain Duration of symptoms (>1 year)	Localised tendon thickening on palpation Range of Motion Plantarflexion strength Silfverskiold test
Crill, Berlet & Hyer (2014)	2014	Prospective	25 (M/F = not specified)	Midportion	Location of pain (2–4 cm above calcaneal insertion)	Pain on palpation Localised tendon thickening on palpation Single-leg heel raise

(Continued)

**Table 7** (continued)

Author	Year	Study design	Sample size	Location	Subjective history	Clinical tests
Murawski <i>et al.</i> (2014)	2014	Retrospective	32 (M/F = 21/11)	Midportion	Location of pain (2–7 cm above calcaneal insertion) Pain with tendon loading Tendon Stiffness	Pain on palpation Localised tendon thickening on palpation
McCormack <i>et al.</i> (2015)	2015	Prospective	15 (M/F = 4/11)	Insertional	Location of pain (distal 2 cm) Duration of symptoms (>6 weeks) Pain with tendon loading VISA-A	Pain on palpation
Oloff <i>et al.</i> (2015)	2015	Prospective	26 (M/F = not specified)	Not specified	Location of pain Duration of symptoms (>6 months)	Not specified
Cheng, Zhang & Cai (2016)	2016	Prospective	42 (M/F = 29/13)	Insertional	Location of pain (distal 2 cm) Duration of symptoms (>6 months) Swelling Pain with tendon loading	Not specified
Syvertson <i>et al.</i> (2017)	2017	Prospective	11 (M/F = 4/7)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Pain with tendon loading Tendon stiffness	Pain on palpation
Wei <i>et al.</i> (2017)	2017	Retrospective	68 (M/F = 53/15)	Insertional	Location of pain Duration of symptoms (>6 months) Pain with tendon loading	Pain on palpation
Alfredson & Spang (2018)	2018	Retrospective	771 (M/F = 481/290)	Insertional Midportion	Pain with tendon loading VISA-A	Not specified
Jowett, Richmond & Bedi (2018)	2018	Prospective	26 (M/F = 13/13)	Midportion	Location of pain Localised swelling Duration of symptoms (>6 months)	Pain on palpation Localised tendon thickening on palpation
Stenson <i>et al.</i> (2018)	2018	Retrospective	664 (M/F = 312/352)	Insertional	Location of pain (insertion) Duration of symptoms (>3 months)	Range of motion
Florit <i>et al.</i> (2019)	2019	Retrospective	110 (M/F = 103/7)	Not specified	Location of pain Pain with tendon loading	Pain on palpation Pain with tendon loading tests (not specified)
Mansur <i>et al.</i> (2019)	2019	Prospective	19 (M/F = 11/8)	Insertional	Location of pain (distal 2 cm)	Pain on palpation
O'Neill <i>et al.</i> (2019)	2019	Prospective	16 (M/F = 11/5)	Midportion	Location of pain Duration of symptoms (>3 months) Pain with tendon loading	Pain on palpation Painful Arc Sign Royal London Hospital Test
Von Wehren <i>et al.</i> (2019)	2019	Retrospective	50 (M/F = 27/23)	Midportion	Location of pain Duration of symptoms (>6 weeks) Pain with tendon loading	Pain on palpation Localised swelling on palpation
Zellers <i>et al.</i> (2019)	2019	Retrospective	56 (M/F = 25/31)	Insertional	Location of pain (insertion)	Pain on palpation
Zhuang <i>et al.</i> (2019)	2019	Prospective	28 (M/F = 17/11)	Insertional	Location of pain (insertion)	Pain on palpation Localised tendon thickening on palpation Pain with resisted plantarflexion Reduced plantarflexion strength Heel raise test
Zhang <i>et al.</i> (2020)	2020	Retrospective	33 (M/F = 31/2)	Insertional	Location of pain (insertion) Duration of symptoms (>3 months)	Not specified

**Note:**

cm, centimetres; M, male; F, female; VISA-A, Victorian Institute of Sport Assessment-Achilles.

three months (*Chester et al., 2008; Knobloch et al., 2008; O'Neill et al., 2019; Stenson et al., 2018; Zhang et al., 2020*), more than six months (*Cheng, Zhang & Cai, 2016; Jowett, Richmond & Bedi, 2018; Lakshmanan & O'Doherty, 2004; Oloff et al., 2015; Wei et al., 2017*), and more than 1 year (*Duthon et al., 2011*). As with the previous studies, the most common objective test for diagnosing Achilles tendinopathy was palpation, with 23 studies utilising it as a diagnostic criteria (*Barge-Caballero et al., 2008; Chester et al., 2008; Crill, Berlet & Hyer, 2014; Duthon et al., 2011; Florit et al., 2019; Jowett, Richmond & Bedi, 2018; Karjalainen et al., 2000; Khan et al., 2003; Knobloch et al., 2008; Maffulli et al., 2008; Mansur et al., 2019; McCormack et al., 2015; Murawski et al., 2014; O'Neill et al., 2019; Paavola et al., 2002; Paavola et al., 2000; Sayana & Maffulli, 2007; Syvertson et al., 2017; Von Wehren et al., 2019; Wei et al., 2017; Welsh & Clodman, 1980; Zellers et al., 2019; Zhuang et al., 2019*). Six studies used only subjective measures for diagnosing Achilles tendinopathy (*Alfredson & Spang, 2018; Cheng, Zhang & Cai, 2016; Knobloch, 2007; Lakshmanan & O'Doherty, 2004; Oloff et al., 2015; Zhang et al., 2020*).

### Case-control studies

Of the 30 case-control studies, one (*Chimenti et al., 2016*) investigated insertional Achilles tendinopathy, 15 studies (*Hutchison et al., 2013; Abate & Salini, 2019; Azevedo et al., 2009; Creaby et al., 2017; Gårdin et al., 2016; Lohrer & Nauck, 2009; Maffulli et al., 2003; Nadeau et al., 2016; Neeter et al., 2003; Pingel et al., 2013; Reid et al., 2012; Romero-Morales et al., 2019a; Ryan et al., 2009; Sengkerij et al., 2009; Romero-Morales et al., 2019b*) investigated midportion Achilles tendinopathy, nine studies (*Chimenti et al., 2020; Coombes et al., 2018; Eckenrode, Kietrys & Stackhouse, 2019; Hernández-Sánchez et al., 2018; Ooi et al., 2015; Rabello et al., 2020; Reiter et al., 2004; Verrall, Schofield & Brustad, 2011; Zhang et al., 2017*) investigated both insertional and midportion Achilles tendinopathy, with five studies (*Cassel et al., 2018; Holmes & Lin, 2006; Jewson et al., 2017; Leung & Griffith, 2008; De Mesquita et al., 2018*) not specifying tendinopathy location (Table 8). As with the previous study types, the most commonly used diagnostic feature was location of pain, which was utilised in 27 of the case-control studies (*Hutchison et al., 2013; Abate & Salini, 2019; Chimenti et al., 2016; Chimenti et al., 2020; Coombes et al., 2018; Creaby et al., 2017; Eckenrode, Kietrys & Stackhouse, 2019; Gårdin et al., 2016; Hernández-Sánchez et al., 2018; Holmes & Lin, 2006; Jewson et al., 2017; Leung & Griffith, 2008; Lohrer & Nauck, 2009; Nadeau et al., 2016; Neeter et al., 2003; De Mesquita et al., 2018; Ooi et al., 2015; Pingel et al., 2013; Rabello et al., 2020; Reid et al., 2012; Reiter et al., 2004; Romero-Morales et al., 2019a; Ryan et al., 2009; Sengkerij et al., 2009; Verrall, Schofield & Brustad, 2011; Zhang et al., 2017; Romero-Morales et al., 2019b*). Insertional tendinopathy was defined as the distal 2 cm of the Achilles tendon in three studies (*Chimenti et al., 2020; Rabello et al., 2020; Verrall, Schofield & Brustad, 2011*). Midportion tendinopathy was defined as 2–6 cm above the calcaneal insertion in eight studies (*Hutchison et al., 2013; Chimenti et al., 2020; Neeter et al., 2003; Rabello et al., 2020; Reid et al., 2012; Ryan et al., 2009; Verrall, Schofield & Brustad, 2011; Zhang et al., 2017*), 2–7 cm above the calcaneal insertion in three studies (*Gårdin et al., 2016; Lohrer & Nauck, 2009; Sengkerij et al., 2009*), and the middle third of the tendon in one study

**Table 8** Case-control studies.

Author	Year	Sample size	Location	Subjective history	Clinical tests
<i>Maffulli et al. (2003)</i>	2003	24 (M/F = 24/0)	Midportion	Not specified	Pain on palpation Painful Arc Sign Royal London Hospital Test
<i>Neeter et al. (2003)</i>	2003	25 (M/F = 15/10)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Duration of symptoms (<3 months)	Pain on palpation Range of motion Single-leg heel raise
<i>Reiter et al. (2004)</i>	2004	35 (M/F = 30/5)	Insertional Midportion	Location of pain Pain with tendon loading VISA-A	Pain on palpation Localised tendon thickening on palpation
<i>Holmes &amp; Lin (2006)</i>	2006	82 (M/F = 44/38)	Not specified	Location of pain Pain with tendon loading	Pain on palpation
<i>Leung &amp; Griffith (2008)</i>	2008	71 (M/F = 31/40)	Not specified	Location of pain Duration of symptoms (>6 months)	Pain on palpation Localised thickening on palpation Localised swelling on palpation
<i>Azevedo et al. (2009)</i>	2009	42 (M/F = 32/10)	Midportion	Gradual onset of pain Tendon stiffness Swelling Pain with tendon loading	Pain on palpation Localised tendon thickening on palpation Painful Arc Sign
<i>Lohrer &amp; Nauck (2009)</i>	2009	119 (M/F = not specified)	Midportion	Location of pain (2–7 cm above calcaneal insertion) Pain with tendon loading	Pain on palpation
<i>Ryan et al. (2009)</i>	2009	48 (M/F = 48/0)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Duration of symptoms (>3 months) Pain with tendon loading	Not specified
<i>Sengkerij et al. (2009)</i>	2009	25 (M/F = 16/9)	Midportion	Location of pain (2–7 cm above calcaneal insertion) Pain with tendon loading	Pain on palpation
<i>Verral, Schofield &amp; Brustad (2011)</i>	2011	190 (M/F = 108/82)	Insertional Midportion	Location of pain (2–6 cm above calcaneal insertion) Location of pain (distal 2 cm) Duration of symptoms (>3 months) Pain with tendon loading Tendon stiffness	Pain on palpation Localised swelling on palpation
<i>Reid et al. (2012)</i>	2012	36 (M/F = not specified)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Pain with tendon loading	Pain on palpation
<i>Hutchison et al. (2013)</i>	2013	21 (M/F = 9/12)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Tendon stiffness	Pain on palpation Localised tendon thickening on palpation Painful Arc Sign Royal London Hospital Test Pain with dorsiflexion Single-leg heel raise Hopping
<i>Pingel et al. (2013)</i>	2013	18 (M/F = 10/8)	Midportion	Location of pain Duration of symptoms (>6 months)	Pain on palpation Localised swelling on palpation
<i>Ooi et al. (2015)</i>	2015	240 (M/F = 180/60)	Insertional Midportion	Location of pain Duration of symptoms (>3 months)	Pain on palpation Localised swelling on palpation
<i>Chimenti et al. (2016)</i>	2016	40 (M/F = 20/20)	Insertional	Location of pain (insertion) Pain with tendon loading	Pain on palpation

**Table 8** (continued)

Author	Year	Sample size	Location	Subjective history	Clinical tests
Gärdin et al. (2016)	2016	30 (M/F = 12/18)	Midportion	Location of pain (2–7 cm above calcaneal insertion) Duration of symptoms (>6 months)	Pain on palpation
Nadeau et al. (2016)	2016	43 (M/F = 30/13)	Midportion	Location of pain (middle third) Duration of symptoms (>4 weeks) VISA-A	Pain on palpation Localised tendon thickening on palpation Pain with passive dorsiflexion Pain with resisted plantarflexion Single-leg heel raise Hopping
Creaby et al. (2017)	2017	25 (M/F = 25/0)	Midportion	Location of pain Pain with tendon loading Tendon stiffness	Pain on palpation Hopping
Jewson et al. (2017)	2017	35 (M/F = 22/13)	Not specified	Location of pain Duration of symptoms (>4 weeks) Pain with tendon loading	Not specified
Zhang et al. (2017)	2017	37 (M/F = 26/11)	Insertional Midportion	Location of pain (2–6 cm above calcaneal insertion) Location of pain (insertion)	Pain on palpation Pain with resisted plantarflexion
Cassel et al. (2018)	2018	182 (M/F = 113/69)	Not specified	Pain with tendon loading	Pain on palpation
Coombes et al. (2018)	2018	67 (M/F = 37/30)	Insertional Midportion	Location of pain Duration of symptoms (>3 months)	Pain on palpation Single-leg heel raise
Hernández-Sánchez et al. (2018)	2018	210 (M/F = 148/62)	Insertional Midportion	Location of pain Pain with tendon loading Tendon stiffness	Not specified
De Mesquita et al. (2018)	2018	67 (M/F = 41/26)	Not specified	Location of pain Duration of symptoms (>2 months)	Pain on palpation
Abate & Salini (2019)	2019	64 (M/F = 40/24)	Midportion	Location of pain Pain with tendon loading	Pain on palpation
Eckenrode, Kietrys & Stackhouse (2019)	2019	41 (M/F = 19/22)	Insertional Midportion	Location of pain Duration of symptoms (>3 months) Pain with tendon loading	Pain on palpation Single-leg heel raise
Romero-Morales et al. (2019a)	2019	141 (M/F = 116/25)	Midportion	Location of pain Duration of symptoms (>3 months) Pain with tendon loading	Pain on palpation
Romero-Morales et al. (2019b)	2019	143 (M/F = not specified)	Midportion	Location of pain Duration of symptoms (>3 months) Pain with tendon loading	Pain on palpation
Chimenti et al. (2020)	2020	46 (M/F = 30/16)	Insertional Midportion	Location of pain (2–6 cm above calcaneal insertion) Location of pain (distal 2 cm) Duration of symptoms (>3 months) Pain with tendon loading Tendon stiffness	Pain on palpation
Rabello et al. (2020)	2020	46 (M/F = 30/16)	Insertional Midportion	Location of pain (2–6 cm above calcaneal insertion) Location of pain (distal 2 cm)	Not specified

**Note:**

cm, centimetres; M, male; F, female; VISA-A, Victorian Institute of Sport Assessment-Achilles.

([Nadeau et al., 2016](#)). Additionally, duration of symptoms was commonly used to diagnose Achilles tendinopathy, with variations in the criteria. Achilles tendinopathy was defined as duration of symptoms of less than three months in one study ([Neeter et al., 2003](#)), greater than four weeks in two studies ([Jewson et al., 2017; Nadeau et al., 2016](#)), greater than two months in one study ([De Mesquita et al., 2018](#)), greater than three months in eight studies ([Chimenti et al., 2020; Coombes et al., 2018; Eckenrode, Kietrys & Stackhouse, 2019; Ooi et al., 2015; Romero-Morales et al., 2019a; Ryan et al., 2009; Verrall, Schofield & Brustad, 2011; Romero-Morales et al., 2019b](#)), and greater than six months in three studies ([Gärdin et al., 2016; Leung & Griffith, 2008; Pingel et al., 2013](#)). Pain with tendon loading was included as a diagnostic criteria in 18 studies ([Abate & Salini, 2019; Azevedo et al., 2009; Cassel et al., 2018; Chimenti et al., 2016; Chimenti et al., 2020; Creaby et al., 2017; Eckenrode, Kietrys & Stackhouse, 2019; Hernández-Sánchez et al., 2018; Holmes & Lin, 2006; Jewson et al., 2017; Lohrer & Nauck, 2009; Reid et al., 2012; Reiter et al., 2004; Romero-Morales et al., 2019a; Ryan et al., 2009; Sengkerij et al., 2009; Verrall, Schofield & Brustad, 2011; Romero-Morales et al., 2019b](#)). One study did not specify a subjective criteria to diagnose Achilles tendinopathy ([Maffulli et al., 2003](#)). Similar to previous study designs, palpation was the most common clinical test to diagnose Achilles tendinopathy, with it being used in 26 studies ([Hutchison et al., 2013; Abate & Salini, 2019; Azevedo et al., 2009; Cassel et al., 2018; Chimenti et al., 2016; Chimenti et al., 2020; Coombes et al., 2018; Creaby et al., 2017; Eckenrode, Kietrys & Stackhouse, 2019; Gärdin et al., 2016; Holmes & Lin, 2006; Leung & Griffith, 2008; Lohrer & Nauck, 2009; Maffulli et al., 2003; Nadeau et al., 2016; Neeter et al., 2003; De Mesquita et al., 2018; Ooi et al., 2015; Pingel et al., 2013; Reid et al., 2012; Reiter et al., 2004; Romero-Morales et al., 2019a; Sengkerij et al., 2009; Verrall, Schofield & Brustad, 2011; Zhang et al., 2017; Romero-Morales et al., 2019b](#)). Four studies relied only on subjective measures to diagnose Achilles tendinopathy ([Hernández-Sánchez et al., 2018; Jewson et al., 2017; Rabello et al., 2020; Ryan et al., 2009](#)).

### Cross-sectional studies

[Table 9](#) provides an overview of the 17 included cross-sectional studies, with 10 studies ([De Jonge et al., 2011; De Marchi et al., 2018; Divani et al., 2010; Finnamore et al., 2019; Maffulli et al., 2008; Praet et al., 2018; Santamato et al., 2019; Scholes et al., 2018; Van der Vlist et al., 2020; Wang et al., 2012](#)) investigating midportion Achilles tendinopathy, four studies ([Docking et al., 2015; Kragsnaes et al., 2014; Turner et al., 2020; Vallance et al., 2020](#)) investigating both insertional and midportion Achilles tendinopathy, and three studies ([Aiyegbusi, Tella & Sanusi, 2020; Longo et al., 2009; Mantovani et al., 2020](#)) not specifying tendinopathy location. Once again, location of pain was the most common subjective measure to diagnose Achilles tendinopathy, with 12 studies utilising it as a diagnostic criteria ([De Jonge et al., 2011; De Marchi et al., 2018; Kragsnaes et al., 2014; Maffulli et al., 2008; Mantovani et al., 2020; Praet et al., 2018; Chester et al., 2008; Scholes et al., 2018; Turner et al., 2020; Vallance et al., 2020; Van der Vlist et al., 2020; Wang et al., 2012](#)). Midportion Achilles tendinopathy was defined as 2–6 cm above the calcaneal insertion in four studies ([De Marchi et al., 2018; Maffulli et al., 2008; Praet et al.,](#)

**Table 9** Cross-sectional studies.

Author	Year	Sample size	Location	Subjective history	Clinical tests
<i>Maffulli et al. (2008)</i>	2008	50 (M/F = 50/0)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Duration of symptoms (>3 months) Pain with tendon loading	Pain on palpation Localised swelling on palpation
<i>Longo et al. (2009)</i>	2009	178 (M/F = 110/68)	Not specified	VISA-A	Pain on palpation Localised tendon thickening on palpation Painful Arc Sign Royal London Hospital Test
<i>Divani et al. (2010)</i>	2010	26 (M/F = 17/9)	Midportion	Not specified	Pain on palpation
<i>De Jonge et al. (2011)</i>	2011	107 (M/F = 51/56)	Midportion	Location of pain (above calcaneal insertion)	Not specified
<i>Wang et al. (2012)</i>	2012	17 (M/F = 17/0)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Duration of symptoms (>3 months) Pain with tendon loading VISA-A	Pain on palpation Royal London Hospital Test
<i>Kragsnaes et al. (2014)</i>	2014	50 (M/F = 27/23)	Insertional Midportion	Location of pain Duration of symptoms (>4 months)	Pain on palpation
<i>Docking et al. (2015)</i>	2015	21 (M/F = 20/1)	Insertional Midportion	Pain with tendon loading	Single-leg heel raise Hopping
<i>De Marchi et al. (2018)</i>	2018	27 (M/F = 19/8)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Duration of symptoms (>6 months) Pain with tendon loading	Not specified
<i>Praet et al. (2018)</i>	2018	20 (M/F = 13/7)	Midportion	Location of pain (2–6 cm above calcaneal insertion) Duration of symptoms (>2 months)	Pain on palpation
<i>Scholes et al. (2018)</i>	2018	21 (M/F = 21/0)	Midportion	Location of pain (midportion) Duration of symptoms (>3 months) Pain with tendon loading Tendon stiffness	Not specified
<i>Finnimore et al. (2019)</i>	2019	25 (M/F = 12/13)	Midportion	Duration of symptoms (>3 months) Pain with tendon loading	Pain on palpation
<i>Santamato et al. (2019)</i>	2019	12 (M/F = 7/5)	Midportion	Location of pain Duration of symptoms (>4 weeks)	Reduced ROM Pain during AROM Painful Arc Sign Royal London Hospital Test
<i>Aiyegbusi, Tella &amp; Sanusi (2020)</i>	2020	85 (M/F = 56/29)	Not specified	VISA-A	Royal London Hospital Test
<i>Mantovani et al. (2020)</i>	2020	19 (M/F = 13/6)	Not specified	Location of pain Duration of symptoms (>3 months) VISA-A (<80)	Pain on palpation
<i>Turner et al. (2020)</i>	2020	15 (M/F = 8/7)	Insertional Midportion	Location of pain Duration of symptoms (>3 months) Gradual onset of pain Pain with tendon loading	Not specified
<i>Vallance et al. (2020)</i>	2020	86 (M/F = 86/0)	Insertional Midportion	Location of pain Duration of symptoms (>3 months) Pain with tendon loading Tendon stiffness	Single-leg heel raise Hopping
<i>Van der Vlist et al. (2020)</i>	2020	28 (M/F = 16/12)	Midportion	Location of pain (2–7 cm above calcaneal insertion) Duration of symptoms (>2 months) Pain with tendon loading	Pain on palpation Localised swelling on palpation

**Note:**

cm, centimetres; M, male; F, female; VISA-A, Victorian Institute of Sport Assessment-Achilles.

2018; Wang et al., 2012), and 2–7 cm above the calcaneal insertion in one study (Van der Vlist et al., 2020). Similarly, 12 studies included duration of symptoms as a diagnostic criteria, with durations of symptoms including greater than four weeks (Santamato et al., 2019), greater than two months (Praet et al., 2018; Van der Vlist et al., 2020), greater than three months (Finnimore et al., 2019; Maffulli et al., 2008; Mantovani et al., 2020; Scholes et al., 2018; Turner et al., 2020; Vallance et al., 2020; Wang et al., 2012), greater than four months (Kratsnaes et al., 2014), and greater than six months (De Marchi et al., 2018). Pain with tendon loading was the next most common subjective diagnostic measure, with nine studies including it as a diagnostic measure (De Marchi et al., 2018; Docking et al., 2015; Finnimore et al., 2019; Maffulli et al., 2008; Scholes et al., 2018; Turner et al., 2020; Vallance et al., 2020; Van der Vlist et al., 2020; Wang et al., 2012). One study (Divani et al., 2010) did not report subjective measures to confirm the diagnosis of Achilles tendinopathy. The most common clinical test included palpation, with nine studies using palpation to clinically diagnose Achilles tendinopathy (Divani et al., 2010; Finnimore et al., 2019; Kratsnaes et al., 2014; Longo et al., 2009; Maffulli et al., 2008; Mantovani et al., 2020; Praet et al., 2018; Van der Vlist et al., 2020; Wang et al., 2012). Four studies (Aiyegbusi, Tella & Sanusi, 2020; Longo et al., 2009; Santamato et al., 2019; Wang et al., 2012) included the Royal London Hospital Test as a clinical measure of Achilles tendinopathy, with four studies (De Jonge et al., 2011; De Marchi et al., 2018; Scholes et al., 2018; Turner et al., 2020) not specifying the clinical tests utilised to confirm the diagnosis.

### Narrative reviews

Of the 43 narrative reviews included in the scoping review, seven studies investigated insertional Achilles tendinopathy (Aldridge, 2004; Benazzo, Todesca & Ceciliani, 1997; Chimenti et al., 2017; Den Hartog, 2009; Hu & Flemister, 2008; Irwin, 2010; Maffulli et al., 2019), 18 studies investigated midportion Achilles tendinopathy (Alfredson, 2003; Alfredson & Cook, 2007; Bains & Porter, 2006; Courville, Coe & Hecht, 2009; Feilmeier, 2017; Järvinen et al., 2001; Kader et al., 2002; Longo, Ronga & Maffulli, 2009; Maffulli & Kader, 2002; Maffulli et al., 2012; Maffulli, Sharma & Luscombe, 2004; Maffulli, Via & Oliva, 2014; McShane, Ostick & McCabe, 2007; Paavola et al., 2002; Scott, Huisman & Khan, 2011; Sharma & Maffulli, 2006; Simpson & Howard, 2009; Tan & Chan, 2008), 13 studies investigated both insertional and midportion Achilles tendinopathy (Aronow, 2005; Asplund & Best, 2013; Bhatty, Khan & Zubairy, 2019; Chazan, 1998; Cook, Khan & Purdam, 2002; Fredericson, 1996; Furia & Rompe, 2007; Horn & McCollum, 2015; Jukes, Scott & Solan, 2020; Maffulli, Giai Via & Oliva, 2015; Maffulli et al., 2020; Pedowitz & Beck, 2017; Saini et al., 2015), and five studies did not specify tendinopathy location (Table 10) (Millar et al., 2021; Baskerville et al., 2018; Maffulli, Giuseppe Longo & Denaro, 2012; Nichols, 1989; Sorosky et al., 2004). The most common subjective diagnostic criteria for diagnosing Achilles tendinopathy was pain with tendon loading, with all 43 included reviews utilising it as a diagnostic criteria (Millar et al., 2021; Aldridge, 2004; Alfredson, 2003; Alfredson & Cook, 2007; Aronow, 2005; Asplund & Best, 2013; Bains & Porter, 2006; Baskerville et al., 2018; Benazzo, Todesca & Ceciliani, 1997; Bhatty, Khan & Zubairy, 2019; Chazan, 1998; Chimenti et al., 2017; Cook, Khan &

**Table 10** Narrative reviews.

Author	Year	Location	Subjective history	Clinical tests
<i>Nichols (1989)</i>	1989	Not specified	Location of pain (Distal 5 cm) Tendon stiffness Pain with tendon loading Gradual onset of pain Change in activity	Pain on palpation Localised tendon thickening on palpation Localised swelling on palpation Reduced ROM Hopping
<i>Fredericson (1996)</i>	1996	Insertional Midportion	Location of pain (2–6 cm above calcaneal insertion) Location of pain (insertion) Pain with tendon loading Tendon stiffness	Pain on palpation Localised tendon thickening on palpation Localised swelling on palpation
<i>Benazzo, Todesca &amp; Ceciliani (1997)</i>	1997	Insertional	Pain with tendon loading Tendon stiffness	Pain on palpation
<i>Chazan (1998)</i>	1998	Insertional Midportion	Gradual onset of pain Tendon stiffness Pain with tendon loading	Pain on palpation Localised tendon thickening on palpation Localised swelling on palpation Pain with passive dorsiflexion Pain with resisted plantarflexion Reduced ROM
<i>Järvinen et al. (2001)</i>	2001	Midportion	Location of pain Pain with tendon loading	Pain on palpation Localised swelling on palpation
<i>Cook, Khan &amp; Purdam (2002)</i>	2002	Insertional Midportion	Gradual onset of pain Location of pain Pain with tendon loading Tendon stiffness Change in activity VISA-A	Pain on palpation Localised tendon thickening on palpation Localised swelling on palpation Single-leg heel raise Hopping
<i>Kader et al. (2002)</i>	2002	Midportion	Gradual onset of pain Location of pain (2–6 cm above calcaneal insertion) Duration of symptoms Tendon stiffness Pain with tendon loading	Pain on palpation Localised tendon thickening on palpation Painful Arc Sign
<i>Maffulli &amp; Kader (2002)</i>	2002	Midportion	Gradual onset of pain Location of pain (2–6 cm above calcaneal insertion) Duration of symptoms Pain with tendon loading	Pain on palpation Localised tendon thickening on palpation Localised swelling on palpation Painful Arc Sign
<i>Paavola et al. (2002)</i>	2002	Midportion	Pain with tendon loading Duration of symptoms	Pain on palpation Localised tendon thickening on palpation Localised swelling on palpation
<i>Alfredson (2003)</i>	2003	Midportion	Location of pain (2–6 cm above calcaneal insertion) Pain with tendon loading Tendon stiffness	Pain on palpation Localised swelling on palpation
<i>Aldridge (2004)</i>	2004	Insertional	Pain with tendon loading	Pain on palpation Pain with passive dorsiflexion
<i>Maffulli, Sharma &amp; Luscombe (2004)</i>	2004	Midportion	Location of pain (2–6 cm above calcaneal insertion) Pain with tendon loading	Pain on palpation Localised tendon thickening on palpation Localised swelling on palpation
<i>Sorosky et al. (2004)</i>	2004	Not specified	Gradual onset of pain Pain with tendon loading Change in training	Pain on palpation Localised tendon thickening on palpation Localised swelling on palpation Pain with passive dorsiflexion Pain with resisted plantarflexion

(Continued)

**Table 10** (continued)

Author	Year	Location	Subjective history	Clinical tests
Aronow (2005)	2005	Insertional Midportion	Pain with tendon loading	Pain on palpation Localised swelling on palpation Pain with passive dorsiflexion (insertional) Painful Arc Sign (midportion) Single-leg Heel Raise
Bains & Porter (2006)	2006	Midportion	Gradual onset of pain Location of pain (2–5 cm above calcaneal insertion) Tendon stiffness Pain with tendon loading Change in training VISA-A	Pain on palpation Localised tendon thickening on palpation Localised swelling on palpation Hopping on the spot Forward hopping 6m hop test
Sharma & Maffulli (2006)	2006	Midportion	Location of pain (2–6 cm above calcaneal insertion) Pain with tendon loading Swelling	Pain on palpation Localised tendon thickening on palpation Localised swelling on palpation
Alfredson & Cook (2007)	2007	Midportion	Pain with tendon loading Tendon stiffness	Localised swelling on palpation Single-leg Heel Raise Hopping on the spot Forward hopping
Furia & Rompe (2007)	2007	Insertional Midportion	Location of pain (2–5 cm above calcaneal insertion) - midportion Location of pain (insertion) Pain with tendon loading	Pain on palpation Localised swelling on palpation Pain with passive dorsiflexion (insertional) Reduced ROM
McShane, Ostick & McCabe (2007)	2007	Midportion	Location of pain Pain with tendon loading Tendon stiffness	Pain on palpation Localised tendon thickening on palpation Single-leg heel raise
Hu & Flemister (2008)	2008	Insertional	Location of pain Pain with tendon loading Tendon stiffness	Pain on palpation Localised tendon thickening on palpation Pain with passive dorsiflexion Reduced ROM Silfverskiold test
Tan & Chan (2008)	2008	Midportion	Location of pain (midportion) Duration of pain (>2 weeks) Pain with tendon loading Tendon stiffness	Pain on palpation Localised tendon thickening on palpation Localised swelling on palpation Single-leg heel raise
Courville, Coe & Hecht (2009)	2009	Midportion	Pain with tendon loading Location of pain Tendon stiffness Change in activity	Pain on palpation Localised tendon thickening on palpation Localised swelling on palpation
Den Hartog (2009)	2009	Insertional	Gradual onset of pain Duration of symptoms Pain with tendon loading	Pain on palpation Localised tendon thickening on palpation Localised swelling on palpation
Longo, Ronga & Maffulli (2009)	2009	Midportion	Location of pain (2–6 cm above calcaneal insertion) Pain with tendon loading Swelling	Pain on palpation Localised tendon thickening on palpation Painful Arc Sign Royal London Hospital Test
Simpson & Howard (2009)	2009	Midportion	Location of pain (2–5 cm above calcaneal insertion) Pain with tendon loading Swelling Change in activity	Pain on palpation Localised swelling on palpation Reduced flexibility in hamstring and calf

**Table 10** (continued)

Author	Year	Location	Subjective history	Clinical tests
<i>Irwin (2010)</i>	2010	Insertional	Location of pain (calcaneal tuberosity) Swelling Pain with tendon loading Tendon stiffness	Pain on palpation Localised swelling on palpation
<i>Scott, Huisman &amp; Khan (2011)</i>	2011	Midportion	Location of pain Pain with tendon loading Swelling	Pain on palpation Localised tendon thickening on palpation
<i>Maffulli et al. (2012)</i>	2012	Midportion	Location of pain (4–6 cm above calcaneal insertion) Pain with tendon loading Swelling Change in activity	Pain on palpation Localised tendon thickening on palpation Localised swelling on palpation Reduced ROM
<i>Maffulli, Giuseppe Longo &amp; Denaro (2012)</i>	2012	Not specified	Gradual onset of pain Tendon stiffness Pain with tendon loading Swelling VISA-A	Pain on palpation Localised tendon thickening on palpation Localised swelling on palpation Painful Arc Sign
<i>Asplund &amp; Best (2013)</i>	2013	Insertional Midportion	Location of pain (2–6 cm above calcaneal insertion) Location of pain (insertion) Pain with tendon loading Tendon stiffness	Pain on palpation Localised swelling on palpation
<i>Maffulli, Via &amp; Oliva (2014)</i>	2014	Midportion	Location of pain (2–6 cm above calcaneal insertion) Pain with tendon loading	Pain on palpation Localised swelling on palpation Single-leg heel raise
<i>Horn &amp; McCollum (2015)</i>	2015	Insertional Midportion	Location of pain (2–6 cm above calcaneal insertion) Location of pain (insertion) Pain with tendon loading	Localised swelling on palpation Painful Arc Sign Royal London Hospital Test
<i>Maffulli, Giai Via &amp; Oliva (2015)</i>	2015	Insertional Midportion	Location of pain (2–6 cm above calcaneal insertion) Location of pain (insertion) Swelling Pain with tendon loading	Pain on palpation
<i>Saini et al. (2015)</i>	2015	Insertional Midportion	Location of pain (midportion) Location of pain (insertion) Pain with tendon loading	Pain on palpation Localised swelling on palpation Pain with dorsiflexion and plantarflexion
<i>Chimenti et al. (2017)</i>	2017	Insertional	Location of pain (distal 2 cm) Pain with tendon loading Tendon stiffness	Pain on palpation Localised swelling on palpation Pain with passive dorsiflexion Pain with resisted plantarflexion Reduced ROM
<i>Feilmeier (2017)</i>	2017	Midportion	Location of pain (2–6 cm above calcaneal insertion) Swelling Pain with tendon loading Tendon stiffness Change in activity VISA-A	Pain on palpation Localised tendon thickening on palpation Pain with passive dorsiflexion Painful Arc Sign Royal London Hospital Test Single-leg Heel Raise Hopping
<i>Pedowitz &amp; Beck (2017)</i>	2017	Insertional Midportion	Location of pain (2–6 cm above calcaneal insertion) Location of pain (insertion) Pain with tendon loading Tendon stiffness	Pain on palpation Single leg heel raise

(Continued)

**Table 10** (continued)

Author	Year	Location	Subjective history	Clinical tests
Baskerville <i>et al.</i> (2018)	2018	Not specified	Gradual onset of pain Pain with tendon loading Tendon stiffness	Pain on palpation Pain with passive and active movement Reduced strength
Bhatti, Khan & Zubairy (2019)	2019	Insertional Midportion	Pain with tendon loading	Pain on palpation Localised swelling on palpation Pain with passive dorsiflexion (insertional) Pain with resisted plantarflexion Reduced ROM
Maffulli <i>et al.</i> (2019)	2019	Insertional	Location of pain (distal 2 cm) Pain with tendon loading Tendon stiffness	Pain on palpation Localised tendon thickening on palpation Localised swelling on palpation
Jukes, Scott & Solan (2020)	2020	Insertional Midportion	Location of pain Pain with tendon loading Tendon stiffness Duration of symptoms	Pain on palpation Localised tendon thickening on palpation Localised swelling on palpation Royal London Hospital Test Silfverskiold test
Maffulli <i>et al.</i> (2020)	2020	Insertional Midportion	Location of pain Pain with tendon loading Localised swelling Tendon stiffness	Pain on palpation Painful Arc Sign Royal London Hospital Test
Millar <i>et al.</i> (2021)	2021	Not specified	Location of pain Pain with tendon loading Tendon stiffness	Pain on palpation Painful Arc Sign Royal London Hospital Test Single leg heel raise Hopping

**Note:**

cm, centimetres; M, male; F, female; VISA-A, Victorian Institute of Sport Assessment-Achilles; ROM, range of motion; m, metres.

Purdam, 2002; Courville, Coe & Hecht, 2009; Den Hartog, 2009; Feilmeier, 2017; Fredericson, 1996; Furia & Rompe, 2007; Horn & McCollum, 2015; Hu & Flemister, 2008; Irwin, 2010; Järvinen *et al.*, 2001; Jukes, Scott & Solan, 2020; Kader *et al.*, 2002; Longo, Ronga & Maffulli, 2009; Maffulli, Giai Via & Oliva, 2015; Maffulli, Giuseppe Longo & Denaro, 2012; Maffulli & Kader, 2002; Maffulli *et al.*, 2020; Maffulli *et al.*, 2012; Maffulli *et al.*, 2019; Maffulli, Sharma & Luscombe, 2004; Maffulli, Via & Oliva, 2014; McShane, Ostick & McCabe, 2007; Nichols, 1989; Paavola *et al.*, 2002; Pedowitz & Beck, 2017; Saini *et al.*, 2015; Scott, Huisman & Khan, 2011; Sharma & Maffulli, 2006; Simpson & Howard, 2009; Sorosky *et al.*, 2004; Tan & Chan, 2008). Location of pain was included as a diagnostic criteria of Achilles tendinopathy in 31 studies, with midportion tendinopathy defined as ‘midportion’ in two studies (Saini *et al.*, 2015; Tan & Chan, 2008), distal 5 cm of the Achilles tendon in one study (Nichols, 1989), 2–5 cm above the calcaneal insertion in three studies (Bains & Porter, 2006; Furia & Rompe, 2007; Simpson & Howard, 2009), 2–6 cm above the calcaneal insertion in 13 studies (Alfredson, 2003; Asplund & Best, 2013; Feilmeier, 2017; Fredericson, 1996; Horn & McCollum, 2015; Kader *et al.*, 2002; Longo, Ronga & Maffulli, 2009; Maffulli, Giai Via & Oliva, 2015; Maffulli & Kader, 2002; Maffulli, Sharma & Luscombe, 2004; Maffulli, Via & Oliva, 2014; Pedowitz & Beck, 2017; Sharma & Maffulli, 2006), and 4–6 cm above the calcaneal insertion in one study (Maffulli *et al.*, 2012). The third most common subjective criteria

**Table 11** Case reports.

Author	Year	Sample size	Location	Subjective history	Clinical tests
<i>Maffulli et al. (2011)</i>	2011	1 (M/F = 1/0)	Midportion	Location of pain (2–4 cm above calcaneal insertion) Duration of symptoms (>3 months) Pain with tendon loading	Pain on palpation Painful Arc Sign Royal London Hospital Test
<i>Van Sterkenburg et al. (2011)</i>	2011	3 (M/F = 1/2)	Midportion	Location of pain (4–7 cm above calcaneal insertion) Tendon stiffness	Pain on palpation
<i>Papa (2012)</i>	2012	1 (M/F = 0/1)	Midportion	Gradual onset of pain Pain with tendon loading	Pain on palpation Localised tendon thickening on palpation Localised swelling on palpation Reduced ROM Single-leg heel raise
<i>Benito (2016)</i>	2016	5 (M/F = 2/3)	Insertional	Location of pain (insertion) Gradual onset of pain Pain with tendon loading	Not specified
<i>Borda &amp; Selhorst (2017)</i>	2017	1 (M/F = 0/1)	Midportion	Gradual onset of pain Pain with tendon loading	Pain on palpation MMT Single-leg heel raise
<i>Jayaseelan, Weber &amp; Jonely (2019)</i>	2019	2 (M/F = 1/1)	Midportion	Location of pain Pain with tendon loading	Pain on palpation Single-leg heel raise Hopping Reduced ROM

**Note:**

cm, centimetres; M, male; F, female; ROM, range of motion; MMT, manual muscle test.

**Table 12** Protocols.

Author	Year	Location	Subjective history	Clinical tests
<i>Barker-Davies et al. (2017)</i>	2017	Insertional Midportion	Location of pain (2–6 cm above calcaneal insertion) Location of pain (insertion) Pain with tendon loading Tendon stiffness Change in training	Pain on palpation Single-leg heel raise Hopping
<i>Habets et al. (2017)</i>	2017	Midportion	Location of pain (2–7 cm above calcaneal insertion) Duration of symptoms (>3 months) Pain with tendon loading	Pain on palpation Localised swelling on palpation
<i>Mansur et al. (2017)</i>	2017	Insertional	Location of pain (distal 2 cm) Duration of symptoms (>3 months)	Pain on palpation
<i>Hasani et al. (2020)</i>	2020	Midportion	Location of pain (2–6 cm above calcaneal insertion) Pain with tendon loading Tendon stiffness VISA-A	Not specified
<i>Post et al. (2020)</i>	2020	Insertional Midportion	Location of pain Pain with tendon loading	Walking Single-leg heel raise Hopping

**Note:**


cm, centimetres; VISA-A, Victorian Institute of Sport Assessment-Achilles.

reported was tendon stiffness, with 24 studies including it as a diagnostic criteria (*Millar et al., 2021; Alfredson, 2003; Alfredson & Cook, 2007; Asplund & Best, 2013; Bains & Porter, 2006; Baskerville et al., 2018; Benazzo, Todesca & Ceciliani, 1997; Chazan, 1998;*

*Chimenti et al., 2017; Cook, Khan & Purdam, 2002; Courville, Coe & Hecht, 2009; Feilmeier, 2017; Fredericson, 1996; Hu & Flemister, 2008; Irwin, 2010; Jukes, Scott & Solan, 2020; Kader et al., 2002; Maffulli, Giuseppe Longo & Denaro, 2012; Maffulli et al., 2020; Maffulli et al., 2019; McShane, Ostick & McCabe, 2007; Nichols, 1989; Pedowitz & Beck, 2017; Tan & Chan, 2008*). As with previous study types, the most common clinical test used to diagnose Achilles tendinopathy was palpation, with all 43 included reviews including it as a clinical measure (*Millar et al., 2021; Aldridge, 2004; Alfredson, 2003; Alfredson & Cook, 2007; Aronow, 2005; Asplund & Best, 2013; Bains & Porter, 2006; Baskerville et al., 2018; Benazzo, Todesca & Ceciliani, 1997; Bhatty, Khan & Zubairy, 2019; Chazan, 1998; Chimenti et al., 2017; Cook, Khan & Purdam, 2002; Courville, Coe & Hecht, 2009; Den Hartog, 2009; Feilmeier, 2017; Fredericson, 1996; Furia & Rompe, 2007; Horn & McCollum, 2015; Hu & Flemister, 2008; Irwin, 2010; Järvinen et al., 2001; Jukes, Scott & Solan, 2020; Kader et al., 2002; Longo, Ronga & Maffulli, 2009; Maffulli, Giai Via & Oliva, 2015; Maffulli, Giuseppe Longo & Denaro, 2012; Maffulli & Kader, 2002; Maffulli et al., 2020; Maffulli et al., 2012; Maffulli et al., 2019; Maffulli, Sharma & Luscombe, 2004; Maffulli, Via & Oliva, 2014; McShane, Ostick & McCabe, 2007; Nichols, 1989; Paavola et al., 2002; Pedowitz & Beck, 2017; Saini et al., 2015; Scott, Huisman & Khan, 2011; Sharma & Maffulli, 2006; Simpson & Howard, 2009; Sorosky et al., 2004; Tan & Chan, 2008*). There was then significant variation in other clinical tests used to diagnose Achilles tendinopathy, with nine studies including the Painful Arc Sign (*Millar et al., 2021; Aronow, 2005; Feilmeier, 2017; Horn & McCollum, 2015; Kader et al., 2002; Longo, Ronga & Maffulli, 2009; Maffulli, Giuseppe Longo & Denaro, 2012; Maffulli & Kader, 2002; Maffulli et al., 2020*), seven studies including reduced range of motion (*Bhatty, Khan & Zubairy, 2019; Chazan, 1998; Chimenti et al., 2017; Furia & Rompe, 2007; Hu & Flemister, 2008; Maffulli et al., 2012; Nichols, 1989*), six studies including the Royal London Hospital Test (*Millar et al., 2021; Feilmeier, 2017; Horn & McCollum, 2015; Jukes, Scott & Solan, 2020; Longo, Ronga & Maffulli, 2009; Maffulli et al., 2020*), and six studies including pain whilst hopping as a clinical diagnostic criteria (*Millar et al., 2021; Alfredson & Cook, 2007; Bains & Porter, 2006; Cook, Khan & Purdam, 2002; Feilmeier, 2017; Nichols, 1989*).

### Case-reports

Table 11 highlights the characteristics of the included case report studies. Five studies (*Borda & Selhorst, 2017; Jayaseelan, Weber & Jonely, 2019; Maffulli et al., 2011; Papa, 2012; Van Sterkenburg et al., 2011*) investigated midportion Achilles tendinopathy and one study (*Benito, 2016*) investigated insertional Achilles tendinopathy. As with the narrative reviews, the most common subjective measure used to diagnose Achilles tendinopathy was pain with tendon loading (*Benito, 2016; Borda & Selhorst, 2017; Jayaseelan, Weber & Jonely, 2019; Maffulli et al., 2011; Papa, 2012*), with the second most common diagnostic criteria being location of pain (*Benito, 2016; Jayaseelan, Weber & Jonely, 2019; Maffulli et al., 2011; Van Sterkenburg et al., 2011*). Midportion Achilles tendinopathy



**Figure 4** Outcome measures used to measure Achilles tendinopathy categorised by purpose.

Full-size DOI: 10.7717/peerj.12166/fig-4

was defined as a location of pain 2–4 cm above the calcaneal insertion in one study ([Maffulli et al., 2011](#)), and 4–7 cm above the calcaneal insertion in another study ([Van Sterkenburg et al., 2011](#)). Five studies ([Borda & Selhorst, 2017](#); [Jayaseelan, Weber & Jonely, 2019](#); [Maffulli et al., 2011](#); [Papa, 2012](#); [Van Sterkenburg et al., 2011](#)) used palpation as an objective measure for diagnosing Achilles tendinopathy, with three studies ([Borda & Selhorst, 2017](#); [Jayaseelan, Weber & Jonely, 2019](#); [Papa, 2012](#)) utilising the pain during single-leg heel raise and one study ([Benito, 2016](#)) not specifying any objective clinical tests.

### Protocols

Of the five included protocol studies, two studies ([Habets et al., 2017](#); [Hasani et al., 2020](#)) investigated midportion Achilles tendinopathy, one study ([Mansur et al., 2017](#)) investigated insertional Achilles tendinopathy and two studies ([Barker-Davies et al., 2017](#); [Post et al., 2020](#)) investigated both insertional and midportion Achilles tendinopathy ([Table 12](#)). The most common reported subjective criteria utilised to diagnose Achilles tendinopathy was location of pain, with midportion Achilles tendinopathy defined as pain 2–6 cm above the calcaneal insertion in two studies ([Barker-Davies et al., 2017](#); [Hasani et al., 2020](#)), and 2–7 cm above the calcaneal insertion in one study ([Habets et al., 2017](#)). Insertional tendinopathy was defined as the distal 2 cm of the Achilles tendon in one study ([Mansur et al., 2017](#)) and the Achilles ‘insertion’ in another study ([Barker-Davies et al., 2017](#)). Clinical diagnostic tests varied with three studies ([Barker-Davies et al., 2017](#); [Habets et al., 2017](#); [Mansur et al., 2017](#)) including palpation, two studies ([Barker-Davies et al., 2017](#); [Post et al., 2020](#)) including pain during a single-leg heel raise and two studies including pain during hopping ([Aggarwal et al., 2015](#); [O’Neill et al., 2019](#)). One study did not specify objective clinical tests ([Hasani et al., 2020](#)).

### Outcome measures

Within the 159 included articles there were 42 different outcome measures in the clinical diagnosis of Achilles tendinopathy, with 49 studies ([Millar et al., 2021](#); [Reiman et al., 2014](#); [Aldridge, 2004](#); [Alfredson & Cook, 2007](#); [Aronow, 2005](#); [Barge-Caballero et al., 2008](#); [Baskerville et al., 2018](#); [Benazzo, Todesca & Ceciliani, 1997](#); [Bhatty, Khan & Zubairy, 2019](#); [Bjordal, Lopes-Martins & Iversen, 2006](#); [Cassel et al., 2018](#); [Chazan, 1998](#); [Chimenti et al., 2017](#); [Courville, Coe & Hecht, 2009](#); [De Jonge et al., 2011](#); [Den Hartog, 2009](#); [Docking et al., 2015](#); [Florit et al., 2019](#); [Fredericson, 1996](#); [Furia & Rompe, 2007](#); [Holmes & Lin, 2006](#); [Horn & McCollum, 2015](#); [Irwin, 2010](#); [Järvinen et al., 2001](#); [Jukes, Scott & Solan, 2020](#); [Kader et al., 2002](#); [Knobloch et al., 2008](#); [Maffulli, Giai Via & Oliva, 2015](#); [Maffulli & Kader, 2002](#); [Maffulli et al., 2003](#); [Maffulli et al., 2011](#); [Maffulli et al., 2020](#); [Maffulli et al., 2012](#); [Maffulli, Sharma & Luscombe, 2004](#); [Maffulli, Via & Oliva, 2014](#); [McShane, Ostick & McCabe, 2007](#); [Nichols, 1989](#); [Paavola et al., 2002](#); [Paavola et al., 2002](#); [Pedowitz & Beck, 2017](#); [Ryan et al., 2009](#); [Saini et al., 2015](#); [Sharma & Maffulli, 2006](#); [Simpson & Howard, 2009](#); [Sorosky et al., 2004](#); [Thomas et al., 2010](#); [Welsh & Clodman, 1980](#); [Xu et al., 2019](#); [Zhang et al., 2017](#)) not reporting any outcome measures. Of the 110 included studies to report on outcome measures ([Hutchison et al., 2013](#); [Abate & Salini, 2019](#); [Aiyegbusi, Tella & Sanusi, 2020](#); [Alfredson, 2003](#); [Alfredson & Spang, 2018](#); [Asplund & Best, 2013](#); [Azevedo et al., 2009](#); [Bains & Porter, 2006](#); [Barker-Davies et al., 2017](#); [Benito, 2016](#); [Boesen et al., 2017](#); [Borda & Selhorst, 2017](#); [Brown et al., 2006](#); [Carcia et al., 2010](#); [Cheng, Zhang & Cai, 2016](#); [Chester et al., 2008](#); [Chimenti et al., 2016](#); [Chimenti et al., 2020](#); [Cook, Khan & Purdam, 2002](#); [Coombes et al., 2018](#); [Creaby et al., 2017](#); [Crill, Berlet & Hyer, 2014](#); [De Marchi et al., 2018](#); [Divani et al., 2010](#); [Duthon et al., 2011](#); [Ebbesen et al., 2018](#); [Eckenrode, Kietrys & Stackhouse, 2019](#); [Feilmeier, 2017](#); [Finnamore et al., 2019](#); [Gärdin et al., 2016](#); [Gatz et al., 2020](#); [Habets et al., 2017](#); [Hasani et al., 2020](#); [Hernández-Sánchez et al., 2018](#); [Hu & Flemister, 2008](#); [Hutchison et al., 2011](#); [Jayaseelan, Weber & Jonely, 2019](#); [Jewson et al., 2017](#); [Jowett, Richmond & Bedi, 2018](#); [Karjalainen et al., 2000](#); [Khan et al., 2003](#); [Knobloch, 2007](#); [Knobloch et al., 2007](#); [Kragsnaes et al., 2014](#); [Krogh et al., 2016](#); [Lakshmanan & O'Doherty, 2004](#); [Leung & Griffith, 2008](#); [Lohrer & Nauck, 2009](#); [Longo et al., 2009](#); [Longo, Ronga & Maffulli, 2009](#); [Maffulli, Giuseppe Longo & Denaro, 2012](#); [Maffulli et al., 2008](#); [Maffulli et al., 2019](#); [Maffulli et al., 2008](#); [Mafi, Lorentzon & Alfredson, 2001](#); [Magnussen, Dunn & Thomson, 2009](#); [Mansur et al., 2019](#); [Mansur et al., 2017](#); [Mantovani et al., 2020](#); [Martin et al., 2018](#); [Mayer et al., 2007](#); [McCormack et al., 2015](#); [Murawski et al., 2014](#); [Nadeau et al., 2016](#); [Neeter et al., 2003](#); [De Mesquita et al., 2018](#); [O'Neill et al., 2019](#); [Oloff et al., 2015](#); [Ooi et al., 2015](#); [Paavola et al., 2000](#); [Paoloni et al., 2004](#); [Papa, 2012](#); [Petersen, Welp & Rosenbaum, 2007](#); [Pingel et al., 2013](#); [Post et al., 2020](#); [Praet et al., 2018](#); [Rabello et al., 2020](#); [Rasmussen et al., 2008](#); [Reid et al., 2012](#); [Reiter et al., 2004](#); [Romero-Morales et al., 2019a](#); [Rompe, Furia & Maffulli, 2009](#); [Rompe et al., 2008](#); [Rompe et al., 2007](#); [Roos et al., 2004](#); [Santamato et al., 2019](#); [Sayana & Maffulli, 2007](#); [Scholes et al., 2018](#); [Scott, Huisman & Khan, 2011](#); [Sengkerij et al., 2009](#); [Silbernagel et al., 2007](#); [Silbernagel et al., 2001](#); [Solomons et al., 2020](#); [Stenson et al., 2018](#); [Stergioulas et al., 2008](#); [Syvertson et al., 2017](#); [Tan & Chan, 2008](#); [Turner et al., 2018](#)

2020; Vallance et al., 2020; Van der Vlist et al., 2020; Van der Vlist et al., 2020; Van Sterkenburg et al., 2011; Verrall, Schofield & Brustad, 2011; Von Wehren et al., 2019; Wang et al., 2012; Wei et al., 2017; Zellers et al., 2019; Zhang et al., 2020; Zhuang et al., 2019; Romero-Morales et al., 2019b), 42 different outcome measures were utilised. Disability was the most commonly measured outcome, with 28 different outcome measures for disability being applied 135 times (Fig. 4). The most common outcome measure for disability was the VISA-A questionnaire, being used in 75% of the studies reporting outcome measures (Abate & Salini, 2019; Aiyegbusi, Tella & Sanusi, 2020; Alfredson & Spang, 2018; Asplund & Best, 2013; Bains & Porter, 2006; Barker-Davies et al., 2017; Benito, 2016; Boesen et al., 2017; Brown et al., 2006; Carcia et al., 2010; Cheng, Zhang & Cai, 2016; Chimenti et al., 2016; Chimenti et al., 2020; Cook, Khan & Purdam, 2002; Coombes et al., 2018; Creaby et al., 2017; Divani et al., 2010; Ebbesen et al., 2018; Eckenrode, Kietrys & Stackhouse, 2019; Feilmeier, 2017; Finnimore et al., 2019; Gårdin et al., 2016; Gatz et al., 2020; Habets et al., 2017; Hasani et al., 2020; Hernández-Sánchez et al., 2018; Hutchison et al., 2011; Jayaseelan, Weber & Jonely, 2019; Jewson et al., 2017; Jowett, Richmond & Bedi, 2018; Khan et al., 2003; Kragsnaes et al., 2014; Krogh et al., 2016; Lakshmanan & O'Doherty, 2004; Leung & Griffith, 2008; Lohrer & Nauck, 2009; Longo et al., 2009; Longo, Ronga & Maffulli, 2009; Maffulli, Giuseppe Longo & Denaro, 2012; Maffulli et al., 2008; Maffulli et al., 2019; Maffulli et al., 2008; Magnussen, Dunn & Thomson, 2009; Mansur et al., 2019; Mansur et al., 2017; Mantovani et al., 2020; Martin et al., 2018; McCormack et al., 2015; Nadeau et al., 2016; De Mesquita et al., 2018; O'Neill et al., 2019; Oloff et al., 2015; Ooi et al., 2015; Pingel et al., 2013; Praet et al., 2018; Rabello et al., 2020; Reid et al., 2012; Reiter et al., 2004; Romero-Morales et al., 2019a; Rompe, Furia & Maffulli, 2009; Rompe et al., 2008; Rompe et al., 2007; Santamato et al., 2019; Sayana & Maffulli, 2007; Scholes et al., 2018; Scott, Huisman & Khan, 2011; Sengkerij et al., 2009; Silbernagel et al., 2007; Solomons et al., 2020; Syvertson et al., 2017; Tan & Chan, 2008; Turner et al., 2020; Vallance et al., 2020; Van der Vlist et al., 2020; Van der Vlist et al., 2020; Van Sterkenburg et al., 2011; Von Wehren et al., 2019; Wang et al., 2012; Wei et al., 2017; Zellers et al., 2019; Zhang et al., 2020; Romero-Morales et al., 2019b). Following, disability, the second most common outcome measure was pain with the VAS (Hutchison et al., 2013; Alfredson, 2003; Alfredson & Spang, 2018; Barker-Davies et al., 2017; Boesen et al., 2017; Chester et al., 2008; Ebbesen et al., 2018; Habets et al., 2017; Hutchison et al., 2011; Knobloch, 2007; Knobloch et al., 2007; Maffulli et al., 2019; Mafi, Lorentzon & Alfredson, 2001; Magnussen, Dunn & Thomson, 2009; Mansur et al., 2019; Mansur et al., 2017; Petersen, Welp & Rosenbaum, 2007; Rasmussen et al., 2008; Reid et al., 2012; Romero-Morales et al., 2019a; Santamato et al., 2019; Scott, Huisman & Khan, 2011; Silbernagel et al., 2001; Stenson et al., 2018; Stergioulas et al., 2008; Turner et al., 2020; Van der Vlist et al., 2020; Van Sterkenburg et al., 2011; Verrall, Schofield & Brustad, 2011; Wei et al., 2017; Zhang et al., 2020; Zhuang et al., 2019; Romero-Morales et al., 2019b) being used in 30% of studies reporting outcome measures and numerical pain rating scale (NPRS) (Borda & Selhorst, 2017; Brown et al., 2006; Chimenti et al., 2016; Chimenti et al., 2020; Coombes et al., 2018; Hasani et al., 2020; Jayaseelan, Weber & Jonely, 2019; Krogh et al., 2016; Mantovani et al., 2020; O'Neill et al., 2019; Paoloni et al.,

2004; Papa, 2012; Post et al., 2020; Rompe, Furia & Maffulli, 2009; Rompe et al., 2008; Rompe et al., 2007; Syvertson et al., 2017; Vallance et al., 2020) being used in 17% of studies reporting outcome measures. Outcome measures related to quality of life were utilised within 12 studies, with the most common outcome measures used being the 12-Item Short Form Survey (SF-12) (Duthon et al., 2011; Habets et al., 2017; Mansur et al., 2017; Murawski et al., 2014; Stenson et al., 2018), EuroQol 5 Dimension 5 Level Questionnaire (EQ-5D-5L) (Chester et al., 2008; De Marchi et al., 2018; Habets et al., 2017; Hasani et al., 2020) and 36-Item Short Form Survey (SF-36) (Hutchison et al., 2011; Maffulli et al., 2019; Petersen, Welp & Rosenbaum, 2007). Similarly, psychosocial outcomes were poorly measured, being utilised on 11 occasions, with the most common outcome measures being the Pain Catastrophizing Scale (PCS) (Chimenti et al., 2020; Eckenrode, Kietrys & Stackhouse, 2019; Hasani et al., 2020; Post et al., 2020; Vallance et al., 2020), Tampa Kinesiophobia Scale (TKS) (Chimenti et al., 2020; Hasani et al., 2020; Post et al., 2020; Vallance et al., 2020), Pain Disability Index (PDI) (Mayer et al., 2007; Scott, Huisman & Khan, 2011) and Pain Efficacy Scale (PES) (Mayer et al., 2007; Scott, Huisman & Khan, 2011).

## DISCUSSION

### Overview

The clinical diagnosis of tendinopathy is commonly determined via both patient history and clinical tests (Cook et al., 2016; Coombes, Bisset & Vicenzino, 2015; Lewis, 2016; Lewis et al., 2015; Maffulli, Khan & Puddu, 1998; Malliaras et al., 2015; Scott et al., 2013).

However, with no consensus on gold standard clinical tests with which to diagnose tendinopathy (Docking, Ooi & Connell, 2015), many research studies utilise a variety of measures to diagnose Achilles tendinopathy (Hutchison et al., 2013). The primary aim of this scoping review was to provide a method for clinically diagnosing Achilles tendinopathy that aligns with the nine core health domains. In order to achieve this, specific objectives were determined that included identifying the most common clinical tests used to diagnose Achilles tendinopathy, identifying the most common outcome measures used to assess Achilles tendinopathy, and summarising the studies to date. This will allow for greater consistency in both research and clinical settings. Additionally, this review aimed to identify the both the areas of strength and weakness.

### Terminology

As highlighted in Fig. 3, ‘Tendinopathy’ was the most commonly term used to describe persistent Achilles tendon pain, particularly in more recent studies. This scoping review aligns with the previous consensus statements advocating the consistent use of the term tendinopathy to describe persistent Achilles tendon pain and associated loss of function in relation to mechanical loading (Scott et al., 2020). There was a noticeable reduction of the use of alternative terms such as tendinitis and tendinosis, particularly since 2018, indicating progression towards unifying the terminology used to describe the clinical condition of persistent pain and dysfunction in the Achilles tendon that is associated with mechanical loading.

A difficulty identified in this scoping review was the inclusion of symptom duration as a measure to diagnose Achilles tendinopathy (Tables 6, 7, 8 and 9). When used as a measure, duration of symptoms varied significantly from four weeks up to 12 months, making identifying a consistent duration of symptoms to diagnose Achilles tendinopathy difficult and potentially contributing to the different terminology used within research and clinical practice. The term tendinitis indicates an inflammatory condition of the Achilles tendon that may develop symptoms in a shorter duration of time, whereas tendinosis indicates a change in tendon structure that would require a longer duration of time for symptoms to develop (Scott *et al.*, 2020). Additionally, the clinical condition of Achilles tendinopathy does not display the characteristics of an inflammatory response such as with tissue tearing (Cook *et al.*, 2016), and the structural changes, as those expected in tendinosis, are not required to be present for pain or dysfunction to develop (Cook *et al.*, 2016).

### The nine core health domains of tendinopathy

Vicenzino *et al.* (2020) reported that the lack of agreed upon tendon health related domains impedes the progress of tendinopathy research. The nine identified domains (patient rating of overall condition, pain on activity or loading, participation, function, psychological factors, disability, physical function capacity, quality of life, and pain over a specified timeframe) should allow for greater consistency in the reporting of tendon research (Vicenzino *et al.*, 2020). This scoping review further highlights the inconsistency in the methods used to diagnose and assess Achilles tendinopathy. There was variation in the methodology used to clinically diagnose and assess Achilles tendinopathy for all key themes; subjective history, clinical tests and outcome measures.

### Subjective history

Multiple measures were identified to determine a diagnosis of Achilles tendinopathy from the subjective interview (Tables 4, 5, 6, 7, 8, 9, 10, 11 and 12). The most commonly used measure was self-reported location of pain, with midportion Achilles tendinopathy most commonly being defined as an area located 2–6 cm above the calcaneal insertion of the Achilles tendon. Insertional tendinopathy was most commonly defined as the distal 2 cm of the Achilles tendon. Additional measures included pain with tendon loading activity, duration of symptoms, and tendon stiffness following tendon loading or at a particular time of the day (*i.e.* morning stiffness). Interestingly, while a change in Achilles tendon loading activity (both an increase and decrease) is considered a catalyst for Achilles tendinopathy (Cook *et al.*, 2016), it was only utilised as a specific criterion in nine of the included studies (Bains & Porter, 2006; Barker-Davies *et al.*, 2017; Cook, Khan & Purdam, 2002; Courville, Coe & Hecht, 2009; Feilmeier, 2017; Maffulli *et al.*, 2012; Nichols, 1989; Simpson & Howard, 2009; Sorosky *et al.*, 2004).

### Objective clinical tests

As with subjective history, numerous clinical tests were identified to diagnose Achilles tendinopathy (Tables 4, 5, 6, 7, 8, 9, 10, 11 and 12). The most commonly identified clinical test for Achilles tendinopathy was tendon palpation (including pain on palpation,

localised tendon thickening or localised swelling). Although, palpation is commonly used to identify the region of pain and is a common clinical measure used to diagnose Achilles tendinopathy (*Hutchison et al., 2013; Reiman et al., 2014; Maffulli et al., 2003; Martin et al., 2018*), studies reported multiple regions of interest for midportion Achilles tendinopathy. Painful regions were described as the ‘midportion’, ‘middle third’, 2 to 4, 2 to 5, 2 to 6, 2 to 7, 4 to 6 and 4 to 7 cm above the calcaneal insertion. Similarly, the region of interest in insertional Achilles tendinopathy was described as the ‘insertion’, calcaneal tuberosity, distal 2 cm, and distal 5 cm.

While there was consistency in the included studies in their use of palpation as a clinical test, there is significant variation in the additional clinical tests used to confirm a diagnosis of Achilles tendinopathy ([Tables 4, 5, 6, 7, 8, 9, 10, 11 and 12](#)). Further clinical tests used to assess Achilles tendinopathy included tendon pain during loading activities (single-leg heel raises and hopping). The most frequently used clinical, tendinopathy specific tests, were the Royal London Hospital Test and the Painful Arc Sign. The Royal London Hospital Test is considered positive when there is a reduction in palpable Achilles tendon pain on ankle dorsiflexion (*Maffulli et al., 2003*). The Painful Arc Sign is considered positive when the area of swelling identified with palpation moves with active ankle plantarflexion and dorsiflexion (*Maffulli et al., 2003*).

### Outcome measures

As was the case for the clinical features, there were significant variations in the outcome measures utilised for making a diagnosis of Achilles tendinopathy ([Fig. 4](#)). While overall disability and participant perceived pain were commonly measured, the impact of Achilles tendinopathy on quality of life and psychological factors were rarely measured. Psychological factors such as pain efficacy, catastrophisation and kinesiophobia scales were identified as important outcome measures in the diagnosis tendinopathy (*Vicenzino et al., 2020*), which aligns with the identified psychological outcome measures (PES, TKS and PCS). Similarly, disability measures that combine patient rated pain and function in relation to tendon-specific activities were identified as integral to monitoring tendinopathy outcomes. This scoping review identified the VISA-A questionnaire as the most commonly used outcome measure to monitor Achilles tendinopathy. In addition to psychological factors and disability, overall quality of life was identified as a core health domain in tendinopathy (*Vicenzino et al., 2020*). The scoping review identified three different outcome measures (SF-12, SF-36 and EQ-5D-5L) that were utilised to assess participant quality of life.

### An example evidence-based method for clinically diagnosing Achilles tendinopathy

While there was significant variation in the methods used to diagnose Achilles tendinopathy, some common themes can be identified. When considering a consistent method for diagnosing and assessing Achilles tendinopathy, it is important to ensure research follows consensus recommendations on both terminology used and reporting outcomes (*Scott et al., 2020; Vicenzino et al., 2020*). Thus, [Table 13](#) provides an

**Table 13** A method for clinically diagnosing Achilles tendinopathy.

Test	Definition of test	Feature	Core health domain
<b>Subjective history</b>			
Self-reported location of pain	Clinician asks patient “Can you point out where you get your pain”	Pain located 2–6 cm above the calcaneal insertion (midportion) Pain located in the distal 2 cm of the Achilles tendon	N/A
Self-reported pain with tendon loading	Patient reported intensity of pain using a VAS or NPRS while performing an Achilles tendon-specific loading task (single-leg heel raise, hopping)	Patient reported increased pain on a VAS or NPRS with Achilles tendon-specific loading task (single-leg heel raise and hopping)	Pain with loading or activity
Self-reported tendon stiffness or pain over a specified time	Clinician asks about pain and stiffness over specified timeframes (e.g. morning, night, 24 hours)	Patient reported morning stiffness or pain Patient reported pain or stiffness at the onset of activity that may “warm-up”	Pain over a specified time
Self-reported overall rating of Achilles tendon	Clinician asks “Can you rate your Achilles tendon where 100% represents no problems and 0% is the worst-case scenario”	Patient reported level of condition	Patient rating of overall condition
<b>Objective tests</b>			
Palpation	Performed by the clinician gently palpating the whole length of the tendon in a proximal to distal direction	Patient reported pain located 2–6 cm above the calcaneal insertion (midportion) with or without subjective opinion of tendon thickening or swelling Patient reported pain located in the distal 2 cm of the Achilles tendon with or without subjective opinion of tendon thickening or swelling	N/A
Single-leg heel raise	Performed by patient rising up on to tip toes and lowering back down in a controlled manner, on both the affected and non-affected leg	Clinician recorded number of completed single-leg heel raises on each leg	Physical function capacity
Hopping	Performed by participant hopping on the spot	Clinician recorded number of completed hops on each leg	Physical function capacity
The Royal London Hospital Test	Performed by the clinician palpating the tendon for any local tenderness with the ankle either in neutral position or in slight plantarflexion. The ankle is then actively dorsiflexed and plantarflexed. With the ankle in maximum dorsiflexion, the portion of the tendon found to be tender is palpated again.	Patient reported pain on palpation reduces significantly or disappears with maximum dorsiflexion	N/A
Painful Arc Sign	Performed by the clinician identifying the intratendinous swelling in the tendon and asking the patient to actively dorsiflex and plantarflex the ankle joint observing the movement of the swelling between the malleoli	The intratendinous swelling moves relative to the malleoli with the Achilles tendon during the ankle movement	N/A
<b>Outcome measures</b>			
VISA-A	The VISA-A questionnaire is a valid and reliable tool to evaluate clinical severity of Achilles tendinopathy that has been translated into multiple languages. Patients can self-administer the questionnaire	The maximum score is 100, with healthy subjects scoring a minimum of 96 ( <a href="#">Malliaras et al., 2015</a> ).	Disability
PCS	Patients are asked to indicate the degree to which they have the above thoughts and feelings when they are experiencing pain using the 0 (not at all) to 4 (all the time) scale.	A total score is yielded (ranging from 0–52), with a score of 30 or below indicating a clinically relevant level of catastrophising ( <a href="#">Scott et al., 2013</a> ).	Psychological factors
SF-12	The SF-12 is a self-reported outcome measure assessing the impact of health on an individual’s everyday life.	The SF-12 creates two summary scores, mental health and physical health ( <a href="#">Pham et al., 2014</a> ).	Quality of Life

**Note:**

N/A, not applicable; cm, centimetres; VAS, Visual Analogue Scale; NPRS, Numerical Pain Rating Scale; VISA-A, Victorian Institute of Sport Assessment-Achilles; PCS, Pain Catastrophising Scale; SF-12, 12-Item Short Form Survey.

amalgamation of the common features used to diagnose Achilles tendinopathy identified in the scoping review and the previously identified nine core health domains for tendinopathy (*Vicenzino et al., 2020*). While the VISA-A is Achilles tendon specific, any validated and reliable pain questionnaire and quality of life questionnaire may be used in place of the Pain Catastrophising Scale and SF-12.

### Limitations

This review was limited to publications in English, which may have excluded key studies published in other languages. Additionally, the screening, inclusion, exclusion and data extraction was performed by one reviewer (WM), which decreases the probability all relevant studies were identified for review and could lead to reviewer bias. The methodological quality of the studies was not assessed as per guidelines for completing scoping reviews (*Peters et al., 2020; Arksey & O'Malley, 2005*), meaning studies of poor design are given equal weighting to those of better quality, however, the descriptive nature of the scoping review limits the potential impact of individual studies' methodological quality on results. The aim of a scoping review is to provide an overview of all literature within a field of evidence (*Pham et al., 2014*), and while there is no specific requirement for methodological quality appraisal, assessing individual literature methodological quality utilising a standardised tool may help authors identify gaps in the literature related to low quality research in addition to lack of research.

### CONCLUSIONS

The specific objectives, including the most common clinical tests used to diagnose Achilles tendinopathy and identifying the most common outcome measures used to assess Achilles tendinopathy were highlighted, with the scoping review identifying the significant variation in the methodology and outcome measures used to diagnose Achilles tendinopathy. This scoping review provides a detailed summary of the current evidence and common themes were identified in the available research to provide an evidence-based method to diagnose Achilles tendinopathy utilising both subjective and objective testing, in addition to recommendations regarding common outcome measures. The primary aim of this scoping review was to identify and provide a method for clinically diagnosing Achilles tendinopathy that aligns with the nine core health domains and a method for diagnosing Achilles tendinopathy is proposed, that includes both results from the scoping review and recent recommendations for reporting results in tendinopathy. The development of a method for the clinical diagnosis of Achilles tendinopathy is key to developing greater homogeneity in future research. By standardising the clinical diagnosis of Achilles tendinopathy, future research is able to investigate other areas of this complex condition and identifying possible subclassifications of Achilles tendinopathy and thus improving tailored individual treatment programmes or monitoring patient progress. Additionally, an evidence-based method for the clinical diagnosis of Achilles tendinopathy will allow clinicians to be more confident with their diagnosis and provide patients with greater certainty.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This research was supported by an Australian Government Research Training Program Scholarship. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:  
Australian Government Research Training Program Scholarship.

### Competing Interests

The authors declare that they have no competing interests.

### Author Contributions

- Wesley Matthews conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Richard Ellis conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- James Furness conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Wayne A. Hing conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.

### Data Availability

The following information was supplied regarding data availability:

The raw data is available in the [Supplementary File](#).

### Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.12166#supplemental-information>.

## REFERENCES

- Abate M, Salini V. 2019. Mid-portion Achilles tendinopathy in runners with metabolic disorders. *European Journal of Orthopaedic Surgery & Traumatology* 29(3):697–703 DOI [10.1007/s00590-018-2336-2](https://doi.org/10.1007/s00590-018-2336-2).
- Aggarwal R, Ringold S, Khanna D, Neogi T, Johnson SR, Miller A, Brunner HI, Ogawa R, Felson D, Oggie A, Aletaha D, Feldman BM. 2015. Distinctions between diagnostic and classification criteria? *Arthritis Care & Research* 67(7):891–897 DOI [10.1002/acr.22583](https://doi.org/10.1002/acr.22583).
- Aiyegbusi AI, Tella BA, Sanusi GA. 2020. Is genu varum a risk factor for the prevalence and severity of achilles tendinopathy? A cross-sectional study of Nigerian elite track and field athletes. *Nigerian Postgraduate Medical Journal* 27(2):87–92 DOI [10.4103/npmj.npmj\\_179\\_19](https://doi.org/10.4103/npmj.npmj_179_19).
- Aldridge T. 2004. Diagnosing heel pain in adults. *American Family Physician* 70(2):332–338.

- Alfredson H.** 2003. Chronic midportion Achilles tendinopathy: an update on research and treatment. *Clinics in Sports Medicine* **22**(4):727–741 DOI [10.1016/S0278-5919\(03\)00010-3](https://doi.org/10.1016/S0278-5919(03)00010-3).
- Alfredson H, Cook J.** 2007. A treatment algorithm for managing Achilles tendinopathy: new treatment options. *British Journal of Sports Medicine* **41**(4):211–216 DOI [10.1136/bjsm.2007.035543](https://doi.org/10.1136/bjsm.2007.035543).
- Alfredson H, Spang C.** 2018. Clinical presentation and surgical management of chronic Achilles tendon disorders—a retrospective observation on a set of consecutive patients being operated by the same orthopedic surgeon. *Foot and Ankle Surgery* **24**(6):490–494 DOI [10.1016/j.fas.2017.05.011](https://doi.org/10.1016/j.fas.2017.05.011).
- Arksey H, O'Malley L.** 2005. Scoping studies: towards a methodological framework. *International Journal of Social Research Methodology* **8**(1):19–32 DOI [10.1080/1364557032000119616](https://doi.org/10.1080/1364557032000119616).
- Aronow MS.** 2005. Posterior heel pain (retrocalcaneal bursitis, insertional and noninsertional Achilles tendinopathy). *Clinics in Podiatric Medicine and Surgery* **22**(1):19–43 DOI [10.1016/j.cpm.2004.08.003](https://doi.org/10.1016/j.cpm.2004.08.003).
- Asplund CA, Best TM.** 2013. Achilles tendon disorders. *BMJ* **346**(7899):29–33 DOI [10.1136/bmj.f1262](https://doi.org/10.1136/bmj.f1262).
- Azevedo LB, Lambert MI, Vaughan CL, O'Connor CM, Schwellnus MP.** 2009. Biomechanical variables associated with Achilles tendinopathy in runners. *British Journal of Sports Medicine* **43**(4):288–292 DOI [10.1136/bjsm.2008.053421](https://doi.org/10.1136/bjsm.2008.053421).
- Bains BS, Porter K.** 2006. Lower limb tendinopathy in athletes. *Trauma* **8**(4):213–224 DOI [10.1177/1460408606078110](https://doi.org/10.1177/1460408606078110).
- Barge-Caballero E, Crespo-Leiro MG, Paniagua-Martín MJ, Muñiz J, Naya C, Bouzas-Mosquera A, Piñón-Esteban P, Marzoa-Rivas R, Pazos-López P, Cursack GC, Cuena-Castillo JJ, Castro-Beiras A.** 2008. Quinolone-related Achilles tendinopathy in heart transplant patients: incidence and risk factors. *The Journal of Heart and Lung Transplantation* **27**(1):46–51 DOI [10.1016/j.healun.2007.09.021](https://doi.org/10.1016/j.healun.2007.09.021).
- Barker-Davies RM, Nicol A, McCurdie I, Watson J, Baker P, Wheeler P, Fong D, Lewis M, Bennett AN.** 2017. Study protocol: a double blind randomised control trial of high volume image guided injections in Achilles and patellar tendinopathy in a young active population. *BMC Musculoskeletal Disorders* **18**(1) DOI [10.1186/s12891-017-1564-7](https://doi.org/10.1186/s12891-017-1564-7).
- Baskerville R, McCartney DE, McCartney SM, Dawes H, Tan GD.** 2018. Tendinopathy in type 2 diabetes: a condition between specialties? *British Journal of General Practice* **68**(677):593–594 DOI [10.3399/bjgp18X700169](https://doi.org/10.3399/bjgp18X700169).
- Benazzo F, Todesca A, Ceciliani L.** 1997. Achilles' tendon tendinitis and heel pain. *Operative Techniques in Sports Medicine* **5**(3):179–188 DOI [10.1016/S1060-1872\(97\)80040-8](https://doi.org/10.1016/S1060-1872(97)80040-8).
- Benito E.** 2016. Physiotherapy's protocol to approach the insertional achilles tendinopathy. *Journal of Human Sport and Exercise* **11**(3):358–366 DOI [10.14198/jhse.2016.113.03](https://doi.org/10.14198/jhse.2016.113.03).
- Bhatty UN, Khan SHM, Zubairy AI.** 2019. Managing the patient with heel pain. *British Journal of Hospital Medicine* **80**(4):196–200 DOI [10.12968/hmed.2019.80.4.196](https://doi.org/10.12968/hmed.2019.80.4.196).
- Bjordal JM, Lopes-Martins RAB, Iversen VV.** 2006. A randomised, placebo controlled trial of low level laser therapy for activated Achilles tendinitis with microdialysis measurement of peritendinous prostaglandin E2 concentrations. *British Journal of Sports Medicine* **40**(1):76–80 DOI [10.1136/bjsm.2005.020842](https://doi.org/10.1136/bjsm.2005.020842).
- Boesen AP, Hansen R, Boesen MI, Malliaras P, Langberg H.** 2017. Effect of high-volume injection, platelet-rich plasma, and sham treatment in chronic midportion Achilles tendinopathy: a randomized double-blinded prospective study. *The American Journal of Sports Medicine* **45**(9):2034–2043 DOI [10.1177/0363546517702862](https://doi.org/10.1177/0363546517702862).

- Borda J, Selhorst M.** 2017. The use of compression tack and flossing along with lacrosse ball massage to treat chronic Achilles tendinopathy in an adolescent athlete: a case report. *Journal of Manual & Manipulative Therapy* **25**(1):57–61 DOI [10.1080/10669817.2016.1159403](https://doi.org/10.1080/10669817.2016.1159403).
- Brown R, Orchard J, Kinchington M, Hooper A, Nalder G.** 2006. Aprotinin in the management of Achilles tendinopathy: a randomised controlled trial. *British Journal of Sports Medicine* **40**(3):275–279 DOI [10.1136/bjsm.2005.021931](https://doi.org/10.1136/bjsm.2005.021931).
- Carcia CR, Martin RL, Houck J, Wukich DK.** 2010. Clinical guidelines—Achilles pain, stiffness, and muscle power deficits: Achilles tendinitis. *Journal of Orthopaedic & Sports Physical Therapy* **40**(9):A1–A26 DOI [10.2519/jospt.2010.0305](https://doi.org/10.2519/jospt.2010.0305).
- Cassel M, Risch L, Intzegiani K, Mueller J, Stoll J, Brecht P, Mayer F.** 2018. Incidence of Achilles and patellar tendinopathy in adolescent elite athletes. *International Journal of Sports Medicine* **39**(9):726–732 DOI [10.1055/a-0633-9098](https://doi.org/10.1055/a-0633-9098).
- Chazan IM.** 1998. Achilles tendinitis part II: clinical examination, differential diagnosis, and approaches to management. *Journal of Manual & Manipulative Therapy* **6**(2):70–77 DOI [10.1179/jmt.1998.6.2.70](https://doi.org/10.1179/jmt.1998.6.2.70).
- Cheng Y, Zhang J, Cai Y.** 2016. Utility of ultrasonography in assessing the effectiveness of extracorporeal shock wave therapy in insertional Achilles tendinopathy. *BioMed Research International* **2016**(2):1–5 DOI [10.1155/2016/2580969](https://doi.org/10.1155/2016/2580969).
- Chester R, Costa ML, Shepstone L, Cooper A, Donell ST.** 2008. Eccentric calf muscle training compared with therapeutic ultrasound for chronic Achilles tendon pain—a pilot study. *Manual Therapy* **13**(6):484–491 DOI [10.1016/j.math.2007.05.014](https://doi.org/10.1016/j.math.2007.05.014).
- Chimenti RL, Chimenti PC, Buckley MR, Houck JR, Flemister AS.** 2016. Utility of ultrasound for imaging osteophytes in patients with insertional Achilles tendinopathy. *Archives of Physical Medicine and Rehabilitation* **97**(7):1206–1209 DOI [10.1016/j.apmr.2015.12.009](https://doi.org/10.1016/j.apmr.2015.12.009).
- Chimenti RL, Cychosz CC, Hall MM, Phisitkul P.** 2017. Current concepts review update: insertional Achilles tendinopathy. *Foot & Ankle International* **38**(10):1160–1169 DOI [10.1177/1071100717723127](https://doi.org/10.1177/1071100717723127).
- Chimenti RL, Hall MM, Dilger CP, Merriwether EN, Wilken JM, Sluka KA.** 2020. Local anesthetic injection resolves movement pain, motor dysfunction, and pain catastrophizing in individuals with chronic Achilles tendinopathy: a nonrandomized clinical trial. *Journal of Orthopaedic & Sports Physical Therapy* **50**(6):334–343 DOI [10.2519/jospt.2020.9242](https://doi.org/10.2519/jospt.2020.9242).
- Cook JL, Khan KM, Purdam C.** 2002. Achilles tendinopathy. *Manual Therapy* **7**(3):121–130 DOI [10.1054/math.2002.0458](https://doi.org/10.1054/math.2002.0458).
- Cook JL, Rio E, Purdam CR, Docking SI.** 2016. Revisiting the continuum model of tendon pathology: what is its merit in clinical practice and research? *British Journal of Sports Medicine* **50**(19):1187–1191 DOI [10.1136/bjsports-2015-095422](https://doi.org/10.1136/bjsports-2015-095422).
- Coombes BK, Bisset L, Vicenzino B.** 2015. Management of lateral elbow tendinopathy: one size does not fit all. *Journal of Orthopaedic & Sports Physical Therapy* **45**(11):938–949 DOI [10.2519/jospt.2015.5841](https://doi.org/10.2519/jospt.2015.5841).
- Coombes BK, Tucker K, Vicenzino B, Vuvan V, Mellor R, Heales L, Nordez A, Hug F.** 2018. Achilles and patellar tendinopathy display opposite changes in elastic properties: a shear wave elastography study. *Scandinavian Journal of Medicine & Science in Sports* **28**(3):1201–1208 DOI [10.1111/sms.12986](https://doi.org/10.1111/sms.12986).
- Courville XF, Coe MP, Hecht PJ.** 2009. Current concepts review: noninsertional Achilles tendinopathy. *Foot & Ankle International* **30**(11):1132–1142 DOI [10.3113/FAI.2009.1132](https://doi.org/10.3113/FAI.2009.1132).

- Creaby MW, Honeywill C, Franettovich Smith MM, Schache AG, Crossley KM.** 2017. Hip biomechanics are altered in male runners with Achilles tendinopathy. *Medicine & Science in Sports & Exercise* **49**(3):549–554 DOI [10.1249/MSS.0000000000001126](https://doi.org/10.1249/MSS.0000000000001126).
- Crill MT, Berlet G, Hyer C.** 2014. Plantar flexor muscle architecture changes as a result of eccentric exercise in patients with Achilles tendinosis. *Foot & Ankle Specialist* **7**(6):460–465 DOI [10.1177/1938640014539812](https://doi.org/10.1177/1938640014539812).
- Daly J, Willis K, Small R, Green J, Welch N, Kealy M, Hughes E.** 2006. A hierarchy of evidence for assessing qualitative health research. *Journal of Clinical Epidemiology* **60**(1):43–49 DOI [10.1016/j.jclinepi.2006.03.014](https://doi.org/10.1016/j.jclinepi.2006.03.014).
- De Jonge S, Van den Berg C, De Vos RJ, Van der Heide HJL, Weir A, Verhaar JAN, Bierma-Zeinstra SMA, Tol JL.** 2011. Incidence of midportion Achilles tendinopathy in the general population. *British Journal of Sports Medicine* **45**(13):1026–1028 DOI [10.1136/bjsports-2011-090342](https://doi.org/10.1136/bjsports-2011-090342).
- De Marchi A, Pozza S, Cenna E, Cavallo F, Gays G, Simbula L, De Petro P, Massè A, Massazza G.** 2018. In Achilles tendinopathy, the neovascularization, detected by contrast-enhanced ultrasound (CEUS), is abundant but not related to symptoms. *Knee Surgery, Sports Traumatology, Arthroscopy* **26**(7):2051–2058 DOI [10.1007/s00167-017-4710-8](https://doi.org/10.1007/s00167-017-4710-8).
- De Mesquita GN, De Oliveira MNM, Rodrigues Matoso AE, De Moura Filho AG, De Oliveira RR.** 2018. Cross-cultural adaptation and measurement properties of the Brazilian Portuguese version of the Victorian Institute of Sport Assessment-Achilles (VISA-A) questionnaire. *Journal of Orthopaedic & Sports Physical Therapy* **48**(7):567–573 DOI [10.2519/jospt.2018.7897](https://doi.org/10.2519/jospt.2018.7897).
- De Vos R-J, Van der Vlist AC, Winters M, Van der Giesen F, Weir A.** 2021. Diagnosing Achilles tendinopathy is like delicious spaghetti carbonara: it is all about key ingredients, but not all chefs use the same recipe. *British Journal of Sports Medicine* **55**(5):247–248 DOI [10.1136/bjsports-2020-102863](https://doi.org/10.1136/bjsports-2020-102863).
- Den Hartog BD.** 2009. Insertional achilles tendinosis: pathogenesis and treatment. *Foot and Ankle Clinics* **14**(4):639–650 DOI [10.1016/j.fcl.2009.08.005](https://doi.org/10.1016/j.fcl.2009.08.005).
- Divani K, Chan O, Padhiar N, Twycross-Lewis R, Maffulli N, Crisp T, Morrissey D.** 2010. Site of maximum neovascularisation correlates with the site of pain in recalcitrant mid-tendon Achilles tendinopathy. *Manual Therapy* **15**(5):463–468 DOI [10.1016/j.math.2010.03.011](https://doi.org/10.1016/j.math.2010.03.011).
- Docking SI, Ooi CC, Connell D.** 2015. Tendinopathy: is imaging telling us the entire story? *Journal of Orthopaedic & Sports Physical Therapy* **45**(11):842–852 DOI [10.2519/jospt.2015.5880](https://doi.org/10.2519/jospt.2015.5880).
- Docking SI, Rosengarten SD, Daffy J, Cook J.** 2015. Structural integrity is decreased in both Achilles tendons in people with unilateral Achilles tendinopathy. *Journal of Science and Medicine in Sport* **18**(4):383–387 DOI [10.1016/j.jsams.2014.06.004](https://doi.org/10.1016/j.jsams.2014.06.004).
- Duthon VB, Lübbeke A, Duc SR, Stern R, Assal M.** 2011. Noninsertional Achilles tendinopathy treated with gastrocnemius lengthening. *Foot & Ankle International* **32**(4):375–379 DOI [10.3113/FAI.2011.0375](https://doi.org/10.3113/FAI.2011.0375).
- Ebbesen BH, Mølgård CM, Olesen JL, Gregersen HE, Simonsen O.** 2018. No beneficial effect of Polidocanol treatment in Achilles tendinopathy: a randomised controlled trial. *Knee Surgery, Sports Traumatology, Arthroscopy* **26**(7):2038–2044 DOI [10.1007/s00167-017-4675-7](https://doi.org/10.1007/s00167-017-4675-7).
- Eckenrode BJ, Kietrys DM, Stackhouse SK.** 2019. Pain sensitivity in chronic Achilles tendinopathy. *International Journal of Sports Physical Therapy* **14**(6):945–956.
- Evans D.** 2003. Hierarchy of evidence: a framework for ranking evidence evaluating healthcare interventions. *Journal of Clinical Nursing* **12**(1):77–84 DOI [10.1046/j.1365-2702.2003.00662.x](https://doi.org/10.1046/j.1365-2702.2003.00662.x).

- Feilmeier M.** 2017. Noninsertional Achilles tendinopathy pathologic background and clinical examination. *Clinics in Podiatric Medicine and Surgery* **34**(2):129–136 DOI [10.1016/j.cpm.2016.10.003](https://doi.org/10.1016/j.cpm.2016.10.003).
- Finnimore E, Waugh C, Solomons L, Ryan M, West C, Scott A.** 2019. Transverse tendon stiffness is reduced in people with Achilles tendinopathy: a cross-sectional study. *PLOS ONE* **14**(2):e0211863 DOI [10.1371/journal.pone.0211863](https://doi.org/10.1371/journal.pone.0211863).
- Florit D, Pedret C, Casals M, Malliaras P, Sugimoto D, Rodas G.** 2019. Incidence of tendinopathy in team sports in a multidisciplinary sports club over 8 seasons. *Journal of Sports Science and Medicine* **18**(4):780–788.
- Fredericson M.** 1996. Common injuries in runners: diagnosis, rehabilitation and prevention. *Sports Medicine* **21**(1):49–72 DOI [10.2165/00007256-199621010-00005](https://doi.org/10.2165/00007256-199621010-00005).
- Furia JP, Rompe JD.** 2007. Extracorporeal shock wave therapy in the treatment of chronic plantar fascitis and Achilles tendinopathy. *Current Opinion in Orthopaedics* **18**(2):102–111 DOI [10.1097/BCO.0b013e328013e594](https://doi.org/10.1097/BCO.0b013e328013e594).
- Gatz M, Betsch M, Dirrichs T, Schrading S, Tingart M, Michalik R, Quack V.** 2020. Eccentric and isometric exercises in Achilles tendinopathy evaluated by the VISA-A score and shear wave elastography. *Sports Healthports Health: A Multidisciplinary Approach* **12**(4):373–381 DOI [10.1177/1941738119893996](https://doi.org/10.1177/1941738119893996).
- Gärdin A, Rasinski P, Berglund J, Shalabi A, Schulte H, Brismar TB.** 2016. T2\* relaxation time in Achilles tendinosis and controls and its correlation with clinical score. *Journal of Magnetic Resonance Imaging* **43**(6):1417–1422 DOI [10.1002/jmri.25104](https://doi.org/10.1002/jmri.25104).
- Habets B, Van Cingel REH, Backx FJG, Huisstede BMA.** 2017. Alfredson versus Silbernagel exercise therapy in chronic midportion Achilles tendinopathy: study protocol for a randomized controlled trial. *BMC Musculoskeletal Disorders* **18**(1):1–9 DOI [10.1186/s12891-017-1656-4](https://doi.org/10.1186/s12891-017-1656-4).
- Hasani F, Haines TP, Munteanu SE, Vicenzino B, Malliaras P.** 2020. Efficacy of different load intensity and time-under-tension calf loading protocols for Achilles tendinopathy (the LOADIT trial): protocol for a randomised pilot study. *Pilot and Feasibility Studies* **6**(1):99 DOI [10.1186/s40814-020-00639-5](https://doi.org/10.1186/s40814-020-00639-5).
- Hernández-Sánchez S, Poveda-Pagán EJ, Alakhdar-Mohmara Y, Hidalgo MD, De-las-Peñas C, Arias-Buría JL.** 2018. Cross-cultural adaptation of the Victorian Institute of Sport Assessment-Achilles (VISA-A) questionnaire for spanish athletes with Achilles tendinopathy. *Journal of Orthopaedic & Sports Physical Therapy* **48**(2):111–120 DOI [10.2519/jospt.2018.7402](https://doi.org/10.2519/jospt.2018.7402).
- Holmes GB, Lin J.** 2006. Etiologic factors associated with symptomatic Achilles tendinopathy. *Foot & Ankle International* **27**(11):952–959 DOI [10.1177/107110070602701115](https://doi.org/10.1177/107110070602701115).
- Horn A, McCollum G.** 2015. Achilles tendinopathy—part 1: aetiology, diagnosis and non-surgical management. *The South African Orthopaedic Journal* **14**(3):24–31 DOI [10.17159/2309-8309/2015/v14n3a2](https://doi.org/10.17159/2309-8309/2015/v14n3a2).
- Hu CT, Flemister AS.** 2008. Insertional Achilles tendinopathy: surgical options. *Operative Techniques in Orthopaedics* **18**(4):247–253 DOI [10.1053/j.oto.2009.02.002](https://doi.org/10.1053/j.oto.2009.02.002).
- Hutchison AM, Beard D, Pallister I, Topliss CJ, Williams P.** 2011. Is physiotherapy effective for patients with a chronic mid-body Achilles tendinopathy? A systematic review of non-surgical and non-pharmacological interventions. *International Musculoskeletal Medicine* **33**(4):152–160 DOI [10.1179/1753615411Y.0000000010](https://doi.org/10.1179/1753615411Y.0000000010).
- Hutchison A-M, Evans R, Bodger O, Pallister I, Topliss C, Williams P, Vannet N, Morris V, Beard D.** 2013. What is the best clinical test for Achilles tendinopathy? *Foot and Ankle Surgery* **19**(2):112–117 DOI [10.1016/j.fas.2012.12.006](https://doi.org/10.1016/j.fas.2012.12.006).

- Irwin TA.** 2010. Current concepts review: insertional Achilles tendinopathy. *Foot & Ankle International* **31**(10):933–939 DOI [10.3113/FAI.2010.0933](https://doi.org/10.3113/FAI.2010.0933).
- Jayaseelan DJ, Weber MJ, Jonely H.** 2019. Potential nervous system sensitization in patients with persistent lower extremity tendinopathies: 3 case reports. *Journal of Orthopaedic & Sports Physical Therapy* **49**(4):272–279 DOI [10.2519/jospt.2019.8600](https://doi.org/10.2519/jospt.2019.8600).
- Jewson JL, Lambert EA, Docking S, Storr M, Lambert GW, Gaida JE.** 2017. Pain duration is associated with increased muscle sympathetic nerve activity in patients with Achilles tendinopathy. *Scandinavian Journal of Medicine & Science in Sports* **27**(12):1942–1949 DOI [10.1111/sms.12820](https://doi.org/10.1111/sms.12820).
- Jowett CRJ, Richmond A, Bedi HS.** 2018. Paratendinous scraping and excision of plantaris for Achilles tendinopathy. *Techniques in Foot & Ankle Surgery* **17**(1):27–30 DOI [10.1097/BTF.0000000000000168](https://doi.org/10.1097/BTF.0000000000000168).
- Jukes CP, Scott G, Solan MC.** 2020. Posterior heel pain. *Orthopaedics and Trauma* **34**(1):3–9 DOI [10.1016/j.mporth.2019.11.001](https://doi.org/10.1016/j.mporth.2019.11.001).
- Järvinen TAH, Kannus P, Paavola M, Järvinen TLN, Józsa L, Järvinen M.** 2001. Achilles tendon injuries. *Current Opinion in Rheumatology* **13**(2):150–155 DOI [10.1097/00002281-200103000-00009](https://doi.org/10.1097/00002281-200103000-00009).
- Kader D, Saxena A, Movin T, Maffulli N.** 2002. Achilles tendinopathy: some aspects of basic science and clinical management. *British Journal of Sports Medicine* **36**(4):239–249 DOI [10.1136/bjsm.36.4.239](https://doi.org/10.1136/bjsm.36.4.239).
- Karjalainen PT, Soila K, Aronen HJ, Pihlajamäki HK, Tynniinen O, Paavonen T, Tirman PFJ.** 2000. MR imaging of overuse injuries of the Achilles tendon. *AJR American Journal of Roentgenology* **175**(1):251–260 DOI [10.2214/ajr.175.1.1750251](https://doi.org/10.2214/ajr.175.1.1750251).
- Khan KM, Forster BB, Robinson J, Cheong Y, Louis L, Maclean L, Taunton JE.** 2003. Are ultrasound and magnetic resonance imaging of value in assessment of Achilles tendon disorders? A two year prospective study. *British Journal of Sports Medicine* **37**(2):149–153 DOI [10.1136/bjsm.37.2.149](https://doi.org/10.1136/bjsm.37.2.149).
- Knobloch K.** 2007. Eccentric training in Achilles tendinopathy: is it harmful to tendon microcirculation? *British Journal of Sports Medicine* **41**(6):e2 DOI [10.1136/bjsm.2006.030437](https://doi.org/10.1136/bjsm.2006.030437).
- Knobloch K, Kraemer R, Jagodzinski M, Zeichen J, Meller R, Vogt PM.** 2007. Eccentric training decreases paratendon capillary blood flow and preserves paratendon oxygen saturation in chronic Achilles tendinopathy. *Journal of Orthopaedic & Sports Physical Therapy* **37**(5):269–276 DOI [10.2519/jospt.2007.2296](https://doi.org/10.2519/jospt.2007.2296).
- Knobloch K, Schreibmueller L, Longo UG, Vogt PM.** 2008. Eccentric exercises for the management of tendinopathy of the main body of the Achilles tendon with or without an AirHeel™ Brace: a randomized controlled trial—B: effects of compliance. *Disability and Rehabilitation* **30**(20–22):1692–1696 DOI [10.1080/09638280701785676](https://doi.org/10.1080/09638280701785676).
- Kragsnaes MS, Fredberg U, Stribolt K, Kjaer SG, Bendix K, Ellingsen T.** 2014. Stereological quantification of immune-competent cells in baseline biopsy specimens from Achilles tendons: results from patients with chronic tendinopathy followed for more than 4 years. *The American Journal of Sports Medicine* **42**(10):2435–2445 DOI [10.1177/0363546514542329](https://doi.org/10.1177/0363546514542329).
- Krogh TP, Ellingsen T, Christensen R, Jensen P, Fredberg U.** 2016. Ultrasound-guided injection therapy of Achilles tendinopathy with platelet-rich plasma or saline. *The American Journal of Sports Medicine* **44**(8):1990–1997 DOI [10.1177/0363546516647958](https://doi.org/10.1177/0363546516647958).
- Lakshmanan P, O'Doherty DP.** 2004. Chronic Achilles tendinopathy: treatment with extracorporeal shock waves. *Foot and Ankle Surgery* **10**(3):125–130 DOI [10.1016/j.fas.2004.04.001](https://doi.org/10.1016/j.fas.2004.04.001).

- Leung JLY, Griffith JF.** 2008. Sonography of chronic achilles tendinopathy: a case-control study. *Journal of Clinical Ultrasound* **36**(1):27–32 DOI [10.1002/\(ISSN\)1097-0096](https://doi.org/10.1002/(ISSN)1097-0096).
- Lewis J.** 2016. Rotator cuff related shoulder pain: assessment, management and uncertainties. *Manual Therapy* **23**:57–68 DOI [10.1016/j.math.2016.03.009](https://doi.org/10.1016/j.math.2016.03.009).
- Lewis J, McCresh K, Roy J-S, Ginn K.** 2015. Rotator cuff tendinopathy: navigating the diagnosis-management conundrum. *Journal of Orthopaedic & Sports Physical Therapy* **45**(11):923–937 DOI [10.2519/jospt.2015.5941](https://doi.org/10.2519/jospt.2015.5941).
- Lohrer H, Nauck T.** 2009. Cross-cultural adaptation and validation of the VISA-A questionnaire for German-speaking Achilles tendinopathy patients. *BMC Musculoskeletal Disorders* **10**(1):134 DOI [10.1186/1471-2474-10-134](https://doi.org/10.1186/1471-2474-10-134).
- Longo UG, Rittweger J, Garau G, Radonic B, Gutwasser C, Gilliver SF, Kusy K, Zieliński J, Felsenberg D, Maffulli N.** 2009. No influence of age, gender, weight, height, and impact profile in achilles tendinopathy in masters track and field athletes. *The American Journal of Sports Medicine* **37**(7):1400–1405 DOI [10.1177/0363546509332250](https://doi.org/10.1177/0363546509332250).
- Longo UG, Ronga M, Maffulli N.** 2009. Achilles tendinopathy. *Sports Medicine and Arthroscopy Review* **17**(2):112–126 DOI [10.1097/JSA.0b013e3181a3d625](https://doi.org/10.1097/JSA.0b013e3181a3d625).
- Maffulli N, Giai Via A, Oliva F.** 2015. Chronic Achilles tendon disorders. *Clinics in Sports Medicine* **34**(4):607–624 DOI [10.1016/j.csm.2015.06.010](https://doi.org/10.1016/j.csm.2015.06.010).
- Maffulli N, Giuseppe Longo U, Denaro V.** 2012. Achilles tendinopathy in dancers. *Journal of Dance Medicine & Science* **16**(3):92–100.
- Maffulli N, Kader D.** 2002. Tendinopathy of tendo achillis. *The Journal of Bone and Joint Surgery. British volume* **84B**(1):1–8 DOI [10.1302/0301-620X.84B1.0840001](https://doi.org/10.1302/0301-620X.84B1.0840001).
- Maffulli N, Kenward MG, Testa V, Capasso G, Regine R, King JB.** 2003. Clinical diagnosis of Achilles tendinopathy with tendinosis. *Clinical Journal of Sport Medicine* **13**(1):11–15 DOI [10.1097/00042752-200301000-00003](https://doi.org/10.1097/00042752-200301000-00003).
- Maffulli N, Khan KM, Puddu G.** 1998. Overuse tendon conditions: time to change a confusing terminology. *Arthroscopy: The Journal of Arthroscopic & Related Surgery* **14**(8):840–843 DOI [10.1016/S0749-8063\(98\)70021-0](https://doi.org/10.1016/S0749-8063(98)70021-0).
- Maffulli N, Longo UG, Campi S, Denaro V.** 2011. Achilles tendinopathy. *Evidence-Based Orthopedics* **2011**:872–878.
- Maffulli N, Longo UG, Kadakia A, Spiezia F.** 2020. Achilles tendinopathy. *Foot and Ankle Surgery* **26**(3):240–249 DOI [10.1016/j.fas.2019.03.009](https://doi.org/10.1016/j.fas.2019.03.009).
- Maffulli N, Longo UG, Petrillo S, Denaro V.** 2012. Management of tendinopathies of the foot and ankle. *Orthopaedics and Trauma* **26**(4):259–264 DOI [10.1016/j.mporth.2012.05.008](https://doi.org/10.1016/j.mporth.2012.05.008).
- Maffulli N, Longo UG, Testa V, Oliva F, Capasso G, Denaro V.** 2008. Italian translation of the VISA-A score for tendinopathy of the main body of the Achilles tendon. *Disability and Rehabilitation* **30**(20–22):1635–1639 DOI [10.1080/09638280701785965](https://doi.org/10.1080/09638280701785965).
- Maffulli N, Saxena A, Wagner E, Torre G.** 2019. Achilles insertional tendinopathy: state of the art. *Journal of ISAKOS: Joint Disorders & Orthopaedic Sports Medicine* **4**(1):48–57 DOI [10.1136/jisakos-2017-000144](https://doi.org/10.1136/jisakos-2017-000144).
- Maffulli N, Sharma P, Luscombe KL.** 2004. Achilles tendinopathy: aetiology and management. *Journal of the Royal Society of Medicine* **97**(10):472–476 DOI [10.1177/0141076809701004](https://doi.org/10.1177/0141076809701004).
- Maffulli N, Via AG, Oliva F.** 2014. Achilles injuries in the athlete: noninsertional. *Operative Techniques in Sports Medicine* **22**(4):321–330 DOI [10.1053/j.otsm.2014.09.001](https://doi.org/10.1053/j.otsm.2014.09.001).

- Maffulli N, Walley G, Sayana M, Longo UG, Denaro V. 2008.** Eccentric calf muscle training in athletic patients with Achilles tendinopathy. *Disability and Rehabilitation* **30**(20–22):1677–1684 DOI [10.1080/09638280701786427](https://doi.org/10.1080/09638280701786427).
- Mafi N, Lorentzon R, Alfredson H. 2001.** Superior short-term results with eccentric calf muscle training compared to concentric training in a randomized prospective multicenter study on patients with chronic Achilles tendinosis. *Knee Surgery, Sports Traumatology, Arthroscopy* **9**(1):42–47 DOI [10.1007/s001670000148](https://doi.org/10.1007/s001670000148).
- Magnussen RA, Dunn WR, Thomson AB. 2009.** Nonoperative treatment of midportion achilles tendinopathy: a systematic review. *Clinical Journal of Sport Medicine* **19**(1):54–64 DOI [10.1097/JSM.0b013e31818ef090](https://doi.org/10.1097/JSM.0b013e31818ef090).
- Malliaras P, Cook J, Purdam C, Rio E. 2015.** Patellar tendinopathy: clinical diagnosis, load management, and advice for challenging case presentations. *Journal of Orthopaedic & Sports Physical Therapy* **45**(11):887–898 DOI [10.2519/jospt.2015.5987](https://doi.org/10.2519/jospt.2015.5987).
- Mansur NSB, Baumfeld T, Villalon F, Aoyama BT, Matsunaga FT, Dos Santos PRD, Dos Santos BS, Tamaoki MJS. 2019.** Shockwave therapy associated with eccentric strengthening for Achilles insertional tendinopathy: a prospective study. *Foot & Ankle Specialist* **12**(6):540–545 DOI [10.1177/1938640019826673](https://doi.org/10.1177/1938640019826673).
- Mansur NSB, Faloppa F, Bellotti JC, Ingham SJMN, Matsunaga FT, Dos Santos PRD, Dos Santos BS, Carrazzone OL, Peixoto G, Aoyama BT, Tamaoki MJS. 2017.** Shock wave therapy associated with eccentric strengthening versus isolated eccentric strengthening for Achilles insertional tendinopathy treatment: a double-blinded randomised clinical trial protocol. *BMJ Open* **7**(1):e013332 DOI [10.1136/bmjopen-2016-013332](https://doi.org/10.1136/bmjopen-2016-013332).
- Mantovani L, Maestroni L, Bettariga F, Gobbo M, Lopomo NF, McLean S. 2020.** Does isometric exercise improve leg stiffness and hop pain in subjects with Achilles tendinopathy? A feasibility study. *Physical Therapy in Sport* **46**(4):234–242 DOI [10.1016/j.ptsp.2020.09.005](https://doi.org/10.1016/j.ptsp.2020.09.005).
- Martin RL, Chimenti R, Cuddeford T, Houck J, Matheson JW, McDonough CM, Paulseth S, Wukich DK, Garcia CR. 2018.** Achilles pain, stiffness, and muscle power deficits: midportion Achilles tendinopathy revision 2018. *Journal of Orthopaedic & Sports Physical Therapy* **48**(5):A1–A38 DOI [10.2519/jospt.2018.0302](https://doi.org/10.2519/jospt.2018.0302).
- Mayer F, Hirschmüller A, Müller S, Schuberth M, Baur H. 2007.** Effects of short-term treatment strategies over 4 weeks in Achilles tendinopathy. *British Journal of Sports Medicine* **41**(7):e6 DOI [10.1136/bjsm.2006.031732](https://doi.org/10.1136/bjsm.2006.031732).
- McCormack J, Underwood F, Slaven E, Cappaert T. 2015.** The minimum clinically important difference on the VISA-A and LEFS for patients with insertional Achilles tendinopathy. *International Journal of Sports Physical Therapy* **10**(5):639–644.
- McShane JM, Ostic B, McCabe F. 2007.** Noninsertional Achilles tendinopathy: pathology and management. *Current Sports Medicine Reports* **6**(5):288–292 DOI [10.1097/01.CSMR.0000306490.15530.b6](https://doi.org/10.1097/01.CSMR.0000306490.15530.b6).
- Merlin T, Weston A, Tooher R. 2009.** Extending an evidence hierarchy to include topics other than treatment: revising the Australian ‘levels of evidence’. *BMC Medical Research Methodology* **9**(1):34 DOI [10.1186/1471-2288-9-34](https://doi.org/10.1186/1471-2288-9-34).
- Millar NL, Silbernagel KG, Thorborg K, Kirwan PD, Galatz LM, Abrams GD, Murrell GAC, McInnes IB, Rodeo SA. 2021.** Tendinopathy (primer). *Nature Reviews Disease Primers* **7**(1):1–21 DOI [10.1038/s41572-020-00234-1](https://doi.org/10.1038/s41572-020-00234-1).
- Murawski CD, Smyth NA, Newman H, Kennedy JG. 2014.** A single platelet-rich plasma injection for chronic midsubstance achilles tendinopathy: a retrospective preliminary analysis. *Foot & Ankle Specialist* **7**(5):372–376 DOI [10.1177/1938640014532129](https://doi.org/10.1177/1938640014532129).

- Nadeau M-J, Desrochers A, Lamontagne M, Larivière C, Gagnon DH.** 2016. Quantitative ultrasound imaging of Achilles tendon integrity in symptomatic and asymptomatic individuals: reliability and minimal detectable change. *Journal of Foot and Ankle Research* **9**(1):1026 DOI [10.1186/s13047-016-0164-3](https://doi.org/10.1186/s13047-016-0164-3).
- Neeter C, Thomeé R, Silbernagel KG, Thomeé P, Karlsson J.** 2003. Iontophoresis with or without dexamethazone in the treatment of acute Achilles tendon pain. *Scandinavian Journal of Medicine and Science in Sports* **13**(6):376–382 DOI [10.1046/j.1600-0838.2003.00305.x](https://doi.org/10.1046/j.1600-0838.2003.00305.x).
- Nichols AW.** 1989. Achilles tendinitis in running athletes. *Journal of the American Board of Family Medicine* **2**(3):196–203.
- Oloff L, Elmi E, Nelson J, Crain J.** 2015. Retrospective analysis of the effectiveness of platelet-rich plasma in the treatment of Achilles tendinopathy: pretreatment and posttreatment correlation of magnetic resonance imaging and clinical assessment. *Foot & Ankle Specialist* **8**(6):490–497 DOI [10.1177/1938640015599033](https://doi.org/10.1177/1938640015599033).
- Ooi CC, Schneider ME, Malliaras P, Chadwick M, Connell DA.** 2015. Diagnostic performance of axial-strain sonoelastography in confirming clinically diagnosed Achilles tendinopathy: comparison with B-Mode ultrasound and color doppler imaging. *Ultrasound in Medicine & Biology* **41**(1):15–25 DOI [10.1016/j.ultrasmedbio.2014.08.019](https://doi.org/10.1016/j.ultrasmedbio.2014.08.019).
- O'Neill S, Radia J, Bird K, Rathleff MS, Bandholm T, Jorgensen M, Thorborg K.** 2019. Acute sensory and motor response to 45-s heavy isometric holds for the plantar flexors in patients with Achilles tendinopathy. *Knee Surgery, Sports Traumatology, Arthroscopy* **27**(9):2765–2773.
- Paavola M, Kannus P, Järvinen TAH, Khan K, Józsa L, Järvinen M.** 2002. Achilles tendinopathy. *The Journal of Bone and Joint Surgery-American Volume* **84**(11):2062–2076 DOI [10.2106/00004623-200211000-00024](https://doi.org/10.2106/00004623-200211000-00024).
- Paavola M, Kannus P, Orava S, Pasanen M, Järvinen M.** 2002. Surgical treatment for chronic Achilles tendinopathy: a prospective seven month follow up study. *British Journal of Sports Medicine* **36**(3):178–182 DOI [10.1136/bjsm.36.3.178](https://doi.org/10.1136/bjsm.36.3.178).
- Paavola M, Kannus P, Paakkala T, Pasanen M, Jarvinen M.** 2000. Long-term prognosis of patients with Achilles tendinopathy: an observational 8-year follow-up study. *The American Journal of Sports Medicine* **28**(5):634–642 DOI [10.1177/03635465000280050301](https://doi.org/10.1177/03635465000280050301).
- Paoloni JA, Appleyard RC, Nelson J, Murrell GAC.** 2004. Topical glyceryl trinitrate treatment of chronic noninsertional Achilles tendinopathy: a randomized, double-blind, placebo-controlled trial. *The Journal of Bone & Joint Surgery* **86**(5):916–922 DOI [10.2106/00004623-200405000-00005](https://doi.org/10.2106/00004623-200405000-00005).
- Papa JA.** 2012. Conservative management of Achilles tendinopathy: a case report. *Journal of the Canadian Chiropractic Association* **56**(3):216–224.
- Pedowitz D, Beck D.** 2017. Presentation, diagnosis, and nonsurgical treatment options of the anterior tibial tendon, posterior tibial tendon, peroneals, and Achilles. *Foot and Ankle Clinics* **22**(4):677–687 DOI [10.1016/j.fcl.2017.07.012](https://doi.org/10.1016/j.fcl.2017.07.012).
- Peters M, Godfrey C, McInerney P, Munn Z, Tricco AC, Khalil H.** 2020. Chapter 11: scoping reviews (2020 version). In: Aromataris E, Munn Z, eds. *JBI Manual for Evidence*. Adelaide: JBI.
- Petersen W, Welp R, Rosenbaum D.** 2007. Chronic Achilles tendinopathy: a prospective randomized study comparing the therapeutic effect of eccentric training, the AirHeel brace, and a combination of both. *The American Journal of Sports Medicine* **35**(10):1659–1667 DOI [10.1177/0363546507303558](https://doi.org/10.1177/0363546507303558).
- Pham MT, Rajić A, Greig JD, Sargeant JM, Papadopoulos A, McEwen SA.** 2014. A scoping review of scoping reviews: advancing the approach and enhancing the consistency. *Research Synthesis Methods* **5**(4):371–385 DOI [10.1002/jrsm.1123](https://doi.org/10.1002/jrsm.1123).

- Pingel J, Harrison A, Simonsen L, Suetta C, Bülow J, Langberg H.** 2013. The microvascular volume of the achilles tendon is increased in patients with tendinopathy at rest and after a 1-hour treadmill run. *The American Journal of Sports Medicine* **41**(10):2400–2408 DOI [10.1177/0363546513498988](https://doi.org/10.1177/0363546513498988).
- Post AA, Rio EK, Sluka KA, Moseley GL, Bayman EO, Hall MM, De Cesar Netto C, Wilken JM, Danielson JF, Chimenti R.** 2020. Effect of pain education and exercise on pain and function in chronic Achilles tendinopathy: protocol for a double-blind, placebo-controlled randomized trial. *JMIR Research Protocols* **9**(11):e19111 DOI [10.2196/19111](https://doi.org/10.2196/19111).
- Praet SFE, Ong JH, Purdam C, Welvaert M, Lovell G, Dixon L, Gaida JE, Anglim J, Manzanero S, Vlahovich N, Hughes D, Waddington G.** 2018. Microvascular volume in symptomatic Achilles tendons is associated with VISA-A score. *Journal of Science and Medicine in Sport* **21**(12):1185–1191 DOI [10.1016/j.jsams.2018.05.013](https://doi.org/10.1016/j.jsams.2018.05.013).
- Rabello LM, Van den Akker-Scheek I, Kuipers IF, Diercks RL, Brink MS, Zwerver J.** 2020. Bilateral changes in tendon structure of patients diagnosed with unilateral insertional or midportion achilles tendinopathy or patellar tendinopathy. *Knee Surgery, Sports Traumatology, Arthroscopy* **28**(5):1631–1638 DOI [10.1007/s00167-019-05495-2](https://doi.org/10.1007/s00167-019-05495-2).
- Rasmussen S, Christensen M, Mathiesen I, Simonson O.** 2008. Shockwave therapy for chronic Achilles tendinopathy: a double-blind, randomized clinical trial of efficacy. *Acta Orthopaedica* **79**(2):249–256 DOI [10.1080/17453670710015058](https://doi.org/10.1080/17453670710015058).
- Reid D, McNair PJ, Johnson S, Potts G, Witvrouw E, Mahieu N.** 2012. Electromyographic analysis of an eccentric calf muscle exercise in persons with and without Achilles tendinopathy. *Physical Therapy in Sport* **13**(3):150–155 DOI [10.1016/j.ptsp.2011.08.003](https://doi.org/10.1016/j.ptsp.2011.08.003).
- Reiman M, Burgi C, Strube E, Prue K, Ray K, Elliott A, Goode A.** 2014. The utility of clinical measures for the diagnosis of Achilles tendon injuries: a systematic review with meta-analysis. *Journal of Athletic Training* **49**(6):820–829 DOI [10.4085/1062-6050-49.3.36](https://doi.org/10.4085/1062-6050-49.3.36).
- Reiter M, Ulreich N, Dirisamer A, Tscholakoff D, Bocek RA.** 2004. Colour and power doppler sonography in symptomatic Achilles tendon disease. *International Journal of Sports Medicine* **25**(4):301–305 DOI [10.1055/s-2004-815828](https://doi.org/10.1055/s-2004-815828).
- Romero-Morales C, Martín-Llantino PJ, César C-L, López-López D, Rén S-G, De-La-Cruz-Torres B, Rodríguez-Sanz D.** 2019a. Ultrasonography features of the plantar fascia complex in patients with chronic non-insertional achilles tendinopathy: a case-control study. *Sensors* **19**(9):2052–2059 DOI [10.3390/s19092052](https://doi.org/10.3390/s19092052).
- Romero-Morales C, Martín-Llantino PJ, César C-L, Palomo-López P, López-López D, Pareja-Galeano H, Rodríguez-Sanz D.** 2019b. Comparison of the sonographic features of the Achilles Tendon complex in patients with and without achilles tendinopathy: a case-control study. *Physical Therapy in Sport* **35**:122–126.
- Rompe JD, Furia J, Maffulli N.** 2009. Eccentric loading versus eccentric loading plus shock-wave treatment for midportion achilles tendinopathy: a randomized controlled trial. *The American Journal of Sports Medicine* **37**(3):463–470 DOI [10.1177/0363546508326983](https://doi.org/10.1177/0363546508326983).
- Rompe JD, Furia J, Maffulli N, Rompe JD, Furia J, Maffulli N.** 2008. Eccentric loading compared with shock wave treatment for chronic insertional achilles tendinopathy: a randomized, controlled trial. *The Journal of Bone and Joint Surgery-American Volume* **90**(1):52–61 DOI [10.2106/JBJS.F.01494](https://doi.org/10.2106/JBJS.F.01494).
- Rompe JD, Nafe B, Furia JP, Maffulli N.** 2007. Eccentric loading, shock-wave treatment, or a wait-and-see policy for tendinopathy of the main body of tendo Achillis: a randomized controlled trial. *The American Journal of Sports Medicine* **35**(3):374–383 DOI [10.1177/0363546506295940](https://doi.org/10.1177/0363546506295940).

- Roos EM, Engström M, Lagerquist A, Söderberg B.** 2004. Clinical improvement after 6 weeks of eccentric exercise in patients with mid-portion Achilles tendinopathy: a randomized trial with 1-year follow-up. *Scandinavian Journal of Medicine and Science in Sports* **14**(5):286–295 DOI [10.1111/j.1600-0838.2004.378.x](https://doi.org/10.1111/j.1600-0838.2004.378.x).
- Ryan M, Grau S, Krauss I, Maiwald C, Taunton J, Hostmann T.** 2009. Kinematic analysis of runners with Achilles mid-portion tendinopathy. *Foot & Ankle International* **30**(12):1190–1195 DOI [10.3113/FAI.2009.1190](https://doi.org/10.3113/FAI.2009.1190).
- Saini SS, Reb CW, Chapter M, Daniel JN.** 2015. Achilles tendon disorders. *Journal of Osteopathic Medicine* **115**(11):670–676 DOI [10.7556/jaoa.2015.138](https://doi.org/10.7556/jaoa.2015.138).
- Santamato A, Beatrice R, Micello MF, Fortunato F, Panza F, Bristogiannis C, Cleopazzo E, Macarini L, Picelli A, Baricich A, Ranieri M.** 2019. Power doppler ultrasound findings before and after focused extracorporeal shock wave therapy for Achilles tendinopathy: a pilot study on pain reduction and neovascularization effect. *Ultrasound in Medicine & Biology* **45**(5):1316–1323 DOI [10.1016/j.ultrasmedbio.2018.12.009](https://doi.org/10.1016/j.ultrasmedbio.2018.12.009).
- Sayana M, Maffulli N.** 2007. Eccentric calf muscle training in non-athletic patients with achilles tendinopathy. *Journal of Science and Medicine in Sport* **10**(1):52–58 DOI [10.1016/j.jsams.2006.05.008](https://doi.org/10.1016/j.jsams.2006.05.008).
- Scholes M, Stadler S, Connell D, Barton C, Clarke RA, Bryant AL, Malliaras P.** 2018. Men with unilateral Achilles tendinopathy have impaired balance on the symptomatic side. *Journal of Science and Medicine in Sport* **21**(5):479–482 DOI [10.1016/j.jsams.2017.09.594](https://doi.org/10.1016/j.jsams.2017.09.594).
- Scott A, Docking S, Vicenzino B, Alfredson H, Zwerver J, Lundgreen K, Finlay O, Pollock N, Cook JL, Fearon A, Purdam CR, Hoens A, Rees JD, Goetz TJ, Danielson P.** 2013. Sports and exercise-related tendinopathies: a review of selected topical issues by participants of the second International Scientific Tendinopathy Symposium (ISTS) Vancouver 2012. *British Journal of Sports Medicine* **47**(9):536–544 DOI [10.1136/bjsports-2013-092329](https://doi.org/10.1136/bjsports-2013-092329).
- Scott A, Huisman E, Khan K.** 2011. Conservative treatment of chronic Achilles tendinopathy. *Canadian Medical Association Journal* **183**(10):1159–1165 DOI [10.1503/cmaj.101680](https://doi.org/10.1503/cmaj.101680).
- Scott A, Squier K, Alfredson H, Bahr R, Cook JL, Coombes B, De Vos R-J, Fu SN, Grimaldi A, Lewis JS, Maffulli N, Magnusson SP, Malliaras P, Mc Auliffe S, Oei EHG, Purdam CR, Rees JD, Rio EK, Gravare Silbernagel K, Speed C, Weir A, Wolf JM, Van den Akker-Scheek I, Vicenzino BT, Zwerver J.** 2020. ICON 2019: international scientific tendinopathy symposium consensus: clinical terminology. *British Journal of Sports Medicine* **54**(5):260–262 DOI [10.1136/bjsports-2019-100885](https://doi.org/10.1136/bjsports-2019-100885).
- Sengkerij PM, De Vos R-J, Weir A, Van Weerde BJG, Tol JL.** 2009. Interobserver reliability of neovascularization score using power doppler ultrasonography in midportion achilles tendinopathy. *The American Journal of Sports Medicine* **37**(8):1627–1631 DOI [10.1177/0363546509332255](https://doi.org/10.1177/0363546509332255).
- Sharma P, Maffulli N.** 2006. Understanding and managing Achilles tendinopathy. *British Journal of Hospital Medicine* **67**(2):64–67 DOI [10.12968/hmed.2006.67.2.20463](https://doi.org/10.12968/hmed.2006.67.2.20463).
- Silbernagel K, Thomeé R, Eriksson BI, Karlsson J.** 2007. Full symptomatic recovery does not ensure full recovery of muscle-tendon function in patients with Achilles tendinopathy. *British Journal of Sports Medicine* **41**(4):276–280 DOI [10.1136/bjsm.2006.033464](https://doi.org/10.1136/bjsm.2006.033464).
- Silbernagel KG, Thomeé R, Thomeé P, Karlsson J.** 2001. Eccentric overload training for patients with chronic Achilles tendon pain: a randomised controlled study with reliability testing of the evaluation methods. *Scandinavian Journal of Medicine & Science in Sports* **11**(4):197–206 DOI [10.1034/j.1600-0838.2001.110402.x](https://doi.org/10.1034/j.1600-0838.2001.110402.x).

- Simpson MR, Howard TM.** 2009. Tendinopathies of the foot and ankle. *American Family Physician* **80**(10):1107–1114.
- Solomons L, Lee JJY, Bruce M, White LD, Scott A.** 2020. Intramuscular stimulation vs sham needling for the treatment of chronic midportion Achilles tendinopathy: a randomized controlled clinical trial. *PLOS ONE* **15**(9):e0238579 DOI [10.1371/journal.pone.0238579](https://doi.org/10.1371/journal.pone.0238579).
- Sorosky B, Press J, Plastaras C, Rittenberg J.** 2004. The practical management of Achilles tendinopathy. *Clinical Journal of Sport Medicine* **14**(1):40–44 DOI [10.1097/00042752-200401000-00007](https://doi.org/10.1097/00042752-200401000-00007).
- Stenson JF, Reb CW, Daniel JN, Saini SS, Albana MF.** 2018. Predicting failure of nonoperative treatment for insertional Achilles tendinosis. *Foot & Ankle Specialist* **11**(3):252–255 DOI [10.1177/1938640017729497](https://doi.org/10.1177/1938640017729497).
- Stergioulas A, Stergioula M, Aarskog R, Lopes-Martins RA, Bjordal JM.** 2008. Effects of low-level laser therapy and eccentric exercises in the treatment of recreational athletes with chronic achilles tendinopathy. *The American Journal of Sports Medicine* **36**(5):881–887 DOI [10.1177/0363546507312165](https://doi.org/10.1177/0363546507312165).
- Syvertson P, Dietz E, Matocha M, McMurray J, Baker R, Nasypyany A, Reordan D, Paddock M.** 2017. A treatment-based classification algorithm to treat achilles tendinopathy: an exploratory case series. *Journal of Sport Rehabilitation* **26**(3):260–268 DOI [10.1123/jsr.2016-0033](https://doi.org/10.1123/jsr.2016-0033).
- Tan SC, Chan O.** 2008. Achilles and patellar tendinopathy: current understanding of pathophysiology and management. *Disability and Rehabilitation* **30**(20–22):1608–1615 DOI [10.1080/09638280701792268](https://doi.org/10.1080/09638280701792268).
- Thomas JL, Christensen JC, Kravitz SR, Mendicino RW, Schuberth JM, Vanore JV, Weil LS Sr., Zlotoff HJ, Bouché R, Baker J.** 2010. The diagnosis and treatment of heel pain: a clinical practice guideline-revision 2010. *Journal of Foot and Ankle Surgery* **49**(3):S1–S19 DOI [10.1053/j.jfas.2010.01.001](https://doi.org/10.1053/j.jfas.2010.01.001).
- Thomas J, Harden A.** 2008. Methods for the thematic synthesis of qualitative research in systematic reviews. *BMC Medical Research Methodology* **8**(1):45 DOI [10.1186/1471-2288-8-45](https://doi.org/10.1186/1471-2288-8-45).
- Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, Moher D, Peters MDJ, Horsley T, Weeks L, Hempel S, Akl EA, Chang C, McGowan J, Stewart L, Hartling L, Aldcroft A, Wilson MG, Garrity C, Lewin S, Godfrey CM, Macdonald MT, Langlois EV, Soares-Weiser K, Moriarty J, Clifford T, Tunçalp Ö, Straus SE.** 2018. PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Annals of Internal Medicine* **169**(7):467–473 DOI [10.7326/M18-0850](https://doi.org/10.7326/M18-0850).
- Turner J, Malliaras P, Goulis J, Mc Auliffe S.** 2020. It's disappointing and it's pretty frustrating, because it feels like it's something that will never go away: a qualitative study exploring individuals' beliefs and experiences of Achilles tendinopathy. *PLOS ONE* **15**(5):e0233459 DOI [10.1371/journal.pone.0233459](https://doi.org/10.1371/journal.pone.0233459).
- Vallance P, Hasani F, Crowley L, Malliaras P.** 2020. Self-reported pain with single leg heel raise or single leg hop offer distinct information as measures of severity in men with midportion and insertional Achilles tendinopathy: an observational cross-sectional study. *Physical Therapy in Sport* **47**(12):23–31 DOI [10.1016/j.ptsp.2020.10.009](https://doi.org/10.1016/j.ptsp.2020.10.009).
- Van der Vlist AC, Van Veldhoven PLJ, Van Oosterom RF, Verhaar JAN, De Vos RJ.** 2020. Isometric exercises do not provide immediate pain relief in Achilles tendinopathy: a quasi-randomized clinical trial. *Scandinavian Journal of Medicine & Science in Sports* **30**(9):1712–1721 DOI [10.1111/sms.13728](https://doi.org/10.1111/sms.13728).
- Van der Vlist AC, Veen JM, Van Oosterom RF, Van Veldhoven PLJ, Verhaar JAN, De Vos RJ.** 2020. Ultrasound doppler flow in patients with chronic midportion Achilles tendinopathy: is

surface area quantification a reliable method? *Journal of Ultrasound in Medicine* **39**(4):731–739  
DOI [10.1002/jum.15152](https://doi.org/10.1002/jum.15152).

- Van Sterkenburg MN, Kerkhoffs GM, Van Dijk CN, Van Sterkenburg MN, Kerkhoffs GMMJ, Van Dijk CN.** 2011. Good outcome after stripping the plantaris tendon in patients with chronic mid-portion Achilles tendinopathy. *Knee Surgery, Sports Traumatology, Arthroscopy* **19**(8):1362–1366 DOI [10.1007/s00167-011-1514-0](https://doi.org/10.1007/s00167-011-1514-0).
- Verrall G, Schofield S, Brustad T.** 2011. Chronic achilles tendinopathy treated with eccentric stretching program. *Foot & Ankle International* **32**(9):843–849 DOI [10.3113/FAI.2011.0843](https://doi.org/10.3113/FAI.2011.0843).
- Vicenzino B, De Vos R-J, Alfredson H, Bahr R, Cook JL, Coombes BK, Fu SN, Gravare Silbernagel K, Grimaldi A, Lewis JS, Maffulli N, Magnusson SP, Malliaras P, Mc Auliffe S, Oei EHG, Purdam C, Rees JD, Rio EK, Scott A, Speed C, Van den Akker-Scheek I, Weir A, Wolf JM, Zwerver J.** 2020. ICON 2019—international scientific tendinopathy symposium consensus: there are nine core health-related domains for tendinopathy (CORE DOMAINS)—delphi study of healthcare professionals and patients. *British Journal of Sports Medicine* **54**(8):444–451 DOI [10.1136/bjsports-2019-100894](https://doi.org/10.1136/bjsports-2019-100894).
- Von Wehren L, Pokorny K, Blanke F, Sailer J, Majewski M.** 2019. Injection with autologous conditioned serum has better clinical results than eccentric training for chronic Achilles tendinopathy. *Knee Surgery, Sports Traumatology, Arthroscopy* **27**(9):2744–2753 DOI [10.1007/s00167-019-05465-8](https://doi.org/10.1007/s00167-019-05465-8).
- Wang HK, Lin KH, Su SC, Shih TTF, Huang YC.** 2012. Effects of tendon viscoelasticity in Achilles tendinosis on explosive performance and clinical severity in athletes. *Scandinavian Journal of Medicine & Science in Sports* **22**(6):e147–e155 DOI [10.1111/j.1600-0838.2012.01511.x](https://doi.org/10.1111/j.1600-0838.2012.01511.x).
- Wei M, Liu Y, Li Z, Wang Z.** 2017. Comparison of clinical efficacy among endoscopy-assisted radio-frequency ablation, extracorporeal shockwaves, and eccentric exercises in treatment of insertional achilles tendinosis. *Journal of the American Podiatric Medical Association* **107**(1):11–16 DOI [10.7547/14-146](https://doi.org/10.7547/14-146).
- Welsh RP, Clodman J.** 1980. Clinical survey of Achilles tendinitis in athletes. *Canadian Medical Association Journal* **122**(2):193–195.
- Xu H, Li H, Hua Y, Bai L, Chang F, Chen S, Chen W, Fang Z, Gui J, Guo Q, Hu Y, Huang H, Jiao C, Li Q, Li W, Liang X, Lu H, Lu L, Miao X, Qu J, Song W, Xu T, Wang Z, Wang X, Wei M, Wei S, Xiang D, Xu H, Yang M, Zhan J, Zhang F, Zhao F, Zhu Y, Tang K.** 2019. Chinese consensus on insertional Achilles tendinopathy. *Orthopaedic Journal of Sports Medicine* **7**(10):232596711987905 DOI [10.1177/2325967119879052](https://doi.org/10.1177/2325967119879052).
- Zellers JA, Bley BC, Pohlig RT, Hamdan AN, Grävare SK.** 2019. Frequency of pathology on diagnostic ultrasound and relationship to patient demographics in individuals with insertional Achilles tendinopathy. *International Journal of Sports Physical Therapy* **14**(5):761–769 DOI [10.26603/ijsp20190761](https://doi.org/10.26603/ijsp20190761).
- Zhang Q, Cai Y, Hua Y, Shi J, Wang Y, Wang Y.** 2017. Sonoelastography shows that Achilles tendons with insertional tendinopathy are harder than asymptomatic tendons. *Knee Surgery, Sports Traumatology, Arthroscopy* **25**(6):1839–1848 DOI [10.1007/s00167-016-4197-8](https://doi.org/10.1007/s00167-016-4197-8).
- Zhang S, Li H, Yao W, Hua Y, Li Y.** 2020. Therapeutic response of extracorporeal shock wave therapy for insertional Achilles tendinopathy between sports-active and nonsports-active patients with 5-year follow-up. *Orthopaedic Journal of Sports Medicine* **8**(1):2325967119898118 DOI [10.1177/2325967119898118](https://doi.org/10.1177/2325967119898118).
- Zhuang Z, Yang Y, Chhantyal K, Chen J, Yuan G, Ni Y, Liu D, Shi D.** 2019. Central tendon-splitting approach and double row suturing for the treatment of insertional Achilles tendinopathy. *BioMed Research International* **2019**:1–10 DOI [10.1155/2019/4920647](https://doi.org/10.1155/2019/4920647).