



Series: The research agenda for general practice/ family medicine and primary health care in Europe. Part 2. Results: Primary care management and community orientation¹

Eva Hummers-Pradier, Martin Beyer, Patrick Chevallier, Sophia Eilat-Tsanani, Christos Lionis, Lieve Peremans, Davorina Petek, Imre Rurik, Jean Karl Soler, Henri Ejh Stoffers, Pinar Topsever, Mehmet Ungan & Paul van Royen

To cite this article: Eva Hummers-Pradier, Martin Beyer, Patrick Chevallier, Sophia Eilat-Tsanani, Christos Lionis, Lieve Peremans, Davorina Petek, Imre Rurik, Jean Karl Soler, Henri Ejh Stoffers, Pinar Topsever, Mehmet Ungan & Paul van Royen (2010) Series: The research agenda for general practice/family medicine and primary health care in Europe. Part 2. Results: Primary care management and community orientation¹, The European Journal of General Practice, 16:1, 42-50, DOI: [10.3109/13814780903563725](https://doi.org/10.3109/13814780903563725)

To link to this article: <https://doi.org/10.3109/13814780903563725>



Published online: 26 Jan 2010.



Submit your article to this journal [↗](#)



Article views: 655



View related articles [↗](#)



Citing articles: 9 View citing articles [↗](#)

BACKGROUND PAPER

Series: The research agenda for general practice/family medicine and primary health care in Europe. Part 2. Results: Primary care management and community orientation¹

EVA HUMMERS-PRADIER¹, MARTIN BEYER², PATRICK CHEVALLIER³, SOPHIA EILAT-TSANANI^{4,5}, CHRISTOS LIONIS⁶, LIEVE PEREMANS^{7,8}, DAVORINA PETEK⁹, IMRE RURIK¹⁰, JEAN KARL SOLER¹¹, HENRI EIJH STOFFERS¹², PINAR TOPSEVER¹³, MEHMET UNGAN¹⁴ & PAUL VAN ROYEN¹⁵

¹Institute of General Practice, Hannover Medical School, Hannover, Germany; ²Institute for General Practice, University of Frankfurt, Frankfurt, Germany; ³Department of General Practice, Saint-Quentin University Versailles, France; ⁴Department of Family Medicine, Emek Medical Center, Faculty of Health Sciences, Ben-Gurion University of the Negev, Israel; ⁵Afula and the valleys, Clalit Health Services, Northern District; ⁶Clinic of Social and Family Medicine, and Department of Social Medicine, University of Crete, Crete, Greece; ⁷Department of Primary, Interdisciplinary University of Antwerp, Antwerp, Belgium; ⁸Department of Youth Health, Vrije Universiteit of Brussels, Brussels, Belgium; ⁹Department of Family Medicine, Medical School, University of Ljubljana, Slovenia; ¹⁰Department of Family Medicine, University Debrecen, Hungary; ¹¹Mediterranean Institute of Primary Care, Attard, Malta; ¹²Department of General Practice, School of Public Health and Primary Care (CAPHRI), Maastricht University Medical Centre, Maastricht, The Netherlands; ¹³Department of Family Medicine, Acibadem University, Istanbul, Turkey; ¹⁴Middle East Technical University Medical Centre, Ankara, Turkey; ¹⁵Department of Primary and Interdisciplinary Care, University of Antwerp, Belgium

Abstract

At the WONCA Europe conference 2009 the recently published 'Research Agenda for General Practice/Family Medicine and Primary Health Care in Europe' was presented. It is a background paper and reference manual, providing advocacy of general practice/family medicine (GP/FM) in Europe. The Research Agenda summarizes the evidence relating to the core competencies and characteristics of the WONCA Europe definition of GP/FM, and its implications for general practitioners/family doctors, researchers and policy makers. *The European Journal of General Practice* publishes a series of articles based on this document. In a first article, background, objectives, and methodology were discussed. In this second article, the results for the core competencies 'primary care management' and 'community orientation' are presented. Though there is a large body of research on various aspects of 'primary care management', it represents a very scattered rather than a meta view. Many studies focus on care for specific diseases, the primary/secondary care interface, or the implications of electronic patient records. Cost efficiency or process indicators of quality are current outcomes. Current literature on community orientation is mainly descriptive, and focuses on either care for specific diseases, or specific patient populations, or on the uptake of preventive services. Most papers correspond poorly to the WONCA concept. For both core competencies, there is a lack of research with a longitudinal perspective and/or relevant health or quality of life outcomes as well as research on patients' preferences and education for organizational aspects of GP/FM.

Key words: *General practice/family medicine, primary care management, community oriented health care, research agenda*

¹Based on: Hummers-Pradier E, et al. Research Agenda for General Practice/Family Medicine and Primary Health Care in Europe. Maastricht: European General Practice Research Network, 2009. pp. 13–15, 27–28.

Background

The 'Research Agenda for General Practice/Family Medicine and Primary Healthcare in Europe' was published in September 2009 by the European




Figure 1. The WONCA tree: Core competencies and characteristics of general practice/family medicine.

General Practice Research Network (EGPRN) (1). It summarizes the evidence relating to the core competencies and characteristics of the WONCA Europe definition of General Practice/Family Medicine (GP/FM) (Figure 1) (2), and indicates evidence gaps and research needs. The European Journal of General Practice presents this document as a series of articles. Background, objectives and methodology were presented and discussed in part 1. Briefly, the Research Agenda is based on international key informant surveys and expert consensus and a comprehensive literature review on research domains related to each of the six core competencies of the European definition, covering health services research, clinical research, educational research and methodology issues (3). In this and the following issues, the results will be presented. This article reflects on the two core competencies which deal primarily with organizational aspects of GP/FM, i.e. 'primary care management' and 'community orientation'.

Definition of the research domains

According to the WONCA Europe definition of GP/FM (2), *primary care management* includes the ability to:

- Manage primary contact with patients, dealing with unselected problems, and providing open and unlimited access;
- Cover the full range of health conditions;
- Make available to the patient the appropriate services within the health care system;

- Coordinate care with other professionals in primary care, other specialists and secondary care;
- Master effective and appropriate care provision and health service utilisation, using resources efficiently;
- Act as an advocate for the patient, i.e. protecting them from harm which may ensue through unnecessary screening, testing and treatment.

Our research domain also includes the clinical effectiveness and health systems effects of models of managing particular health problems in primary care, i.e. defined disease management programmes, and ways of organizing care within a practice or primary health care team. Educational research in this field comprises management skills at a health system and practice level, as well as education for collaborating medical professionals with a range of backgrounds and expertise.

The core competency of *community orientation* includes the ability to reconcile the health needs of individual patients and the health needs of the community in balance with available resources (2). Presentation of the paradigm of community oriented medicine started with work of Kark in the 1950s and 1960s (4), and received a more structured definition during the 1980s. According to this definition, the following topics can be included in the research domain: health needs reflecting individual health needs in the context of a person's environment, as well as community health needs, and possible conflicts between these two. It also includes the specific context-related decision making process, and cooperation with other professionals and agencies according to these health needs.

Table I. Search strategies: Primary care management.

-
- ‘organization and administration’ [MeSH Terms] combined with ‘primary health care’ [Majr MeSH] and/or ‘family practice’ [MeSH]
 - ‘practice management’ combined with ‘primary health care’ [Majr MeSH] and/or ‘family practice’ [MeSH], ‘health services’ [MeSH], ‘education, medical’ [Majr MeSH]
 - ‘health services accessibility’ [Majr MeSH] combined with ‘primary health care’ [Majr MeSH] and/or ‘family practice’ [MeSH]
 - ‘medical records systems, computerized’ [MeSH] combined with ‘primary health care’ [Majr MeSH] and/or ‘family practice’ [MeSH]
-

The research domains of primary care management and community orientation overlap with each other, and to some extent also with the competencies of ‘patient centred care’, ‘specific problem solving skills’, and ‘comprehensive approach’. These research domains also reflect three of Starfield’s four central components of primary care, i.e. accessibility, coordination (defined as the degree to which the primary care provider manages all the patient’s health care and possesses the necessary infrastructure to do so) and comprehensiveness (there defined as the provision of a range of services broad enough to meet all common needs in the population) (5–7). Her fourth component, longitudinality or continuity, will be considered with the core competency of ‘person-centred care’ in this research agenda.

Methodology—overview of search strategies

A general description of the methodology of our evaluation—key informant surveys, a comprehensive literature review and expert consensus—was presented in the first part of this series (3).

Literature on primary care management was sought using the MeSH terms and combinations shown in Table I.

As there is no explicit MeSH term for community oriented primary care, combinations of several terms were used, as shown in Table II.

Additional searches using ‘seek related articles’ options, MeSH terms of relevant articles, free text searches or search strings not limited to ‘family practice’ or ‘primary health care’ were used to extend the overview. Literature was reviewed and consented conclusions were drawn according to the procedure described in part 1 of this series (3).

Results

Primary care management

The research field of primary care management is very large. The retrieved literature gave very scattered results, rather than a meta-view. Although there have been few systematic comparisons, it seems that there is little evidence in favour of any particular *organizational, funding or workforce model*. However, it seems obvious that the organization and workforce of general practice has to be developed further in order to meet current and future requirements of primary care management better. Evidence shows advantages for health systems that rely relatively more on primary health care and general practice in comparison to those systems tending towards specialist care, in terms of better population health outcomes, improved equity, access and continuity and lower cost (8,9).

Common *outcome measures* in research on primary care management aspects included effectiveness with regard to quality aspects/quality indicators, often with a benchmarking approach, or efficiency with regard to costs (10–19). Outcomes which reliably reflect patients’ health or well-being or Starfield’s central components or indicators (5) were rarely used.

Many of the retrieved papers were related to the management in primary care of patients with a specific disease (very often depression or other mental conditions, or diabetes) or of a defined patient population, i.e. geriatric care (20–24). The effect of various *primary care management models or interventions*, such as outreach preventive visits or care by nurse practitioners, was studied in different patient populations. Several studies suggest that for some well defined conditions, quality of care provided by appropriately trained nurses is as high as care provided by

Table II. Search strategies: Community orientation.

-
- (‘community networks’ [MeSH] OR ‘community health services’ [MeSH] OR ‘community health planning’ [MeSH] OR ‘community-institutional relations’ [MeSH] OR ‘community health aides’ [MeSH] OR ‘community health nursing’ [MeSH] OR ‘community health centres’ [MeSH] OR ‘community medicine’ [MeSH] OR ‘consumer participation’ [MeSH] OR ‘delivery of health care’ [MeSH]) AND (‘primary health care’ [Majr MeSH] OR ‘family practice’ [MeSH] NOT ‘public health’ [MeSH])
 - (‘minority groups’ [MeSH] AND ‘health services needs and demand’ [MeSH] OR ‘community networks/utilization’) AND (‘primary health care’ [Majr MeSH] OR ‘family practice’ [MeSH])
 - ‘residence characteristics’ [MeSH] AND (‘primary health care’ [Majr MeSH] OR ‘family practice’ [MeSH])
 - ‘community’ AND (‘family practice’ [MeSH] OR ‘physicians, family’ [MeSH]) AND ‘education’ [Majr MeSH]
-

doctors and health outcomes for patients are comparable. However, primary care by nurse practitioners is likely to cost as much as care provided by (salaried) GPs according to currently available data. Interventions on practice organization seem to influence service uptake, but the effect on health outcomes was rarely studied (25,26).

The impact of *consultation length* has been studied in observational studies, but without conclusive findings. Further trials are needed focusing on health outcomes and cost effectiveness.

Access to primary care is differently organized across various countries both inside and outside Europe. Many of the retrieved papers were disease related studies or nursing research papers. Improving access is a key policy issue in improving quality of (primary) care and in guaranteeing equity in health care, but until now the topic has mostly been studied from a local point of view rather than as a general characteristic or in a comparative way (27–32).

An important focus of research was *collaborative care* and the *interface between primary and secondary care*. However, this interface is organized very differently in various European countries, implying that studies have to be interpreted in their local context and cannot really be generalized to another setting. Much research has been conducted with regard to referral rates and it shows a wide variation between individual general practitioners. Local educational interventions actively involving GPs and secondary care specialists, and structured referral sheets are the only intervention shown to have an impact on referral rates based on current evidence. The effects of an ‘in-house’ second opinion and other intermediate primary care based alternatives to outpatient referral appear promising in settings with otherwise strict gate-keeping by GPs (33–44). Cooperation with pharmacists (i.e. for control or coordination of prescriptions) may possibly reduce medication-related adverse events. More randomized controlled trials of primary care based pharmacist-led interventions are needed, to decide on the effectiveness of this (45).

There has been quite a lot of research on the role and potential effects of *electronic medical records (EMR)* in primary care. The use of *ICPC* and coding of GPs’ daily practice activity based on patients’ reasons for encounter was a central concept of many papers (46). These show the usefulness, potential and possibilities for further development of primary care epidemiology based on the electronic medical records coded with *ICPC* and structured according to episodes of care (47–50). However, in many European countries medical record utilisation and quality are less than ideal from the perspectives of primary care epidemiology or data collection for national and

international research databases. This is caused by either the lack of standard classifications, or by national legislation imposing use of ICD based coding. Often, this results in incomplete records. For research purposes, a pragmatic approach to ‘everyday’ EMR or other routine data, i.e. health insurance or billing data used for secondary analyses, must be adopted (51,52). There is a rapidly increasing body of literature both on methodological and quality issues of research on EMR/routine data, and on research projects using such records as a data source, for example eHID, QRESEARCH and other morbidity/EMR databases in the UK, Belgium and the Netherlands (53–57). A substantial proportion of research papers dealt with the potential of these databases to generate data for quality management (including audits) and of EMRs to represent a platform for implementing guidelines or recommendations, to identify patients eligible for treatment or preventive services, or to prompt drug warnings. Results obtained were mostly positive, but not overwhelmingly so, which may depend on the EMR system used. Effectiveness with regard to health outcomes is rarely studied.

There has been a limited amount of educational research on primary care management aspects. Of the studies performed, the vast majority focuses on *educational interventions aimed at doctors* to implement specific service approaches, for example, prevention activities, inter-professional collaboration, or care strategies for specific diseases, detection of disease, and prescribing. Most studies showed either small or insignificant effects; sustainability of these effects has not been studied (58–62).

Community orientation

Community orientation seems to be a rather new competence. Several articles from recent decades were descriptive, explanatory opinion papers. They attempted to define the concept of community oriented primary care and described its development (4,63–68). The English language concept of ‘community’ includes both small entities such as a family, and large communities such as a school, a city or a country. This renders the concept difficult to study and results in considerable overlap between public health and primary care research.

Not unexpectedly, many narrative and descriptive reports were retrieved. A lot of research literature was related to nursing rather than GP/FM. There were few research articles on community oriented primary care (COPC), and most lacked control groups or comparators. Research considering relevant, GP related outcomes was missing. There were some studies exploring health needs with a combination of qualitative and quantitative methods, but in general

there is a lack of qualitative research. Most studies focused on very specific issues, either care for defined diseases (mental conditions, common chronic diseases like diabetes or arthritis), or patient/population groups (geriatric patients, mothers and newborns, minorities or deprived groups) or on preventive services (vaccinations, screening, smoking cessation, dentistry) (69–91). They were community or population based, but were not community orientated from a primary care point of view. Thus, most of the current research did not really cover the concept of a community oriented approach as described in the European definition of GP/FM (92–97).

Some articles described methods of disease management in COPC involving cooperation between primary care and institutions in the community, i.e. GPs' referral to several community programmes, or collaborative care (98–104), or evaluated the implementation of a COPC model in local communities (105,106). Regarding educational research, there were several articles describing educational programmes on community related themes, or teaching experiences with students, but very few evaluations (107–114). The majority of studies were not controlled.

Implications

Research

Given these results, further research should focus on:

- Developing research instruments and outcome measures reflecting and measuring the different aspects of primary care management and community orientation, and their convergence;
- Patient and doctor's perceptions, perspectives and preferences on practice management issues (such as open access, telephone consultations, telemedicine);
- Comparing different models of care and evaluating effectiveness of different primary care management strategies or interventions, not only at the level of patient satisfaction and/or service uptake, but also on the health outcome level;
- Validity and utility of electronic patient records in a general practice; and use of information technologies in COPC;
- Routine collection and the feasibility/validity of data from GPs' electronic medical records, their use for studies of morbidity and GP care (incl. appropriate denominators), and as a means for recruitment, data collection and data management in research;
- Comparing different approaches/models of primary health care in the community, regarding outcomes with respect to both individual health and community needs (115);
- Community based care models for specific areas of clinical work (for example palliative care, drug addiction programmes);
- Effective methods of inter-professional education and teaching management skills to (future) GPs;
- Education for COPC, including the evaluation of programmes with a clear methodology.

Research methodology

The following methodological needs can be formulated:

- Instrumental research to develop and validate measures for practice management issues and aspects of community orientation;
- Longitudinal observational studies, i.e. on epidemiology of GP/FM, including specific aspects and outcomes of care, and looking at both individuals and the community;
- Interventional research (controlled trials—comparing different primary care management strategies, or comparing innovative strategies with 'care as usual');
- Implementation studies of effective strategies (observational);
- Observational cohort studies comparing different approaches and models, also on education;
- Mixed design studies.

It can be concluded that much of the current research focussed on specific diseases rather than a generalist perspective. Outcomes are usually process indicators or cost efficiency, research with regard to health outcomes is rare. Though there is a large body of scientific literature on organizational aspects of primary care, many essential topics are not or not sufficiently well studied.

Acknowledgments

The authors acknowledge the support and contribution of many organizations and persons. The full text can be read in the first article of this series (*Eur J Gen Pract.* 2009;15:243–50).

Declaration of interest: The authors report having no conflicts of interest. The authors alone are responsible for the content and writing of the paper. All authors are members of EGPRN and active in its committees. Additionally, EHP is member of the WONCA Europe Executive Board. The Research Agenda was supported solely by EGPRN and grants from WONCA Europe.

Full text versions of the research agenda

Electronic versions (pdf) are available from: <http://www.egprn.org>. Paper versions can be requested from the Coordinating Centre of EGPRN, Mrs Hanny Prick. E-mail: hanny.prick@hag.unimaas.nl

References

- Hummers-Pradier E, Beyer M, Chevallier P, Eilat-Tsanani S, Lionis C, Peremans L, et al. Research agenda for general practice / family medicine and primary health care in Europe. Maastricht: European General Practice Research Network EGPRN, 2009.
- WONCA-Europe definition of Family Medicine. 2005. <http://www.woncaeurope.org/> (accessed 11 December 2009).
- Hummers-Pradier E, Beyer M, Chevallier P, Eilat-Tsanani S, Lionis C, Peremans L, et al. The research agenda for general practice/family medicine and primary health care in Europe. Part 1. Background and methodology. *Eur J Gen Pract.* 2009;15:243–50.
- Kark SL, Kark E. An alternative strategy in community health care: community-oriented primary health care. *Isr J Med Sci.* 1983;19:707–13.
- Starfield B. Is primary care essential? *Lancet* 1994;344:1129–33.
- Forrest CB, Starfield B. The effect of first-contact care with primary care clinicians on ambulatory health care expenditures. *J Fam Pract.* 1996;43:40–8.
- Haggerty JL, Pineault R, Beaulieu MD, Brunelle Y, Gauthier J, Goulet F, et al. Practice features associated with patient-reported accessibility, continuity, and coordination of primary health care. *Ann Fam Med.* 2008;6:116–23.
- Starfield B, Shi L, Macinko J. Contribution of primary care to health systems and health. *Milbank Q.* 2005;83:457–502.
- WHO Europe. What are the advantages and disadvantages of restructuring a health care system to be more focused on primary care services? WHO 2004 <http://www.euro.who.int/Document/E82997.pdf> (accessed 11 December 2009).
- Bunn F, Byrne G, Kendall S. Telephone consultation and triage: Effects on health care use and patient satisfaction. *Cochrane Database Syst Rev.* 2004;(4):CD004180.
- Ward D, Severs M, Dean T, Brooks N. Care home versus hospital and own home environments for rehabilitation of older people. *Cochrane Database Syst Rev.* 2008;(4):CD003164.
- Gruen RL, Weeramanthri TS, Knight SE, Bailie RS. Specialist outreach clinics in primary care and rural hospital settings. *Cochrane Database Syst Rev.* 2004;(1):CD003798.
- Boult C, Reider L, Frey K, Leff B, Boyd CM, Wolff JL, et al. Early effects of 'Guided Care' on the quality of health care for multimorbid older persons: A cluster-randomized controlled trial. *J Gerontol A Biol Sci Med Sci.* 2008;63:321–7.
- Lauritzen T, Jensen MS, Thomsen JL, Christensen B, Engberg M. Health tests and health consultations reduced cardiovascular risk without psychological strain, increased healthcare utilization or increased costs. An overview of the results from a 5-year randomized trial in primary care. The Ebeltoft Health Promotion Project (EHPP). *Scand J Public Health* 2008;36:650–61.
- Kendrick D, Fielding K, Bentley E, Miller P, Kerslake R, Pringle M. The role of radiography in primary care patients with low back pain of at least 6 weeks duration: a randomised (unblinded) controlled trial. *Health Technol Assess.* 2001;5:1–69.
- Richards SH, Bankhead C, Peters TJ, Austoker J, Hobbs FD, Brown J, et al. Cluster randomised controlled trial comparing the effectiveness and cost-effectiveness of two primary care interventions aimed at improving attendance for breast screening. *J Med Screen* 2001;8:91–8.
- van Wijk MA, van der Lei J, Mosseveld M, Bohnen AM, van Bommel JH. Assessment of decision support for blood test ordering in primary care. A randomized trial. *Ann Intern Med* 2001;134:274–81.
- Shannon GR, Wilber KH, Allen D. Reductions in costly healthcare service utilization: findings from the care advocate program. *J Am Geriatr Soc.* 2006;54:1102–7.
- Black MM, Nair P, Kight C, Wachtel R, Roby P, Schuler M. Parenting and early development among children of drug-abusing women: Effects of home intervention. *Pediatrics* 1994;94:440–8.
- Kendrick T, Burns T, Freeling P. Randomised controlled trial of teaching general practitioners to carry out structured assessments of their long term mentally ill patients. *Br Med J.* 1995;311:93–8.
- Thompson C, Kinmonth AL, Stevens L, Peveler RC, Stevens A, Ostler KJ, et al. Effects of a clinical-practice guideline and practice-based education on detection and outcome of depression in primary care: Hampshire depression project randomised controlled trial. *Lancet* 2000;355:185–91.
- Downs M, Turner S, Bryans M, Wilcock J, Keady J, Levin E, et al. Effectiveness of educational interventions in improving detection and management of dementia in primary care: cluster randomised controlled study. *Br Med J.* 2006;332:692–6.
- Chiba N, Veldhuijzen Van Zanten SJ, Escobedo S, Grace E, Lee J, Sinclair P, et al. Economic evaluation of Helicobacter pylori eradication in the CADET-Hp randomized controlled trial of Helicobacter pylori-positive primary care patients with uninvestigated dyspepsia. *Aliment Pharmacol Ther.* 2004;19:349–58.
- Reynolds CF 3rd, Degenholtz H, Parker LS, Schulberg HC, Mulsant BH, Post E, et al. Treatment as usual (TAU) control practices in the PROSPECT Study: managing the interaction and tension between research design and ethics. *Int J Geriatr Psychiatry* 2001;16:602–8.
- Kendrick T, Simons L, Mynors-Wallis L, Gray A, Lathlean J, Pickering R, et al. A trial of problem-solving by community mental health nurses for anxiety, depression and life difficulties among general practice patients. The CPN-GPstudy. *Health Technol Assess.* 2005;9:1–104.
- Laurant M, Reeves D, Hermens R, Braspenning J, Grol R, Sibbald B. Substitution of doctors by nurses in primary care. *Cochrane Database Syst Rev.* 2005;(2):CD001271.
- Forrest CB, Starfield B. Entry into primary care and continuity: The effects of access. *Am J Public Health* 1998;88:1330–6.
- Rohrer JE, Bernard M, Naessens J, Furst J, Kircher K, Adamson S. Impact of open-access scheduling on realized access. *Health Serv Manage Res.* 2007;20:134–9.
- Lasser KE, Mintzer IL, Lambert A, Cabral H, Bor DH. Missed appointment rates in primary care: The importance of site of care. *J Health Care Poor Underserved* 2005;16:475–86.
- Bundy DG, Randolph GD, Murray M, Anderson J, Margolis PA. Open access in primary care: Results of a North Carolina pilot project. *Pediatrics* 2005;116:82–7.
- Solberg LI, Maciosek MV, Sperl-Hillen JM, Crain AL, Engebretson KI, Asplin BR, et al. Does improved access to care affect utilization and costs for patients with chronic conditions? *Am J Manag Care* 2004;10:717–22.

32. Smits FT, Brouwer HJ, Ter Riet G, van Weert HH. Epidemiology of frequent attenders: A 3-year historic cohort study comparing attendance, morbidity and prescriptions of one-year and persistent frequent attenders. *BMC Public Health* 2009;9:36.
33. Akbari A, Mayhew A, Al-Alawi MA, Grimshaw J, Winkens R, Glidewell E, et al. Interventions to improve outpatient referrals from primary care to secondary care. *Cochrane Database Syst Rev.* 2008;(4):CD005471.
34. Slade M, Gask L, Leese M, McCrone P, Montana C, Powell R, et al. Failure to improve appropriateness of referrals to adult community mental health services—lessons from a multi-site cluster randomized controlled trial. *Fam Pract.* 2008;25:181–90.
35. Gruen RL, Weeramanthri TS, Knight SE, Bailie RS. Specialist outreach clinics in primary care and rural hospital settings. *Cochrane Database Syst Rev.* 2004;(1):CD003798.
36. Smith SM, Allwright S, O'Dowd T. Effectiveness of shared care across the interface between primary and specialty care in chronic disease management. *Cochrane Database Syst Rev.* 2007;(3):CD004910.
37. Smith SM, Allwright S, O'Dowd T. Does sharing care across the primary-specialty interface improve outcomes in chronic disease? A systematic review. *Am J Manag Care* 2008;14:213–24.
38. Faulkner A, Mills N, Bainton D, Baxter K, Kinnersley P, Peters TJ, et al. Systematic review of the effect of primary care-based service innovations on quality and patterns of referral to specialist secondary care. *Br J Gen Pract.* 2003;53:878–84.
39. Liu CF, Hedrick SC, Chaney EF, Heagerty P, Felker B, Hasenberg N, et al. Cost-effectiveness of collaborative care for depression in a primary care veteran population. *Psychiatr Serv.* 2003;54:698–704.
40. Rossignol M, Abenheim L, Séguin P, Neveu A, Collet JP, Ducruet T, et al. Coordination of primary health care for back pain. A randomized controlled trial. *Spine* 2000;25:251–8.
41. Kunz R, Wegscheider K, Guyatt G, Zielinski W, Rakowsky N, Donner-Banzhoff N, et al. Impact of short evidence summaries in discharge letters on adherence of practitioners to discharge medication. A cluster-randomised controlled trial. *Qual Saf Health Care* 2007;16:456–61.
42. Morrison J, Carroll L, Twaddle S, Cameron I, Grimshaw J, Leyland A, et al. Pragmatic randomised controlled trial to evaluate guidelines for the management of infertility across the primary care-secondary care interface. *Br Med J.* 2001;322:1282–4.
43. Casas A, Troosters T, Garcia-Aymerich J, Roca J, Hernández C, Alonso A, et al. Members of the CHRONIC project. Integrated care prevents hospitalisations for exacerbations in COPD patients. *Eur Respir J.* 2006;28:123–30.
44. Vlek JF, Vierhout WP, Knottnerus JA, Schmitz JJ, Winter J, Wesselingh-Megens AM. A randomised controlled trial of joint consultations with general practitioners and cardiologists in primary care. *Br J Gen Pract.* 2003;53:108–12.
45. Watson MC, Bond CM, Grimshaw JM, Mollison J, Ludbrook A, Walker AE. Educational strategies to promote evidence-based community pharmacy practice: a cluster randomized controlled trial (RCT). *Fam Pract.* 2002;19:529–36.
46. ICPC-2 international classification of primary care, second edition. Prepared by the international classification committee of WONCA. Oxford: Oxford University Press; 1998.
47. Okkes I, Jamoulle M, Lamberts H, Bentzen N. ICPC-2-E: the electronic version of ICPC-2. Differences from the printed version and the consequences. *Fam Pract.* 2000;17:101–7.
48. Okkes IM, Polderman GO, Fryer GE, Yamada T, Bujak M, Oskam SK, et al. The role of family practice in different health care systems. A comparison of reasons for encounter, diagnoses, and interventions in primary care populations in the Netherlands, Japan, Poland, and the United States. *J Fam Pract.* 2002;51:72–3.
49. Soler JK, Okkes I, Lamberts H, Wood M. The coming of age of ICPC: celebrating the 21st birthday of the international classification of primary care. *Fam Pract.* 2008;25:312–7.
50. Woodwell DA. National ambulatory medical care survey: 1997 summary. *Adv Data;* 1999. p. 1–28.
51. Wood L, Martinez C. The general practice research database: Role in pharmacovigilance. *Drug Saf.* 2004;27:871–81.
52. Hippisley-Cox J, Pringle M, Cater R, Wynn A, Hammersley V, Coupland C, et al. The electronic patient record in primary care—regression or progression? A cross sectional study. *Br Med J.* 2003;326:1439–43.
53. Fleming DM, Elliott C, Pringle M. Electronic health indicator data eHID 2008 http://ec.europa.eu/eahc/projects/linkedocument/sanco/2003/2003129_1_en.pdf (accessed 11 December 2009).
54. Hippisley-Cox J, Stables D, Pringle M. QRESEARCH: A new general practice database for research. *Inform Prim Care* 2004;12:49–50.
55. McCormick A, Fleming D, Carlton J. Morbidity statistics from general practice. fourth national study 1991–1992. London: Office of Population Censuses and Surveys; 1995.
56. Van Weel, C. The continuous morbidity registration Nijmegen: Background and history of a Dutch general practice database. *Eur J Gen Pract.* 2008;14(Suppl.1):5–12.
57. Bartholomeeusen S, Kim CY, Mertens R, Faes C, Buntinx F. The denominator in general practice, a new approach from the Intego database. *Fam Pract.* 2005;22:442–7.
58. Shuval K, Berkovits E, Netzer D, Hekselman I, Linn S, Brezis M, et al. Evaluating the impact of an evidence-based medicine educational intervention on primary care doctors' attitudes, knowledge and clinical behaviour: A controlled trial and before and after study. *J Eval Clin Pract.* 2007;13:581–98.
59. Bahn TJ, Cronau HR, Way DP. A comparison of family medicine and internal medicine experiences in a combined clerkship. *Fam Med.* 2003;35:499–503.
60. McNulty CA, Thomas M, Bowen J, Buckley C, Charlett A, Gelb D, et al. Improving the appropriateness of laboratory submissions for urinalysis from general practice. *Fam Pract.* 2008;25:272–8.
61. Altiner A, Brockmann S, Sielk M, Wilm S, Wegscheider K, Abholz HH. Reducing antibiotic prescriptions for acute cough by motivating GPs to change their attitudes to communication and empowering patients: a cluster-randomized intervention study. *J Antimicrob Chemother.* 2007;60:638–44.
62. Hogg W, Baskerville N, Lemelin J. Cost savings associated with improving appropriate and reducing inappropriate preventive care: Cost-consequences analysis. *BMC Health Serv Res.* 2005;5:20.
63. Nutting PA, Wood M, Conner EM. Community-oriented primary care in the United States. A status report. *JAMA* 1985;253:1763–6.
64. Nutting PA. Community-oriented primary care: an integrated model for practice, research, and education. *Am J Prev Med.* 1986;2:140–7.
65. Tollman S. Community oriented primary care: origins, evolution, applications. *Soc Sci Med.* 1991;32:633–42.
66. Nevin JE, Gohel MM. Community-oriented primary care. *Prim Care* 1996;23:1–15.

67. Longlett SK, Kruse JE, Wesleuy RM. Community-oriented primary care: Historical perspective. *J Am Board Fam Pract.* 2001;14:54–63.
68. Mullan F, Epstein L. Community-oriented primary care: New relevance in a changing world. *Am J Public Health* 2002;92:1748–55.
69. Vickrey BG, Mittman BS, Connor KI, Pearson ML, Della Penna RD, Ganiats TG, et al. The effect of a disease management intervention on quality and outcomes of dementia care: A randomized, controlled trial. *Ann Intern Med.* 2006;145:713–26.
70. Callahan CM, Boustani MA, Unverzagt FW, Austrom MG, Damush TM, Perkins AJ, et al. Effectiveness of collaborative care for older adults with Alzheimer disease in primary care: A randomized controlled trial. *JAMA* 2006;295:2148–57.
71. Counsell SR, Callahan CM, Clark DO, Tu W, Buttar AB, Stump TE, et al. Geriatric care management for low-income seniors: A randomized controlled trial. *JAMA* 2007;298:2623–33.
72. Chen PH, Rovi S, Washington J, Jacobs A, Vega M, Pan KY, et al. Randomized comparison of 3 methods to screen for domestic violence in family practice. *Ann Fam Med.* 2007;5:430–5.
73. Hay EM, Foster NE, Thomas E, Peat G, Phelan M, Yates HE, et al. Effectiveness of community physiotherapy and enhanced pharmacy review for knee pain in people aged over 55 presenting to primary care: Pragmatic randomised trial. *Br Med J.* 2006;333:995.
74. Thomas KS, Muir KR, Doherty M, Jones AC, O'Reilly SC, Basseij EJ. Home based exercise programme for knee pain and knee osteoarthritis: Randomised controlled trial. *Br Med J.* 2002;325:752.
75. van 't Veer-Tazelaar N, van Marwijk H, van Oppen P, Nijpels G, van Hout H, Cuijpers P, et al. Prevention of anxiety and depression in the age group of 75 years and over: A randomised controlled trial testing the feasibility and effectiveness of generic stepped care programme among elderly community residents at high risk of developing anxiety and depression versus usual care. *BMC Public Health* 2006; 6:186.
76. Short LM, Surprenant ZJ, Harris JM Jr. A community-based trial of an online intimate partner violence CME program. *Am J Prev Med.* 2006;30:181–5.
77. Clancy DE, Yeager DE, Huang P, Magruder KM. Further evaluating the acceptability of group visits in an uninsured or inadequately insured patient population with uncontrolled type 2 diabetes. *Diabetes Educ.* 2007;33:309–14.
78. Parker DR, Assaf AR. Community interventions for cardiovascular disease. *Prim Care.* 2005;32:865–81.
79. Széles G, Vokó Z, Jenei T, Kardos L, Pocsai Z, Bajtay A, et al. A preliminary evaluation of a health monitoring programme in Hungary. *Eur J Public Health.* 2005;15: 26–32.
80. Wiesemann A, Metz J, Nuessel E, Scheidt R, Scheuermann W. Four years of practice-based and exercise-supported behavioural medicine in one community of the German CINDI area. (Countrywide integrated non-communicable diseases intervention). *Int J Sports Med.* 1997;18: 308–15.
81. Kreling BA, Cañar J, Catipon E, Goodman M, Pallesen N, Pomeroy J, et al. Latin American cancer research coalition. Community primary care/academic partnership model for cancer control. *Cancer* 2006;107:2015–22.
82. Murray SA. Experiences with 'rapid appraisal' in primary care: involving the public in assessing health needs, orientating staff, and educating medical students. *Br Med J.* 1999;318:440–4.
83. Hopton JL, Dlugolecka M. Need and demand for primary health care: A comparative survey approach. *Br Med J.* 1995;310:1369–73.
84. Pepall E, Earnest J, James R. Understanding community perceptions of health and social needs in a rural Balinese village: Results of a rapid participatory appraisal. *Health Promot.* 2007;22:44–52.
85. Fisher B, Neve H, Heritage Z. Community development, user involvement, and primary health care. *Br Med J.* 1999;318:749–50.
86. Wilkinson JR, Murray SA. Health needs assessment: Assessment in primary care: practical issues and possible approaches. *Br Med J.* 1998;316:1524–8.
87. Murray SA, Graham LJ. Practice based health needs assessment: Use of four methods in a small neighbourhood. *Br Med J.* 1995;310:1443–8.
88. Jackson AK. Cultural competence in health visiting practice: A baseline survey. *Community Pract.* 2007;80:17–22.
89. Gorin SS, Ashford AR, Lantigua R, Hossain A, Desai M, Troxel A, et al. Effectiveness of academic detailing on breast cancer screening among primary care physicians in an underserved community. *J Am Board Fam Med.* 2006;19:110–21.
90. Flores G, Fuentes-Afflick E, Barbot O, Carter-Pokras O, Claudio L, Lara M, et al. The health of Latino children: Urgent priorities, unanswered questions, and a research agenda. *JAMA* 2002;288:82–90.
91. Giachello AL, Arrom JO, Davis M, Sayad JV, Ramirez D, Nandi C, et al. Chicago Southeast Diabetes Community Action Coalition. Reducing diabetes health disparities through community-based participatory action research: The Chicago Southeast Diabetes Community Action Coalition. *Public Health Rep.* 2003;118:309–23.
92. Mettee TM, Martin KB, Williams RL. Tools for community-oriented primary care: a process for linking practice and community data. *J Am Board Fam Pract.* 1998;11:28–33.
93. Britt H, Scahill S, Miller G. ICPC PLUS for community health? A feasibility study. *Health Inf Manag.* 1997–1998; 27:171–5.
94. Harzheim E, Duncan BB, Stein AT, Cunha CR, Goncalves MR, Trinidad TG, et al. Quality and effectiveness of different approaches in primary care in Brazil. *BMC Health Serv Res.* 2006;6:156.
95. Fletcher AE, Price GM, Ng ES, Stirling SL, Bulpitt CJ, Breeze E, et al. Population-based multidimensional assessment of older people in UK general practice: A cluster-randomised factorial trial. *Lancet* 2004;364:1667–77.
96. Westfall J, Stevenson J. A guided tour of community-based participatory research: an annotated bibliography. *Ann Fam Med.* 2007;5:185–6.
97. Stevenson J. Recent publications regarding community-based participatory research. <http://www.fndrl.org/879> (accessed 11 December 2009).
98. Proser M. Deserving the spotlight: health centers provide high-quality and cost-effective care. *J Ambulat Care Manage.* 2005;28:321–30.
99. Proser MJ, Christianson JB, Grogan CM. Alternative models for the delivery of rural health services. *Rural Health* 1990;6:419–36.
100. Giesen P, van Lin N, Mookink H, van den Bosch W, Grol R. General practice cooperatives: long waiting times for home visits due to long distances? *BMC Health Serv Res.* 2007;7:19.
101. van Uden CJ, Zwietering PJ, Hobma SO, Ament AJ, Wesseling G, et al. Follow-up care by patient's own general practitioner after contact with out-of-hours care. A descriptive study. *BMC Fam Pract.* 2005;6:23.
102. Connor A, Rainer LP, Simcox JB, Thomisee K. Increasing the delivery of health care services to migrant farm worker

- families through a community partnership model. *Public Health Nurs.* 2007;24:355-60.
103. Cunningham CO, Sohler NL, Wong MD, Relf M, Cunningham WE, Drainoni ML, et al. Utilization of health care services in hard-to-reach marginalized HIV-infected individuals. *AIDS Patient Care STDS* 2007;21:177-86.
 104. Knightbridge SM, King R, Rolfe TJ. Using participatory action research in a community-based initiative addressing complex mental health needs. *Aust NZ J Psychiatry* 2006;40:325-32.
 105. Gillam S, Schamroth A. The community-oriented primary care experience in the United Kingdom. *Am J Public Health* 2002;92:1721-5.
 106. Iliffe S, Lenihan P. Integrating primary care and public health: learning from the community-oriented primary care model. *Int J Health Serv.* 2003;33:85-98.
 107. Klevens J, Valderrama C, Restrepo O, Vargas P, Casasbuenas M, Avella MM. Teaching community oriented primary care in a traditional medical school: a two year progress report. *J Community Health* 1992;17:231-45.
 108. Brill JR, Ohly S, Stearns MA. Training community-responsive physicians. *Acad Med.* 2002;77:747.
 109. Albritton TA, Wagner PJ. Linking cultural competency and community service: a partnership between students, faculty, and the community. *Acad Med.* 2002;77:738-9.
 110. Dobbie A, Kelly P, Sylvia E, Freeman J. Evaluating family medicine residency COPC programs: meeting the challenge. *Fam Med.* 2006;38:399-407.
 111. Dornan T, Littlewood S, Margolis SA, Scherpbier A, Spencer J, Ypinazar V. How can experience in clinical and community settings contribute to early medical education? A BEME systematic review. *Med Teach.* 2006;28:3-18.
 112. Longlett SK, Kruse JE, Wesley RM. Community-oriented primary care: critical assessment and implications for resident education. *J Am Board Fam Pract.* 2001;14:141-7.
 113. Tamblyn R, Abrahamowicz M, Dauphinee D, Girard N, Bartlett G, Grand'Maison P, et al. Effect of a community oriented problem based learning curriculum on quality of primary care delivered by graduates: Historical cohort comparison study. *Br Med J.* 2005;331:1002.
 114. Peleg R, Biderman A, Polaceck Y, Tandeter H, Scvartzman P. The family medicine clerkship over the past 10 years at Ben Gurion University of the Negev. *Teach Learn Med.* 2005; 17:258-61.
 115. van Weel C, De Maeseneer J, Roberts R. Integration of personal and community health care. *Lancet* 2008;372:871-2.