

# Characterising brines in deep Mesozoic sandstone reservoirs, Denmark

Hanne D. Holmslykke<sup>\*1</sup>, Niels H. Schovsbo<sup>1</sup>, Lars Kristensen<sup>1</sup>, Rikke Weibel<sup>1</sup> and Lars Henrik Nielsen<sup>1</sup>

RESEARCH ARTICLE | OPEN ACCESS

GEUS Bulletin Vol 43 | e2019430104 | Published online: 17 July 2019

<https://doi.org/10.34194/GEUSB-201943-01-04>

The Danish subsurface contains several sandstone units, which represent a large geothermal resource (Vosgerau *et al.* 2016). Currently, only three geothermal plants are operating in Denmark, but several exploration licences are expected to be awarded in 2019. Geothermal energy is exploited from deeply buried porous sandstones by bringing warm formation water (brine) to the surface, extracting the heat and

returning the cooled water to the same sandstones. The reduced temperature of the brine during this process implies a risk of scaling, which may reduce reservoir permeability and hence injectivity. Predicting the chemical composition of formation waters, however, could help to reduce the risk associated with scaling in planned geothermal facilities.




Fig. 1. Location of the wells used for brine type characterisation and distribution of geothermal reservoirs: Skagerrak (yellow – orange), Bunter Sandstone (red), and Gassum (blue) formations. Modified from Weibel *et al.* (2017a).

Here, we present a regional overview of the geochemistry of brines from deep Mesozoic sandstones in the Danish Basin and North German Basin that supplements previous studies, notably by Laier (2002, 2008). The brine composition at shallow burial typically reflects the original (connate) formation water chemistry, which is determined by the original depositional environment of the sandstone, for example fluvial or marine. However, the mineralogical composition of the sandstone changes during burial, whereby some minerals may dissolve or precipitate when exposed to higher temperatures. These mineral changes are reflected in the brine composition, which typically becomes more saline with increased burial (e.g. Laier 2008; Kharaka & Hanor 2003).

The brine chemistry reported here shows a distinct depth trend, which reflects original connate formation waters that are modified through burial diagenesis. We have classified the brines into brine types, which are shown to be related to their depositional environment, depth, geological formation and geographical domains.

## Methods

We collected new samples from the production wells at each of the three Danish geothermal sites (Margrethholm, Sønderborg and Thisted) in 2017. The samples were analysed for pH, anions, cations and trace elements. Cations were analysed by ICP-MS (PerkinElmer Elan6100DRC Quadrupol) with a standard deviation of 3–15% depending on the element measured. Samples for anion analysis were frozen for ion-chromatography (LC50-CD50, Dionex, CA, USA) with a quantification limit of 0.05 mg/l. Total dissolved species (TDS) were calculated as the weight sum of analysed ions per weight of saltwater.




Fig. 2. Stratigraphy of reservoirs and location of brine types. Brine types are defined in Fig. 3. The broken line indicates present-day depth of 2 km below surface. **Tø**: Tønder. **Sø**: Sønderborg. **Aa**: Aars. **Fa**: Farsø. **Ti**: Thisted. **St**: Stenlille. **Ma**: Margrethholm. **Palaeo.**: Palaeozoic. **Cen.**: Cenozoic.

These new data are combined with previously published data from the northern North German Basin (Tønder-4, -5 wells, Sønderborg-1A, -2 wells) and from the Danish Basin (Fig. 1, Farsø-1, Aars-1, Stenlille-1, -19 wells, Thisted -2, -3 wells, Margrethholm-1, -2 wells; Laier, 2002, 2008; Hjuler *et al.* 2019). Collectively, these samples span the Triassic Skagerrak, Bunter Sandstone and Falster formations, the Upper Triassic – Lower Jurassic Gassum Formation and the Jurassic Fjerritslev and Haldager Sand formations (Fig. 2). Contaminated samples picked during test pumping were deselected and only samples in the late phase with stable water chemistry (notably  $K^+$  and  $Cl^-$ ) were used. A principal component analysis (PCA) was conducted on the combined dataset to classify the formation water chemistry into one of three brine types (Table 1). Altogether, the new samples and the samples from the literature total 39 samples (Fig. 3). In the PCA, present day vertical depth is used with no correction for Cenozoic uplift.

## Results and discussion

To classify the brine types we applied a PCA, which transforms a matrix of measured data ( $X$ ; comprised of  $N$  samples and  $P$  variables) into sets of projection subspaces delineated by principal components. Each component is a linear combination of all  $P$  variables, which displays variance-maximised interrelationships between the variables (Schovsbo *et al.* 2016). PCA results are presented as a score plot (Fig. 3A), which displays groupings, or clusters, of samples based on compositional similarities, alongside a loading plot (Fig. 3B), which shows the variable correlations. Finally, we quantify the proportion of the total dataset variance that can be modelled by each component (as shown in each of the axes in Fig. 3).

### Brine type classification

In the PCA model, the first two principal component axes resolve 85% of the total data variance (Fig. 3). The main trend expressed on the PCA-1 axis is high ion concentrations in water from deeply buried reservoirs, expressed as high positive PCA-1 loadings at depth (Fig. 3A) for all ions except  $SO_4^{2-}$  (Fig. 3B). The PCA-2 axis displays high positive loadings of  $Na^+$ ,  $Cl^-$ ,  $K^+$ ,  $SO_4^{2-}$  and  $Br^-$ , as well as high negative loadings of elements like  $Mg^{2+}$ ,  $Ca^{2+}$  and also at depth (Fig. 3B).

From the PCA, three main brine types can be identified based on natural groupings in the PCA-2 versus PCA-1 plot (Fig. 3A). The characteristics and occurrence of each type are presented below.

*Brine type 1* plots with negative PCA-1 and with slightly positive (type 1A) or negative (type 1B) PCA-2 score (Fig.




Fig. 3. Principal component analysis (PCA). A: Score plot. B: Loading plot of PCA on regional data. The plots model 85% of the total data variance. Variance proportions are shown along each component axis. Brine types are classified according to their groupings in A.

3A) and is characterised by relative low chloride content ( $Cl^- < 110\,000$  ppm) and low TDS (Table 1). Type 1A is  $SO_4^{2-}$  enriched, and type 1B is  $SO_4^{2-}$  depleted (Table 1). Type 1 occurs mainly in Gassum reservoirs from the Stenlille, Thisted (type 1B) and Sønderborg wells (type 1A). These three reservoirs are buried between 1.2 and 1.6 km depth.

*Brine type 2* is characterised by positive PCA-1 and negative (type 2A) or neutral (type 2B) PCA-2 scores and can be characterised compositionally by medium to high salinities ( $Cl^- > 130\,000$  ppm), and high to very high  $Ca^{2+}$  and  $Mg^{2+}$  concentrations (Fig. 3). Type 2A is  $Cl^-$  and  $K^+$  depleted relative to 2B, whereas type 2B is  $K^+$  enriched (Table 1). Brine type 2 occurs at depths greater than 2 km in a broad range of

reservoirs belonging to the Haldager Sand, Gassum, Bunter Sandstone and Skagerrak formations. Type 2A occurs exclusively in the Margretheholm area (Fig. 3A).

*Brine type 3* has intermediate PCA-1 and highly positive PCA-2 scores (Fig. 3A), reflecting high salinities ( $Cl^- > 190\,000$  ppm, Table 1). Type 3 waters are restricted to the Bunter Sandstone reservoir in the Tønder area overlying a Zechstein salt dome. In addition, halite (NaCl) is also present in the Triassic sequence at Tønder (Laier & Nielsen 1989). The brine is halite saturated and it is estimated that *c.* 5 g/L halite has precipitated at surface conditions during production tests (Hjuler *et al.* 2019).

Table 1. Brine type characteristics.

Brine Type	Depth m	$Cl^-$ ppm	$Br^-$ ppm	$SO_4^{2-}$ ppm	$Na^+$ ppm	$K^+$ ppm	$Ca^{2+}$ ppm	$Mg^{2+}$ ppm	Ca:Cl mol/mol	TDS ppm	pH	$Sr^{2+}$ ppm	$Fe_{tot}$ ppm	$Mn_{tot}$ ppm	$NH_4^+$ ppm	$Zn^{2+}$ ppm	$Ba^{2+}$ ppm	$Li^+$ ppm	$SiO_2$ ppm
1A	1250	97 264	310	869	53 382	502	4372	1167	00.04	157 866	06.06	196	19	6	-	0	1	2	-
1B	1376	105 000	372	26	56 667	421	8033	1480	00.08	171 998	06.03	498	39	15	27	11	15	10	13
3	1769	195 250	1275	595	115 000	1425	8750	1400	00.04	323 695	-	-	-	-	-	-	-	-	-
2B	2511	177 667	993	163	77 333	1838	27 950	2820	00.16	288 765	05.09	815	281	-	80	20	28	9	28
2A	2565	135 407	824	261	53 758	717	22 276	2820	00.16	216 064	06.03	862	6	20	21	5	11	11	6

TDS: Total dissolved species.  $Fe_{tot}$ : Total iron.  $Mn_{tot}$ : Total manganese.

**Table 2. Saturation index (SI) for selected minerals calculated with PHREEQC.**

Brine Type	Depth (m)	Anhydrite	Halite	Barite	Celestite
1A	1250	-0.42	-0.90	00.40	00.08
1B	1376	-1.69	-0.79	-0.06	-1.08
3	1769	00.35	00.39	-	-
2B	2511	-0.29	00.03	-0.20	-0.09
2A	2565	-0.20	-0.55	00.03	00.04

Minerals with a saturation index in the band  $-0.4 \leq SI \leq 0.4$  are assumed to be in equilibrium, within the uncertainties of analyses and thermodynamic parameters in the PHREEQC database.

### Saturation index from PHREEQC simulations

To further interpret the brine types, we performed a chemical speciation analysis using the numerical code PHREEQC and its Pitzer database. The *in situ* reservoir temperature was estimated from the regional temperature gradient (e.g. Balling *et al.* 1981). The fluid pressure of the reservoirs was assumed to be hydrostatic (9.79 KPa/m) assuming a water density of 1.1 g/cm<sup>3</sup>. The saturation state of the brines with respect to selected minerals is indicated by the Saturation Index (SI) whereby positive and negative values indicate super-saturation and undersaturation, respectively. Equilibrium with respect to the mineral is assumed for  $-0.4 \leq SI \leq +0.4$ . This accounts for the uncertainties associated with the difficulties of sampling brines at high temperature and pressure, the analytical uncertainty and the application of thermodynamic equilibrium constants on mineral phases in saline systems. Formation water with a SI within this band is assumed to be saturated, and thus in equilibrium for this mineral. Carbonate minerals are not included in Table 2, due to the difficulty of correctly measuring components of the carbonate system (pH, HCO<sub>3</sub><sup>-</sup> etc.) in heated pressurised samples.

The salinity of the brine types increases with the average depth (e.g. Cl<sup>-</sup> in brine types 1A, 1B, 2B, Table 1). This relationship was previously explained by the diffusion of Cl<sup>-</sup> from the underlying Zechstein salt deposits occurring in most of the Danish Basin and the North German Basin (Laier 2002). For the deepest reservoirs (type 2B) the brine is saturated with respect to halite (SI = 0.03; Table 2). Even though types 2A and 2B are sampled from the same depth, type 2A has a significantly lower Cl-concentration and is undersaturated with respect to halite (SI = -0.55, Table 2). This may be because brine type 2A, from Margrethholm geothermal plant, was sampled from reservoir sandstones at the basin margin, where Zechstein salt deposits are absent. Brine type 3, sampled in Tønder has a high salt content com-

pared to deeper brines (types 2A and B), and the brine is saturated with respect to halite. Brine type 3 is the only sample located in an area where halite-cemented sandstone intervals are present and salt deposits exist both above (Røth salt) and below (Zechstein salt).

Brine type 3 is saturated with respect to anhydrite (CaSO<sub>4</sub> SI = 0.35; Table 2), which makes sense as anhydrite is a common mineral in the Bunter Sandstone Formation (Weibel & Friis 2004). Despite a high SI for barite (BaSO<sub>4</sub>) in brine type 1A (SI = 0.40, Table 2), we did not observe any barite precipitation in the Margrethholm geothermal plant. The high saturation index probably reflects the uncertainties associated with sampling and analysing from deep reservoirs and most likely the formation water is in equilibrium with barite in the reservoir.

The Ca:Cl ratio (Table 1) is significantly higher in the deeper types (2A and 2B) compared to the shallower brines (types 1A, 1B and 3). Calcite (CaCO<sub>3</sub>) is common in the shallow part of the Gassum Formation, but ankerite (Ca(Mg, Fe, Mn)(CO<sub>3</sub>)<sub>2</sub>) is more abundant in the deeper parts (Weibel *et al.* 2017b). Hence, replacement of calcite with a Fe- and Mg-rich carbonate would liberate Ca<sup>2+</sup> to the formation water. Similarly, Ca<sup>2+</sup> may be liberated to the pore fluid, as dolomite (CaMg(CO<sub>3</sub>)<sub>2</sub>) becomes more stable than calcite in the deeply buried parts of the Skagerrak Formation (Weibel *et al.* 2017a).

The K<sup>+</sup> content generally increases with burial depth in the Gassum Formation (Table 1). K<sup>+</sup> may have been liberated by albite (NaAlSi<sub>3</sub>O<sub>8</sub>) replacement (albitisation) of K-feldspar (KAlSi<sub>3</sub>O<sub>8</sub>), which has been documented in Gassum sandstones from both the Aars-1 and Farsø-1 cores (Weibel *et al.* 2017b). Formation water from the Bunter Sandstone Formation generally has a similarly high or higher K<sup>+</sup> content as the deeply buried Gassum Formation. This may relate to a generally higher abundance of K-feldspar and rock fragments (Weibel & Friis 2004), or it may originate from KCl in the under or overlying evaporites.

### Conclusions

Here, we have identified and characterised three brine types present in Danish geothermal reservoirs. Reservoir depth and the occurrence of salt in the subsurface layer appear to be a dominant control. Brine type 1 occurs in reservoirs shallower than 2 km, whereas Brine type 2 occurs in both Jurassic and Triassic sandstones buried to more than 2 km. Among the deeply buried reservoirs, type 2A brine is only found in the Margrethholm wells and is interpreted to reflect the absence of salt in the subsurface around these wells in contrast to all other analysed deep wells. Brine type 3 is highly saline and occurs at less than 2 km depth in the Tønder area. Clas-

sification of brine types according to chemical composition, highlights variable risks in potential scale and scale types and shows that local conditions must be considered prior to any new planned geothermal facility. Geological information, depth and geographical domains can serve as a rough predictive tool, and will be further refined as additional data are collected from new geothermal wells.

## Acknowledgments

This contribution is part of the project GEOTHERM (“Geothermal energy from sedimentary reservoirs – Removing obstacles for large scale utilization”) (No. 6154-00011B) funded by the Innovation Fund Denmark (IFD). We thank the two reviewers, Ida Fabricius and Nicolas Marty, for their comments, which improved the manuscript.

## References

- Balling, N., Kristensen, J.I., Breiner, N., Poulsen, K.D., Rasmussen, R. & Saxov, S. 1981: Geothermal measurements and subsurface temperature modelling in Denmark. *GeoSkrifter* **16**, 173 pp.
- Hjuler, M.L., Olivarius, M., Boldreel, L.O., Kristensen, L., Laier, T., Mathiesen, A., Nielsen, C.M. & Nielsen, L.H. 2019: Multidisciplinary approach to assess geothermal potential, Tønder area, North German Basin. *Geothermics* **78**, 211–223. <https://doi.org/10.1016/j.geothermics.2018.12.001>
- Kharaka, Y.K. & Hanor, J.S. 2003: Deep fluids in the continents: I. Sedimentary Basins. In: Drevor, J.I. (ed.): *Treatise on Geochemistry* **5**, 499–540. <https://doi.org/10.1016/b0-08-043751-6/05085-4>
- Laier, T. 2002: Vurdering af udfældningsrisici ved geotermisk produktion fra Margretheholmboringen MAH-1A. Beregning af mætningsindeks for mineraler i saltvand fra Danmarks dybere undergrund. Danmarks og Grønlands Geologiske Undersøgelser Rapport **2002/95**. 48pp
- Laier, T. 2008: Chemistry of Danish saline formation waters relevant for core fluid experiments. Fluid chemistry data for lab experiments related to CO<sub>2</sub> storage in deep aquifers. Danmarks og Grønlands Geologiske Undersøgelser Rapport **2008/48**.
- Laier, T. & Nielsen, B.L. 1989: Cementing halite in Triassic Sandstone (Tønder Southwest Denmark) as a result of hyperfiltration of brines. *Chemical Geology* **76**, 353–363. [https://doi.org/10.1016/0009-2541\(89\)90103-4](https://doi.org/10.1016/0009-2541(89)90103-4)
- Schovsbo, N.H., Hedegaard, K., Holmslykke, H.D., Kjøller, C., Kristensen, L., Thomsen, E. & Esbensen, K.H. 2016: Formation water and produced water types in Danish oil and gas fields: Implications for enhanced oil recovery by “smart” water. *Geological Survey of Denmark and Greenland Bulletin* **35**, 43–46.
- Vosgerau, H. *et al.* 2016: A WebGIS portal for exploration of deep geothermal energy based on geological and geophysical data. *Geological Survey of Denmark and Greenland Bulletin* **35**, 23–26.
- Weibel, R. & Friis, H. 2004: Opaque minerals as keys for distinguishing oxidising and reducing diagenetic conditions in the Lower Triassic Bunter Sandstone, North German Basin. *Sedimentary Geology* **169**, 129–149. <https://doi.org/10.1016/j.sedgeo.2004.05.004>
- Weibel, R. *et al.* 2017a: The influence of climate on early and burial diagenesis of Triassic and Jurassic sandstones from the Norwegian – Danish Basin. *The Depositional Record* **3**, 60–91. <https://doi.org/10.1002/dep2.27>
- Weibel, R., Olivarius, M., Kristensen, L., Friis, H., Hjuler, M.L., Kjøller, C., Mathiesen, A. & Nielsen, L.H. 2017b: Predicting permeability of low enthalpy geothermal reservoirs: A case study from the Upper Triassic – Lower Jurassic Gassum Formation, Norwegian – Danish Basin. *Geothermics* **65**, 135–157. <https://doi.org/10.1016/j.geothermics.2016.09.003>

## How to cite

Holmslykke, H.D., Schovsbo, N.H., Kristensen, L., Weibel, R. & Nielsen, L.H. 2019: Characterising brines in deep Mesozoic sandstone reservoirs, Denmark. *Geological Survey of Denmark and Greenland Bulletin* **43**, e2019430104. <https://doi.org/10.34194/GEUSB-201943-01-04>

---

\*Corresponding author: Hanne D. Holmslykke | *E-mail*: [hdb@geus.dk](mailto:hdb@geus.dk)

<sup>1</sup> *Geological Survey of Denmark and Greenland (GEUS), Øster Voldgade 10, DK-1350, Copenhagen K, Denmark.*